Article Type: Spotlight

Running title: Whole-genome phylogenomics of the Tinamous

Whole-genomes illuminate the drivers of gene tree discordance and the tempo of

tinamou diversification (Aves: Tinamidae)

Authors:

Lukas J. Musher[1], Therese A. Catanach[1], Thomas Valqui[2], Robb T. Brumfield[3], Alexandre

Aleixo[4], Kevin P. Johnson[5], Jason D. Weckstein[1,6]

Author affiliations:

[1] The Academy of Natural Sciences of Drexel University, Department of Ornithology

Philadelphia, PA, 19103, USA.

[2] Facultad de Ciencias Forestales, Universidad Nacional Agraria La Molina, Lima, Peru and

CORBIDI-Centro de Ornitología y Biodiversidad, Lima, Peru.

[3] Department of Biological Sciences and Museum of Natural Science, Louisiana State

University, Baton Rouge, LA, 70803, USA.

[4] Instituto Tecnológico Vale - ITV, Belém, Brazil.

[5] Illinois Natural History Survey, Prairie Research Institute, University of Illinois Urbana-

Champaign, Champaign, IL, USA.

[6] Department of Biodiversity, Earth, and Environmental Sciences, Drexel University,

Philadelphia, PA, 19103, USA.

Abstract:
As an old group that has diversified in South America over millions of years, the tinamous (Palaeognathae: Tinamidae) are of high interest for understanding the evolution of birds and the assembly of the Neotropical biota. However, there are currently no complete species-level phylogenies of this group. Most prior work has been based on either morphological data or a small number of molecular markers, each of which has limited capability for reconstructing the tinamou phylogeny. Therefore, the interrelationships of most tinamou species are uncertain. We analyzed 80 whole-genomes from a mix of historical study skins and frozen tissues, including all 46 recognized species of tinamous to (1) reconstruct their interrelationships, (2) estimate the timeframe of tinamou evolution, and (3) examine for the effects of incomplete lineage sorting (ILS) and ancestral introgression on genome evolution. We compared results for coding (BUSCO) and ultraconserved element (UCE) loci, as well as sex-linked and autosomal markers, and used fossil-calibrated tip-dating to estimate divergence times. Tinamous diverged from their sister-group, the extinct Moas, 50-60 mya, and their crown divergence occurred roughly 30-40 mya, followed by constant diversification rates until the present. Phylogenetic reconstructions were largely robust across methods and datasets. Only one clade in the genus *Crypturellus* displayed substantial species-tree discordance across the different data sets. To investigate the impacts of introgression on this discordance, we quantified introgression for 100kb non-overlapping windows across the genome, and identified pervasive genome-wide introgression. The distribution of this introgression across the genome was dependent on the assumed phylogeny applied to the f-branch model. When assuming one of these topologies in the f-branch model, patterns of introgression matched theoretical predictions about genome architecture. Overall, we present the most complete phylogeny for tinamous to date, identify an unrecognized species, and provide a case study for species-level phylogenomic analysis using whole-genomes.

Keywords: BUSCO; Divergence dating; Fossilized Birth-Death; Gene tree; Incomplete lineage sorting; Introgression; Phylogenomic; Sex chromosome; Species tree; Stochastic error; Tinamous; Ultraconserved elements; Whole-genome sequencing; Z-chromosome

Introduction

Neotropical organisms have been at the center of discussions about the drivers of the

diversification and assembly of tropical biotas (Hughes and Eastwood 2006; Antonelli et al.

2018; Albert et al. 2020; Capurucho et al. 2023). However, most groups lack complete species-

level phylogenies, which are crucial for testing questions in these fields. To this end, the

tinamous (Palaeognathae: Tinamidae) are of key interest. First, as the only flighted group of

palaeognathous birds, their phylogenetic placement among this group is an enigma (Cracraft

1974; Harshman et al. 2008; Cloutier et al. 2019). Second, they are relatively old, having been

isolated in Neotropical regions for at least 25 million years (Claramunt and Cracraft 2015; Prum

et al. 2015; Yonezawa et al. 2017). However, there has been limited work on tinamou

phylogenomics to date. There are no complete species-level tinamou phylogenies, and prior

phylogenetic studies utilized either phenotypic data with few-to-no genetic markers (Bertelli and

Porzecanski; Bertelli et al. 2002; Valqui 2009; Bertelli 2017; Almeida et al. 2022), or

phylogenomic data with minimal taxonomic sampling (Cloutier et al. 2019). Thus, the species-

level relationships of the tinamous remain uncertain, hampering efforts to understand the

diversification of this enigmatic group.

Unfortunately, obtaining even a near-complete and robust phylogeny for tinamous

represents a considerable challenge. Although there has been a pronounced effort to collect

modern voucher-linked fresh tissues of Neotropical birds (Moncrieff et al. 2019, 2021; Del-Rio et

al. 2021), by comparison to other groups, tinamous are poorly represented in modern tissue

collections. Thus, a mix of historical samples with degraded DNA and intraspecific sampling is

needed to better document tinamou diversity and phylogenetic interrelationships (McCormack et

al. 2016). This is important because some studies have suggested that currently recognized

genera are non-monophyletic, warranting further study to reevaluate their taxonomic treatment

(Bertelli 2017; Almeida et al. 2022). Moreover, and in line with other Neotropical bird groups,

current taxonomic treatments may also misrepresent true species-level diversity (Krabbe et al.

2020; Freeman and Pennell 2021; Musher et al. 2023). Deep divergences and non-reciprocal

monophyly (i.e., paraphyly/polyphyly) of intraspecific populations suggest that some widespread

species likely include multiple species-level taxa by some criteria (Valqui 2009).

The lack of species-level phylogenies for tinamous also hampers efforts to understand

the timeframe of diversification of this group. Past studies have concluded that the crown

divergence within tinamous resulted in two major clades (subfamilies), the forest dwelling

Tinaminae and open country Nothurinae. However, current estimates for this node vary from

roughly 20 mya to as much as ~50 mya (Mitchell et al. 2014; Prum et al. 2015; Bertelli 2017;

Berv and Field 2018; Almeida et al. 2022; Stiller et al. 2024). Moreover, the origins of all extant

tinamou genera lack sufficient divergence time estimates. Despite these limitations, there exist

multiple crown fossils from both major clades, making this group a good candidate for tip-based

dating approaches (Stadler 2010; Didier et al. 2012). An advantage of tip-dating over standard

node-calibration is that it does not require the precise phylogenetic placement of fossils into the

tree. This is important because the phylogenetic positions of multiple tinamou fossils, such as

Crypturellus reai Chandler 2012 and *Nothura parvula* Brodkorb 1963, have been equivocal

(Bertelli et al. 2014; Almeida et al. 2022). Thus, tip-dating methods allow for the inclusion of

these fossils for calibration despite uncertainty about their phylogenetic affinities.

When combined with fossil data, high-throughput sequencing of historical and modern

material offers a path forward for obtaining a complete and robust dated phylogeny. Yet even

these large datasets, although highly informative, present unique problems (Reddy et al. 2017;

Zhao et al. 2023). For example, analyses that utilize alternative datasets or methodological

frameworks to answer the same phylogenetic questions have often yielded conflicting, well-

supported results (Prum et al. 2015; Simion et al. 2017; Franz et al. 2019; Schultz et al. 2023;

Stiller et al. 2024). Some uncertainty arises in the presence of rapid diversification that

increases the rate of incomplete lineage sorting (ILS) and likelihood of ancestral introgression,

two major biological sources of discordance among data partitions in phylogenetic datasets.

These processes increase discordance of gene histories, posing challenges for phylogenetic

reconstruction but also offering insights into complex evolutionary processes (Burbrink and

Gehara 2018; Rivas-González et al. 2023). An additional challenge for reconstructing rapidly

diversifying clades is the paucity of polymorphic nucleotide sites needed for reconstructing short

internodes (Leaché et al. 2015; Mclean et al. 2019). Briefly, rapid divergence events can be

difficult to reconstruct because the short timeframe also means there is limited phylogenetic

signal for those divergence events within individual gene alignments. Interestingly, the use of

sex-linked markers may help alleviate these issues as they are known to (1) evolve faster, thus

accumulating more substitutions that may be useful for reconstructing recalcitrant nodes, (2)

have smaller effective population sizes, thereby reducing rates of gene tree discordance (ILS),

and (3) experience lower rates of recombination and introgression (Charlesworth et al. 1987;

Meisel and Connallon 2013; Fontaine et al. 2015; Li et al. 2019; Musher et al. 2024). Thus, if

genealogical signal across much of the genome is associated with introgression due to

ancestral hybridization, but not the bifurcating history of species divergence (i.e., the species

tree), sex-chromosomes may be enriched with the signal of the species tree topology (Fontaine

et al. 2015; Li et al. 2019; Burbrink et al. 2025).

Here, we sequenced whole genomes for all described tinamou species and many

subspecies using a combination of frozen tissues and antique DNA sampled from historical

museum skins. Specifically, we reconstruct the evolutionary history in this group while

examining and testing for the impacts ILS and introgression on phylogenetic inference. Our

objectives are to (1) reconstruct the species-level interrelationships of tinamous, (2) estimate the

timeframe of tinamou diversification, and (3) test for the effects of ILS and introgression on

genome evolution, including differences between sex-linked and autosomal markers.

Materials and methods

Sampling

There are 46 currently recognized species of tinamous classified into nine genera

(Clements et al. 2023). Twenty-two of these species are monotypic and the remaining species

contain two or more subspecies, totaling 175 named taxa (monotypic species plus subspecies)

recognized in Tinamidae. We sampled 55 frozen tissues, 14 toe pads from museum study skins,

and downloaded 11 whole-genome assemblies from NCBI Assembly Archive, spanning a total

of 71 named taxa and all 46 recognized tinamou species (Table S1).

Tinamous belong to the infraclass, Palaeognathae which includes many extant flightless

ratites such as ostriches (Struthionidae) and rheas (Rheidae) along with extinct forms such as

Moas (Dinornithiformes). Recent work has corroborated that moas are the sister group to

tinamous and that rheas belong to a clade that is sister to tinamous plus moas (Phillips et al.

2010; Baker et al. 2014; Cloutier et al. 2019). Thus, as outgroup taxa for tree rooting, we also

downloaded whole genome assemblies for *Rhea pennata* (greater rhea) and *Anomalopteryx*

didiformis (little bush moa) from the NCBI Assembly Archive. Table S1 lists the 82 (79 Tinamous

+ two outgroup taxa) samples included in this study.

Whole-genome assembly and ortholog identification

The details of whole-genome sequencing and assembly can be found in the

Supplementary Materials document. After sequencing, we cleaned and mapped raw reads for

each sample (Table S1) to a closely related scaffold-level assembly from NCBI. Decisions about

which genome to use for reference mapping were made based on a previously published

tinamou phylogeny (Almeida et al. 2022). We used (1) fastp (Chen et al. 2018) to clean raw

reads, trim adapters and remove low quality reads, (2) Burrows-Wheeler Aligner (bwa) v0.7.17

(Li and Durbin 2009) to map the cleaned reads to reference genomes, (3) samtools v1.6 to

convert the resulting sam-files into sorted bam-files (Li et al. 2009), and (4) Analysis of Next

Generation Sequencing Data (ANGSD) v1.2.13 (Korneliussen et al. 2014) to convert bam-files

into fasta format genomes. To obtain pseudo-chromosome assemblies for each sample, we

then used ragtag v2.1.0 (Alonge et al. 2022) to scaffold each assembly against the

chromosome-level *Rhea pennata* genome (NCBI refseq assembly: GCF_028389875.1) using

default settings.

We identified and extracted two types of orthologous markers from the whole-genomes.

First, we utilized nucleotide sequences of protein-coding genes harvested using Benchmarking

of Universal Single Copy Orthologs (BUSCO) v5.3.0 (Simão et al. 2015). We used custom

scripts to append orthologous genes from all samples into the same text file before aligning.

Second, we harvested ultraconserved elements (UCEs) from each genome assembly, including

the entire conserved UCE core plus 1,000 bp of flanking sequence on both 5' and 3' ends.

UCEs are conserved, putatively single-copy, sequences in the genome that are flanked by sites

that become more variable moving away from the conserved core (Faircloth et al. 2012;

McCormack et al. 2012). We followed the pipeline outlined in Phyluce v1.7.1 (Faircloth 2016) for

harvesting loci from whole genomes using the Tetrapods-UCE-5kv1 probeset of 5,060 UCE's

(see Supplementary Materials). To examine differences among sex-linked and autosomal

markers, we also created two datasets of UCE's (see Supplementary Materials), one dataset of

sequences mapping to the Z-chromosome (an avian sex chromosome with idealized N_e of 0.75

autosomal N_e) and one that mapped to autosomes. Some analyses suggested that UCE's were

more reliable markers than BUSCO's, which had a potential impact of systematic error (see

Supplemental Materials); for this reason, we only separated the UCE dataset, and not the

BUSCO's, into sex-linked and autosomal datasets. All datasets were aligned using MAFFT

v7.453 (Katoh and Standley 2013).

To evaluate impacts of missing data, we filtered UCE and BUSCO alignments twice,

once to keep only those loci with a minimum of 75% of samples represented in each alignment

(75% complete datasets) and once to keep only those with complete sampling (100% complete

datasets). However, for autosomal and Z-linked datasets, we kept only 75% complete

alignments to maximize alignment inclusion in these two smaller datasets. Finally, each

alignment was trimmed using TrimAL v1.4 (Capella-Gutierrez et al. 2009) with a gap threshold

of 97.5% to trim out potential BUSCO mis-annotations unique to a small number of samples

(see Supplementary Materials). This resulted in six primary datasets of sequence alignments

used in our analyses for the entire Tinamou dataset (four additional datasets are discussed in

the Supplemental Materials document): (1) 75% complete BUSCO, (2) 100% complete BUSCO,

(3) 75% complete UCE, (4) 100% complete UCE, (5) 75% complete autosomal UCE, and (6)

75% complete Z-linked UCE.

Phylogenomic analyses

For each of the six datasets, we employed both a concatenation approach using IQ-

TREE v2.0.3 (Minh et al. 2020) and a multi-species coalescent (MSC) approach using ASTRAL-

III v5.7.7 (Mirarab et al. 2014; Zhang et al. 2018). For our concatenation approach, we used IQ-

TREE to concatenate all alignments and used the edge-linked proportional partition model to

partition by locus (Chernomor et al. 2016). In IQ-TREE, we implemented ModelFinder

(Kalyaanamoorthy et al. 2017) to choose the best substitution model from the standard model

set for each partition from all possible models. We then used IQ-TREE to reconstruct the

phylogeny and perform 1,000 ultrafast bootstrap replicates (Hoang et al. 2018).

Next, we estimated phylogenies for each alignment (gene trees) within each of the six

datasets, again using IQ-TREE with ModelFinder to identify the best-fit substitution model, and

again performed 1,000 ultrafast bootstraps on each gene tree. We then collapsed all low-

support (bootstrap<75) nodes within each gene tree (Simmons and Gatesy 2021). Using these

gene trees, we implemented a gene tree summarization quartet-based MSC approach in

ASTRAL-III. This approach assumes that all gene tree discordance is due to ILS without

significant ancestral introgression, recombination, or gene tree estimation error.

In addition to estimating node bootstrap support in IQ-TREE and local posterior

probability in ASTRAL-III, we assessed node support using gene concordance factors (gCF's)

for all species trees using IQ-TREE. gCF's are the percentage of gene trees that contain a given

node in a species tree.

Estimating the tempo of tinamou diversification

To estimate the timeframe of tinamou diversification, we used a fossilized birth-death

model with six crown Tinamidae tip-fossil calibrations using an optimized relaxed clock in Beast

v2.7.7 with the Sampled Ancestors and Optimized Relaxed Clock packages (Drummond et al.

2006; Gavryushkina et al. 2014; Bouckaert et al. 2019; Zhang and Drummond 2020; Douglas et

al. 2021), and conditioning on the root and rho parameters. Specifically, we filtered the

autosomal UCE dataset for the most clock-like loci that also had low RF distances between

gene and species trees using modified R scripts from a prior study (Musher et al. 2019). The

details of these analyses, along with our justification of priors, node constraints, and fossil

calibrations can be found in the Supplementary Materials (Alvarenga 1983; Bertelli and Chiappe

2005; Almeida et al. 2022). We also modeled speciation and extinction rates using the R-

package 'TESS' (Höhna et al. 2016) (see Supplementary Materials).

Evaluation of gene tree discordance

Gene tree discordance arises from multiple sources, including biological processes like

ILS and introgression and gene tree estimation error. To examine the drivers of gene tree

discordance and thereby better interpret phylogenetic results, we first used R v4.3.1 (R Core

Team 2023) to assess the relationship between the number of parsimony informative sites (PIS)

at each locus and and gene tree discordance. PIS were defined as variable sites in the

alignment wherein each variant nucleotide is represented by at least two samples. We then

measured the RF distances (Robinson and Foulds 1981) between each collapsed gene tree

and the inferred species tree for the 100% complete UCE dataset only (an examination of the

remaining datasets can be found in the Supplemental Materials), and used RF as a measure of

gene tree discordance. RF distances were quantified using the 'RF.dist' function in phangorn

v2.11.1 (Schliep 2011). To estimate RF distances, we compared the gene trees from each

dataset to their corresponding MSC tree. To test for differences in gene tree discordance, as

measured by RF distance, between autosomal and Z-linked UCE's, we used Kruskal-Wallis

tests in base R (R Core Team 2023). We then ran generalized linear models using PIS, a metric

of information content, as the independent variable and RF distance as the dependent variable

for the 100% complete UCE dataset. To characterize the relationship between information

content and gene tree topology, we compared AIC values for linear and logarithmic fits for each

regression and chose the model with the lowest AIC for each dataset. This enabled us to

examine the potential impact of stochastic error (error introduced by insufficient data) on gene

tree discordance, as the R-squared from this analysis measures the proportion of variation in

gene tree topology (RF distance) that is explained by alignment information content (PIS). If

stochastic error is absent, the slope should not significantly differ from zero and/or R^2 should be

low. From this, we could also see if Z-linked markers behaved differently from autosomal

markers, with the expectation that Z-linked UCE's should have reduced levels of ILS and

introgression relative to autosomal markers, and thus reduced gene-tree-to-species-tree

discordance as measured using RF distance. Specifically, we expect reduced RF distances for

Z-linked gene trees relative to autosomal gene trees given comparable levels of phylogenetic

information content (PIS). In other words, we expect that, while accounting for potential impacts

of stochastic error driven by insufficient data in the alignments, RF distances will be reduced for

Z-linked trees.

Because we found conflicting phylogenetic relationships between datasets for a sub-

clade in the genus *Crypturellus* (Clade A, see Results), we assessed the extent to which this

gene tree discordance could be explained by ILS due to rapid divergences or introgression

between non-sister taxa. To do so, we first filtered all UCE alignments to only include one

member of each species within Clade A (10 total plus one outgroup, as above for sliding window

datasets), and then re-filtered for 100% completeness (including fewer samples resulted in more

alignment with complete representation; complete sampling is a prerequisite for the quartet-

based ILS analysis outlined below). We then reran IQ-TREE, using the same model selection,

bootstrapping, and node collapsing as outlined above, to produce gene trees for the Clade A

datasets. We then evaluated the impacts of ILS and introgression on the relationships in Clade

A in multiple ways. Analyses found in the Supplemental Materials document, consistent with

examples in the literature (Reddy et al. 2017; Stiller et al. 2024), suggested that BUSCO's might

be enriched for systematic error, so we therefore only applied these analyses to the UCE

dataset.

First, we measured the relative quartet frequencies for two internal branches in Clade A

on both Z-linked and Autosomal UCE's. Quartets are unrooted four-taxon statements for which

only three alternative topologies exist. The MSC model assumes that all gene tree discordance

arises via ILS (i.e., no introgression, no recombination, etc.), and it makes predictions about the

relative frequencies of alternative quartet topologies for all internal branches in a phylogeny.

Specifically, the MSC predicts one majority topology consistent with the species tree at a

relative frequency of >⅓ and two minority topologies of equal frequency at <⅓ (Pamilo and Nei

1988). As ILS increases, the frequencies of all three alternative quartets approach the ⅓

threshold. Deviations from these expectations (i.e., minority topologies are not equivalent in

frequency) suggest that additional processes may be influencing gene tree discordance. For this

method, we utilized the R package 'MSCquartets' (Rhodes et al. 2021; Allman et al. 2023).

Because ILS is expected at relatively short internal branches, we evaluated the relative

frequencies of alternative quartet topologies at two internal branches of the difficult-to-resolve

clade (Clade A). To test for deviations from MSC predictions, we used a multinomial test in the

R-package EMT (Menzel 2024) using the Monte Carlo approach to perform 10,000,000

simulated withdrawals based on expected quartet frequencies. We calculated the expected

quartet frequencies for these branches using the 'expectedCFs' function in MSCQuartets on the

UCE MSC species tree, which was congruent with the most common Clade A topology among

datasets (see Results and Supplementary Materials).

To test for putative introgression among taxa in Clade A, we first estimated f_{DM} for 100

kb windows across the whole genome. F_{DM} is an f-branch statistic derived from the ABBA-BABA

test (D-statistic) that can quantify the magnitude of introgression assuming a species tree

topology. To do so, we first obtained genome-wide single nucleotide polymorphism (SNP) data

for one member of each species within Clade A (*C. c. goldmani*, *C. erythropus*, *C. atrocapillus*,

C. duidae, *C. kerriae*, *C. noctivagus*, *C. boucardi*, *C. strigulosus*, *C. transfasciatus, and C. u.*

undulatus) plus one outgroup, *C. variegatus*. We again mapped cleaned reads for these

samples to a reference genome, following the same protocol as outlined above, but this time

using the pseudo-chromosome assembly from the Clade A ingroup with a relatively complete

genome assembly, *C. u. undulatus* (LSUMNS 34614), as the same reference genome for all

samples. We then used bcftools to call SNPs for all variants across the genome of each

species. Finally, we filtered this merged vcf file using vcftools v1.16 (Danecek et al. 2011) to

obtain SNPs with a minimum allele frequency = 0.05, maximum missing data = 0.5, minimum

quality = 30, minimum depth = 10, and maximum depth = 100. Using these SNP data, we then

estimated F_{DM} for 100 kb non-overlapping windows across the genome using the

'ABBABABAwindows.py' script located at

https://github.com/simonhmartin/genomics_general#trees-for-sliding-windows (Martin et al.

2014). We specifically tested for introgression at branches whose phylogenetic positions

differed among analyses and datasets. Because our datasets and phylogenetic analyses did not

agree on a single most probable topology for Clade A, we ran this script four times, each time

assuming one of the three hypothesized topologies (T1, T2, or T3; T4 had low support and was

considered unresolved; see Results, Figure 2) as our guide tree.

323
Figure 1: Time-calibrated Phylogeny of tinamous based on Autosomal UCEs with 1,000 bp of flanking
sequence. Blue node bars represent 95% Highest Posterior Densities for node ages. White circles denote
nodes constrained for monophyly based on maximum likelihood and MSC results. Tinamou illustrations
by TAC.

Finally, we estimated the species tree topology while accounting for gene flow using a

Bayesian approach implemented in PhyloNet v3.8.0 (the 'MCMC_GT' algorithm) (Wen et al.

2016). This method employs a reverse-jump Markov Chain Monte Carlo (rjMCMC) to sample

the posterior distribution of phylogenetic networks under a multi-species network coalescent

(MSNC) model using rooted gene trees as input. The MSNC is similar to the MSC, but relaxes

the assumption of no introgression by modeling genome evolution as a network rather than

bifurcating phylogeny. We ran the analysis twice, again using the gene trees with collapsed low-

support nodes from the Z-linked and Autosomal Clade A UCE's varying the maximum number

of reticulate nodes in each run to assess variability in the results. We ran PhyloNet 11 times for

both the autosomal and Z-linked datasets restricted to Clade A, once each for m-values

between zero and ten (m defines the maximum number of reticulate nodes allowed in the

MSNC model). We ran the rjMCMC for 5×10^7 generations with a burn-in of 10%, using the

pseudo-likelihood calculation. Because likelihood scores tended to increase with each

successive increase in m-value, we chose the optimal m-value using a breakpoint analysis in

the R package 'segmented' (Muggeo et al. 2014). The optimal m-value was chosen by fitting a

segmented linear model to our likelihoods for each m-value. This allowed us to identify

breakpoints in the slope of the regression, where increases in m-value resulted in diminishing

gains in likelihood.

Results

Phylogenomics of Tinamous

The concatenated and MSC phylogenies for tinamous were well-supported and broadly

congruent across datasets (Figures 1,2, S1–S21). The 75% and 100% complete datasets were

congruent except for in the concatenated UCE results, implying that missing data had limited

impact on phylogenetic reconstruction. However, there was slight variation in bootstrap and

gCF's support values, with 75% complete datasets tending to have higher support (Figure 2).

The phylogenies from all datasets were congruent except for relationships within a single clade

containing ten species of *Crypturellus,* we refer to as Clade A (Figure 2). The placement of *C.*

atrocapillus was most unstable but was often recovered as sister to a clade containing *C.*

boucardi, C. kerriae, C. erythropus, C. duidae, and *C. strigulosus* (Figure 2 "T1"). Overall, we

recovered four different topologies for this clade based on the primary datasets (Figure 2: T1–

T4). However, the Z-linked UCE dataset had low support for the placement of *C. atrocapillus* in

the ASTRAL-III analysis (posterior probability = 0.46), and therefore failed to resolve the

relationships of Clade A.

Our results were broadly consistent with recognized taxonomic classifications. We

recovered monophyletic subfamilies Nothurinae and Tinaminae and most genera and species

were also monophyletic. However, two genera were non-monophyletic: the monotypic genus

Taoniscus was embedded within *Nothura*, and *Rhynchotus* was embedded within *Nothoprocta*,

rendering *Nothura* and *Nothoprocta* paraphyletic, respectively. We also recovered *Nothoprocta*

pentlandii as polyphyletic; one specimen from Bolivia (voucher LSUMZ 123403) was sister to *N.*

perdicaria, whereas the remaining samples from Peru formed a clade that was sister to all

remaining *Nothoprocta*. All other species that were represented by multiple samples were

recovered as monophyletic.

The tempo of tinamou diversification

Our tip-dated phylogeny revealed relatively constant diversification of tinamous, which

diverged from Moas roughly 55mya (Figure 1). The crown divergence of tinamous, separating

forest (Tinaminae) from non-forest (Nothurinae) lineages, occurred roughly 37mya. A list of

divergence times for genera and major clades can be found in Table S2. Speciation and

extinction rates were stable through time, without any significant rate changes (Figure S25). We

recovered a high rate of molecular evolution for the Tinamidae stem lineage (0.0034

substitutions/million years) compared to the mean for all lineages (ORCucldMean = 0.0019;

95% HPD = 0.0016–0.0023).

Figure 2: Phylogenetic topologies for a difficult-to-resolve clade (Clade A) of *Crypturellus*
tinamous. The phylogenetic placement of *C. atrocapillus* (pink branch) was discordant among datasets
and species tree approaches. The top four topologies are shown at the top (T1–T4). The heatmap at the
bottom shows node support and/or concordance for each dataset and phylogenetic approach
(concatenation and MSC). From top to bottom, these are (1) gene concordance factors (gCF) for a
"universal" node shared among all three topologies, (2) gCF for a "unique" node in each topology
associated with the position of *C. atrocapillus*, (3) bootstrap (IQ-TREE) or posterior probability (ASTRAL-
III) for the "universal" node, and (4) bootstrap or PP for the "unique" node. The "universal" (node 1) and
"unique" (node 2) nodes on each of the four topologies are marked on trees at the top of the figure. The
numbers on the heatmap denote the rounded values, and the colors correspond to normalized values,
such that the darkest red in each row is the highest value and the darkest blue is the lowest for that row.
PP and bootstrap values are both shown as percentages for easy comparison. Results for additional
datasets can be found in the Supplemental Materials.
396

Gene tree discordance arises from stochastic error, ILS, and introgression

We examined gene tree discordance for each dataset using RF distances between gene

and species trees. To quantify the relationship between alignment information content, as

measured using PIS, and gene tree discordance, we modeled RF distances as a function of the

number of PIS per locus using a generalized linear model (Figure 3A & S23). This model was

overall consistent with an expected negative association between PIS and RF distance, with

lower RF distances in gene trees built from more informative alignments. This negative

association (R^2=0.29, *P*<0.0001) indicated that nearly one third of the variation in RF distance in

UCE gene trees was explained by the number of PIS. This relationship was also logarithmic,

suggesting that the effect of PIS on gene tree discordance was greatest for the least informative

alignments but this effect quickly declined. When comparing autosomal and Z-linked UCE's in

this context, Z-linked UCE's primarily fell below the model trendline, suggesting that for the

same values of PIS, Z-linked UCE's more closely resembled the species tree. Indeed, overall,

Z-linked UCE gene trees averaged fewer differences from the species tree (mean RF distance

between gene and species tree = 32.23.68 ± 9.67) than autosomal gene trees (mean=38.58 ±

9.67), and this difference was significant (X^2=26.65, df=1, P<0.0001). Taken together, these

findings suggest lower rates of ILS and/or introgression on the Z-chromosome, even when

accounting for potential stochastic error.

We found evidence of ILS and deviations from the MSC model using the quartet

approach (Figure 3B). For one branch (branch 3), we recovered frequencies for three alternative

quartets that approached the ⅓ threshold when using autosomal UCE's, implying ILS has

promoted phylogenetic discordance at these branches. However, two of the alternative quartets

exceeded the ⅓ threshold and were roughly equal in frequency, indicating a deviation from the

MSC. When assessing quartet frequencies for Z-linked gene trees at branch 3, the results

closely matched expectations of ILS, wherein the relative frequencies of the three alternative

quartets were approximately equal to the ⅓ threshold, and we did not reject a null model where

Z-linked quartets had equivalent frequencies based on the multinomial test (P=0.88). For

another branch (branch 5), we found weaker evidence for ILS, with a single majority topology

(34|67) and two minority topologies. For both autosomal and Z-linked gene trees, we also found

deviations from pure ILS, as the two minority topologies were not equal in frequency. However,

we again found that Z-linked gene trees matched expectations of reduced ILS on the sex

chromosomes, with a majority topology becoming more frequent in the Z-linked quartet analysis

(i.e., less gene tree discordance). In all cases, quartet frequencies differed from MSC

expectations given the UCE ASTRAL-III species tree (P<0.0001).

433
Figure 3: Gene tree discordance for autosomal and sex-linked UCEs. (A) Linear regression model
exploring the relationship between information content (# parsimony informative sites) and gene tree
discordance (RF distance between gene and species tree). The negative correlation suggests that
alignments with less information content result in larger RF distances, but that Z-linked UCE's (yellow
points), fall below the trendline and thus have reduced RF distances given the same *x*-axis values. (B)
Relative quartet frequencies for alternative unrooted topologies in Clade A. T1–T4 correspond to quartets
consistent with topologies shown in Figure 2. Numbers on the branches of the unrooted network at the
left coincide with numbers at the x-axis of each bar plot. For example, 12|45 denotes an unrooted quartet
consistent with Topology 1 (T1; Figure 2) wherein *C. noctivagus* (branch 1) is sister to *C. transfasciatus* +
C. cinnamomeus (branch 2) and *C. atrocapillus* (branch 4) is sister to branch 5. "Tx" refers to a quartet
that does not match one of the four primary topologies in Figure 2.

Figure 4: Analysis of genome-wide introgression. (A) Introgression (f_{dM}) mapping for 100 kb non-overlapping windows across the genome assuming each of the three well supported topologies (Figure 2, T1–T3). The trees at the left of each plot show the assumed background topology and red arrows indicate positive or negative values of f_{dM}. Tips on the trees show taxa assigned to each population in the introgression model (from left to right: P1, P2, P3, and P4 assigned populations in the model). (B) The results from the multi-species network coalescent model in PhyloNet. The Z-linked tree was weakly supported.

454    Our genome-wide analysis of f_{DM} for 100 kb sliding windows provided contrasting signals

455 for introgression across the genome, but identified non-zero signals of introgression across

456 most windows of the genome (Figure 4A). When assuming T1 as the phylogenetic background

457 in the model, we found slight negative values of f_{dM} across autosomes indicating introgression

458 between the *C. cinnamomeus/transfasciatus* branch and the *C.*

459 *strigulosus/duidae/erythropus/kerriae/boucardi* branch (P1 and P3 in this model). These values

460 of f_{dM} only slightly deviated from zero (mean autosomal f_{dM} = -0.021 ± 0.00022), whereas the

461 analysis recovered very strong signals of introgression for many windows along much of the Z-

462 chromosome (mean Z-linked f_{dM} = -0.117 ± 0.0052), and this difference was significant (X^2 =

463 311.21, *df* = 1, *P* < 0.0001). When assuming T2 as the phylogenetic background, we saw the

464 opposite pattern; autosomes again showed consistent negative values of f_{dM} indicating

465 introgression between *C. atrocapillus* and the *C. strigulosus/duidae/erythropus/kerriae/boucardi*

466 branch (mean autosomal f_{dM} = -0.027 ± 0.00029), but the windows on the Z-chromosome

467 averaged much closer to zero (mean Z-linked f_{dM} = -0.010 ± 0.00098). The difference between

468 mean autosomal and Z-linked introgression was significant (X^2 = 293.49, *df* = 1, *P* < 0.0001).

469 When assuming T3 as the phylogenetic background, we again found evidence of genome-wide

470 introgression, this time mostly positive values indicating introgression between *C. atrocapillus*

471 and the *C. strigulosus/duidae* branch. There were several obvious negative peaks across the

472 genome in this scenario, including much of the Z-chromosome (mean autosomal f_{dM} = 0.027 ±

473 0.00037; mean Z-linked f_{dM} = -0.022 ± 0.0019), indicating introgression between *C. atrocapillus*

474 and the *C. erythropus/kerriae/boucardi* branch.

475    As a final test of the Clade A topology and impacts of gene flow, our phylogenomic

476 network analysis revealed multiple nodes involved in reticulation that may explain gene tree

477 discordance in Clade A (Figure 4B). A breakpoint analysis indicated that the optimal m-value

478 (number of reticulate nodes) was m=3 for autosomal UCE's and was m=1 for Z-linked UCE's

479 (Figure S26); thus, we recovered fewer introgression events for the Z-linked than autosomal

UCE's. For autosomal UCE's, we found a network with a reticulate node involving *C.*

atrocapillus (posterior probability = 1.0), suggesting that uncertainty around the placement of

this species could conceivably be driven by historical introgression. For the Z-linked loci, we

recovered a network most closely resembling T1 with a signal of introgression between *C.*

undulatus and *C. atrocapillus,* but this specific network topology was not supported (posterior

probability = 0.46).

Discussion

Here we used whole-genome sequencing to explore the sources of phylogenomic

conflict and reconstruct the species-level phylogenetic history of tinamous, a relatively old

Neotropical avian group of broad interest in evolutionary biology (Sackton et al. 2019; Li et al.

2023). Our results based on different analytical approaches (concatenated and MSC methods)

and datasets (UCE's and BUSCO's) were largely congruent. The topology of only a single clade

in the genus *Crypturellus* (Clade A), which contained multiple successive short internodal

branches, varied among approaches. These results indicate that species tree estimation in this

group was, in general, robust, but that gene tree estimation error, ILS, and introgression may

have been key drivers of gene tree discordance that hampered phylogenetic inference for one

recalcitrant node in the genus *Crypturellus* (in Clade A).

Indeed, both ILS and introgression likely drove uncertainty in the position of *C.*

atrocapillus, the most unstable node in the phylogeny, which had relatively low concordance

factors and reduced node support in many datasets (Figure 2). The relative quartet frequencies

for this node closely approached the ⅓ threshold predicted by high ILS, especially from Z-

chromosome alignments (Figure 3B). However, given that we could not eliminate the impacts of

stochastic error, it remains uncertain how much of the gene tree discordance described in the

quartet analysis was associated with ILS versus gene tree estimation error. In addition to ILS

and introgression, rapid divergence events can be difficult to reconstruct because the short

timeframe also means there is limited phylogenetic signal for those divergence events within

individual gene trees (Leaché et al. 2015; Mclean et al. 2019). Still, multiple methods pointed

towards genome-wide introgression in this clade, potentially depleting signal for the species tree

across much of the genome and further complicating phylogenetic reconstruction in Clade A.

Additional analysis and intraspecific sampling may be necessary to determine the precise

phylogenetic relationships of taxa in Clade A (Arcila et al. 2017; Shen et al. 2017; Remsen et al.

2024).

Phylogeny and taxonomy of the Tinamous

Our phylogenies based on multiple datasets all resulted in highly similar and well-

supported relationships among tinamou species. For example, past studies, based primarily on

morphological data or few molecular loci, recovered high uncertainty in the interrelationships of

Crypturellus species (Bertelli et al. 2014; Almeida et al. 2022), but our reconstructions for this

genus were mostly robust. Moreover, concatenation and MSC approaches largely agreed on

the overarching relationships at both shallow and deeper timescales. Importantly, our

phylogenetic results imply that multiple taxonomic changes are necessary to appropriately

classify taxa within Tinamidae. First, *Nothoprocta cinerascens* was recovered as the sister to

the genus *Rhychotus* (Spix 1825). *Rhynchotus* (Spix 1825) has nomenclatural priority over

Nothoprocta (Sclater and Salvin 1873)*,* and we therefore suggest moving *N. cinerascens* to the

genus *Rhynchotus* (Bertelli and Porzecanski 2004; Valqui 2009). Second, *Taoniscus nanus* was

nested within the genus *Nothura*, and thus, *Taoniscus* (Gloger 1842) is a junior synonym of

Nothura (Wagler 1827)*.* We therefore suggest transferring *T. nanus* to *Nothura.* Third, *C.*

duidae, often considered a close relative of *C. noctivagus* or *C. erythropus* (Remsen et al.

2024), was found not to have close affinities to either of these; instead, it was consistently

placed as sister to *C. strigulosus.* Finally, we found evidence for what is likely an unrecognized

species of *Nothoprocta* (Valqui 2009); specimens ascribed to *N. pentlandii* were polyphyletic,

with a specimen of the nominate subspecies from Bolivia recovered as sister to *N. perdicaria*,

and multiple specimens from Peru recovered as a clade sister to all other *Nothoprocta.* Thus,

there appear to be at least two species-level taxa within *N. pentlandii*, and more work is needed

to properly circumscribe the taxa in this species complex.

Tip-dating enables approximation of the relationships of fossils even when phylogenetic data

are equivocal

Our estimates of divergence dates for tinamous were broadly consistent with a recent

study (Almeida et al. 2022). Although we used many of the same fossils for calibration, Almeida

et al. (2022) used node calibrations. Here we used tip-dating, which explicitly accounts for

uncertainty in fossil placement, enabling us to include additional crown fossils never before

included in a dating analysis. Although the placement of some fossils in our dataset have been

recovered with strong support in previous studies, others have not (Bertelli et al. 2014; Almeida

et al. 2022). For example, prior work has found that an extinct *Eudromia*-like species is sister to

the clade containing extant *Eudromia* and *Tinamotis,* and two undescribed fossil fragments

(MACN-SC-3610 and MACN-SC-3613) belong to stem *Crypturellus*. On the other hand, the

placement of the extinct *C. reai*, was equivocal in morphological phylogenies despite sharing

clear synapomorphies with extant *Crypturellus*. Our analysis allowed the placement of *C. reai* to

vary between crown and stem *Crypturellus*, and showed that this species either belongs to an

early crown lineage or late stem lineage, but is not likely closely related to any extant species

(Figure 1). For example, the posterior probability of crown *Crypturellus* in our tip-dated tree was

0.4, indicating a possibility that *C. reai* could be nested within the crown. However, all other

nodes in this clade were strongly supported as monophyletic (PP≥0.98; supplemental tree files).

In line with other macroevolutionary studies for Neotropical birds, we found relatively

constant rates of diversification for tinamous (Patel et al. 2011; d'Horta et al. 2013; Harvey et al.

2020). Our results agree with those of Almeida et al. (2022) that tinamous diverged from their

sister group, the extinct moas, during the Paleocene or early Eocene with crown tinamous

diverging into forest (Tinaminae) and open-country (Nothurinae) clades during late Eocene or

early Oligocene (Figure 1; Table S6). All extant genera of tinamous arose in the mid-to-late

Miocene, except *Eudromia* (Late Miocene to Pliocene) and *Crypturellus* (early-to-mid Miocene).

Using our unique approach, we also recapitulated the high substitution rate of the stem tinamid

lineage, one of the highest in modern birds (Berv and Field 2018), a finding that remains

unexplained.

The drivers of gene tree discordance in Clade A include pervasive signals of introgression

We examined orthologous data harvested from whole genome assemblies for the

potential impacts of stochastic error to better inform our tests of ILS and introgression. We found

that, as expected, as parsimony informative sites (PIS) are added to alignments, gene-tree-to-

species-tree discordance is reduced. This relationship is well known (Meiklejohn et al. 2016;

Burbrink et al. 2020), but confirming it for our dataset allowed us to understand baseline levels

of potential stochastic error in the data (i.e., 29% of the variance in RF distances is explained by

alignment information content), and thus evaluate levels of ILS in light of this relationship. Z-

linked gene trees fell below the trendline for the PIS x RF distance regression model, implying

that for the same number of PIS as autosomal alignments, Z-linked alignments had lower RF

distances. We argue that this is likely because Z-linked UCE's should have reduced rates of

ILS, though disentangling stochastic and other forms of error from legitimate biological signals

of ILS in these gene tree summarization methods is admittedly difficult.

In Clade A, the placement of *C. atrocapillus* varied among datasets and analytical

approaches, but one topology was most common (T1). This topology was recovered for most

UCE trees as well as the concatenated Z-linked UCE tree (Figure 2). Sex-linked markers are

expected to experience reduced rates of ILS and faster coalescent times than autosomes due to

smaller effective population size, as well as reduced rates of recombination and introgression

(Charlesworth et al. 1987; Meisel and Connallon 2013; Li et al. 2019; Rivas-González et al.

2023; Musher et al. 2024; Burbrink et al. 2025). These expectations are consistent with our

findings that (1) as laid out above, RF distances between gene and species trees were lower for

Z-linked UCE's than for autosomal UCE's given comparable values of PIS, suggesting reduced

ILS on the Z-chromosome, (2) the quartet frequencies of alternative topologies for relatively

short internal branch lengths more closely matched MSC model expectations for Z-linked than

autosomal UCE's, suggesting reduced introgression on the Z-chromosome, and (3) the optimal

number of reticulation events was lower for Z-linked phylogenetic network than for the

autosomal network, though this last observation could be driven, in part, by the large difference

in the number of alignments obtained for each dataset (3,942 autosomal UCE's vs. 200 Z-linked

UCE's). Nevertheless, the gene trees from the Z-linked dataset failed to resolve the species tree

and network for Clade A on their own. It thus seems likely that given the small number of Z-

linked genes analyzed, and rapid divergence at the *C. atrocapillus* node, a larger Z-linked

dataset may be needed to resolve the relationships in Clade A. Future work might generate Z-

chromosome alignments to obtain more data that could help resolve Clade A.

Interestingly, our analysis of introgression using f_{dM}, which does not rely on gene tree

estimation, may provide some insight into the species tree of Clade A. When assuming T1, the

most commonly recovered topology in our study, as the species tree, we found low levels of

introgression for most sliding windows across the autosomes but significant peaks of

introgression (negative f_{DM} values) along those on the Z-chromosome, a finding that is opposed

to theoretical expectations that the Z-chromosome should have reduced introgression (Burbrink

et al. 2025). This result is striking and could be interpreted to mean that T1 is the wrong

topology despite being the obvious candidate for the species tree. Contrastingly, when

assuming T2 as the species tree, we recovered levels of introgression (negative f_{DM} values) that

were consistent with such expectations: pervasive signals of introgression across all the

autosomes, with values closer to zero on the Z-chromosome. These results raise the possibility

that T2, despite only being recovered in one analysis, might be the correct species tree, but the

signal for T2 has been depleted in the genome by the overwriting impacts of pervasive

autosomal introgression (Fontaine et al. 2015). T2 was also the most frequent autosomal

quartet for branch 3 (Figure 3B). Pervasive introgression between *C. atrocapillus* and the

ancestor of *C. erythropus, C. boucardi*, and *C. kerriae* could explain both the difficulty in

determining the phylogenetic placement of *C. atrocapillus*, while also explaining why the T1 was

the most commonly recovered topology (i.e., it may be a signal of introgression). Indeed, our

phylogenomic network from autosomes, which estimated the phylogenetic history while

accounting for introgression, recovered a network involving reticulation between *C. atrocapillus*,

the *C. strigulosus + duidae* clade, and the *C. erythropus + boucardi + kerriae* clade. However,

we would argue that more work is needed to corroborate this hypothesis. Instead, we

conservatively suggest that the phylogeny of Clade A remains uncertain, with evidence that

rapid divergence and introgression have promoted this phylogenetic uncertainty.

Acknowledgements

We thank Alvaro Hernandez and Chris Wright at the University of Illinois Roy J. Carver

Biotechnology Center for assistance with Illumina sequencing. We thank Kimberly K. O. Walden

for assistance in depositing Illumina reads to NCBI SRA. We also thank Jorge Doña for

providing R code to assist with breakpoint analysis. Dan Lane, Steve Cardiff, and Christopher

Witt helped confirm voucher identities for multiple samples. We thank B. T. Smith, J. Cracraft, P.

Sweet, T. Trombone, S. Katanova (AMNH), J. Bates, S. Hackett, and B. Marks (FMNH), F.

Sheldon and D. Dittman (LSUMNS), H.F. James and B. Schmidt (USNM), and N. Rice (ANSP)

for loaning material that was crucial for this work. Finally, we thank the members of ANSP

Ornithology Department (A. Del Grosso, E. Griffith, K. Kuabara, and J. Merwin) for comments

on early versions of the manuscript. Comments from S. Ruane, Isaac Overcast, and five

anonymous referees greatly improved this work. This study was funded by NSF grant DEB-

1855812 to JDW and KPJ. TAC was partially supported by NSF grant DEB-2203228.

Data Availability

639   All raw reads are available on the NCBI Sequence Read Archive (project and sample accession

640   numbers can be found in Table S1). All the original data and scripts necessary to reproduce the

641   analyses reported in this study can be accessed through LJM's github page:

642   https://github.com/lukemusher/Tinamou-phylogenomics.git.

Literature Cited

Albert J.S., Tagliacollo V.A., Dagosta F. 2020. Diversification of Neotropical Freshwater Fishes.
Annu. Rev. Ecol. Evol. Syst. 51:27–53.

Allman E.S., Baños H., Mitchell J.D., Rhodes J.A. 2023. MSCquartets: analyzing gene tree
quartets under the multi-species coalescent. R package version 1.3.1.

Almeida F.C., Porzecanski A.L., Cracraft J.L., Bertelli S. 2022. The evolution of tinamous
(Palaeognathae: Tinamidae) in light of molecular and combined analyses. Zool. J. Linn.
Soc. 195:106–124.

Alonge M., Lebeigle L., Kirsche M., Jenike K., Ou S., Aganezov S., Wang X., Lippman Z.B.,
Schatz M.C., Soyk S. 2022. Automated assembly scaffolding using RagTag elevates a new
tomato system for high-throughput genome editing. Genome Biol. 23:258.

Alvarenga H.M.F. 1983. Uma ave ratitae do paleoceno brasileiro: bacia calcaŕia de itaboraí,
estado do Rio de Janeiro, Brasil. Boletim do Museu Nacional, Nova Série, Geologia. 41:1–
8.

Antonelli A., Zizka A., Carvalho F.A., Scharn R., Bacon C.D., Silvestro D., Condamine F.L.
2018. Amazonia is the primary source of Neotropical biodiversity. Proc. Natl. Acad. Sci.
115:6034–6039.

Arcila D., Ortí G., Vari R., Armbruster J.W., Stiassny M.L.J., Ko K.D., Sabaj M.H., Lundberg J.,
Revell L.J., Betancur-R R. 2017. Genome-wide interrogation advances resolution of
recalcitrant groups in the tree of life. Nat. Ecol. Evol. 1:0020.

Baker A.J., Haddrath O., McPherson J.D., Cloutier A. 2014. Genomic support for a moa-
tinamou clade and adaptive morphological convergence in flightless ratites. Mol. Biol. Evol.
31:1686–1696.

Bertelli S. 2017. Advances on tinamou phylogeny: an assembled cladistic study of the volant

palaeognathous birds. Cladistics. 33:351–374.

Bertelli S.B., Chiappe L.M. 2005. Earliest tinamous (Aves: Palaeognathae) from the Miocene of
Argentina and their phylogenetic position. Contributions in science. 502:1–20.

Bertelli S., Chiappe L.M., Mayr G. 2014. Phylogenetic interrelationships of living and extinct
Tinamidae, volant palaeognathous birds from the New World. Zool. J. Linn. Soc. 172:145–
184.

Bertelli S., Giannini N.P., Goloboff P.A. 2002. A phylogeny of the tinamous (aves:
palaeognathiformes) based on integumentary characters. Syst. Biol. 51:959–979.

Bertelli S., Porzecanski A.L. 2004. Tinamou (tinamidae) systematics: A preliminary combined
analysis of morphology and molecules. Ornitologia neotropical 15:293–299.

Berv J.S., Field D.J. 2018. Genomic Signature of an Avian Lilliput Effect across the K-Pg
Extinction. Syst. Biol. 67:1–13.

Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A.,
Heled J., Jones G., Kühnert D., De Maio N., Matschiner M., Mendes F.K., Müller N.F.,
Ogilvie H.A., du Plessis L., Popinga A., Rambaut A., Rasmussen D., Siveroni I., Suchard
M.A., Wu C.-H., Xie D., Zhang C., Stadler T., Drummond A.J. 2019. BEAST 2.5: An
advanced software platform for Bayesian evolutionary analysis. PLoS Comput. Biol.
15:e1006650.

Burbrink F.T., DeBaun D., Foley N.M., Murphy W.J. 2025. Recombination-aware
phylogenomics. Trends Ecol. Evol. 0:0 (*in press*)

Burbrink F.T., Gehara M. 2018. The Biogeography of Deep Time Phylogenetic Reticulation.
Syst. Biol. 67:743–744.

Capella-Gutierrez S., Silla-Martinez J.M., Gabaldon T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 25:1972–1973.

Capurucho J.M., Musher L.J., Lees A., Rego M.A., Del-Rio G., Aleixo A., Luzuriaga-Aveiga
V.E., Ferreira M., Ribas C., Thom G. 2023. Amazonian avian biogeography: Broadscale
patterns, microevolutionary processes, and habitat-specific models revealed by
multidisciplinary approaches. Ornithology. 141:1–17.

Charlesworth B., Coyne J.A., Barton N.H. 1987. The Relative Rates of Evolution of Sex
Chromosomes and Autosomes. Am. Nat. 130:113–146.

Chen S., Zhou Y., Chen Y., Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 34:i884–i890.

Chernomor O., von Haeseler A., Minh B.Q. 2016. Terrace Aware Data Structure for
Phylogenomic Inference from Supermatrices. Syst. Biol. 65:997–1008.

Claramunt S., Cracraft J. 2015. A new time tree reveals Earth history's imprint on the evolution
of modern birds. Sci Adv. 1:e1501005.

Clements J.F., Rasmussen P.C., Schulenberg T.S., Iliff M.J., Fredericks T.A., Gerbracht J.A.,
Lepage D., Spencer A., Billerman S.M., Sullivan B.L., Wood C.L. 2023. The eBird/Clements

checklist of Birds of the World: v2023.

Cloutier A., Sackton T.B., Grayson P., Clamp M., Baker A.J., Edwards S.V. 2019. Whole-
genome analyses resolve the phylogeny of flightless birds (Palaeognathae) in the presence
of an empirical anomaly zone. Syst. Biol. 68:937–955.

Cracraft J. 1974. Phylogeny and evolution of the ratite birds. Ibis. 116:494–521.

Danecek P., Auton A., Abecasis G., Albers C.A., Banks E., DePristo M.A., Handsaker R.E.,
Lunter G., Marth G.T., Sherry S.T., McVean G., Durbin R., 1000 Genomes Project Analysis
Group. 2011. The variant call format and VCFtools. Bioinformatics. 27:2156–2158.

Del-Rio G., Mutchler M.J., Costa B., Hiller A.E., Lima G. 2021. Birds of the Juruá River:
extensive várzea forest as a barrier to terra firme birds. Journal of Ornithology 162:565–
577.

Didier G., Royer-Carenzi M., Laurin M. 2012. The reconstructed evolutionary process with the
fossil record. J. Theor. Biol. 315:26–37.

Douglas J., Zhang R., Bouckaert R. 2021. Adaptive dating and fast proposals: Revisiting the
phylogenetic relaxed clock model. PLoS Comput. Biol. 17:e1008322.

Drummond A.J., Ho S.Y.W., Phillips M.J., Rambaut A. 2006. Relaxed phylogenetics and dating
with confidence. PLoS Biol. 4:e88.

Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic
loci. Bioinformatics. 32:786–788.

Faircloth B.C., McCormack J.E., Crawford N.G., Harvey M.G., Brumfield R.T., Glenn T.C. 2012.
Ultraconserved elements anchor thousands of genetic markers spanning multiple
evolutionary timescales. Syst. Biol. 61:717–726.

Fontaine M.C., Pease J.B., Steele A., Waterhouse R.M., Neafsey D.E., Sharakhov I.V., Jiang
X., Hall A.B., Catteruccia F., Kakani E., Mitchell S.N., Wu Y.-C., Smith H.A., Love R.R.,
Lawniczak M.K., Slotman M.A., Emrich S.J., Hahn M.W., Besansky N.J. 2015. Mosquito
genomics. Extensive introgression in a malaria vector species complex revealed by
phylogenomics. Science. 347:1258524.

Franz N.M., Musher L.J., Brown J.W., Yu S., Ludäscher B. 2019. Verbalizing phylogenomic
conflict: Representation of node congruence across competing reconstructions of the
neoavian explosion. PLoS Comput. Biol. 15:e1006493.

Freeman B.G., Pennell M.W. 2021. The latitudinal taxonomy gradient. Trends in Ecology &
Evolution. 36:778–786.

Gavryushkina A., Welch D., Stadler T., Drummond A. 2014. Bayesian inference of sampled
ancestor trees for epidemiology and fossil calibration. PLoS Comput. Biol. 10:e1003919.

Harshman J., Braun E.L., Braun M.J., Huddleston C.J., Bowie R.C.K., Chojnowski J.L., Hackett
S.J., Han K.-L., Kimball R.T., Marks B.D., Miglia K.J., Moore W.S., Reddy S., Sheldon F.H.,
Steadman D.W., Steppan S.J., Witt C.C., Yuri T. 2008. Phylogenomic evidence for multiple
losses of flight in ratite birds. Proc. Natl. Acad. Sci. U. S. A. 105:13462–13467.

Harvey M.G., Bravo G.A., Claramunt S., Cuervo A.M., Derryberry G.E., Battilana J., Seeholzer
G.F., McKay J.S., O'Meara B.C., Faircloth B.C., Edwards S.V., Pérez-Emán J., Moyle R.G.,
Sheldon F.H., Aleixo A., Smith B.T., Chesser R.T., Silveira L.F., Cracraft J., Brumfield R.T.,
Derryberry E.P. 2020. The evolution of a tropical biodiversity hotspot. Science. 370:1343–
1348.

Hoang D.T., Chernomor O., von Haeseler A., Minh B.Q., Vinh L.S. 2018. UFBoot2: Improving
the Ultrafast Bootstrap Approximation. Mol. Biol. Evol. 35:518–522.

Höhna S., May M.R., Moore B.R. 2016. TESS: an R package for efficiently simulating
phylogenetic trees and performing Bayesian inference of lineage diversification rates.
Bioinformatics. 32:789–791.

d'Horta F.M., Cuervo A.M., Ribas C.C., Brumfield R.T., Miyaki C.Y. 2013. Phylogeny and
comparative phylogeography of Sclerurus (Aves: Furnariidae) reveal constant and cryptic
diversification in an old radiation of rain forest understory specialists. J. Biogeogr. 40:37–
49.

Hughes C., Eastwood R. 2006. Island radiation on a continental scale: exceptional rates of plant
diversification after uplift of the Andes. Proc. Natl. Acad. Sci. U. S. A. 103:10334–10339.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.
14:587–589.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Korneliussen T.S., Albrechtsen A., Nielsen R. 2014. ANGSD: Analysis of Next Generation
Sequencing Data. BMC Bioinformatics. 15:356.

Krabbe N.K., Schulenberg T.S., Hosner P.A., Rosenberg K.V., Davis T.J., Rosenberg G.H.,
Lane D.F., Andersen M.J., Robbins M.B., Cadena C.D., Valqui T., Salter J.F., Spencer A.J.,
Angulo F., Fjeldså J. 2020. Untangling cryptic diversity in the High Andes: Revision of the
Scytalopus [magellanicus] complex (Rhinocryptidae) in Peru reveals three new species.
Auk. 137:ukaa003.

Leaché A.D., Chavez A.S., Jones L.N., Grummer J.A., Gottscho A.D., Linkem C.W. 2015.
Phylogenomics of phrynosomatid lizards: conflicting signals from sequence capture versus
restriction site associated DNA sequencing. Genome Biol. Evol. 7:706–719.

Li G., Figueiró H.V., Eizirik E., Murphy W.J. 2019. Recombination-Aware Phylogenomics
Reveals the Structured Genomic Landscape of Hybridizing Cat Species. Mol. Biol. Evol.
36:2111–2126.

Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 25:1754–1760.

Li H., Handsaker B., Wysoker A., Fennell T., Ruan J., Homer N., Marth G., Abecasis G., Durbin
R., 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map
format and SAMtools. Bioinformatics. 25:2078–2079.

Li Q., Chen D., Wang S. 2023. Character displacement of egg colors during tinamou speciation.
Evolution. 77:1874–1881.

Martin S.H., Davey J.W., Jiggins C.D. 2014. Evaluating the Use of ABBA–BABA Statistics to
Locate Introgressed Loci. Mol. Biol. Evol. 32:244–257.

McCormack J.E., Faircloth B.C., Crawford N.G., Gowaty P.A., Brumfield R.T., Glenn T.C. 2012.
Ultraconserved elements are novel phylogenomic markers that resolve placental mammal
phylogeny when combined with species-tree analysis. Genome Res. 22:746–754.

McCormack J.E., Tsai W.L.E., Faircloth B.C. 2016. Sequence capture of ultraconserved
elements from bird museum specimens. Mol. Ecol. Resour. 16:1189–1203.

Mclean B.S., Bell K.C., Allen J.M., Helgen K.M., Cook J.A. 2019. Impacts of Inference Method
and Data set Filtering on Phylogenomic Resolution in a Rapid Radiation of Ground
Squirrels (Xerinae: Marmotini). Syst. Biol. 68:298–316.

Meisel R.P., Connallon T. 2013. The faster-X effect: integrating theory and data. Trends Genet.
29:537–544.

Menzel U. 2024. EMT: exact multinomial test: goodness-of-fit test for discrete multivariate data.
R package version 1.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,
Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mirarab S., Reaz R., Bayzid M.S., Zimmermann T., Swenson M.S., Warnow T. 2014. ASTRAL:
genome-scale coalescent-based species tree estimation. Bioinformatics. 30:i541–8.

Mitchell K.J., Llamas B., Soubrier J., Rawlence N.J., Worthy T.H., Wood J., Lee M.S.Y., Cooper
A. 2014. Ancient DNA reveals elephant birds and kiwi are sister taxa and clarifies ratite bird
evolution. Science. 344:898–900.

Moncrieff A.E., Johnson O., Felix C., Hiller A.E., Corbett E.C., Brady M.L., Seeholzer G.F.,
Bautista E., Lane D.F., Harvey M.G. 2021. Avifaunal surveys in the central Peruvian
Amazon clarify range limits and highlight links between avian and habitat diversity. Wilson
J. Ornithol. 132:934–951.

Moncrieff A.E., Johnson O., Lane D.F., Alonso J.Á., Balta K., Eckhardt K., Armenta J., Valqui T.,
Hernández F., Huaira M.S., Mur C., Harvey M.G., Verde-Guerra K., Ramírez S.F. 2019.
Avifaunal surveys along the lower Huallaga River, Region of Loreto, Peru: New
distributional records, collection of topotypes, and taxonomic implications. Wilson J.
Ornithol. 131:486–501.

Muggeo V.M.R., Atkins D.C., Gallop R.J., Dimidjian S. 2014. Segmented mixed models with
random changepoints: a maximum likelihood approach with application to treatment for
depression study. Stat. Modelling. 14:293–313.

Musher L.J., Del-Rio G., Marcondes R.S., Brumfield R.T., Bravo G.A., Thom G. 2024.
Geogenomic predictors of genetree heterogeneity explain phylogeographic and
introgression history: a case study in an Amazonian bird (Thamnophilus aethiops). Syst.

Biol. 73:36–52.

Musher L.J., Ferreira M., Auerbach A.L., Cracraft J. 2019. Why is Amazonia a "source"of
biodiversity? Climate-mediated dispersal and synchronous speciation across the Andes in
an avian group (Tityrinae). Proc. Roy. Soc. B. 286:20182343.

Musher L.J., Krabbe N.K., Areta J.I. 2023. Underestimated Neotropical diversity: Integrative
taxonomy reveals two unrelated look-alike species in a suboscine bird (Pachyramphus
albogriseus). Ornithology 140:ukac047.

Pamilo P., Nei M. 1988. Relationships between gene trees and species trees. Mol. Biol. Evol.
5:568–583.

Patel S., Weckstein J.D., Patané J.S.L., Bates J.M., Aleixo A. 2011. Temporal and spatial
diversification of Pteroglossus aracaris (Aves: Ramphastidae) in the Neotropics: constant
rate of diversification does not support an increase in radiation during the Pleistocene. Mol.
Phylogenet. Evol. 58:105–115.

Phillips M.J., Gibb G.C., Crimp E.A., Penny D. 2010. Tinamous and moa flock together:
mitochondrial genome sequence analysis reveals independent losses of flight among
ratites. Syst. Biol. 59:90–107.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Lemmon E.M., Lemmon A.R.
2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA
sequencing. Nature. 526:569–573.

R Core Team. 2023. R: A language and environment for statistical computing. Foundation for
Statistical Computing, Vienna, Austria.

Reddy S., Kimball R.T., Pandey A., Hosner P.A., Braun M.J., Hackett S.J., Han K.-L., Harshman
J., Huddleston C.J., Kingston S., Marks B.D., Miglia K.J., Moore W.S., Sheldon F.H., Witt
C.C., Yuri T., Braun E.L. 2017. Why Do Phylogenomic Data Sets Yield Conflicting Trees?
Data Type Influences the Avian Tree of Life more than Taxon Sampling. Syst. Biol. 66:857–
879.

Remsen J.V. Jr, Areta J.I., Bonaccorso E., Claramunt S., Del-Rio G., Jaramillo A., Lane D.F.,
Robbins M.B., Stiles F.G., Zimmer K.J. Version 2024. A classification of the bird species of
South America. Museum of Natural Science, Louisiana State University.
http://www.museum.lsu.edu/~Remsen/SACCBaseline.htm.

Rhodes J.A., Baños H., Mitchell J.D., Allman E.S. 2021. MSCquartets 1.0: quartet methods for
species trees and networks under the multispecies coalescent model in R. Bioinformatics.
37:1766–1768.

Rivas-González I., Rousselle M., Li F., Zhou L., Dutheil J.Y., Munch K., Shao Y., Wu D.,
Schierup M.H., Zhang G. 2023. Pervasive incomplete lineage sorting illuminates speciation
and selection in primates. Science. 380:eabn4409.

Robinson D.F., Foulds L.R. 1981. Comparison of phylogenetic trees. Math. Biosci. 53:131–147.

Sackton T.B., Grayson P., Cloutier A., Hu Z., Liu J.S., Wheeler N.E., Gardner P.P., Clarke J.A.,
Baker A.J., Clamp M., Edwards S.V. 2019. Convergent regulatory evolution and loss of

flight in paleognathous birds. Science. 364:74–78.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.

Schultz D.T., Haddock S.H.D., Bredeson J.V., Green R.E., Simakov O., Rokhsar D.S. 2023.
Ancient gene linkages support ctenophores as sister to other animals. Nature. 618:110–
117.

Shen X.-X., Hittinger C.T., Rokas A. 2017. Contentious relationships in phylogenomic studies
can be driven by a handful of genes. Nat. Ecol. Evol. 1:126.

Simão F.A., Waterhouse R.M., Ioannidis P., Kriventseva E.V., Zdobnov E.M. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 31:3210–3212.

Simion P., Philippe H., Baurain D., Jager M., Richter D.J., Di Franco A., Roure B., Satoh N.,
Quéinnec É., Ereskovsky A., Lapébie P., Corre E., Delsuc F., King N., Wörheide G.,
Manuel M. 2017. A Large and Consistent Phylogenomic Dataset Supports Sponges as the
Sister Group to All Other Animals. Curr. Biol. 27:958–967.

Simmons M.P., Gatesy J. 2021. Collapsing dubiously resolved gene-tree branches in
phylogenomic coalescent analyses. Mol. Phylogenet. Evol. 158:107092.

Stadler T. 2010. Sampling-through-time in birth-death trees. J. Theor. Biol. 267:396–404.

Stiller J., Feng S., Chowdhury A.-A., Rivas-González I., Duchêne D.A., Fang Q., Deng Y.,
Kozlov A., Stamatakis A., Claramunt S., Nguyen J.M.T., Ho S.Y.W., Faircloth B.C., Haag J.,
Houde P., Cracraft J., Balaban M., Mai U., Chen G., Gao R., Zhou C., Xie Y., Huang Z.,
Cao Z., Yan Z., Ogilvie H.A., Nakhleh L., Lindow B., Morel B., Fjeldså J., Hosner P.A., da
Fonseca R.R., Petersen B., Tobias J.A., Székely T., Kennedy J.D., Reeve A.H., Liker A.,
Stervander M., Antunes A., Tietze D.T., Bertelsen M.F., Lei F., Rahbek C., Graves G.R.,
Schierup M.H., Warnow T., Braun E.L., Gilbert M.T.P., Jarvis E.D., Mirarab S., Zhang G.
2024. Complexity of avian evolution revealed by family-level genomes. Nature 629:851–
860.

Valqui T. 2009. Phylogeogaphy of Nothoprocta Tinamous and the The Phylogeny of the
Tinamidae. Diss. Louisiana State Univesity and Agricultural & Mechanical College.

Wen D., Yu Y., Nakhleh L. 2016. Bayesian Inference of Reticulate Phylogenies under the
Multispecies Network Coalescent. PLoS Genet. 12:e1006006.

Yonezawa T., Segawa T., Mori H., Campos P.F., Hongoh Y., Endo H., Akiyoshi A., Kohno N.,
Nishida S., Wu J., Jin H., Adachi J., Kishino H., Kurokawa K., Nogi Y., Tanabe H.,
Mukoyama H., Yoshida K., Rasoamiaramanana A., Yamagishi S., Hayashi Y., Yoshida A.,
Koike H., Akishinonomiya F., Willerslev E., Hasegawa M. 2017. Phylogenomics and
morphology of extinct paleognaths reveal the origin and evolution of the ratites. Curr. Biol.
27:68–77.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

Zhang R., Drummond A. 2020. Improving the performance of Bayesian phylogenetic inference

under relaxed clock models. BMC Evol. Biol. 20:54.

Zhao M., Kurtis S.M., White N.D., Moncrieff A.E., Leite R.N., Brumfield R.T., Braun E.L., Kimball
R.T. 2023. Exploring Conflicts in Whole Genome Phylogenetics: A Case Study Within
Manakins (Aves: Pipridae). Syst. Biol. 72:161–178.