1    **Article Type:** Research Article
2
3    **Whole-genome phylogenomics of the tinamous (Aves: Tinamidae): comparing gene tree**
4    **estimation error between BUSCOs and UCEs illuminates rapid divergence with**
5    **introgression**
6
7    **Running title:** Whole-genome phylogenomics of the tinamous
8

9    **Authors:**
10   Lukas J. Musher[1], Therese A. Catanach[1], Thomas Valqui[2], Robb T. Brumfield[3], Alexandre
11   Aleixo[4], Kevin P. Johnson[5], Jason D. Weckstein[1,6]
12
**Author affiliations:**
[1] The Academy of Natural Sciences of Drexel University, Department of Ornithology
Philadelphia, PA, 19103, USA.
[2] Facultad de Ciencias Forestales, Universidad Nacional Agraria La Molina, Lima, Peru and
CORBIDI-Centro de Ornitología y Biodiversidad, Lima, Peru.
[3] Department of Biological Sciences and Museum of Natural Science, Louisiana State
University, Baton Rouge, LA, 70803, USA.
[4] Instituto Tecnológico Vale - ITV, Belém, Brazil.
[5] Illinois Natural History Survey, Prairie Research Institute, University of Illinois Urbana-
Champaign, Champaign, IL, USA.
[6] Department of Biodiversity, Earth, and Environmental Sciences, Drexel University,
Philadelphia, PA, 19103, USA.
25

**Abstract:**
Incomplete lineage sorting (ILS) and introgression increase genealogical discordance across the
genome, which complicates phylogenetic inference. In such cases, identifying orthologs that
result in gene trees with low estimation error is crucial because phylogenomic methods rely on
accurate gene histories. We sequenced whole genomes of tinamous (Aves: Tinamidae) to
reconstruct their interrelationships and dissect the sources of gene tree and species-tree
discordance.  We compared results based on five ortholog sets: (1) coding genes (BUSCOs), (2)
ultraconserved elements (UCEs) with short flanking regions, (3) UCEs with intermediate flanks,
(4) UCEs with long flanks, and (5) UCEs mapped to the Z-chromosome. We hypothesized that
orthologs with more phylogenetically informative sites would result in more accurate species
trees because the resulting gene trees contain lower stochastic error. Consistent with our
hypothesis (and a large body of theory), we found that long UCEs had the most informative sites
and lowest rates of error. Surprisingly, BUSCO gene trees contained high error compared to long
UCEs, despite having many informative sites. Unlike UCEs, BUSCO gene sequences showed a
positive association between the proportion of informative sites and gene tree error. Thus, the
underlying properties of molecular evolution differ between BUSCO and UCE datasets, and these
differences should be considered when selecting loci for phylogenomic analysis. Importantly, these
results indicate stochastic error is not driving inaccurate gene tree estimation for BUSCO loci,
instead suggesting a more problematic impact of systematic error in this data-type. Still, species
trees from different datasets were mostly congruent. Only one clade, which has a history of ILS
and introgression, exhibited substantial species-tree discordance across the different data sets.
We suggest that agreement between the Z-chromosome dataset and that of long UCEs lends

support to this topology because the Z-chromosome is expected to contain low rates of ILS and
faster coalescent times due its relatively smaller effective population size. Overall, we present
the most complete phylogeny for tinamous to date, identify an unrecognized species, and provide
a case study for species-level phylogenomic analysis using whole-genomes.
52
53 **Keywords:** BUSCO; Protein-coding genes; Concatenation; Data biases; Divergence; Estimation
54 error; Gene tree; Incomplete Lineage Sorting; Introgression; Phylogenomics; Species Tree;
55 Tinamous; UCEs; Ultraconserved elements; Whole-genome sequencing
56
57 **Introduction**
58
Despite the growth of high-throughput sequencing approaches over the past decade,
reconstructing the tree of life remains challenging. Analyses that utilize alternative datasets,
methodological frameworks, or substitution models to answer the same phylogenetic questions
often yield conflicting results, which is surprising given that modern phylogenomic datasets
typically contain hundreds of thousands or even millions of phylogenetically informative
characters (Dunn et al. 2008; Jarvis et al. 2014; Prum et al. 2015; Reddy et al. 2017; Simion et al.
2017; Franz et al. 2019; Schultz et al. 2023; Stiller et al. 2024). In many cases, different methods
or data yield equally well-supported but conflicting phylogenetic topologies. Thus, two key
questions for systematists are, (1) why do phylogenomic datasets yield discordant topologies?
and (2) how should one choose among conflicting well-supported topologies?
One way in which discordant phylogenies can result from the same or similar datasets is
from the use of different methods for reconstructing species trees. One frequently employed
method is to concatenate all alignments into a single supermatrix and treat the resulting
phylogeny as a genome-wide average (henceforth, concatenation). Another common approach is
to estimate a gene tree for each molecular marker independently and summarize their histories to
estimate the species trees based on expectations from a multi-species coalescent (MSC) model
(Zhang and Mirarab 2022). Unlike the concatenation approach, MSC methods are designed to
account for expected variation in gene histories due to incomplete lineage sorting (ILS). ILS
causes some gene trees to differ from the species tree because ancestral variation is retained
during speciation, resulting in alleles that coalesce before speciation events (Maddison 1997).
Each of these two methods, concatenation and MSC, has theoretical advantages, but can
result in erroneous species trees under certain conditions. For example,concatenation may be the
best method when ILS is weak, but is likely to be statistically inconsistent when ILS is strong
(Mendes and Hahn 2018; Bryant and Hahn 2020). This is because when ILS is strong, gene
genealogies are expected to differ more from the species tree compared to situations when ILS is
weak, thus reducing the probability that concatenation will find the true tree. ILS is more likely
to occur when successive divergence times are short and effective population sizes are large,
such as during rapid or adaptive radiations (Maddison 1997; Mclean et al. 2019; Lescroart et al.
2023; Tan et al. 2023). Thus, discordance among concatenation and MSC methods may be in
part driven by the presence of rapid diversification that leads to ILS and high gene tree
heterogeneity that biases concatenation results.
When applying MSC methods, identifying orthologous markers that lead to accurate,
unbiased gene tree estimation is crucial to properly infer phylogeny (Meiklejohn et al. 2016;
Springer and Gatesy 2016; Simmons and Gatesy 2021; Tea et al. 2021). MSC methods that
utilize gene trees as input (gene tree summarization methods) may fail if gene tree heterogeneity

is driven by estimation error rather than coalescence events. In other words, the MSC model is
intended to incorporate gene tree heterogeneity due to ILS, but erroneous gene trees resulting
from low quality, biased, or inconsistent sequence data will increase species tree estimation error
(Xi et al. 2015). Thus, dissecting biological and artifactual sources of gene tree discordance to
more accurately infer species trees is a major challenge in the age of phylogenomics, where more
data does not automatically translate to improved inferences (Rodríguez-Ezpeleta et al. 2007;
Meiklejohn et al. 2016; Blom et al. 2017; Smith et al. 2023). Because MSC methods assume that
all gene tree discordance is driven by ILS, eliminating erroneous gene trees becomes important
(Roch and Warnow 2015; Springer and Gatesy 2016). This point has led to a variety of methods
for filtering genomic datasets for phylogenetic efficacy (Doyle et al. 2015; Kuang et al. 2018;
Smith et al. 2018a, 2018b; Zhao et al. 2023), but has generally lacked consensus on the types of
data that contain low error to begin with.
Given the complexities associated with phylogenetic inference, identifying datasets of
gene orthologs containing minimal bias is a principal goal of systematics research and has
implications across diverse fields in molecular biology. One widely employed data type for
phylogenomics includes coding sequences obtained via anchored hybrid enrichment,
transcriptomics, or low coverage whole-genome sequencing (Lemmon et al. 2012; Prum et al.
2015; Allen et al. 2017; Waterhouse et al. 2017; Johnson 2019; Zhang et al. 2019; Burbrink et al.
2020; Boyd et al. 2022; Van Damme et al. 2022). Another data type includes ultraconserved
elements (UCEs), which target conserved genomic regions to identify orthology, but also yield
sequences of their more variable flanking regions for phylogenetic reconstruction across a range
of timescales (Faircloth et al. 2012, 2013; McCormack et al. 2012; Musher and Cracraft 2018;
Gueuning et al. 2020; Catanach et al. 2021; Ostrow et al. 2023). Both data types are widely
applied in phylogenomics but can sometimes result in different phylogenetic topologies (Reddy
et al. 2017). Nevertheless, there have been few studies comparing the efficacy of these marker
types to one another. Thus, the ability of coding versus primarily non-coding markers to resolve
phylogenomic trees remains an open question.
One method for evaluating ortholog quality and gene tree error is through quantification
of alignment information content, such as the number of parsimony informative sites
(henceforth, PIS), or by comparing levels of gene tree heterogeneity among data types (Fong and
Fujita 2011; Harris et al. 2014; Burbrink et al. 2020; Leite et al. 2021; Smith et al. 2023). For
example, alignments with few PIS often result in erroneous or poorly resolved gene trees
because of a lack of phylogenetic signal in the data, a problem referred to as stochastic error (Xi
et al. 2015; Meiklejohn et al. 2016; Kapli et al. 2021). Here phylogenetic resolution increases as
phylogenetically informative characters are added due to a reduction in stochastic error driven by
insufficient sampling. For example, alignments of coding genes, or UCEs that include long
flanking regions, should contain many PIS, and therefore result in lower rates of stochastic error
than UCEs with short flanking regions that contain relatively few variable sites. Some studies
have found that loci with more PIS result in more "clock-like" gene trees, suggesting that more
informative loci result in gene trees with branch lengths that represent real biological signal
rather than data-driven artifacts (Musher et al. 2019). Other studies have shown a negative
correlation between the number of PIS and Robinson-Foulds distance (a measure of phylogenetic
dissimilarity; henceforth, RF distance) between gene and species trees (Burbrink et al. 2020).
This relationship implies that more informative alignments result in reduced stochastic error.
However, as an alternative, one might also expect that phylogenetic markers that are too variable

can result in erroneous gene trees, if phylogenetic signal becomes lost due to sequence saturation
(Felsenstein 1978; Brinkmann et al. 2005).
Another, more troublesome source of inaccurate gene trees in phylogenetic analysis is
systematic error, which is driven by data biases that cause model violations (e.g., GC content
biases or multiple fast-evolving branches) (Rodríguez-Ezpeleta et al. 2007). For instance,
analyses of coding versus non-coding markers have resulted in conflicting topologies for the
backbone of Neoaves (McCormack et al. 2013; Jarvis et al. 2014; Prum et al. 2015). Some
authors have suggested that this discordance can be attributed to data biases associated with
coding genes (e.g., elevated GC content and model misspecification) that also bias downstream
species tree inferences (Reddy et al. 2017; Stiller et al. 2024). Although an observed negative
association between alignment information content and gene tree error can be viewed as an
inverse correlation between information content and stochastic estimation error, deviations from
this relationship may suggest an effect of systematic error. Thus, documenting the relationship
between alignment information content and gene tree error, as well as the variation in overall
gene tree heterogeneity among different datasets, should inform future phylogenomic study
design and illuminate data-type efficacy.
Whole-genome sequencing offers a way to test the robustness of a phylogenetic topology
given multiple types of orthologous markers, including both coding and non-coding sequences,
from the same underlying data. For example, whole genomes are often used to harvest large
numbers of single-copy orthologous protein-coding markers called BUSCO (Benchmarking
Universal Single Copy Orthologs) genes (Waterhouse et al. 2017; Alaei Kakhki et al. 2023).
Given their conservation of function, BUSCO genes can theoretically tackle both relatively deep
and shallow phylogenetic problems (Timilsena et al. 2022; Van Damme et al. 2022; Alaei
Kakhki et al. 2023). Similarly, there are multiple pipelines that can easily be used to obtain
UCEs and other conserved elements for phylogenomics from whole-genome datasets (Faircloth
2016; Edwards et al. 2017). Thus, sequencing whole genomes enables a robust test of the relative
influences of stochastic and systematic error on phylogenetic reconstruction, and thus the
efficacy of different data types.
In this study, we apply whole-genome sequencing to reconstruct a species-level
phylogeny of the tinamous (Aves: Tinamidae). There has been limited work on tinamou
phylogenomics to date. Most past phylogenetic studies utilized either morphological data with
relatively few genetic markers (Bertelli et al. 2002; Bertelli and Porzecanski 2004; Valqui 2008;
Bertelli 2017; Almeida et al. 2022), or phylogenomic data with large-scale molecular sampling
but minimal taxonomic sampling (Cloutier et al. 2019). Thus, the species-level phylogenetic
relationships of the tinamous remain uncertain. Here, we build on past studies by sampling
thousands of orthologous markers across 45 out of 46 described species and a total of 65 named
taxa (monotypic species plus subspecies). Specifically, we compare and contrast levels of PIS
and gene tree estimation error in two types of orthologous markers: complete single-copy
protein-coding genes (BUSCOs) and UCEs. In doing so, our objectives are to (1) examine the
effect of alignment information content on gene tree estimation error to dissect sources of
stochastic and systematic gene tree estimation error, (2) use this information to identify high
quality (low error) orthologs, (3) identify the biological drivers of phylogenetic discordance
among data types and methods, and (4) use these inferences to robustly reconstruct the
evolutionary history of the tinamous.
183
184 **Materials and methods**

185
*Sampling*

187
A total of 46 species of tinamous classified into nine genera are currently recognized (Clements et al. 2023). Of these, 22 are monotypic and the remaining species contain two or more subspecies. Altogether there are 175 named taxa (monotypic species plus subspecies) recognized in Tinamidae. We sampled 50 fresh tissues, 12 toe pads from historical museum study skins, and downloaded 10 whole-genome assemblies from the NCBI Assembly Archive, spanning a total of 66 named taxa (subspecies plus monotypic species) and 45 recognized tinamou species. Our sampling included all recognized species except one; a sample initially identified as *Crypturellus boucardi* turned out to be *C. soui meserythrus* (Voucher LSUMZ Birds 180663, Tissue LSUMNS B-53413).

Tinamous belong to the infraclass, Palaeognathae, which includes many extant flightless ratites such as ostriches (Struthionidae) and rheas (Rheidae) along with extinct forms such as Moas (Dinorninithiformes). Recent work has indicated that moas are the sister group to tinamous and that rheas belong to a clade that is sister to tinamous plus moas (Cloutier et al. 2019). Thus, as outgroup taxa for tree rooting, we also downloaded whole genome assemblies for *Rhea pennata* (Greater Rhea) and *Anomalopteryx didiformis* (Little Bush Moa) from the NCBI Assembly Archive. Table S1 lists the 74 samples (72 tinamous plus two outgroups) included in this study.

*DNA extraction, library preparation, and whole genome sequencing*

For fresh tissues, we extracted genomic DNA (gDNA) using the MagAttract High Molecular Weight kit from Qiagen (Valencia, California). Toe pads extraction of historical study skins was carried out in a dedicated historical DNA extraction laboratory at the Academy of Natural Sciences of Drexel University to reduce contamination risk. Toe pad samples were first washed in a brief bath of EtOH to help remove superficial contaminants, and then soaked in $H_2O$ for three hours to hydrate the desiccated flesh for DNA lysis. Samples were then digested using 180 μL ATL and 30 μL proteinase K for each sample and incubated at 56º C overnight. DNA isolation was then performed using the QiaQuick spin columns and extraction kit from Qiagen (Valencia, CA).

Shotgun sequencing libraries were prepared for each extract using the Hyper library construction kit (Kapa Biosystems). These libraries were sequenced using 150 bp paired-end reads on an S4 lane of an Illumina NovaSeq 6000. These libraries were pooled and tagged with unique dual-end adaptors, and pooling consisted of between 13 and 18 samples per lane aimed to achieve at least 30X coverage of the nuclear genome. Adapters were trimmed and demultiplexed using bcl2fastq v.2.20. We deposited raw reads in the NCBI SRA (Table S1).

*Whole genome assembly*

We cleaned and then mapped raw reads for each sample (Table S1) to a closely-related scaffold-level assembly from NCBI. Decisions about which downloaded genome to use for read mapping were made based on a previous tinamou phylogeny (Almeida et al. 2022). Specifically, we used fastp (Chen et al. 2018) to clean fastq files, trim adapters and remove low quality reads.

Next, we used Burrows-Wheeler Aligner (BWA) version 0.7.17 (Li and Durbin 2009) to map the
cleaned reads to reference genomes. Once reads were mapped, we used samtools version 1.6 to
convert the resulting sam-files into sorted bam-files (Li et al. 2009) and Analysis of Next
Generation Sequencing Data (ANGSD) version 1.2.13 (Korneliussen et al. 2014) to convert
bam-files into fasta format. We used '-doFasta 2' to ensure that the consensus nucleotide was
written for each polymorphic site.
To assess genome quality, completeness, and redundancy for each assembly, we used
Benchmarking Universal Single Copy Orthologs (BUSCO) version 5.3.0 (Simão et al. 2015).
BUSCO searches genome assemblies and identifies genes present in single copy using a database
of known single-copy orthologs from a clade-specific database of genes. We used the
'aves_odb10' lineage dataset, which utilizes 8,338 genes in the chicken genome. We used the '--
augustus' flag to obtain nucleotide sequences for genes in addition to amino acid sequences. This
setting uses augustus version 3.2 (Hoff et al. 2019) to annotate each assembly, and outputs full
nucleotide gene sequences for all complete single-copy orthologs in fasta format. This step was
necessary to obtain our coding gene dataset for phylogenomic analysis. We also used samtools to
estimate mean and standard deviation of sequence coverage for each genome.
247
248  *Ortholog identification and alignment*
249
We identified and extracted two types of orthologous markers from the WGS scaffolds.
First, we utilized nucleotide sequences for the 8,338 single copy orthologs obtained from the
BUSCO 'aves_odb10' dataset (henceforth, BUSCO dataset). We used custom scripts to append
orthologous genes into the same text file and then used MACSE (Multiple Alignment of Coding
SEquences) version 2.06 to align nucleotide sequences (Ranwez et al. 2018). MACSE is a
codon-aware alignment algorithm that aligns nucleotide sequences with respect to their protein
translation, but allows nucleotide sequences to contain exceptions to this underlying codon
structure. This allowed us to accurately align each of the 8,338 gene sequences for all 74
samples.
Second, we harvested ultraconserved elements (UCEs) from each genome assembly.
UCEs are highly conserved, typically single-copy, sequences in the genome that are flanked by
neutral sites that become more variable moving away from the conserved core (McCormack et
al. 2012). These may overlap with coding or non-coding sequences, but the general pattern of
conserved UCE core with variable flanking region should result in very different distribution of
variation within a locus than the BUSCO genes, which are purely coding and are most likely to
vary at third codon positions due to the redundant nature of the genetic code. To identify and
extract UCEs from the genome we followed the pipeline outlined in Phyluce version 1.7.1
(Faircloth 2016) for harvesting and loci from whole genomes. We specifically harvested UCE
loci using the Tetrapods-UCE-5kv1 dataset, which includes 5,060 UCEs.
In this pipeline, phyluce scripts are used to align probe sequences (corresponding to
conserved UCE cores) to the genome and then extract those sequences from the genome with a
user-specified length of flanking region. Because we were interested in understanding whether
and how information content covaried with gene tree estimation error, we harvested UCEs three
times, varying the length of the flanking region included with the UCE core each time. We
harvested UCEs from each genome that included 100 bp (similar to sequence lengths obtained
with target capture from degraded DNA datasets such as historical museum skins; henceforth
UCE100Flank dataset), 300 bp (similar to sequence lengths obtained with target capture from

standard fresh tissue; e.g., Musher and Cracraft, 2018, Tea et al. 2021; henceforth UCE300Flank
dataset), and 1,000 bp (longer than typical of target capture datasets; henceforth UCE1000Flank
dataset) of flanking region. We then used MAFFT (Katoh and Standley 2013) to align
orthologous UCE loci for downstream phylogenomic analysis.
Because missing data can be an additional source of phylogenetic noise, we evaluated
each of the four datasets (three UCE and one BUSCO) using only complete (alignments for
which all samples are included) alignments. This helped to eliminate gene tree discordance and
other sources of error deriving from variation in missing data content.
Finally, to further test a difficult-to-resolve clade (*Crypturellus* Clade A, see Results), we
created a dataset of UCEs that mapped to the Z-chromosome (an avian sex chromosome with
idealized $N_e$ of 0.75 autosomal $N_e$) in *Rhea pennata* genome (NCBI refseq assembly:
GCF_028389875.1). To do so, we first used the above phyluce commands to locate UCEs in the
Rhea genome. We then made a list of UCE loci mapping to the Z-chromosome by using "grep"
in UNIX extract lines in the '.lastz' file containing UCEs on the Z-chromosome. Next, we
filtered alignments to only include taxa in the difficult-to-resolve clade of interest. Thus, our
final dataset consisted of alignments of UCEs with 1000 bp of flanking region that mapped to the
Z-chromosome in *Rhea* (henceforth, Z-chromosome dataset) and only include 15 samples in one
clade in the genus *Crypturellus*.
295
296 *Phylogenomic analyses*
297
For each of the five datasets, we employed both a concatenation approach using iqtree
v2.0.3 (Minh et al. 2020) and an MSC approach using ASTRAL version 5.7.7 (Mirarab et al.
2014; Zhang et al. 2018). First, for each of the datasets, we used iqtree to concatenate all
alignments and used the edge-linked proportional partition model to partition by locus
(Chernomor et al. 2016). In iqtree, we also implemented ModelFinder (Kalyaanamoorthy et al.
2017) to choose the best substitution model for each partition from all possible models. We then
used iqtree to reconstruct the phylogeny and perform 1000 ultrafast bootstrap replicates (Hoang
et al. 2018). Next, for our MSC approach, we estimated gene trees (phylogenies derived from
presumed independently sorting loci) for each alignment within each of the five datasets again
using iqtree, ModelFinder to identify the best-fit substitution model, and again performed 1000
ultrafast bootstraps on each gene tree. We then collapsed all low support (bootstrap<75) nodes
within each gene tree (Simmons and Gatesy 2021). Using these gene trees, we implemented a
quartet-based MSC approach in ASTRAL. ASTRAL approximates the MSC model by
estimating the proportions of all possible four-taxon statements (quartets) among all estimated
gene trees and then summarizing across the genome. This approach assumes that all gene tree
heterogeneity (discordance among estimated gene histories) is due to ILS without significant
ancestral introgression or gene tree estimation error. Finally, in addition to estimating node
bootstrap support (iqtree) and posterior probability (Astral), we assessed node support by
calculating gene concordance factors for all trees using iqtree. Gene concordance factors are the
percentage of gene trees that contain a given node in a species tree. The collapsed gene trees
obtained at this phase of data analysis were used as the input gene trees for all subsequent
analyses outlined below.
320
321 *Assessing ortholog quality and gene tree error*
322

To evaluate general differences among dataset variability, we first estimated the average
variability among sites for each alignment within each dataset (excluding the Z-chromosome
dataset, which is overlapping with the UCE1000Flank dataset). To do so, we quantified the
average variability for each nucleotide site relative to the center of the alignment using the
'phyluce_align_get_smilogram_from_alignments' script available in the Phyluce package
(Faircloth 2016). Then, to assess the relationships between alignment information content, data-
type, and gene tree estimation error, we wrote custom scripts in R version 4.3.1 (R Core Team
2023) to estimate the number and proportion (informative sites/alignment length) of parsimony
informative sites at each locus, as well as the RF distances (Robinson and Foulds 1981) between
each estimated gene tree and the inferred species tree. RF distances were quantified using the
'RF.dist' function in phangorn version 2.11.1 (Schliep 2011). Parsimony informative sites were
defined as variable sites in the alignment wherein each variant nucleotide is represented by at
least two samples. To estimate RF distances, we compared the gene trees from each dataset
inferred using iqtree to the UCE1000Flank MSC tree based on the UCE1000Flank dataset. To
test for differences in these measures among datasets, we used Kruskall-Wallace and pairwise
Wilcoxon rank sum tests with a Benjamini-Hochberg p-correction implemented in base R (R
Core Team 2023). We then ran generalized linear models using information content as the
independent variable and RF distance as the dependent variable for each of the four datasets.
Because we found no differences in RF distance when using MSC and concatenated species
trees, for these models, we only used RF distances relative to the MSC species tree resulting
from our ASTRAL analysis. We compared AIC values for linear and logarithmic fits for each
regression and chose the model with the lowest AIC for each dataset. These regression analyses
were done once for each dataset, and once with all three UCE datasets combined. Although the
latter analysis contains the same UCE locus multiple times, albeit with additional flanking
regions, we believe this analysis helps illuminate how the information content of loci from across
the spectrum (i.e., very uninformative to very informative) influences rates of stochastic gene
tree error in UCEs compared with BUSCOs. We also explored these relationships after filtering
the BUSCO for GC content (by removing genes with GC content greater than two standard
deviations from the mean) and for genes resembling UCE datasets (removing genes greater than
two standard deviations above the mean proportion of PIS in the UCE1000Flank dataset and
below the mean of the UCE100Flank dataset).
354
355     *Tests of incomplete lineage sorting and introgression*
356
To examine the effects of ILS and introgression on the phylogenetic discordance of Clade
A, we first looked at the alternative quartet topologies for five short internal branches and then
used a phylogenomic network approach to test for reticulation (phylogenomic non-bifurcation).
Quartets are unrooted four taxon statements for which only three alternative sets of relationships
(topologies) exist. Because we found conflicting phylogenetic relationships between datasets for
a clade of species in the genus *Crypturellus* (Figure 3), we wanted to assess the extent to which
this conflict could be explained by ILS due to rapid divergences or introgression between non-
sister taxa. The MSC model assumes that all gene tree heterogeneity arises via ILS (i.e., no
introgression, no gene tree error, no recombination, etc.), and it makes explicit predictions about
the relative frequencies of alternative quartet topologies for all internal branches in a phylogeny.
Specifically, the MSC model predicts one majority topology that is consistent with the species
tree at a relative frequency $>\frac{1}{3}$ and two minority topologies of equal relative frequencies $<\frac{1}{3}$

(Pamilo and Nei 1988). As ILS increases, the relative frequencies of all three alternative quartets
approach the ⅓ threshold. Deviations from these expectations (i.e., minority topologies are not
equivalent in frequency) suggests that processes such as introgression may be influencing gene
tree heterogeneity. For the quartet method, we used custom scripts that utilize the R package
'MSCquartets' to identify the relative quartet frequencies for five nodes of interest (Rhodes et al.
2021; Allman et al. 2023). Because ILS is expected at relatively short internal branches, we
qualitatively evaluated the relative frequencies of alternative topologies at these branches to
assess whether ILS might be involved in increasing genealogical heterogeneity, and thus
phylogenetic disagreement of a single difficult-to-resolve Clade (Clade A).
To test for putative introgression, we applied a Bayesian approach implemented in
PhyloNet 3.8.0 (the 'MCMC_GT' algorithm) (Wen et al. 2016). This method employs reverse-
jump Markov Chain Monte Carlo (rjMCMC) to sample the posterior distribution of phylogenetic
networks under a multi-species network coalescent (MSNC) model using only rooted gene trees
as input data. The MSNC is similar to the MSC, but relaxes the assumption of no introgression
by modeling genome evolution as a network rather than bifurcating phylogeny. In doing so, it
accounts for both incomplete lineage sorting and reticulation (multiple ancestral lineages that
contribute alleles to a single daughter lineage). Such reticulation is typically attributed to
historical introgression. We used the gene trees with collapsed low-support nodes from the
UCE1000Flank dataset as these contained reduced rates of both stochastic and systematic gene
tree estimation error. We ran the analysis three times, varying the maximum number of reticulate
nodes in each run to assess variability in the results. Specifically, we ran MCMC_GT allowing
for a maximum of between zero (m=0) and ten (m=10) reticulate nodes. We assigned all samples
to the species level, and ran the analysis using species as tips in the network rather than samples
(i.e., some species were represented by multiple taxa). We ran the rjMCMC for $5 \times 10^7$
generations with a burn-in of $5 \times 10^6$ generations, using the pseudo-likelihood calculation to
reduce computation time. Because likelihood scores tended to increase with each successive
increase in m-value, and the rjMCMC typically found the maximum allowed in each run, we
chose the optimal m-value using a breakpoint analysis. Specifically, using the R package
'segmented' (Muggeo et al. 2014), we chose the network with the optimal m-value by fitting a
segmented linear model to our likelihoods for each m-value (Muggeo et al. 2014). This allowed
us to identify breakpoints in the slope of the regression, where increases in m-value resulted in
diminishing gains in likelihood. The m-value at the breakpoint in the segmented model was
chosen as the optimal m-value.
402
403 **Results**
404 *Assembly metrics, completeness, and ortholog metrics*
405
We successfully assembled 61 of 62 newly sequenced genomes and extracted BUSCO
and UCE targets from these assemblies plus 10 publicly available tinamou and 2 publicly
available outgroup whole genome assemblies (Table S1). Genome wide sequence coverage
varied within and among samples, but was generally high; the mean of average coverage across
samples was 40.41x (standard deviation = 13.55x). Most genomes were also relatively complete,
containing an average of 89.69% (standard deviation = 10.88%) of 8,338 complete single-copy
BUSCO genes.
From these whole genomes, we extracted and aligned two types of orthologous markers,
BUSCO genes and UCEs. Specifically, we compiled five different ortholog sets, which after

filtering for complete occupancy (all 72 samples represented in each alignment) included (1)
2,507 BUSCOs (6,368,028 bp in concatenated alignment), (2) 2,887 UCEs with 100 bp flanking
regions (969,279 bp in concatenated alignment), (3) 2,879 UCEs with 300 bp flanking regions
(2,227,921 bp in the concatenated alignment), (4) 2,803 UCEs with 1000 bp flanks (6,902,014 bp
in the concatenated alignment), and (5) 206 UCEs with 1000 bp flanks mapped to the Z-
chromosome (504,750 bp in concatenated alignment).
421
*Phylogenomics of Tinamous*
We reconstructed the phylogenetic history of tinamous using each of the five ortholog datasets,
applying both concatenation and a gene tree summarization approach (ASTRAL; henceforth MSC phylogeny).
The reconstructed concatenated and MSC phylogenies for tinamous were well-supported and
broadly congruent across data types (Figures 1, 2, S1–S10). The phylogenies recovered from all
five datasets were entirely congruent except for relationships within a single clade (henceforth,
Clade A) of *Crypturellus* containing 13 named taxa across nine recognized species. The
placement of *C. atrocapillus* was most difficult, but was most often recovered as sister to a clade
containing *C. kerriae, C. erythropus, C. duidae,* and *C. strigulosus* (Figure 3). However, the
UCE1000Flank MSC tree and concatenate Z-chromosome tree recovered *C. atrocapillus* as
sister to a clade that also included *C. transfasciatus* and *C. cinnamomeus.*
433

434
**Figure 1: Concatenated Phylogeny of tinamous based on UCEs with 1,000 bp of flanking sequence.** All nodes
have bootstrap value of 100% except where noted. Note relatively short internodal branch lengths in Clade A.
Tinamou illustrations by TAC.

438
439  **Figure 2: Multi-species Coalescent (ASTRAL) Phylogeny of tinamous based on UCEs with 1,000 bp of**
440  **flanking sequence.** All nodes have posterior probability of 1. Branches labeled B1–B5 denote rapidly diverging
441  branches evaluated using quartet analysis (see Figure 6). Tinamou illustrations by TAC.
442

Figure 3: Phylogenetic topologies for a difficult to resolve clade (Clade A) of *Crypturellus* tinamous. The phylogenetic placement of *C. atrocapillus* (pink branch) was discordant among datasets and species tree approaches. Node support is shown for internal branches of interest, including bootstrap value or posterior probability (first value) and gene concordance factor (second value). Bottom right heatmap shows Robinson-Foulds distances between all pairs of trees. Detailed node support values can be found in the electronic supplementary material.

One genome, downloaded from NCBI (GCA 013389825) was labeled as *C. undulatus*, but did not cluster with other members of that species, instead clustering with either *C. strigulosus* (most trees) or *C. erythropus* (concatenated BUSCO tree). We could not confirm the species identity of this sample because the voucher was unavailable. Moreover, another downloaded genome, GCA 013398335, was listed as *Nothoprocta ornata* on NCBI but after including newly sequenced material, this *N. ornata* clustered with *N. pentlandii* samples from Peru. After examining the voucher specimen (CORBIDI 168605), the sample was indeed confirmed as *N. pentlandii oustaleti* and not *N. ornata* as indicated in NCBI.

Our results were broadly consistent with recognized taxonomic classifications. For example, we recovered monophyletic subfamilies Nothurinae and Tinaminae and most genera and species were also monophyletic. However, we recovered two genera as non-monophyletic: the monotypic genus *Taoniscus* was embedded within *Nothura* and *Rhynchotus* was embedded within *Nothoprocta*, rendering *Nothura* and *Nothoprocta* paraphyletic. We also recovered *Nothoprocta pentlandii* as polyphyletic; one specimen from Bolivia (voucher LSUMZ 123403) was sister to *N. perdicaria*, whereas the remaining samples from Peru formed a clade that was sister to all remaining *Nothoprocta*. All other species that were represented by multiple samples were recovered as monophyletic.

*Assessment of ortholog information content and gene tree estimation error*

We recovered significant differences in the number and proportion of PIS per alignment as well as levels of gene tree estimation error among the four datasets. As expected, the UCE100Flank dataset had the fewest PIS, averaging just 45 PIS per alignment (standard deviation = 28.84), the UCE300Flank dataset averaged an intermediate number of PIS with a mean of 207 per alignment (standard deviation = 88.37), and the UCE1000Flank dataset contained the most PIS with a mean of 1036 per alignment (standard deviation = 250.57). The BUSCO alignments also contained a high number of PIS, though with very high variance, indicating a range of informative and uninformative loci (mean = 923, standard deviation = 690.08 PIS per locus) (Table 1; Figure 4). Kruskall-Wallis tests confirmed that differences in both the number ($X^2 = 578$, $df = 3$, $P < 0.00001$) and proportion ($X^2 = 6408.9$, $df = 3$, $P < 0.00001$) of PIS significantly differed among datasets. Wilcoxon rank sum tests also indicated that differences between pairwise comparisons of these datasets were all significantly different ($P < 0.00001$ for all). GC content also differed among datasets (Kruskal-Wallace $X^2 = 4079.4$, $df = 3$, $P < 0.00001$), with BUSCOs showing the highest mean and variance (mean = 0.51, standard deviation = 0.062). A Wilcoxan rank sum test confirmed that all pairwise comparisons of datasets differed in GC content ($P < 0.0001$) except between UCE100Flank and UCE300Flank datasets ($P=0.39$).

| Dataset | Mean #PIS | StDev #PIS | Mean prop. PIS | StDev prop. PIS | Mean RF | StDev RF | Mean GC | StDev GC |
|---|---|---|---|---|---|---|---|---|
| BUSCOs | 922.51 | 690.08 | 0.37 | 0.10 | 48.00 | 16.70 | 0.51 | 0.06 |
| UCE100Flank | 45.20 | 28.84 | 0.13 | 0.08 | 107.96 | 20.16 | 0.39 | 0.06 |
| UCE300Flank | 206.87 | 88.37 | 0.26 | 0.10 | 46.59 | 13.05 | 0.39 | 0.06 |
| UCE1000Flank | 1035.71 | 250.57 | 0.42 | 0.08 | 26.66 | 7.13 | 0.40 | 0.05 |

**Table 1:** Means and standard deviations of the number and proportion of parsimony informative sites (PIS), Robinson-Foulds Distances between gene and species trees (RF), and GC content (GC) for each dataset analyzed for the study.

To measure gene tree estimation error, we examined gene tree heterogeneity across datasets using Robinson-Foulds (RF) distances between gene and species trees. Although some heterogeneity is expected due to the stochastic nature of the coalescent process, we assumed that increases in the mean and variance of RF distances between datasets were attributable to increased gene tree estimation error. The UCE100Flank dataset had the highest RF distances between gene and species trees, whereas the UCE1000Flank dataset had the lowest mean and variance in RF distances (Figure 4; Table 1). The BUSCOs and UCE300Flank datasets had intermediate RF distances, with similar means and variances (Table 1). Interestingly, RF distances from the BUSCO dataset were highly variable, with estimated gene trees from some loci showing relatively low (akin to the UCE1000Flank dataset) and some showing exceptionally high (akin to the UCE100Flank dataset) RF distances between gene and species trees. A Kruskall-Wallis test confirmed that RF-distances differed significantly among datasets ($X^2 = 3051.8$, $df = 3$, $P < 0.00001$), and a Wilcoxon rank sum test also revealed that differences between pairwise comparisons of these datasets were significantly different ($P < 0.01$ for all

comparisons). Finally, UCEs mapping to the Z-chromosome had lower average RF distance
between gene and species tree (mean=22.68, standard deviation=8.77) than those on autosomes
(mean=26.77, standard deviation=7.05), and this difference was significant (Kruskal-Wallace
$X^2$=29.299, df=1, P<0.0001). However, there was no difference in RF distances between Z-
chromosome UCEs and autosomal UCEs for UCE300Flank (Kruskal-Wallace $X^2$=0.27, df=1,
P=0.60) and UCE100Flank (Kruskal-Wallace $X^2$=0.36, df=1, P=0.55) datasets (Figure 4e).

512
**Figure 4: Information content, GC content, and gene tree heterogeneity for each dataset.** The top row of violin
plots shows information content for each dataset measured as (a) the number of parsimony informative sites (PIS)
and (b) the percentage of parsimony informative sites along with (c) GC content for each dataset. The bottom row of
violin plots shows gene tree heterogeneity for each dataset or data subset measured by the (d–e) Robinson-Foulds
distance between each gene tree and the inferred species tree (assuming the UCE1000Flank dataset) and (f) gene
concordance factors (gCF) for all nodes within each tree. Here, concordance factors were quantified for each
dataset's phylogeny (either concatenated iqtree or MSC Astral) using the collapsed gene trees from each dataset.
520
To quantify the relationship between alignment information content and gene tree
estimation error and test the hypothesis that increased information content led to reduced
stochastic gene tree estimation error, we modeled RF distances as a function of the number of
parsimony informative sites (PIS) per locus using generalized linear models. These models were
overall consistent with our hypothesized pattern of a negative association between both the
number and proportion of PIS and RF distance, with lower RF distances in gene trees built from
more informative alignments (Figure 5; Table S2). However, there was notable variation in the
slope and tightness of fit of these models among datasets. For example, the UCE300Flank
dataset, with intermediate numbers of PIS overall, had a tight negative association between RF
distance and both the number of PIS and percentage of PIS per locus (Table S2). The same
negative associations were revealed for UCE100Flank and UCE1000Flank datasets, but these
relationships were much noisier with low $R^2$, indicating a weak relationship. However, these
differences were better visualized when the three UCE datasets were combined, revealing a
strong and tightly fitting negative logarithmic relationship between the number of PIS per
alignment and RF distance between gene and species tree. Although this relationship was similar
for the BUSCO dataset using the number of PIS as the predictor variable, we found the opposite
relationship when using the percentage of PIS. Thus, BUSCOs and UCEs seem to behave
somewhat differently in relation to how similar the gene trees are to the species tree and patterns
of molecular variation. Specifically, stochastic error associated with insufficient phylogenetic
information seems to inflate RF distances in UCE gene trees, but does not show the same clear

relationship for BUSCOs. This result was robust, holding true for BUSCO datasets that were
filtered to resemble UCE-like datasets and after filtering outliers in GC content (Figure S11).

543
544 **Figure 5: Covariation between information content and gene tree estimation error.** Linear regression models
545 exploring the relationship between information content and gene tree heterogeneity. The negative correlation in most
546 instances suggests that alignments with less information content result in increased gene tree estimation error. UCEs
547 that map to the Z-chromosome in each dataset are shown in yellow.

548
549 *Tests of incomplete lineage sorting and introgression*

550
We found evidence of both ILS and introgression in driving gene tree heterogeneity in
*Crypturellus* Clade A. All five branches that we examined showed evidence of either ILS or
deviations from the MSC model (Figures 6A & S12). For branches 1 and 2 (two of the three
shortest branches within Clade A) we recovered frequencies for three alternative quartets that
closely approached the ⅓ threshold, implying ILS has promoted phylogenetic conflict at these
branches. For branches 3–5, we found possible deviations from the MSC model, though this
could be an artifact of sample size. To test whether the deviations from the MSC model observed
in Clade A could be attributed to introgression, we applied a phylogenomic network analysis,
which revealed multiple nodes involved in reticulation that may explain gene tree discordance in
*Crypturellus* Clade A (Figures 6B; electronic supplementary material). Our breakpoint analysis
revealed that m=2 was the optimal m-value for our PhyloNet analysis (Figures 6C & S13), with
diminishing gains or reduced likelihood for successively larger m-values. Specifically, the slope
of the segmented model was significant and tightly fitting (Breakpoint = 4.31, Adj. $R^2$ = 0.957, β
= 2811.1, P = 0.00019). In the first reticulate node, *C. cinnamomeus* was recovered as reticulate
between *C. transfasciatus* (inheritance probability = 0.92) and the unidentified species
(inheritance probability = 0.08). In the second, *C. undulatus* was reticulate between the basal
node of the clade (inheritance probability = 0.84) and a node containing *C. cinnamomeus* and the
unidentified taxon (inheritance probability = 0.16). In the third reticulate node, the unidentified
taxon was recovered as reticulate between *C. cinnamomeus* (inheritance probability = 0.41) and
*C. strigilosus* (inheritance probability = 0.59). And finally, a node containing *C. cinnamomeus*
and the unidentified taxon was recovered as reticulate between *C. erythropus* (inheritance

probability = 0.50) and an internal node containing all taxa except *C. noctivagus* and *C.*
*undulatus* (inheritance probability = 0.50). Thus, some gene tree heterogeneity within the low-
error UCE1000Flank dataset may be attributable to historical introgression between non-sister
taxa.

576
**Figure 6**: Incomplete lineage sorting and genomic reticulation in the tinamou phylogeny. (A) Relative quartet
frequencies for five short internal branches in Clade A based on the UCE1000Flank dataset. Bar graphs depict the
relative frequencies of each of three alternative unrooted quartet topologies (for any unrooted four taxon statement
there exist only three alternative sets of relationships). The stippled red lines indicate the ⅓ threshold for the
frequency of gene trees given a multispecies coalescent model, which predicts a single majority quartet topology
consistent with the true species tree with a frequency >⅓, and two minority topologies with equivalent relative
frequencies < ⅓. As quartet frequencies approach the ⅓ threshold, they indicate stronger ILS. Major deviations from
the expectations of the MSC model indicate violations of the model may be present, such as gene tree estimation
bias or introgression. (B) Phylogenetic network results for the network with the optimal m-value (m=4) from
PhyloNet showing non-bifurcating relationships among a difficult to resolve clade of *Crypturellus* tinamous (Clade
A). Red branches demarcate reticulation, multiple ancestors contributing to a single daughter branch. Inheritance
probability, or the proportion of the discordant gene trees that can be attributed to introgression, are marked on each
reticulate branch. Branches are labeled 1–5 and correspond to labeled branches in Figure 2. (C) The optimal m-value
was chosen using a breakpoint analysis with a segmented model.

591
592 **Discussion**
593

Here we used whole-genome sequencing to explore the biological and artifactual sources
of phylogenomic conflict and to reconstruct the species-level history of a relatively old
Neotropical avian group of broad interest in evolutionary biology (Bertelli and Porzecanski
2004; Prum et al. 2015; Altimiras et al. 2017; Sackton et al. 2019; Li et al. 2023), the tinamous.
Although we found that alignment information content and gene tree error varied considerably
within and among datasets, our results based on different analytical approaches (concatenated
and MSC methods) and datasets (UCEs and BUSCOs) were largely congruent. The topology of
only a single clade (Clade A) varied among methods and datasets (Figures 1, 2, and 3).
Specifically, the placement of *C. atrocapillus* was the most difficult node to resolve. These
results indicate that species tree estimation is, in general, robust to gene tree estimation error
when biological sources of gene tree heterogeneity are limited. Nevertheless, the concatenated
tree shows that Clade A contained multiple successive short internodal branch lengths (Figure 1),
which suggests ILS may be a key driver of gene tree heterogeneity in this clade, and thus an
important factor driving phylogenetic conflict. Indeed, quartet analysis revealed multiple internal

branches with relative quartet frequencies approaching the ⅓ threshold predicted by high ILS
(Figures 6A and S12), and relatively low gene concordance factors (Figure 3).
Cases such as that of Clade A, where ILS is strong and divergences rapid, necessitate the
use of datasets without significant gene tree estimation error to accurately infer the species tree
because concatenation is expected to fail when ILS is strong (Roch and Warnow 2015; Xi et al.
2015; Springer and Gatesy 2016). However, in addition to ILS, rapid divergence events can be
difficult to reconstruct because the short timeframe also means there is limited phylogenetic
signal for those divergence events within individual gene trees (Leaché et al. 2015; Mclean et al.
2019). For this reason, we suggest the MSC tree based on the UCE1000Flank dataset is likely the
most reliable. We suggest this dataset is reliable, first, because it had the lowest rates of gene tree
estimation error (lowest median and variance in RF distances between gene and species trees)
(Xi et al. 2015), and second because longer sequences with more PIS would be more likely to
harbor rare PIS that support rapid divergence events. Interestingly, the UCE1000Flank MSC tree
was congruent with the concatenated Z-chromosome topology. Sex-linked markers are expected
to experience reduced rates of ILS and faster coalescent times than autosomes due to a smaller
effective population size, as well as reduced rates of recombination and introgression
(Charlesworth et al. 1987; Meisel and Connallon 2013; Li et al. 2019; Moreira et al. 2023;
Musher et al. 2023). This notion is consistent with our finding that RF distances were lower for
long UCEs mapping to the Z-chromosome than for Autosomal UCEs. In both the concatenated
Z-chromosome and UCE1000Flank MSC trees, gene concordance factors were slightly higher
than for alternative topologies given the same data. Still, given the lack of agreement on the
placement of *C. atrocapillus*, further analysis may be needed to confirm the relationships within
this rapidly diversifying clade, which shows evidence of ILS and introgression between non-
sister taxa.
632
633  *Increasing alignment information content reduces estimation error for UCEs but not BUSCOs*
634
We examined the effects of data type (coding versus primarily non-coding datasets) and
information content (number and proportion of PIS) on gene tree estimation error and found that
coding sequences and genes with relatively few PIS contained higher rates of gene tree
estimation error. Although UCEs conformed to the expected negative association between PIS
and gene tree estimation error as measured using RF distance (Figure 5), BUSCOs did not.
Instead, we recovered a positive association between the proportion of PIS per alignment and RF
distance, possibly because coding genes with proportionally more variable sites are evolving
faster and thus are more prone to multiple substitutions (i.e. saturation effects), especially at third
codon positions which are less constrained. This implies that stochastic error is less impactful in
driving inaccurate gene tree estimation for BUSCO loci, and instead suggests a more problematic
impact of systematic error in this data-type.
Other studies have similarly found that coding sequences perform relatively poorly
compared with putatively neutral loci due to potential model misspecification, poor absolute
model fit, or skewed GC content (Reddy et al. 2017; Stiller et al. 2024). We therefore agree with
past studies that these differences in data properties and gene tree estimation error are associated
with biases in coding sequences that result in model misspecifications and thus systematic error.
Overall, these results show that data type and alignment information content have the potential to
compromise phylogenomic inference from gene tree summarization methods that rely on the
MSC model, even for datasets that contain relatively large amounts of data (Roch and Warnow

2015; Springer and Gatesy 2016). Interestingly, MSC results seemed most impacted by stochastic error because the UCE100Flank MSC tree was quite different from other topologies, and the BUSCO MSC tree converged on a commonly recovered topology. On the other hand, systematic error may be more likely to affect concatenation analyses, since the concatenated BUSCO tree was also quite different than other topologies (Figure 3).

We measured rates of gene tree estimation error using RF distances between gene trees and the inferred species tree, under the assumption that increases in RF distance were indicative of higher rates of phylogenetic error. Although gene histories are not expected to be congruent with the species tree in many cases (Tajima 1983; Pamilo and Nei 1988; Maddison 1997), the assumption that RF distances are good indicators of error rates is valid because different datasets contained different means and variances in RF distances among gene trees. Although the $R^2$ was small for the UCE1000Flank dataset, the overall effect was most evident when the three UCE datasets were combined into a single analysis, which showed asymptotic convergence on relatively low mean and variance RF distances after reaching about 500 PIS (Figure 5). The asymptotic shape of this relationship implies that as PIS are added to an alignment, the resulting gene trees converge on a level of heterogeneity that is representative of or approaching real biological signal (i.e., ILS and/or introgression) rather than methodological or data-driven artifacts. Moreover, if all gene tree heterogeneity was due to biological processes such as ILS, one would expect the mean and variance in RF distances to be constant among datasets because these are derived from the same underlying samples. Although some level of gene tree heterogeneity is expected in *all* phylogenomic datasets, even if one assumes the impossible, where empirical gene tree estimation error is absent (Maddison 1997; Gatesy and Springer 2014; Edwards et al. 2016), it is reasonable to assume that differences in the mean and variance in RF distances among datasets are good proxies for differences in gene tree estimation error.

*Comparisons of estimation error between data types and the benefits of whole genome phylogenomics*

We scrutinized orthologous data harvested from whole genome assemblies to identify the characteristics of phylogenomic datasets that might lead to lower bias. We found that short alignments with relatively few PIS had increased error (measured as difference between gene tree and species tree), a finding consistent with other studies (Meiklejohn et al. 2016; Burbrink et al. 2020). Likewise, we found that long UCEs tend to have low rates of gene tree estimation error and are efficacious markers to use in phylogenomic studies. We also found that the gene trees from the protein-coding dataset derived from BUSCO harvesting and splicing of exons from the tinamou genomes were very noisy. Despite containing large numbers and proportions of PIS (Figure 4) available across many genes, we found high variance in RF distances for these genes and thus conclude that such datasets might be more prone to systematic error than datasets of UCEs, the longest of which (i.e., UCE1000Flank) had lower gene tree estimation error and behaved more predictably (Figures 4–5). As a comparison, the UCE300Flank dataset also had high variance in RF distances (i.e., it contained a mix of low and high error trees), but this variance was well explained by the number of PIS in each locus. Thus, coding and UCE datasets have different underlying properties of molecular evolution and these differences should be considered when selecting loci for study and in data analysis. Despite these differences, BUSCO and UCE datasets converged on a very similar phylogenetic topology for Clade A.

Although we derived datasets from whole genome sequences, the UCE100Flank and
UCE300Flank datasets are likely to most closely match datasets obtained from typical target
capture approaches (Smith et al. 2014; Musher and Cracraft 2018; Tea et al. 2021), and are
therefore indicative of rates of error in datasets derived from those protocols. For example, many
studies utilize sequences from degraded DNA such as historical museum specimens, which may
result in many relatively short UCE alignments, but even datasets sequenced from fresh tissues
typically only capture about 100-300 bp of flanking region around the conserved UCE core.
Thus, we offer multiple recommendations for future phylogenomic studies. First, if available,
WGS data are highly preferred to sequence capture datasets because they allow for analysis of a
much wider array of data types that can be compared and contrasted as done herein (Jarvis et al.
2014; Zhang et al. 2019). Such data also allow studies to achieve longer flanking regions around
conserved UCE cores that significantly reduce bias during gene tree estimation (Figures 4–5),
and enable analyses of many other processes of interest, such as evolution of gene families.
Moreover, we did not include intron or intergenic data in the current study, but these data would
also be available from WGS and may behave more like UCE data rather than like the nearby
exons themselves (Stiller et al. 2024).
Second, if a very large genome size is cost prohibitive for WGS, then target capture
datasets such as UCEs will likely contain high variance in RF distances (i.e., a mix of "good"
and "bad" loci), but this variance is likely to be well explained by the number of PIS in each
alignment (Figure 5). We therefore suggest that these datasets should be scrutinized and filtered,
especially if a rapid radiation is involved (Doyle et al. 2015; Smith et al. 2018b, 2023; Mclean et
al. 2019). Because the number and proportion of PIS were strongly predictive of gene tree
estimation error, we suggest that simply removing alignments with too few PIS may be useful in
many cases, as has been suggested elsewhere (Harris et al. 2014; Hosner et al. 2016; Blom et al.
2017; Leite et al. 2021). Given these findings, common phylogenomic approaches that sequence
loci with intermediate or low levels of information content such as sequence capture of UCEs
may be insufficient to resolve the relationships of taxa that evolved rapidly, such as adaptive
radiations, where rates of ILS due to rapid diversification are expected to be intense (Gatesy and
Springer 2014; Mclean et al. 2019). In these situations, the number of variable sites may also be
too few to resolve gene trees involving short branches. Mapping loci to sex chromosomes, as
done here, may help clarify the relationships of these difficult nodes.
Finally, although our BUSCO dataset was noisy and included a range of high- and low-
error gene trees, we do not suggest that datasets of coding genes are inherently poor quality.
Indeed, protein-coding datasets such as BUSCOs, transcriptomes, and others are likely to remain
useful for a range of phylogenomic problems. For example, given their conservation of function,
datasets of protein-coding genes will be important for reconstructing phylogenetic relationships
at deeper timescales using either nucleotide or amino acid sequences (Dunn et al. 2008; Allen et
al. 2017; Zhang et al. 2019; Boyd et al. 2022). Still, given that our BUSCO dataset was noisy
compared to longer UCEs, we suggest exercising caution when applying protein-coding genes to
difficult phylogenomic problems (Reddy et al. 2017). Careful fitting of substitution models, data
filtering, and perhaps even manual inspection of alignments may reduce bias in these and other
datasets of protein-coding sequence (Anisimova and Kosiol 2009; Springer and Gatesy 2016).
However, our results do not suggest that BUSCOs, or protein-coding genes in general,
should be filtered by information content or RF distances. Information content was a weak and
inconsistent predictor of gene tree estimation error for BUSCOs. Alignments with fewer PIS or a
higher proportion of PIS were both associated with increased gene tree error. Likewise, some

amount of gene tree heterogeneity is expected due to the coalescent process. Thus, assuming that all high RF gene trees are erroneous and removing them from a dataset could mislead MSC analyses. More research may be needed to understand how best to filter BUSCO datasets, and perhaps protein-coding datasets in general, for phylogenomics using gene tree summarization methods.

*Phylogeny and taxonomy of the Tinamous*

Our phylogenies based on multiple datasets with varying amounts of gene tree estimation error and information content all resulted in highly similar and well-supported relationships among tinamou species (Figures 1–3, S1–S10). Moreover, concatenation and MSC approaches largely agreed on the overarching relationships at both shallow and deeper timescales. Importantly, our phylogenetic results imply that multiple taxonomic changes are necessary to appropriately classify taxa within Tinamidae. First, *Nothoprocta cinerascens* was recovered as sister to the genus *Rhynchotus*. *Rhynchotus* (Spix 1825) has nomenclatural priority over *Nothoprocta* (Sclater and Salvin 1873)*,* and therefore we suggest moving *N. cinerascens* to the genus *Rhynchotus (Bertelli and Porzecanski 2004; Valqui 2008).* Second, *Taoniscus nanus* was recovered as embedded within the genus *Nothura*, and thus, *Taoniscus* (Gloger 1842) is a junior synonym of *Nothura* (Wagler 1827)*.* We therefore suggest transferring *T. nanus* to *Nothura.* Finally, we found evidence for what is likely an unrecognized species of *Nothoprocta (Valqui 2008)*; specimens ascribed to *N. pentlandii* were polyphyletic, with a specimen of the nominate subspecies from Bolivia recovered as sister to *N. perdicaria*, and multiple specimens from Peru recovered as a clade sister to all other *Nothoprocta.* Thus, there appear to be at least two species-level taxa within *N. pentlandii*.

**Acknowledgements:**
We thank Alvaro Hernandez and Chris Wright at the University of Illinois Roy J. Carver Biotechnology Center for assistance with Illumina sequencing. We thank Kimberly K. O. Walden for assistance in depositing Illumina reads to NCBI SRA. We also thank Jorge Doña for providing R code to assist with breakpoint analysis. Dan Lane, Steve Cardiff, and Christopher Witt helped confirm voucher ID's for multiple samples. We thank B. T. Smith, J. Cracraft, P. Sweet, T. Trombone, S. Katanova (AMNH), J. Bates, S. Hackett, and B. Marks (FMNH), F. Sheldon and D. Dittman (LSUMNS), H.F. James and B. Schmidt (USNM), and N. Rice (ANSP) for loaning material that was crucial for this work. Finally, we thank the members of ANSP Ornithology Department (A. Del Grosso, E. Griffith, K. Kuabara, and J. Merwin) for comments on early versions of the manuscript. This study was funded by NSF grant DEB-1855812 to JDW and KPJ. TAC was partially supported by NSF grant DEB-2203228.

**Data Availability:**
All raw reads are available on the NCBI Sequence Read Archive (project and sample accession numbers can be found in Table S1). All data and code needed to replicate the analyses found herein can be found at LJM's github page https://github.com/lukemusher/Tinamou-phylogenomics.git or on Dryad at https://datadryad.org/stash/share/Me-IQt7EojLEAL6TpteI8zsy8sX6FWVALe0zLFkepcA.

## Literature Cited

Alaei Kakhki N., Schweizer M., Lutgen D., Bowie R.C.K., Shirihai H., Suh A., Schielzeth H., Burri R. 2023. A Phylogenomic Assessment of Processes Underpinning Convergent Evolution in Open-Habitat Chats. Mol. Biol. Evol. 40.

Allen J.M., Boyd B., Nguyen N.-P., Vachaspati P., Warnow T., Huang D.I., Grady P.G.S., Bell K.C., Cronk Q.C.B., Mugisha L., Pittendrigh B.R., Leonardi M.S., Reed D.L., Johnson K.P. 2017. Phylogenomics from Whole Genome Sequences Using aTRAM. Syst. Biol. 66:786–798.

Allman E.S., Baños H., Mitchell J.D., Rhodes J.A. 2023. MSCquartets: analyzing gene tree quartets under the multi-species coalescent. R package version 1.3.1.

Almeida F.C., Porzecanski A.L., Cracraft J.L., Bertelli S. 2022. The evolution of tinamous (Palaeognathae: Tinamidae) in light of molecular and combined analyses. Zool. J. Linn. Soc. 195:106–124.

Altimiras J., Lindgren I., Giraldo-Deck L.M., Matthei A., Garitano-Zavala Á. 2017. Aerobic performance in tinamous is limited by their small heart. A novel hypothesis in the evolution of avian flight. Sci. Rep. 7:15964.

Anisimova M., Kosiol C. 2009. Investigating protein-coding sequence evolution with probabilistic codon substitution models. Mol. Biol. Evol. 26:255–271.

Bertelli S. 2017. Advances on tinamou phylogeny: an assembled cladistic study of the volant palaeognathous birds. Cladistics. 33:351–374.

Bertelli S., Giannini N.P., Goloboff P.A. 2002. A phylogeny of the tinamous (aves: palaeognathiformes) based on integumentary characters. Syst. Biol. 51:959–979.

Bertelli S., Porzecanski A.L. 2004. Tinamou (tinamidae) systematics: A preliminary combined analysis of morphology and molecules. Ornitol. Neotrop. 15 (Suppl):293–299.

Blom M.P.K., Bragg J.G., Potter S., Moritz C. 2017. Accounting for Uncertainty in Gene Tree Estimation: Summary-Coalescent Species Tree Inference in a Challenging Radiation of Australian Lizards. Syst. Biol. 66:352–366.

Boyd B.M., Nguyen N.-P., Allen J.M., Waterhouse R.M., Vo K.B., Sweet A.D., Clayton D.H., Bush S.E., Shapiro M.D., Johnson K.P. 2022. Long-distance dispersal of pigeons and doves generated new ecological opportunities for host-switching and adaptive radiation by their parasites. Proc. Biol. Sci. 289:20220042.

Brinkmann H., van der Giezen M., Zhou Y., Poncelin de Raucourt G., Philippe H. 2005. An empirical assessment of long-branch attraction artefacts in deep eukaryotic phylogenomics. Syst. Biol. 54:743–757.

Bryant D., Hahn M.W. 2020. The Concatenation Question. Pages 3.4: 1–3.4: 23 in Phylogenetics in the Genomic Era (C. Scornavacca, F. Delsuc, and N. Galtier, eds.). .

Burbrink F.T., Grazziotin F.G., Pyron R.A., Cundall D., Donnellan S., Irish F., Keogh J.S.,
Kraus F., Murphy R.W., Noonan B., Raxworthy C.J., Ruane S., Lemmon A.R., Lemmon
E.M., Zaher H. 2020. Interrogating Genomic-Scale Data for Squamata (Lizards, Snakes, and
Amphisbaenians) Shows no Support for Key Traditional Morphological Relationships. Syst.
Biol. 69:502–520.

Catanach T.A., Halley M.R., Allen J.M., Johnson J.A., Thorstrom R., Palhano S., Thunder C.P.,
Gallardo J.C., Weckstein J.D. 2021. Systematics and conservation of an endemic radiation
of Accipiter hawks in the Caribbean islands. The Auk. 138:1–23.

Charlesworth B., Coyne J.A., Barton N.H. 1987. The Relative Rates of Evolution of Sex
Chromosomes and Autosomes. Am. Nat. 130:113–146.

Chen S., Zhou Y., Chen Y., Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 34:i884–i890.

Chernomor O., von Haeseler A., Minh B.Q. 2016. Terrace Aware Data Structure for
Phylogenomic Inference from Supermatrices. Syst. Biol. 65:997–1008.

Clements J.F., Rasmussen P.C., Schulenberg T.S., Iliff M.J., Fredericks T.A., Gerbracht J.A.,
Lepage D., Spencer A., Billerman S.M., Sullivan B.L., Wood C.L. 2023. The
eBird/Clements checklist of Birds of the World: v2023. .

Cloutier A., Sackton T.B., Grayson P., Clamp M., Baker A.J., Edwards S.V. 2019. Whole-
genome analyses resolve the phylogeny of flightless birds (Palaeognathae) in the presence of
an empirical anomaly zone. Syst. Biol. 68:937–955.

Doyle V.P., Young R.E., Naylor G.J.P., Brown J.M. 2015. Can We Identify Genes with
Increased Phylogenetic Reliability? Syst. Biol. 64:824–837.

Dunn C.W., Hejnol A., Matus D.Q., Pang K., Browne W.E., Smith S.A., Seaver E., Rouse G.W.,
Obst M., Edgecombe G.D., Sørensen M.V., Haddock S.H.D., Schmidt-Rhaesa A., Okusu
A., Kristensen R.M., Wheeler W.C., Martindale M.Q., Giribet G. 2008. Broad
phylogenomic sampling improves resolution of the animal tree of life. Nature. 452:745–749.

Edwards S.V., Cloutier A., Baker A.J. 2017. Conserved Nonexonic Elements: A Novel Class of
Marker for Phylogenomics. Syst. Biol. 66:1028–1044.

Edwards S.V., Xi Z., Janke A., Faircloth B.C., McCormack J.E., Glenn T.C., Zhong B., Wu S.,
Lemmon E.M., Lemmon A.R., Leaché A.D., Liu L., Davis C.C. 2016. Implementing and
testing the multispecies coalescent model: A valuable paradigm for phylogenomics. Mol.
Phylogenet. Evol. 94:447–462.

Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic
loci. Bioinformatics. 32:786–788.

Faircloth B.C., McCormack J.E., Crawford N.G., Harvey M.G., Brumfield R.T., Glenn T.C.
2012. Ultraconserved elements anchor thousands of genetic markers spanning multiple
evolutionary timescales. Syst. Biol. 61:717–726.

Faircloth B.C., Sorenson L., Santini F., Alfaro M.E. 2013. A Phylogenomic Perspective on the
Radiation of Ray-Finned Fishes Based upon Targeted Sequencing of Ultraconserved
Elements (UCEs). PLoS One. 8:e65923.

Felsenstein J. 1978. Cases in which Parsimony or Compatibility Methods will be Positively
Misleading. Syst. Biol. 27:401–410.

Fong J.J., Fujita M.K. 2011. Evaluating phylogenetic informativeness and data-type usage for
new protein-coding genes across Vertebrata. Mol. Phylogenet. Evol. 61:300–307.

Franz N.M., Musher L.J., Brown J.W., Yu S., Ludäscher B. 2019. Verbalizing phylogenomic
conflict: Representation of node congruence across competing reconstructions of the
neoavian explosion. PLoS Comput. Biol. 15:e1006493.

Gatesy J., Springer M.S. 2014. Phylogenetic analysis at deep timescales: unreliable gene trees,
bypassed hidden support, and the coalescence/concatalescence conundrum. Mol.
Phylogenet. Evol. 80:231–266.

Gueuning M., Frey J.E., Praz C. 2020. Ultraconserved yet informative for species delimitation:
Ultraconserved elements resolve long-standing systematic enigma in Central European bees.
Mol. Ecol. 29:4203–4220.

Harris R.B., Carling M.D., Lovette I.J. 2014. The influence of sampling design on species tree
inference: a new relationship for the New World chickadees (Aves: Poecile). Evolution.
68:501–513.

Hoang D.T., Chernomor O., von Haeseler A., Minh B.Q., Vinh L.S. 2018. UFBoot2: Improving
the Ultrafast Bootstrap Approximation. Mol. Biol. Evol. 35:518–522.

Hoff K.J., Lomsadze A., Borodovsky M., Stanke M. 2019. Whole-Genome Annotation with
BRAKER. Methods Mol. Biol. 1962:65–95.

Hosner P.A., Faircloth B.C., Glenn T.C., Braun E.L., Kimball R.T. 2016. Avoiding Missing Data
Biases in Phylogenomic Inference: An Empirical Study in the Landfowl (Aves:
Galliformes). Mol. Biol. Evol. 33:1110–1125.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C., Nabholz
B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li H., Zhou L.,
Narula N., Liu L., Ganapathy G., Boussau B., Bayzid M.S., Zavidovych V., Subramanian S.,
Gabaldón T., Capella-Gutiérrez S., Huerta-Cepas J., Rekepalli B., Munch K., Schierup M.,
Lindow B., Warren W.C., Ray D., Green R.E., Bruford M.W., Zhan X., Dixon A., Li S., Li
N., Huang Y., Derryberry E.P., Bertelsen M.F., Sheldon F.H., Brumfield R.T., Mello C.V.,
Lovell P.V., Wirthlin M., Schneider M.P.C., Prosdocimi F., Samaniego J.A., Vargas
Velazquez A.M., Alfaro-Núñez A., Campos P.F., Petersen B., Sicheritz-Ponten T., Pas A.,
Bailey T., Scofield P., Bunce M., Lambert D.M., Zhou Q., Perelman P., Driskell A.C.,
Shapiro B., Xiong Z., Zeng Y., Liu S., Li Z., Liu B., Wu K., Xiao J., Yinqi X., Zheng Q.,
Zhang Y., Yang H., Wang J., Smeds L., Rheindt F.E., Braun M., Fjeldsa J., Orlando L.,
Barker F.K., Jønsson K.A., Johnson W., Koepfli K.-P., O'Brien S., Haussler D., Ryder

O.A., Rahbek C., Willerslev E., Graves G.R., Glenn T.C., McCormack J., Burt D., Ellegren H., Alström P., Edwards S.V., Stamatakis A., Mindell D.P., Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert M.T.P., Zhang G. 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. Science. 346:1320–1331.

Johnson K.P. 2019. Putting the genome in insect phylogenomics. Curr Opin Insect Sci. 36:111–117.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods. 14:587–589.

Kapli P., Flouri T., Telford M.J. 2021. Systematic errors in phylogenetic trees. Curr. Biol. 31:R59–R64.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Korneliussen T.S., Albrechtsen A., Nielsen R. 2014. ANGSD: Analysis of Next Generation Sequencing Data. BMC Bioinformatics. 15:356.

Kuang T., Tornabene L., Li J., Jiang J., Chakrabarty P., Sparks J.S., Naylor G.J.P., Li C. 2018. Phylogenomic analysis on the exceptionally diverse fish clade Gobioidei (Actinopterygii: Gobiiformes) and data-filtering based on molecular clocklikeness. Mol. Phylogenet. Evol. 128:192–202.

Leaché A.D., Chavez A.S., Jones L.N., Grummer J.A., Gottscho A.D., Linkem C.W. 2015. Phylogenomics of phrynosomatid lizards: conflicting signals from sequence capture versus restriction site associated DNA sequencing. Genome Biol. Evol. 7:706–719.

Leite R.N., Kimball R.T., Braun E.L., Derryberry E.P., Hosner P.A., Derryberry G.E., Anciães M., McKay J.S., Aleixo A., Ribas C.C., Brumfield R.T., Cracraft J. 2021. Phylogenomics of manakins (Aves: Pipridae) using alternative locus filtering strategies based on informativeness. Mol. Phylogenet. Evol. 155:107013.

Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored hybrid enrichment for massively high-throughput phylogenomics. Syst. Biol. 61:727–744.

Lescroart J., Bonilla-Sánchez A., Napolitano C., Buitrago-Torres D.L., Ramírez-Chaves H.E., Pulido-Santacruz P., Murphy W.J., Svardal H., Eizirik E. 2023. Extensive Phylogenomic Discordance and the Complex Evolutionary History of the Neotropical Cat Genus Leopardus. Mol. Biol. Evol. 40.

Li G., Figueiró H.V., Eizirik E., Murphy W.J. 2019. Recombination-Aware Phylogenomics Reveals the Structured Genomic Landscape of Hybridizing Cat Species. Mol. Biol. Evol. 36:2111–2126.

Li H., Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 25:1754–1760.

Li H., Handsaker B., Wysoker A., Fennell T., Ruan J., Homer N., Marth G., Abecasis G., Durbin R., 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 25:2078–2079.

Li Q., Chen D., Wang S. 2023. Character displacement of egg colors during tinamou speciation. Evolution. 77:1874–1881.

Maddison W.P. 1997. Gene Trees in Species Trees. Syst. Biol. 46:523–536.

McCormack J.E., Faircloth B.C., Crawford N.G., Gowaty P.A., Brumfield R.T., Glenn T.C. 2012. Ultraconserved elements are novel phylogenomic markers that resolve placental mammal phylogeny when combined with species-tree analysis. Genome Res. 22:746–754.

McCormack J.E., Harvey M.G., Faircloth B.C., Crawford N.G., Glenn T.C., Brumfield R.T. 2013. A phylogeny of birds based on over 1,500 loci collected by target enrichment and high-throughput sequencing. PLoS One. 8:e54848.

Mclean B.S., Bell K.C., Allen J.M., Helgen K.M., Cook J.A. 2019. Impacts of Inference Method and Data set Filtering on Phylogenomic Resolution in a Rapid Radiation of Ground Squirrels (Xerinae: Marmotini). Syst. Biol. 68:298–316.

Meiklejohn K.A., Faircloth B.C., Glenn T.C., Kimball R.T., Braun E.L. 2016. Analysis of a Rapid Evolutionary Radiation Using Ultraconserved Elements: Evidence for a Bias in Some Multispecies Coalescent Methods. Syst. Biol. 65:612–627.

Meisel R.P., Connallon T. 2013. The faster-X effect: integrating theory and data. Trends Genet. 29:537–544.

Mendes F.K., Hahn M.W. 2018. Why Concatenation Fails Near the Anomaly Zone. Syst. Biol. 67:158–169.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A., Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mirarab S., Reaz R., Bayzid M.S., Zimmermann T., Swenson M.S., Warnow T. 2014. ASTRAL: genome-scale coalescent-based species tree estimation. Bioinformatics. 30:i541–8.

Moreira L.R., Klicka J., Smith B.T. 2023. Demography and linked selection interact to shape the genomic landscape of codistributed woodpeckers during the Ice Age. Mol. Ecol. 32:1739–1759.

Muggeo V.M.R., Atkins D.C., Gallop R.J., Dimidjian S. 2014. Segmented mixed models with random changepoints: a maximum likelihood approach with application to treatment for depression study. Stat. Modelling. 14:293–313.

Musher L.J., Cracraft J. 2018. Phylogenomics and species delimitation of a complex radiation of Neotropical suboscine birds (Pachyramphus). Mol. Phylogenet. Evol. 118:204–221.

Musher L.J., Del-Rio G., Marcondes R.S., Brumfield R.T., Bravo G.A., Thom G. 2023.
Geogenomic predictors of genetree heterogeneity explain phylogeographic and introgression
history: a case study in an Amazonian bird (Thamnophilus aethiops). Syst. Biol.

Musher L.J., Ferreira M., Auerbach A.L., Cracraft J. 2019. Why is Amazonia a "source"of
biodiversity? Climate-mediated dispersal and synchronous speciation across the Andes in an
avian group (Tityrinae). Proc. Roy. Soc. B.

Ostrow E.N., Catanach T.A., Bates J.M., Aleixo A., Weckstein J.D. 2023. Phylogenomic
analysis confirms the relationships among toucans, toucan-barbets, and New World barbets
but reveals paraphyly of Selenidera toucanets and evidence for mitonuclear discordance.
Ornithology. 140:ukad022.

Pamilo P., Nei M. 1988. Relationships between gene trees and species trees. Mol. Biol. Evol.
5:568–583.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Lemmon E.M., Lemmon A.R.
2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA
sequencing. Nature. 526:569–573.

Ranwez V., Douzery E.J.P., Cambon C., Chantret N., Delsuc F. 2018. MACSE v2: Toolkit for
the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. Mol.
Biol. Evol. 35:2582–2584.

R Core Team. 2023. R: A language and environment for statistical computing. Foundation for
Statistical Computing, Vienna, Austria.: .

Reddy S., Kimball R.T., Pandey A., Hosner P.A., Braun M.J., Hackett S.J., Han K.-L.,
Harshman J., Huddleston C.J., Kingston S., Marks B.D., Miglia K.J., Moore W.S., Sheldon
F.H., Witt C.C., Yuri T., Braun E.L. 2017. Why Do Phylogenomic Data Sets Yield
Conflicting Trees? Data Type Influences the Avian Tree of Life more than Taxon Sampling.
Syst. Biol. 66:857–879.

Rhodes J.A., Baños H., Mitchell J.D., Allman E.S. 2021. MSCquartets 1.0: quartet methods for
species trees and networks under the multispecies coalescent model in R. Bioinformatics.
37:1766–1768.

Robinson D.F., Foulds L.R. 1981. Comparison of phylogenetic trees. Math. Biosci. 53:131–147.

Roch S., Warnow T. 2015. On the Robustness to Gene Tree Estimation Error (or lack thereof) of
Coalescent-Based Species Tree Methods. Syst. Biol. 64:663–676.

Rodríguez-Ezpeleta N., Brinkmann H., Roure B., Lartillot N., Lang B.F., Philippe H. 2007.
Detecting and overcoming systematic errors in genome-scale phylogenies. Syst. Biol.
56:389–399.

Sackton T.B., Grayson P., Cloutier A., Hu Z., Liu J.S., Wheeler N.E., Gardner P.P., Clarke J.A.,
Baker A.J., Clamp M., Edwards S.V. 2019. Convergent regulatory evolution and loss of
flight in paleognathous birds. Science. 364:74–78.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.

Schultz D.T., Haddock S.H.D., Bredeson J.V., Green R.E., Simakov O., Rokhsar D.S. 2023.
    Ancient gene linkages support ctenophores as sister to other animals. Nature. 618:110–117.

Simão F.A., Waterhouse R.M., Ioannidis P., Kriventseva E.V., Zdobnov E.M. 2015. BUSCO:
    assessing genome assembly and annotation completeness with single-copy orthologs.
    Bioinformatics. 31:3210–3212.

Simion P., Philippe H., Baurain D., Jager M., Richter D.J., Di Franco A., Roure B., Satoh N.,
    Quéinnec É., Ereskovsky A., Lapébie P., Corre E., Delsuc F., King N., Wörheide G.,
    Manuel M. 2017. A Large and Consistent Phylogenomic Dataset Supports Sponges as the
    Sister Group to All Other Animals. Curr. Biol. 27:958–967.

Simmons M.P., Gatesy J. 2021. Collapsing dubiously resolved gene-tree branches in
    phylogenomic coalescent analyses. Mol. Phylogenet. Evol. 158:107092.

Smith B.T., Harvey M.G., Faircloth B.C., Glenn T.C., Brumfield R.T. 2014. Target capture and
    massively parallel sequencing of ultraconserved elements for comparative studies at shallow
    evolutionary time scales. Syst. Biol. 63:83–95.

Smith B.T., Mauck W.M., Benz B., Andersen M.J. 2018a. Uneven missing data skews
    phylogenomic relationships within the lories and lorikeets [Internet]. .

Smith B.T., Merwin J., Provost K.L., Thom G., Brumfield R.T., Ferreira M., Mauck W.M.,
    Moyle R.G., Wright T.F., Joseph L. 2023. Phylogenomic Analysis of the Parrots of the
    World Distinguishes Artifactual from Biological Sources of Gene Tree Discordance. Syst.
    Biol. 72:228–241.

Smith S.A., Brown J.W., Walker J.F. 2018b. So many genes, so little time: A practical approach
    to divergence-time estimation in the genomic era. PLoS One. 13:e0197433.

Springer M.S., Gatesy J. 2016. The gene tree delusion. Mol. Phylogenet. Evol. 94:1–33.

Stiller J., Feng S., Chowdhury A.-A., Rivas-González I., Duchêne D.A., Fang Q., Deng Y.,
    Kozlov A., Stamatakis A., Claramunt S., Nguyen J.M.T., Ho S.Y.W., Faircloth B.C., Haag
    J., Houde P., Cracraft J., Balaban M., Mai U., Chen G., Gao R., Zhou C., Xie Y., Huang Z.,
    Cao Z., Yan Z., Ogilvie H.A., Nakhleh L., Lindow B., Morel B., Fjeldså J., Hosner P.A., da
    Fonseca R.R., Petersen B., Tobias J.A., Székely T., Kennedy J.D., Reeve A.H., Liker A.,
    Stervander M., Antunes A., Tietze D.T., Bertelsen M.F., Lei F., Rahbek C., Graves G.R.,
    Schierup M.H., Warnow T., Braun E.L., Gilbert M.T.P., Jarvis E.D., Mirarab S., Zhang G.
    2024. Complexity of avian evolution revealed by family-level genomes. Nature.

Tajima F. 1983. Evolutionary relationship of DNA sequences in finite populations. Genetics.
    105:437–460.

Tan X., Qi J., Liu Z., Fan P., Liu G., Zhang L., Shen Y., Li J., Roos C., Zhou X., Li M. 2023.
    Phylogenomics Reveals High Levels of Incomplete Lineage Sorting at the Ancestral Nodes
    of the Macaque Radiation. Mol. Biol. Evol. 40.

Tea Y.-K., Xu X., DiBattista J.D., Lo N., Cowman P.F., Ho S.Y.W. 2021. Phylogenomic
Analysis of Concatenated Ultraconserved Elements Reveals the Recent Evolutionary
Radiation of the Fairy Wrasses (Teleostei: Labridae: Cirrhilabrus). Syst. Biol. 71:1–12.

Timilsena P.R., Wafula E.K., Barrett C.F., Ayyampalayam S., McNeal J.R., Rentsch J.D.,
McKain M.R., Heyduk K., Harkess A., Villegente M., Conran J.G., Illing N., Fogliani B.,
Ané C., Pires J.C., Davis J.I., Zomlefer W.B., Stevenson D.W., Graham S.W., Givnish T.J.,
Leebens-Mack J., dePamphilis C.W. 2022. Phylogenomic resolution of order- and family-
level monocot relationships using 602 single-copy nuclear genes and 1375 BUSCO genes.
Front. Plant Sci. 13:876779.

Valqui T. 2008. Phylogeogaphy of Nothoprocta Tinamous and the The Phylogeny of the
Tinamidae. Doctoral thesis. Louisiana State University.

Van Damme K., Cornetti L., Fields P.D., Ebert D. 2022. Whole-Genome Phylogenetic
Reconstruction as a Powerful Tool to Reveal Homoplasy and Ancient Rapid Radiation in
Waterflea Evolution. Syst. Biol. 71:777–787.

Waterhouse R.M., Seppey M., Simão F.A., Manni M., Ioannidis P., Klioutchnikov G.,
Kriventseva E.V., Zdobnov E.M. 2017. BUSCO applications from quality assessments to
gene prediction and phylogenomics. Mol. Biol. Evol.

Wen D., Yu Y., Nakhleh L. 2016. Bayesian Inference of Reticulate Phylogenies under the
Multispecies Network Coalescent. PLoS Genet. 12:e1006006.

Xi Z., Liu L., Davis C.C. 2015. Genes with minimal phylogenetic information are problematic
for coalescent analyses when gene tree estimation is biased. Mol. Phylogenet. Evol. 92:63–
71.

Zhang C., Mirarab S. 2022. Weighting by Gene Tree Uncertainty Improves Accuracy of Quartet-
based Species Trees. Mol. Biol. Evol. 39.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

Zhang F., Ding Y., Zhu C.-D., Zhou X., Orr M.C., Scheu S., Luan Y.-X. 2019. Phylogenomics
from low☐coverage whole☐genome sequencing. Methods Ecol. Evol. 10:507–517.

Zhao M., Kurtis S.M., White N.D., Moncrieff A.E., Leite R.N., Brumfield R.T., Braun E.L.,
Kimball R.T. 2023. Exploring Conflicts in Whole Genome Phylogenetics: A Case Study
Within Manakins (Aves: Pipridae). Syst. Biol. 72:161–178.

1079