Adenoviral Delivery of the CIITA Transgene Induces T-Cell-Mediated Killing in Glioblastoma
Organoids

Ilaria Salvato[1,2,3,*], Eliane Klein[1,3,*], Aurélie Poli[4], Mahsa Rezaeipour[1], Luca Ermini[1], Bakhtiyor
Nosirov[1,5], Anuja Lipsa[1], Anais Oudin[1], Virginie Baus[1], Gian Mario Dore[2], Antonio Cosma[6], Anna
Golebiewska[1], Simone P. Niclou[1,3,§], Antonio Marchini[2,7,8,§].

[1]NORLUX Neuro-Oncology Laboratory, Department of Cancer Research, Luxembourg Institute of
Health (LIH), Luxembourg;

[2]Laboratory of Oncolytic Virus Immuno-Therapeutics (LOVIT), Department of Cancer Research,
Luxembourg Institute of Health (LIH), Luxembourg;

[3]Department of Life Sciences and Medicine, Faculty of Science, Technology and Medicine (FSTM),
University of Luxembourg, Luxembourg;

[4]Neuro-Immunology Group, Department of Cancer Research, Luxembourg Institute of Health
(LIH), Luxembourg;

[5]Multiomics Data Science Research Group, Department of Cancer Research, Luxembourg Institute
of Health (LIH), Luxembourg;

[6]National Cytometry Platform, Translational Medicine Operation Hub, Luxembourg Institute of
Health, Esch-sur-Alzette, Luxembourg;

[7]Laboratory of Oncolytic Virus Immuno-Therapeutics, German Cancer Research Center,
Heidelberg, Germany.

[8]Current address: European Commission, Joint Research Centre (JRC), Geel, Belgium;

* Equal contribution

[§] Co-senior authors and corresponding authors

WORD COUNT (Excluding References): 10.989

ABSTRACT

The immunosuppressive nature of the microenvironment poses significant challenges in
developing effective immunotherapies against glioblastoma (GBM). Enhancing immune-mediated
responses is a prerequisite for a successful therapy. Here, we present a cancer gene therapy
approach utilizing a replication-deficient adenovirus (AdV) armed with the Class II Major
Histocompatibility Complex (MHC-II) Transactivator *CIITA* gene (Ad-CIITA) to induce antigen-
presenting properties in GBM cells. Successful induction of MHC-II molecules at the cell
membrane was achieved in infected GBM cell lines and primary human GBM organoids. Infection

with an AdV carrying a mutant form of *CIITA* resulted in cytoplasmic accumulation of CIITA

without subsequent MHC-II expression. Intriguingly, both wild-type and mutant Ad-CIITA

triggered prominent immune-mediated tumor cell death in a co-culture system with primary

human GBM organoids, suggesting a partially MHC-II-independent process. We further show that

the observed cancer cell killing requires the presence of T-cells and direct contact between GBM

and immune cells. Overall, these findings highlight that AdV-mediated *CIITA* delivery enhances T-

cell-mediated immunity against GBM, the precise mechanism of which remains to be elucidated.

KEYWORDS

Adenovirus/Gene therapy/Glioblastoma/MHC-II/Tumor Organoids

INTRODUCTION

Malignant brain tumors including glioblastoma (GBM) remain a major clinical and scientific

challenge. In GBM patients the median overall survival is limited to 12-15 months after tumor

diagnosis (Ostrom, Cioffi et al. 2021). The immunosuppressive tumor microenvironment (TME) as

well as systemic immunosuppression pose multiple challenges to novel immunotherapies. GBMs

recruit immunosuppressive cell types such as tumor-associated microglia and macrophages

(TAMs) and regulatory T-cells (Tregs), while only a limited number of T-cells are present, the latter

often displaying exhaustion markers that inhibit immune cell function (Quail and Joyce 2017,

Woroniecka, Chongsathidkiet et al. 2018, Pires-Afonso, Niclou and Michelucci 2020). GBM tumors

may also downregulate the expression of major histocompatibility complex (MHC) molecules,

surface glycoproteins that play a central role in presenting cancer-derived antigens to the immune

system (Burster, Gärtner et al. 2021).

Several immunotherapeutic approaches have been investigated for the treatment of GBM,

including immune checkpoint inhibitors, vaccines, adoptive cell therapy, as well as virus-based

therapy, utilizing oncolytic viruses and replication-deficient virus-mediated cancer gene therapy

(Agosti, Zeppieri et al. 2023). Despite promising results in preclinical studies, these treatments,

whether tested individually or in combination with standard care, have so far failed to translate in

clinically meaningful outcomes for GBM patients (Rainov 2000, Immonen, Vapalahti et al. 2004,

Weller, Butowski et al. 2017, Narita, Arakawa et al. 2019, Cloughesy, Brenner et al. 2020,

Cloughesy, Petrecca et al. 2020, Reardon, Brandes et al. 2020, Lim, Weller et al. 2022, Liau,

Ashkan et al. 2023, Omuro, Brandes et al. 2023). The discrepancy between clinical and preclinical

results can be largely attributed to the inadequacy of efficacy studies based on oversimplified

preclinical models not recapitulating the complex biology of GBM tumors.

Replication-deficient adenoviruses (AdVs) have demonstrated safety and tolerability as gene therapy and vaccination vectors, e.g. in the treatment of various cancer types, including GBM. Current GBM treatments using AdV derived vectors focus on the delivery of herpes simplex virus (HSV) thymidine kinase (TK), p*53*, interleukin (IL)-12, interferon (IFN)-β, and a chimeric death receptor (VB-111) (Banerjee, Nunez et al. 2021).

A promising approach is to arm non-lytic AdV vectors with novel therapeutic transgenes, able to boost immune responses against GBM. Class II Major Histocompatibility Complex Transactivator (CIITA) is the master regulator of MHC class II (MHC-II) molecules, expression of which is largely limited to professional antigen presenting cells (APCs) (Forlani, Shallak et al. 2023). Upon expression and translocation to the nucleus, CIITA acts as a non-DNA binding transcriptional coactivator by interacting with the enhanceosome complex. This complex consists of several constitutively-expressed DNA-bound factors preassembled at MHC-II promoters (Leon Machado and Steimle 2021). CIITA then nucleates various transcription factors and chromatin modifiers necessary to activate the transcription of the cell surface "classical" human leukocyte antigen (HLA)-DR, HLA-DP and HLA-DQ glycoproteins, which physically present antigens to CD4[+] T helper cells, as well as HLA-DM, HLA-DO and invariant chain (Ii) molecules, which are involved in intracellular antigen processing (Leon Machado and Steimle 2021). Previous work on carcinoma cell lines (Sartoris, Valle et al. 1998, Hillman, Kallinteris et al. 2003, Mortara, Castellani et al. 2006, Thompson, Dissanayake et al. 2006, Palma, Duangkhae et al. 2017), and more recently on murine glioma cells (Celesti, Gatta et al. 2023), found that ectopic stable expression of CIITA can transform tumor cells into MHC-II-positive cells with enhanced antigen processing and presentation capabilities. In these models, CIITA-transfected/infected MHC-II-positive cells were capable of priming and activating CD4[+] T helper cells, resulting in effective tumor rejection or growth retardation in mice with subcutaneous tumors (Hillman, Kallinteris et al. 2003, Meazza, Comes et al. 2003, Thompson, Dissanayake et al. 2006, Frangione, Mortara et al. 2010, Bou Nasser Eddine, Forlani et al. 2017, Palma, Duangkhae et al. 2017, Celesti, Gatta et al. 2023).

Here, we engineered a replication-deficient serotype 5 adenovirus (AdV) with the *CIITA* transgene ("Ad-CIITA") with the aim of converting infected primary human GBM organoids into "surrogate" APCs. The primary objective was to design a gene therapy that would enable the infected tumor cells to prime and activate T helper cells, thereby fostering antitumor immune responses. A second objective was to establish an advanced human GBM organoid system, based on primary organoids co-cultured with human immune cells, to assess the functional impact of the virus in a complex model system. We find that in our model system, AdV-mediated delivery of the *CIITA* transgene effectively induces cancer cell death in the presence of immune cells. While the exact

mechanism remains to be elucidated, our data suggest that the effect is at least partially

independent of MHC-II expression.

RESULTS

MHC-II expression in GBM is limited to non-neoplastic cells of the tumor microenvironment.

To assess the diversity of MHC-II molecules in GBM, we examined gene expression profiles at bulk

and single cell level in patient tumors and preclinical models. At bulk RNA level, expression of

CIITA and genes coding for MHC-II antigen presenting molecules (*HLA-DQA1*, *HLA-DQB1*, *HLA-*

DRA, HLA-DRB1, HLA-DRB5) was higher in GBM patient tumors than in normal brain tissue (**Fig.**

1A). However, analysis of single cell RNA-seq profiles of GBM patient tumors revealed that tumor

cells do not express *CIITA* nor genes encoding for MHC-II antigen processing (*CD74*, *HLA-DMA*,

HLA-DOA) and presentation (*HLA-DQA1*, *HLA-DQB1*, *HLA-DRA, HLA-DRB1, HLA-DRB5)* (**Fig. 1B**).

High expression of these genes was detected in immune cells of the tumor microenvironment,

explaining the high transcript levels observed by bulk RNA-seq. The highest MHC-II expression was

detected in well-known APCs, including dendritic cells, macrophages, microglia, and B cells (**Fig.**

1B). Limited cancer cell expression of MHC-II genes was confirmed in GBM patient-derived

orthotopic xenografts (PDOXs), where bulk RNA-seq profiles across 28 PDOXs showed low

expression of human genes involved in MHC-II antigen processing and presentation and

moderately high levels of enhanceosome-associated genes (**Fig. 1C**). Similar expression profiles

were seen in human GBM cell lines, including adherent and 3D sphere cultures of GBM stem-like

cells (GSCs) (**Supplementary Fig. 1A**). Only the U87 cell line presented high expression of certain

MHC-II associated genes (e.g., *CD74*, *HLA-DPA1*, *HLA-DRA*, *HLA-DRB4*, and *HLA-DRB5*).

Interestingly, the genes associated with the MHC-II enhanceosome complex (*CREB1*, *NFYA*, *NFYB*,

NFYC, *RFX5*, *RFXANK*, and *RFXAP)* showed sustained expression throughout patient tumors, PDOX

and cell lines (**Supplementary Fig. 1A**), suggesting that GBM cells have the capacity to express

MHC-II molecules upon activation of the MHC-II master regulator CIITA. Moreover, most of the

preclinical models exhibited transcriptional expression of the Coxsackie and Adenovirus Receptor

(*CXADR*), the specific receptor for AdV serotype 5, as well as the integrin αvβ5 necessary for AdV

infection (*ITGAV* and *ITGB5*) (Nemerow and Stewart 1999) (**Fig. 1C, Supplementary Fig. 1A**). This

observation implies that GBM tumor cells are likely to be susceptible to AdV infection, highlighting

the suitability of non-lytic AdVs for cancer gene therapy applications targeting GBM.

Next, we assessed the presence of MHC-II proteins at the cell surface of GBM cells by flow

cytometry. The majority of GBM cells showed no MHC-II protein at the cell surface under regular

culture conditions (**Fig. 1D, Supplementary Fig.1B**). Out of 10 GBM cell lines analyzed, only U87

and BG5 cells showed some positive cells (18.7% and 10.8% respectively). Interestingly, IFN-γ cytokine stimulus did not activate MHC-II expression in GBM cells, except for U87, BG5, and T16NS (**Fig. 1D**).

In summary, we confirmed that MHC-II molecules are largely absent from GBM tumor cells but are expressed in tumor-associated APCs. Based on the presence of transcripts of the MHC-II enhanceosome complex, ectopic expression of CIITA appears as a valid strategy to deliver MHC-II molecules at the cell membrane. Moreover, the expression of the necessary AdV receptor and cell adhesion molecules confers infectivity to GBM cells to serotype 5 AdVs.

Human GBM stem-like cells infected with Ad-CIITA express CIITA and present MHC-II molecules at the cell surface.

To convert MHC-II-negative GBM tumor cells into MHC-II-positive cells with APC-like properties, we generated a replication-deficient AdV carrying the wild-type *CIITA* gene (Ad-CIITA) (**Fig. 2A**). The serotype 5 AdV backbone is widely employed in cancer gene therapy for its ability to efficently transduce large transgenes (up to 8 Kbases) in a vast range of cell and tissue types, including GBM. Importantly, AdVs can infect both proliferating and quiescent cells and their DNA does not integrate into the host genome, but rather remains episomally in the host nucleus, precluding the risk of insertional mutagenesis (Bulcha, Wang et al. 2021). Three different AdV control vectors were generated, harboring either: i) no transgene (Ad-null); ii) a green fluorescent protein (GFP)-encoding transgene (Ad-GFP); or (iii) a mutated *CIITA* transgene (Ad-CIITA mutant) (**Fig. 2A**). The latter was newly generated in our laboratory, carrying a novel missense point mutation (guanine-to-adenosine) (**Supplementary Fig. 2A**) that translated into a full-length CIITA protein with a glycine (G) to arginine (A) substitution at amino acid position 1088 (**Supplementary Fig. 2B**). We found that this mutation impaired the translocation of the MHC-II transactivator to the cell nucleus (**Supplementary Fig. 2C**).

To determine optimal viral infection, we infected mouse glioma GL261 cells at various multiplicities of infection (MOIs) and assessed MHC-II expression via flow cytometry at 72 hours post-infection. The Ad-CIITA vector successfully delivered the human *CIITA* transgene into the murine cell line, leading to a dose-dependent increase in IA-IE (MHC-II molecules) levels at the cell membrane compared to non-infected controls (**Supplementary Fig. 2D**). Specifically, at a virus MOI of 50, GL261 tumor cells expressed the CIITA transactivator at both mRNA and protein levels (**Supplementary Fig. 2E**). Notably, the expression of IA-IE (MHC-II) molecules peaked at 72 hours post-infection, with nearly all tumor cells displaying them at the cell membrane (**Supplementary Fig. 2F**).

Next, we assessed AdV-mediated expression of MHC-II molecules in human 3D GSC lines (P3NS and T16NS) by infecting the cells with a virus MOI of 50. High *CIITA* mRNA (**Fig. 2B**) and protein (**Fig. 2C**) were detected at 72 hours post-infection with both Ad-CIITA and Ad-CIITA mutant viruses. Since only wild-type CIITA translocates to the nucleus (**Supplementary Fig. 2C**), HLA-DRα mRNA and protein were only detected in GBM cells infected with Ad-CIITA but not with Ad-CIITA mutant virus (**Fig. 2B-C**). Surface MHC-II molecules were confirmed by flow cytometry using a pan antibody targeting all surface MHC-II antigen presenting molecules (HLA-DR/DP/DQ epitopes) (**Fig. 2D**). As expected, GBM cells infected with Ad-CIITA mutant virus did not display MHC-II molecules at the cell surface (**Fig. 2D**). In summary, we show efficient infection of human GBM cells leading to MHC-II molecules at the cell membrane, a necessary requirement for functional antigen presentation. Moreover, a mutant CIITA construct has been generated which allows to effectively distinguish the effects of nuclear versus non-nuclear CIITA expression.

Infection with Ad-CIITA induces MHC-II expression in human primary GBM organoids.

Following the results in adherent and 3D GBM cell lines, we next determined the efficacy of Ad-CIITA virus in primary GBM organoids, obtained from mechanical dissociation of patient tissue, followed by serial transplantation as PDOXs (Golebiewska, Hau et al. 2020). Following our well-established protocol (Oudin, Baus et al. 2021), GBM organoids of uniform size can be cultured short term *ex vivo* without repeatable passaging *in vitro*, which allows for preservation of complex molecular features and a proliferative rate similar to *in vivo* tumors. We selected three GBM (P3, T16 and T188) representing tumors with classical genetic and epigenetic features, including gain of Chr7 and loss of Chr10 (**Table S1**).

To determine the optimal viral MOI, we infected P3 organoids with increasing MOI of Ad-CIITA for up to 7 days post-infection. An MOI between 25 and 100 led to strong ectopic expression of CIITA and HLA-DRα at 72 hours post-infection (**Fig. 3A**). GBM organoids infected with Ad-CIITA at MOI 75 exhibited a tissue structure comparable to that of non-infected GBM organoids, albeit with a somewhat lower cell density (**Fig. 3B**). Again, HLA-positive cells were only detected in Ad-CIITA-infected cells (**Fig. 3B** lower panel). Flow cytometry confirmed that Ad-CIITA infection successfully translated into expression of MHC-II molecules at the cell membrane (**Fig. 3C**).

To evaluate the cytotoxicity of virus infection in primary GBM cells, dsRed-expressing P3 organoids were infected with Ad-CIITA or control viruses (Ad-null and Ad-CIITA mutant) at increasing MOIs and monitored real-time with the Incucyte S3 live cell imaging system. Tumor cell viability was assessed by the dsRed fluorescence intensity signal over time. Consistent with previous studies showing that first-generation AdV vectors can lead to cell cycle arrest at elevated

MOIs (Zheng, Goldsmith et al. 2000), AdV infected organoids exhibited a dose-dependent reduction in growth rate compared to non-infected organoids (**Fig. 3D**). This effect appeared more evident with the wild-type Ad-CIITA vector (**Fig. 3D**). However, viral infection per se did not result in cell death, as deduced from the dsRed fluorescence signal (**Fig. 3D**) and the H&E stained sections (**Fig. 3B**). Similar results were seen in T16 GBM organoids, while in T188 organoids the proliferation rate did not seem to be affected by viral infection (**Supplementary Fig. 3**).

In conclusion, our AdV-CIITA constructs effectively infect primary GBM organoids with only a small impact on tumor growth. Robust MHC-II expression is obtained with Ad-CIITA, but not with the control Ad-null or Ad-CIITA mutant vectors.

Ad-CIITA triggers immune-mediated GBM tumor cell death primarily mediated by CD3$^+$ T-cells.

To evaluate the potential of CIITA-expressing human GBM to elicit adaptive immune responses, we established a protocol involving the co-culture of virally-infected primary GBM organoids with allogenic peripheral blood mononuclear cells (PBMCs). PBMCs were obtained from healthy donors or from partially HLA-matched donors (**Fig. 4A**). For efficient monitoring and quantification of viability, we employed fluorescently-labeled GBM organoids, either through dsRed expression (P3 and T16) or labeling with cytolight rapid red dye (T188). Prior to adding immune cells, GBM organoids were infected with Ad-CIITA for 72 hours, at MOI 75 (P3 and T188) or MOI 25 (T16) to allow adequate MHC-II surface expression. Control GBM organoids were left uninfected or infected with the control viruses Ad-null or Ad-CIITA mutant. Ad-null-infected and non-infected GBM organoids enabled assessment of viral backbone intrinsic toxicity and immunogenicity in co-culture conditions. GBM organoids were then co-cultured with non-labeled PBMCs or purified individual immune cell populations of interest (CD3$^+$, CD4$^+$, CD8$^+$ T-cells) at different target:effector (T:E) ratios (1:1, 1:5, 1:10) for up to 6 days. The purity of PBMCs and isolated cell types were assessed by flow cytometry before co-culture (**Supplementary Fig. 4A, Table S2**). Co-cultures were continuously monitored using the Incucyte S3 live cell imaging system (**Fig. 4A**) and tumor cell death was monitored by tracking the decline in dsRed signal over time.

Based on previous tumor-immune cell co-culture studies (Zhou, Lieshout et al. 2022), we decided to first use immune cells from healthy human donors for our co-culture setup. In the absence of immune cells, P3 GBM organoids retained their structure, their proliferation being slightly affected by viral infection (Ad-null) and further by CIITA expression (Ad-CIITA) (**Fig. 4B**), in line with the experiments without co-culture conditions shown in **Fig. 3D**. Upon the addition of allogenic PBMCs, we observed a pronounced organoid disruption and tumor cell killing in Ad-CIITA-infected organoids (86,5%±10,2), contrasting with a lower effect in Ad-null-infected

organoids (42,2%±7,3) (**Fig. 4B, Video S1**). Throughout the text, the reported percentages in parentheses represent a reduction in dsRed signal compared to non-infected controls at the end-point of the co-culture. These findings imply that the induction of PBMC killing capacity is primarely associated with the expression of CIITA by the tumor cells rather than solely through the acquired immunogenicity linked to AdV infection itself.

To further investigate the association between the observed killing and MHC-II surface expression, we conducted co-cultures using allogenic PBMCs and organoids infected with the Ad-CIITA mutant virus. Interestingly, the Ad-CIITA mutant also led to robust tumor cell killing in the presence of PBMCs (78,3%±21,7) (**Fig 4B, Video S1**). This was unexpected as we have shown that this construct did not induce MHC-II molecules expression (**Fig. 2B-C**). Similar outcomes were obtained in T16 GBM organoids at different MOIs (25 and 75), where no cell death was observed in uninfected or Ad-null-infected T16 organoids. Pronounced tumor cell death was induced by Ad-CIITA and again, though to a lesser extent, by Ad-CIITA mutant (**Supplementary Fig. 4B**). These data suggest that CIITA expression potentiates the immunogenicity of GBM organoids for alloreactive PMBCs, potentially mediated by a cumulative effect of MHC-II molecules-dependent and independent mechanisms.

We next sought to determine whether the observed killing phenotype was orchestrated by CD3$^+$ T-cells. Similar to PBMCs, purified CD3$^+$ T-cells alone efficiently induced tumor cell killing in P3 organoids infected with Ad-CIITA (76,4%±21,9) or Ad-CIITA mutant (75,1%±29,4), suggesting a primary T-cells involvement (**Fig. 4C vs 4B**). An additional effect from other immune cells cannot be completely ruled out. To assess the specificity of Ad-CIITA induced T-cell killing, we performed co-culture experiments with purified CD4$^+$ and CD8$^+$ T-cells, which rely on MHC-II and MHC-I molecules, respectively. Interestingly, efficient tumor cell killing was maintained when GBM organoids were co-cultured solely with either CD4$^+$ T-cells (Ad-CIITA: 75,6%±16,8) or CD8$^+$ T-cells (Ad-CIITA: 67,5%±28,8) (**Fig. 4C**). Similar results were obtained with T16 and T188 organoids (**Supplementary Fig. 4C**), reinforcing the involvement of a MHC-II independent mechanism in GBM cell killing.

To further delineate the role of MHC-II molecules in the observed increased immunogenicity mediated by Ad-CIITA and Ad-CIITA mutant for alloreative T-cells, we set up a co-culture system with effector cells sharing a partial match at HLA alleles with the target GBM organoids. Specifically, PBMCs were chosen to match the HLA typing of the donor GBM tumors at least at one allele of each MHC class I (MHC-I) (*HLA-A*, *HLA-B*, *HLA-C*) and each MHC-II (*HLA-DRB1/3/4/5*, *HLA-DQA1*, *HLA-DQB1*, *HLA-DPA1*, *HLA-DPB1*) locus. The matching degree varied depending on the donor, as presented in **Table S3**. The Ad-CIITA-mediated killing phenotype was retained in the P3 co-culture model (**Fig. 4D)** and confirmed in T16 tumors (**Supplementary Fig. 4D**). Although the

cytotoxic activity of HLA-matched PBMCs and isolated CD3$^+$ cells was reduced compared to

allogenic PBMC counterpart (**Fig. 4C vs Fig. 4D**), these data indicate that the main killing activity

observed could not be solely attributed to an alloreactive immune reaction linked to mismatching

MHC molecules.

In summary, our findings demonstrate that both Ad-CIITA and Ad-CIITA mutant enhance the

immunogenicity of GBM for allogenic T-cells, facilitating efficient tumor cell killing in the presence

of CD3+ T-cells. These data suggest that CIITA activates a T-cell mediated immune response that is

at least partially independent of MHC-II antigen presentation.

Immune-mediated tumor cell killing of GBM organoids requires CIITA expression and cell-to-cell

contact with T-cells.

To address the transgene specificity of the observed phenotype, we used an AdV carrying the *GFP*

transgene as a control. GFP expression was detected in P3 GBM organoids via live cell imaging at

24, 48, and 72 hours post-infection (**Fig. 5A**). Unlike Ad-CIITA, Ad-GFP did not lead to tumor cell

death upon co-culture with PBMCs (**Fig. 5B**). The same results were obtained in the T16 co-culture

model (**Supplementary Fig. 5A-B**), thus confirming the selectivity of the immune-mediated

phenotype toward CIITA.

Next, we aimed to determine whether the immune-mediated tumor killing required cell-to-cell

contact. We performed co-cultures using a two-layer transwell system, allowing for physical

separation of GBM organoids and immune cells, while maintaining the crosstalk via soluble

chemical mediators. In this setup, we observed the expected reduction in organoid growth due to

AdV infection, however organoid breakdown in the presence of Ad-CIITA and Ad-CIITA mutant

was completely abolished (**Fig. 5C**). This was replicated in T16 organoids (**Supplementary Fig. 5C**),

indicating that immune cells require direct cell-to-cell interaction with target GBM cells to

effectively induce tumor cell death.

To investigate whether AdV infection triggered other APC-like features in addition to MHC-II

molecules in infected GBM organoids, we conducted flow cytometry analysis of cell membrane

expression of costimulatory molecules CD80 and CD86, along with MHC-I. P3 GBM organoids

displayed positive MHC-I expression, which remained unaltered by AdV infection or CIITA

expression. Other surface co-stimulators were not expressed in GBM organoids either at baseline

or upon AdV /Adv-CIITA infection (**Supplementary Fig. 5D**).

We next aimed to assess different modes of contact-dependent T-cell-mediated cell killing.

Cytotoxic T-cells conventionally release lytic granules, such as perforin and granzymes, through

calcium-dependent exocytosis into the immunological synapse between cytotoxic T-cells and

target cells. However, degranulating T-cells did not appear to be involved in CIITA-mediated
killing, as evidenced by the absence of cell surface presentation of the degranulating marker
CD107a (**Fig. 5D**). Cell death can also be induced via activation of death receptor signaling in
target cells, including tumor necrosis factor alpha (TNF-α)/TNF-receptor (TNF-R) and Fas/Fas
ligand (FasL) (Barry and Bleackley 2002). We therefore selectively blocked these death pathways
via anti-human TNF-R and FasL monoclonal antibodies. Blocking either TNF- or Fas-mediated
signaling did not impact the ability of CIITA to induce the killing of GBM organoids (**Fig. 5E**), while
tumor cell killing induced by Actinomycin D and dimethyl sulfoxide (DMSO) was successfully
blocked by neutralizing TNF- and Fas-mediated signalling, respectively (**Supplementary Fig. 5E**).
As we did not detect classical T-cell activation mechanisms, we next investigated the differential
secretion of cytokines upon tumor cell killing mediated by Ad-CIITA (wild-type CIITA) (**Fig. 5F**).
Using conditioned media collected at 72 hours post co-culture, we detected increased secretion
of the extracellular matrix metalloproteinase inducer (EMMPRIN) and RANTES (Regulated upon
Activation, Normal T-cell Expressed, and Secreted) in Ad-CIITA co-culture medium. The secretion
of EMMPRIN, and consequently, the increased activity of metalloproteinases in degrading the
tumor extracellular matrix, might be linked with the observed CIITA-induced immune-mediated
organoid disruption preceeding tumor cell death. Moreover, compared to controls, the Ad-CIITA
co-culture medium exhibited decreased levels of IL-1alpha, IL-18, VEGF, and IL-6. This may
indicate a reduction of pro-tumorigenic, pro-angiogenic, and immunosuppressive potential of the
GBM tumor microenvironment upon Ad-CIITA infection. Whether these changes in cytokine levels
are functionally linked to tumor cell killing remains to be determined.

DISCUSSION

MHC-II molecules are the restricting elements of antigen presentation and CD4$^+$ T helper cell
activation. Their expression is usually confined to professional APCs, including dendritic cells, B
cells, macrophages, but they can be induced upon cytokine stimuli in other cell types, including
endothelial and epithelial cells (Drozina, Kohoutek et al. 2005). CIITA is the master regulator of
MHC-II expression, and this biological function can be exploited to stimulate adequate triggering
of the T helper arm of adaptive immunity against the tumor, a necessary requirement for effector
immune responses (Accolla, Ramia et al. 2019).
Given the need to develop new strategies against GBM tumors, we equipped a non-lytic AdV with
the wild-type *CIITA* transgene (Ad-CIITA). This proof-of-concept study employed a serotype 5
AdV, which could face reduced gene delivery efficiency due to pre-existing immunity to AdVs. For
in vivo application, various AdV vectors with genetic modifications could be developed to mitigate

immune responses and enhance gene delivery efficiency. At basal level, GBM tumor cells generally lacked MHC-II expression, while other components of the MHC-II transcriptional machinery were present. We show that Ad-CIITA infection of GBM cells resulted in robust MHC-II expression at the surface of infected GBM cells.

Initially planned as a control vector, we generated a novel *CIITA* mutant harboring a point mutation that encodes a full-length protein in which the glycine at position 1088 is replaced by an arginine. This mutation resides within the carboxy-terminal region of CIITA. The replacement of the non-polar, small amino acid glycine with arginine, a larger and positively charged amino acid, may lead to alterations in the overall structure and electrostatic interactions within the protein, impacting its folding, stability and binding properties (Molnár, Szakács and Tusnády 2016). Structurally, a fully functional MHC-II transactivator consists of an amino-terminal acidic domain, a proline/serine/threonine-rich region, a central GTP binding domain and a string of four leucine-rich repeats (LRRs), which is essential for CIITA nuclear localization, enhanceosome assembly and CIITA transactivator activity (Leon Machado and Steimle 2021). The positively charged mutation presented in this study was found to reside in the fourth LRR, spanning from amino acid position 1073 to 1093. This novel mutation impeded the translocation of the CIITA master regulator to the cell nucleus, thereby abrogating its role as a transcriptional coactivator. While a wild-type CIITA protein typically exhibits a rather balanced distribution between the nucleus and cytoplasm (roughly 50:50 ratio) (Leon Machado and Steimle 2021), the G1088A mutant protein was effectively excluded from the nucleus. A previous study showed that the carboxy-terminal region of CIITA is essential for its functionality as substitution of phenylalanine with serine at position 962 also abrogated MHC-II expression (Towey and Kelly 2002). Our novel point-mutated CIITA variant may thus effectively serve to unravel MHC-II-dependent and -independent activities of CIITA.

We here demonstrate that while Ad-CIITA infection has limited impact on tumor cell growth, effective killing of infected GBM organoids is activated and maintained only in the presence of immune cells. Our study reveals a new ability of CIITA to activate T-cells which is independent of MHC-II molecules and antigen presentation. Firstly, we find that the CIITA mutant, unable to translocate into the nucleus and therefore to induce MHC-II is still able to trigger a T-cell mediated GBM tumor cell death. Secondly, in our model, CIITA wild type is also able to activate a MHC-II independent immunity, as depletion of CD4[+] cells only partly impaired the ability of the protein to activate immune cell death. A direct influence of the viral infection could be ruled out since control empty and GFP-containing vectors had a significantly lesser effect on T-cell activation or tumor cell death induction. Our findings indicate that the tumor killing phenotype relies on direct cell-to-cell interaction between target GBM and immune cells. Interestingly, we

did not detect activation of canonical T-cell granule exocytosis or of known death signaling

pathways like Fas/FasL or TNF-α/TNF-R. The mechanism by which CIITA induces a MHC-II

independent immune-mediated tumor cell death remains to be characterized.

We show that the cytotoxicity induced by both Ad-CIITA and Ad-CIITA mutant in patient-derived

GBM cells was primarily mediated by CD3$^+$ T-cells, which is in line with *in vivo* results in mice of

CIITA transfection in the TS/A murine mammary adenocarcinoma model (Meazza, Comes et al.

2003, Frangione, Mortara et al. 2010) and GL261 glioma model (Celesti, Gatta et al. 2023).

Nevertheless, we cannot currently exclude the involvement of other immune components.

Further investigation should be directed in validating the anticancer role of cytoplasmic CIITA in

humanized GBM animal models, by examining whether T-cell-mediated immune recognition can

lead to tumor rejection or growth retardation following orthotopic injection of Ad-CIITA mutant-

infected GBM cells.

It has already been shown that CIITA not only governs the transcription of MHC-II genes but also

modulates the expression of other immune-related target genes, such as MHC-I, Plexin-A1, and

collagen type I α2, among others (Leon Machado and Steimle 2021). Our findings suggest that

CIITA may have additional roles in the cytoplasm, next to its well-established nuclear

transcriptional activation function. Previous research has described cytoplasmic effects of CIITA,

such as the L1035P CIITA mutant enhancing human immunodeficiency virion maturation through

Gag processing by the viral protease (Porter, Kelley et al. 2010).

T-cell activation and proliferation commonly require at least two key events: (i) T-cell receptor

(TCR) binding to a specific MHC-restricted antigen, and (ii) the interaction of costimulatory

molecules on the surface of T-cells and APCs. We have shown that the killing phenotype relies on

direct cell-to-cell interaction between the target tumor organoids and the immune cells and was

observed despite the absence of CD80 and CD86 costimulatory molecules, which is in line with

previous reports (Meazza, Comes et al. 2003, Mortara, Castellani et al. 2006, Frangione, Mortara

et al. 2010).

A protein cytokine array revealed increased secretion of EMMPRIN and RANTES along with a

decrease in various pro-inflammatory cytokines in Ad-CIITA co-culture medium. EMPPRIN is a

glycoprotein expressed and secreted by various cell types, including lymphocytes and tumor cells,

and is known to play a role in metalloproteinase induction, energy metabolism, T-cell activation,

co-stimulation, and proliferation (Kaushik, Yong et al. 2016). RANTES, also known as CCL5, is a

potent proinflammatory chemokine that acts either as a potent chemoattractant or as a mitogen-

like immune cell activator (Appay, Dunbar et al. 2000). RANTES-induced T-cell activation has

various effects, including lymphocyte proliferation or apoptotic cell death, as well as the release

of pro-inflammatory cytokines (Appay, Dunbar et al. 2000). Further studies are warranted to gain

a comprehensive understanding of the implications of soluble EMMPRIN and RANTES in the

observed T-cell-mediated killing phenotype.

In conclusion, our findings highlight the potential of viral delivery of the *CIITA* transgene in human

GBM as a promising immunotherapeutic approach for enhancing adaptive responses against

GBM. The observed T-cell-mediated cytotoxicity, partially independent of MHC-II antigen

presentation, opens up new avenues for further investigation into CIITA function beyond its

canonical nuclear transcriptional activation and MHC-II antigen presentation. Further elucidation

of the mechanism of T-cell-mediated tumor cell death is needed to understand the specific effects

induced by CIITA in the tumor cell cytoplasm. Additionally, it would be interesting to explore the

involvement of other immune cell subsets, such as natural killer cells and macrophages, in CIITA-

induced immune responses and to study potential immunosuppressive molecules induced by

CIITA treatment, as prolonged MHC-II molecules expression is commonly associated with

increased PD-1 expression (Chih, Sahm et al. 2021).

MATHERIALS AND METHODS

Patient-derived preclinical models and cell culture.

Human GBM patient tumor tissues were provided by the Department of Neurosurgery of the

Centre Hospitalier de Luxembourg, Luxembourg and the Neurosurgical Department of Haukeland

University Hospital, Bergen, Norway. Patient provided their informed consent, and the study was

accepted by the local ethical committees (the National Committee for Ethics in Research (CNER)

Luxembourg, under the protocol number 201201/06, and the Haukeland University Hospital,

under the protocol number 2009/117). PDOX models were generated and maintained by cycles of

short-term organoid cultures and orthotopic xenografting as previously described (Oudin, Baus et

al. 2021) (Golebiewska, Hau et al. 2020).

Patient-derived models expressing dsRed (P3dsRed and T16dsRed) were created from the original

PDOXs by stable transfection of a dsRed-expressing lentiviral vector (a three-vector system:

transfer vector pRRLsin-PPThCMV, helper plasmids pCMVdeltaR8.74, and VSV-G pMD.G.2) in GBM

organoids *ex vivo* as previously described (Niclou, Danzeisen et al. 2008). To perform *ex vivo*

experiments in GBM organoids, tumor cells were purified with use of previously published

protocol, which includes MACS-based purification of GBM cells from enzymatically-digested PDOX

tumor tissue and subsequent recreation of uniform organoids *ex vivo* (Oudin, Baus et al. 2021).

PDOX-derived organoids were cultured in Dulbecco's modified Eagle's medium (DMEM)

containing 4.5 g/L glucose w/o glutamine supplemented with 10% fetal bovine serum (FBS), 2 mM

UltraGlutamine, 0.4 mM nonessential amino acids (NEAAs) and 100 U/mL penicillin–streptomycin

("Organoid medium").

Human GBM stem-like cell (GSC) lines, BG5, BG7, GG6 and GG16, and adherent GBM cell lines

U87 and U251 were kindly provided by Prof. Rolf Bjerkvig from University of Bergen. GSC lines

NCH421, NCH644 and NCH465 were kindly provide by Prof. Christel Herold-Mende from

University of Heidelberg. Human GSC lines P3NS and T16NS were generated from PDOX models as

previously described (Golebiewska, Hau et al. 2020). GSC lines were cultured as 3D spheres in

serum-free Neurobasal medium supplemented with 1X B-27™ Supplement, 2 mM UltraGlutamine,

0,1 U/mL heparin, 20 ng/mL bFGF, 20 ng/mL EGF and 100 U/mL penicillin–streptomycin ("NCS

medium"). Human embryonic kidney 293 T (293T) cells were obtained from ATCC. Murine GL261

glioma were kindly provided by Prof. Roberto Accolla from University of Insubria. All cell lines

were grown in DMEM high glucose, GlutaMAX™ supplemented with 10% FBS and 100 U/mL

penicillin–streptomycin ("DMEM medium"). All cells were maintained at 37°C in a 5% CO^2

atmosphere and 90% humidity and regularly tested for mycoplasma.

HLA typing.

Genomic DNA was extracted from GBM organoids with the commercially available DNeasy Blood

& Tissue Kit (Qiagen) according to the manufacturer's instructions. A total of 1200 ng was used for

ultrahigh resolution six-digits sequencing at 11 HLA loci (MHC-I *HLA-A*, *-B*, *-C*; MHC-II *HLA-*

DRB1/3/4/5, *-DQA1*, *-DQB1*, *-DPA1*, *-DPB1*) using the Illumina TruSight HLA v2 Sequencing panel

(**Table S3**).

Gene expression analysis.

The expression of MHC-II associated genes in GBM patient tumors and control normal brain tissue

was investigated with the GEPIA2 tool (Tang, Kang et al. 2019), which allows for exploration of

large-scale gene expression datasets from The Cancer Genome Atlas (TCGA) and the Genotype-

Tissue Expression (GTEx) projects. To examine gene expression in single cells isolated from GBM

patient tumors, we exploited the GBmap dataset (Ruiz-Moreno, Salas et al. 2022). The DotPlot

function in the Seurat package for R was applied to visualize gene expression changes in selected

cell populations. Gene expression in preclinical models was assessed with the Gene-Chip® Human

Gene 1.0ST Arrays (Affimetrix) as described previously (Golebiewska, Hau et al. 2020) or RNA-

sequencing (RNA-seq) on a bulk tissue. For bulk RNA-seq, total RNA was isolated from

cryopreserved PDOX tumor tissue using direct-zol RNA miniprep kit (Zymo Research). RNA quality

was checked with the HS RNA (15 nt) Kit (Agilent). mRNA-seq profiling was carried out with 50

million reads/sample, 2x150bp, using the Illumina Stranded mRNA prep kit on an Illumina

NovaSeq 6000 sequencer. RNA-seq quality control was performed using FastQC. Adaptors were

trimmed from raw sequencing reads using Cutadapt version 3.7 (Martin 2011), and the trimmed

reads were aligned to both the human reference genome GRCh38 and the mouse reference

genome GRCm39 by STAR 2.7.10b aligner (Dobin, Davis et al. 2013). Deconvolution of mouse and

human sequencing reads was performed with XenofilterR (Kluin, Kemper et al. 2018), which

separates mouse from human sequence reads. Uniquely human-mapped reads were ultimately

used to quantify gene expression using FeatureCounts (Liao, Smyth and Shi 2014). The R package

DGEobj.utils was used to normalize gene expression controlling for both library size and the gene

lengths, by converting raw counts to TPM (transcripts per million). Heatmap was generated using

the R package Pheatmap. All analyses were performed with R version 4.2.0.

Peripheral blood lymphocytes.

Allogeneic PBMCs were obtained from healthy human donors from the Red Cross of Luxembourg,

without controlling for HLA genotype. Allogeneic HLA-matched PBMCs were purchased from

Immunospot based on their HLA matching at MHC-I and MHC-II loci with the GBM organoids

(**Table S3**). The PBMC fraction was isolated from peripheral blood by density gradient

centrifugation and cryopreserved until later use. Cryopreserved human PBMCs were thawed at

37°C for 10-15 seconds in RPMI 1640 Medium, GlutaMAX™ supplemented with 9% FBS, 1%

human serum, 1 mM Sodium Pyruvate, 55 μM 2-Mercaptoethanol, 0,1 mM NEAA, 5 mM HEPES

and 100 U/mL penicillin-streptomycin ("T-cell medium"). After thawing, cells were incubated for

10 minutes with 25 U/mL benzonase (Sigma–Aldrich). Allogeneic HLA-matched PBMCs were

instead thawed, according to Immunospot's protocol.

To establish co-cultures involving specific T-cell populations, we first isolated human CD3[+] T-cells

from PBMCs using a CD3[+] T-cell negative isolation kit (Biolegend), according to the manufacturer's

instructions. Subsequently, we further separated CD4[+] and CD8[+] T-cell subsets from the

negatively-selected CD3[+] T-cells using the respective CD4[+] and CD8[+] T-cell negative isolation kits

(Miltenyi), according to the manufacturer's instructions. The purity of the isolated lymphocyte

fractions was evaluated using multi-color flow cytometry analysis. **Table S2** presents the

composition of PBMCs and the purity of the isolated sub-populations prior to co-culturing as

percentage of CD45[+] cells.

AdV vectors.

The plasmid pcFG3CIITA8 was originally created by J. Ting (Chin, Li and Ting 1997). The human

CIITA cDNA contained in pcFG3CIITA8 was first cloned into an intermediate pShuttle vector

(pShuttle-CIITA) and then homologous recombined into an AdV backbone, according to the Ad-
Easy-1 protocol (He, Zhou et al. 1998). The resulting Ad-CIITA vector encodes MHC-II
TransActivator isoform III (NP_000237.2, 1130 aa). AdVs containing a mutated version of CIITA
(Ad-CIITA mutant) and a CMV promoter without transgene (Ad-null) were used as controls. Sanger
sequencing revealed that Ad-CIITA mutant contains a mutation at amino acid position 1088
(glycine to arginine). This mutation was visualized using PyMOL molecular visualization tool. For
all vectors, the AdV backbone represents a first-generation replication-deficient (ΔE1-E3)
adenovirus serotype 5, and transgene expression is regulated via the CMV promoter.

AdV production and titration.

AdV plasmids were digested with PacI restriction enzyme to linearize the DNA and therefore
expose the two ITRs necessary for virus production. Once linearized, the AdV vector DNA was
transfected into adherent 293T-cells with Lipofectamine™ LTX Reagent with PLUS™ Reagent
(Thermo Fisher) according to the manufacturer's instructions. Virus was harvested at 70-80%
cytopathic effect (CPE), submitted to three rounds of freeze (liquid nitrogen)/thaw (37°C) cycles,
and then centrifuged at 2000 rpm for 10 minutes at RT. After two rounds of amplification in 293T-
cells, the AdV particles were harvested at 50% CPE and purified via a chromatography-based
system via an Adeno-X™ Maxi Purification Kit (Takara), following the manufacturer's instructions.
All recombinant AdVs were titrated at the same time using the Adeno-X Rapid Titer kit (Takara)
according to the manufacturer's protocol. Briefly, 293T-cells were infected with 10-fold serial
dilutions of the viral stock from 10^{-2} to 10^{-6} and, 48 hours post-infection, were fixed in ice-cold
methanol and stained with a primary antibody specific for the AdV hexon protein. Virus titer is
expressed as infectious units (IU)/mL.

AdV infection.

As a standard procedure, AdV infection was conducted on the same day as cell seeding. To
prevent potential inactivation of viral particles due to proteins of the fetal bovine serum, AdVs
were diluted in serum-free media. Non-infected cells were used as internal controls and treated
with virus-free serum-free media. The choice of medium depended on the target cells to be
infected. For murine (GL261) and human (U251) cell lines, seeding was performed in DMEM
medium, and virus infection was carried out in serum-free DMEM medium. In the case of GSC
lines (P3NS and T16NS), cell seeding and virus infection were both performed using NSC medium,
as this medium does not contain serum. Afterward, both non-infected and AdV-infected cells
were gently shaken every 15-20 minutes for an hour to facilitate virus adhesion to the host cell

membrane and subsequent entry. Subsequently, they were incubated at 37°C for a maximum of six days post-infection. At day 6, cells were gently transferred in a bigger well and the medium replenished.

As for the infection of primary GBM organoids, cryopreserved GBM single cells (P3, T16, and T188), purified from PDOX tumor tissue *in vivo*, were thawed and seeded in 50 μL of organoid medium in 96-well, Nunclon Sphera-Treated, U-Shaped-Bottom, ultra-low attachment microplates (Thermo Fisher). Tumor cells were plated at a density of $2E10^4$ cells/well and then directly infected with 50 μL of AdVs diluted in serum-free organoid medium. Non-infected GBM organoids were instead treated with 50 μL of virus-free serum-free organoid medium. Cells were then gently shacked every 15-20 minutes for an hour, centrifuged at 300 x g for 5 minutes and then incubated at 37°C on an orbital shaker at 40-50 rpm for up to six days post-infection, depending on the read-out.

Assessment of virus cytotoxicity.

To assess virus cytotoxicity, mouse cell lines (GL261) were infected at different AdV multiplicity of infection (MOIs, infectious unit/cell, IU/cell): 0 (non-infected control), 1,56, 3,1, 6,2, 12,5, 25, 50 and 100. 2.5% DMSO (Sigma–Aldrich) was used as a positive control to induce cell death. Growth of AdV-infected cells was assessed via xCELLigence Real-Time Cell Analyzer over a period of 166 hours post-infection.

To assess virus cytotoxicity, human GBM organoids (P3dsRed, T16dsRed and cytolight rapid red dye-labeled T188 organoids) were infected at different AdV MOIs: 0 (non-infected control), 1, 5, 10, 25, 50 and 75. Growth of AdV-infected organoids was assessed via Incucyte S3 live cell imaging system for at least seven days post-infection.

AdV-infected GBM organoid - immune cell co-culture.

Three days prior co-culture, cryopreserved GBM single cells (P3dsRed, T16dsRed, and cytolight red rapid dye-labeled T188) were infected with AdV particles at the desired virus MOI in a final volume of 100 μL organoid medium. AdV infection was performed as described in the "AdV infection" section. The 96-well, Nunclon Sphera-Treated, U-Shaped-Bottom, ultra-low attachment (Thermo Fisher) were continuously agitated for three days and centrifuged daily at 300 × g for 3 minutes. GBM organoid formation and red fluorescence were monitored daily via Incucyte S3 live cell imaging system. On the day of the co-culture, cryopreserved human immune cells were thawed, counted and resuspended at $2\text{-}3E10^6$ cells/mL in T-cell medium. For all the three GBM models, we co-cultured the infected GBM organoids with three allogeneic healthy donors (CR014,

CR015, and CR017). Moreover, for P3 model, we co-cultured the infected GBM organoids with

three allogeneic HLA-matched donors (CTL573, CTL500, and CTL640). Co-cultures were either

performed with PBMCs, $CD3^+$, $CD4^+$, or $CD8^+$ T-cell fractions. Immune cells were diluted in 100 µL

of T-cell medium to reach different target:effector (T:E) ratios: 1:1, 1:5, 1:10, and added to the

GBM organoids, reaching a final volume of 200 µL. Co-cultures were performed for up to 3-4 days

in the Incucyte S3 live cell imaging system in the presence of 25 U/mL IL-2 (Miltenyi) and 5 µg/mL

anti-CD28 antibody (Biolegend). Depending on the read-out and the specific patient-derived

model, the co-culture was stopped at 1, 2, 3 or 4 days after co-culture. At the co-culture endpoint,

100ul of culture supernatant was collected from each experimental well and stored it at -20°C.

GBM organoids were dissociated for flow cytometry analysis.

Incucyte S3 live cell imaging.

The Incucyte S3 live cell imaging system was used to monitor red fluorescent protein-expressing

GBM organoids in real time, which can provide valuable information on tumor cell viability over

time. GBM organoids were imaged daily for the first 3 days post seeding and/or infection and,

after the addition of the immune cells, every 4 hours for a maximum of 96 hours after co-culture,

depending on the tumor model. GBM organoids were imaged with 4X magnification for phase

contrast, bright-field, and red channels. Data were then analyzed using Incucyte 2022A software.

In the case of P3dsRed and T16dsRed tumor models, wherein tumor cells exhibited stable

expression of the dsRed fluorescent protein, we quantified tumor cell viability by continuously

monitoring the total integrated red fluorescence intensity (RCU x µm²/Image) over time. Dying

tumor cells lose dsRed, enabling effective tracking of cell death and achieving precise

quantification through monitoring changes in red fluorescence intensity. As for T188 tumor

model, we transiently labeled T188 single tumor cells with 0.3 µM of Incucyte® Cytolight Rapid

Red Dye (Sartorius) according to the manufacturer's instructions. As for Cytolight-stained GBM

organoids, the red fluorescent signal persisted post-cell death. To assess tumor cell viability, the

focus was on evaluating the structural integrity of the organoids by measuring the red total area

($µm^2$/Image) over time. An increase in the red fluorescent area served as an indicator of GBM

organoid disruption.

Immunohistochemistry and immunofluorescence on paraffin sections.

AdV-infected and non-infected GBM organoids (300-450 µm in size) were carefully transferred

with wide-bore pipette tips from the 96-well, Nunclon Sphera-Treated, U-Shaped-Bottom, ultra-

low attachment (Thermo Fisher) into 15 mL Falcon tubes. They were gently washed with ice-cold

1X PBS and fixed in 4% paraformaldehyde for 20-30 minutes at RT. Fixation time depends on the

organoid size. Organoid sedimentation was preferred over centrifugation to preserve organoid

structure. Organoids were then gently washed, resuspended in 2.5% agar diluted in water and

transferred into Tissue-Tek cryomolds (Sakura) kept on ice for 5-10 min until agar solidification. If

present, bubbles were removed with a needle. With the help of forceps, agar blocks were then

moved to cassettes for automated paraffin embedding. Immunohistochemical and

immunofluorescence staining was performed following standard procedures with the antibodies

listed in **Table S4**.

Two-layer transwell co-culture assay.

AdV-infected GBM organoid – PBMC co-cultures were carried out as described in "AdV-infected

GBM organoid – immune cell co-culture" section using a two-layer transwell model to investigate

the cell contact dependency of immune cell-mediated tumor killing. Transwell-96 with 0.4 μm

pore polyester membrane inserts (StemCell Technologies) were used to physically separate the

AdV-infected and non-infected GBM organoids from the immune cells, which were plated on top

of the transwell inserts. Immune cell-mediated tumor killing was assessed in real time via the

Incucyte S3 live cell imaging system.

Degranulation and T-cell activation assay.

To assess the degranulation activity of the T lymphocytes in the co-culture, flow cytometry was

performed to measure the surface expression of the degranulation marker CD107a (**Table S4**).

AdV-infected GBM organoid – CD3$^+$ T-cell co-cultures were carried out as described in the "AdV-

infected GBM organoid – immune cell co-culture" section. At 24 and 48 hours post co-culture,

with a T:E ratio of 1:10, the GBM organoid – CD3$^+$ T-cells were incubated with anti-CD107a

antibody for 1 hour at 37°C. As a positive control, freshly thawed CD3$^+$ T-cells were stimulated

with a combination of 50 ng/mL phorbol 12-myristate 13-acetate (PMA) (Enzo Life Sciences) and 1

μg/mL ionomycin (Enzo Life Sciences) and incubated with BV785-conjugated anti-CD107a

antibody for 1 hour at 37°C. Both co-culture samples and positive controls were then incubated

for 4 hours at 37°C with BD GolgiStop™ (dilution 1:1000), containing monensin, and BD GolgiPlug

(dilution 1:500), containing brefeldin A. These protein transport inhibitors block intracellular

protein transport processes, leading to the accumulation of cytokines in the Golgi complex. CD3$^+$

T-cells from five different wells (~1 million cells) were pooled together, fixed, and acquired using

the NovoCyte Quanteon Flow Cytometer, and the resulting data were analyzed using FlowJo

version 10 software.

Neutralization assay.

AdV-infected GBM organoid – PBMC co-cultures were carried out as described in the "AdV-infected GBM organoid – immune cell co-culture" section. At 72 hours post-infection and prior to the addition of immune cells, organoids were treated with increasing concentrations of anti-human TNF-RI/TNFRSF1A (0.16 µg/mL, 0.08 µg/mL, 0.04 µg/mL, and 0.02 µg/mL) and anti-human Fas Ligand/TNFSF6 (20 ng/mL, 10 ng/mL, 5 ng/mL, and 2.5 ng/mL) monoclonal antibodies (**Table S4**). Immune cell-mediated tumor killing was assessed in real time via the Incucyte S3 live cell imaging system.

Measurement of cytokine secretion.

At 72 hours post co-culture, conditioned media from GBM organoid – PBMC co-cultures were collected, followed by centrifugation to remove particulates, and then stored at -20°C. To avoid protein degradation, repeated freeze-thaw cycles were avoided. These co-culture supernatants were subsequently screened using the Proteome Profiler Human XL Cytokine Array Kit (R&D), capable of detecting 105 different cytokines simultaneously, following the manufacturer's instructions. For each infection condition (Non-infected, Ad-null, Ad-CIITA), a total volume of 380 µL of co-culture supernatant was used per membrane of the cytokine array kit. Once the complete protocol was executed, each membrane was simultaneously exposed to X-Ray films with multiple exposure times (1, 5, and 10 minutes). The average signal (pixel density) per analyte was quantified using Quick Spots software.

Western blot.

Total proteins from AdV-infected and non-infected mouse cell lines and human GSCs were extracted in RIPA lysis buffer (Thermo Fisher) supplemented with protease and phosphatase inhibitors (Thermo Fisher) for 45 minutes at 4°C and then quantified using a bicinchoninic acid (BCA) protein assay. Heat-denatured protein samples were then separated on NuPage 4-12% BisTris gels (Thermo Fisher) and blotted onto PVDF membranes. To prevent nonspecific antibody binding, membranes were then blocked in 4% (wt/vol) skimmed milk in 1X Tris-Buffered Saline with 0.1% Tween 20 (1X TBS-T) for 1 hour at room temperature (RT). Membranes were first incubated with anti-CIITA and anti-HLA-DR primary antibodies diluted in blocking solution overnight at 4°C and, after washing steps in 1X TBS-T, with horseradish peroxidase (HRP)-conjugated secondary antibodies diluted in blocking solution for 1 hour at RT. Anti-GAPDH and anti-LaminB1 primary antibodies were used as internal loading controls. A detailed list of the antibodies and dilutions is available in **Table S4**. The chemiluminescent signal was revealed using the Pierce™ ECL Western Blotting Substrate (Thermo Fisher) using the ImageQuant LAS4010 analyzer.

Quantitative RT–PCR.

Total RNA from AdV-infected and non-infected mouse cell lines and human GSCs was extracted using the RNeasy Mini Kit (Qiagen) and reverse-transcribed to cDNA to measure CIITA and MHC-II gene expression using Fast SYBR Green (Thermo Fisher) and QuantStudio 5 Real-Time PCR System. It was run in biological triplicates, and the relative gene expression levels were calculated through the ΔCT method by using *elongation factor 1 alpha (EF1α)* as a housekeeping gene. The primers used were as follows: CIITA (Fw: TTATGCCAATATCGCGGAACTG, Rv: CATCTGGTCCTATGTGCTTGAAA), HLA-DRα (Fw: AGTCCCTGTGCTAGGATTTTTCA, Rv: ACATAAACTCGCCTGATTGGTC), EF1α (Fw: TTGTCGTCATTGGACACGTAG, Rv: TGCCACCGCATTTATAGATCAG), and GAPDH (Fw: CTCCCACTCTTCCACCTTCG, Rv: CCACCACCCTGTTGCTGTAG).

Immunofluorescence.

U251 cells were seeded at a density of 5E10^4 cells/well in 8-well μ-slides (Ibidi). Cells were kept overnight at 37°C and then either infected with AdVs or left uninfected. Three days post-infection, cells were fixed with 4% PFA for 15 minutes at RT and, after washing steps in 1X PBS, blocked in 1X PBS with 0.5% Triton X-100 and 10% FBS for 1 hour at RT. Cells were then incubated with anti-CIITA primary antibodies diluted in 1X PBS with 0.5% Triton X-100 overnight at 4°C. After three washes in 1X PBS, the cells were incubated with fluorescently labeled secondary antibodies and Hoechst dye diluted in 1X PBS with 0.01% BSA for 1 hour at RT. A detailed list of the antibodies and dilutions used can be found in **Table S4**. Cells were then imaged with a Zeiss LSM 880 Confocal Laser Scanning Microscope and Zen Blue software.

Flow cytometry.

Single cells were stained with fluorochrome-conjugated antibodies diluted in ice-cold 1X Phosphate-Buffered Saline (PBS) with 0,2% BSA ("FACS buffer") for 30 minutes at 4°C in the dark. Alternatively, antibodies were diluted in ice-cold BD Horizon Brilliant Stain Buffer ("BV buffer") for multicolor immunofluorescent staining using three or more brilliant violet antibodies. After wash, cells were re-suspended in 100 μL of FACS buffer prior acquisition (1 million cells in 100 μL, we scaled down accordingly for a smaller number of cells). Stained samples were then acquired with NovoCyte Quanteon Flow Cytometer and the resulting data were analyzed using FlowJo version 10. A detailed list of all the antibodies and dilutions used can be found in **Table S4**.

Statistical analysis.

The experimental results were analyzed using GraphPad Prism 9 and reported as the mean ±

standard deviation (SD) or, for co-culture experiments, as mean ± standard error of mean (SEM).

One-way ANOVA was employed to compare multiple groups, while two-way ANOVA was used to

compare two or more factors simultaneously. All means were calculated based on a minimum of

three independent experiments. For tumor organoid-immune cell co-culture experiments, all

means were calculated based on a minimum of three immune cell donors, representing three

different biological replicates. A P value < 0.05 was used to determine statistical significance.

ACKNOWLEDGEMENTS: We are grateful for the use of the LIH PRECISION-PDX Brain Tumor Bank

(www.precision-pdx.lu), supported by the patients, the Neurosurgery Department of the Centre

Hospitalier de Luxembourg, the NORLUX Neuro-Oncology Laboratory and the Clinical and

Epidemiological Investigation Center of the LIH. We thank Prof. Roberto Accolla and Prof. Greta

Forlani for providing cell lines and critically reviewing the manuscript, Pilar M. Moreno-Sanchez

for technical assistance in the management of PDOXs, and the Luxembourg Croix Rouge and its

donors for generously providing healthy donor blood samples. We acknowledge the support from

LIH's core facilities (Animal facility, In vivo imaging facility, National Cytometry Platform, LUXGEN

sequencing platform).

AUTHOR CONTRIBUTION: Conceptualization (I. S., A.P. A.G., S.P.N., A.M.), methodology (I.S., E.K.,

A.P., M.R., A.C., A.G.), investigation (I.S., E.K., A.P., M.R., A.L., A.O., V.B., G.M.D.), formal analysis

(L.E., B.N., A.G.), supervision (A.P., A.G., S.P.N., A.M.) writing - original draft (I.S., A.G., S.P.N.,

A.M.), and writing – review & editing (all authors). The authors approved the final version of the

manuscript as submitted and agreed to be accountable for all the aspects of the work.

CONFLICT OF INTEREST: The authors have nothing to disclose.

THE PAPER EXPLAINED:

Problem: Glioblastoma (GBM) is a highly aggressive brain tumor with a poor prognosis. Current

immunotherapies for GBM have been largely unsuccessful due to the tumor's highly

immunosuppressive microenvironment. Consequently, there is an urgent need for new strategies

to boost immune responses against GBM.

Results: We successfully engineered and produced a non-lytic adenovirus (AdV) for the delivery to

cancer cells of CIITA, the master regulator of MHC-II antigen processing and presenting molecules.

The resulting AdV-induced MHC-II expression on GBM cell surfaces for at least six days post-

infection. Interestingly, infection with both an AdV carrying a wild-type nuclear CIITA and a
mutant cytoplasmic version of CIITA, but not with a GFP-armed AdV, resulted in substantial T-cell-
mediated tumor cell death. The latter was mediated via MHC-II-independent pathways, in
conjunction with MHC-II antigen presentation. Moreover, tumor cell death relied on direct
contact between GBM and immune cells, without involving canonical death pathways.

Impact: The study introduces a novel approach to combat the immunosuppressive GBM tumor
environment through AdV-mediated delivery of the *CIITA* transgene. This method shows promise
in enhancing T-cell-mediated immunity against GBM, providing a hopeful avenue for future GBM
treatment strategies. Additionally, it unveils an unexpected role of CIITA in triggering anti-tumor
immune responses that appears to be at least in part independent of MHC-II antigen
presentation.

FUNDING INFORMATION: We acknowledge the financial support by the Luxembourg Institute of
Health, FNRS-Télévie (GBModImm no. 7.8513.18, GBModImm2 no. 7651720F, ImmoGBM
7.8505.20, ImmoGBM2 7.6603.22) and the Luxembourg National Research Fund
(C20/BM/14646004/GLASSLUX, C21/BM/15739125/DIOMEDES).

742
743

Figure 1. Correlation between CIITA and MHC-II expression in human GBM and associated
models. A. Bulk RNA-seq gene expression of CIITA and selected MHC-II molecules in GBM patient
tumors (n=163) versus control normal brain (n=207). *p<0.01 (one-way ANOVA). **B.** Single cell
RNA-seq gene expression profiles in GBM patient tumors. Detected cells were classified into
tumor and non-neoplastic cells of the tumor microenvironment (top dot plot). Gene expression is
depicted for major cell types of the tumor microenvironment in GBM (bottom dot plot). The size
and color of each dot represent the percentage of cells within a class and the average expression
of a gene within a class, respectively. **C.** Heatmap representing bulk RNA-seq (human specific)
gene expression in GBM patient-derived orthotopic xenografts (PDOXs): selected MHC-II
molecules associated with enhanceosome, antigen processing, and presentation are shown.
CXADR, *ITGAV* and *ITGB5* are associated with AdV infection capacity. A lack of expression is
represented by the grey color; in every PDOX, the associated replicates are indicated by the
numbers. **D.** Flow cytometric analysis of MHC-II molecules at the cell membrane (anti-HLA-
DR/DP/DQ-FITC) in various GBM cell lines in control conditions (cells unstained or mock-
stimulated) and upon stimulation with IFN-γ.

Figure 2. Characterization of Ad-CIITA in human glioblastoma stem-like cells (GSCs). **A**.
Schematic depicting the serotype 5 AdV (ΔE1-E3) vectors used in the study (see results for
details). **B-D**. Virus quality check in human GSC lines P3NS and T16NS infected at an MOI of 50.
CIITA and HLA-DRα expression at 72 hours post-infection. mRNA (**B**) and protein levels (**C**) of CIITA
and HLA-DRα were analyzed by quantitative RT–PCR and Western blot analysis, respectively. qRT–
PCR data were normalized to the amount of Ef1a, while Western blot data were normalized to the
amount of GAPDH. Mean ± SD. ***p<0.001 and ****p<0.0001 (one-way ANOVA). **D**. Surface
expression of MHC-II (anti-HLA-DR/DP/DQ-FITC) molecules was analyzed by flow cytometry at 72
hours and 6 days post-infection.

Figure 3. Successful infection of human primary GBM organoids by Ad-CIITA vectors. A. CIITA
and HLA-DRα expression levels in P3 GBM organoids infected with Ad-CIITA or control virus at
increasing MOIs (5, 10, 25, 50, 75) were analyzed by Western blot analysis at 72 hours post-
infection. **B-C.** P3 GBM organoids at MOI 50: Representative H&E (top) and anti-HLA-DRα
(bottom) staining at 72 hours post-infection (**B**). Flow cytometry analysis of cell surface expression
of MHC-II (anti-HLA-DR/DP/DQ-FITC) at 72 hours and 6 days post-infection (**C**). **D.** P3dsRed GBM
organoids infected with Ad-CIITA or control virus at increasing MOIs (5, 10, 25, 50, 75). Evaluation
of virus-mediated cytotoxicity as determined via the Incucyte S3 live cell imaging system. Graphs
represent tumor cell viability as total red integrated intensity (RCU x µm²/Image) over a period of
186 hours post-infection. Results are expressed as mean ± SEM.

Figure 4. Immune cell-mediated tumor cell killing in human primary GBM organoids induced by
Ad-CIITA and Ad-CIITA mutant A. Schematic of the co-culture setup of Ad-infected GBM
organoids with immune cells. Red fluorescent GBM organoids were seeded and infected for 72
hours. Co-culture with effector cells (PBMCs, CD3$^+$, CD4$^+$, CD8$^+$) was performed in the presence of
IL-2 and anti-CD28 antibody for 3 days. Tumor cell viability was assessed using the IncucyteS3 live
cell imaging system. For the dsRed-expressing GBM organoids, viability was determined by
measuring the dsRed fluorescence levels, namely total red integrated intensity (RCU x µm²/Image)
over time. **B-D.** Evaluation of viability of P3dsRed organoids infected at an MOI of 75. **B.** GBM

organoids alone (no co-culture) or in co-culture with PBMCs. **C.** GBM organoids in co-culture with
either purified CD3$^+$ T-cells, CD4$^+$ T-cells or CD8$^+$ T-cells. **D.** HLA-matched co-culture with either
PBMCs or purified CD3$^+$ T-cells.

Each co-culture panel includes: i) a graph displaying curves of tumor cell viability (measured as
total red integrated intensity [RCU x μm²/Image]) over a period of 134 hours after infection, ii) a
quantification of tumor cell viability at the co-culture endpoint (134 hours) at the tested T:E ratios
(indicated on the x-axis), and iii) representative phase contrast and red fluorescence images of
GBM organoids alone or in co-culture (T:E ratio of 1:10) with immune cells at the co-culture
endpoint (134 hours). Results are expressed as mean ± SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and
****$p < 0.0001$ (two-way ANOVA).

Figure 5. Mechanisms of CIITA-mediated immune cell killing in human primary GBM organoids.
A-C. P3dsRed organoids infected at a virus MOI of 75 and in co-culture with healthy donor PBMCs
at a 1:10 ratio. **A.** Representative live images of phase contrast and green fluorescence in Ad-GFP-
infected and non-infected organoids. Side quantification shows the total green integrated
intensity (GCU x μm²/Image) at 24, 48, and 72 hours post-infection compared to non-infected
control. Mean ± SD. ****$p < 0.0001$ (two-way ANOVA). **B.** Immune-mediated tumor cell killing in
Ad-GFP- or Ad-CIITA-infected GBM organoids compared to non-infected control. Mean ± SEM.
****$p < 0.0001$ (one-way ANOVA). **C.** Immune-mediated tumor cell killing in two-layer transwell
co-culture. Graphs display curves of tumor cell viability (measured as total red integrated intensity
[RCU x μm²/Image] over a period of 134 hours post-infection. Side quantifications show tumor cell
viability at the co-culture endpoint (134 hours). Representative live images show phase contrast
and red fluorescence in GBM organoids at the co-culture endpoint (134 hours). Mean ± SEM.
$p < 0.01$ (one-way ANOVA). **D. Surface expression of the degranulating marker CD107a in CD3$^+$ T-
cells at 24 hours post co-culture with infected organoids, as analyzed via flow cytometry assay.
The data are shown as overlay histograms (left panel) and graphs quantifying the mean
fluorescence intensity (MFI) (right panel) of CD107a in CD4$^+$ and CD8$^+$ T-cell populations. As
controls for T-cell activation, unstimulated CD3$^+$ T-cells and CD3$^+$ T-cells stimulated for 4 hours
with PMA and ionomycin were used as negative and positive controls, respectively. Mean ± SD.
$p < 0.01$ and ***$p < 0.001$ (one-way ANOVA). **E. Neutralization of death signaling pathways.
Organoids were treated with increasing concentrations of anti-human TNF-R (MAB225) (left
panel) and anti-human FasL (MAB126) (right panel) monoclonal antibodies and co-cultured with
healthy donor PBMCs. Mean ± SEM. **F.** Screening of conditioned media from GBM organoid-PBMC
co-cultures (T:E of 1:10) using a protein cytokine array (400 μL of cell supernatants, 10-minute
exposure). On the bottom left, heatmap of the most upregulated and downregulated proteins
detected via protein array based on mean pixel density. Each row represents a specific cytokine
with its array coordinates, while each column represents a co-culture condition. On the bottom
right, graphs show levels of secreted EMMPRIN and RANTES protein, expressed as mean pixel
density. Mean ± SD.

 SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1. MHC-II gene expression analysis in GBM patient tumors and human preclinical models. A. Heatmap representing gene expression levels of selected MHC-II molecules associated with enhanceosome, antigen processing, presentation, and AdV viral infection capacity in normal human brain, GBM patient tumors, patient-derived orthotopic xenografts (PDOXs), GSC lines (NCH421k, NCH644), and classical glioma lines (U87, U251) grown *in vitro* or *in vivo* as xenograft ('X'). A lack of expression is represented by the grey color. Human specific arrays were applied for transcriptome analysis. **B.** Flow cytometric analysis of MHC-II cell membrane expression (anti-HLA-DR/DP/DQ-FITC) in GBM cells in control conditions (cells unstained or mock-stimulated) and upon stimulation with IFN-γ.

Supplementary Figure 2. Characterization of viruses in adherent human (U251) and murine (GL261) cell lines. A. Sanger sequencing results of the *CIITA* mutant DNA show a guanine (g) to adenine (a) genetic mutation in the *CIITA* gene that results in the substitution from a glycine (Gly) to an arginine (Arg) at amino acid position 1088 of a full-length CIITA protein. **B.** Visual representation of the amino acid string surrounding amino acid position 1088 (highlighted in yellow) in wild-type (left panel) and mutated (right panel) CIITA protein. Molecular figures were prepared via PyMOL. **C.** Representative confocal microscopy images of CIITA immunofluorescence staining (Alexa Fluor 488) in human U251 cells at 72 hours post-infection. Nuclei were counterstained in blue (Hoechst dye) (magnification 63X). The distribution of CIITA proteins between the cytoplasm and nucleus was quantified by measuring the ratio of cytoplasmic to nuclear fluorescence intensity. Mean ± SD. ****$p<0.0001$ (two-way ANOVA). **D-F.** Ad-CIITA characterization in murine GL261 cells **D.** MHC-II (anti-IA-IE-PerCP) expression as a result of Ad-CIITA infection at increasing virus MOIs (1.56, 3.1, 6.2, 12.5, 25, 50, 100, as analyzed by flow cytometry. **E.** mRNA (left) and protein (right) levels of CIITA at 48 and 72 hours post-infection, as analyzed by quantitative RT–PCR and by Western blot analysis, respectively. RT-PCR results were normalized to the amount of GAPDH. Mean ± SD. ***$p<0.001$, ****$p<0.0001$ (two-way ANOVA). Western Blot data were normalized to the amount of GAPDH. **F.** Kinetics of the expression of MHC-II (anti-IA-IE-PerCP) molecules, as analyzed by flow cytometry.

Supplementary Figure 3. Assessment of virus-mediated cytotoxicity in human primary GBM organoids. T16dsRed (top) and Cytolight Rapid Red Dye-labeled T188 (bottom) GBM organoids infected with Ad-CIITA or control viruses at increasing MOIs (5, 10, 25, 50, 75). Evaluation of virus-mediated cytotoxicity as determined via the Incucyte S3 live cell imaging system. Graphs represent tumor cell viability as total red integrated intensity (RCU x μm²/image) for T16dsRed organoids, and red total area (μm²/Image) for cytolight rapid red dye-labeled T188 organoids, over a 186-hour post-infection period. Results are expressed as mean ± SEM.

Supplementary Figure 4. Evaluation of immune cell-mediated tumor cell killing in human primary GBM organoids. A. Flow cytometry gating strategy to phenotype PBMCs and MACS-purified isolated T-cell fractions. **B-C.** Evaluation of viability of T16dsRed (left panel) or of cytolight rapid red dye-labeled T188 (right panel) organoids infected at an MOI of 25 or 75, respectively. Deviations from the virus MOI are explicitly indicated on the graph. **B.** GBM organoids alone (no co-culture) or in co-culture with PBMCs. In the case of cytolight rapid red dye-labeled T188 organoids, viability was assessed by measuring organoid integrity, namely the red total area

(μm^2/Image) over time. **C.** GBM organoids in co-culture with either purified CD3[+] T-
cells or CD8[+] T-cells. **D.** Evaluation of viability of T16dsRed organoids infected at an MOI of 25 in
HLA-matched co-culture with either PBMCs or purified CD3[+] T-cells. T16dsRed co-culture panels
include: i) a graph displaying curves of tumor cell viability (measured as total red integrated
intensity [RCU x μm^2/Image]) over a period of 170 hours after infection, ii) a quantification of
tumor cell viability at the co-culture endpoint (170 hours) at the tested T:E ratios (indicated on the
x-axis), and iii) representative Incucyte S3 phase contrast and red fluorescence images of GBM
organoids in co-culture (T:E ratio of 1:10) with immune cells at the co-culture endpoint. Mean ±
SEM. *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001 (two-way ANOVA). Cytolight-stained
T188 co-culture panels include: i) a quantification of tumor cell viability (measured as total red
area [μm^2/Image]) at the co-culture endpoint (134 hours) at the tested T:E ratios (indicated on the
x-axis), and ii) representative Incucyte S3 phase contrast and red fluorescence images of GBM
organoids in co-culture (T:E ratio of 1:10) with immune cells at the co-culture endpoint. Mean ±
SEM. *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001 (two-way ANOVA).

Supplementary Figure 5. Mechanisms of CIITA-mediated immune cell killing in human primary
GBM organoids. A-C. T16dsRed organoids infected at a virus MOI of 25 and in co-culture with
healthy donor PBMCs at a 1:10 ratio. **A.** Representative Incucyte S3 images show phase contrast
and green fluorescence in Ad-GFP-infected and non-infected organoids. Side quantification shows
the total green integrated intensity (GCU x μm^2/Image) at 24, 48, and 72 hours post-infection
compared to non-infected control. Mean ± SD. ****p<0.0001 (two-way ANOVA). **B.** Immune-
mediated tumor cell killing in Ad-GFP- or Ad-CIITA-infected GBM organoids compared to non-
infected control. Mean ± SEM. *p<0.05, **p<0.01 (one-way ANOVA). **C.** Immune-mediated tumor
cell killing in two-layer transwell co-culture. Graphs display curves of tumor cell viability
(measured as total red integrated intensity [RCU x μm^2/Image] over a period of 170 hours post-
infection. Side quantifications show tumor cell viability at the co-culture endpoint (170 hours).
Representative Incucyte S3 images show phase contrast and red fluorescence in GBM organoids
at the co-culture endpoint (170 hours). Mean ± SEM. **D.** Flow cytometry analysis of cell surface
expression of MHC-I (anti-HLA-A/B/C-BV510), CD80 (anti-CD80-PE), CD86 (anti-CD86-BV421) as
analyzed by flow cytometry. **E.** Positive controls of death signaling pathway neutralization.
P3dsRed organoids were treated with (left) actinomycin D (1 µg/mL) and increasing
concentrations of anti-human TNF-R (MAB225) or with (right) 2.5% DMSO and anti-human FasL
(MAB126). Mean ± SEM.

900
901

REFERENCES

Accolla, R. S., E. Ramia, A. Tedeschi and G. Forlani (2019). "CIITA-Driven MHC Class II Expressing Tumor Cells
as Antigen Presenting Cell Performers: Toward the Construction of an Optimal Anti-tumor Vaccine." Front
Immunol **10**: 1806.

Agosti, E., M. Zeppieri, L. De Maria, C. Tedeschi, M. M. Fontanella, P. P. Panciani and T. Ius (2023).
"Glioblastoma Immunotherapy: A Systematic Review of the Present Strategies and Prospects for
Advancements." Int J Mol Sci **24**(20).

Appay, V., P. R. Dunbar, V. Cerundolo, A. McMichael, L. Czaplewski and S. Rowland-Jones (2000). "RANTES
activates antigen-specific cytotoxic T lymphocytes in a mitogen-like manner through cell surface
aggregation." International Immunology **12**(8): 1173-1182.

Banerjee, K., F. J. Nunez, S. Haase, B. L. McClellan, S. M. Faisal, S. V. Carney, J. Yu, M. S. Alghamri, A. S. Asad,
A. J. N. Candia, M. L. Varela, M. Candolfi, P. R. Lowenstein and M. G. Castro (2021). "Current Approaches for
Glioma Gene Therapy and Virotherapy." Front Mol Neurosci **14**: 621831.

Barry, M. and R. C. Bleackley (2002). "Cytotoxic T lymphocytes: all roads lead to death." Nat Rev Immunol
2(6): 401-409.

Bou Nasser Eddine, F., G. Forlani, L. Lombardo, A. Tedeschi, G. Tosi and R. S. Accolla (2017). "CIITA-driven
MHC class II expressing tumor cells can efficiently prime naive CD4(+) TH cells in vivo and vaccinate the host
against parental MHC-II-negative tumor cells." Oncoimmunology **6**(1): e1261777.

Bulcha, J. T., Y. Wang, H. Ma, P. W. L. Tai and G. Gao (2021). "Viral vector platforms within the gene therapy
landscape." Signal Transduct Target Ther **6**(1): 53.

Burster, T., F. Gärtner, C. Bulach, A. Zhanapiya, A. Gihring and U. Knippschild (2021). "Regulation of MHC I
Molecules in Glioblastoma Cells and the Sensitizing of NK Cells." Pharmaceuticals (Basel) **14**(3).

Celesti, F., A. Gatta, M. Shallak, A. M. Chiaravalli, M. Cerati, F. Sessa, R. S. Accolla and G. Forlani (2023).
"Protective anti-tumor vaccination against glioblastoma expressing the MHC class II transactivator CIITA."
Front Immunol **14**: 1133177.

Chih, Y., K. Sahm, A. Sadik, T. Bunse, N. Trautwein, S. Pusch, S. Stevanovic, C. Opitz, A. Deimling, W. Wick, F.
Sahm, L. Bunse and M. Platten (2021). "KS01.3.A Tumoral MHC class II expression in gliomas drives T cell
exhaustion." Neuro-Oncology **23**(Supplement_2): ii3-ii3.

Chin, K.-C., G. G.-X. Li and J. P.-Y. Ting (1997). "Importance of acidic, proline/serine/threonine-rich, and
GTPbinding regions in the major histocompatibility complex class II transactivator: Generation
of transdominant-negative mutants." Proceedings of the National Academy of Sciences **94**(6): 2501-2506.

Cloughesy, T. F., A. Brenner, J. F. de Groot, N. A. Butowski, L. Zach, J. L. Campian, B. M. Ellingson, L. S.
Freedman, Y. C. Cohen, N. Lowenton-Spier, T. Rachmilewitz Minei, S. Fain Shmueli and P. Y. Wen (2020). "A
randomized controlled phase III study of VB-111 combined with bevacizumab vs bevacizumab monotherapy
in patients with recurrent glioblastoma (GLOBE)." Neuro Oncol **22**(5): 705-717.

Cloughesy, T. F., K. Petrecca, T. Walbert, N. Butowski, M. Salacz, J. Perry, D. Damek, D. Bota, C. Bettegowda,
J.-J. Zhu, F. Iwamoto, D. Placantonakis, L. Kim, B. Elder, G. Kaptain, D. Cachia, Y. Moshel, S. Brem, D. Piccioni,
J. Landolfi, C. C. Chen, H. Gruber, A. R. Rao, D. Hogan, W. Accomando, D. Ostertag, T. T. Montellano, T.
Kheoh, F. Kabbinavar and M. A. Vogelbaum (2020). "Effect of Vocimagene Amiretrorepvec in Combination
With Flucytosine vs Standard of Care on Survival Following Tumor Resection in Patients With Recurrent
High-Grade Glioma: A Randomized Clinical Trial." JAMA Oncology **6**(12): 1939-1946.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson and T. R. Gingeras
(2013). "STAR: ultrafast universal RNA-seq aligner." Bioinformatics **29**(1): 15-21.

Drozina, G., J. Kohoutek, N. Jabrane-Ferrat and B. M. Peterlin (2005). "Expression of MHC II genes." Curr Top
Microbiol Immunol **290**: 147-170.

Forlani, G., M. Shallak, A. Gatta, A. K. B. Shaik and R. S. Accolla (2023). "The NLR member CIITA: Master
controller of adaptive and intrinsic immunity and unexpected tool in cancer immunotherapy." Biomed J
46(5): 100631.

Frangione, V., L. Mortara, P. Castellani, A. De Lerma Barbaro and R. S. Accolla (2010). "CIITA-driven MHC-II
positive tumor cells: preventive vaccines and superior generators of antitumor CD4+ T lymphocytes for
immunotherapy." Int J Cancer **127**(7): 1614-1624.

Golebiewska, A., A.-C. Hau, A. Oudin, D. Stieber, Y. A. Yabo, V. Baus, V. Barthelemy, E. Klein, S. Bougnaud, O.
Keunen, M. Wantz, A. Michelucci, V. Neirinckx, A. Muller, T. Kaoma, P. V. Nazarov, F. Azuaje, A. De Falco, B.
Flies, L. Richart, S. Poovathingal, T. Arns, K. Grzyb, A. Mock, C. Herold-Mende, A. Steino, D. Brown, P. May,
H. Miletic, T. M. Malta, H. Noushmehr, Y.-J. Kwon, W. Jahn, B. Klink, G. Tanner, L. F. Stead, M. Mittelbronn,
A. Skupin, F. Hertel, R. Bjerkvig and S. P. Niclou (2020). "Patient-derived organoids and orthotopic
xenografts of primary and recurrent gliomas represent relevant patient avatars for precision oncology."
Acta Neuropathologica **140**(6): 919-949.

He, T.-C., S. Zhou, L. T. da Costa, J. Yu, K. W. Kinzler and B. Vogelstein (1998). "A simplified system for
generating recombinant adenoviruses." Proceedings of the National Academy of Sciences **95**(5): 2509-2514.

Hillman, G. G., N. L. Kallinteris, J. Li, Y. Wang, X. Lu, Y. Li, S. Wu, J. L. Wright, P. Slos, J. V. Gulfo, R. E.
Humphreys and M. Xu (2003). "Generating MHC Class II+/Ii- phenotype after adenoviral delivery of both an
expressible gene for MHC Class II inducer and an antisense Ii-RNA construct in tumor cells." Gene Ther
10(17): 1512-1518.

Immonen, A., M. Vapalahti, K. Tyynelä, H. Hurskainen, A. Sandmair, R. Vanninen, G. Langford, N. Murray
and S. Ylä-Herttuala (2004). "AdvHSV-tk gene therapy with intravenous ganciclovir improves survival in
human malignant glioma: a randomised, controlled study." Mol Ther **10**(5): 967-972.

Kaushik, D. K., H. Y. Yong, J. N. Hahn, C. Silva, S. Casha, R. J. Hurlbert, F. H. Jacques, R. Lisak, O. Khan, C.
Ionete, C. Larochelle, A. Prat, A. Bar-Or and V. W. Yong (2016). "Evaluating Soluble EMMPRIN as a Marker of
Disease Activity in Multiple Sclerosis: Studies of Serum and Cerebrospinal Fluid." PLoS One **11**(10):
e0163802.

Kluin, R. J. C., K. Kemper, T. Kuilman, J. R. de Ruiter, V. Iyer, J. V. Forment, P. Cornelissen-Steijger, I. de Rink,
P. Ter Brugge, J. Y. Song, S. Klarenbeek, U. McDermott, J. Jonkers, A. Velds, D. J. Adams, D. S. Peeper and O.
Krijgsman (2018). "XenofilteR: computational deconvolution of mouse and human reads in tumor xenograft
sequence data." BMC Bioinformatics **19**(1): 366.

Leon Machado, J. A. and V. Steimle (2021). "The MHC Class II Transactivator CIITA: Not (Quite) the Odd-
One-Out Anymore among NLR Proteins." Int J Mol Sci **22**(3).

Liao, Y., G. K. Smyth and W. Shi (2014). "featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features." Bioinformatics **30**(7): 923-930.

Liau, L. M., K. Ashkan, S. Brem, J. L. Campian, J. E. Trusheim, F. M. Iwamoto, D. D. Tran, G. Ansstas, C. S.
Cobbs, J. A. Heth, M. E. Salacz, S. D'Andre, R. D. Aiken, Y. A. Moshel, J. Y. Nam, C. P. Pillainayagam, S. A.
Wagner, K. A. Walter, R. Chaudhary, S. A. Goldlust, I. Y. Lee, D. A. Bota, H. Elinzano, J. Grewal, K. Lillehei, T.
Mikkelsen, T. Walbert, S. Abram, A. J. Brenner, M. G. Ewend, S. Khagi, D. S. Lovick, J. Portnow, L. Kim, W. G.
Loudon, N. L. Martinez, R. C. Thompson, D. E. Avigan, K. L. Fink, F. J. Geoffroy, P. Giglio, O. Gligich, D. Krex, S.
M. Lindhorst, J. Lutzky, H. J. Meisel, M. Nadji-Ohl, L. Sanchin, A. Sloan, L. P. Taylor, J. K. Wu, E. M. Dunbar, A.
B. Etame, S. Kesari, D. Mathieu, D. E. Piccioni, D. S. Baskin, M. Lacroix, S. A. May, P. Z. New, T. J. Pluard, S. A.
Toms, V. Tse, S. Peak, J. L. Villano, J. D. Battiste, P. J. Mulholland, M. L. Pearlman, K. Petrecca, M. Schulder,
R. M. Prins, A. L. Boynton and M. L. Bosch (2023). "Association of Autologous Tumor Lysate-Loaded

Dendritic Cell Vaccination With Extension of Survival Among Patients With Newly Diagnosed and Recurrent Glioblastoma: A Phase 3 Prospective Externally Controlled Cohort Trial." JAMA Oncol **9**(1): 112-121.

Lim, M., M. Weller, A. Idbaih, J. Steinbach, G. Finocchiaro, R. R. Raval, G. Ansstas, J. Baehring, J. W. Taylor, J. Honnorat, K. Petrecca, F. De Vos, A. Wick, A. Sumrall, S. Sahebjam, I. K. Mellinghoff, M. Kinoshita, M. Roberts, R. Slepetis, D. Warad, D. Leung, M. Lee, D. A. Reardon and A. Omuro (2022). "Phase III trial of chemoradiotherapy with temozolomide plus nivolumab or placebo for newly diagnosed glioblastoma with methylated MGMT promoter." Neuro Oncol **24**(11): 1935-1949.

Martin, M. (2011). "CUTADAPT removes adapter sequences from high-throughput sequencing reads." EMBnet.journal **17**.

Meazza, R., A. Comes, A. M. Orengo, S. Ferrini and R. S. Accolla (2003). "Tumor rejection by gene transfer of the MHC class II transactivator in murine mammary adenocarcinoma cells." Eur J Immunol **33**(5): 1183-1192.

Molnár, J., G. Szakács and G. E. Tusnády (2016). "Characterization of Disease-Associated Mutations in Human Transmembrane Proteins." PLoS One **11**(3): e0151760.

Mortara, L., P. Castellani, R. Meazza, G. Tosi, A. De Lerma Barbaro, F. A. Procopio, A. Comes, L. Zardi, S. Ferrini and R. S. Accolla (2006). "CIITA-induced MHC class II expression in mammary adenocarcinoma leads to a Th1 polarization of the tumor microenvironment, tumor rejection, and specific antitumor memory." Clin Cancer Res **12**(11 Pt 1): 3435-3443.

Narita, Y., Y. Arakawa, F. Yamasaki, R. Nishikawa, T. Aoki, M. Kanamori, M. Nagane, T. Kumabe, Y. Hirose, T. Ichikawa, H. Kobayashi, T. Fujimaki, H. Goto, H. Takeshima, T. Ueba, H. Abe, T. Tamiya, Y. Sonoda, A. Natsume, T. Kakuma, Y. Sugita, N. Komatsu, A. Yamada, T. Sasada, S. Matsueda, S. Shichijo, K. Itoh and M. Terasaki (2019). "A randomized, double-blind, phase III trial of personalized peptide vaccination for recurrent glioblastoma." Neuro Oncol **21**(3): 348-359.

Nemerow, G. R. and P. L. Stewart (1999). "Role of alpha(v) integrins in adenovirus cell entry and gene delivery." Microbiol Mol Biol Rev **63**(3): 725-734.

Niclou, S. P., C. Danzeisen, H. P. Eikesdal, H. Wiig, N. H. C. Brons, A. M. F. Poli, A. Svendsen, A. Torsvik, P. Ø. Enger, J. A. Terzis and R. Bjerkvig (2008). "A novel eGFP-expressing immunodeficient mouse model to study tumor-host interactions." The FASEB Journal **22**(9): 3120-3128.

Omuro, A., A. A. Brandes, A. F. Carpentier, A. Idbaih, D. A. Reardon, T. Cloughesy, A. Sumrall, J. Baehring, M. van den Bent, O. Bähr, G. Lombardi, P. Mulholland, G. Tabatabai, U. Lassen, J. M. Sepulveda, M. Khasraw, E. Vauleon, Y. Muragaki, A. M. Di Giacomo, N. Butowski, P. Roth, X. Qian, A. Z. Fu, Y. Liu, V. Potter, A. G. Chalamandaris, K. Tatsuoka, M. Lim and M. Weller (2023). "Radiotherapy combined with nivolumab or temozolomide for newly diagnosed glioblastoma with unmethylated MGMT promoter: An international randomized phase III trial." Neuro Oncol **25**(1): 123-134.

Ostrom, Q. T., G. Cioffi, K. Waite, C. Kruchko and J. S. Barnholtz-Sloan (2021). "CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2014-2018." Neuro Oncol **23**(12 Suppl 2): iii1-iii105.

Oudin, A., V. Baus, V. Barthelemy, C. Fabian, E. Klein, M. Dieterle, M. Wantz, A. C. Hau, C. Dording, A. Bernard, A. Michelucci, Y. A. Yabo, G. Kanli, O. Keunen, R. Bjerkvig, S. P. Niclou and A. Golebiewska (2021). "Protocol for derivation of organoids and patient-derived orthotopic xenografts from glioma patient tumors." STAR Protoc **2**(2): 100534.

Palma, M. L., P. Duangkhae, B. Douradinha, I. F. T. Viana, P. O. Rigato, R. Dhalia, R. B. Mailliard, S. M. Barratt-Boyes, E. J. M. Nascimento, T. M. Oshiro, A. J. da Silva Duarte and E. T. A. Marques (2017). "Development of potent class II transactivator gene delivery systems capable of inducing de novo MHC II expression in human cells, in vitro and ex vivo." Gene Ther **24**(6): 342-352.

Pires-Afonso, Y., S. P. Niclou and A. Michelucci (2020). "Revealing and Harnessing Tumour-Associated
Microglia/Macrophage Heterogeneity in Glioblastoma." Int J Mol Sci **21**(3).

Porter, K. A., L. N. Kelley, A. George, J. A. Harton and K. M. Duus (2010). "Class II Transactivator (CIITA)
Enhances Cytoplasmic Processing of HIV-1 Pr55Gag." PLOS ONE **5**(6): e11304.

Quail, D. F. and J. A. Joyce (2017). "The Microenvironmental Landscape of Brain Tumors." Cancer Cell **31**(3):
326-341.

Rainov, N. G. (2000). "A phase III clinical evaluation of herpes simplex virus type 1 thymidine kinase and
ganciclovir gene therapy as an adjuvant to surgical resection and radiation in adults with previously
untreated glioblastoma multiforme." Hum Gene Ther **11**(17): 2389-2401.

Reardon, D. A., A. A. Brandes, A. Omuro, P. Mulholland, M. Lim, A. Wick, J. Baehring, M. S. Ahluwalia, P.
Roth, O. Bahr, S. Phuphanich, J. M. Sepulveda, P. De Souza, S. Sahebjam, M. Carleton, K. Tatsuoka, C. Taitt,
R. Zwirtes, J. Sampson and M. Weller (2020). "Effect of Nivolumab vs Bevacizumab in Patients With
Recurrent Glioblastoma: The CheckMate 143 Phase 3 Randomized Clinical Trial." JAMA Oncol **6**(7): 1003-
1010.

Ruiz-Moreno, C., S. M. Salas, E. Samuelsson, S. Brandner, M. E. G. Kranendonk, M. Nilsson and H. G.
Stunnenberg (2022). "Harmonized single-cell landscape, intercellular crosstalk and tumor architecture of
glioblastoma." bioRxiv: 2022.2008.2027.505439.

Sartoris, S., M. T. Valle, A. L. Barbaro, G. Tosi, T. Cestari, A. D'Agostino, A. M. Megiovanni, F. Manca and R. S.
Accolla (1998). "HLA class II expression in uninducible hepatocarcinoma cells after transfection of AIR-1
gene product CIITA: acquisition of antigen processing and presentation capacity." J Immunol **161**(2): 814-
820.

Tang, Z., B. Kang, C. Li, T. Chen and Z. Zhang (2019). "GEPIA2: an enhanced web server for large-scale
expression profiling and interactive analysis." Nucleic Acids Research **47**(W1): W556-W560.

Thompson, J. A., S. K. Dissanayake, B. R. Ksander, K. L. Knutson, M. L. Disis and S. Ostrand-Rosenberg (2006).
"Tumor cells transduced with the MHC class II Transactivator and CD80 activate tumor-specific CD4+ T cells
whether or not they are silenced for invariant chain." Cancer Res **66**(2): 1147-1154.

Towey, M. and A. P. Kelly (2002). "Nuclear localisation of CIITA is controlled by a carboxy terminal leucine-
rich repeat region." Molecular Immunology **38**(8): 627-634.

Weller, M., N. Butowski, D. D. Tran, L. D. Recht, M. Lim, H. Hirte, L. Ashby, L. Mechtler, S. A. Goldlust, F.
Iwamoto, J. Drappatz, D. M. O'Rourke, M. Wong, M. G. Hamilton, G. Finocchiaro, J. Perry, W. Wick, J. Green,
Y. He, C. D. Turner, M. J. Yellin, T. Keler, T. A. Davis, R. Stupp and J. H. Sampson (2017). "Rindopepimut with
temozolomide for patients with newly diagnosed, EGFRvIII-expressing glioblastoma (ACT IV): a randomised,
double-blind, international phase 3 trial." Lancet Oncol **18**(10): 1373-1385.

Woroniecka, K., P. Chongsathidkiet, K. Rhodin, H. Kemeny, C. Dechant, S. H. Farber, A. A. Elsamadicy, X. Cui,
S. Koyama, C. Jackson, L. J. Hansen, T. M. Johanns, L. Sanchez-Perez, V. Chandramohan, Y. A. Yu, D. D.
Bigner, A. Giles, P. Healy, G. Dranoff, K. J. Weinhold, G. P. Dunn and P. E. Fecci (2018). "T-Cell Exhaustion
Signatures Vary with Tumor Type and Are Severe in Glioblastoma." Clin Cancer Res **24**(17): 4175-4186.

Zheng, C., C. M. Goldsmith, B. C. O'Connell and B. J. Baum (2000). "Adenoviral Vector Cytotoxicity Depends
in Part on the Transgene Encoded." Biochemical and Biophysical Research Communications **274**(3): 767-
771.

Zhou, G., R. Lieshout, G. S. van Tienderen, V. de Ruiter, M. E. van Royen, P. P. C. Boor, L. Magré, J. Desai, K.
Köten, Y. Y. Kan, Z. Ge, L. Campos Carrascosa, C. Geuijen, D. Sprengers, L. J. W. van der Laan, M. M. A.
Verstegen and J. Kwekkeboom (2022). "Modelling immune cytotoxicity for cholangiocarcinoma with
tumour-derived organoids and effector T cells." British Journal of Cancer **127**(4): 649-660.

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 4

Figure 5

Supplementary Figure 5

Table S1. Molecular profiling of the PDOX models used in the study.

		Model		
		P3	**T16**	**T188**
Matched Patient information	Patient code	BER0003	LIH0016	LIH0188
	Age at collection	64	52	68
	Sex	Male	Female	Male
	Tumor localization	N/A	Fronto-parietal right	N/A
	Surgery	N/A	Recurrence	Primary
	Treatment received prior PDOX	N/A	TMZ	No prior treatment
	Histological diagnosis	GBM Grade IV	GBM Grade IV	GBM Grade IV
PDOX characteristics	Original stain of mouse	NOD.CB17-Prkdc^{scid}/J	NOD.CB17-Prkdc^{scid}/J	NOD.CB17-Prkdc^{scid}/J
	Site of implantation	Brain	Brain	Brain
	Implantation	Organoids	Organoids	Organoids
	PDOX survival time (days)	42.5 +/- 3.6	75 +/- 6.5	64 +/- 10
	Proliferation index in PDOX (% Ki67 +/- SD)	54.7 +/- 3.7	18.9 +/- 5.8	37.1+/- 2.9
	Ploidy PDOX	2.1N	(2.1N) + 3.4N	2.3N
Molecular features in PDOXs	Copy number variations (CNV)	+[Chr7, 19p, 20q], -[1p36-1p34.1, 1q21.1-q44, -5p15.33-31, Chr9, Chr10, 11p15-14, 20p] --CDKN2A/B	++[EGFR, MDM4], +7q, - [6q, Chr10, 13q11-q31.1], --CDKN2A/B	[++EGFR], +[Chr7, Chr 19, Chr 20], -[1p36.23, 6p21.32, 6q27, Chr 10, 11q24.2], --CDKN2A/B
	Glioma-specific gene mutations	PTEN (chr10:g.87957923del, ENSP00000361021.3:p.Asp236ThrfsTer20) TP53 (chr17:g.7675085C>T, ENSP00000269305.4:p.Cys176Tyr)	EGFR ex25-27del	EGFR ex-6-7del, ex25-26del, MDM4 (chr1:g.204549329A>C, ENSP00000356150.3:p.Lys374Gln) PTCH1 (chr9:g.95459693C>T, ENSP00000332353.6:p.Val932Ile)
	DNA methylation profiles	GBM, IDHwt, RTK II	GBM, IDHwt, RTK II	GBM, IDHwt, RTK II/I
	MGMT promoter methylation	Methylated	Unmethylated	Unmethylated
	Verhaak glioma subtypes classification	Classic-like	Classical	Classic-like

Table S2. Composition of PBMC and isolated sub-populations before co-culture (shown as % of CD45+).

CR014	% CD14+	% CD19+	% CD3+	% CD4+	% CD8+	% CD56+
PBMC	29,9 ± 0,64	3,72 ± 0,28	49 ± 1,56	38,2 ± 1,41	8,68 ± 0,25	5,5 ± 0,06
Isolated CD3+	0,39 ± 0,06	0,12 ± 0,01	91,33 ± 1,23	70,73 ± 1,39	16,47 ± 0,14	0,2 ± 0,05
Isolated CD4+	0,02 ± 0,01	0,01 ± 0	98,2 ± 0,58	96,37 ± 1,02	0,02 ± 0,01	0,01 ± 0
Isolated CD8+	0,04 ± 0,02	0,01 ± 0	92,47 ± 1,93	0,48 ± 0,37	84,67 ± 3,84	0,04 ± 0,01

CR015	% CD14+	% CD19+	% CD3+	% CD4+	% CD8+	% CD56+
PBMC	20,2 ± 1,04	3,6 ± 0,33	61,33 ± 4,86	41,67 ± 2,26	18,03 ± 2,34	6 ± 0,03
Isolated CD3+	0,295 ± 0,1	0,049 ± 0,01	93,8 ± 1,83	66,53 ± 1,09	24 ± 3,07	0,42 ± 0,09
Isolated CD4+	0,02 ± 0,01	0,01 ± 0	99,27 ± 0,15	96,9 ± 0,41	0,61 ± 0,47	0,01 ± 0
Isolated CD8+	0,1 ± 0,05	0,03 ± 0,01	96,63 ± 1,16	1,23 ± 1,59	90,07 ± 3,64	0,02 ± 0,01

CR017	% CD14+	% CD19+	% CD3+	% CD4+	% CD8+	% CD56+
PBMC	6.85	7.2	61.8	29.5	29.8	3.05
Isolated CD3+	0.03	0.03	90.40	43.90	42.7	0.086
Isolated CD4+	0.01	0.03	98.80	92.30	4.17	0.00168
Isolated CD8+	0.01	0.01	96.30	0.01	94.2	0.011

CTL500	% CD14+	% CD19+	% CD3+	% CD4+	% CD8+	% CD56+
PBMCs	27.3	3.65	52.4	28.8	18.7	7.51
Isolated CD3+	0.11	0.034	93.7	50.6	34.6	1.44

CTL573	% CD14+	% CD19+	% CD3+	% CD4+	% CD8+	% CD56+
PBMC	21.6	6.29	61.5	33.9	24.5	4.53
Isolated CD3+	0.66	0.18	92	51.2	35.2	2.49

CTL640	% CD14+	% CD19+	% CD3+	% CD4+	% CD8+	% CD56+
PBMC	14.9	3.38	56.2	38.9	14.7	14.1
Isolated CD3+	0.14	1.50E-01	86.4	59.8	21.8	6.92

Table S3. Tumor/Immune cells typing at MHC-I and MHC-II loci.

P3 tumor

		Tumor cells	Immune cells					
		P3	CTL573	CTL500	CTL640	CR014	CR015	CR017
MHC-I	HLA-A	2:01:01	2:01	1:01	2:01	N/A	N/A	N/A
		~	3:01	2:01	29:02:00	N/A	N/A	N/A
	HLA-B	27:05:02G	7:02	44:02:00	39:01:00	N/A	N/A	N/A
		44:02:01G	44:02:00	57:01:00	44:02:00	N/A	N/A	N/A
	HLA-C	1:02:01	5:01	5:01	5:01	N/A	N/A	N/A
		5:01:01	7:02	6:02	12:03	N/A	N/A	N/A
MHC-II	HLA-DRB1	4:01:01	4:01	4:01	4:01	N/A	N/A	N/A
		4:08:01	15:01	13:05	11:01	N/A	N/A	N/A
	HLA-DQB1	3:01:01	3:01	3:01	3:01	N/A	N/A	N/A
		~	6:02	3:01	~	N/A	N/A	N/A
	HLA-DPB1	2:01:02	4:01	04:01G	4:01	N/A	N/A	N/A
		4:01:01	11:01	04:01G	23:01	N/A	N/A	N/A
	HLA-DQA1	3:03:01	1:02	not tested	3:01	N/A	N/A	N/A
		~	3:01	not tested	5:01	N/A	N/A	N/A
	HLA-DRB3/4/5	DRB4*01:03:01	DRB4*01:01	DRB3*02:02	DRB3*02:02	N/A	N/A	N/A
		~	DRB5*01:01	DRB4*01:03	DRB4*01:01	N/A	N/A	N/A
	HLA-DPA1	1:03:01	1:03	1:03	1:03	N/A	N/A	N/A
		~	2:01	1:03	~	N/A	N/A	N/A
	Match		7/9	8/9	7/9	0/9	0/9	0/9

T16 tumor

		Tumor cells	Immune cells			
		T16	CTL572	CR014	CR015	CR017
MHC-I	HLA-A	2:01:01	2:01	N/A	N/A	N/A
		68:02:01	3:01:00	N/A	N/A	N/A
	HLA-B	8:01:01	7:02:00	N/A	N/A	N/A
		35:01:01G	8:01	N/A	N/A	N/A
	HLA-C	4:01:01	7:01	N/A	N/A	N/A
		7:01:01	7:02	N/A	N/A	N/A
MHC-II	HLA-DRB1	03:01:01G	3:01	N/A	N/A	N/A
		11:01:01G	15:01	N/A	N/A	N/A
	HLA-DQB1	2:01:01	2:01	N/A	N/A	N/A
		3:01:01	6:02	N/A	N/A	N/A
	HLA-DPB1	1:01:01	2:01	N/A	N/A	N/A
		4:01:01	4:01	N/A	N/A	N/A
	HLA-DQA1	5:01:01	1:02	N/A	N/A	N/A
		5:05:01	5:01	N/A	N/A	N/A
	HLA-DRB3/4/5	DRB3*01:01:02G	DRB3 01:01	N/A	N/A	N/A
		DRB3*02:02:01	DRB5 01:01	N/A	N/A	N/A
	HLA-DPA1	1:03:01	1:03	N/A	N/A	N/A
		2:01:02	~	N/A	N/A	N/A
	Match		9/9	0/9	0/9	0/9

T188 tumor		Tumor cells	Immune cells		
		T188	CR014	CR015	CR017
MHC-I	HLA-A	2:01:01	N/A	N/A	N/A
		3:01:01	N/A	N/A	N/A
	HLA-B	7:02:01	N/A	N/A	N/A
		15:01:01G	N/A	N/A	N/A
	HLA-C	1:02:01	N/A	N/A	N/A
		7:02:01	N/A	N/A	N/A
MHC-II	HLA-DRB1	15:01:01	N/A	N/A	N/A
		13:01:01	N/A	N/A	N/A
	HLA-DQB1	6:02:01	N/A	N/A	N/A
		6:03:01	N/A	N/A	N/A
	HLA-DPB1	4:01:01	N/A	N/A	N/A
		13:01:01G	N/A	N/A	N/A
	HLA-DQA1	1:02:01	N/A	N/A	N/A
		1:03:01	N/A	N/A	N/A
	HLA-DRB3/4/5	DRB3*02:02:01	N/A	N/A	N/A
		DRB5*01:01:01	N/A	N/A	N/A
	HLA-DPA1	1:03:01	N/A	N/A	N/A
		2:01:04	N/A	N/A	N/A
	Match		0/9	0/9	0/9

~   Homozygous

G   ambigous allele typing

Table S4. List of all antibodies used in the study.

Antibody	Fluorochrome	Clone	Application	Dilution	Incubation	Vendor	Cat. No.
CIITA	-	7-1H	WB/IF	1:200	O/N at 4°C	Santa Cruz	sc-13556
Alexa Fluor 488	-	Polyclonal	IF	1:10000	1 hour at RT	ThermoFisher	A11001
CD107a	BV785	H4A3	FACS	1:20	30 min at 4°C	BioLegend	328644
CD11b	BV785	ICRF44	FACS	1:20	30 min at 4°C	Biolegend	301346
CD14	PE-Cy7	63D3	FACS	1:20	30 min at 4°C	Biolegend	367112
CD19	BB770	SJ25C1	FACS	1:20	30 min at 4°C	BD Biosciences	566396
CD3	BV510	UCHT1	FACS	1:20	30 min at 4°C	BioLegend	300448
CD4	PE-Cy5	RPA-T4	FACS	1:166	30 min at 4°C	BioLegend	300510
CD45	FITC	2D1	FACS	1:20	30 min at 4°C	BioLegend	368508
CD56	PE-Dazzle CF594	NCAM16.2	FACS	1:166	30 min at 4°C	BD Biosciences	564849
CD8	APC	RPA-T8	FACS	1:60	30 min at 4°C	BioLegend	301014
CD80	PE	L307.4	FACS	1:5	30 min at 4°C	BD Biosciences	560925
CD86	BV421	2331 (FUN-1)	FACS	1:20	30 min at 4°C	BD Biosciences	562433
Fas Ligand/TNFSF6	-	100419	Neutralization	-	-	R&D systems	MAB126
GAPDH	-	D16H11	WB	1:10000	O/N at 4°C	Cell Signalling	5174S
HLA-A,B,C	BV510	W6/32	FACS	1:20	30 min at 4°C	BioLegend	311435
HLA-DR	-	EPR3692	IHC	1:200	3 hour at RT	Abcam	ab92511
HLA-DR,DP,DQ	FITC	Tu39	FACS	1:5	30 min at 4°C	BD Biosciences	555558
HLA-DRα	-	G-7	WB	1:100	O/N at 4°C	Santa Cruz	sc-55593
Hoechst 33258	-	-	IF	1:10000	1 hour at RT	ThermoFisher	H3569
HRP secondary	-	Polyclonal	WB	1:10000	1 hour at RT	Jackson ImmunoResearch	111-035-003
I-A/I-E	PerCP/Cy5.5	M5/114.15.2	FACS	1:40	30 min at 4°C	BioLegend	107626
Lamin B1	-	Polyclonal	WB	1:10000	O/N at 4°C	Abcam	ab16048
Live/Dead fixable Near-IR	APC-Cy7	-	FACS	1:1000	30 min at 4°C	Invitrogen	L34975
TNF RI/TNFRSF1A	-	16803	Neutralization	-	-	R&D systems	MAB225