Title: Dimensions of early life adversity are differentially associated with patterns of delayed and

accelerated brain maturation

Authors: Dani Beck[1,2,3,*], Lucy Whitmore[4], Niamh MacSweeney[1,2,3], Alexis Brieant[5], Valerie Karl[1,2],

Ann-Marie G. de Lange[2,6,7], Lars T. Westlye[2,8,9], Kathryn L. Mills[4], Christian K. Tamnes[1,2,3]

[1] PROMENTA Research Center, Department of Psychology, University of Oslo, Oslo, Norway

[2] NORMENT, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical

Medicine, University of Oslo, Oslo, Norway

[3] Division of Mental Health and Substance Abuse, Diakonhjemmet Hospital, Oslo, Norway

[4] Department of Psychology, University of Oregon, Eugene, OR, USA

[5] Department of Psychological Science, University of Vermont, Vermont, USA

[6] Department of Clinical Neurosciences, CHUV and University of Lausanne, Lausanne, Switzerland

[7] Department of Psychiatry, University of Oxford, Oxford, UK

[8] Department of Psychology, University of Oslo, Oslo, Norway

[9] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Oslo, Norway

23 [*] Correspondence: Dani Beck (dani.beck@psykologi.uio.no)

Abstract

Background

Different types of early-life adversity have been associated with children's brain structure and

function. However, understanding the disparate influence of distinct adversity exposures on

the developing brain remains a major challenge.

Methods

This study investigates the neural correlates of 10 robust dimensions of early-life adversity

identified through exploratory factor analysis in a large community sample of youth from the

Adolescent Brain Cognitive Development (ABCD) Study. Brain age models were trained,

validated, and tested separately on T1-weighted (T1; N = 9524), diffusion tensor (DTI; N =

8834), and resting-state functional (rs-fMRI; N = 8233) magnetic resonance imaging (MRI)

data from two time points (mean age = 10.7 years, SD = 1.2, range = 8.9-13.8 years).

Results

Bayesian multilevel modelling supported distinct associations between different types of

early-life adversity exposures and younger- and older-looking brains. Dimensions generally

related to emotional neglect, such as lack of primary and secondary caregiver support, and

lack of caregiver supervision, were associated with lower brain age gaps (BAGs), i.e.,

younger-looking brains. In contrast, dimensions generally related to caregiver

psychopathology, trauma exposure, family aggression, substance use and separation from

biological parent, and socio-economic disadvantage and neighbourhood safety were

associated with higher BAGs, i.e., older-looking brains.

Conclusions

The findings suggest that dimensions of early-life adversity are differentially associated with

distinct neurodevelopmental patterns, indicative of dimension-specific delayed and

accelerated brain maturation.

Keywords: ABCD Study, Adolescence, Brain age, Development, Early-life Adversity, MRI

1. Introduction

Early-life adversity (ELA) such as exposure to abuse, violence, neglect, separation from caregivers, and chronic poverty, among others, can have widespread effects on youth neurodevelopment (1) and increase risk for mental disorders (2,3). Previous studies investigating how ELA influences neural development have adopted varied theoretical frameworks, reflecting the complex and multifaceted nature of the field (4,5). Traditionally, research has focused on a single type of adversity; for example, reporting associations between heightened amygdala activation and early exposure to violence (6), and lower volumes of gray matter and low income (7) and exposure to institutionalisation (8). A critical limitation of this approach however, is the fact that most children are exposed to numerous types of adversities concurrently (9). The cumulative risk approach overcomes this by aggregating different forms of adversity into a singular risk factor. Research has found that early cumulative risk was prospectively associated with lower total gray matter volume, cortex volume, and right superior parietal and inferior parietal cortical thickness (10). While the cumulative risk approach is useful for identifying children at greatest risk for intervention and can thus serve as a vital public health tool, aggregating risk factors may obscure potentially diverging effects of different adverse experiences on the developing brain.

More nuanced perspectives that have emerged differentiate between adverse experiences related to threat versus deprivation (11), drawing support for the neural basis of this distinction from studies on fear learning and sensory deprivation. For children exposed to threat, studies have reported lower cortical thickness, surface area, volume of the amygdala and hippocampus (12–14) and ventro-medial prefrontal cortex (vmPFC) (15,16), in addition to reduced resting-state amygdala–vmPFC connectivity (17). The observed effects may be indicative of accelerated maturation and are consistent with life history and evolutionary-biology theories proposing accelerated development as an adaptation to harsh or stressful environments (4,18–20). Similarly, another conceptual model, the stress acceleration hypothesis, argues that adversity may expedite neural development as a means of compensating for the absence of species-expectant maternal buffering of emotional reactivity (21).

For children exposed to deprivation, such as institutional rearing, lack of social and cognitive stimulation, and other forms of parental absence, research has revealed altered structure and function in the frontoparietal network, amygdala-hippocampal-PFC connectivity (11,22), and reductions in cortical gray matter and total brain volume, in addition to widespread cortical thinning (8,23,24). Moreover, electroencephalogram (EEG)

studies have shown associations between spectral profiles indicative of delayed patterns of

functional and cortical maturation and neglect (25), poverty (26), and parental stress (27).

While previous research has contributed to an understanding of how different features

of ELA are associated with unique brain outcomes (28), real-world occurrences of adversity

are multifaceted and often co-occur in a complex manner, making it a considerable challenge

to precisely account for the heterogeneity in ELA. To address this, several data-driven

methods have been applied (29–33), albeit on small or homogenous samples. Recent research

by Brieant et al. (2023) capitalised on big data from the Adolescent Brain Cognitive

Development (ABCD) Study and identified 10 dimensions of adversity co-occurrence

pertaining to conceptual domains reflecting 1) caregiver psychopathology, 2) socioeconomic

disadvantage and lack of neighbourhood safety, 3) secondary caregiver lack of support, 4)

primary caregiver lack of support, 5) child report of family conflict, 6) caregiver substance

use and biological parent separation, 7) family anger and arguments, 8) family aggression, 9)

trauma exposure, and 10) caregiver lack of supervision. Yet, to date, the neural correlates of

these dimensions of ELA have not been investigated.

Brain age prediction offers a framework that can combine multiple imaging features,

alleviating the need for selection of specific metrics or regions, and yields an individualised

surrogate marker of brain maturation (35). Brain age involves estimation of biological age

based on brain MRI characteristics, which may differ from an individual's chronological age

(36). This difference, termed the brain age gap (BAG), could reflect deviation from typical

neurodevelopmental patterns, and has been validated in several neurodevelopmental studies

(37–40). Previous literature has also validated the stability of brain age models across early

adolescence, with evidence of BAG scores tracking with metrics of maturation (40). Studies

have also linked lower BAG in youth to attention-deficit hyperactivity disorder (ADHD),

lower socio-economic status, higher anxiety and depression, as well as greater general

psychopathology symptom severity (37,41–44). In the context of ELA, where different

dimensions of adversity may be associated with unique brain outcomes (4,28,45–49), brain

age can probe individualised markers of delayed or accelerated maturation by means of

younger- or older-looking brains.

To this end, using ABCD data, our primary aim was to test for associations between

MRI-based estimates of brain maturation and 10 previously characterised dimensions of ELA

co-occurrence (34). Based on theoretical accounts and the empirical studies reviewed above,

we hypothesised ELA dimensions of 1) caregiver psychopathology, 2) socioeconomic

disadvantage and lack of neighbourhood safety, 5) child report of family conflict, 6)

caregiver substance use and biological parent separation, 7) family anger and arguments, 8)

family aggression, and 9) trauma to be associated with older-looking brains and accelerated

maturation between the two time points. We hypothesised ELA exposures of 3) secondary

caregiver lack of support, 4) primary caregiver lack of support, and 10) caregiver lack of

supervision to be associated with younger-looking brains and delayed maturation over time.

2. Methods and Materials

2.1. Sample and ethical approval

The Adolescent Brain Cognitive Development (ABCD) Study [®] (50) comprises of children

and adolescents part of an ongoing longitudinal study. Participants were excluded using the

ABCD Study exclusion criteria listed elsewhere (51). Data used in the present study were

drawn from the ABCD curated annual release 5.0, containing data from baseline up until the

second-year visit (https://data-archive.nimh.nih.gov/abcd). All ABCD Study data is stored in

the NIMH Data Archive Collection #2573, which is available for registered and authorised

users (Request #7474, PI: Westlye). The 5.0 release will be permanently available as a

persistent dataset defined in the NDA Study 1299 and has been assigned the DOI

10.15154/8873-zj65. The Institutional Review Board (IRB) at the University of California,

San Diego, approved all aspects of the ABCD Study (52). Parents or guardians provided

written consent, while the child provided written assent. The current study was conducted in

line with the Declaration of Helsinki and was approved by the Norwegian Regional

Committee for Medical and Health Research Ethics (REK 2019/943).

2.2. Demographic information and data quality assurance

The initial sample consisted of ~11,800 participants (52% male) at mean age 10.75 (SD =

1.18, range 8.92-13.83) years, with baseline and two-year follow-up observations (obs) of T1

(obs = 19,048), DTI (obs = 17,672), and rs-fMRI (obs = 16,495) data. Quality control

procedures followed a standard protocol described in Hagler et al. (2019). Briefly,

participants with excessive head motion or poor data quality were excluded from the curated

data release by the ABCD Study team. Additional quality assurance was carried out

following extraction of data using the recommendations for data cleaning provided by the

ABCD Study team (using data structure abcd_imgincl01). Following quality assurance, the

final sample included T1 obs of 19,047, DTI obs of 17,668, and rs-fMRI obs of 16,466 used

for the current study. Demographic information for each brain MRI modality-specific sample

can be found in SI Table 1, while the T1 sample is illustrated in Figure 1.

Figure 1. Age distribution split by sex and time point. Time point (1 and 2) represent baseline and
two-year-follow-up data from the ABCD Study cohort. Female data in teal blue; male in lavender.
Dotted gray line is drawn between mean age at time point 1 and 2.

2.3. MRI acquisition, processing, and segmentation

Neuroimaging data were acquired at 21 different sites (using 31 scanners) and processed by
the ABCD Study team. A 3-T Siemens Prisma, General Electric 750 or Phillips scanner was
used for data acquisition. Protocols used for data acquisition and processing are described in
detail elsewhere (50,53) and available in SI Section 1. Three modalities of brain structural
and functional measures were used in the present study: structural grey matter measures (T1),
diffusion white matter microstructural measures (DTI), and resting-state functional
connectivity measures (rs-fMRI) (53). Cortical surface reconstruction and subcortical
segmentation was performed with FreeSurfer v7.1.1 (54,55). White matter microstructural
measures were generated using AtlasTrack, a probabilistic atlas-based method for automated
segmentation of white matter fibre tracts (Hagler et al., 2009). Measures of functional
connectivity were computed using a seed-based, correlational approach (57), where average
time courses were calculated for cortical surface-based ROIs using a functionally-defined
parcellation based on resting-state functional connectivity patterns (58) and subcortical ROIs

(55). A detailed description of the processing and extraction of brain imaging data used is

provided in full in SI Section 1.

Briefly, For T1, we extracted tabulated total and regional measures of cortical surface area,

thickness, volume, sulcal depth, intensity-gray-white contrast, and subcortical volume (397

measures). For DTI, full and inner (multi) shell tissue properties including functional

anisotropy (FA) and mean (MD), longitudinal (or axial, AD), and transverse (or radial, RD)

diffusivity were extracted for total and regional features (576 measures). For rs-fMRI,

functional connectivity within and between parcellations from Gordon network, including

subcortical data (58) were extracted (416 measures). Following procedures of quality

assurance (Section 2.2), harmonisation of multi-scanner effects was carried out using

longCombat (59) (see SI Section 2 and SI Figures 1-3).

2.4. Brain age prediction

Brain age prediction was carried out using the eXtreme Gradient Boosting (XGBoost)

regression model (60), a machine learning algorithm that a gradient boosting library by

combining multiple decision trees to create predictive models. Parameters were tuned using

ten-fold cross-validation, stratified by age. The models were fitted using the best estimators

and optimised models were applied to the (hold-out) test sample. R2, RMSE, and MAE were

calculated to evaluate prediction accuracy in the test set. For each brain modality (T1, DTI,

rs-fMRI), 50% of the data was used as the hold-out test sample and 50% was used for model

training and validation. Here, data was split ensuring an equal distribution of cross-sectional

and longitudinal data across training and testing samples, whereby no two datapoints from the

same individual (longitudinal) were separated.

215   Consistent with a recent brain age paper using the ABCD Study sample (61),

confounding effects from complex family-related factors were minimised using a group

shuffle split with family ID as the group indicator to ensure that no siblings were split across

training and test sets. A detailed overview of T1, DTI, and rs-fMRI training and test samples,

including demographic information are provided in SI Section 3, SI Table 1, and SI Figures

4-6. A complete list of all the extracted measures used for brain age prediction per modality

is provided in SI Tables 2-4. Feature importance scores for each model is provided in SI

Figures 7-9. To adjust for commonly observed age-bias (overestimated predictions for

younger participants and underestimated predictions for older participants) (62), we applied a

statistical correction as previously described in (63). The difference between an individual's

predicted brain age and their chronological age (BAG) was calculated by subtracting the

participants chronological age from the age bias corrected predicted age (BAG = predicted

age - chronological age) for each of the models, providing T1, DTI, and rs-fMRI-based BAG

values for all participants. The resulting BAG can be either positive or negative, indicating an

older- or younger-looking brain than the individual's actual age.

2.5. Early life adversity

The current study utilised 10 previously obtained factor scores from 60 measures of early-life

adversity (ELA) as detailed in (34). Briefly, Brieant and colleagues identified 139 potential

ELA items from the ABCD Study baseline measures, which encompass a spectrum of ELA

constructs including caregiving disruption, caregiver psychopathology, maltreatment,

neighbourhood safety/violence, among others. These variables were sourced from child and

parent reports, as well as researcher assessments, and originated from modified versions of

validated scales. To identify dimensions, 60 ELA variables which were binary, polytomous,

and continuous in nature were entered into an exploratory factor analysis (EFA) conducted in

Mplus version 8.7 (64), resulting in 10 dimensions (F1 to F10; see SI Table 5 and SI Figure

10 for correlation matrix). To obtain factor scores, an exploratory structural equation model

was carried out specifying the number of factors identified in the EFA. Further details of the

variable selection process, identification of ELA dimensions, and calculation of factor

loadings can be found in Brieant et al. (2023).

2.6. Statistical analysis

All analyses were carried out using R version 4.2.1 (65). To investigate the association (main

effect) between each ELA dimension (F1:F10) and deviation from expected age patterns (i.e.,

BAG), and whether the effect of each ELA dimension on BAG varies across time points

(interaction effect), Bayesian multilevel models were carried out using the *brms* (66,67) R-

package. Here, multivariate models are fitted in familiar syntax to comparable frequentist

approaches such as a linear mixed effects model using the *lme4* (68). We assessed the

relationship between each ELA dimension at baseline and residualised (age-bias corrected)

BAG, where modality-specific BAG (T1, DTI, rs-fMRI) was first entered as the dependent

variable, each ELA dimension (F1:F10) and interaction term (TP:F1-F10) between time point

(TP) and ELA dimensions (F1:F10) were separately entered as independent fixed effects

variables with sex entered as a covariate, and with subject ID as the random effect.

258   To prevent false positives and to regularise the estimated associations, we defined a

standard prior around zero with a standard deviation of 0.1 for all regression coefficients,

reflecting a baseline expectation of effects being small but allowing for sufficient flexibility

in estimation. Each model was run with 8000 iterations, including 4000 warmup iterations,

across four chains. This setup was chosen to ensure robust convergence and adequate

sampling from the posterior distributions. For each coefficient of interest, we report the mean

estimated value of the posterior distribution (b) and its 95% credible interval (the range of

values that with 95% confidence contains the true value of the association), and calculated

the Bayes Factor (BF) – provided as evidence ratios in the presented figures – using the

Savage-Dickey method (69). Briefly, BF can be interpreted as a measure of the strength of

evidence (*extreme, very strong, strong, moderate, anecdotal, none*) in favour of the null or

alternative hypothesis. For a pragmatic guide on BF interpretation, see SI Table 6.

3. Results

3.1. Descriptive statistics

Descriptive statistics can be found in SI Table 1. Table 1 summarises descriptive and model

validation statistics pertaining to each brain age prediction model, following age-correction.

Figure 2 shows predicted age as a function of chronological age for each age prediction

model. SI Figure 11 shows a correlation matrix including each modality-specific predicted

brain age and BAG.

Table 1. Average R^2, root mean square error (RMSE), mean absolute error (MAE), and Pearson's correlations between predicted and chronological age (*r*) for each brain age prediction model, with 95% confidence intervals.

	T1	**DTI**	**rs-fMRI**
r	0.59 [0.57-0.60]	0.66 [0.65-0.67]	0.41 [0.39-0.43]
R^2	0.34	0.43	0.17
RMSE	0.96	0.89	1.08
MAE	0.79	0.71	0.89

Figure 2. Predicted age as a function of age. Scatter plots demonstrating the correlation between

chronological and predicted ages for three brain imaging modalities: T1, DTI, and rs-fMRI. Each plot illustrates

the Pearson correlation coefficient (r) and statistical significance (*p*-value), with data points representing

individual samples at baseline and follow-up time points. The plots reveal varying degrees of performance

accuracy across modalities, with T1 and DTI models showing higher correlations compared to rs-fMRI.

3.2. Bayesian multilevel modelling

Bayesian multilevel modelling tested the association between each ELA dimension and

modality-specific BAG. The full results are available in SI Tables 7-9 and are visualised

below in Figure 3. For estimated credible intervals, see SI Figures 12 and 13.

3.2.1. T1 BAG relations

For T1 BAG, the test revealed evidence of a positive association between ELA dimension F2

(socioeconomic disadvantage and neighbourhood safety) and T1 BAG (BF < 0.01, β = 0.13),

indicating that this dimension was associated with older-looking brains. Further, the tests

revealed evidence of negative associations between ELA dimensions F3 (secondary caregiver

lack of support) (BF = 0.9, β = -0.05) and F4 (primary caregiver lack of support) (BF = 0.17,

β = -0.08) and T1 BAG, indicating that dimensions related to caregiver emotional neglect are

associated with younger-looking brains. In terms of interaction effects of time point, we

found a negative association between F10 (lack of supervision) and T1 BAG (BF = 0.08, β =

-0.09), indicating that unsupervised youth diverge more from normative age patterns

throughout the course of the study period.

3.2.2. DTI BAG relations

For DTI, the tests revealed evidence of a positive association between F2 and DTI BAG (BF

= 0.27, β = 0.06), aligning with the T1 BAG findings, and indicating that those living in more

Early life adversity and brain age in youth

disadvantaged and less safe environments may have older-looking brains. In terms of

interaction effects of time point, the tests revealed evidence supporting a positive association

between ELA dimension F1 (caregiver psychopathology) and DTI BAG (BF = 0.09, β =

0.07), indicating that the brain ages of these youth will diverge more (accelerate) from

normative age patterns over time.

3.2.3. rs-fMRI BAG relations

For rs-fMRI, the tests revealed evidence of a positive association between F2 and rs-fMRI

BAG (BF < 0.01, β = 0.18), aligning with findings from DTI and T1 BAG. The tests also

revealed positive associations between F1 (BF = 0.63, β = 0.07), F6 (caregiver substance

abuse and separation from biological parents) (BF = 0.03, β = 0.15), F8 (family aggression)

(BF = 0.66, β = 0.08), and F9 (trauma exposure) (BF = 0.07, β = 0.15) and rs-fMRI BAG.

These positive associations largely indicate that dimensions linked to a threatening

environment relate to older-looking brains. Further, the tests revealed a negative association

between F4 and rs-fMRI BAG (BF = 0.27, β = -0.10), in line with findings from T1 BAG. In

terms of interaction effects of time point, we found evidence supporting negative associations

between both F3 (BF = 0.62, β = -0.08) and F5 (family conflict) (BF = 0.58, β = -0.08) and

rs-fMRI BAG, and a positive association between F6 (BF = 0.51, β = 0.09) and rs-fMRI

BAG.

327
Figure 3. Associations between ELA dimensions and BAG for DTI, T1, and rs-fMRI. The figure shows
posterior distributions of the estimates of the standardised coefficient. Estimates for each ELA dimension on
BAG (main effect) on the left, and ELA dimension interaction effect of time point (TP:ELA) on BAG on the
right. Colour scale follows directionality of evidence, with positive (red) values indicating evidence in favour of
positive associations (greater adversity linked to older-looking brains) and negative (blue) values indicating
evidence in favour of negative associations (greater adversity linked to younger-looking brains) for each ELA
dimension. The width of the distributions represent the uncertainty of the parameter estimates. F1: Caregiver
psychopathology; F2: Socio-economic disadvantage and neighbourhood safety; F3: Secondary caregiver lack of
support; F4: Primary caregiver lack of support; F5: Youth report of family conflict; F6: Caregiver substance use
and separation from biological parent; F7: Family anger and arguments; F8: Family aggression; F9: Trauma
exposure; F10: Lack of supervision.

4. Discussion

Research indicates that approximately half of all children will experience at least one form of
adversity by the time they reach adulthood (2,70). Different forms of early-life adversity co-

occur and may uniquely impact child brain structure and function. This work sought to

delineate the potentially unique brain outcomes of 10 co-occurring dimensions of early-life

adversity in a longitudinal sample of 9–14-year-old youth. Our main findings indicate that

ELA dimensions of caregiver lack of support and supervision are associated with younger-

looking brains and that ELA dimensions of caregiver psychopathology, socioeconomic

disadvantage and neighbourhood safety, caregiver substance abuse and separation from

biological parents, family aggression, and trauma exposure are associated with older-looking

brains. Our findings largely align with and extend the current literature on the differential

impact of different co-occurring patterns of environmental exposures and adversity on brain

development (45,49).

4.1. Links to accelerated brain maturation

Families living in lower socioeconomic status neighbourhoods are exposed to more harms,

such as interpersonal violence (71), and are more likely to have concerns about

neighbourhood safety (72). Positive associations between the dimension representing

socioeconomic disadvantage and neighbourhood safety (F2) and BAG were found with all

three brain age models, and are consistent with research showing socioeconomic

disadvantage and neighbourhood violence linked to smaller cortical volumes and greater

cortical thinning (73–75). Positive associations between exposure to trauma (F9) and BAG is

also consistent with previous research showing that children exposed to trauma are more

likely to be misclassified as adults by means of older DNA methylation age compared to

chronological age, and earlier pubertal maturation (76–78).

For caregiver psychopathology (F1), we found both positive BAG associations as

main effects and as interaction effects of time, suggesting that parent psychopathology-

related deviations from expected age-patterns accelerate over time. Our findings are

challenging to interpret in the context of mixed results from previous research. ABCD Study

findings have revealed smaller volume in the right putamen (79), left hippocampus (80), and

bilateral hippocampi (81) in relation to parental psychopathology. These findings reflect age

patterns of hippocampi and putamen usually not seen until late adolescents and young

adulthood – whereby subcortical volumes are expected to reduce over time (82,83) –

suggesting the results may indicate accelerated ageing in this younger sample, and thus

aligning with the directionality of our results. However, studies using other datasets have

found that parental history of depression is associated with larger volume in the bilateral

amygdala (84), and others have reported no effects (85). ELA dimensions that indicated the

potential presence of household hostility including family aggression (e.g., throwing things,

hitting) (F8) revealed a positive main effect for rs-fMRI BAG, but no effects for family

arguments (e.g., expressing anger, fighting, raising voices) (F7), with the latter finding going

against our hypothesis.

Co-occurrence of caregiver substance abuse and separation from biological parents in

one dimension (F6) might reflect child custody issues related to caregiver substance use

disorders or arrests (34,86). Moreover, this dimension also has factor loadings from domestic

violence. Our results revealed positive rs-fMRI BAG associations both as a main effect and

an interaction effect of time. Drawing from concepts of stress acceleration, parental

deprivation accelerates the functional development of the mPFC in children, such that

amygdala–mPFC interactions are more adultlike following deprivation experiences (21).

Children in this group may have accelerated maturation as a means of adapting from a state

of parent-regulated to self-regulated emotional processing due to absent or inconsistent

parental care (4,19–21). Further research is needed to understand the underlying mechanisms

at play. In summary, dimensions of ELA co-occurrence related to older-looking brains

support research reporting accelerated brain maturation in children exposed to potentially

more hostile or dangerous environments.

4.2. Links to delayed brain maturation

Neurobiological studies of brain development have long assumed a deficit model in which

lack of input to the development of a child will result in delay of certain skills (87). In the

child brain, this may be reflected by a delay in pruning and thus larger brain volumes and

younger-looking brains. Our results support this, with lack of primary (F4) and secondary

(F3) caregiver support as well as lack of caregiver supervision (F10) all revealing negative

associations with T1 and rs-fMRI BAG, with main effects for F3 and F4, and interaction

effects for F3 and F10. Importantly, these factors each share elements of emotional neglect.

Previous research investigating more severe forms of neglect (physical and emotional)

have found that children reared in institutions demonstrate EEG patterns suggestive of a

delay in cortical maturation in frontal, temporal, and occipital regions (25). However, a

wealth of studies also report conflicting results indicative of advanced maturation in similar

samples (8,24). A caveat of research carried out on neglect is that children are often in

environments enriched for several co-occurring ELAs, making it difficult to rely on the

stability of the neglect scores while considering other stressors. Importantly, our results

specifically capture emotional forms of neglect in terms of lack of household emotional

support in a community with relatively lower levels of risk, which may explain divergence

from prior studies on extreme physical neglect or institutionalisation.

Lastly, a negative interaction effect of time for rs-fMRI BAG and family conflict (F5)

was also found, indicating that youth exposed to family conflict diverge more from normative

age patterns (i.e., delayed maturation) over time. This contradicted our hypothesis and

findings from previous research also utilising the ABCD Study cohort which reported high

family conflict associations with smaller cortical surface areas of the orbitofrontal cortex,

anterior cingulate cortex, and middle temporal gyrus (88). In summary, ELA dimensions

related to emotional neglect were associated with younger-looking brains and delayed

maturation.

4.3. Strengths and limitations

Our study adds new insight into the neural correlates of co-occurring ELA dimensions in a

large-scale longitudinal community sample. Using a sample not enriched for adversity

exposure has the added value of demonstrating that even less severe ELA exposure is

associated with changes in the developing brain, facilitating broader generalisation. However,

there is trade-off in that our findings cannot necessarily be generalised to interpret neural

correlates of children exposed to more severe forms of adversity, as these groups may not be

well-represented in the ABCD Study. Future research is required to address this without

losing adequate power. There remain also additional challenges such as accounting for

differences in chronicity of adversity events, interindividual differences in resilience, and

overlap in adversity types.

Building on data driven methods applied in Brieant et al. (2023), the current study

benefitted from ELA dimensions accounting for more variability in ELA patterns and a

dimensional structure replicated in an independent sample, suggesting some stability. The

current study utilised three estimates of brain age based on different MRI modalities in an

attempt to capture potentially tissue-specific effects of ELA dimensions. While the

multimodal (with longitudinal data) approach is a strength, deep learning methods have

shown greater accuracy in recent years (39,89). Moreover, focusing on different regional

rather than global brain metrics and their association with adversity dimensions represents an

opportunity for future research.

4.4. Conclusion

The current study supports notions that brain MRI outcomes related to early-life adversity are

differentially associated with accelerated and delayed brain maturation. Neurodevelopmental

processes influenced by experiences of trauma, parental psychopathology, socio-economic

disadvantage and neighbourhood safety, caregiver substance abuse and separation from

biological parents, and family aggression, are at least partially distinct from those influenced

by experiences of emotional neglect, with brain age deviations indicating differential

maturational patterns. Future research should build on this work by investigating, for

example, how brain age patterns mediate the associations between dimensions of early life

adversity and child behavioural and symptom measures. Such studies could elucidate whether

indices of brain maturation serve as an underlying biological mechanism linking early

adversity to later child outcomes, thereby offering a more comprehensive understanding of

these developmental processes and helping guide intervention strategies that aim to mitigate

the impact of early-life adversity and supporting healthy development of at-risk youth.

5. Funding

This work was supported by the Research Council of Norway (#223273, #288083, #323951,

#276082, #249795, #248238, #298646, #300767), the South-Eastern Norway Regional

Health Authority (#2019069, #2021070, #500189, #2014097, #2015073, #2016083,

#2018076, #2019101), KG Jebsen Stiftelsen, the European Union-funded Horizon Europe

project 'environMENTAL' (101057429); and the UK Research and Innovation (UKRI) under

the UK government's Horizon Europe (10041392 and 10038599), the Swiss National Science

Foundation (#PZ00P3_193658), and the European Research Council under the European

Union's Horizon 2020 Research and Innovation program (ERC StG, #802998 and RIA

#847776). The work was performed on the Service for Sensitive Data (TSD) platform, owned

by the University of Oslo, operated, and developed by the TSD service group at the

University of Oslo IT-Department (USIT). Computations were performed on resources

provided by Sigma2 - the National Infrastructure for High Performance Computing and Data

Storage in Norway.

6. Acknowledgments

Data used in the preparation of this article were obtained from the Adolescent Brain

Cognitive DevelopmentSM (ABCD) Study (https://abcdstudy.org), held in the NIMH Data

Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000

children aged 9-10 and follow them over 10 years into early adulthood. The ABCD Study® is

supported by the National Institutes of Health and additional federal partners under award

numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018,

U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117,

U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156,

U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093,

U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at

https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete

listing of the study investigators can be found at https://abcdstudy.org/consortium_members/.

ABCD consortium investigators designed and implemented the study and/or provided data

but did not necessarily participate in the analysis or writing of this report. This manuscript

reflects the views of the authors and may not reflect the opinions or views of the NIH or

ABCD consortium investigators. This manuscript has been posted on the preprint server

bioRxiv: https://www.biorxiv.org/content/10.1101/2024.01.22.576780v1. We thank the co-

authors of Brieant et al. (2023) for their contributions to the development of the factor scores

used in this paper (Anna Vannucci, Hajer Nakua, Jenny Harris, Jack Lovell, Divya

Brundavanam, Nim Tottenham, and Dylan Gee).

7. Financial disclosures

None of the authors report financial conflict or potential conflict of interest.

8. References

1. McLaughlin KA, Sheridan MA, Gold AL, Duys A, Lambert HK, Peverill M, *et al.* (2016):

Maltreatment Exposure, Brain Structure, and Fear Conditioning in Children and

Adolescents [no. 8]. *Neuropsychopharmacology* 41: 1956–1964.

2. Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, *et al.*

(2010): Childhood adversities and adult psychopathology in the WHO World Mental

Health Surveys. *Br J Psychiatry* 197: 378–385.

3. Wade M, Wright L, Finegold KE (2022): The effects of early life adversity on children's

mental health and cognitive functioning [no. 1]. *Transl Psychiatry* 12: 1–12.

4. Ellis BJ, Sheridan MA, Belsky J, McLaughlin KA (2022): Why and how does early

adversity influence development? Toward an integrated model of dimensions of

environmental experience. *Dev Psychopathol* 34: 447–471.

5. Smith KE, Pollak SD (2021): Rethinking Concepts and Categories for Understanding the

Neurodevelopmental Effects of Childhood Adversity. *Perspect Psychol Sci* 16: 67–

93.

6. White SF, Voss JL, Chiang JJ, Wang L, McLaughlin KA, Miller GE (2019): Exposure to

violence and low family income are associated with heightened amygdala

responsiveness to threat among adolescents. *Dev Cogn Neurosci* 40: 100709.

7. Hanson JL, Hair N, Shen DG, Shi F, Gilmore JH, Wolfe BL, Pollak SD (2013): Family

Poverty Affects the Rate of Human Infant Brain Growth. *PLOS ONE* 8: e80954.

8. Sheridan MA, Fox NA, Zeanah CH, McLaughlin KA, Nelson CA (2012): Variation in

neural development as a result of exposure to institutionalization early in childhood.

Proc Natl Acad Sci U S A 109: 12927–12932.

9. McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, Kessler RC (2012):

Childhood adversities and first onset of psychiatric disorders in a national sample of

adolescents. *Arch Gen Psychiatry* 69: 1151–1160.

10. Chad-Friedman E, Botdorf M, Riggins T, Dougherty LR (2021): Early childhood

cumulative risk is associated with decreased global brain measures, cortical thickness,

and cognitive functioning in school-age children. *Dev Psychobiol* 63: 192–205.

11. Sheridan MA, McLaughlin KA (2014): Dimensions of early experience and neural

development: deprivation and threat. *Trends Cogn Sci* 18: 580–585.

12. Colich NL, Platt JM, Keyes KM, Sumner JA, Allen NB, McLaughlin KA (2020): Earlier

age at menarche as a transdiagnostic mechanism linking childhood trauma with

multiple forms of psychopathology in adolescent girls. *Psychol Med* 50: 1090–1098.

13. Gold AL, Sheridan MA, Peverill M, Busso DS, Lambert HK, Alves S, *et al.* (2016):

Childhood abuse and reduced cortical thickness in brain regions involved in

emotional processing. *J Child Psychol Psychiatry* 57: 1154–1164.

14. Lim L, Radua J, Rubia K (2014): Gray Matter Abnormalities in Childhood Maltreatment:

A Voxel-Wise Meta-Analysis. *Am J Psychiatry* 171: 854–863.

15. De Brito SA, Viding E, Sebastian CL, Kelly PA, Mechelli A, Maris H, McCrory EJ

(2013): Reduced orbitofrontal and temporal grey matter in a community sample of

maltreated children. *J Child Psychol Psychiatry* 54: 105–112.

16. Hanson JL, Chung MK, Avants BB, Shirtcliff EA, Gee JC, Davidson RJ, Pollak SD

(2010): Early stress is associated with alterations in the orbitofrontal cortex: A tensor-

based morphometry investigation of brain structure and behavioral risk. *J Neurosci*

30: 7466–7472.

17. Herringa RJ, Birn RM, Ruttle PL, Burghy CA, Stodola DE, Davidson RJ, Essex MJ

(2013): Childhood maltreatment is associated with altered fear circuitry and increased

internalizing symptoms by late adolescence. *Proc Natl Acad Sci U S A* 110: 19119–

19124.

18. Belsky J (2012): The Development of Human Reproductive Strategies: Progress and

Prospects. *Curr Dir Psychol Sci* 21: 310–316.

19. Frankenhuis WE, de Weerth C (2013): Does Early-Life Exposure to Stress Shape or

Impair Cognition? *Curr Dir Psychol Sci* 22: 407–412.

20. McEwen BS, Gianaros PJ (2010): Central role of the brain in stress and adaptation: Links

to socioeconomic status, health, and disease. *Ann N Y Acad Sci* 1186: 190–222.

21. Callaghan BL, Tottenham N (2016): The Stress Acceleration Hypothesis: effects of early-

life adversity on emotion circuits and behavior. *Curr Opin Behav Sci* 7: 76–81.

22. Silvers JA, Lumian DS, Gabard-Durnam L, Gee DG, Goff B, Fareri DS, *et al.* (2016):

Previous Institutionalization Is Followed by Broader Amygdala-Hippocampal-PFC

Network Connectivity during Aversive Learning in Human Development. *J Neurosci*

Off J Soc Neurosci 36: 6420–6430.

23. McLaughlin KA, Sheridan MA, Winter W, Fox NA, Zeanah CH, Nelson CA (2014):

Widespread Reductions in Cortical Thickness Following Severe Early-Life

Deprivation: A Neurodevelopmental Pathway to Attention-Deficit/Hyperactivity

Disorder. *Biol Psychiatry* 76: 629–638.

24. Mehta MA, Golembo NI, Nosarti C, Colvert E, Mota A, Williams SCR, *et al.* (2009):

Amygdala, hippocampal and corpus callosum size following severe early institutional

deprivation: The English and Romanian Adoptees Study Pilot. *J Child Psychol*

Psychiatry 50: 943–951.

25. McLaughlin KA, Fox NA, Zeanah CH, Sheridan MA, Marshall P, Nelson CA (2010):

Delayed Maturation in Brain Electrical Activity Partially Explains the Association

Between Early Environmental Deprivation and Symptoms of Attention-

Deficit/Hyperactivity Disorder. *Biol Psychiatry* 68: 329–336.

26. Pietto ML, Kamienkowski JE, Lipina SJ (2017): Electrophysiological Approaches in the

Study of the Influence of Childhood Poverty on Cognition. In: Ibáñez A, Sedeño L,

García AM, editors. *Neuroscience and Social Science: The Missing Link*. Cham:

Springer International Publishing, pp 349–381.

27. Pierce LJ, Thompson BL, Gharib A, Schlueter L, Reilly E, Valdes V, *et al.* (2019):

Association of Perceived Maternal Stress During the Perinatal Period With

Electroencephalography Patterns in 2-Month-Old Infants. *JAMA Pediatr* 173: 561–

570.

28. McLaughlin KA, Sheridan MA, Humphreys KL, Belsky J, Ellis BJ (2021): The Value of

Dimensional Models of Early Experience: Thinking Clearly About Concepts and

Categories. *Perspect Psychol Sci* 16: 1463–1472.

29. Cohen-Cline H, Jones KG, Kulkarni-Rajasekhara S, Polonsky HM, Vartanian KB (2019):

Identifying underlying constructs of childhood adversity in a low-income population.

Child Abuse Negl 91: 1–11.

30. Mersky JP, Janczewski CE, Topitzes J (2017): Rethinking the Measurement of Adversity:

Moving Toward Second-Generation Research on Adverse Childhood Experiences.

Child Maltreat 22: 58–68.

31. Nikolaidis A, Heleniak C, Fields A, Bloom PA, VanTieghem M, Vannucci A, *et al.*

(2022): Heterogeneity in caregiving-related early adversity: Creating stable

dimensions and subtypes. *Dev Psychopathol* 34: 621–634.

32. van Zyl T-L, O'Neill T, Rushe T (2023): Distinct Psychological Profiles Linking

Childhood Adversity and Substance Misuse in High-Risk Young Adults. *J Fam*

Violence 38: 633–645.

33. Zinn ME, Huntley ED, Keating DP (2020): Resilience in adolescence: Prospective Self

moderates the association of early life adversity with externalizing problems. *J*

Adolesc 81: 61–72.

34. Brieant A, Vannucci A, Nakua H, Harris J, Lovell J, Brundavanam D, *et al.* (2023):

Characterizing the dimensional structure of early-life adversity in the Adolescent

Brain Cognitive Development (ABCD) Study. *Dev Cogn Neurosci* 61: 101256.

35. Cole, Franke K (2017): Predicting Age Using Neuroimaging: Innovative Brain Ageing

Biomarkers. *Trends Neurosci* 40: 681–690.

36. Franke K, Ziegler G, Klöppel S, Gaser C (2010): Estimating the age of healthy subjects

from T1-weighted MRI scans using kernel methods: Exploring the influence of

various parameters. *NeuroImage* 50: 883–892.

37. Drobinin V, Van Gestel H, Helmick CA, Schmidt MH, Bowen CV, Uher R (2022): The

Developmental Brain Age Is Associated With Adversity, Depression, and Functional

Outcomes Among Adolescents. *Biol Psychiatry Cogn Neurosci Neuroimaging* 7:

406–414.

38. Franke K, Luders E, May A, Wilke M, Gaser C (2012): Brain maturation: Predicting

individual BrainAGE in children and adolescents using structural MRI. *NeuroImage*

63: 1305–1312.

39. Holm MC, Leonardsen EH, Beck D, Dahl A, Kjelkenes R, de Lange A-MG, Westlye LT

(2023): Linking brain maturation and puberty during early adolescence using

longitudinal brain age prediction in the ABCD cohort. *Dev Cogn Neurosci* 60:

101220.

40. Whitmore LB, Weston SJ, Mills KL (2023): BrainAGE as a measure of maturation

during early adolescence. *Imaging Neurosci* 1: 1–21.

41. Lund MJ, Alnæs D, de Lange A-MG, Andreassen OA, Westlye LT, Kaufmann T (2022):

Brain age prediction using fMRI network coupling in youths and associations with

psychiatric symptoms. *NeuroImage Clin* 33: 102921.

42. Cohen JW, Ramphal B, DeSerisy M, Zhao Y, Pagliaccio D, Colcombe S, *et al.* (2024):

Relative brain age is associated with socioeconomic status and anxiety/depression

problems in youth. *Dev Psychol* 60: 199–209.

43. Cropley VL, Tian Y, Fernando K, Mansour L. S, Pantelis C, Cocchi L, Zalesky A (2021):

Brain-Predicted Age Associates With Psychopathology Dimensions in Youths. *Biol*

Psychiatry Cogn Neurosci Neuroimaging 6: 410–419.

44. Kurth F, Levitt JG, Gaser C, Alger J, Loo SK, Narr KL, *et al.* (2022): Preliminary

evidence for a lower brain age in children with attention-deficit/hyperactivity

disorder. *Front Psychiatry* 13. Retrieved June 8, 2023, from

https://www.frontiersin.org/articles/10.3389/fpsyt.2022.1019546

45. Alnæs D, Kaufmann T, Marquand AF, Smith SM, Westlye LT (2020): Patterns of

sociocognitive stratification and perinatal risk in the child brain. *Proc Natl Acad Sci*

117: 12419–12427.

46. Holz NE, Zabihi M, Kia SM, Monninger M, Aggensteiner P-M, Siehl S, *et al.* (2023): A

stable and replicable neural signature of lifespan adversity in the adult brain. *Nat*

Neurosci 26: 1603–1612.

47. Jeong HJ, Durham EL, Moore TM, Dupont RM, McDowell M, Cardenas-Iniguez C, *et al.*

(2021): The association between latent trauma and brain structure in children [no. 1].

Transl Psychiatry 11: 1–9.

48. Malave L, van Dijk MT, Anacker C (2022): Early life adversity shapes neural circuit

function during sensitive postnatal developmental periods [no. 1]. *Transl Psychiatry*

12: 1–14.

49. Modabbernia A, Reichenberg A, Ing A, Moser DA, Doucet GE, Artiges E, *et al.* (2021):

Linked patterns of biological and environmental covariation with brain structure in

adolescence: a population-based longitudinal study [no. 9]. *Mol Psychiatry* 26: 4905–

4918.

50. Casey BJ, Cannonier T, Conley MI, Cohen AO, Barch DM, Heitzeg MM, *et al.* (2018):

The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition

across 21 sites. *Dev Cogn Neurosci* 32: 43–54.

51. Karcher NR, Barch DM, Avenevoli S, Savill M, Huber RS, Simon TJ, *et al.* (2018):

Assessment of the Prodromal Questionnaire–Brief Child Version for Measurement of

Self-reported Psychoticlike Experiences in Childhood. *JAMA Psychiatry* 75: 853–861.

52. Auchter AM, Hernandez Mejia M, Heyser CJ, Shilling PD, Jernigan TL, Brown SA, *et al.*

(2018): A description of the ABCD organizational structure and communication

framework. *Dev Cogn Neurosci* 32: 8–15.

53. Hagler DJ, Hatton SeanN, Cornejo MD, Makowski C, Fair DA, Dick AS, *et al.* (2019):

Image processing and analysis methods for the Adolescent Brain Cognitive

Development Study. *NeuroImage* 202: 116091.

54. Dale AM, Fischl B, Sereno MI (1999): Cortical Surface-Based Analysis. 16.

55. Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, *et al.* (2002): Whole

Brain Segmentation. *Neuron* 33: 341–355.

56. Hagler Jr. DJ, Ahmadi ME, Kuperman J, Holland D, McDonald CR, Halgren E, Dale AM

(2009): Automated white-matter tractography using a probabilistic diffusion tensor

atlas: Application to temporal lobe epilepsy. *Hum Brain Mapp* 30: 1535–1547.

57. Van Dijk KRA, Hedden T, Venkataraman A, Evans KC, Lazar SW, Buckner RL (2010):

Intrinsic Functional Connectivity As a Tool For Human Connectomics: Theory,

Properties, and Optimization. *J Neurophysiol* 103: 297–321.

58. Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE (2016):

Generation and Evaluation of a Cortical Area Parcellation from Resting-State

Correlations. *Cereb Cortex N Y NY* 26: 288–303.

59. Beer JC, Tustison NJ, Cook PA, Davatzikos C, Sheline YI, Shinohara RT, Linn KA

(2020): Longitudinal ComBat: A method for harmonizing longitudinal multi-scanner

imaging data. *NeuroImage* 220: 117129.

60. Chen T, Guestrin C (2016): XGBoost: A Scalable Tree Boosting System. *Proceedings of*

the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data

Mining 785–794.

61. Dehestani N, Whittle S, Vijayakumar N, Silk TJ (2023): Developmental brain changes

during puberty and associations with mental health problems. *Dev Cogn Neurosci* 60:

101227.

62. Liang H, Zhang F, Niu X (2019): Investigating systematic bias in brain age estimation

with application to post☐traumatic stress disorders. *Hum Brain Mapp* hbm.24588.

63. de Lange, Cole (2020): Commentary: Correction procedures in brain-age prediction.

NeuroImage Clin 26: 102229.

64. Muthén L, Muthén B (1998): Mplus User Guide. Retrieved November 21, 2023, from

https://www.statmodel.com/HTML_UG/introV8.htm

65. R Core Team (2022): *R: A Language and Environment for Statistical Computing*. Vienna,

Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-

project.org/

66. Bürkner P-C (2017): **brms**☐: An *R* Package for Bayesian Multilevel Models Using *Stan*.

J Stat Softw 80. https://doi.org/10.18637/jss.v080.i01

67. Bürkner P-C (2018): Advanced Bayesian Multilevel Modeling with the R Package brms.

R J 10: 395.

68. Bates D, Mächler M, Bolker B, Walker S (2015): Fitting Linear Mixed-Effects Models

Using **lme4**. *J Stat Softw* 67. https://doi.org/10.18637/jss.v067.i01

69. Wagenmakers E-J, Lodewyckx T, Kuriyal H, Grasman R (2010): Bayesian hypothesis

testing for psychologists: A tutorial on the Savage–Dickey method. *Cognit Psychol*

60: 158–189.

70. Green JG, McLaughlin KA, Berglund PA, Gruber MJ, Sampson NA, Zaslavsky AM,

Kessler RC (2010): Childhood adversities and adult psychiatric disorders in the

national comorbidity survey replication I: associations with first onset of DSM-IV

disorders. *Arch Gen Psychiatry* 67: 113–123.

71. Chong VE, Lee WS, Victorino GP (2015): Neighborhood socioeconomic status is

associated with violent reinjury. *J Surg Res* 199: 177–182.

72. Meyer OL, Castro-Schilo L, Aguilar-Gaxiola S (2014): Determinants of Mental Health

and Self-Rated Health: A Model of Socioeconomic Status, Neighborhood Safety, and

Physical Activity. *Am J Public Health* 104: 1734–1741.

73. Butler O, Yang X, Laube C, Kühn S, Immordino☐Yang MH (2018): Community

violence exposure correlates with smaller gray matter volume and lower IQ in urban

adolescents. *Hum Brain Mapp* 39: 2088–2097.

74. Noble KG, Houston SM, Brito NH, Bartsch H, Kan E, Kuperman JM, *et al.* (2015):

Family income, parental education and brain structure in children and adolescents [no.

5]. *Nat Neurosci* 18: 773–778.

75. Whittle S, Vijayakumar N, Simmons JG, Dennison M, Schwartz O, Pantelis C, *et al.*

(2017): Role of Positive Parenting in the Association Between Neighborhood Social

Disadvantage and Brain Development Across Adolescence. *JAMA Psychiatry* 74:

824–832.

76. Mendle J, Leve LD, Van Ryzin M, Natsuaki MN, Ge X (2011): Associations Between

Early Life Stress, Child Maltreatment, and Pubertal Development Among Girls in

Foster Care. *J Res Adolesc Off J Soc Res Adolesc* 21: 871–880.

77. Sumner JA, Colich NL, Uddin M, Armstrong D, McLaughlin KA (2019): Early

Experiences of Threat, but Not Deprivation, Are Associated With Accelerated

Biological Aging in Children and Adolescents. *Biol Psychiatry* 85: 268–278.

78. Wolf EJ, Maniates H, Nugent N, Maihofer AX, Armstrong D, Ratanatharathorn A, *et al.*

(2018): Traumatic Stress and Accelerated DNA Methylation Age: A Meta-Analysis.

Psychoneuroendocrinology 92: 123–134.

79. Pagliaccio D, Alqueza KL, Marsh R, Auerbach RP (2020): Brain Volume Abnormalities

in Youth at High Risk for Depression: Adolescent Brain and Cognitive Development

Study. *J Am Acad Child Adolesc Psychiatry* 59: 1178–1188.

80. Chen MC, Hamilton JP, Gotlib IH (2010): Decreased Hippocampal Volume in Healthy

Girls at Risk of Depression. *Arch Gen Psychiatry* 67: 270–276.

81. Rao U, Chen L-A, Bidesi AS, Shad MU, Thomas MA, Hammen CL (2010):

Hippocampal Changes Associated with Early-Life Adversity and Vulnerability to

Depression. *Biol Psychiatry* 67: 357–364.

82. Herting MM, Johnson C, Mills KL, Vijayakumar N, Dennison M, Liu C, *et al.* (2018):

Development of subcortical volumes across adolescence in males and females: A

multisample study of longitudinal changes. *NeuroImage* 172: 194–205.

83. Dima D, Modabbernia A, Papachristou E, Doucet GE, Agartz I, Aghajani M, *et al.*

(2022): Subcortical volumes across the lifespan: Data from 18,605 healthy individuals

aged 3–90 years. *Hum Brain Mapp* 43: 452–469.

84. Lupien SJ, Parent S, Evans AC, Tremblay RE, Zelazo PD, Corbo V, *et al.* (2011): Larger

amygdala but no change in hippocampal volume in 10-year-old children exposed to

maternal depressive symptomatology since birth. *Proc Natl Acad Sci* 108: 14324–

14329.

85. Mattoni M, Hopman HJ, Dadematthews A, Chan SSM, Olino TM (2023): Specificity of

associations between parental psychopathology and offspring brain structure.

Psychiatry Res Neuroimaging 334: 111684.

86. Freisthler B, Weiss RE (2008): Using Bayesian Space-Time Models to Understand the

Substance Use Environment and Risk for Being Referred to Child Protective Services.

Subst Use Misuse 43: 239–251.

87. Glaser D (2000): Child Abuse and Neglect and the Brain—A Review. *J Child Psychol*

Psychiatry 41: 97–116.

88. Gong W, Rolls ET, Du J, Feng J, Cheng W (2021): Brain structure is linked to the

association between family environment and behavioral problems in children in the

ABCD study [no. 1]. *Nat Commun* 12: 3769.

89. Leonardsen EH, Peng H, Kaufmann T, Agartz I, Andreassen OA, Celius EG, *et al.*

(2022): Deep neural networks learn general and clinically relevant representations of

the ageing brain. *NeuroImage* 256: 119210.