**Evolution of glucuronoxylan side chain variability in vascular plants and**

**the counter-adaptation of pathogenic cell-wall-degrading hydrolases**

Li Yu[1#], Louis F.L. Wilson[1#], Oliver M. Terrett[1#], Joel Wurman-Rodrich[1], Jan J.

Lyczakowski[1], Xiaolan Yu[1], Kristian B.R.M. Krogh[2], Paul Dupree[1*]

[1] Department of Biochemistry, University of Cambridge, Hopkins Building, The Downing

Site, Tennis Court Road, Cambridge CB2 1QW, UK

[2] Novozymes A/S, Krogshøjvej 36, 2880 Bagsværd, Denmark

[#] These authors contributed equally to this paper.

[*] Corresponding author: Paul Dupree   Phone number: (44)1223 333340

Email: pd101@cam.ac.uk

This manuscript includes: 6498 words; 8 Coloured Figures, 12 Supplemental Figures and 1

Supplemental Table.

**<u>Summary</u>**

• Polysaccharide structural complexity not only influences cell wall strength and
extensibility, but also hinders pathogenic and biotechnological attempts to saccharify the
wall. In certain species and tissues, glucuronic acid side chains on xylan exhibit
arabinopyranose or galactose decorations whose genetic and evolutionary basis is
completely unknown, impeding efforts to understand their function and engineer wall
digestibility.

• Genetics and polysaccharide profiling were used to identify the responsible loci in
Arabidopsis and Eucalyptus from proposed candidates, while phylogenies uncovered a
shared evolutionary origin. GH30-family *endo*-glucuronoxylanase activities were
analysed by electrophoresis and their differing specificities were rationalised by
phylogeny and structural analysis.

• The newly identified xylan arabinopyranosyltransferases comprise an overlooked
subfamily in the GT47-A family of Golgi glycosyltransferases, previously assumed to
comprise mainly xyloglucan galactosyltransferases, highlighting an unanticipated
adaptation of both donor and acceptor specificities. Further neofunctionalisation has
produced a Myrtaceae-specific xylan galactosyltransferase. Simultaneously, GH30 endo-
glucuronoxylanases have convergently adapted to overcome these decorations, suggesting
a role for these structures in defence. The differential expression of glucuronoxylan-
modifying genes across Eucalyptus tissues, however, hints at further functions.

• Our results demonstrate the rapid adaptability of biosynthetic and degradative
carbohydrate-active enzyme activities, providing insight into a plant-pathogen arms race
and facilitating plant cell wall biotechnological utilisation.

**<u>Key words:</u>** evolution, cell wall, glycosyltransferase, neofunctionalisation, XAPT, xylan,
xylanase

## Introduction

Surrounding every plant cell, the cell wall is an extracellular matrix that generates turgor pressure and affords protection from abiotic and biotic stresses. Cell walls are composed of a complex combination of polysaccharides, phenolic polymers, and proteins; however, despite a universal and fundamental role in cell biology, their structural compositions have diverged dramatically and rapidly throughout evolution (Burton *et al*., 2010). Given the cell wall's importance both for morphology and as a complex physical and metabolic barrier to pathogens, this diversity likely results not only from adaptation to different physical environments, but also from a complex ongoing arms race between plants and their enemies (Malinovsky *et al.*, 2014).

Forming a common hemicellulosic component of plant cell walls, xylan polysaccharides are thought to bind to and thus chaperone cellulose microfibrils (Busse-Wicher *et al.*, 2014; Simmons *et al.*, 2016) as well as mediating cross-links to other cell wall components (Terrett & Dupree, 2019; Tryfona *et al.*, 2023). Xylans vary in structure between different species and tissues, but invariably they possess a backbone of β1,4-linked xylosyl residues (Scheller & Ulvskov, 2010). In Arabidopsis and other eudicots, the xylan backbone is typically decorated with 2/3-linked acetate and α1,2-linked glucuronic acid (GlcA), which is often 4-*O*-methylated. The spacing of these decorations along the xylan backbone is critical for regulating its interaction with cellulose microfibrils (Simmons *et al.*, 2016; Grantham *et al.*, 2017; Tryfona *et al.*, 2023). While most xylan in Arabidopsis is present in the secondary cell wall (SCW), it also makes up part of the primary cell wall (PCW) (Mortimer *et al.*, 2015). Although the function of PCW xylan has not been determined, plants appear to allocate dedicated machinery to synthesise it: whereas the SCW xylan backbone is synthesised by IRX9, IRX10, and IRX14/14L, PCW xylan is synthesised specifically by IRX9L, IRX10L, and IRX14 (Mortimer *et al.*, 2015; Anders *et al.*, 2022). Furthermore, the pattern of GlcA decoration differs substantially between PCW and SCW xylan. SCW xylan glucuronidation is achieved by two enzymes: GUX1, which produces an even-spaced pattern, and GUX2, which produces a more random pattern of GlcA side chains (Bromley *et al.*, 2013). In contrast, PCW xylan is glucuronidated consistently at every sixth backbone xylosyl residue by GUX3 (Mortimer *et al.*, 2015). Notably, in at least some PCW-rich Arabidopsis tissues, the GlcA moieties of PCW xylan can be further decorated with a pentose that has been assigned as a putative α1,2-linked L-arabinopyranose (Ara*p*) (Chong *et al.*, 2015; Mortimer *et al.*, 2015).

The glycosyltransferase responsible for this decoration has not yet been identified, hindering

investigations into the functions of these Ara$p$-α1,2-GlcA-α1,2- (U$^{Arap}$) disaccharide

sidechains.

Prior to its discovery in Arabidopsis, decoration of xylan GlcA residues with additional

sugars was previously observed for several species in the *Eucalyptus* genus of the Myrtaceae

family (Shatalov *et al.*, 1999; Evtuguin *et al.*, 2003; Pinto *et al.*, 2005). Interestingly, further

experiments clearly showed that the GlcA residues in Eucalyptus xylan can not only be

substituted with α1,2-linked L-Ara$p$ (as thought to be the case in Arabidopsis) but also with a

sterically similar moiety: β1,2-linked D-galactose (Gal) (Togashi *et al.*, 2009; Pena *et al.*,

2016). Furthermore, the U$^{Arap}$ sidechain has also been shown to be widely present in xylans

from non-commelinid monocots (Pena *et al.*, 2016). However, whether the Gal-β1,2-GlcA-

α1,2- (U$^{Gal}$) sidechain is present in plants outside the *Eucalyptus* genus is unknown.

Disaccharide side chains terminating in 2-linked sugars, particularly β1,2-linked galactose,

can also be found decorating other PCW hemicelluloses, namely xyloglucan and β-

galactoglucomannan (β-GGM). Arabidopsis mutants lacking the galactosyl residues of

xyloglucan's Gal-β1,2-Xyl-α1,6- side chains exhibit stunted growth (Tamura *et al.*, 2005),

while further removal of the equivalent residue in β-GGM's Gal-β1,2-Gal-α1,6- side chains

exacerbates the phenotype (Yu *et al.*, 2022). These side chain structures may influence the

ability of these hemicelluloses to achieve their presumed primary function in chaperoning and

cross-linking cellulose microfibrils. However, as well as contributing to the material

properties of the cell wall, PCW hemicelluloses pose an obstacle to invading pathogens.

Pathogens must therefore secrete cell wall-degrading glycosyl hydrolases (GHs) in order to

gain access to the cell, and the added structural complexity afforded by cell wall

polysaccharide decorations likely increase the difficulty in doing so.

In particular, to overcome the barrier posed by xylan, fungal and bacterial pathogens typically

secrete a variety of xylanolytic enzymes. Xylanase activity is thought to be an important part

of infection by disease-causing microbes (Bellincampi *et al*., 2014) , and cereals such as

wheat produce xylanase-inhibitory proteins in response to fungal infection (Chmelová *et al.*,

2019). Xylan degradation begins with backbone cleavage by *endo*-xylanases, the best studied

being from CAZy families GH5, 8, 10, 11, and 30 (Paes *et al.*, 2012). Enzymes from GH8,

10, and 11 tend to cut preferentially at unsubstituted regions of the xylan backbone. Such

enzymes are impeded by side chain decorations, which also require later deconstruction by

dedicated *exo*-glycosidases (terminal GlcA residues are removed by α-glucuronidases from

families GH67 and GH115, for instance). However, some *endo*-xylanases have specifically

evolved to recognise certain side chains: most characterised GH30 xylanases require a GlcA-

substituted xylosyl residue at the −2 subsite, while GH5 xylanases can be arabinoxylan-

specific. Nevertheless, binding of such enzymes is inhibited by incompatible or clustered side

chains. For these reasons, additional structural complexity provides increased resistance to

degradation, and plants are presumably under a constant selection pressure to innovate new

and more complex xylan structures. An example of such adaptation may lie in corn bran

xylan, whose unique complexity provides exceptional recalcitrance to enzymatic digestion

(Rogowski *et al.*, 2015). In turn, pathogens are under a selection pressure to develop

hydrolases that can accommodate these more complex xylan structures (Beri *et al.*, 2020).

Investigating the true functions of such cell wall structural features, as well as the

evolutionary forces behind their continuing diversification, is impossible without knowing

the relevant biosynthetic genes. In the case of xylan, the genes necessary for

arabinopyranosylating and galactosylating glucuronic acid side chains are still unknown.

Here we report that a family of mainly xyloglucan-modifying glycosyltransferases have been

remodelled through evolution for this purpose, requiring reprogramming of both donor sugar

and backbone acceptor specificities to generate the new xylan-arabinopyranosylating activity.

We characterise the activities of three family members through heterologous expression in

Arabidopsis, and show that, moreover, this family has itself undergone even more recent

neofunctionalisation in the Myrtaceae family to produce xylan galactosyltransferases,

explaining the previously observed xylan structures in Eucalyptus. Furthermore, we show

that the relative expression of xylan arabinopyranosyltransferases and galactosyltransferases

across Eucalyptus tissues defines two contrasting distributions of $U^{Arap}$ and $U^{Gal}$ structures,

suggesting a potential functional role in specialised cell wall architectures. Finally, we show

not only that the decoration of GlcA with Ara*p*/Gal impedes the action of some GH30

glucuronoxylanases (suggesting a role in passive defence) but also that a minority of GH30

family members have convergently acquired amino acid substitutions to accommodate such

structures. As well as providing insight into the synthesis and function of PCW xylan and the

arms race between plants and pathogens, our findings are therefore of benefit in improving

industrial hemicellulosic biomass degradation.

## **Materials and Methods**

### **Plant Material**

All Arabidopsis (*Arabidopsis thaliana* (L.) Heynh.) T-DNA insertion mutant lines were in the Col-0 ecotype. *At1g68470* (*Atgt17/xapt1*): WiscDsLox437H04 (*xapt1-1*), *at2g29040* (*Atgt11*): Salk_108349, *at2g32750* (*Atgt12*): Salk_057536, *at2g32740* (*Atgt13*): Salk_053593C, *at4g22580* (*Atgt19*): Salk_075020 and *at5g41250*: Salk_085915 (*Atgt20*) were obtained from the Nottingham Arabidopsis Stock Centre (NASC, Nottingham, UK). Homozygous lines were isolated by PCR. The *irx9l* (SALK_037323) and *mur3-3* mutants have been described previously (Wu *et al.*, 2010; *Kong et al.*, 2015). All the primers used are shown on Table S1. Arabidopsis young stem was harvested at 30 days and bottom stem was harvested at 8 weeks.

Actively growing *Eucalyptus dalrympleana* Maiden branch and leaves were harvested at Jesus College, University of Cambridge in April 2018. Young, actively growing *Myrtus communis* L. and *Metrosideros excelsa* Sol. ex Gaertn. branchlets were harvested at Cambridge University Botanic Garden, and *Plinia cauliflora* (Mart.) Kausel, *Psidium guajava* L., and *Myrcianthes pungens* (O.Berg) D.Legrand material was a gift of Pedro Araújo (University of Campinas). Combined phloem and periderm cortex material was separated from xylem/pith by peeling or scraping. After collection, all the materials were put into 96% ethanol. For all samples, alcohol insoluble residue (AIR) was prepared as described previously (Goubet *et al.*, 2009).

### **Plant Growth and Arabidopsis transformation**

Arabidopsis plants were grown on a 9:1 mix of Levington M3 compost and vermiculite and grown at 21°C under a 16 h/8 h photoperiod. Constructs used for *A. thaliana* Xylan ArabinoPyranosylTransferase 1 (*At*XAPT1), *Eucalyptus grandis* W.Hill ex Maiden (*Eg*XAPT), and Xylan GalactoPyranosylTransferase (*Eg*XLPT) overexpression in *xapt1* and WT plants were prepared using OpenPlant common syntax GoldenGate assembly (Patron *et al.*, 2015). *AtXAPT1* (*At1g68470*), *EgXAPT* (*Eucgr.D00738*; GenBank: XM_010054097.3), and *EgXLPT* (*Eucgr.H00343*; GenBank: XM_010070979.3) coding sequences and the *CESA3* and *IRX3* promoter sequences were synthesised *de novo* by IDT or GeneWiz after removal of

165    *BsaI* and *BpiI* restriction sites. Arabidopsis transformation was performed using the floral dip

166    method as previously described (Clough & Bent, 1998).

**Phylogenetic and bioinformatics analysis**

For the XAPT phylogeny, XAPT orthologues (GT47-A$_V$) from a wide range of tracheophyte

species were extracted from our recent phylogeny of GT47-A enzymes (Yu *et al.*, 2022). To

obtain further XAPT orthologues from plants in the Myrtales order, the genomes of

*Metrosideros polymorpha* (Izuno *et al.*, 2016; Izuno *et al.*, 2019), *Rhodamnia argantea*,

*Syzygium oleosum*, and *Punica granatum* (https://www.ncbi.nlm.nih.gov/genome/) and the

transcriptome of *Oenothera rosea* (Carpenter et al., 2019; One Thousand Plant

Transcriptomes, 2019), were searched using TBLASTN (Altschul *et al.*, 1990) using

*At*XAPT1 as a query with a cut-off at $E = 1 \times 10^{-125}$. Sequences were aligned using MUSCLE

(Edgar, 2004a; Edgar, 2004b) and truncated to the GT47 domain. ProtTest3 (Darriba *et al.*,

2011) was then used to select an appropriate substitution model, and a phylogeny was

calculated using RAxML (Stamatakis, 2014) with 100 rapid bootstraps. For the main text

figure, the tree was then trimmed to species of interest using ape 5.0 (Paradis & Schliep,

2019).

For the GH30 subfamily 8 tree, CAZy sequences were downloaded from the dbCAN2 server

(http://bcb.unl.edu/dbCAN2/) (Zhang *et al.*, 2018). Using bespoke Python scripts invoking

HMMER (Eddy, 2011), GH30_8 domain-containing sequences were extracted and trimmed

to the GH30 domain. Sequences with a GH30 domain sequence shorter than 200 residues

were omitted. Extremely similar sequences were removed using CD-HIT (Li & Godzik,

2006; Fu *et al.*, 2012) with a 99 % similarity threshold. The remaining sequences were

aligned using MAFFT (Katoh & Standley, 2013); ProtTest3 was then used to select an

appropriate substitution model, and a phylogeny was calculated using RAxML with 100 rapid

bootstraps.

**Genomic DNA extraction and sequencing of native *XAPT* and *XLPT* loci**

*E. dalrympleana* leaves were ground in liquid nitrogen and genomic DNA was extracted as

previously described (James *et al.*, 2001), supplementing the extraction buffer with 3 %

polyvinylpyrrolidone. PCR amplification was achieved using Q5 High-Fidelity DNA

Polymerase (Biolab); primers are listed in Table S1. PCR products were purified by gel

extraction and isopropanol precipitation before Sanger sequencing.

**Enzyme Digestions**

Soluble hemicelluloses were prepared from AIR as previously described (Lyczakowski *et al.*,

2017). For Arabidopsis bottom stem and *E. dalrympleana* xylan analysis, 0.5 mg AIR was

used for one reaction. For Arabidopsis young stem, 1 mg AIR was used for one reaction.

Soluble hemicellulose fraction was digested with an excess of Xyn11A from *Neocallimastix*

*patriciarum* (*Np*Xyn11A, Genbank: CAA46498.1; from Megazyme) in 50 mM ammonium

acetate, pH 6.0 at 30°C overnight.

Alternatively, xylan was digested with GH30 xylanases with a 70 μg ml$^{-1}$ concentration from

different bacteria at 25°C for 30 min in 50 mM ammonium acetate, pH 6.0. The GH30

xylanases included: XynC from *Bacteroides ovatus* (*Bo*XynC, GenBank: ALJ48333.1; a gift

of Harry Gilbert, Newcastle University); Xyn30A from *Hungateiclostridium thermocellum*

(*Ct*Xyn30A, GenBank: ABN54208.1; from NZYTech). XynA from *Dickeya chrysanthemi*

strain D1 (*Ec*$_D$Xyn30A, GenBank: AAB53151.1), XynA from *Dickeya chrysanthemi* strain

P860219 (*Ec*$_P$Xyn30A, GenBank: AAL16415.1), and the two mutagenized enzymes

*Ec*$_D$Xyn30A$_{Y255L}$ and *Ec*$_P$Xyn30A$_{L255Y}$ were provided by Novozymes A/S. For mass

spectrometry and galactosidase sensitivity experiments, xylanase products were then digested

overnight with 10 μL of 1 mg ml$^{-1}$ *B. ovatus* GH115 glucuronidase (*Bo*Agu115A, GenBank:

EDO10816.1; a gift of Harry Gilbert, Newcastle University) to remove the non-decorated

GlcA side chains.

For the galactosidase sensitivity assay, the xylanase products were passed through a 3-kDa

Nanosep centrifugal device. Next, ethanol was added to a final concentration to 65% before

incubation at −20°C for 2 h. The samples were then centrifuged at 12,000g for 10 min; the

supernatant was subsequently dried 60°C *in vacuo* and resuspended in 200 μl of 50 mM

ammonium acetate, pH 6.0. To this, 0.5 μl of 4 kU ml$^{-1}$ GH35 β-galactosidase from

*Aspergillus niger* (Megazyme) was added and incubated at 37°C for 18 h. Samples were then

dried at 60°C *in vacuo*.

 **Oligosaccharides Fingerprint Analysis by PACE**

Samples and $(Xyl)_{1-6}$ standards (Megazyme, Ireland) were derivatized with 8-
aminonapthalene-1,3,6-tresulphonic acid (ANTS; Invitrogen) as described previously
(Goubet et al., 2002). After drying, the samples were resuspended in 100 µl of 3 M urea, of
which 2 µl was loaded onto the PACE gels. The samples were run and visualized as
described previously (Goubet *et al.*, 2009).

**Preparation of Oligosaccharides for MS**

Following the enzyme digestion, released peptides and enzymes were removed using reverse-
phase Sep-Pak C18 cartridges (Waters) as previously described. Oligosaccharides were
reductively aminated with 2-aminobenzoic acid (2-AA) and cleaned using a GlycoClean S
cartridge (ProZyme) as previously described (Tryfona & Stephens, 2010).

## **Results**

*At1g68470* **encodes** *XYLAN ARABINOPYRANOSYLTRANSFERASE 1* (*XAPT1*) **in**
**Arabidopsis**

Despite an almost-complete genetic understanding of xylan biosynthesis, the gene(s)
responsible for arabinopyranosylating and galactosylating glucuronic acid side chains remain
unknown, thwarting efforts to determine their function and manipulate xylan structure. To
remedy this, we sought to obtain candidate arabinopyranosyltransferase mutants in
Arabidopsis, which normally exhibits putative $U^{Arap}$ (Ara$p$-α1,2-GlcA-α1,2-) xylan side
chains in its primary cell wall. However, according to the CAZy database (Drula *et al.*, 2022),
the Arabidopsis genome encodes 558 glycosyltransferases (GTs) distributed amongst at least
43 GT families, of which only a minority are functionally characterised; too large a number
to be screened experimentally. Instead, we attempted to predict likely candidates through
biochemical plausibility. Firstly, we observed that the α1,2-linked arabinopyranosyl moiety
of the $U^{Arap}$ structure is sterically almost identical to the β1,2-linked galactosyl moiety of the
Eucalyptus $U^{Gal}$ structure (Gal-β1,2-GlcA-α1,2-), differing only in the presence or absence of
a C6 hydroxymethyl group. Therefore, we hypothesised that the two corresponding
glycosyltransferases may be closely related. Secondly, owing to our recent work on

xyloglucan and β-GGM synthesis, we observed that all subfamily A members of the CAZy

GT47 family (GT47-A) characterised so far exhibit an inverting mechanism and affix the

donor sugar, most commonly galactose, using a 1,2-glycosidic linkage (MUR3, for instance,

creates the Gal-β1,2-Xyl-α1,6- 'L' side chain of xyloglucan). Furthermore, all known

acceptors are α-linked monosaccharide decorations of a β1,4-linked hemicellulose backbone.

These intriguing similarities to the theoretical activities of xylan arabinopyranosyltransferases

and galactosyltransferases led us to hypothesise that the responsible genes may reside in the

same family.

We recently calculated a large-scale phylogeny of the GT47-A subfamily in plants (Yu *et al.*,

2022), which revealed that the subfamily consists of at least seven subclades (GT47-A$_{I-VII}$).

Importantly, xyloglucan-modifying activities are spread almost universally throughout these

clades, which has always given the impression that all unidentified members share this

activity. Nevertheless, several Arabidopsis members—as well as two subclades (A$_{IV}$ and

A$_V$)—are completely uncharacterised; hence, we hypothesised that these previously

overlooked enzymes could harbour unanticipated xylan-modifying activities. To test this

hypothesis, we attempted to detect the loss of the PCW xylan U$^{Ara\textit{p}}$ structure in T-DNA

knockout lines for the six remaining Arabidopsis GT47-A subclade members with unknown

functions (*At5g41250/AtGT20* from A$_{II}$, *At4g22580/AtGT19* from A$_{IV}$, *At1g68470/AtGT17*

from A$_V$, and *At2g29040/AtGT11*, *At2g32750/AtGT12*, and *At2g32740/AtGT13* from A$_{VII}$).

*At2g29040/AtGT11* has since been implicated in galactosylation of pollen tube xyloglucan

(Wei *et al.*, 2021). We used the *irx9l* mutant, which lacks PCW-type xylan (and therefore

does not exhibit any U$^{Ara\textit{p}}$ structure), as a control (Mortimer *et al.*, 2015). AIR from the

young stems of these mutants was incubated with alkali to solubilize the hemicelluloses, and

then digested with *Np*Xyn11A *endo*-xylanase. The resultant oligosaccharides were analysed

by PACE. As previously reported (Mortimer *et al.*, 2015), wild-type (WT) young stem

yielded a series of products including xylose (X), xylobiose (X$_2$), aldopentaouronic acid with

(XU$^m$XX) or without (XUXX) 4-O-methyl group on the GlcA (generically termed

'XU$^{[m]}$XX'), and xylopentaose substituted with the putative Ara*p*-GlcA- disaccharide

(XU$^{Ara\textit{p}}$XXX). In contrast, digestion of material from the *irx9l* mutant released xylose,

xylobiose (X$_2$), and XU$^{[m]}$XX, but not XU$^{Ara\textit{p}}$XXX. Interestingly, although we could detect

XU$^{Ara\textit{p}}$XXX in five of the six GT47-A mutants, this product was also conspicuously absent

for *at1g68470* (A$_V$), suggesting that xylan in this mutant lacks the U$^{Ara\textit{p}}$ decoration (Fig. 1a,

S1). To help confirm this, WT young stem was hydrolysed with the GH30-family xylanase

Xyn30A from *Bacteroides ovatus* (*Bo*XynC), which requires a GlcA substitution at its $-2$

subsite. As previously described (Mortimer *et al.*, 2015), digestion of WT material with

*Bo*XynC produced XXXXU$^{[m]}$X (U$_1^{[m]}$X$_6$) and XXXXU$^{Arap}$X (U$_1^{Arap}$X$_6$) as major products,

in accordance with the consistent GlcA spacing of PCW xylan. In contrast, digestion of

material from the *at1g68470* mutant released U$_1^{[m]}$X$_6$ but not the U$_1^{Arap}$X$_6$ oligosaccharide

(Fig. 1b). These results suggest that, unexpectedly, *At1g68470* is required for the synthesis of

the putative Ara*p*-GlcA- disaccharide substitution on Arabidopsis PCW xylan.

To further support our conclusion, we complemented the *at1g68470* mutant with *At1g68470*

under the control of the PCW-synthesis-specific *CESA3* promoter. AIR from young stems

was digested by xylanase *Np*Xyn11A and analysed by PACE. In all three independent

transgenic lines, the XU$^{Arap}$XXX product was detected at substantially higher levels than in

WT (Fig. 1a). Based on these data, and the previous arabinopyranose assignment, we

concluded that *At1g68470* likely encodes a xylan-specific arabinopyranosyltransferase that

catalyses the formation of an Ara*p*-α1,2-GlcA linkage. We therefore propose that *At1g68470*

is named ***XYLAN ARABINOPYRANOSYLTRANSFERASE 1*** (***XAPT1***).

**XAPT underwent gene duplication in the Myrtales**

Having successfully predicted the identity of *XAPT1* in Arabidopsis, we also investigated our

hypothesis that xylan galactosyltransferases could be closely related. Importantly, whereas

the U$^{Arap}$ xylan branch structure has been reported in a wide range of species, including

Arabidopsis, *Eucalyptus* spp., and non-commelinid monocots, the U$^{Gal}$ structure has so far

only been detected in *Eucalyptus* spp. (Togashi *et al.*, 2009; Pena *et al.*, 2016). This suggests

that xylan galactosyltransferases have in fact evolved more recently than XAPTs. We

hypothesised that: *a)* XAPT1's parent subclade, GT47-A$_V$, comprises (mainly) xylan

arabinopyranosyltransferases, and *b)* that xylan galactosyltransferases also derive from this

subclade. To explore this idea, we constructed a phylogeny of XAPT homologues from 56

species of vascular plant, including *Eucalyptus grandis*, three fellow members of the

Myrtaceae family (*Metrosideros polymorpha*, *Syzygium oleosum*, and *Rhodamnia argentea*),

and two members of the wider Myrtales order (*Punica granatum* and *Oenothera rosea*). Fig.

2 presents the rooted phylogeny, which has been trimmed for brevity (the full phylogeny can

be found in Fig. S2). We found that most of the plant genomes—including *P. granatum* and

*O. rosea*—exhibited a single gene (or cluster of near-identical copies) in this subclade. In

contrast, all four Myrtaceae genomes exhibited two paralogues that appear to have arisen
from a single, recent duplication event that likely occurred in the Myrtales order. In the case
of *E. grandis*, these two paralogues correspond to the genes *Eucgr.D00738* and
*Eucgr.H00343*. Hence, we hypothesised that Eucgr.D00738 (*Eg*XAPT, hereafter) and
Eucgr.H00343 (*Eg*XLPT) could each constitute either a xylan GlcA
arabinopyranosyltransferase or potentially a xylan GlcA galactosyltransferase.

## *Eucgr.D00738* and *Eucgr.H00343* encode *Eg*XAPT and *Eg*XLPT

To investigate the activities of the two Myrtaceae XAPT-related paralogues, we expressed
*Eg*XAPT and *Eg*XLPT under the strong, PCW-synthesis-specific *CESA3* promoter in the
*xapt1* Arabidopsis mutant. Xylan digestion with *Np*Xyn11A xylanase revealed that
expression of either gene could restore the presence of disaccharide side chain structures (Fig.
3a), suggesting that both proteins are active and catalyse the formation of Ara$p$-α1,2-GlcA or
a similar linkage on xylan.

PCW xylan is the presumed natural substrate of XAPT in Arabidopsis, but PCW xylan
scarcity *in planta* makes its structure difficult to analyse. For this reason, we expressed the
two *E. grandis* enzymes in SCW-producing tissues, which synthesise much more xylan.
Furthermore, $^{[Me]}$GlcA sidechains on SCW xylan are completely unsubstituted (Bromley *et*
*al.*, 2013; Mortimer *et al.*, 2015), which produces an ideal background for characterisation of
*Eg*XAPT and *Eg*XLPT. The *Eg*XAPT and *Eg*XLPT coding sequences were placed under the
control of the SCW-synthesis-specific *IRX3* promoter and expressed in WT Arabidopsis, and
xylan from mature bottom-stems was digested with the glucuronoxylanase *Bo*XynC and
analysed by PACE. As described previously (Bromley *et al.*, 2013), digestion of the WT
xylan released products decorated with unsubstituted GlcA ($U_1^{[m]}X_n$), whose sizes
corresponded to expected $^{[Me]}$GlcA spacings (Fig. 3b). In the *Eg*XAPT- and *Eg*XLPT-
expressing lines, however, similar analysis revealed very different oligosaccharide products,
suggesting that most of the $^{[Me]}$GlcA had been modified, perhaps into $U^{Arap/Gal,[m]}$ structures.
The spacing between GlcA decorations along the xylan backbone did not appear to be
affected in these transgenic plants. Interestingly, oligosaccharides with substituted GlcA in
*Eg*XLPT-expressing lines migrated marginally slower than those in *Eg*XAPT-expressing
lines, suggesting that the products of *Eg*XAPT and *Eg*XLPT might differ subtly in structure.

To further characterise the products of $Eg$XAPT and $Eg$XLPT, xylan from the relevant transgenic lines was digested simultaneously with $Np$Xyn11A and a *Bacteroides ovatus* GH115 ($Bo$GH115) α-glucuronidase, which removes unsubstituted GlcA side chains. As expected, PACE analysis indicated that WT bottom-stem xylan was fully digested into xylose and xylobiose (Fig. S3). In the $Eg$XAPT- and $Eg$XLPT-expressing lines, however, a band of higher molecular weight ($XU^{Arap,[m]}XXX/XU^{Gal,[m]}XXX$) was produced. To distinguish between any arabinopyranosylated and galactosylated oligosaccharides, we then treated these products with a β-galactosidase ($An$GH35). Whereas the higher-molecular-weight band from $Eg$XAPT-expressing lines was completely insensitive to $An$GH35 (revealing it to be entirely $XU^{Arap,[m]}XXX$), almost all of the equivalent product from $Eg$XLPT-expressing lines shifted to $XU^{[m]}XXX$ after β-galactosidase treatment—revealing its identity as $XU^{Gal,[m]}XXX$ and indicating the presence of the $U^{Gal}$ structure in these plants. These results suggest that $Eg$XLPT, but not $Eg$XAPT, is capable of transferring galactose to GlcA side chains.

To confirm these assignments, we analysed $Bo$XynC glucuronoxylanase products by mass spectrometry (MS), which can distinguish between pentose and hexose substituents. To prevent further ambiguity between $U_1^{[m]}X_n$ and $U_1^{Arap,[m]}X_{n-1}$, $Bo$XynC products were treated with α-glucuronidase $Bo$GH115 to remove unsubstituted GlcA residues, thereby leaving $U_1^{Arap/Gal}X_n$ oligosaccharides as the only acidic products. The resulting oligosaccharides were derivatised with 2-aminobenzoic acid (2-AA) and then analysed by MALDI-TOF MS (Fig. 4, S4). As predicted, only xylo-oligosaccharides were produced from WT, confirming full removal of unsubstituted $^{[Me]}$GlcA by the α-glucuronidase. Products from plants expressing $At$XAPT under the same promoter were analysed in parallel for comparison. Lines expressing $At$XAPT and $Eg$XAPT exhibited abundant amounts of $U_1^{Arap}X_n$ and low amounts of $U_1^{Arap,m}X_n$, confirming that $Eg$XAPT is a xylan arabinopyranosyltransferase. In lines expressing $Eg$XLPT, however, the dominant sugars released could be assigned as $U_1^{Gal}X_n$, with a smaller presence of $U_1^{Gal,m}X_n$. Interestingly, this approach also detected low amounts of $U_1^{Arap}X_n$ and $U_1^{Arap,m}X_n$ in all $Eg$XLPT independent lines (Fig. 4, S4). These results suggest that $Eg$XLPT is primarily a xylan galactosyltransferase with a low level of arabinopyranosyltransferase activity, at least when expressed in Arabidopsis.

**Different *Eucalyptus dalrympleana* tissues contain structurally distinct xylans**

Although $U^{Arap}$ and $U^{Gal}$ side chains are known to exist in plants from the *Eucalyptus* genus (Togashi *et al.*, 2009; Pena *et al.*, 2016), the tissue distribution of these structures is unknown, limiting our insight into their function. To investigate where the enzymes might be active, we studied expression data available for *EgXAPT* and *EgXLPT* at https://EucGenIE.org (Hefer *et al.*, 2011), which provides mRNA-Seq datasets for both *E. grandis* and the hybrid *E. grandis* × *E. urophylla*. Despite some potential disagreement between the two datasets, a clear pattern was still apparent: whereas *EgXAPT* expression is relatively constant and lowest in xylem tissues, the expression of *EgXLPT* is much stronger in xylem tissues than in leaves, flowers, and shoot tips (Fig. S5, S6). Interestingly, the expression of *EgXLPT* is dramatically higher in *E. grandis* × *E. urophylla* tension wood xylem than control xylem, as has been previously noted (Mizrachi *et al.*, 2015).

To study the predicted xylan structure distributions *in planta*, we harvested an actively growing branch of *Eucalyptus dalrympleana* (mountain gum). To confirm that the *XAPT* and *XLPT* genes of *E. dalrympleana* do not differ substantially to those of *E. grandis*, we PCR-amplified them from genomic DNA using homology-based primers and sequenced the two intronless coding sequences, revealing 98% (XAPT) and 99% (XLPT) protein identity between the *E. dalrympleana* and *E. grandis* sequences (Fig. S7, S8). Next, to determine the extent of GlcA substitution amongst *E. dalrympleana* tissues, we sectioned the branch into cortex, phloem, sapwood/xylem, and leaf. Major leaf veins were removed to avoid contamination from vascular tissue (Fig. 5a). Alkali-treated AIR from each tissue was digested with *Bo*XynC xylanase and analysed with PACE. Based on the PACE xylo-oligosaccharide profiles, the tissues broadly fell into two groups with very different xylan structures (Fig. 5b). In the first group, which comprises leaf and phloem, $U_1^{[m]}X_6$ and $U_1^{Arap/Gal,[m]}X_6$ were the most abundant xylan oligosaccharides, with traces of $U_1^{[m]}X_5$ and $U_1^{Arap/Gal,[m]}X_5$. In the other group, which comprises cortex and xylem, the oligosaccharides were dominated by xylo-oligosaccharides decorated with single $^{[Me]}$GlcA. The cortex and xylem xylan had a GlcA spacing pattern very similar to SCW xylan from Arabidopsis (Bromley *et al.*, 2013). Small amounts of $U_1^{Arap/Gal,[m]}X_5$ and $U_1^{Arap/Gal,[m]}X_6$ oligosaccharide could also be detected in cortex and xylem.

In order to discriminate $U^{Arap}$ and $U^{Gal}$ structures in different Eucalyptus tissues, we again treated *Bo*XynC xylanase products with α-glucuronidase GH115 and analysed the remaining

acidic products by MS. The MS profile of xylan showed that neutral and acidic

oligosaccharides were released from all tissues but with different ratios (Fig. 6). Consistent

with the expression data (Fig. S5, S6), leaf tissue released $U_1^{\text{Ara}p,m}X_5$ and $U_1^{\text{Ara}p,m}X_6$

oligosaccharides but no $U_1^{\text{Gal},m}X_n$ oligosaccharides. In contrast, the stem tissues yielded

detectable levels of $U_1^{\text{Gal},m}X_5$ and $U_1^{\text{Gal},m}X_6$. From cortex to phloem to xylem, the percentage

of GlcA residues that were substituted decreased, while at the same time, the ratio of Gal to

Ara$p$ decorations increased. In summary, both the patterning and substitution of xylan GlcA

side chains varies in different *E. dalrympleana* tissues and appears to correspond with the

previously observed expression patterns of *Eg*XAPT and *Eg*XLPT in *E. grandis* / *E. grandis*

*x E. urophylla*. More specifically, XAPT appears to be responsible for arabinosylating

glucuronoxylan mainly in certain PCW-rich tissues, whereas glucuronoxylan galactosylation

by XLPT appears to be enriched in woody tissues but is less abundant overall.

**Other Myrtaceae plants contain $U^{\text{Gal}}$ and $U^{\text{Ara}p}$ structures in their xylan**

The results of our phylogenetic analysis suggested that other genera in the Myrtaceae family

might possess XLPT orthologues, and therefore might exhibit $U^{\text{Gal}}$ structures on their xylan.

To investigate this hypothesis, we sectioned and analysed tissue from five additional

Myrtaceae members: *Myrtus communis* (myrtle), *Metrosideros excelsa* (pohutukawa), *Plinia*

*cauliflora* (jaboticaba), *Psidium guajava* (guava), and *Myrcianthes pungens* (guabiju). Xylan

extracted from both cortex/phloem (combined) and xylem/pith was treated with xylanase

*Np*Xyn11A and α-glucuronidase *Bo*GH115, and as before, the products were subsequently

digested with *An*GH35 β-galactosidase. In xylem/pith, $XU^{\text{Ara}p,[m]}XXX/XU^{\text{Gal},[m]}XXX$

oligosaccharides were detectable in all species, with *M. communis* exhibiting remarkably

high levels of GlcA substitution (Fig. S9). Notably, for *M. communis*, *P. cauliflora*, *P.*

*guajava*, and *M. pungens* xylan, but not *M. excelsa*, a large proportion of this

$XU^{\text{Ara}p,[m]}XXX/XU^{\text{Gal},[m]}XXX$ product was sensitive to *An*GH35, indicating that this band

was primarily composed of the $XU^{\text{Gal},[m]}XXX$ oligosaccharide with a smaller amount of

$XU^{\text{Ara}p}XXX$ in xylem (Fig. S9). In cortex/phloem, $XU^{\text{Ara}p,[m]}XXX/XU^{\text{Gal},[m]}XXX$

oligosaccharides were detectable in all species except in *P. cauliflora* (Fig. S10). However,

only *M. communis* yielded products that were sensitive to *An*GH35, suggesting a lack of

$U^{\text{Gal},[m]}$ structures in the cortex/phloem of the other four species. Our results therefore support

the idea that many Myrtaceae plants exhibit $U^{\text{Gal}}$ side chains and that this structure is enriched

in woodier tissues.

**GH30-family glucuronoxylanases have convergently evolved to overcome the obstacle of glucuronyl side chain modification**

The distinct patterns of xylan arabinopyranosylation and galactosylation in Myrtaceae-family plants hint at functional importance for physical cell wall properties. We were therefore surprised to observe that Arabidopsis *xapt1* mutant plants exhibit no obvious growth phenotype; nor was any new phenotype revealed when *xapt1* plants were crossed with *mur3* plants lacking the xyloglucan β-galactosyltransferase MUR3 (Fig. S11). Nevertheless, we considered the possibility that xylan arabinopyranosylation could be important for other aspects of Arabidopsis biology. For example, cell wall polysaccharides constitute a barrier to potential pathogens and plant pathogens therefore rely on polysaccharide-degrading enzymes for their virulence. In this work, we have used two *endo*-xylanases to analyse xylan structure: *Np*Xyn11A, from CAZy family GH11, and *Bo*XynC, a member of GH30 subfamily 8 from the gut bacterium *B. ovatus*. In order to act on xylan, the latter of these specifically recognises and binds to GlcA substitutions. Curiously, previously solved crystal structures of related GH30 xylanases from plant pathogen *Dickeya chrysanthemi* (previously *Erwinia chrysanthemi*) strain D1 (*Ec*$_D$Xyn30A) and *Bacillus subtilis* (*Bs*XynC) show that both enzymes make a hydrogen bond to the C2 hydroxyl of GlcA using the phenol group of Tyr$_{255}$ or Tyr$_{231}$ respectively (Fig. 7a) (St John *et al.*, 2011; Urbanikova *et al.*, 2011). We reasoned that the 1,2-linked Ara*p*/Gal substitution that decorates GlcA residues at this very hydroxyl should therefore impede *Ec*$_D$Xyn30A and *Bs*XynC from binding their substrate. In spite of this, our results clearly showed that the related *Bo*XynC is able to bind and cleave at the U$^{Arap/Gal}$ decorations (Figs. 1b, 3b). To determine whether *Ec*$_D$Xyn30A and other GH30 xylanases might have the same activity, we digested xylan from Arabidopsis *gux1,2* young stem with *Ec*$_D$Xyn30A or *Ct*Xyn30A (from *Clostridium thermocellum*). PACE analysis of the products revealed that, whereas *Bo*XynC is capable of cleaving the xylan backbone adjacent to both substituted and unsubstituted $^{[Me]}$GlcA, *Ec*$_D$Xyn30A and *Ct*Xyn30A are only able to cleave effectively at unsubstituted $^{[Me]}$GlcA decorations (Fig. 8a). These results demonstrate a hitherto unknown substrate specificity variant amongst GH30 glucuronoxylanases.

To help understand the difference in substrate specificity between these enzymes, we aligned their protein sequences. The alignment showed that the tyrosine residue involved in binding the C2 hydroxyl of GlcA is also present in *Ct*Xyn30A. Interestingly, however, the corresponding residue in *Bo*XynC is not tyrosine but leucine (Fig. 7b).

To investigate the extent of amino acid variety at this position, we conducted a phylogenetic

analysis of the entire GH30 subfamily 8. We then labelled the sequences in the tree according

to the amino acid at the position of interest. The results showed that *Bo*XynC belongs to a

distinct subgroup of enzymes with leucine rather than tyrosine at this position (Fig. 7c).

Interestingly, we also identified an enzyme from *D. chrysanthemi* P860219 that diverged

much more recently from $Ec_\mathrm{D}$Xyn30A than *Bo*XynC, but has independently acquired a

leucine at this position (we henceforth refer to this enzyme as $Ec_\mathrm{P}$Xyn30A) (Fig. 7b, c).

Although $Ec_\mathrm{D}$Xyn30A and $Ec_\mathrm{P}$Xyn30A otherwise share 82% protein sequence identity, like

*Bo*XynC, $Ec_\mathrm{P}$Xyn30A (but not $Ec_\mathrm{D}$Xyn30A) was able to cut xylan adjacent to $U^{\mathrm{Ara}p/\mathrm{Gal}}$

decorations (Fig. 8b).

To investigate whether this single residue difference is sufficient to explain the discrepancy

in substrate specificity, we performed site-directed mutagenesis on the $Ec_\mathrm{D}$Xyn30A and

$Ec_\mathrm{P}$Xyn30A enzymes. $\mathrm{Leu}_{255}$ from $Ec_\mathrm{P}$Xyn30A was mutated to tyrosine to produce

$Ec_\mathrm{P}$Xyn30A$_{\mathrm{L255Y}}$, while $\mathrm{Tyr}_{255}$ from $Ec_\mathrm{D}$Xyn30A was mutated to leucine to produce

$Ec_\mathrm{D}$Xyn30A$_{\mathrm{Y255L}}$. Initially, we used the purified enzymes to digest WT bottom-stem xylan

without Ara*p*/Gal substitutions. All four enzymes released identical products of equal

abundance, suggesting that all four variants exhibit glucuronoxylanase activity (Fig. S12).

We used the purified enzymes to hydrolyse alkali-extracted xylan from plants expressing the

$_{pro}$*IRX3*:*EgXLPT*/*XAPT* constructs. As predicted, both $Ec_\mathrm{D}$Xyn30A and $Ec_\mathrm{P}$Xyn30A$_{\mathrm{L255Y}}$

produced primarily xylo-oligosaccharides with GlcA residues, whereas the enzymes

$Ec_\mathrm{D}$Xyn30A$_{\mathrm{Y255L}}$ and $Ec_\mathrm{P}$Xyn30A could release many more products with $U^{\mathrm{Ara}p/\mathrm{Gal}}$

decorations (Fig. 8b). These data suggest that a tyrosine residue at this position prevents

efficient activity on xylan substrates with substituted GlcA decorations (or that leucine is

required at this position for activity), and that multiple GH30 glucuronoxylanase subfamilies

have independently adapted to accommodate substituted GlcA decorations. Importantly,

these mutations appear to constitute counter-adaptations to the evolution of xylan

arabinopyranosylation, implicating $U^{\mathrm{Ara}p}$ side chains in passive defence against pathogens.

## **Discussion**

In this work, we successfully predicted and characterised the enzymes responsible for

decorating GlcA xylan branches in Arabidopsis and *Eucalyptus* spp., identifying GT47-A$_\mathrm{V}$ as

a novel gene subfamily. While the two new activities share the most characteristic feature of

others in GT47-A—adding a 2-linked sugar to an α-linked decoration of a hemicellulose
backbone—it is notable that both donor (arabinopyranose) and acceptor (xylan) contrast with
those of their closest characterised sisters (galactose/xyloglucan for both GT47-A$_{III}$ and
GT47-A$_{VI}$) (Yu *et al.*, 2022). This significant change in activity over a comparatively short
timespan contributes to an emerging picture of rapid neofunctionalisation within the GT47-A
subfamily (Yu *et al.*, 2022; Wilson *et al.*, 2023), whose driving forces we are just beginning
to understand.

The identity of the pentosyl residue decorating GlcA branches in Arabidopsis PCW xylan has
not been fully confirmed: previous attempts to identify it by NMR could not distinguish
between α1,2-linked arabinopyranose and β1,2-linked xylose. Nevertheless, the former is
favoured because the decoration is insensitive to β-xylosidase (Mortimer *et al.*, 2015).
Furthermore, the equivalent pentosyl decorations in *Eucalyptus grandis* and non-commelinid
monocots have been unambiguously determined to be α1,2-linked arabinopyranose (Pena *et*
*al.*, 2016). The close evolutionary relationship between the Arabidopsis and *E. grandis* XAPT
enzymes further strengthens the idea that the Arabidopsis pentosyl decoration is indeed
arabinopyranose. Hence, we are confident in assigning *At*XAPT1 as an
arabinopyranosyltransferase.

The identification of XAPT and XLPT enzymes in GT47-A$_V$ helps to explain the distribution
of substituted GlcA xylan side chains amongst plant species. For instance, our previous
phylogeny (Yu *et al.*, 2022) demonstrated that GT47-A$_V$ genes have been completely lost
from grass (Poaceae) genomes, explaining why U$^{Arap}$ side chains are present in non-
commelinid monocot orders but have never been seen in grasses (we can also now predict the
presence of U$^{Arap}$ structures in non-grass commelinids such as pineapple). We also observed
candidate XAPT orthologues in lycophytes and gymnosperms; it will be interesting to
determine if these species also carry decorated GlcA side chains. In addition, our work
highlights the emergence of U$^{Gal}$ side chains in the Myrtaceae family, likely arising from
recent duplication and neofunctionalisation of the *XAPT* locus during the evolution of the
Myrtales order.

Unlike arabinoxylan, glucuronoxylan is present in virtually every land plant cell wall,
suggesting a specific and important role for GlcA in cell wall structure—perhaps as a
covalent bridge to lignin polymers in secondary cell walls (Giummarella *et al.*, 2019; Terrett
& Dupree, 2019) or as a factor in xylan solubility and assembly onto cellulose (Schultink *et*

*al.*, 2015). Further modification of GlcA by 2-*O*-arabinopyranosylation, 2-*O*-galactosylation,

and/or 4-*O*-methylation could therefore potentially modulate xylan incorporation into the cell

wall. At the same time, the $U^{Arap}$ and $U^{Gal}$ disaccharide side chains resemble the functionally

important side chains of other PCW hemicelluloses: namely, the Gal-β1,2-Xyl-α1,6- and Gal-

β1,2-Gal-α1,6- side chains of xyloglucan and β-GGM, respectively. For xyloglucan at least,

the galactosyl decorations may influence its binding to cellulose (Zhao *et al.*, 2014) and/or

increase its solubility during biosynthesis (Tamura *et al.*, 2005; Kong *et al.*, 2015). Whether

decoration of GlcA could have an effect on xylan–cellulose interactions will require

extensive computational and solid state NMR work to establish.

Despite these possibilities, but consistent with GlcA methyltransferase mutants (Urbanowicz

*et al.*, 2012; Li *et al.*, 2013) and glucuronyltransferase mutants (Mortimer *et al.*, 2015), we

did not observe any obvious phenotype of *xapt* or *xapt mur3-3* plants. Hence, any critical role

for $U^{Arap}$ structures in Arabidopsis vegetative growth is unclear. Nevertheless, we observed

an interesting interplay between GlcA arabinopyranosylation/galactosylation and

methylation: whereas all $U^{Arap}$ and $U^{Gal}$ structures in *E. dalrympleana* were methylated (i.e.

$U^{Arap,m}$ and $U^{Gal,m}$), the equivalent structures in transgenic Arabidopsis lines expressing

*Eg*XAPT and *Eg*XLPT were almost completely unmethylated. This might suggest that

Arabidopsis GlcA 4-*O*-methyltransferases do not accept substituted GlcA residues.

Furthermore, some clues for the function of $U^{Arap}$ and $U^{Gal}$ structures may lie in their uneven

tissue distribution. Based on previous comparisons between 'young' and 'old' stems in

Arabidopsis, $U^{Arap}$ structures have so far been assumed to be a general feature of 'PCW

xylan' (Mortimer *et al.*, 2015). However, in the Myrtaceae family at least, we have shown

that the situation is likely more complicated. For example, while $U^{Arap}$ structures could be

found in typically PCW-rich organs and tissues such as leaf, cortex, and phloem, $U^{Gal}$

structures were specifically enriched in xylem/pith but completely absent from leaves.

Furthermore, given that our dissections lacked the precision to separate individual cell types,

it is possible that substitution of GlcA is limited to specialised cells within the tissues of

interest. Single-cell expression atlases for Arabidopsis stem (Shi *et al.*, 2021) and leaf (Kim

*et al.*, 2021) lend support to this idea by suggesting that *At*XAPT expression may be

selectively enriched in cell types such as cambium, phloem parenchyma, and/or stem

epidermis. Moreover, although we did not investigate the xylan structures *in planta*, *Eg*XLPT

appears to be specifically upregulated in tension wood xylem tissues—perhaps by as much as

a 50-fold increase (Mizrachi *et al.*, 2015)—hinting at a specific functional role in the cell

wall. As previously proposed, it is possible that xylan structure could be remodelled in

tension wood, perhaps in a way that affects its incorporation into the wall. Whether any

significant biochemical differences exist between $U^{Arap}$ and $U^{Gal}$ structures is unclear, but it is

possible that they differ in sensitivity to *exo*-glycosidases—be they microbial or cell wall

remodelling enzymes (such as the β-galactosidase MUM2).

Whether or not the GlcA branches on xylan play a structural role in the cell wall, their

absence leads to a major reduction in biomass resistance to enzymatic saccharification

(Lyczakowski *et al.*, 2017), suggesting that they may constitute a direct or indirect factor in

plant-pathogen interaction. Here, we have uncovered that the substitution of GlcA residues

likely affects the ability of GH30-family glucuronoxylanases to degrade xylan. Remarkably,

different lineages of this family appear to have convergently overcome the obstacle posed by

C2 modification through a single point mutation. Interestingly, the strain of *D. chrysanthemi*

whose glucuronoxylanase has recently acquired tolerance for these decorations, P860219,

was apparently cultured from *Anthurium* sp.: a non-commelinid from the Alismatales order

whose close relatives exhibit abundant $U^{Arap}$ xylan side chains (Hurlbert & Preston, 2001).

This is in contrast to substitution-intolerant strain D1, which was cultured from corn

(Vroemen *et al.*, 1995), a member of the Poaceae that lacks any substitutions. This hints that

cell wall structures could be applying a direct selection pressure on xylanase evolution.

In summary, we have identified one of the missing components of xylan synthesis in higher

plants. The identification of XAPT1 permits engineering of existing xylan polysaccharides

using synthetic biology approaches. Furthermore, by revealing previously unknown substrate

specificities in GH30 glucuronoxylanases, our results could facilitate optimisation of

industrial lignocellulosic biomass degradation. These results hint at the constant evolutionary

battle between plants and pathogens.

**<u>Acknowledgements and funding</u>**

We thank Paul Stearn (Jesus College, Cambridge), Ángela Cano and Sam Brockington

(Cambridge University Botanic Garden), and Pedro Araújo (University of Campinas) for

providing plant material.

This work was supported by the Leverhulme Trust Centre for Natural Material Innovation,

the Biotechnology and Biological Sciences Research Council (BBSRC) of the UK as part of

the OpenPlant Synthetic Biology Research Centre (Reference BB/L014130/1), The UKRI
grant underwriting the ERC advanced grant (EVOCATE Function and evolution of plant cell
wall architecture for sustainable technologies. EP/X027120/1), Innovate UK, the Cambridge
BBSRC-DTP Programme (Reference BB/J014540/1), a Novo Nordisk Foundation grant
Oxymist (Grant Number NNF20OC0059697), and a BBSRC iCASE studentship (Reference
BB/M015432/1).

**Competing interests:** K.B.R.M.K. is an employee of Novozymes, an enzyme company.
The remaining authors declare no conflict of interests.

## Author contributions

LY conceived the original screen, designed the study, performed most polysaccharide
analyses and plant transformations, and co-wrote the paper; LFLW conducted the
phylogenetic analyses, contributed to the biochemical analysis, designed the xylanase point
mutations, and co-wrote the paper; OMT carried out the molecular biology, conducted
xylanase expression and purification, and co-wrote the paper; JWR identified differences in
xylanase activities and supported the biochemical analysis and identification of Arabidopsis
T-DNA insertion lines; JJL supported the biochemical analysis and molecular biology; XY
made the double mutant; KBRMK contributed to the design and production of mutated
xylanases; PD designed and supervised the study, interpreted data and co-wrote the paper.

## Data Availability

The data that support the findings of this study are available in the main text and in the
Supporting Information for this article. Sequences for newly identified *Eucalyptus*
*dalrympleana* XAPT and XLPT will be deposited in GenBank after acceptance. All plant
material descried in this manuscript is available on a reasonable request from the
corresponding authors.

## References

**Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990.** Basic local alignment
search tool. *Journal of Molecular Biology* **215**(3): 403-410.

**Anders N, Wilson LFL, Sorieul M, Nikolovski N, Dupree P. 2022.** β-1, 4-Xylan backbone
synthesis in higher plants: How complex can it be? *Frontiers in Plant Science* **13**.
**Bellincampi D, Cervone F, Lionetti V. 2014.** Plant cell wall dynamics and wall-related
susceptibility in plant–pathogen interactions. *Frontiers in Plant Science* **5**: 228.
**Beri D, York WS, Lynd LR, Pena MJ, Herring CD. 2020.** Development of a thermophilic
coculture for corn fiber conversion to ethanol. *Nature Communications* **11**(1): 1937.
**Bromley JR, Busse-Wicher M, Tryfona T, Mortimer JC, Zhang ZN, Brown DM,**
**Dupree P. 2013.** GUX1 and GUX2 glucuronyltransferases decorate distinct domains
of glucuronoxylan with different substitution patterns. *Plant Journal* **74**(3): 423-434.
**Brown DM, Zhang Z, Stephens E, Dupree P, Turner SR. 2009.** Characterization of IRX10
and IRX10-like reveals an essential role in glucuronoxylan biosynthesis in
Arabidopsis. *Plant Journal* **57**(4): 732-746.
**Burton RA, Gidley MJ, Fincher GB. 2010.** Heterogeneity in the chemistry, structure and
function of plant cell walls. *Nature Chemical Biology* **6**(10): 724-732.
**Busse-Wicher M, Gomes TCF, Tryfona T, Nikolovski N, Stott K, Grantham NJ, Bolam**
**DN, Skaf MS, Dupree P. 2014.** The pattern of xylan acetylation suggests xylan may
interact with cellulose microfibrils as a twofold helical screw in the secondary plant
cell wall of Arabidopsis thaliana. *Plant Journal* **79**(3): 492-506.
**Carpenter EJ, Matasci N, Ayyampalayam S, Wu S, Sun J, Yu J, Jimenez Vieira FR,**
**Bowler C, Dorrell RG, Gitzendanner MA, et al. 2019.** Access to RNA-sequencing
data from 1,173 plant species: The 1000 Plant transcriptomes initiative (1KP).
*Gigascience* **8**(10).
**Chmelová D, Škulcová D, Ondrejovic M. 2019.** Microbial xylanases and their inhibition by
specific proteins in cereals. *Kvasny Prumysl* **65**(4): 127–133-127–133.
**Chong SL, Koutaniemi S, Juvonen M, Derba-Maceluch M, Mellerowicz EJ, Tenkanen**
**M. 2015.** Glucuronic acid in Arabidopsis thaliana xylans carries a novel pentose
substituent. *International Journal of Biological Macromolecules* **79**: 807-812.
**Clough SJ, Bent AF. 1998.** Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant Journal* **16**(6): 735-743.
**Darriba D, Taboada GL, Doallo R, Posada D. 2011.** ProtTest 3: fast selection of best-fit
models of protein evolution. *Bioinformatics* **27**(8): 1164-1165.
**Drula E, Garron ML, Dogan S, Lombard V, Henrissat B, Terrapon N. 2022.** The
carbohydrate-active enzyme database: functions and literature. *Nucleic Acids Res*
**50**(D1): D571-D577.
**Eddy SR. 2011.** Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**(10): e1002195.
**Edgar RC. 2004a.** MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. *BMC Bioinformatics* **5**: 113.
**Edgar RC. 2004b.** MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Research* **32**(5): 1792-1797.
**Evtuguin DV, Tomas JL, Silva AMS, Neto CP. 2003.** Characterization of an acetylated
heteroxylan from Eucalyptus globulus Labill. *Carbohydrate Research* **338**(7): 597-
604.
**Fu L, Niu B, Zhu Z, Wu S, Li W. 2012.** CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**(23): 3150-3152.
**Giummarella N, Pu YQ, Ragauskas AJ, Lawoko M. 2019.** A critical review on the
analysis of lignin carbohydrate bonds. *Green Chemistry* **21**(7): 1573-1595.
**Goubet F, Barton CJ, Mortimer JC, Yu XL, Zhang ZN, Miles GP, Richens J, Liepman**
**AH, Seffen K, Dupree P. 2009.** Cell wall glucomannan in Arabidopsis is synthesised
by CSLA glycosyltransferases, and influences the progression of embryogenesis.
*Plant Journal* **60**(3): 527-538.

**Grantham NJ, Wurman-Rodrich J, Terrett OM, Lyczakowski JJ, Stott K, Iuga D,**
**Simmons TJ, Durand-Tardif M, Brown SP, Dupree R, et al. 2017.** An even
pattern of xylan substitution is critical for interaction with cellulose in plant cell walls.
*Nature Plants* **3**(11): 859-865.
**Han MH, Liu YT, Zhang FL, Sun DF, Jiang JX. 2020.** Effect of galactose side-chain on
the self-assembly of xyloglucan macromolecule. *Carbohydrate Polymers* **246**.
**Hefer C, Mizrachi E, Joubert F, Myburg A 2011**. The Eucalyptus genome integrative
explorer (EucGenIE): a resource for Eucalyptus genomics and transcriptomics. *BMC*
*Proceedings*: Springer. 1-1.
**Hurlbert JC, Preston JF, 3rd. 2001.** Functional characterization of a novel xylanase from a
corn strain of Erwinia chrysanthemi. *Journal of Bacteriology* **183**(6): 2093-2100.
**Immelmann R, Gawenda N, Ramirez V, Pauly M. 2023.** Identification of a xyloglucan
beta-xylopyranosyltransferase from Vaccinium corymbosum. *Plant Direct* **7**(7): e514.
**Izuno A, Hatakeyama M, Nishiyama T, Tamaki I, Shimizu-Inatsugi R, Sasaki R,**
**Shimizu KK, Isagi Y. 2016.** Genome sequencing of Metrosideros polymorpha
(Myrtaceae), a dominant species in various habitats in the Hawaiian Islands with
remarkable phenotypic variations. *Journal of Plant Research* **129**(4): 727-736.
**Izuno A, Wicker T, Hatakeyama M, Copetti D, Shimizu KK. 2019.** Updated Genome
Assembly and Annotation for Metrosideros polymorpha, an Emerging Model Tree
Species of Ecological Divergence. *G3 (Bethesda)* **9**(11): 3513-3520.
**James CM, Barrett JA, Russell SJ, Gibby M. 2001.** A rapid PCR based method to
establish the potential for paternal inheritance of chloroplasts in Pelargonium. *Plant*
*Molecular Biology Reporter* **19**: 163-167.
**Katoh K, Standley DM. 2013.** MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular Biology and Evolution* **30**(4):
772-780.
**Kim JY, Symeonidi E, Pang TY, Denyer T, Weidauer D, Bezrutczyk M, Miras M,**
**Zollner N, Hartwig T, Wudick MM, et al. 2021.** Distinct identities of leaf phloem
cells revealed by single cell transcriptomics. *Plant Cell* **33**(3): 511-530.
**Kong YZ, Pena MJ, Renna L, Avci U, Pattathil S, Tuomivaara ST, Li XM, Reiter WD,**
**Brandizzi F, Hahn MG, et al. 2015.** Galactose-depleted xyloglucan is dysfunctional
and leads to dwarfism in Arabidopsis. *Plant Physiology* **167**(4): 1296-U1294.
**Li W, Godzik A. 2006.** Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**(13): 1658-1659.
**Li XF, Jackson P, Rubtsov DV, Faria-Blanc N, Mortimer JC, Turner SR, Krogh KB,**
**Johansen KS, Dupree P. 2013.** Development and application of a high throughput
carbohydrate profiling technique for analyzing plant cell wall polysaccharides and
carbohydrate active enzymes. *Biotechnology for Biofuels* **6**:94.
**Li XM, Cordero I, Caplan J, Molhoj M, Reiter WD. 2004.** Molecular analysis of 10
coding regions from arabidopsis that are homologous to the MUR3 xyloglucan
galactosyltransferase. *Plant Physiology* **134**(3): 940-950.
**Lyczakowski JJ, Wicher KB, Terrett OM, Faria-Blanc N, Yu X, Brown D, Krogh K,**
**Dupree P, Busse-Wicher M. 2017.** Removal of glucuronic acid from xylan is a
strategy to improve the conversion of plant biomass to sugars for bioenergy.
*Biotechnology for Biofuels* **10**: 224.
**Malinovsky FG, Fangel JU, Willats WG. 2014.** The role of the cell wall in plant immunity.
*Frontiers in Plant Science* **5**: 178.
**Mizrachi E, Maloney VJ, Silberbauer J, Hefer CA, Berger DK, Mansfield SD, Myburg**
**AA. 2015.** Investigating the molecular underpinnings underlying morphology and

changes in carbon partitioning during tension wood formation in Eucalyptus. *New Phytologist* **206**(4): 1351-1363.

Mortimer JC, Faria-Blanc N, Yu XL, Tryfona T, Sorieul M, Ng YZ, Zhang ZN, Stott K, Anders N, Dupree P. 2015. An unusual xylan in Arabidopsis primary cell walls is synthesised by GUX3, IRX9L, IRX10L and IRX14. *Plant Journal* **83**(3): 413-426.

One Thousand Plant Transcriptomes I. 2019. One thousand plant transcriptomes and the phylogenomics of green plants. *Nature* **574**(7780): 679-685.

Paes G, Berrin JG, Beaugrand J. 2012. GH11 xylanases: Structure/function/properties relationships and applications. *Biotechnology Advances* **30**(3): 564-592.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**(3): 526-528.

Patron NJ, Orzaez D, Marillonnet S, Warzecha H, Matthewman C, Youles M, Raitskin O, Leveau A, Farre G, Rogers C, et al. 2015. Standards for plant synthetic biology: a common syntax for exchange of DNA parts. *New Phytologist* **208**(1): 13-19.

Pena MJ, Kulkarni AR, Backe J, Boyd M, O'Neill MA, York WS. 2016. Structural diversity of xylans in the cell walls of monocots. *Planta* **244**(3): 589-606.

Pinto PC, Evtuguin DV, Neto CP. 2005. Structure of hardwood glucuronoxylans: modifications and impact on pulp retention during wood kraft pulping. *Carbohydrate Polymers* **60**(4): 489-497.

Rogowski A, Briggs JA, Mortimer JC, Tryfona T, Terrapon N, Lowe EC, Basle A, Morland C, Day AM, Zheng HJ, et al. 2015. Glycan complexity dictates microbial resource allocation in the large intestine. *Nature Communications* **6**: 7481.

Scheller HV, Ulvskov P. 2010. Hemicelluloses. *Annual Review of Plant Biol*ogy **61**: 263-289.

Schultink A, Cheng K, Park YB, Cosgrove DJ, Pauly M. 2013. The identification of two arabinosyltransferases from tomato reveals functional equivalency of xyloglucan side chain substituents. *Plant Physiology* **163**(1): 86-94.

Schultink A, Naylor D, Dama M, Pauly M. 2015. The Role of the Plant-Specific ALTERED XYLOGLUCAN9 Protein in Arabidopsis Cell Wall Polysaccharide O-Acetylation. *Plant Physiology* **167**(4): 1271-1283.

Shatalov AA, Evtuguin DV, Neto CP. 1999. (2-O-alpha-D-galactopyranosyl-4-O-methyl-alpha-D-glucurono)-D-xylan from Eucalyptus globulus Labill. *Carbohydrate Research* **320**(1-2): 93-99.

Shi D, Jouannet V, Agusti J, Kaul V, Levitsky V, Sanchez P, Mironova VV, Greb T. 2021. Tissue-specific transcriptome profiling of the Arabidopsis inflorescence stem reveals local cellular signatures. *Plant Cell* **33**(2): 200-223.

Simmons TJ, Mortimer JC, Bernardinelli OD, Pöppler AC, Brown SP, Deazevedo ER, Dupree R, Dupree P. 2016. Folding of xylan onto cellulose fibrils in plant cell walls revealed by solid-state NMR. *Nature Communications* **7**: 13902.

St John FJ, Hurlbert JC, Rice JD, Preston JF, Pozharski E. 2011. Ligand bound structures of a glycosyl hydrolase family 30 glucuronoxylan xylanohydrolase. *Journal of Molecular Biology* **407**(1): 92-109.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**(9): 1312-1313.

Tamura K, Shimada T, Kondo M, Nishimura M, Hara-Nishimura I. 2005. KATAMARI1/MURUS3 is a novel Golgi membrane protein that is required for endomembrane organization in Arabidopsis. *Plant Cell* **17**(6): 1764-1776.

Terrett OM, Dupree P. 2019. Covalent interactions between lignin and hemicelluloses in plant secondary cell walls. *Current Opinion in Biotechnology* **56**: 97-104.

**Togashi H, Kato A, Shimizu K. 2009.** Enzymatically derived aldouronic acids from Eucalyptus globulus glucuronoxylan. *Carbohydrate Polymers* **78**(2): 247-252.

**Tryfona T, Bourdon M, Delgado Marques R, Busse-Wicher M, Vilaplana F, Stott K, Dupree P. 2023.** Grass xylan structural variation suggests functional specialization and distinctive interaction with cellulose and lignin. *Plant Journal* **113**(5): 1004-1020.

**Tryfona T, Liang HC, Kotake T, Kaneko S, Marsh J, Ichinose H, Lovegrove A, Tsumuraya Y, Shewry PR, Stephens E, et al. 2010.** Carbohydrate structural analysis of wheat flour arabinogalactan protein. *Carbohydrate Research* **345**(18): 2648-2656.

**Tryfona T, Stephens E. 2010.** Analysis of carbohydrates on proteins by offline normal-phase liquid chromatography MALDI-TOF/TOF-MS/MS. *Plant Cell Wall: Methods and Protocols* **658**: 137-151.

**Urbanikova L, Vrsanska M, Morkeberg Krogh KB, Hoff T, Biely P. 2011.** Structural basis for substrate recognition by Erwinia chrysanthemi GH30 glucuronoxylanase. *FEBS Journal* **278**(12): 2105-2116.

**Urbanowicz BR, Pena MJ, Ratnaparkhe S, Avci U, Backe J, Steet HF, Foston M, Li HJ, O'Neill MA, Ragauskas AJ, et al. 2012.** 4-O-methylation of glucuronic acid in Arabidopsis glucuronoxylan is catalyzed by a domain of unknown function family 579 protein. *Proceedings of the National Academy of Sciences of the United States of America* **109**(35): 14253-14258.

**Van Bel M, Diels T, Vancaester E, Kreft L, Botzki A, Van de Peer Y, Coppens F, Vandepoele K. 2018.** PLAZA 4.0: an integrative resource for functional, evolutionary and comparative plant genomics. *Nucleic Acids Research* **46**(D1): D1190-D1196.

**Vroemen S, Heldens J, Boyd C, Henrissat B, Keen NT. 1995.** Cloning and characterization of the bgxA gene from Erwinia chrysanthemi D1 which encodes a beta-glucosidase/xylosidase enzyme. *Molecular Genetics and Genomics* **246**(4): 465-477.

**Wei QQ, Yang Y, Li H, Liu ZW, Fu R, Feng HQ, Li C. 2021.** The xyloglucan galactosylation modulates the cell wall stability of pollen tube. *Planta* **254**(6): 133-140.

**Wilson LF, Neun S, Yu L, Tryfona T, Stott K, Hollfelder F, Dupree P. 2023.** The biosynthesis, degradation, and function of cell wall β☐xylosylated xyloglucan mirrors that of arabinoxyloglucan. *New Phytologist* **240** (6): 2353-2371.

**Wu AM, Hornblad E, Voxeur A, Gerber L, Rihouey C, Lerouge P, Marchant A. 2010.** Analysis of the Arabidopsis IRX9/IRX9-L and IRX14/IRX14-L pairs of glycosyltransferase genes reveals critical contributions to biosynthesis of the hemicellulose glucuronoxylan. *Plant Physiology* **153**(2): 542-554.

**Yu L, Yoshimi Y, Cresswell R, Wightman R, Lyczakowski JJ, Wilson LFL, Ishida K, Stott K, Yu XL, Charalambous S, et al. 2022.** Eudicot primary cell wall glucomannan is related in synthesis, structure, and function to xyloglucan. *Plant Cell* **34**(11): 4600-4622.

**Zhang H, Yohe T, Huang L, Entwistle S, brown P, Yang Z, Busk PK, Xu Y, Yin Y. 2018.** dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Research* **46**(W1): W95-W101.

**Zhao Z, Crespi VH, Kubicki JD, Cosgrove DJ, Zhong LH. 2014.** Molecular dynamics simulation study of xyloglucan adsorption on cellulose surfaces: effects of surface hydrophobicity and side-chain variation. *Cellulose* **21**(2): 1025-1039.

**Zhu L, Dama M, Pauly M. 2018.** Identification of an arabinopyranosyltransferase from *Physcomitrella patens* involved in the synthesis of the hemicellulose xyloglucan. *Plant Direct* **2**(3): e00046.

## **Figure Legends**

**Figure 1**. AT1G68470/XAPT1 from GT47 subclade A is a functional xylan arabinopyranosyltransferase. (a), PACE analysis of *Np*Xyn11A xylanase digests of alkali-extracted xylan from AIR of WT, the *xapt1* T-DNA insertion mutant, and three independent transgenic lines expressing *XAPT1* under the strong primary wall-specific promoter of *CESA3* in the *xapt1* background. (b), PACE analysis of *Bo*XynC xylanase digests of WT and *xapt1* mutant young stem. Xylo-oligosaccharide standard: $X$–$X_6$.

**Figure 2**. Trimmed phylogenetic tree of the XAPT subclade of GT47-$A_V$. Sequences grouped in the XAPT subclade of our previous GT47-A tree (Yu et al., 2022) were supplemented with HMMER and TBLASTN hits for XAPT-like sequences in Myrtales genomes. Following alignment, the phylogeny was inferred using RAxML. For rooting, *At*MUR3 was included as the outgroup. Due to the large number of sequences (106), we show only an abridged form of the tree here (displaying the more pertinent sequences), hence bootstrap values (from 100 pseudoreplicates) may underestimate the confidence of the splits shown. The full phylogeny can be viewed in Fig. S3.

**Figure 3**. Overexpression of *Eg*XLPT or *Eg*XAPT can decorate the glucuronoxylan with an additional sugar. (a), Expression of *Eg*XLPT and *Eg*XAPT under primary cell wall *CESA3* promoter can complement the xapt1 xylan structure phenotype. Alkali-extracted xylan from 5-week-old young stem was digested with *Np*Xyn11A xylananase and analyzed by PACE. (b), Expression of *Eg*XLPT and *Eg*XAPT under secondary wall *IRX3* promoter in Arabidopsis WT can decorate most of the glucuronoxylan with an addition sugar. Alkali-extracted xylan from 8-week-old bottom stem was digested with *Bo*XynC xylanase and analysed by PACE. Xylo-oligosaccharide standard: $X$–$X_6$.

**Figure 4**. *Eg*XAPT and *Eg*XLPT exhibit different donor sugar specificities. AIR was extracted from 8-week-old bottom stems of either Arabidopsis WT plants or transgenic plants expressing *Eg*XAPT or *Eg*XLPT underthe secondary cell wall-specific promoter of IRX3. Alkali-extracted xylan was then digested with *Bo*Xyn30C xylanase followed by GH115 glucuronidase. The products were labelled with 2-AA and analysed by MALDI-TOF MS.

**Figure 5**. Xylan structure in different *Eucalyptus dalrympleana* tissues. (a), Eucalyptus stems

were sectioned into cortex, phloem, and xylem/sapwood. Major leaf vein tissue was not

included for the leaf xylan analysis. Scale bar = 0.9 cm. (b), PACE analysis of *Bo*XynC

xylanase digests of AIR from different tissues. Xylo-oligosaccharide standard: $X–X_6$.

$U^{Arap,Gal,[m]}X_n$ structures are labelled with arrows.

**Figure 6**. Distribution of xylan with decorated GlcA in different *Eucalyptus dalrympleana*

tissues. AIR from different tissues was digested with *Bo*XynC xylanase followed by GH115

glucuronidase to remove the undecorated GlcA. The products were labelled with 2-AA and

then analysed by MS.

**Figure 7**. Differences in GH30 xylanase active site residues may reflect differences in ability

to accommodate substituted glucuronic acid sidechains. (a), Previously characterised active

site of GH30 xylanase from plant pathogen *Dickeya chrysanthemi* (previously *Erwinia*

*chrysanthemi*) strain D1 (*Ec*$_D$Xyn30A; PDB: 2Y24). Bound aldotetraouronic acid is shown in

stick representation. The side chain of Tyr255, which forms a hydrogen bond with the GlcA

C2 hydroxyl, is highlighted in orange. (b), Alignment segment of five GH30 xylanases,

highlighting residues aligned to Tyr255 in *Ec*$_D$Xyn30A. (c), Molecular phylogeny of GH30

subfamily 8 sequences downloaded from dbCAN2 (http://bcb.unl.edu/dbCAN2/). Sequences

that display a tyrosine residue in the column aligned to Tyr255 are highlighted in orange;

those that instead display leucine are highlighted in blue, while those with asparagine are

highlighted in purple.

**Figure 8**. PACE gels to show the activities of different GH30 xylanases towards substituted

glucuronoxylan. (a), PACE analysis of three GH30 xylanases-digests of xylan from *gux1*

*gux2* double mutant, which only contains primary cell wall glucuronoxylan. (b), AIR from

the transgenic plants expressing *Eg*XAPT or *Eg*XLPT under the secondary cell wall-specific

promoter of *IRX3* were used as the substrates. Alkali-extracted xylan was then digested with

different GH30 and the corresponding modified GH30 enzymes. Xylo-oligosaccharide

standard: $X–X_6$.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

(a)

Xyl

Xyl

GlcA

Xyl

Me

6
5
1
4
2   3

Tyr255

2.7 Å

PDB: 2Y24

(b)

$Ec_D$ Xyn30A    QLWMTEH**Y**VDSKQSAN[263]
$Ec_P$ Xyn30A    QVWMTEH**L**VDSKQSAN[263]
$Bs$XynC         DLWMTEV**Y**YPNSDTNS[270]
$Ct$Xyn30A       ELWMTEV**Y**VPNSDSNS[268]
$Bo$XynC         TWWMTEH**L**FNDGENSD[382]
                 ****      . :. .

(c)

Residue aligned to
Tyr255 in EcDXyn30A:
tyrosine (Y)
leucine (L)
asparagine (N)
other

0.3

Figure 7.

Figure 8