1   **XAPT and XLPT enzymes modify the glucuronic acid side chains of tissue-**
2   **specific xylans in Arabidopsis and Eucalyptus**

Li Yu[1#], Louis F.L. Wilson[1#], Oliver M. Terrett[1#], Joel Wurman-Rodrich[1], Jan J.
Lyczakowski[1], Xiaolan Yu[1], Kristian B.R.M. Krogh[2], Paul Dupree[1*]

[1] Department of Biochemistry, University of Cambridge, Hopkins Building, The Downing

Site, Tennis Court Road, Cambridge CB2 1QW, UK

[2] Novozymes A/S, Krogshøjvej 36, 2880 Bagsværd, Denmark

[#] These authors contributed equally to this paper.

[*] Corresponding author: Paul Dupree   Phone number: (44)1223 333340

Email: pd101@cam.ac.uk

This manuscript includes: 6495 words; 8 Coloured Figures, 12 Supplemental Figures and 1
Supplemental Table.

Summary

- Plant cell wall polysaccharides such as xylan not only contribute to cell shape, but also

hinder pathogenic and biotechnological attempts to saccharify the wall. This recalcitrance

derives substantially from polysaccharide structural complexity, driven by biosynthetic

glycosyltransferases. Glucuronic acid sidechains along the xylan backbone can be further

decorated with arabinopyranose or galactose in certain tissues and species. We sought to

identify the responsible glycosyltransferases.

- We identified the relevant locus from candidate Arabidopsis mutants by polysaccharide

profiling and reverse genetics. We demonstrated different activities of two paralogues in

eucalyptus by heterologous expression and investigated this neofunctionalisation by

molecular phylogeny. Finally, we measured the activities of various GH30-family *endo*-

glucuronoxylanases on modified xylans *in vitro*.

- *At*XAPT1 (AT1G68470) and *Eg*XAPT (Eucgr.D00738) encode Xylan

ArabinoPyranosylTransferases, whereas *Eg*XLPT (Eucgr.H00343) encodes a Xylan

GalactoPyranosylTransferase. While arabinopyranosyl decorations in native eucalyptus

appear to be enriched in primary walls, overexpression of the eucalyptus enzymes in

Arabidopsis can accomplish substantial changes in secondary wall xylan structure. Only

some GH30 glucuronoxylanases have evolved to accommodate substituted glucuronic

acid, perhaps through a single active site substitution.

- Our results demonstrate the adaptability of biosynthetic and degradative carbohydrate-

active enzyme activities, providing insight into a plant-pathogen arms race and facilitating

wall biotechnological utilisation.

Key words: biosynthesis, cell wall, glycosyltransferase, neofunctionalisation, XAPT,

xylan, xylanase

Introduction

The plant cell wall is an extracellular matrix that surrounds every plant cell (Burton *et al.*,

2010). Depending on the cell type, the cell wall supports the upright growth of the plant and

acts as a complex physical barrier that protects the cell from abiotic and biotic stress. Cell

walls are composed of a mixture of polysaccharides, phenolic polymers, and proteins whose

composition varies considerably between different plants (Burton *et al.*, 2010). The structural

complexity of cell walls is thought to contribute to cell wall properties, including their

resistance to degradation by pathogens, and the different structures of polysaccharides may

shape the properties of different cell types and tissues.

Forming a common hemicellulosic component of plant cell walls, xylan polysaccharides are

thought to bind to and thus chaperone cellulose microfibrils (Busse-Wicher *et al.*, 2014;

Simmons *et al.*, 2016) and mediate cross-links to other cell wall components (Terrett &

Dupree, 2019; Tryfona *et al.*, 2023). Xylans vary in structure between different species and

tissues, but invariably they possess a backbone of β1,4-linked xylosyl residues (Scheller &

Ulvskov, 2010). In Arabidopsis and other eudicots, the xylan backbone is typically decorated

with side groups of 2/3-linked acetate and α1,2-linked glucuronic acid (GlcA), which is often

4-*O*-methylated. The spacing of these decorations along the xylan backbone is critical for

regulating its interaction with cellulose microfibrils (Simmons *et al.*, 2016; Grantham *et al.*,

2017; Tryfona *et al.*, 2023). While most xylan in Arabidopsis is present in the secondary cell

wall (SCW), it also makes up part of the primary cell wall (PCW) (Mortimer *et al.*, 2015).

Although the function of PCW xylan has not been determined, plants appear to allocate

dedicated machinery to synthesise it: whereas the SCW xylan backbone is synthesised by

IRX9, IRX10, and IRX14/14L, PCW xylan is synthesised specifically by IRX9L, IRX10L,

and IRX14 (Mortimer *et al.*, 2015; Anders *et al.*, 2022). Furthermore, the pattern of GlcA

decoration differs substantially between PCW and SCW xylan. SCW xylan glucuronidation is

achieved by two enzymes: GUX1, which produces an even-spaced pattern, and GUX2, which

produces a more random pattern of GlcA side chains (Bromley *et al.*, 2013). In contrast,

PCW xylan is glucuronidated consistently at every sixth backbone xylosyl residue by GUX3

(Mortimer *et al.*, 2015). Notably, in at least some PCW-rich Arabidopsis tissues, the GlcA

moieties of PCW xylan can be further decorated with a pentose that has been assigned as a

putative α1,2-linked L-arabinopyranose (Ara*p*) (Chong *et al.*, 2015; Mortimer *et al.*, 2015).

The glycosyltransferase responsible for this decoration has not yet been identified, hindering

investigations into the functions of these Ara$p$-α1,2-GlcA-α1,2- (U$^{\text{Ara}p}$) disaccharide
sidechains.

Prior to its discovery in Arabidopsis, decoration of xylan GlcA residues with additional
sugars was previously observed for several species in the *Eucalyptus* genus of the Myrtaceae
family (Shatalov *et al.*, 1999; Evtuguin *et al.*, 2003; Pinto *et al.*, 2005). Interestingly, further
experiments showed unequivocally that the GlcA residues in Eucalyptus xylan can not only
be substituted with α1,2-linked L-Ara$p$ (as thought to be the case in Arabidopsis) but also
with a sterically similar moiety, β1,2-linked D-galactose (Gal) (Togashi *et al.*, 2009; Pena *et*
al., 2016). Furthermore, the U$^{\text{Ara}p}$ sidechain has also been shown to be widely present in
xylans from non-commelinid monocots (Pena *et al.*, 2016). However, whether the Gal-β1,2-
GlcA-α1,2- (U$^{\text{Gal}}$) sidechain is present in plants outside the *Eucalyptus* genus is unknown.

Similar disaccharide side chains with terminal β1,2-linked galactose can be found in several
other widespread PCW hemicelluloses. For example, the glucan backbone of xyloglucan—
the most abundant hemicellulose in eudicot PCWs—is frequently substituted with a Gal-
β1,2-Xyl-α1,6- disaccharide (Scheller & Ulvskov, 2010). Similarly, the backbone mannosyl
residues of β-galactoglucomannan (β-GGM) can be substituted with a Gal-β1,2-Gal-α1,6-
disaccharide (Yu *et al.*, 2022). The galactosyl residue on xyloglucan side chains is essential
for normal morphology, and mutant plants lacking these residues exhibit severely stunted
growth (Tamura *et al.*, 2005); furthermore, this phenotype can be exacerbated by removing
the analogous galactosyl residues from β-GGM (Yu *et al.*, 2022). Although the mechanism
for this phenotype has not been completely clarified, these phenotypes may result from self-
aggregation of xyloglucan in the absence of the galactosyl side chains (Kong *et al.*, 2015;
Han *et al.*, 2020). However, galactose itself is not specifically required for xyloglucan
function, as normal growth of these galactosylation mutants can be restored by replacing the
missing galactosyl residues with α-L-arabinofuranosyl, α-L-arabinopyranosyl, or β-
xylopyranosyl moieties (Schultink *et al.*, 2013; Zhu *et al.*, 2018; Immelmann *et al.*, 2023;
Wilson *et al.*, 2023).

As well as contributing to the material properties of the cell wall, PCW hemicelluloses pose
an obstacle to invading pathogens. Pathogens—as well as saprophytes and xylan-degrading
gut micro-organisms—must therefore secrete cell wall-degrading glycosyl hydrolases (GHs)
in order to gain access to the cell. To overcome the barrier posed by xylan, fungal and
bacterial pathogens typically secrete *endo*-xylanases—the best studied being from CAZy

families GH5, 8, 10, 11, and 30 (Paes *et al.*, 2012). Different families differ in their substrate

specificities: whereas GH8, 10, and 11 tend to cut preferentially at unsubstituted regions of

the xylan backbone, GH5 xylanases are typically arabinoxylan-specific, and most

characterised GH30 xylanases require a GlcA-substituted xylosyl residue at the -2 subsite.

Efficient xylan digestion is dependent on additional glycosidases that remove obstructive

backbone decorations and/or help to break down the resultant oligosaccharides (α-

glucuronidases from family GH115, for instance); therefore, a pathogen's ability to digest

xylan is limited by the GH activities it can produce. For these reasons, the additional

structural complexity provided by side chains in xylan and other PCW hemicelluloses may

limit degradation. Indeed, xylanases active on simpler birch wood xylans have been shown to

be less effective at hydrolysing complex corn xylan (Rogowski *et al.*, 2015). Xylanase

activity is thought to be an important part of infection by disease-causing microbes

(Bellincampi *et al.*, 2014). Supporting this, cereals such as wheat produce xylanase-inhibitory

proteins in response to fungal infection (Chmelová *et al.*, 2019). Thus, the structural

complexity of xylan, and its effects on xylan digestibility may have important effects during

infection by pathogens.

In this work, we identified the glycosyltransferases responsible for decorating xylan GlcA

residues in both Arabidopsis and *Eucalyptus grandis*. Using a synthetic biology approach, we

engineered Arabidopsis SCW xylan so that the majority of its GlcA branches exhibited Ara*p*

or Gal substitutions. These highly substituted novel xylans may have useful biotechnological

properties, and may affect cell wall degradation by plant pathogens. We demonstrate the

difference in tissue distribution between U^{Arap} and U^{Gal} structures in *Eucalyptus* spp., and, by

analysing the evolution of the GT47-A$_V$ family (Yu *et al.*, 2022), explain how the U^{Gal}

structure has arisen specifically within the Myrtaceae family. Furthermore, we show not only

that the decoration of GlcA with Ara*p*/Gal impedes the action of some GH30

glucuronoxylanases, but also that a convergently acquired amino acid substitution allows

others to accommodate such structures. As well as providing insight into the synthesis and

function of PCW xylan and the arms race between plants and pathogens, our results could

therefore help increase efficiency in industrial hemicellulosic biomass degradation.

Materials and Methods

Plant Material

All Arabidopsis (*Arabidopsis thaliana* (L.) Heynh.) T-DNA insertion mutant lines were in the Col-0 ecotype. *At1g68470* (*Atgt17/xapt1*): WiscDsLox437H04 (*xapt1-1*), *at2g29040* (*Atgt11*): Salk_108349, *at2g32750* (*Atgt12*): Salk_057536, *at2g32740* (*Atgt13*): Salk_053593C, *at4g22580* (*Atgt19*): Salk_075020 and *at5g41250*: Salk_085915 (*Atgt20*) were obtained from the Nottingham Arabidopsis Stock Centre (NASC, Nottingham, UK). Homozygous lines were isolated by PCR. The *irx9l* (SALK_037323) and *mur3-3* mutants have been described previously (Wu *et al.*, 2010; *Kong et al.*, 2015). All the primers used are shown on Table S1. Arabidopsis young stem was harvested at 30 days and bottom stem was harvested at 8 weeks.

Actively growing *Eucalyptus dalrympleana* Maiden branch and leaves were harvested at Jesus College, University of Cambridge in April 2018. Young, actively growing *Myrtus communis* L. and *Metrosideros excelsa* Sol. ex Gaertn. branchlets were harvested at Cambridge University Botanic Garden, and *Plinia cauliflora* (Mart.) Kausel, *Psidium guajava* L., and *Myrcianthes pungens* (O.Berg) D.Legrand material was a gift of Pedro Araújo (University of Campinas). Combined phloem and periderm cortex material was separated from xylem/pith by peeling or scraping. After collection, all the materials were put into 96% ethanol. For all samples, alcohol insoluble residue (AIR) was prepared as described previously (Goubet *et al.*, 2009).

Plant Growth and Arabidopsis transformation

Arabidopsis plants were grown on a 9:1 mix of Levington M3 compost and vermiculite and grown at 21°C under a 16 h/8 h photoperiod. Constructs used for *A. thaliana* Xylan ArabinoPyranosylTransferase 1 (*At*XAPT1), *Eucalyptus grandis* W.Hill ex Maiden (*Eg*XAPT), and Xylan GalactoPyranosylTransferase (*Eg*XLPT) overexpression in *xapt1* and WT plants were prepared using OpenPlant common syntax GoldenGate assembly (Patron *et al.*, 2015). *AtXAPT1* (*At1g68470*), *EgXAPT* (*Eucgr.D00738*; GenBank: XM_010054097.3), and *EgXLPT* (*Eucgr.H00343*; GenBank: XM_010070979.3) coding sequences and the *CESA3* and *IRX3* promoter sequences were synthesised *de novo* by IDT or GeneWiz after removal of

BsaI and *BpiI* restriction sites. Arabidopsis transformation was performed using the floral dip

method as previously described (Clough & Bent, 1998).

Phylogenetic and bioinformatics analysis

For the XAPT phylogeny, XAPT orthologues from a wide range of tracheophyte species

were extracted from our recent phylogeny of GT47-A enzymes (Yu *et al.*, 2022), which

contained protein sequences from Plaza Dicots 4.5, Plaza Monocots 4.5, and Plaza

Gymnosperms 1.0 (Van Bel *et al.*, 2018), as well as from several independent genomes. To

obtain further XAPT orthologues from plants in the Myrtales order, the genomes of

Metrosideros polymorpha (Izuno et al., 2016; Izuno et al., 2019), *Rhodamnia argantea*,

Syzygium oleosum, and *Punica granatum* (all available from NCBI Genome,

https://www.ncbi.nlm.nih.gov/genome/) and the transcriptome of *Oenothera rosea* (Carpenter

et al., 2019; One Thousand Plant Transcriptomes, 2019), were searched using TBLASTN

(Altschul *et al.*, 1990) using *At*XAPT1 as a query with a cut-off at $E = 1 \times 10^{-125}$. Sequences

were aligned using MUSCLE (Edgar, 2004a; Edgar, 2004b) and truncated to the GT47

domain. ProtTest3 (Darriba *et al.*, 2011) was then used to select an appropriate substitution

model, and a phylogeny was calculated using RAxML (Stamatakis, 2014) with 100 rapid

bootstraps. For the main text figure, the tree was then trimmed to species of interest using ape

5.0 (Paradis & Schliep, 2019).

For the GH30 subfamily 8 tree, CAZy sequences were downloaded from the dbCAN2 server

(http://bcb.unl.edu/dbCAN2/) (Zhang *et al.*, 2018). Using bespoke Python scripts invoking

HMMER (Eddy, 2011), GH30_8 domain-containing sequences were extracted and trimmed

to the GH30 domain. Sequences with a GH30 domain sequence shorter than 200 residues

were omitted. Extremely similar sequences were removed using CD-HIT (Li & Godzik,

2006; Fu *et al.*, 2012) with a 99 % similarity threshold. The remaining sequences were

aligned using MAFFT (Katoh & Standley, 2013); ProtTest3 was then used to select an

appropriate substitution model, and a phylogeny was calculated using RAxML with 100 rapid

bootstraps.

Genomic DNA extraction and sequencing of native *XAPT* and *XLPT* loci

E. dalrympleana leaves were ground in liquid nitrogen and genomic DNA was extracted as

previously described (James *et al.*, 2001), supplementing the extraction buffer with 3 %

polyvinylpyrrolidone. PCR amplification was achieved using Q5 High-Fidelity DNA

Polymerase (New England Biolabs, Ipswich, MA, USA) according to manufacturer's

instructions. Primers are listed in Table S1. PCR products were purified by gel extraction and

isopropanol precipitation before Sanger sequencing by the DNA Sequencing Facility of the

Department of Biochemistry, University of Cambridge.

Enzyme Digestions

Soluble hemicelluloses were prepared from AIR as previously described (Lyczakowski *et al.*,

2017). For Arabidopsis bottom stem and *E. dalrympleana* xylan analysis, 0.5 mg AIR was

used for one reaction. For Arabidopsis young stem, 1 mg AIR was used for one reaction.

Soluble hemicellulose fraction was digested with an excess of Xyn11A from *Neocallimastix*

patriciarum (*Np*Xyn11A, Genbank: CAA46498.1; from Megazyme, Ireland) in 50 mM

ammonium acetate, pH 6.0 at 30°C overnight.

Alternatively, xylan was digested with GH30 xylanases with a 70 μg ml^{-1} concentration from

different bacteria at 25°C for 30 min in 50 mM ammonium acetate, pH 6.0. The GH30

xylanases included: XynC from *Bacteroides ovatus* (*Bo*XynC, GenBank: ALJ48333.1; a gift

of Harry Gilbert, Newcastle University); Xyn30A from *Hungateiclostridium thermocellum*

(*Ct*Xyn30A, GenBank: ABN54208.1; from NZYTech, Portugal). XynA from *Dickeya*

chrysanthemi strain D1 (*Ec*$_D$Xyn30A, GenBank: AAB53151.1), XynA from *Dickeya*

chrysanthemi strain P860219 (*Ec*$_P$Xyn30A, GenBank: AAL16415.1), and the two

mutagenized enzymes *Ec*$_D$Xyn30A$_{Y255L}$ and *Ec*$_P$Xyn30A$_{L255Y}$ were provided by Novozymes

A/S (Bagsværd, Denmark). For mass spectrometry and galactosidase sensitivity experiments,

xylanase products were then digested overnight with 10 μL of 1 mg ml^{-1} *B. ovatus* GH115

glucuronidase (*Bo*Agu115A, GenBank: EDO10816.1; a gift of Harry Gilbert, Newcastle

University) to remove the non-decorated GlcA side chains.

For the galactosidase sensitivity assay, the xylanase products were passed through a 3 kDa

Nanosep centrifugal device. Next, ethanol was added to a final concentration to 65% before

incubation at −20°C for 2 h. The samples were then centrifuged at 12,000g for 10 min; the

supernatant was subsequently dried 60°C *in vacuo* and resuspended in 200 µl of 50 mM
ammonium acetate, pH 6.0. To this, 0.5 µl of 4 kU ml^{-1} GH35 β-galactosidase from
Aspergillus niger (Megazyme, Ireland) was added and incubated at 37^{o}C for 18 h. Samples
were then dried at 60°C *in vacuo*.

Oligosaccharides Fingerprint Analysis by PACE

224 Samples and (Xyl)$_{1-6}$ standards (Megazyme, Ireland) were derivatized with 8-
225 aminonapthalene-1,3,6-tresulphonic acid (ANTS; Invitrogen) as described previously
226 (Goubet et al., 2002). After drying, the samples were resuspended in 100 µl of 3 M urea (final
227 concentration of standards = 20 µM), of which 2 µl was loaded onto the PACE gels. The
228 samples were run and visualized as described previously (Goubet *et al.*, 2009).

Preparation of Oligosaccharides for MS

Following the enzyme digestion, released peptides and enzymes were removed using reverse-
phase Sep-Pak C18 cartridges (Waters, MA, USA) as previously described. Oligosaccharides
were reductively aminated with 2-aminobenzoic acid (2-AA), using optimized labelling
conditions as previously described (Tryfona & Stephens, 2010). The labelled samples were
then purified from reductive amination reagents using a GlycoClean S cartridge (ProZyme,
CA, USA) as described previously (Tryfona *et al.*, 2010).

Results

At1g68470 **encodes** *XYLAN ARABINOPYRANOSYLTRANSFERASE 1* (*XAPT1*) **in Arabidopsis**

We recently established that PCW xyloglucan and β-galactoglucomannan (β-GGM)
polysaccharides in Arabidopsis have significant similarities in their structure and biosynthesis
(Yu *et al.*, 2022). In particular, both polysaccharides exhibit β1,2-galactosylation of their
α1,6-linked backbone decorations (to produce Gal-β1,2-Xyl-α1,6- and Gal-β1,2-Gal-α1,6-
side chains, respectively). By comparison with previously identified xyloglucan
galactosyltransferases from the GT47-A glycosyltransferase family, MUR3 and XLT2, we
successfully predicted that the equivalent galactosyltransferase for β-GGM, MBGT1, would

 reside in the same family. Since verifying the activity of MBGT1, we were curious to see

whether other GT47-A members might work on further polysaccharides. Seeing as α-L-Ara$p$

differs from β-D-Gal by only a hydroxymethyl group, we hypothesised that α1,2-

arabinopyranosylation of PCW xylan glucuronic acid decorations (Ara$p$-α1,2-GlcA-α1,2-)

might also be mediated by a member of this family.

To test this hypothesis, we attempted to detect the loss of the PCW xylan U^{Arap} structure in T-

DNA knockout lines for the six remaining GT47-A subclade members with unknown

functions (*At2g29040/AtGT11*, *At2g32750/AtGT12*, *At2g32740/AtGT13*, *At1g68470/AtGT17*,

At4g22580/AtGT19, and *At5g41250/AtGT20*). *At2g29040/AtGT11* has since been implicated

in galactosylation of pollen tube xyloglucan (Wei *et al.*, 2021). We used the *irx9l* mutant,

which lacks PCW-type xylan (and therefore does not exhibit any U^{Arap} structure), as a control

(Mortimer *et al.*, 2015). AIR from the young stems of these mutants was incubated with

alkali to solubilize the hemicelluloses, and then digested with *Np*Xyn11A *endo*-xylanase. The

resultant oligosaccharides were analysed by PACE. As previously reported (Mortimer *et al.*,

2015), wild-type (WT) young stem yielded a series of products including xylose (X),

xylobiose (X_2), aldopentaouronic acid with (XU^mXX) or without ($XUXX$) 4-O-methyl group

on the GlcA (generically termed '$XU^{[m]}XX$'), and xylopentaose substituted with the putative

Ara$p$-GlcA- disaccharide ($XU^{Arap}XXX$). In contrast, digestion of material from the *irx9l*

mutant released xylose, xylobiose (X_2), and $XU^{[m]}XX$, but not $XU^{Arap}XXX$. Interestingly,

although we could detect $XU^{Arap}XXX$ in five of the six GT47-A mutants, this product was

also conspicuously absent for *at1g68470*, suggesting that xylan in this mutant lacks the U^{Arap}

decoration (Fig. 1a, S1). To help confirm this, WT young stem was hydrolysed with the

GH30-family xylanase Xyn30A from *Bacteroides ovatus* (*Bo*XynC), which requires a GlcA

substitution at its −2 subsite. As previously described (Mortimer *et al.*, 2015), digestion of

WT material with *Bo*XynC produced $XXXXU^{[m]}X$ ($U_1^{[m]}X_6$) and $XXXXU^{Arap}X$ ($U_1^{Arap}X_6$) as

major products, in accordance with the consistent GlcA spacing of PCW xylan. In contrast,

digestion of material from the *at1g68470* mutant released $U_1^{[m]}X_6$ but not the $U_1^{Arap}X_6$

oligosaccharide (Fig. 1b). These results suggest that *At1g68470* is required for the synthesis

of the putative Ara$p$-GlcA- disaccharide substitution on Arabidopsis PCW xylan.

To further support our conclusion, we complemented the *at1g68470* mutant with *At1g68470*

under the control of the PCW-synthesis-specific *CESA3* promoter. AIR from young stems

was digested by xylanase *Np*Xyn11A and analysed by PACE. In all three independent

transgenic lines, the XU$^{Ara p}$XXX product was detected at substantially higher levels than in
WT (Fig. 1a). Based on these data, and the previous arabinopyranose assignment, we
concluded that *At1g68470* likely encodes a xylan-specific arabinopyranosyltransferase that
catalyses the formation of an Ara$p$-α1,2-GlcA linkage. We therefore propose that *At1g68470*
is named ***XYLAN ARABINOPYRANOSYLTRANSFERASE 1*** (*XAPT1*).

xapt1 mutant plants do not exhibit a growth phenotype

We investigated the importance of U$^{Ara p}$ disaccharide substitutions on Arabidopsis PCW
xylan for plant growth through observation of the *xapt1* growth phenotype. *xapt1* plants grew
similarly to WT plants and exhibited no obvious morphological differences (Fig. S2).
However, since XAPT1 activity is similar to that of xyloglucan β-galactosyltransferase
MUR3, we reasoned that there could be some functional redundancy between the two
enzymes, as we recently showed for MUR3 and its paralogue glucomannan β-
galactosyltransferase MBGT1 (Yu *et al.*, 2022). Hence, we crossed *xapt1* with the *mur3-3*
mutant to produce double mutant plants. Nevertheless, there was no clear difference in the
growth of the *xapt1 mur3-3* double mutant and the *mur3-3* single mutant. Hence, XAPT1
function does not appear to have a clear role in plant growth.

XAPT underwent gene duplication in the Myrtales

Whereas the U$^{Ara p}$ xylan branch structure has been reported in a wide range of species,
including Arabidopsis, *Eucalyptus* spp., and non-commelinid monocots, the U^{Gal} structure
has so far only been detected in *Eucalyptus* spp. (Togashi *et al.*, 2009; Pena *et al.*, 2016).
Hence, we were interested to know if *Eucalyptus* spp. possess a unique galactosyltransferase
responsible for these decorations. In our previous study, we showed that within GT47-A,
XAPT1 belongs to a distinct subclade from xyloglucan- (MUR3, XLT2, XUT1) and β-GGM-
modifying (MBGT) enzymes (Yu *et al.*, 2022). This putative XAPT subclade (GT47-A$_V$)
contains representatives from Selaginella, gymnosperms, dicots, and non-Poaceae monocots.
Because the U$^{Ara p}$ and U^{Gal} decorations are highly similar in structure (with Ara$p$ lacking only
the C6 hydroxymethyl group of Gal), we hypothesised that the galactosyltransferase
responsible for U^{Gal} side chains might belong to the same subclade. To explore this idea, we
constructed a phylogeny of XAPT homologues from 56 species of vascular plant, including
Eucalyptus grandis, three fellow members of the Myrtaceae family (*Metrosideros*

polymorpha, *Syzygium oleosum*, and *Rhodamnia argantea*), and two members of the wider

Myrtales order (*Punica granatum* and *Oenothera rosea*). Fig. 2 presents the rooted

phylogeny, which has been trimmed for brevity (the full phylogeny can be found in Fig. S3).

We found that most of the plant genomes—including *P. granatum* and *O. rosea*—exhibited a

single gene in this subclade. In contrast, all four Myrtaceae genomes exhibited two

paralogues that appear to have arisen from a single, recent duplication event in the Myrtales

order. In the case of *E. grandis*, these two paralogues correspond to the genes *Eucgr.D00738*

and *Eucgr.H00343*. Hence, we hypothesised that Eucgr.D00738 (*Eg*XAPT, hereafter) and

Eucgr.H00343 (*Eg*XLPT) could each constitute either a xylan GlcA

arabinopyranosyltransferase or potentially a xylan GlcA galactosyltransferase.

Eucgr.D00738 and *Eucgr.H00343* encode *Eg*XAPT and *Eg*XLPT

To investigate the activities of the two Myrtaceae XAPT-related paralogues in GT47-A, we

expressed *Eg*XAPT and *Eg*XLPT under the strong, PCW-synthesis-specific *CESA3* promoter

in the *xapt1* mutant. Xylan digestion with *Np*Xyn11A xylanase revealed that disaccharide

side chain structures were restored in the young stems of two transgenic lines for each

construct (Fig. 3a), suggesting that both proteins are active and catalyse the formation of

Ara$p$-α1,2-GlcA or a similar linkage on xylan.

PCW xylan is the presumed natural substrate of XAPT in Arabidopsis, but PCW xylan

scarcity *in planta* makes its structure difficult to analyse. For this reason, we expressed the

two *E. grandis* enzymes in SCW-producing tissues, which synthesise much more xylan.

Furthermore, [Me]GlcA sidechains on SCW xylan are completely unsubstituted (Bromley *et*

al., 2013; Mortimer *et al.*, 2015), which produces an ideal background for characterisation of

*Eg*XAPT and *Eg*XLPT. The *Eg*XAPT and *Eg*XLPT coding sequences were placed under the

control of the SCW-synthesis-specific *IRX3* promoter and expressed in WT Arabidopsis, and

xylan from mature bottom-stems was digested with the glucuronoxylanase *Bo*XynC and

analysed by PACE. As described previously (Bromley *et al.*, 2013), digestion of the WT

xylan released products decorated with unsubstituted GlcA ($U_1^{[m]}X_n$), whose sizes

corresponded to expected [Me]GlcA spacings (Fig. 3b). In the *Eg*XAPT- and *Eg*XLPT-

expressing lines, however, similar analysis revealed very different oligosaccharide products,

suggesting that most of the [Me]GlcA had been modified, perhaps into $U^{Arap/Gal,[m]}$ structures.

The spacing between GlcA decorations along the xylan backbone did not appear to be

affected in these transgenic plants. Interestingly, oligosaccharides with substituted GlcA in *Eg*XLPT-expressing lines migrated marginally slower than those in *Eg*XAPT-expressing lines, suggesting that the products of *Eg*XAPT and *Eg*XLPT might differ subtly in structure.

To further characterise the products of *Eg*XAPT and *Eg*XLPT, xylan from the relevant transgenic lines was digested simultaneously with *Np*Xyn11A and a *Bacteroides ovatus* GH115 (*Bo*GH115) α-glucuronidase, which removes unsubstituted GlcA side chains. As expected, PACE analysis indicated that WT bottom-stem xylan was fully digested into xylose and xylobiose (Fig. S4). In the *Eg*XAPT- and *Eg*XLPT-expressing lines, however, a band of higher molecular weight ($XU^{Arap,[m]}XXX/XU^{Gal,[m]}XXX$) was produced. To distinguish between any arabinopyranosylated and galactosylated oligosaccharides, we then treated these products with a β-galactosidase (*An*GH35). Whereas the higher-molecular-weight band from *Eg*XAPT-expressing lines was completely insensitive to *An*GH35 (revealing it to be entirely $XU^{Arap,[m]}XXX$), almost all of the equivalent product from *Eg*XLPT-expressing lines shifted to $XU^{[m]}XXX$ after β-galactosidase treatment—revealing its identity as $XU^{Gal,[m]}XXX$ and indicating the presence of the U^{Gal} structure in these plants. These results suggest that *Eg*XLPT, but not *Eg*XAPT, is capable of transferring galactose to GlcA side chains.

To confirm these assignments, we analysed *Bo*XynC glucuronoxylanase products by mass spectrometry (MS), which can distinguish between pentose and hexose substituents. To prevent further ambiguity between $U_1^{[m]}X_n$ and $U_1^{Arap,[m]}X_{n-1}$, *Bo*XynC products were treated with α-glucuronidase *Bo*GH115 to remove unsubstituted GlcA residues, thereby leaving $U_1^{Arap/Gal}X_n$ oligosaccharides as the only acidic products. The resulting oligosaccharides were derivatised with 2-aminobenzoic acid (2-AA) and then analysed by MALDI-TOF MS (Fig. 4, S5). As predicted, only xylo-oligosaccharides were produced from WT, confirming full removal of unsubstituted $^{[Me]}$GlcA by the α-glucuronidase. Products from plants expressing *At*XAPT under the same promoter were analysed in parallel for comparison. Lines expressing *At*XAPT and *Eg*XAPT exhibited abundant amounts of $U_1^{Arap}X_n$ and low amounts of $U_1^{Arap,m}X_n$, confirming that *Eg*XAPT is a xylan arabinopyranosyltransferase. In lines expressing *Eg*XLPT, however, the dominant sugars released could be assigned as $U_1^{Gal}X_n$, with a smaller presence of $U_1^{Gal,m}X_n$. Interestingly, this approach also detected low amounts of $U_1^{Arap}X_n$ and $U_1^{Arap,m}X_n$ in all *Eg*XLPT independent lines (Fig. 4, S5). These results suggest that *Eg*XLPT is primarily a xylan galactosyltransferase with a low level of arabinopyranosyltransferase activity, at least when expressed in Arabidopsis.

Different *Eucalyptus dalrympleana* tissues contain structurally distinct xylans

Although U^{Arap} and U^{Gal} side chains are known to exist in plants from the *Eucalyptus* genus (Togashi *et al.*, 2009; Pena *et al.*, 2016), the tissue distribution of these structures is unknown, limiting our insight into their function. To investigate where the enzymes might be active, we studied expression data available for *EgXAPT* and *EgXLPT* at https://EucGenIE.org (Hefer et al., 2011), which provides mRNA-Seq datasets for both *E. grandis* and the hybrid *E. grandis* × *E. urophylla*. Despite some potential disagreement between the two datasets, a clear pattern was still apparent: whereas *EgXAPT* expression is relatively constant and lowest in xylem tissues, the expression of *EgXLPT* is much stronger in xylem tissues than in leaves, flowers, and shoot tips (Fig. S6, S7). Interestingly, the expression of *EgXLPT* is dramatically higher in *E. grandis* × *E. urophylla* tension wood xylem than control xylem, as has been previously noted (Mizrachi *et al.*, 2015).

To study the predicted xylan structure distributions *in planta*, we harvested an actively growing branch of *Eucalyptus dalrympleana* (mountain gum). To confirm that the *XAPT* and *XLPT* genes of *E. dalrympleana* do not differ substantially to those of *E. grandis*, we PCR-amplified them from genomic DNA using homology-based primers and sequenced the two intronless coding sequences, revealing 98% (XAPT) and 99% (XLPT) protein identity between the *E. dalrympleana* and *E. grandis* sequences (Fig. S8, S9). Next, to determine the extent of GlcA substitution amongst *E. dalrympleana* tissues, we sectioned the branch into cortex, phloem, sapwood/xylem, and leaf. Major leaf veins were removed to avoid contamination from vascular tissue (Fig. 5a). Alkali-treated AIR from each tissue was digested with *Bo*XynC xylanase and analysed with PACE. Based on the PACE xylo-oligosaccharide profiles, the tissues broadly fell into two groups with very different xylan structures (Fig. 5b). In the first group, which comprises leaf and phloem, $U_1^{[m]}X_6$ and $U_1^{Arap/Gal,[m]}X_6$ were the most abundant xylan oligosaccharides, with traces of $U_1^{[m]}X_5$ and $U_1^{Arap/Gal,[m]}X_5$. In the other group, which comprises cortex and xylem, the oligosaccharides were dominated by xylo-oligosaccharides decorated with single $^{[Me]}$GlcA. The cortex and xylem xylan had a GlcA spacing pattern very similar to SCW xylan from Arabidopsis (Bromley *et al.*, 2013). Small amounts of $U_1^{Arap/Gal,[m]}X_5$ and $U_1^{Arap/Gal,[m]}X_6$ oligosaccharide could also be detected in cortex and xylem.

In order to discriminate U^{Arap} and U^{Gal} structures in different Eucalyptus tissues, we again treated *Bo*XynC xylanase products with α-glucuronidase GH115 and analysed the remaining

acidic products by MS. The MS profile of xylan showed that neutral and acidic

oligosaccharides were released from all tissues but with different ratios (Fig. 6). Consistent

with the expression data (Fig. S6, S7), leaf tissue released $U_1^{\mathrm{Ara}p,\mathrm{m}}X_5$ and $U_1^{\mathrm{Ara}p,\mathrm{m}}X_6$

oligosaccharides but no $U_1^{\mathrm{Gal},\mathrm{m}}X_n$ oligosaccharides. In contrast, the stem tissues yielded

detectable levels of $U_1^{\mathrm{Gal},\mathrm{m}}X_5$ and $U_1^{\mathrm{Gal},\mathrm{m}}X_6$. From cortex to phloem to xylem, the percentage

of GlcA residues that were substituted decreased, while at the same time, the ratio of Gal to

Ara$p$ decorations increased. We did not detect any structures containing unmethylated GlcA.

In summary, both the patterning and substitution of xylan GlcA side chains varies in different

E. dalrympleana tissues and appears to correspond with the previously observed expression

patterns of *Eg*XAPT and *Eg*XLPT in *E. grandis* / *E. grandis x E. urophylla*. More

specifically, XAPT appears to be responsible for arabinosylating glucuronoxylan mainly in

certain PCW-rich tissues, whereas glucuronoxylan galactosylation by XLPT appears to be

enriched in woody tissues but is less abundant overall.

Other Myrtaceae plants contain U^{Gal} and $U^{\mathrm{Ara}p}$ structures in their xylan

The results of our phylogenetic analysis suggested that other genera in the Myrtaceae family

might possess XLPT orthologues, and therefore might exhibit U^{Gal} structures on their xylan.

To investigate this hypothesis, we sectioned and analysed tissue from five additional

Myrtaceae members: *Myrtus communis* (myrtle), *Metrosideros excelsa* (pohutukawa), *Plinia*

cauliflora (jaboticaba), *Psidium guajava* (guava), and *Myrcianthes pungens* (guabiju). Xylan

extracted from both cortex/phloem (combined) and xylem/pith was treated with xylanase

*Np*Xyn11A and α-glucuronidase *Bo*GH115, and as before, the products were subsequently

digested with *An*GH35 β-galactosidase. In xylem/pith, $XU^{\mathrm{Ara}p,[\mathrm{m}]}XXX/XU^{\mathrm{Gal},[\mathrm{m}]}XXX$

oligosaccharides were detectable in all species, with *M. communis* exhibiting remarkably

high levels of GlcA substitution (Fig. S10). Notably, for *M. communis*, *P. cauliflora*, *P.*

guajava, and *M. pungens* xylan, but not *M. excelsa*, a large proportion of this

$XU^{\mathrm{Ara}p,[\mathrm{m}]}XXX/XU^{\mathrm{Gal},[\mathrm{m}]}XXX$ product was sensitive to *An*GH35, indicating that this band

was primarily composed of the $XU^{\mathrm{Gal},[\mathrm{m}]}XXX$ oligosaccharide with a smaller amount of

$XU^{\mathrm{Ara}p}XXX$ in xylem (Fig. S10). In cortex/phloem, $XU^{\mathrm{Ara}p,[\mathrm{m}]}XXX/XU^{\mathrm{Gal},[\mathrm{m}]}XXX$

oligosaccharides were detectable in all species except in *P. cauliflora* (Fig. S11). However,

only *M. communis* yielded products that were sensitive to *An*GH35, suggesting a lack of

$U^{\mathrm{Gal},[\mathrm{m}]}$ structures in the cortex/phloem of the other four species. Our results therefore support

the idea that many Myrtaceae plants exhibit U^{Gal} side chains and that this structure is enriched

in woodier tissues.

GH30-family glucuronoxylanases have convergently evolved to overcome the obstacle of

glucuronyl side chain modification

Cell wall polysaccharides constitute a barrier to potential pathogens. Plant pathogens

therefore rely on polysaccharide-degrading enzymes for their virulence. In this work, we have

used two *endo*-xylanases to analyse xylan structure: *Np*Xyn11A, from CAZy family GH11,

and *Bo*XynC, a member of GH30 subfamily 8 from the gut bacterium *B. ovatus*. In order to

act on xylan, the latter of these specifically recognises and binds to GlcA substitutions.

Curiously, previously solved crystal structures of related GH30 xylanases from plant

pathogen *Dickeya chrysanthemi* (previously *Erwinia chrysanthemi*) strain D1 (Ec_DXyn30A)

and *Bacillus subtilis* (*Bs*XynC) show that both enzymes make a hydrogen bond to the C2

hydroxyl of GlcA using the phenol group of Tyr_{255} or Tyr_{231} respectively (Fig. 7a) (St John *et*

al., 2011; Urbanikova *et al.*, 2011). We reasoned that the 1,2-linked Ara*p*/Gal substitution

that decorates GlcA residues at this very hydroxyl should therefore impede Ec_DXyn30A and

*Bs*XynC from binding their substrate. In spite of this, our results clearly showed that the

related *Bo*XynC is able to bind and cleave at the $U^{Arap/Gal}$ decorations (Figs. 1b, 3b). To

determine whether Ec_DXyn30A and other GH30 xylanases might have the same activity, we

digested xylan from Arabidopsis *gux1,2* young stem with Ec_DXyn30A or *Ct*Xyn30A (from

Clostridium thermocellum). PACE analysis of the products revealed that, whereas *Bo*XynC is

capable of cleaving the xylan backbone adjacent to both substituted and unsubstituted

[Me]GlcA, Ec_DXyn30A and *Ct*Xyn30A are only able to cleave effectively at unsubstituted

[Me]GlcA decorations (Fig. 8a). These results demonstrate a hitherto unknown substrate

specificity variant amongst GH30 glucuronoxylanases.

To help understand the difference in substrate specificity between these enzymes, we aligned

their protein sequences. The alignment showed that the tyrosine residue involved in binding

the C2 hydroxyl of GlcA is also present in *Ct*Xyn30A. Interestingly, however, the

corresponding residue in *Bo*XynC is not tyrosine but leucine (Fig. 7b).

To investigate the extent of amino acid variety at this position, we conducted a phylogenetic

analysis of the entire GH30 subfamily 8. We then labelled the sequences in the tree according

to the amino acid at the position of interest. The results showed that *Bo*XynC belongs to a

distinct subgroup of enzymes with leucine rather than tyrosine at this position (Fig. 7c).
Interestingly, we also identified an enzyme from *D. chrysanthemi* P860219 that diverged
much more recently from Ec_DXyn30A than *Bo*XynC, but has independently acquired a
leucine at this position (we henceforth refer to this enzyme as Ec_PXyn30A) (Fig. 7b, c).
Although Ec_DXyn30A and Ec_PXyn30A otherwise share 82% protein sequence identity, like
*Bo*XynC, Ec_PXyn30A (but not Ec_DXyn30A) was able to cut xylan adjacent to $U^{Arap/Gal}$
decorations (Fig. 8b).

To investigate whether this single residue difference is sufficient to explain the discrepancy
in substrate specificity, we performed site-directed mutagenesis on the Ec_DXyn30A and
Ec_PXyn30A enzymes. Leu_{255} from Ec_PXyn30A was mutated to tyrosine to produce
Ec_PXyn30A$_{L255Y}$, while Tyr_{255} from Ec_DXyn30A was mutated to leucine to produce
Ec_DXyn30A$_{Y255L}$. Initially, we used the purified enzymes to digest WT bottom-stem xylan
without Ara*p*/Gal substitutions. All four enzymes released identical products of the same
quantities, suggesting that all four variants exhibit glucuronoxylanase activity (Fig. S12). We
used the purified enzymes to hydrolyse alkali-extracted xylan from plants expressing the
$_{pro}IRX3$:*EgXLPT*/*XAPT* constructs. As predicted, both Ec_DXyn30A and Ec_PXyn30A$_{L255Y}$
produced primarily xylo-oligosaccharides with GlcA residues, whereas the enzymes
Ec_PXyn30A and Ec_DXyn30A$_{Y255L}$ could release many more products with $U^{Arap/Gal}$
decorations (Fig. 8b). These data suggest that a tyrosine residue at this position prevents
efficient activity on xylan substrates with substituted GlcA decorations (or that leucine is
required at this position for activity), and that multiple GH30 glucuronoxylanase subfamilies
have independently adapted to accommodate substituted GlcA decorations.

Discussion

In this work, we have identified the enzymes responsible for decorating GlcA xylan side
branches in Arabidopsis and *Eucalyptus* spp.. All three enzymes belong to subclade V of the
GT47-A family (Yu *et al.*, 2022). Interestingly, all known activities from this family involve
the formation of a 1,2 glycosidic bond between the donor sugar and an α-linked backbone
substitution of a hemicellulose. Furthermore, XAPT1 is the third GT47-family enzyme found
to be involved in xylan synthesis, with IRX10 and XYS1/IRX10L already belonging to
GT47-D (Li *et al.*, 2004; Brown *et al.*, 2009).

The identity of the pentosyl residue decorating GlcA branches in Arabidopsis PCW xylan has

not been fully confirmed: previous attempts to identify it by NMR could not distinguish

between α1,2-linked arabinopyranose and β1,2-linked xylose. Nevertheless, the former is

favoured because the decoration is insensitive to β-xylosidase (Mortimer *et al.*, 2015).

Furthermore, the equivalent pentosyl decorations in *Eucalyptus grandis* and non-commelinid

monocots have been unambiguously determined to be α1,2-linked arabinopyranose (Pena *et*

al., 2016). The close evolutionary relationship between the Arabidopsis and *E. grandis* XAPT

enzymes further strengthens the idea that the Arabidopsis pentosyl decoration is indeed

arabinopyranose.   Hence,   we   are   confident   in   assigning   *At*XAPT1   as   an

arabinopyranosyltransferase.

The identification of XAPT and XLPT enzymes in GT47-A$_V$ helps to explain the distribution

of substituted GlcA xylan side chains amongst plant species. Our previous phylogeny (Yu *et*

al., 2022) demonstrated that GT47-A$_V$ genes have been completely lost from grass (Poaceae)

genomes. This explains why U$^{\text{Ara}p}$ side chains have been observed in non-commelinid

monocot orders (Asparagales, for instance), but never in grasses. Traditionally, a distinction

has been made between commelinid and non-commelinid plants in terms of their cell wall

structures; however, our phylogeny shows that even some members of the commelinid Poales

order (*Ananas comosus*, for example) exhibit a XAPT orthologue. In addition, our work

explains the emergence of U$^{\text{Gal}}$ side chains in the Myrtaceae family. We conclude that such

structures arose from a recent duplication and neofunctionalisation of the *XAPT* locus in a

subgroup of the Myrtales or Myrtaceae. Whether galactosyl decorations have any functional

distinction from arabinopyranosyl decorations is unclear, especially as we did not observe a

difference in preference between the two by any GH30 glucuronoxylanase. It is possible that

the difference in sugar substitution could have consequences for the action of individual *exo*-

glycosidases—be they microbial or cell wall remodelling enzymes (such as the β-

galactosidase MUM2).

Unlike arabinoxylan, glucuronoxylan is present to at least some extent in virtually every land

plant cell wall. Therefore, the properties of GlcA may have a specific and important role in

cell wall structure. For instance, in secondary cell walls, GlcA has been reported to form the

covalent bridge to lignin polymers *via* its C6 carboxylate (Giummarella *et al.*, 2019; Terrett

& Dupree, 2019). The GlcA may also be important, alongside acetylation, for xylan solubility

and assembly onto cellulose (Schultink *et al.*, 2015). Further modification of GlcA by 2-*O*-

arabinopyranosylation, 2-*O*-galactosylation, and/or 4-*O*-methylation could therefore

potentially modulate xylan incorporation into the cell wall. At the same time, the U^{Arap} and

U^{Gal} disaccharide side chains resemble the functionally important side chains of other PCW

hemicelluloses: namely, the Gal-β1,2-Xyl-α1,6- and Gal-β1,2-Gal-α1,6- side chains of

xyloglucan and β-GGM, respectively. Computational studies of xyloglucan oligosaccharides

have suggested that the β1,2-linked galactosyl residues interact with the polysaccharide

backbone and weaken its binding to cellulose microfibrils (Zhao *et al.*, 2014). The galactosyl

residues may also influence xyloglucan solubility during biosynthesis (Tamura *et al.*, 2005;

Kong *et al.*, 2015). Whether decoration of GlcA could have an effect on xylan–cellulose

interactions will require extensive computational and solid state NMR work to establish.

Despite these possibilities, but consistent with GlcA methyltransferase mutants (Urbanowicz

et al., 2012; Li *et al.*, 2013) and glucuronyltransferase mutants (Mortimer *et al.*, 2015), we

did not observe any obvious phenotype of *xapt* or *xapt mur3-3* plants, which fails to support

an important function for such decorations in normal plant growth. Nevertheless, we

observed an interesting interplay between the two types of GlcA modification: whereas all

GlcA residues in the *E. dalrympleana* disaccharide side chains were methylated, the

equivalent GlcA residues in our *Eg*XAPT and *Eg*XLPT transgenic Arabidopsis lines were

almost completely unmethylated. This might suggest that Arabidopsis GlcA 4-*O*-

methyltransferases do not accept substituted GlcA residues. Furthermore, some clues for the

function of U^{Arap} and U^{Gal} structures may lie in their uneven tissue distribution. Based on

previous comparisons between 'young' and 'old' stems in Arabidopsis, U^{Arap} structures have

so far been assumed to be a general feature of 'PCW xylan' (Mortimer *et al.*, 2015).

However, in the Myrtaceae family at least, we have shown that the distribution of decorations

is likely to be more complicated than that. For example, while U^{Arap} structures could be found

in typically PCW-rich organs and tissues such as leaf, cortex, and phloem, U^{Gal} structures

were specifically enriched in xylem/pith but completely absent from leaves. Furthermore,

given that our dissections lacked the precision to separate individual cell types, it is possible

that substitution of GlcA is limited to specialised cells within the tissues of interest. Single-

cell expression atlases for Arabidopsis stem (Shi *et al.*, 2021) and leaf (Kim *et al.*, 2021) lend

support to this idea by suggesting that *At*XAPT expression may be selectively enriched in cell

types such as cambium, phloem parenchyma, and/or stem epidermis. Moreover, although we

did not investigate the xylan structures *in planta*, *Eg*XLPT appears to be specifically

upregulated in tension wood xylem tissues—perhaps by as much as a 50-fold increase

(Mizrachi *et al.*, 2015)—hinting at a specific structural role in the cell wall. As previously
proposed, it is possible that xylan structure could be remodelled in tension wood, perhaps in a
way that affects its incorporation into the wall.

Whether or not the GlcA branches on xylan play a structural role in the cell wall, their
absence leads to a major reduction in biomass resistance to enzymatic saccharification
(Lyczakowski *et al.*, 2017), suggesting that they may constitute a direct or indirect factor in
plant-pathogen interaction. Here, we have uncovered that the substitution of GlcA residues
likely affects the ability of GH30-family glucuronoxylanases to degrade xylan. Remarkably,
different lineages of this family appear to have convergently overcome the obstacle posed by
C2 modification through a single point mutation. Interestingly, the strain of *D. chrysanthemi*
whose glucuronoxylanase has recently acquired tolerance for these decorations, P860219,
was apparently cultured from *Anthurium* sp.: a non-commelinid from the Alismatales order
whose close relatives exhibit abundant $U^{Ara\textit{p}}$ xylan side chains (Hurlbert & Preston, 2001).
This is in contrast to substitution-intolerant strain D1, which was cultured from corn
(Vroemen *et al.*, 1995), a member of the Poaceae that lacks any substitutions. This hints that
cell wall structures could be applying a direct selection pressure on xylanase evolution.

In summary, we have identified one of the missing components of xylan synthesis in higher
plants. The identification of XAPT1 permits engineering of existing xylan polysaccharides
using synthetic biology approaches. Furthermore, by revealing previously unknown substrate
specificities in GH30 glucuronoxylanases, our results could facilitate optimisation of
industrial lignocellulosic biomass degradation. These results hint at the constant evolutionary
battle between plants and pathogens.

<u>Acknowledgements and funding</u>

We thank Paul Stearn (Jesus College, Cambridge), Ángela Cano and Sam Brockington
(Cambridge University Botanic Garden), and Pedro Araújo (University of Campinas) for
providing plant material.

This work was supported by the Leverhulme Trust Centre for Natural Material Innovation,
the Biotechnology and Biological Sciences Research Council (BBSRC) of the UK as part of
the OpenPlant Synthetic Biology Research Centre (Reference BB/L014130/1), The UKRI
grant underwriting the ERC advanced grant (EVOCATE Function and evolution of plant cell

wall architecture for sustainable technologies. EP/X027120/1), Innovate UK, the Cambridge

BBSRC-DTP Programme (Reference BB/J014540/1), a Novo Nordisk Foundation grant

Oxymist (Grant Number NNF20OC0059697), and a BBSRC iCASE studentship (Reference

BB/M015432/1).

Competing interests: K.B.R.M.K. is an employee of Novozymes, a major enzyme

producing company. The remaining authors declare no conflict of interests.

Author contributions

LY conceived the original screen, designed the study, performed most polysaccharide

analyses and plant transformations, and co-wrote the paper; LFLW conducted the

phylogenetic analyses, contributed to the biochemical analysis, designed the xylanase point

mutations, and co-wrote the paper; OMT carried out the molecular biology, conducted

xylanase expression and purification, and co-wrote the paper; JWR identified differences in

xylanase activities and supported the biochemical analysis and identification of Arabidopsis

T-DNA insertion lines; JJL supported the biochemical analysis and molecular biology; XY

made the double mutant; KBRMK contributed to the design and production of mutated

xylanases; PD designed and supervised the study, interpreted data and co-wrote the paper.

Data Availability

607   The data that support the findings of this study are available in the main text and in the

608   Supporting Information for this article. Sequences for newly identified *Eucalyptus*

609   *dalrympleana* XAPT and XLPT will be deposited in GenBank after acceptance. All plant

610   material descried in this manuscript is available on a reasonable request from the

611   corresponding authors.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
search tool. *Journal of Molecular Biology* **215**(3): 403-410.
Anders N, Wilson LFL, Sorieul M, Nikolovski N, Dupree P. 2022. β-1, 4-Xylan backbone
synthesis in higher plants: How complex can it be? *Frontiers in Plant Science* **13**.
Bellincampi D, Cervone F, Lionetti V. 2014. Plant cell wall dynamics and wall-related
susceptibility in plant–pathogen interactions. *Frontiers in Plant Science* **5**: 228.

Bromley JR, Busse-Wicher M, Tryfona T, Mortimer JC, Zhang ZN, Brown DM,
Dupree P. 2013. GUX1 and GUX2 glucuronyltransferases decorate distinct domains
of glucuronoxylan with different substitution patterns. *Plant Journal* **74**(3): 423-434.
Brown DM, Zhang Z, Stephens E, Dupree P, Turner SR. 2009. Characterization of IRX10
and IRX10-like reveals an essential role in glucuronoxylan biosynthesis in
Arabidopsis. *Plant Journal* **57**(4): 732-746.
Burton RA, Gidley MJ, Fincher GB. 2010. Heterogeneity in the chemistry, structure and
function of plant cell walls. *Nature Chemical Biology* **6**(10): 724-732.
Busse-Wicher M, Gomes TCF, Tryfona T, Nikolovski N, Stott K, Grantham NJ, Bolam
DN, Skaf MS, Dupree P. 2014. The pattern of xylan acetylation suggests xylan may
interact with cellulose microfibrils as a twofold helical screw in the secondary plant
cell wall of Arabidopsis thaliana. *Plant Journal* **79**(3): 492-506.
Carpenter EJ, Matasci N, Ayyampalayam S, Wu S, Sun J, Yu J, Jimenez Vieira FR,
Bowler C, Dorrell RG, Gitzendanner MA, et al. 2019. Access to RNA-sequencing
data from 1,173 plant species: The 1000 Plant transcriptomes initiative (1KP).
Gigascience **8**(10).
Chmelová D, Škulcová D, Ondrejovic M. 2019. Microbial xylanases and their inhibition by
specific proteins in cereals. *Kvasny Prumysl* **65**(4): 127–133-127–133.
Chong SL, Koutaniemi S, Juvonen M, Derba-Maceluch M, Mellerowicz EJ, Tenkanen
M. 2015. Glucuronic acid in Arabidopsis thaliana xylans carries a novel pentose
substituent. *International Journal of Biological Macromolecules* **79**: 807-812.
Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant Journal* **16**(6): 735-743.
Darriba D, Taboada GL, Doallo R, Posada D. 2011. ProtTest 3: fast selection of best-fit
models of protein evolution. *Bioinformatics* **27**(8): 1164-1165.
Eddy SR. 2011. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**(10): e1002195.
Edgar RC. 2004a. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. *BMC Bioinformatics* **5**: 113.
Edgar RC. 2004b. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Research* **32**(5): 1792-1797.
Evtuguin DV, Tomas JL, Silva AMS, Neto CP. 2003. Characterization of an acetylated
heteroxylan from Eucalyptus globulus Labill. *Carbohydrate Research* **338**(7): 597-
604.
Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**(23): 3150-3152.
Giummarella N, Pu YQ, Ragauskas AJ, Lawoko M. 2019. A critical review on the
analysis of lignin carbohydrate bonds. *Green Chemistry* **21**(7): 1573-1595.
Goubet F, Barton CJ, Mortimer JC, Yu XL, Zhang ZN, Miles GP, Richens J, Liepman
AH, Seffen K, Dupree P. 2009. Cell wall glucomannan in Arabidopsis is synthesised
by CSLA glycosyltransferases, and influences the progression of embryogenesis.
Plant Journal **60**(3): 527-538.
Grantham NJ, Wurman-Rodrich J, Terrett OM, Lyczakowski JJ, Stott K, Iuga D,
Simmons TJ, Durand-Tardif M, Brown SP, Dupree R, et al. 2017. An even
pattern of xylan substitution is critical for interaction with cellulose in plant cell walls.
Nature Plants **3**(11): 859-865.
Han MH, Liu YT, Zhang FL, Sun DF, Jiang JX. 2020. Effect of galactose side-chain on
the self-assembly of xyloglucan macromolecule. *Carbohydrate Polymers* **246**.
Hefer C, Mizrachi E, Joubert F, Myburg A 2011. The Eucalyptus genome integrative
explorer (EucGenIE): a resource for Eucalyptus genomics and transcriptomics. *BMC*
Proceedings: Springer. 1-1.

Hurlbert JC, Preston JF, 3rd. 2001. Functional characterization of a novel xylanase from a
corn strain of Erwinia chrysanthemi. *Journal of Bacteriology* **183**(6): 2093-2100.
Immelmann R, Gawenda N, Ramirez V, Pauly M. 2023. Identification of a xyloglucan
beta-xylopyranosyltransferase from Vaccinium corymbosum. *Plant Direct* **7**(7): e514.
Izuno A, Hatakeyama M, Nishiyama T, Tamaki I, Shimizu-Inatsugi R, Sasaki R,
Shimizu KK, Isagi Y. 2016. Genome sequencing of Metrosideros polymorpha
(Myrtaceae), a dominant species in various habitats in the Hawaiian Islands with
remarkable phenotypic variations. *Journal of Plant Research* **129**(4): 727-736.
Izuno A, Wicker T, Hatakeyama M, Copetti D, Shimizu KK. 2019. Updated Genome
Assembly and Annotation for Metrosideros polymorpha, an Emerging Model Tree
Species of Ecological Divergence. *G3 (Bethesda)* **9**(11): 3513-3520.
James CM, Barrett JA, Russell SJ, Gibby M. 2001. A rapid PCR based method to
establish the potential for paternal inheritance of chloroplasts in Pelargonium. *Plant*
Molecular Biology Reporter **19**: 163-167.
Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular Biology and Evolution* **30**(4):
772-780.
Kim JY, Symeonidi E, Pang TY, Denyer T, Weidauer D, Bezrutczyk M, Miras M,
Zollner N, Hartwig T, Wudick MM, et al. 2021. Distinct identities of leaf phloem
cells revealed by single cell transcriptomics. *Plant Cell* **33**(3): 511-530.
Kong YZ, Pena MJ, Renna L, Avci U, Pattathil S, Tuomivaara ST, Li XM, Reiter WD,
Brandizzi F, Hahn MG, et al. 2015. Galactose-depleted xyloglucan is dysfunctional
and leads to dwarfism in Arabidopsis. *Plant Physiology* **167**(4): 1296-U1294.
Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**(13): 1658-1659.
Li XF, Jackson P, Rubtsov DV, Faria-Blanc N, Mortimer JC, Turner SR, Krogh KB,
Johansen KS, Dupree P. 2013. Development and application of a high throughput
carbohydrate profiling technique for analyzing plant cell wall polysaccharides and
carbohydrate active enzymes. *Biotechnology for Biofuels* **6**:94.
Li XM, Cordero I, Caplan J, Molhoj M, Reiter WD. 2004. Molecular analysis of 10
coding regions from arabidopsis that are homologous to the MUR3 xyloglucan
galactosyltransferase. *Plant Physiology* **134**(3): 940-950.
Lyczakowski JJ, Wicher KB, Terrett OM, Faria-Blanc N, Yu X, Brown D, Krogh K,
Dupree P, Busse-Wicher M. 2017. Removal of glucuronic acid from xylan is a
strategy to improve the conversion of plant biomass to sugars for bioenergy.
Biotechnology for Biofuels **10**: 224.
Mizrachi E, Maloney VJ, Silberbauer J, Hefer CA, Berger DK, Mansfield SD, Myburg
AA. 2015. Investigating the molecular underpinnings underlying morphology and
changes in carbon partitioning during tension wood formation in Eucalyptus. *New*
Phytologist **206**(4): 1351-1363.
Mortimer JC, Faria-Blanc N, Yu XL, Tryfona T, Sorieul M, Ng YZ, Zhang ZN, Stott K,
Anders N, Dupree P. 2015. An unusual xylan in Arabidopsis primary cell walls is
synthesised by GUX3, IRX9L, IRX10L and IRX14. *Plant Journal* **83**(3): 413-426.
One Thousand Plant Transcriptomes I. 2019. One thousand plant transcriptomes and the
phylogenomics of green plants. *Nature* **574**(7780): 679-685.
Paes G, Berrin JG, Beaugrand J. 2012. GH11 xylanases: Structure/function/properties
relationships and applications. *Biotechnology Advances* **30**(3): 564-592.
Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**(3): 526-528.

Patron NJ, Orzaez D, Marillonnet S, Warzecha H, Matthewman C, Youles M, Raitskin
O, Leveau A, Farre G, Rogers C, et al. 2015. Standards for plant synthetic biology:
a common syntax for exchange of DNA parts. *New Phytologist* **208**(1): 13-19.
Pena MJ, Kulkarni AR, Backe J, Boyd M, O'Neill MA, York WS. 2016. Structural
diversity of xylans in the cell walls of monocots. *Planta* **244**(3): 589-606.
Pinto PC, Evtuguin DV, Neto CP. 2005. Structure of hardwood glucuronoxylans:
modifications and impact on pulp retention during wood kraft pulping. *Carbohydrate*
Polymers **60**(4): 489-497.
Rogowski A, Briggs JA, Mortimer JC, Tryfona T, Terrapon N, Lowe EC, Basle A,
Morland C, Day AM, Zheng HJ, et al. 2015. Glycan complexity dictates microbial
resource allocation in the large intestine. *Nature Communications* **6**: 7481.
Scheller HV, Ulvskov P. 2010. Hemicelluloses. *Annual Review of Plant Biol*ogy **61**: 263-
289.
Schultink A, Cheng K, Park YB, Cosgrove DJ, Pauly M. 2013. The identification of two
arabinosyltransferases from tomato reveals functional equivalency of xyloglucan side
chain substituents. *Plant Physiology* **163**(1): 86-94.
Schultink A, Naylor D, Dama M, Pauly M. 2015. The Role of the Plant-Specific
ALTERED XYLOGLUCAN9 Protein in Arabidopsis Cell Wall Polysaccharide O-
Acetylation. *Plant Physiology* **167**(4): 1271-1283.

Shatalov AA, Evtuguin DV, Neto CP. 1999. (2-O-alpha-D-galactopyranosyl-4-O-methyl-
alpha-D-glucurono)-D-xylan from Eucalyptus globulus Labill. *Carbohydrate*
Research **320**(1-2): 93-99.
Shi D, Jouannet V, Agusti J, Kaul V, Levitsky V, Sanchez P, Mironova VV, Greb T.
2021. Tissue-specific transcriptome profiling of the Arabidopsis inflorescence stem
reveals local cellular signatures. *Plant Cell* **33**(2): 200-223.
Simmons TJ, Mortimer JC, Bernardinelli OD, Pöppler AC, Brown SP, Deazevedo ER,
Dupree R, Dupree P. 2016. Folding of xylan onto cellulose fibrils in plant cell walls
revealed by solid-state NMR. *Nature Communications* **7**: 13902.
St John FJ, Hurlbert JC, Rice JD, Preston JF, Pozharski E. 2011. Ligand bound
structures of a glycosyl hydrolase family 30 glucuronoxylan xylanohydrolase. *Journal*
of Molecular Biology **407**(1): 92-109.
Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**(9): 1312-1313.
Tamura K, Shimada T, Kondo M, Nishimura M, Hara-Nishimura I. 2005.
KATAMARI1/MURUS3 is a novel Golgi membrane protein that is required for
endomembrane organization in Arabidopsis. *Plant Cell* **17**(6): 1764-1776.
Terrett OM, Dupree P. 2019. Covalent interactions between lignin and hemicelluloses in
plant secondary cell walls. *Current Opinion in Biotechnology* **56**: 97-104.
Togashi H, Kato A, Shimizu K. 2009. Enzymatically derived aldouronic acids from
Eucalyptus globulus glucuronoxylan. *Carbohydrate Polymers* **78**(2): 247-252.
Tryfona T, Bourdon M, Delgado Marques R, Busse-Wicher M, Vilaplana F, Stott K,
Dupree P. 2023. Grass xylan structural variation suggests functional specialization
and distinctive interaction with cellulose and lignin. *Plant Journal* **113**(5): 1004-1020.
Tryfona T, Liang HC, Kotake T, Kaneko S, Marsh J, Ichinose H, Lovegrove A,
Tsumuraya Y, Shewry PR, Stephens E, et al. 2010. Carbohydrate structural
analysis of wheat flour arabinogalactan protein. *Carbohydrate Research* **345**(18):
2648-2656.

Tryfona T, Stephens E. 2010. Analysis of carbohydrates on proteins by offline normal-
phase liquid chromatography MALDI-TOF/TOF-MS/MS. *Plant Cell Wall: Methods*
and Protocols **658**: 137-151.

Urbanikova L, Vrsanska M, Morkeberg Krogh KB, Hoff T, Biely P. 2011. Structural
basis for substrate recognition by Erwinia chrysanthemi GH30 glucuronoxylanase.
FEBS Journal **278**(12): 2105-2116.

Urbanowicz BR, Pena MJ, Ratnaparkhe S, Avci U, Backe J, Steet HF, Foston M, Li HJ,
O'Neill MA, Ragauskas AJ, et al. 2012. 4-O-methylation of glucuronic acid in
Arabidopsis glucuronoxylan is catalyzed by a domain of unknown function family
579 protein. *Proceedings of the National Academy of Sciences of the United States of*
America **109**(35): 14253-14258.

Van Bel M, Diels T, Vancaester E, Kreft L, Botzki A, Van de Peer Y, Coppens F,
Vandepoele K. 2018. PLAZA 4.0: an integrative resource for functional,
evolutionary and comparative plant genomics. *Nucleic Acids Research* **46**(D1):
D1190-D1196.

Vroemen S, Heldens J, Boyd C, Henrissat B, Keen NT. 1995. Cloning and characterization
of the bgxA gene from Erwinia chrysanthemi D1 which encodes a beta-
glucosidase/xylosidase enzyme. *Molecular Genetics and Genomics* **246**(4): 465-477.

Wei QQ, Yang Y, Li H, Liu ZW, Fu R, Feng HQ, Li C. 2021. The xyloglucan
galactosylation modulates the cell wall stability of pollen tube. *Planta* **254**(6): 133-
140.

Wilson LF, Neun S, Yu L, Tryfona T, Stott K, Hollfelder F, Dupree P. 2023. The
biosynthesis, degradation, and function of cell wall β☐xylosylated xyloglucan mirrors
that of arabinoxyloglucan. *New Phytologist* **240** (6): 2353-2371.

Wu AM, Hornblad E, Voxeur A, Gerber L, Rihouey C, Lerouge P, Marchant A. 2010.
Analysis of the Arabidopsis IRX9/IRX9-L and IRX14/IRX14-L pairs of
glycosyltransferase genes reveals critical contributions to biosynthesis of the
hemicellulose glucuronoxylan. *Plant Physiology* **153**(2): 542-554.

Yu L, Yoshimi Y, Cresswell R, Wightman R, Lyczakowski JJ, Wilson LFL, Ishida K,
Stott K, Yu XL, Charalambous S, et al. 2022. Eudicot primary cell wall
glucomannan is related in synthesis, structure, and function to xyloglucan. *Plant Cell*
34(11): 4600-4622.

Zhang H, Yohe T, Huang L, Entwistle S, brown P, Yang Z, Busk PK, Xu Y, Yin Y.
2018. dbCAN2: a meta server for automated carbohydrate-active enzyme annotation.
Nucleic Acids Research **46**(W1): W95-W101.

Zhao Z, Crespi VH, Kubicki JD, Cosgrove DJ, Zhong LH. 2014. Molecular dynamics
simulation study of xyloglucan adsorption on cellulose surfaces: effects of surface
hydrophobicity and side-chain variation. *Cellulose* **21**(2): 1025-1039.

Zhu L, Dama M, Pauly M. 2018. Identification of an arabinopyranosyltransferase from
Physcomitrella patens involved in the synthesis of the hemicellulose xyloglucan.
Plant Direct **2**(3): e00046.

Figure Legends

Figure 1. AT1G68470/XAPT1 from GT47 subclade A is a functional xylan arabinopyranosyltransferase. (a), PACE analysis of *Np*Xyn11A xylanase digests of alkali-extracted xylan from AIR of WT, the *xapt1* T-DNA insertion mutant, and three independent transgenic lines expressing *XAPT1* under the strong primary wall-specific promoter of *CESA3* in the *xapt1* background. (b), PACE analysis of *Bo*XynC xylanase digests of WT and *xapt1* mutant young stem. Xylo-oligosaccharide standard: X–X_6.

Figure 2. Trimmed phylogenetic tree of the XAPT subclade of GT47-A. Sequences grouped in the XAPT subclade of our previous GT47-A tree (Yu et al., 2022) were supplemented with HMMER and TBLASTN hits for XAPT-like sequences in Myrtales genomes. Following alignment, the phylogeny was inferred using RAxML. For rooting, *At*MUR3 was included as the outgroup. Due to the large number of sequences (106), we show only an abridged form of the tree here (displaying the more pertinent sequences), hence bootstrap values (from 100 pseudoreplicates) may underestimate the confidence of the splits shown. The full phylogeny can be viewed in Fig. S3.

Figure 3. Overexpression of *Eg*XLPT or *Eg*XAPT can decorate the glucuronoxylan with an additional sugar. (a), Expression of *Eg*XLPT and *Eg*XAPT under primary cell wall *CESA3* promoter can complement the *xapt1* xylan structure phenotype. Alkali-extracted xylan from 5-week-old young stem was digested with *Np*Xyn11A xylananase and analyzed by PACE. (b), Expression of *Eg*XLPT and *Eg*XAPT under secondary wall *IRX3* promoter in Arabidopsis WT can decorate most of the glucuronoxylan with an addition sugar. Alkali-extracted xylan from 8-week-old bottom stem was digested with *Bo*XynC xylanase and analysed by PACE. Xylo-oligosaccharide standard: X–X_6.

Figure 4. *Eg*XAPT and *Eg*XLPT exhibit different donor sugar specificities. AIR was extracted from 8-week-old bottom stems of either Arabidopsis WT plants or transgenic plants expressing *Eg*XAPT or *Eg*XLPT under the secondary cell wall-specific promoter of IRX3. Alkali-extracted xylan was then digested with *Bo*Xyn30C xylanase followed by GH115 glucuronidase. The products were labelled with 2-AA and analysed by MALDI-TOF MS.

Figure 5. Xylan structure in different *Eucalyptus dalrympleana* tissues. (a), Eucalyptus stems were sectioned into cortex, phloem, and xylem/sapwood. Major leaf vein tissue was not included for the leaf xylan analysis. Scale bar = 0.9 cm. (b), PACE analysis of *Bo*XynC

xylanase digests of AIR from different tissues. Xylo-oligosaccharide standard: $X–X_6$.

$U^{Arap,Gal,[m]}X_n$ structures are labelled with arrows.

Figure 6. Distribution of xylan with decorated GlcA in different *Eucalyptus dalrympleana*

tissues. AIR from different tissues was digested with *Bo*XynC xylanase followed by GH115

glucuronidase to remove the undecorated GlcA. The products were labelled with 2-AA and

then analysed by MS.

Figure 7. Differences in GH30 xylanase active site residues may reflect differences in ability

to accommodate substituted glucuronic acid sidechains. (a), Previously characterised active

site of GH30 xylanase from plant pathogen *Dickeya chrysanthemi* (previously *Erwinia*

chrysanthemi) strain D1 (*Ec*$_D$Xyn30A; PDB: 2Y24). Bound aldotriuronic acid is shown in

stick representation. The side chain of Tyr255, which forms a hydrogen bond with the GlcA

C2 hydroxyl, is highlighted in orange. (b), Alignment segment of five GH30 xylanases,

highlighting residues aligned to Tyr255 in *Ec*$_D$Xyn30A. (c), Molecular phylogeny of GH30

subfamily 8 sequences downloaded from dbCAN2 (http://bcb.unl.edu/dbCAN2/). Sequences

that display a tyrosine residue in the column aligned to Tyr255 are highlighted in orange;

those that instead display leucine are highlighted in blue, while those with asparagine are

highlighted in purple.

Figure 8. PACE gels to show the activities of different GH30 xylanases towards substituted

glucuronoxylan. (a), PACE analysis of three GH30 xylanases-digests of xylan from *gux1*

gux2 double mutant, which only contains primary cell wall glucuronoxylan. (b), AIR from

the transgenic plants expressing *Eg*XAPT or *Eg*XLPT under the secondary cell wall-specific

promoter of *IRX3* were used as the substrates. Alkali-extracted xylan was then digested with

different GH30 and the corresponding modified GH30 enzymes. All the plant materials are

30-day-old young stem. Xylo-oligosaccharide standard: $X–X_6$.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

(a)

Xyl

Xyl

Xyl

GlcA

Me

Tyr255

2.7 Å

PDB: **2Y24**

(b)

Ec_D Xyn30A	QLWMTEH**Y**VDSKQSAN	263
Ec_P Xyn30A	QVWMTEH**L**VDSKQSAN	263
BsXynC	DLWMTEV**Y**YPNSDTNS	270
CtXyn30A	ELWMTEV**Y**VPNSDSNS	268
BoXynC	TWWMTEH**L**FNDGENSD	382

**** . : .

(c)

Residue aligned to
Tyr255 in EcDXyn30A:
tyrosine (Y)
leucine (L)
asparagine (N)
other

0.3

Figure 7.

Figure 8