Genome data artifacts and functional studies of deletion repair in the BA.1 SARS-CoV-2 spike protein

Miguel Álvarez-Herrera[a], Paula Ruiz-Rodriguez[a], Beatriz Navarro-Domínguez[ab], Joao Zulaica[a], Brayan Grau[a], María Alma Bracho[ac], Manuel Guerreiro[de], Cristóbal Aguilar-Gallardo[e], Fernando González-Candelas[ac], Iñaki Comas[cf], Ron Geller[a], Mireia Coscollá[a]*

[a]*Institute for Integrative Systems Biology (I²SysBio, University of Valencia-CSIC), FISABIO Joint Research Unit "Infection and Public Health", Paterna, Spain*

[b]*Department of Genetics, Universtiy of Granada, Granada, Spain*

[c]*CIBER in Epidemiology and Public Health (CIBERESP), Madrid, Spain*

[d]*Haematology Department, La Fe University and Polytechnic Hospital, Valencia, Spain*

[e]*La Fe Health Research Institute (IIS-La Fe), Valencia, Spain*

[f]*Tuberculosis Genomics Unit, Institute of Biomedicine of Valencia (IBV-CSIC), Valencia, Spain*

* Email: mireia.coscolla@csic.es. Address: Institute for Integrative Systems Biology, C/ Catedrático Agustín Escardino 9, Building 4, 46980 Paterna, Spain

Keywords: SARS-CoV-2, spike, transmission, deletion, fusogenicity, antibody neutralization

Abstract

Mutations within the N-terminal domain (NTD) of the spike (S) protein are critical for the emergence of successful SARS-CoV-2 viral lineages. The NTD has been repeatedly impacted by deletions, often exhibiting complex and dynamic patterns, such as the recurrent emergence and disappearance of deletions in dominant variants. This study investigates the influence of repair of NTD lineage-defining deletions found in the BA.1 lineage (Omicron variant) on viral success. We performed comparative genomic analyses of more than 10 million SARS-CoV-2 genomes from GISAID to decipher the transmission success of viruses lacking S:ΔH69/V70, S:ΔV143/Y145, or both. These findings were contrasted against a screening of publicly available raw sequencing data, revealing substantial discrepancies between data repositories, suggesting that spurious deletion repair observations in GISAID may result from systematic artifacts. Specifically, deletion repair events were approximately an order of magnitude less frequent in the read-run survey. Our results suggest that deletion repair events are rare, isolated events with limited direct influence on SARS-CoV-2 evolution or transmission. Nevertheless, such events could facilitate the emergence of fitness-enhancing mutations. To explore potential drivers of NTD deletion repair patterns, we characterized the viral phenotype of such markers in a surrogate *in vitro* system. Repair of the S:ΔH69/V70 deletion reduced viral infectivity, while simultaneous repair with S:ΔV143/Y145 led to lower fusogenicity. In contrast, individual S:ΔV143/Y145 repair enhanced both fusogenicity and susceptibility to neutralization by sera from vaccinated individuals. This work underscores the complex genotype-phenotype landscape of the spike NTD in SARS-CoV-2, which impacts viral biology, transmission efficiency, and immune escape potential, offering insights with direct relevance to public health, viral surveillance, and the adaptive mechanisms driving emerging variants.

Introduction

As we navigate the constantly shifting landscape of the COVID-19 pandemic, the remarkable potential of SARS-CoV-2 for genetic adaptation has taken centre stage. Global turnover of SARS-CoV-2 lineages happened several times, with three variants of concern (VOCs) displacing the previous predominant lineages just in 2021: first, Alpha (B.1.1.7 and Q sublineages), then Delta (B.1.617.2 and AY sublineages), and lastly, Omicron (B.1.1.529 and BA sublineages, among others) (Rambaut et al. 2020). This successive replacement of lineages throughout the pandemic suggests that the newer had a higher adaptive value than the previous ones (Z. Chen et al. 2021; da Silva Francisco Junior et al. 2021), and thus, these three lineages are supposed to carry mutations associated with higher transmissibility or immune evasion.

The spike (S) protein —a type I transmembrane N-linked glycosylated protein— is a hotspot for mutations with high adaptive value. Spike proteins are located on the surface of SARS-CoV-2, and their main role is to mediate viral cell entry (Letko et al. 2020). The spike protein forms a homotrimer, which is cleaved post-transcriptionally into two subunits: S1 and S2. The S1 consists of the amino or N-terminal domain (NTD) and the receptor-binding domain (RBD), and it is responsible for binding to the host cell-surface angiotensin converting enzyme-2 (ACE2) receptors. The S2 subunit includes the trimeric core of the protein and is responsible for membrane fusion (Harvey et al. 2021; V'kovski et al. 2021). Therefore, some amino acid changes in the S protein may confer an advantage for transmission, considering its role in mediating viral cell entry. Additionally, most antibodies target sites either at the NTD or RBD and, therefore, mutations in these regions may enhance immune escape.

Specifically, deletions may play a decisive role in SARS-CoV-2 adaptive

evolution, particularly in deletion-tolerant genome regions such as the S gene, as they

can hardly be corrected by the proofreading activity of its RNA-dependent RNA

polymerase (RdRP) (Denison et al. 2011; McCarthy et al. 2021; Minskaia et al. 2006).

Notably, the NTD has been extensively impacted by deletions, which have emerged

independently in multiple variants, often exhibiting complex and dynamic patterns,

exemplified by the recurrent emergence and disappearance of the S:ΔH69/V70 deletion

in dominant variants (McMillen et al. 2022; Meng et al. 2021). Acquisition of deletions

in the NTD of the spike glycoprotein during long-term infections of

immunocompromised patients has been reported and identified as an evolutionary

pattern defined by recurrent deletion patterns that alter defined antibody epitopes

(Avanzato et al. 2020; Choi et al. 2020). For example, different deletions are observed

in Delta (S:ΔE156/F157-R158G), Alpha (S:ΔY144) and BA.1 (S:ΔV143/Y145)

variants, all mapping to the same surface area, indicating a convergent function (Mishra

et al. 2022; Planas et al. 2021). All these mutations are within the NTD site targeted by

many anti-NTD neutralizing antibodies (McCallum, De Marco, et al. 2021). In the

Alpha variant, S:ΔH69/V70 and S:ΔY144 are directly associated with antibody escape

(Kemp et al. 2021; McCallum, De Marco, et al. 2021) and increased infectivity (Cantoni

et al. 2022). More recent instances exist as well. Despite S:ΔY144 not being found in

BA.2, its descendant lineage BJ.1 seems to have independently acquired this deletion

(see cov-lineages/pango-designation issues #915 and #922 on GitHub). Then, it was

passed on to XBB viruses through recombination with a BA.2-descended lineage, which

was associated with increased immune escape regardless of the vaccination status or

infection history (Mykytyn et al. 2023; Tamura et al. 2023; Zhang et al. 2022).

Subsequently, in January 2023, the newly emerging XBB.1.5 lineage was shown to

display an increased receptor-binding affinity and infectivity with respect to its parental

lineage (Uriu et al. 2023; Yue et al. 2023).

The BA.1 lineage exhibited a growth advantage over previous variants of

concern (Puhach et al. 2022; Ward et al. 2022) and soon took over the Delta variant.

During this time, we became intrigued by the lack of overlap in NTD mutations

between them. This is not the case for the Alpha variant and the BA.1 lineage, which

share NTD deletions. If mutations in the NTD increased immune escape without

compromising binding to ACE2, one might expect mutations in NTD to have a

cumulative ("the more the better") effect. However, that would not be the case if

epistatic interactions between sites prevented the fixation of particular mutations in

different genomic and genetic backgrounds. Our primary objective in this study was to

investigate the effect of NTD deletion repair in BA.1 background. We examined the

differences in success among three BA.1 haplotypes lacking the S:ΔH69/V70 and

S:ΔV143/Y145 deletions, individually and in combination, which had been recurrently

identified in immunocompromised individuals with chronic infections before

widespread vaccination against COVID-19 (Avanzato et al. 2020; Choi et al. 2020;

Kemp et al. 2021; McCarthy et al. 2021). Our approach integrates public data surveys

with functional evidence to better understand how rare mutational events might shape

viral evolution and pandemic dynamics.

Materials and Methods

GISAID survey data retrieval and processing

We performed two surveys to track the absence of NTD deletions S:ΔH69/V70 and

S:ΔV143/Y145 in sequences assigned to the Omicron lineage BA.1 or any of its

sublineages (hereafter referred to collectively as "BA.1"). The first survey included

11,334,504 available SARS-CoV-2 sequences fetched from the GISAID (Khare et al.

2021) EpiCoV database on June 15, 2022. The GISAID EPI_SET is available at DOI:

10.55876/gis8.230801ex. We ran Nextalign CLI v1.9.0 (Hadfield et al. 2018) with

default scoring settings to obtain aligned genomes and peptides corresponding to each

gene. Genomes from non-human hosts were discarded. Then, we filtered out genomes

containing more than 5 % ambiguous bases and 1,000 gaps, or at least one

indeterminate position among the lineage-defining sites of the spike protein. 10,353,158

genomes passed these filters. Then, BA.1 genomes were selected and classified into one

of three haplotypes, depending on their lack of S:ΔH69/V70 (RepΔ69/70),

S:ΔV143/Y145 (RepΔ143/145), or both (RepBoth). After running Nextclade CLI v3.8.2

(Aksamentov et al. 2021), we applied a BA.1 lineage filter alongside specific mutation

and motif inclusion and exclusion criteria. Manual curation followed. For RepΔ69/70,

sequences were selected if they carried S:ΔV143, S:ΔY144, either S:ΔG142 or

S:ΔY145, and a lack of S:ΔH69 and S:ΔV70. Sequences were required to match the

S:69-70:HV motif. For RepΔ143/145, sequences were selected if they carried S:ΔH69,

S:ΔV70, either S:G142D or S:Y145D, and a lack of S:ΔV143, S:ΔY144, and either

S:ΔG142 or S:ΔY145. The required motif was S:142-145:DVYY. For RepBoth,

sequences were selected if they carried S:G142D, and a lack of S:ΔH69, S:ΔV70,

S:ΔG142, S:ΔV143, S:ΔY144 and S:ΔY145. Sequences were required to match both

the S:69-70:HV and S:142-145:DVYY motifs. We introduced an additional quality

control step based on known control mutations: because BA.1 viruses typically exhibit

the ORF1a:ΔS3675/F3677 (NSP6) deletion, sequences lacking this deletion were

discarded. We used R v4.3.3 along with *tidyverse* v2.0.0 (Wickham et al. 2019) to

conduct, manage and visualize these analyses.

To assess potential artifacts in the quality of GISAID data, we performed an

initial exploratory analysis of raw sequencing reads obtained via the European

Nucleotide Archive (ENA). First, we identified GISAID genomes with corresponding

raw reads deposited in the ENA by cross-referencing GISAID identifiers (virus name

and accession number). Raw reads were downloaded and reprocessed to test the

congruence between the GISAID consensus sequences and the raw read mappings. The

mappings were manually inspected to confirm genotypes, and we calculated two

metrics for each haplotype: the ratio of fixed deletion repairs (allele frequency >0.95)

and the ratio of cases with significant deletion repair read support (allele

frequency >0.10). Both the exploratory analysis and full ENA survey workflow

described hereunder employed the same workflow outline and software.

ENA survey data retrieval and processing

Following this exploratory comparison, we extended the analysis to independently

contrast the GISAID survey. We performed a second survey of combinations of NTD

repaired deletions using SARS-CoV-2 "read run" records via the ENA Portal API. This

survey tracked the absence of deletions S:ΔH69/V70 and S:ΔV143/Y145 (separately

and combined) in sequences assigned to lineage BA.1 (Omicron variant). We searched

for runs from samples collected from 1 November 2021 to 1 August 2022, filtering by

NCBI taxonomy code (2697049 for SARS-CoV-2) and host scientific name (*Homo*

sapiens). Instrument platforms matching DNBseq, Element and capillary sequencing

were discarded. We also discarded runs with an RNAseq library strategy, as well as

transcriptomic, metagenomic and metatranscriptomic library sources. The survey

brought out 1,541,892 matching records as of 10 December 2024 (Supplemental Table

1). After filtering for data URL availability, 1,250,257 records remained. The date of

submission of these records ranged between 10 November 2021 and 1 December 2024.

We did a near 3-fold reduction of the dataset size, keeping 464,859 records selected at

random for further analysis (see also Supplemental Table 1). Then, we ran an in-house

pipeline implemented as a Snakemake (Köster & Rahmann 2012) workflow to detect

NTD repaired deletions. We ran Snakemake v8.25.3 with Python v3.12.7.

For each sample, the following steps were carried out. FASTQ files were

downloaded using the FTP URLs found in the ENA metadata. Then, read preprocessing

and quality control (QC) were performed using *fastp* v0.23.4 (S. Chen 2023) with

default settings. Filtered reads were mapped to a complete genome sequence from a

BA.1 SARS-CoV-2 isolate (GenBank accession number: OZ070629.1) with *minimap2*

v2.28 (H. Li 2018, 2021), using the recommended presets according to the software

documentation (*sr* for Illumina and Ion Torrent reads, *map-ont* for Oxford Nanopore

reads, and *map-hifi* for PacBio reads). After the mapping step, pileups were calculated

using *samtools* v1.20 (Danecek et al. 2021) and consensus genomes were obtained with

iVar v1.4.3 (Grubaugh et al. 2019) with a minimum insertion frequency of 0.9, a

minimum depth of 40 and otherwise default settings. These sequences were fed to

pangolin v4.3 (O'Toole et al. 2021) to perform lineage assignment. The output was

used to further filter the dataset, keeping records with passing *pangolin* QC and an

assignment of BA.1 lineage or Omicron (BA.1-like) constellation. Then, variant calling

was performed on the remaining samples with *iVar* v1.4.3 as well, with a minimum

frequency threshold of 0.05, a minimum depth of 40 and otherwise default settings.

Results were further annotated using *SnpEff* v5.2 and filtered with *SnpSift* v5.2

(Cingolani et al. 2012). We used a custom *SnpEff* database based on the mapping

reference. Finally, matching records were classified into one of the deletion repair

haplotypes —RepΔ69/70, RepΔ143/145 or RepBoth— while records with no evidence

of repaired deletions were discarded. We established a minimum allele frequency

threshold of 0.75 for each repaired deletion allele. For the two single repair haplotypes,

we set a maximum frequency threshold of 0.25 non-repaired deletion.

Additionally, we generated graphic visualizations of the genome region between

sites 21694 and 22010 for visual inspection using R v4.3.3 with *ape* v5.8 (Paradis &

Schliep 2019), *Rsamtools* v2.18.0 (Morgan et al. 2023), *tidyverse* v2.0.0 (Wickham et

al. 2019) and *ggpubr* v0.6.0 (Kassambara 2023). The repository at

https://github.com/PathoGenOmics-Lab/ena-spike-ntd-repdel-analysis (v1.0.0) contains

the Snakemake workflow and the code to perform the initial screening.

Number of emergences and transmission events

To infer the success of each combinatorial haplotype, we assessed the number of

minimum emergence events and whether these originated subsequent transmission

events. We placed genomes matching deletion repair haplotypes on a global-scale

mutation-annotated tree (MAT) of public sequences (McBroome et al. 2021), under a

maximum parsimony criterion, using UShER v0.6.2 (Turakhia et al. 2021). For GISAID

data, genomes were placed on a tree dated to the survey date (15 June 2022), while for

ENA data, consensus sequences inferred through our pipeline were placed on a tree

dated to the last input record (11 March 2022).

To quantify the transmission of each haplotype, we developed *tcfinder* v1.0.0, a

lightweight and efficient tool that performs an exhaustive breadth-first search of the

samples on a phylogeny. This phylogeny must be provided using the tabular *phylo4*

structure, as defined in *phylobase* v0.8.10 (https://github.com/fmichonneau/phylobase).

The tool is available in Bioconda (Grüning et al. 2018). We selected clusters containing

at least 2 members and at least 90 % target samples. This requirement was reduced from

100 % to account for potential sequencing errors and ambiguous sample placements on

the phylogenetic tree.

The minimum number of independent emergences for each haplotype was

derived from the phylogenies, by adding up the number of transmission clusters to the

number of samples that were not transmitted. To adequately compare the relative

epidemiological success of the most prevalent haplotypes, we developed a method to

estimate their transmission fitness within each cluster. This was calculated as the ratio

between the size of each transmission cluster and the number of sequences that were

collected between the first and last cases of the cluster, with a 7-day padding added to

both ends of the time window to account for missing data and short-lived clusters. This

approach effectively standardizes the size of each transmission cluster, correcting

differences in surveillance efforts and viral prevalence at different geographic regions

and time periods. In essence, it provides a specific rate of emergence and transmission

success relative to the broader population background. Consequently, for clusters

exhibiting cross-border transmission (i.e., involving samples from different countries),

we further divided the cluster time window into country-specific sub-windows. Then,

the denominator of our transmission fitness estimate was calculated as the sum of the

number of records in the data repository for each country-specific time window. For

GISAID data, the full survey was used as background. For ENA data, the 464,859

records selected at random were used. Regardless, there were no significant differences

between the distribution of fitness estimates within each haplotype, whether using

country-specific sub-windows ("slicing" method) or not ("adding" method)

(Supplemental Figure 1). Results in the main text use the first method. Differences in

the distribution of the estimated transmission fitness between haplotypes and host age

were evaluated using Wilcoxon rank-sum tests. To determine age fold changes, the ratio

of group average values was calculated. Statistical analyses were performed and

visualized using R v4.1.2 along with *tidyverse* v2.0.0 (Wickham et al. 2019) and *ggpubr*

v0.4.0 (Kassambara 2023).

The complete pipeline (including the phylogenetic placement, transmission

quantification, fitness estimation and data visualizations) is available as a Snakemake

workflow at https://github.com/PathoGenOmics-Lab/transcluster (v2.1.0). We also

studied the location and time span of the transmission clusters using the associated

sample metadata from GISAID.

Biological characterization of BA.1 deletion repair

Deletion repairs of S:ΔH69/V70 and S:ΔV143/ΔY145 were introduced individually and

in combination into a pCG1 plasmid encoding a codon-optimized BA.1 spike protein

(Giménez et al. 2022) by site-directed mutagenesis. All the constructs were verified by

Sanger sequencing. Pseudotyped vesicular stomatitis virus (VSV) encoding a GFP

reporter gene and carrying the different spike proteins was produced as previously

reported (Gozalbo-Rovira et al. 2020). To assess the effects on virus production,

pseudotyped VSV carrying each construct were produced independently three times.

The resulting viruses were then titrated by infecting Vero E6 cells (kindly provided by

Dr. Luis Enjuanes; CNB-CSIC, Spain) or Vero E6-TMPRSS2 cells (JCRB Cell bank

catalogue code: JCRB1819) for 16 hours, followed by quantification of GFP-expressing

infected cells using a live cell microscope (Incucyte SX5; Sartorius) to obtain the

number of focus forming units (FFU) per millilitre. To assess thermal stability, 500

FFU of these pseudotyped viruses were incubated for 15 min at a range of temperature

in a thermal cycler (30.4, 31.4, 33.0, 35.2, 38.2, 44.8, 47.0, 48.6 and 49.6 °C; Biometra

T one Gradient, Analytik Jena) and the surviving virus was used to infect VeroE6-

TMPRSS2 cells for 16 h. The GFP signal in each well was then determined using a live-

cell microscope (Incucyte SX5, Sartorius). The average GFP signal observed in mock-

infected wells was subtracted from all infected wells, followed by standardization of the

GFP signal to the average GFP signal from wells incubated at 30.4 °C. Finally, we fitted

a three-parameter log-logistic function to the data using the *drc* v3.0-1 R package (*LL.3*

function) and calculated the temperature resulting in 50 % reduction in virus infection

(*ED* function). To assess the effects on neutralization by polyclonal sera, we used six

sera from convalescent patients from the first COVID-19 wave in Spain and six sera

from individuals that had been administered two doses of the BioNTech-Pfizer

Comirnaty COVID-19 vaccine. The neutralization capacity of the sera (NT50) was

obtained as previously described on VeroE6-TMPRSS2 cells (Giménez et al. 2022). We

used a previously described flow cytometry assay based on the use of polyclonal sera to

examine surface expression (Grzelak et al. 2020). Briefly, HEK293T cells were

transfected with the different S mutants using the calcium chloride method. 24 h later,

cells were detached using PBS with 1 mM EDTA, washed, and incubated in ice with

different polyclonal sera (three from convalescent patients from the 1st COVID-19

wave in Spain and one from individuals that had been administered two doses of the

BioNTech-Pfizer Comirnaty COVID-19 vaccine) at a 1:300 dilution in PBS containing

0.5 % BSA and 2 mM EDTA for 30 min. Next, cells were washed three times with

PBS, stained with anti-IgG Alexa Fluor 647 (Thermo Fisher Scientific) at a 1:400

dilution, and analyzed by flow cytometry similarly treated un-transfected controls to set

the threshold for positive cells. For cell-cell fusion assays we used a split Venus

fluorescent protein system (García-Murria et al. 2019). HEK293T cells were grown

overnight in 24 well plates (1.5 x 105 cells/well) using DMEM supplemented with 10 %

FBS. After 24 hours, cells were transfected using Lipofectamine 2000 (Invitrogen) with

0.5 µg of either a 1:1 mixture of the S plasmids and a Jun-Nt Venus fragment (Addgene

22012) plasmid or a mixture of hACE2 plasmid (kindly provided by Dr. Markus

Hoffman; German Primate Center, Goettingen/Germany) (Hoffmann et al. 2020) and

the Fos-Ct Venus fragment (Addgene 22013). After 24 h, cells were counted, and the S-

transfected cells were mixed at a 1:1 ratio with ACE2-transfected cells and seeded in 96

well plates (3 x 104 cells/well) in 100 µL of media. Cells transfected with the Wuhan-

Hu-1 served as a positive control, while cells transfected with hACE2 and Jun-Nt Venus

were used as negative control. We obtained the GFP Integrated Intensity

(GCU·µm²/image) in each condition using a live-cell imaging platform (Incucyte SX5,

Sartorius) at 24 hours post mixing and standardized to the signal obtained from the

positive control (Wuhan-Hu-1 spike protein). All experiments were performed at least

three times in triplicates.

Statistical analyses were performed and visualized using R v4.1.2, along with

tidyverse v2.0.0 (Wickham et al. 2019) and *ggpubr* v0.4.0 (Kassambara 2023) to

facilitate the analysis and enhance the visualization of these results. Comparisons were

conducted utilizing t-tests. We used unpaired tests for all assays but neutralization (log-

transformed NT50 values) and surface expression, for which we used paired tests, as we

used the same sets of polyclonal sera in each experiment. To determine fold change

values, the ratio of group average values was calculated.

Results

Detection and transmission dynamics of spike NTD deletion repair in
the BA.1 lineage using GISAID genomes

We first examined the differences in success within the Omicron BA.1 lineage

depending on the presence of ΔH69/V70 and ΔV143/Y145 in the NTD of the spike

protein through a global survey of viral genome data. This domain is characterized by

recurrent deletions occurring in distinct independent lineages (McCallum, De Marco, et

al. 2021; McCallum, Walls, et al. 2021; McCarthy et al. 2021; Meng et al. 2021; Planas

et al. 2021; Tamura et al. 2023). We interrogated the possibility of different patterns of

deletion repair of epidemiological significance and the drivers behind their emergence

through a survey of 11.3 million samples from the GISAID's EpiCov platform (Figure

1A). We identified 5,499 samples that carried repaired S:ΔH69/V70, S:ΔV143/Y145, or

both (Figure 2A, Table 1; full details in Supplemental Table 2). The most frequently

observed group consisted of samples exhibiting the repaired S:ΔH69/V70 in BA.1

background, while the remaining repair patterns were less frequently observed.

321
Figure 1. Distribution of collection dates of the two survey datasets. Bars show the weekly number of records in
each survey dataset, excluding those with missing collection dates. Records identified as deletion repair-bearing were
extracted from the BA.1 data partition in both surveys, which is highlighted in a lighter shade. (A) GISAID dataset,
with lineages extracted from metadata. (B) ENA dataset (3-fold random subset), with lineages assigned running
pangolin v4.3 on consensus sequences reconstructed from read mappings (see Materials and Methods), as no lineage
metadata is provided in database records. Although lineage filtering prior to data processing was not possible for the
ENA dataset, the chosen time frame ensured adequate representation of BA.1 records.

329
Figure 2. Summary of GISAID survey results for BA.1 genomes with repaired S:ΔH69/V70, S:ΔV143/Y145 or
both. A) Sample collection timelines broken down by sampling location. The five countries with the highest number
of observations globally are displayed. B) Differences in the estimated transmission fitness between deletion repair
haplotypes. This metric corrects the size of each transmission cluster regarding local spatial and temporal sequencing
efforts and lineage abundances. The differences were assessed using Wilcoxon rank-sum tests, and P-values are
displayed above each line connecting a pair of haplotypes. Each data point corresponds to a transmission cluster. Group
sizes (number of transmission clusters, n) are displayed next to the p-values.

Table 1. Description of Omicron BA.1 haplotypes with repaired NTD deletions detected through the GISAID
survey. The ratio to BA.1 observations is calculated as the fraction of observations over the number of BA.1
observations found in the survey (n = 2,420,866).

Haplotype	Observations	Ratio to BA.1	Transmission clusters	Minimum independent emergences
BA.1 RepΔ69/70	4,643	$1.92 \cdot 10^{-3}$	159	3,578
BA.1 RepΔ143/145	557	$2.30 \cdot 10^{-4}$	29	499
BA.1 RepBoth	299	$1.24 \cdot 10^{-4}$	26	215

Nevertheless, estimates of prevalence can be distorted by several factors,

including the geographical location and temporal specificity associated with both

sequencing efforts and the distribution of lineages throughout the pandemic.

Transmission, on the other hand, is usually considered as a proxy of viral fitness,

because it is related to its basic reproductive number, reflecting the ability of the virus

to replicate, persist and spread within hosts and in the population (Domingo 2016).

Therefore, we identified the transmission clusters of each haplotype through a

phylogenetic estimation using worldwide SARS-CoV-2 genomes. We measured an

average within-cluster collection date window of 24 ± 27 days, with 70 % of clusters

showing only within-border transmission (Supplemental Figure 2). The haplotype

bearing the S:ΔH69/V70 single repair had the highest median transmission fitness,

followed by the double repair and, finally, by the S:ΔV143/Y145 single repair (Figure

2B). We then investigated host age as a potential adaptive determinant of NTD deletion

repair in BA.1 background. No significant differences in host age were observed

between deletion repair haplotypes (Supplemental Figure 3). However, transmitted

samples with repaired S:ΔH69/V70 were associated with higher host age compared to

non-transmitted samples (P = 4.7·10^{-3}; Supplemental Figure 4), an effect likely

independent of sampling location (Supplemental Figures 5 and 6). These findings

should be interpreted cautiously due to substantial missing patient metadata

(Supplemental Figure 7).

To evaluate the reliability of GISAID-derived data, we compared its consensus

sequences to corresponding raw reads from the European Nucleotide Archive (ENA).

We cross-referenced the available ENA entries with all GISAID identifiers of genomes

with deletion repairs to identify matching records (Supplemental Table 3). Among 18

genomes with matching ENA runs, less than 25 % confirmed fixed deletion repairs as

indicated by the GISAID consensus, and fewer than 60 % had significant read support

of the deletion repair genotype. Specifically, for repaired S:ΔH69/V70, seven genomes

matched ENA runs, with only one exhibiting a fixed repaired deletion. For repaired

S:ΔV143/Y145, one genome matched ENA runs, with none showing fixed repairs. For

genomes with both repaired deletions, ten matched ENA runs, with three displaying

fixed repairs. These findings suggest that GISAID sequences may not reliably represent

deletion repair genotypes, highlighting the need for further validation of such analyses.

Detection and transmission dynamics of spike NTD deletion repair in
the BA.1 lineage using raw sequencing data

In our survey of viral genome data from GISAID we were able to identify repairs to

deletion mutations, which are theoretically rare events. However, their validity may be

confounded by sequencing artifacts. Public consensus genome sequences often suffer

from low coverage, poor sequencing quality, and imputed data, hindering the detection

of genuine deletion repair events. To address potential artifacts, we analyzed 464,859

read run records from the ENA (Figure 1B), selected out of 1.54 million collected

between 1 November 2021 and 1 August 2022. This allowed us to reassess haplotype

assignments with greater reliability.

We observed substantial discrepancies compared to the initial GISAID genome

survey. We identified 24 samples that carried repaired S:ΔH69/V70, S:ΔV143/Y145 or

both (Figure 3A, Table 2; full details in Supplemental Table 4) with an average

haplotype-defining allele frequency of 0.918 (see Figure 3B). The ratio of repaired

deletions relative to BA.1 observations was around 3- to 51-fold lower (depending on

the haplotype) than in the GISAID dataset, underscoring a significant drop in detection.

This suggests that the earlier survey was influenced by artifacts or biases in sequence

processing.

Figure 3. Summary of ENA survey results for BA.1 records with repaired S:ΔH69/V70, S:ΔV143/Y145 or both.
A) Sample collection timelines broken down by sampling location. The five countries with at least one observation are displayed. Crossed markers indicate observations belonging to the single detected transmission cluster. The timeframe aligns with Figure 2A for consistency. B) Allele frequency of deletion repairs in each run record. For each allele, each point represents the allele frequency in a different record. Point bins have a width of 0.05. Note that, for haplotype RepΔV143/Y145, some samples bear repaired S:ΔH69/V70 with a frequency under the 0.25 threshold (see Materials and Methods). No differences in the estimated transmission fitness between deletion repair haplotypes are reported, as we did not detect transmission for all haplotypes (see Table 2).

Table 2. Description of Omicron BA.1 haplotypes with repaired NTD deletions detected through the ENA survey. The ratio to BA.1 observations is calculated as the fraction of observations over the number of BA.1 read runs found in the survey analysis (n = 186,387). The average and standard deviation (SD) of the frequency of the deletion repair allele(s) associated with each haplotype are reported as well.

Haplotype	Observations	Ratio to BA.1	Average repair allele frequency	Transmission clusters	Minimum independent emergences
BA.1 RepΔ69/70	7	$3.76 \cdot 10^{-5}$	0.925 ± 0.073	1	6
BA.1 RepΔ143/145	15	$8.05 \cdot 10^{-5}$	0.884 ± 0.064	0	15
BA.1 RepBoth	2	$1.07 \cdot 10^{-5}$	0.945 ± 0.021	0	2

We then performed a phylogenetic estimation of transmission clusters using worldwide SARS-CoV-2 genomes. A single transmission cluster was detected (see Table 2), involving two records with repaired S:ΔH69/V70 (accession numbers SRR19566289 and SRR20496144) collected 20 days apart in California and New Jersey, U.S. The allele frequency of the deletion repair declined 12 % between infections. Notably, the assigned haplotype exhibited the highest estimated transmission fitness in the earlier GISAID survey (see Figure 2B). In short, generally, each observation was detected as an independent emergence with limited epidemiological impact, suggesting that these NTD repaired deletion events might be rare, isolated, and unlikely to drive SARS-CoV-2 evolution or transmission dynamics by themselves.

NTD deletion repair in Omicron BA.1 background has a non-accumulative effect on viral phenotype

NTD deletion repair events could have still contributed to viral diversity in a broader evolutionary context. Even isolated events might provide new opportunities for fitness-enhancing mutations to emerge, particularly in high-transmission scenarios. To investigate whether certain viral characteristics act as drivers of certain patterns of deletion repair compared to others, we analysed the effect of these events in the Omicron BA.1 spike using a pseudotyped VSV system. We generated VSV pseudotyped with the BA.1 spike proteins with repaired S:ΔH69/V70, repaired S:ΔV143/Y145, or both deletions repaired. We then used these pseudotyped viruses to examine possible phenotypic effects on spike function, including the efficiency of virus production, thermal stability, surface expression, susceptibility to antibody neutralization, and fusogenicity.

To assess possible effects on virus production, which encompasses both viral egress and infection of the next cell, VSV was pseudotyped with all spike constructs under identical conditions at the same time, and the amount of virus produced titrated on Vero E6 cells (Figure 4A). Virus production was unaffected by S:ΔV143/Y145 repair (0.77-fold change; $P = 0.06$) but was significantly reduced upon repairing S:ΔH69/V70 (0.28-fold; $P = 6.6 \cdot 10^{-4}$) or both deletions (0.34-fold; $P = 1.6 \cdot 10^{-4}$). A similar effect was observed when the titre of the virus was evaluated in cells expressing the TMPRSS2 co-receptor, indicating a substantial negative effect of S:ΔH69/V70 repair by itself on virus production (0.26-fold; $P = 0.017$).Reduced viral production can stem from lower stability of the Spike proteins or lower expression on the cell surface. To test the former, we obtained the temperature resulting in a 50 % reduction of virus

titre for each spike variant (i.e., 50 % inactivation temperature; Figure 4B). No

significant differences were observed, suggesting that spike protein stability was not the

driver of these differences. We therefore examined whether the different constructs had

altered cell expression of the different constructs by flow cytometry, using polyclonal

sera from four individuals (Figure 4C). We did not detect differences in the median cell

surface expression of the spike protein between the canonical BA.1 protein and any

deletion repair haplotype. However, the repair of S:ΔV143/Y145 resulted in higher

expression than S:ΔH69/V70 (3.2-fold; P = 0.0030) and the double repair (1.9-fold; P =

0.032), resembling the effect observed with virus production.

446
Figure 4. Experimental assessment of the effect of repairing NTD deletions in Omicron BA.1 background. (A)
Comparison of viral titres obtained for each spike variant using pseudotyped VSV on the indicated cell line. (B)
Comparison of the temperature resulting in 50 % inactivation of pseudotyped VSV carrying the indicated S protein.
(C) Surface expression of the spike protein in transfected HEK293T cells quantified using flow cytometry. (D)
Reciprocal 50 % neutralization titres (NT50) of sera from convalescent and vaccinated individuals. (E) Relative ability
of each spike variant to drive cell-cell fusion relative to the Wuhan-Hu-1 spike. (F) Summary of the log2-transformed
fold change for each spike variant relative to that of BA.1 in terms of average virus production in VERO E6 cells,
average fusogenicity, median NT50, average surface expression and average 50 % inactivating temperature. Subfigures
A-E represent the median and interquartile range of at least three replicates. Differences were assessed using Wilcoxon
rank-sum tests.

As neutralizing antibodies can be key drivers of viral evolution, we questioned

whether deletion repair affected neutralization by polyclonal sera from convalescent

donors from the first epidemic wave in Spain and those dually vaccinated with the

Comirnaty mRNA vaccine (n = 6 each; Figure 4D). Significant increases in

susceptibility to neutralization in sera from vaccinated individuals against viruses with

repaired S:ΔV143/Y145 (2.1-fold; P = 0.011) or both S:ΔH69/V70 and S:ΔV143/Y145

repaired (1.5-fold; P = 0.020) were observed. A similar trend was observed with

convalescent sera, but statistical significance was not reached (P = 0.19 and P = 0.42,

respectively) possibly due to a higher intra-sample variability. Thus, these results

indicate that increased neutralization is driven by the repair of S:ΔV143/Y145 in BA.1

background.

Cell to cell spread via fusion of the plasma membrane could potentially reduce

exposure to neutralizing antibodies, compensating for increased susceptibility to

neutralization. Hence, we also examined whether deletion repair could alter the ability

of the spike protein to fuse cells (Figure 4E). Interestingly, the repair of S:ΔV143/Y145

increased cell fusion relative to the BA.1 spike protein (1.5-fold; P = 0.026), while the

repair of both S:ΔH69/V70 and S:ΔV143/Y145 led to a decrease of more than 50 % in

the average fusogenicity (0.42-fold; P = 0.021). The presence of S:ΔH69/V70 by itself

did not seem to play a role in cell-cell fusion in a BA.1 background.

Discussion

Deletions in the SARS-CoV-2 genome have a significant impact on viral adaptation and

fitness, often surpassing the effects of single nucleotide variants (Cantoni et al. 2022;

McCarthy et al. 2021; Meng et al. 2021; Mishra et al. 2022; Venkatakrishnan et al.

2023). In fact, deletions in the NTD region of the spike protein are fixed in many

prominent viral variants. Our understanding of the repair patterns of deletions is crucial

for elucidating the genetic and phenotypic characteristics of the variants of concern. In

this work, we demonstrate that repairing these deletions can alter viral characteristics

and potentially influence the success, emergence and transmission dynamics of specific

viral haplotypes.

The *in vitro* assessment of the viral characteristics driving different patterns of

deletion repair in BA.1 showed differences in spike virus production and infectivity,

fusogenicity, and sera neutralization, but no differences in thermal stability. Repair of

S:ΔH69/V70 led to reduced virus titres individually and in combination with

S:ΔV143/Y145 repair. We did not observe differences in thermal stability with the

BA.1 protein, suggesting that the effect of S:ΔH69/V70 and S:ΔV143/Y145 repair did

not affect full-protein stability.

Cell-cell fusion and subsequent formation of syncytia can be mediated by viral

membrane glycoproteins (W. Li et al. 2003; Tseng et al. 2005) such as the spike protein

found in SARS-CoV-2, and it is a key feature of SARS-CoV-2 infection (Buchrieser et

al. 2020; Bussani et al. 2020; Hoffmann et al. 2020). In line with prior research, our

results point to a lower fusogenicity of the unaltered BA.1 spike protein compared to

that of Wuhan-Hu-1, which has been attributed to its inefficient spike cleavage and

unfavoured TMPRSS2-mediated cell entry (Meng et al. 2021). Interestingly, deletion

repair patterns exerted a non-accumulative influence on the fusogenicity of the BA.1

spike protein: repair of S:ΔH69/V70 alone did not have a significant impact and repair

of S:ΔV143/Y145 promoted higher fusogenicity, while repair of both deletions led to a

significant decrease in fusogenicity.

Deletion repair did also affect sera neutralization. In the BA.1 background,

repaired S:ΔV143/Y145 was associated with a significant increase in sensitivity to

neutralization by polyclonal sera obtained from vaccinated individuals. Conversely, we

did not detect a significant association of repaired S:ΔV143/Y145 with sensitivity to

sera from convalescent, "first-wave" patients. Repair of S:ΔH69/V70 did not affect

neutralization with either serum group. Although our surrogate system might not fully

reflect the complex interactions of the spike protein with host cells and the immune

system *in vivo*, and other regions of the spike protein or the viral genome might also

contribute to SARS-CoV-2 fitness and adaptation, our results point to the variant-

independent influence of NTD variability on immune effects. These findings point to

deletion repair events being rare and isolated, as their context-dependent and often

deleterious effects generally suggest limited adaptive advantage. This may also indicate

epistatic incompatibilities that constrain their contribution to viral fitness and

transmission. In agreement with the *in vitro* results, sequence surveys suggest that

deletion repair events were rare and likely isolated, with limited direct impact on SARS-

CoV-2 evolution or transmission dynamics. This aligns with the hypothesis that

recurrent deletion and apparent repair events in early variants of concern (VOC) may

have been driven by persistent viral reservoirs, where prolonged infections or animal

reservoirs allow for cycles of mutation accumulation and reversion before

reintroduction. In contrast, such dynamics appear less relevant in second-generation

dominant lineages (e.g., BA.2 sublineages), as reviewed by Roemer et al. (Roemer et al.

2023).

All survey efforts are expected to be biased by geographic and temporal

differences in sequencing efforts and lineage prevalence. Sequence analysis could be

affected by sampling bias and technical artifacts. However, despite our attempts to

account for demographic differences between variants, our analysis could be affected by

biases in genome sequencing and publication practices.

In our study, we initially observed instances of apparent genetic reversions that

could, in fact, be attributable to missing information in poorly sequenced regions or

polymorphic sites. The ENA survey revealed a significantly lower frequency of deletion

repair events compared to GISAID, and the majority of one-to-one comparisons of

GISAID genomes against raw reads data did not confirm the deletion repair genotype.

This suggests potential artifacts in GISAID sequences, likely arising from a replacement

of problematic genomic regions with the homologous ancestral sequence. Such

undocumented adjustments can create reference bias, distorting the interpretation of

genetic events and complicating genomic analyses, particularly given the quality control

mechanisms employed by repositories like GISAID. In fact, key global resources such

as the daily-updated mutation-annotated trees we used for transmission estimation

(McBroome et al. 2021) could well be impacted by these issues. We highlight the need

for greater scrutiny of sequence deposition and quality control mechanisms. Ongoing

efforts such as the Viridian initiative (Hunt et al. 2024) have already pointed out these

issues and are working toward reducing their impact. Incidentally, the limitations of the

FASTA format itself also contribute to this problem, since it restricts the amount of

metadata and quality information that can directly accompany sequences. This choice,

while historically convenient, may no longer align with the need for richer data

representations. Accordingly, submission of raw sequencing data from genome studies

to public repositories should be prioritized.

We also acknowledge that our study does not specifically address the mutational

mechanism that led to deletion repair. We speculate that recombination between

different infecting variants could be behind the emergence deletion-repairing

haplotypes, at least in some cases. We further speculate that template switching at

wrong loci during replication ("imperfect homologous recombination", as hypothesized

by Chrisman et al. 2021) could explain the emergence of rare indel events such as

deletion repairs in the absence of coinfections. Recombination might have played a role

in the evolution of SARS-CoV-2 from the ancestral coronavirus before diversification

in humans (Andersen et al. 2020; Boni et al. 2020; Kirtipal et al. 2020; MacLean et al.

2021; Wells et al. 2021), and studies using focused datasets have also pointed to

ongoing recombination and proposed insights into its dynamics (e.g., Ignatieva et al.

2022; Jackson et al. 2021; Perez-Florido et al. 2023; VanInsberghe et al. 2021). Still, the

emergence of the first demonstrably recombinant lineages of epidemiological relevance

was not reported until years into the pandemic (Tamura et al. 2023; Yue et al. 2023).

Due to the poor phylogenetic resolution of the datasets, our efforts ultimately led to an

inconclusive result, rendering us unable to articulate a systematic verification of the

recombination hypothesis.

In summary, our study describes the emergence of rare deletion repair events in

the SARS-CoV-2 spike NTD, and investigates their impact on viral fitness,

transmission, and clinical characteristics depending on the genetic context. In doing so,

we have provided novel insights into the repair patterns of NTD deletions in Omicron

BA.1 background and their phenotypic consequences. While significant progress has

been recently made in estimating the fitness effects of individual mutations across the

SARS-CoV-2 genome (Bloom & Neher 2023), these approaches do not focus on how

mutations interact within different genomic contexts. There have also been efforts to

specifically characterize the variability of the S protein NTD through full swap assays,

comparing the ancestral (Wuhan-Hu-1) spike protein with that of Alpha and Omicron

BA.1 (Cantoni et al. 2022), Delta and Omicron BA.1 and BA.2 (Meng et al. 2021) and

even with more distant virus like SARS-CoV (Qing et al. 2021). However, to our best

knowledge, no previous study has undertaken a combinatorial approach to characterize

each separate marker. Our work reveals that repair of specific deletions might be driven

by distinct phenotypic traits, such as changes in viral characteristics, including

fusogenicity, infectivity and susceptibility to neutralization. Understanding these

genotype-phenotype relationships can inform the evolutionary dynamics and adaptation

of SARS-CoV-2 variants. Future studies building upon these findings will contribute to

the development of effective strategies to monitor and mitigate the impact of NTD

deletions in emerging SARS-CoV-2 variants.

Data availability

The GISAID EPI_SET underlying this article is available via

https://dx.doi.org/10.55876/gis8.230801ex. Raw reads data is publicly available via the

European Nucleotide Archive (ENA) under the accession numbers provided in

Supplemental Table 1. The software we developed and used to find clusters in a global

phylogeny, *tcfinder* v1.0.0, is a command line tool released as free software under the

GNU GPLv3 license. Its source code is available in GitHub

(https://github.com/PathoGenOmics-Lab/tcfinder). It is also available as a package in

the Bioconda *conda* channel (https://anaconda.org/bioconda/tcfinder). The pipeline used

for the transmission analysis, *transcluster* v2.1.0, is a Snakemake workflow also

released as free software under the GNU GPLv3 license, and its source code is available

in GitHub (https://github.com/PathoGenOmics-Lab/transcluster). The processed data,

code, configuration files and results of the ENA survey and both transmission analyses

are available via https://dx.doi.org/10.20350/digitalCSIC/17032. The lists of sample

identifiers assigned to each haplotype under study are also incorporated into the article

and its online supplemental material. Code for performing the ENA survey and the

pipeline for processing the resulting raw sequencing data is available as free software

under the GNU GPLv3 license, and its source code is available in GitHub

(https://github.com/PathoGenOmics-Lab/ena-spike-ntd-repdel-analysis; v1.0.0).

Acknowledgements

We gratefully acknowledge all data contributors, i.e., the Authors and their Originating

laboratories responsible for obtaining the specimens, and their Submitting laboratories

for generating the genetic sequence and metadata and sharing via the GISAID Initiative,

on which this research is based. We also extend our acknowledgement to all

contributors and submitters of raw reads hosted by the European Nucleotide Archive

(ENA) and other members of the International Nucleotide Sequence Database

Collaboration (INSDC), which are also integral to this study. We thank Dr. Óscar

González Recio (INIA-CSIC, Spain) for providing sequencing data from a BA.1 sample

with S:142G and repaired NTD deletions (GISAID accession code: EPI_ISL_9805648)

for an initial exploratory analysis related to S gene target failure (SGTF), Dr. Luis

Enjuanes (CNB-CSIC, Spain) for providing Vero E6 cells, and Dr. Markus Hoffman

(German Primate Center, Germany) for providing the hACE2 plasmid. We thank Dr.

Irving Cancino Muñoz for his insightful feedback and recommendations on sequence

data processing. We are grateful to Jimena Solana González for reviewing the

manuscript and offering thoughtful feedback. We also thank Francisco José Martínez

Martínez for comments on phylogeny visualization and analysis of transmission

clusters. The computations were performed on the HPC cluster Garnatxa at the Institute

for Integrative Systems Biology (I^2SysBio, Spain). The I^2SysBio is a joint collaborative

research institute involving the University of Valencia (UV) and the Spanish National

Research Council (CSIC).

Funding details

This research work was funded by the European Commission – NextGenerationEU/

PRTR (Regulation EU 2020/2094), through CSIC's Global Health Platform (PTI+ Salud

Global) to MC, RG, IC and FGC. MAH is supported by the Generalitat Valenciana and

the European Social Fund "ESF Investing in your future" through grant

CIACIF/2022/333. This work was also a part of projects CNS2022-135116 (MC) and

CNS2022-135100 (RG) funded by MCIN/AEI/10.13039/501100011033 and the

European Union NextGenerationEU/PRTR.

Ethics statement

Sera samples for the biological characterization of BA.1 deletion repair were obtained

from the La Fe University and Polytechnic Hospital of Valencia and were collected

after informed written consent had been obtained, with approval by the ethical

committee and institutional review board (registration number 2020-123-1).

Contributions

MAH, BND, PRR and MC conceived the theoretical framework. MAH and BND

devised the initial idea. MAH and MC conceived the final study design. MAH

implemented and performed the data retrieval, data curation and computations. JZ, BG,

and RG designed the experiments. JZ and BG performed the experiments. MAH carried

646   out the statistical analyses of the experiments. MG and CAG contributed biological

647   samples. MAB did project management. MAH drafted the manuscript with support

648   from BDN, PRR, RG and MC. FGC aided in interpreting the results. MAH, PRR, BND,

649   RG, FGC, IC and MC discussed the results and commented on the manuscript. MC

650   supervised the project.

Declaration of interest statement

652   The authors report there are no competing interests to declare.

References

Aksamentov, I., Roemer, C., Hodcroft, E. B., & Neher, R. A. (2021). 'Nextclade: clade
assignment, mutation calling and quality control for viral genomes', *Journal of
Open Source Software*, 6/67: 3773. DOI: 10.21105/joss.03773
Andersen, K. G., Rambaut, A., Lipkin, W. I., Holmes, E. C., & Garry, R. F. (2020).
'The proximal origin of SARS-CoV-2', *Nature Medicine*, 1–3. DOI:
10.1038/s41591-020-0820-9
Avanzato, V. A., Matson, M. J., Seifert, S. N., Pryce, R., Williamson, B. N., Anzick, S.
L., Barbian, K., et al. (2020). 'Case Study: Prolonged Infectious SARS-CoV-2
Shedding from an Asymptomatic Immunocompromised Individual with
Cancer', *Cell*, 183/7: 1901-1912.e9. DOI: 10.1016/j.cell.2020.10.049
Bloom, J. D., & Neher, R. A. (2023). 'Fitness effects of mutations to SARS-CoV-2
proteins', *Virus Evolution*, 9/2: vead055. DOI: 10.1093/ve/vead055
Boni, M. F., Lemey, P., Jiang, X., Lam, T. T.-Y., Perry, B. W., Castoe, T. A., Rambaut,
A., et al. (2020). 'Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage
responsible for the COVID-19 pandemic', *Nature Microbiology*, 5/11: 1408–17.
Nature Publishing Group. DOI: 10.1038/s41564-020-0771-4
Buchrieser, J., Dufloo, J., Hubert, M., Monel, B., Planas, D., Rajah, M. M., Planchais,
C., et al. (2020). 'Syncytia formation by SARS-CoV-2-infected cells', *The
EMBO journal*, 39/23: e106267. DOI: 10.15252/embj.2020106267
Bussani, R., Schneider, E., Zentilin, L., Collesi, C., Ali, H., Braga, L., Volpe, M. C., et
al. (2020). 'Persistence of viral RNA, pneumocyte syncytia and thrombosis are
hallmarks of advanced COVID-19 pathology', *EBioMedicine*, 61: 103104. DOI:
10.1016/j.ebiom.2020.103104
Cantoni, D., Murray, M. J., Kalemera, M. D., Dicken, S. J., Stejskal, L., Brown, G.,
Lytras, S., et al. (2022). 'Evolutionary remodelling of N-terminal domain loops
fine-tunes SARS-CoV-2 spike', *EMBO reports*, 23/10: e54322. John Wiley &
Sons, Ltd. DOI: 10.15252/embr.202154322
Chen, S. (2023). 'Ultrafast one-pass FASTQ data preprocessing, quality control, and
deduplication using fastp', *iMeta*, 2/2: e107. DOI: 10.1002/imt2.107
Chen, Z., Chong, K. C., Wong, M. C. S., Boon, S. S., Huang, J., Wang, M. H., Ng, R.

684   W. Y., et al. (2021). 'A global analysis of replacement of genetic variants of
685    SARS-CoV-2 in association with containment capacity and changes in disease
686    severity', *Clinical Microbiology and Infection*, 27/5: 750–7. DOI:
687    10.1016/j.cmi.2021.01.018
688 Choi, B., Choudhary, M. C., Regan, J., Sparks, J. A., Padera, R. F., Qiu, X., Solomon, I.
689    H., et al. (2020). 'Persistence and Evolution of SARS-CoV-2 in an
690    Immunocompromised Host', *New England Journal of Medicine*, 383/23: 2291–
691    3. Massachusetts Medical Society. DOI: 10.1056/NEJMc2031364
692 Chrisman, B. S., Paskov, K., Stockham, Nate., Tabatabaei, K., Jung, J.-Y., Washington,
693    P., Varma, M., et al. (2021). 'Indels in SARS-CoV-2 occur at template-
694    switching hotspots', *BioData Mining*, 14: 20. DOI: 10.1186/s13040-021-00251-
695    0
696 Cingolani, P., Patel, V. M., Coon, M., Nguyen, T., Land, S. J., Ruden, D. M., & Lu, X.
697    (2012). 'Using Drosophila melanogaster as a Model for Genotoxic Chemical
698    Mutational Studies with a New Program, SnpSift', *Frontiers in Genetics*, 3: 35.
699    DOI: 10.3389/fgene.2012.00035
700 Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,
701    Whitwham, A., et al. (2021). 'Twelve years of SAMtools and BCFtools',
702    *GigaScience*, 10/2: giab008. DOI: 10.1093/gigascience/giab008
703 Denison, M. R., Graham, R. L., Donaldson, E. F., Eckerle, L. D., & Baric, R. S. (2011).
704    'Coronaviruses: an RNA proofreading machine regulates replication fidelity and
705    diversity', *RNA biology*, 8/2: 270–9. DOI: 10.4161/rna.8.2.15013
706 Domingo, E. (2016). 'Chapter 7 - Long-Term Virus Evolution in Nature'. Domingo E.
707    (ed.) *Virus as Populations*, pp. 227–62. Academic Press: Boston. DOI:
708    10.1016/B978-0-12-800837-9.00007-1
709 García-Murria, M. J., Expósito-Domínguez, N., Duart, G., Mingarro, I., & Martinez-
710    Gil, L. (2019). 'A Bimolecular Multicellular Complementation System for the
711    Detection of Syncytium Formation: A New Methodology for the Identification
712    of Nipah Virus Entry Inhibitors', *Viruses*, 11/3: 229. Multidisciplinary Digital
713    Publishing Institute. DOI: 10.3390/v11030229
714 Giménez, E., Albert, E., Zulaica, J., Torres, I., Rusu, L., Moreno, A. R., Burgos, J. S., et
715    al. (2022). 'Severe Acute Respiratory Syndrome Coronavirus 2 Adaptive
716    Immunity in Nursing Home Residents Following a Third Dose of the Comirnaty
717    Coronavirus Disease 2019 Vaccine', *Clinical Infectious Diseases*, 75/1: e865–8.
718    DOI: 10.1093/cid/ciac223
719 Gozalbo-Rovira, R., Gimenez, E., Latorre, V., Francés-Gómez, C., Albert, E., Buesa, J.,
720    Marina, A., et al. (2020). 'SARS-CoV-2 antibodies, serum inflammatory
721    biomarkers and clinical severity of hospitalized COVID-19 patients', *Journal of
722    Clinical Virology*, 131: 104611. DOI: 10.1016/j.jcv.2020.104611
723 Grubaugh, N. D., Gangavarapu, K., Quick, J., Matteson, N. L., De Jesus, J. G., Main, B.
724    J., Tan, A. L., et al. (2019). 'An amplicon-based sequencing framework for
725    accurately measuring intrahost virus diversity using PrimalSeq and iVar',
726    *Genome Biology*, 20/1: 8. DOI: 10.1186/s13059-018-1618-7
727 Grüning, B., Dale, R., Sjödin, A., Chapman, B. A., Rowe, J., Tomkins-Tinch, C. H.,
728    Valieris, R., et al. (2018). 'Bioconda: sustainable and comprehensive software
729    distribution for the life sciences', *Nature Methods*, 15/7: 475–6. Nature
730    Publishing Group. DOI: 10.1038/s41592-018-0046-7
731 Grzelak, L., Temmam, S., Planchais, C., Demeret, C., Tondeur, L., Huon, C., Guivel-

Benhassine, F., et al. (2020). 'A comparison of four serological assays for
detecting anti–SARS-CoV-2 antibodies in human serum samples from different
populations', *Science Translational Medicine*, 12/559: eabc3103. American
Association for the Advancement of Science. DOI:
10.1126/scitranslmed.abc3103

Hadfield, J., Megill, C., Bell, S. M., Huddleston, J., Potter, B., Callender, C.,
Sagulenko, P., et al. (2018). 'Nextstrain: real-time tracking of pathogen
evolution', *Bioinformatics*, 34/23: 4121–3. DOI: 10.1093/bioinformatics/bty407

Harvey, W. T., Carabelli, A. M., Jackson, B., Gupta, R. K., Thomson, E. C., Harrison,
E. M., Ludden, C., et al. (2021). 'SARS-CoV-2 variants, spike mutations and
immune escape', *Nature Reviews Microbiology*, 19/7: 409–24. DOI:
10.1038/s41579-021-00573-0

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S.,
Schiergens, T. S., et al. (2020). 'SARS-CoV-2 Cell Entry Depends on ACE2 and
TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor', *Cell*,
181/2: 271-280.e8. DOI: 10.1016/j.cell.2020.02.052

Hunt, M., Hinrichs, A. S., Anderson, D., Karim, L., Dearlove, B. L., Knaggs, J.,
Constantinides, B., et al. (2024). 'Addressing pandemic-wide systematic errors
in the SARS-CoV-2 phylogeny'. bioRxiv. DOI: 10.1101/2024.04.29.591666

Ignatieva, A., Hein, J., & Jenkins, P. A. (2022). 'Ongoing Recombination in SARS-
CoV-2 Revealed through Genealogical Reconstruction', *Molecular Biology and
Evolution*, 39/2: msac028. DOI: 10.1093/molbev/msac028

Jackson, B., Boni, M. F., Bull, M. J., Colleran, A., Colquhoun, R. M., Darby, A. C.,
Haldenby, S., et al. (2021). 'Generation and transmission of interlineage
recombinants in the SARS-CoV-2 pandemic', *Cell*, 184/20: 5179-5188.e8. DOI:
10.1016/j.cell.2021.08.014

Kassambara, A. (2023). 'ggpubr: "ggplot2" Based Publication Ready Plots'. Retrieved
from CRAN.

Kemp, S. A., Collier, D. A., Datir, R. P., Ferreira, I. A. T. M., Gayed, S., Jahun, A.,
Hosmillo, M., et al. (2021). 'SARS-CoV-2 evolution during treatment of chronic
infection', *Nature*, 592/7853: 277–82. Nature Publishing Group. DOI:
10.1038/s41586-021-03291-y

Khare, S., Gurry, C., Freitas, L., Schultz, M. B., Bach, G., Diallo, A., Akite, N., et al.
(2021). 'GISAID's Role in Pandemic Response', *China CDC Weekly*, 3/49:
1049–51. China CDC Weekly. DOI: 10.46234/ccdcw2021.255

Kirtipal, N., Bharadwaj, S., & Kang, S. G. (2020). 'From SARS to SARS-CoV-2,
insights on structure, pathogenicity and immunity aspects of pandemic human
coronaviruses', *Infection, Genetics and Evolution: Journal of Molecular
Epidemiology and Evolutionary Genetics in Infectious Diseases*, 85: 104502.
DOI: 10.1016/j.meegid.2020.104502

Köster, J., & Rahmann, S. (2012). 'Snakemake—a scalable bioinformatics workflow
engine', *Bioinformatics*, 28/19: 2520–2. DOI: 10.1093/bioinformatics/bts480

Letko, M., Marzi, A., & Munster, V. (2020). 'Functional assessment of cell entry and
receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses', *Nature
Microbiology*, 5/4: 562–9. DOI: 10.1038/s41564-020-0688-y

Li, H. (2018). 'Minimap2: pairwise alignment for nucleotide sequences', (I. Birol,
Ed.)*Bioinformatics*, 34/18: 3094–100. DOI: 10.1093/bioinformatics/bty191

——. (2021). 'New strategies to improve minimap2 alignment accuracy', (C. Alkan,

780   Ed.)*Bioinformatics*, 37/23: 4572–4. DOI: 10.1093/bioinformatics/btab705

781 Li, W., Moore, M. J., Vasilieva, N., Sui, J., Wong, S. K., Berne, M. A., Somasundaran,
782   M., et al. (2003). 'Angiotensin-converting enzyme 2 is a functional receptor for
783   the SARS coronavirus', *Nature*, 426/6965: 450–4. Nature Publishing Group.
784   DOI: 10.1038/nature02145

785 MacLean, O. A., Lytras, S., Weaver, S., Singer, J. B., Boni, M. F., Lemey, P.,
786   Kosakovsky Pond, S. L., et al. (2021). 'Natural selection in the evolution of
787   SARS-CoV-2 in bats created a generalist virus and highly capable human
788   pathogen', *PLoS biology*, 19/3: e3001115. DOI: 10.1371/journal.pbio.3001115

789 McBroome, J., Thornlow, B., Hinrichs, A. S., Kramer, A., De Maio, N., Goldman, N.,
790   Haussler, D., et al. (2021). 'A Daily-Updated Database and Tools for
791   Comprehensive SARS-CoV-2 Mutation-Annotated Trees', *Molecular Biology*
792   *and Evolution*, 38/12: 5819–24. DOI: 10.1093/molbev/msab264

793 McCallum, M., De Marco, A., Lempp, F. A., Tortorici, M. A., Pinto, D., Walls, A. C.,
794   Beltramello, M., et al. (2021). 'N-terminal domain antigenic mapping reveals a
795   site of vulnerability for SARS-CoV-2', *Cell*, 184/9: 2332-2347.e16. DOI:
796   10.1016/j.cell.2021.03.028

797 McCallum, M., Walls, A. C., Sprouse, K. R., Bowen, J. E., Rosen, L. E., Dang, H. V.,
798   De Marco, A., et al. (2021). 'Molecular basis of immune evasion by the Delta
799   and Kappa SARS-CoV-2 variants', *Science (New York, N.Y.)*, 374/6575: 1621–
800   6. DOI: 10.1126/science.abl8506

801 McCarthy, K. R., Rennick, L. J., Nambulli, S., Robinson-McCarthy, L. R., Bain, W. G.,
802   Haidar, G., & Duprex, W. P. (2021). 'Recurrent deletions in the SARS-CoV-2
803   spike glycoprotein drive antibody escape', *Science (New York, N.Y.)*, 371/6534:
804   1139–42. DOI: 10.1126/science.abf6950

805 McMillen, T., Jani, K., Robilotti, E. V., Kamboj, M., & Babady, N. E. (2022). 'The
806   spike gene target failure (SGTF) genomic signature is highly accurate for the
807   identification of Alpha and Omicron SARS-CoV-2 variants', *Scientific Reports*,
808   12/1: 18968. Nature Publishing Group. DOI: 10.1038/s41598-022-21564-y

809 Meng, B., Kemp, S. A., Papa, G., Datir, R., Ferreira, I. A. T. M., Marelli, S., Harvey,
810   W. T., et al. (2021). 'Recurrent emergence of SARS-CoV-2 spike deletion
811   H69/V70 and its role in the Alpha variant B.1.1.7', *Cell Reports*, 35/13: 109292.
812   DOI: 10.1016/j.celrep.2021.109292

813 Minskaia, E., Hertzig, T., Gorbalenya, A. E., Campanacci, V., Cambillau, C., Canard,
814   B., & Ziebuhr, J. (2006). 'Discovery of an RNA virus 3'->5' exoribonuclease
815   that is critically involved in coronavirus RNA synthesis', *Proceedings of the*
816   *National Academy of Sciences of the United States of America*, 103/13: 5108–
817   13. DOI: 10.1073/pnas.0508200103

818 Mishra, T., Dalavi, R., Joshi, G., Kumar, A., Pandey, P., Shukla, S., Mishra, R. K., et al.
819   (2022). 'SARS-CoV-2 spike E156G/Δ157-158 mutations contribute to increased
820   infectivity and immune escape', *Life Science Alliance*, 5/7: e202201415. DOI:
821   10.26508/lsa.202201415

822 Morgan, M., Pagès, H., Obenchain, V., & Hayden, N. (2023). 'Rsamtools'.
823   Bioconductor. DOI: 10.18129/B9.BIOC.RSAMTOOLS

824 Mykytyn, A. Z., Rosu, M. E., Kok, A., Rissmann, M., Amerongen, G. van,
825   Geurtsvankessel, C., Vries, R. D. de, et al. (2023). 'Antigenic mapping of
826   emerging SARS-CoV-2 omicron variants BM.1.1.1, BQ.1.1, and XBB.1', *The*
827   *Lancet Microbe*, 4/5: e294–5. Elsevier. DOI: 10.1016/S2666-5247(22)00384-6

O'Toole, Á., Scher, E., Underwood, A., Jackson, B., Hill, V., McCrone, J. T.,
Colquhoun, R., et al. (2021). 'Assignment of epidemiological lineages in an
emerging pandemic using the pangolin tool', *Virus Evolution*, 7/2: veab064.
DOI: 10.1093/ve/veab064

Paradis, E., & Schliep, K. (2019). 'ape 5.0: an environment for modern phylogenetics
and evolutionary analyses in R', *Bioinformatics*, 35/3: 526–8. DOI:
10.1093/bioinformatics/bty633

Perez-Florido, J., Casimiro-Soriguer, C. S., Ortuño, F., Fernandez-Rueda, J. L., Aguado,
A., Lara, M., Riazzo, C., et al. (2023). 'Detection of High Level of Co-Infection
and the Emergence of Novel SARS CoV-2 Delta-Omicron and Omicron-
Omicron Recombinants in the Epidemiological Surveillance of Andalusia',
International Journal of Molecular Sciences, 24/3: 2419. DOI:
10.3390/ijms24032419

Planas, D., Veyer, D., Baidaliuk, A., Staropoli, I., Guivel-Benhassine, F., Rajah, M. M.,
Planchais, C., et al. (2021). 'Reduced sensitivity of SARS-CoV-2 variant Delta
to antibody neutralization', *Nature*, 596/7871: 276–80. Nature Publishing
Group. DOI: 10.1038/s41586-021-03777-9

Puhach, O., Adea, K., Hulo, N., Sattonnet, P., Genecand, C., Iten, A., Jacquérioz, F., et
al. (2022). 'Infectious viral load in unvaccinated and vaccinated individuals
infected with ancestral, Delta or Omicron SARS-CoV-2', *Nature Medicine*,
28/7: 1491–500. Nature Publishing Group. DOI: 10.1038/s41591-022-01816-0

Qing, E., Kicmal, T., Kumar, B., Hawkins, G. M., Timm, E., Perlman, S., & Gallagher,
T. (2021). 'Dynamics of SARS-CoV-2 Spike Proteins in Cell Entry: Control
Elements in the Amino-Terminal Domains', *mBio*, 12/4. American Society for
Microbiology. DOI: 10.1128/mbio.01590-21

Rambaut, A., Holmes, E. C., O'Toole, Á., Hill, V., McCrone, J. T., Ruis, C., du Plessis,
L., et al. (2020). 'A dynamic nomenclature proposal for SARS-CoV-2 lineages
to assist genomic epidemiology', *Nature Microbiology*, 5/11: 1403–7. DOI:
10.1038/s41564-020-0770-5

Roemer, C., Sheward, D. J., Hisner, R., Gueli, F., Sakaguchi, H., Frohberg, N.,
Schoenmakers, J., et al. (2023). 'SARS-CoV-2 evolution in the Omicron era',
Nature Microbiology, 8/11: 1952–9. Nature Publishing Group. DOI:
10.1038/s41564-023-01504-w

da Silva Francisco Junior, R., Lamarca, A. P., de Almeida, L. G. P., Cavalcante, L.,
Machado, D. T., Martins, Y., Brustolini, O., et al. (2021). 'Turnover of SARS-
CoV-2 Lineages Shaped the Pandemic and Enabled the Emergence of New
Variants in the State of Rio de Janeiro, Brazil', *Viruses*, 13/10: 2013.
Multidisciplinary Digital Publishing Institute. DOI: 10.3390/v13102013

Tamura, T., Ito, J., Uriu, K., Zahradnik, J., Kida, I., Anraku, Y., Nasser, H., et al.
(2023). 'Virological characteristics of the SARS-CoV-2 XBB variant derived
from recombination of two Omicron subvariants', *Nature Communications*,
14/1: 2800. Nature Publishing Group. DOI: 10.1038/s41467-023-38435-3

Tseng, C.-T. K., Tseng, J., Perrone, L., Worthy, M., Popov, V., & Peters, C. J. (2005).
'Apical Entry and Release of Severe Acute Respiratory Syndrome-Associated
Coronavirus in Polarized Calu-3 Lung Epithelial Cells', *Journal of Virology*,
79/15: 9470–9. American Society for Microbiology. DOI:
10.1128/jvi.79.15.9470-9479.2005

Turakhia, Y., Thornlow, B., Hinrichs, A. S., De Maio, N., Gozashti, L., Lanfear, R.,

Haussler, D., et al. (2021). 'Ultrafast Sample placement on Existing tRees
(UShER) enables real-time phylogenetics for the SARS-CoV-2 pandemic',
Nature Genetics, 53/6: 809–16. Nature Publishing Group. DOI:
10.1038/s41588-021-00862-7
Uriu, K., Ito, J., Zahradnik, J., Fujita, S., Kosugi, Y., Schreiber, G., & Sato, K. (2023).
'Enhanced transmissibility, infectivity, and immune resistance of the SARS-
CoV-2 omicron XBB.1.5 variant', *The Lancet Infectious Diseases*, 23/3: 280–1.
Elsevier. DOI: 10.1016/S1473-3099(23)00051-8
VanInsberghe, D., Neish, A. S., Lowen, A. C., & Koelle, K. (2021). 'Recombinant
SARS-CoV-2 genomes circulated at low levels over the first year of the
pandemic', *Virus Evolution*, 7/2: veab059. DOI: 10.1093/ve/veab059
Venkatakrishnan, A. J., Anand, P., Lenehan, P. J., Ghosh, P., Suratekar, R., Silvert, E.,
Pawlowski, C., et al. (2023). 'Expanding repertoire of SARS-CoV-2 deletion
mutations contributes to evolution of highly transmissible variants', *Scientific
Reports*, 13/1: 257. Nature Publishing Group. DOI: 10.1038/s41598-022-26646-
5
V'kovski, P., Kratzel, A., Steiner, S., Stalder, H., & Thiel, V. (2021). 'Coronavirus
biology and replication: implications for SARS-CoV-2', *Nature Reviews.
Microbiology*, 19/3: 155–70. DOI: 10.1038/s41579-020-00468-6
Ward, T., Glaser, A., Overton, C. E., Carpenter, B., Gent, N., & Seale, A. C. (2022).
'Replacement dynamics and the pathogenesis of the Alpha, Delta and Omicron
variants of SARS-CoV-2', *Epidemiology and Infection*, 151: e32. DOI:
10.1017/S0950268822001935
Wells, H. L., Letko, M., Lasso, G., Ssebide, B., Nziza, J., Byarugaba, D. K., Navarrete-
Macias, I., et al. (2021). 'The evolutionary history of ACE2 usage within the
coronavirus subgenus Sarbecovirus', *Virus Evolution*, 7/1: veab007. DOI:
10.1093/ve/veab007
Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R.,
Grolemund, G., et al. (2019). 'Welcome to the tidyverse', *Journal of Open
Source Software*, 4/43: 1686. DOI: 10.21105/joss.01686
Yue, C., Song, W., Wang, L., Jian, F., Chen, X., Gao, F., Shen, Z., et al. (2023). 'ACE2
binding and antibody evasion in enhanced transmissibility of XBB.1.5', *The
Lancet Infectious Diseases*, 23/3: 278–80. Elsevier. DOI: 10.1016/S1473-
3099(23)00010-5
Zhang, X., Chen, L.-L., Ip, J. D., Chan, W.-M., Hung, I. F.-N., Yuen, K.-Y., Li, X., et
al. (2022). 'Omicron sublineage recombinant XBB evades neutralising
antibodies in recipients of BNT162b2 or CoronaVac vaccines', *The Lancet
Microbe*, 4/3: e131. Elsevier. DOI: 10.1016/S2666-5247(22)00335-4