Larval zebrafish maintain elevation with multisensory control of posture and locomotion

Samantha N. Davis [ORCID],[1] **Yunlu Zhu** [ORCID],[1] **David Schoppik**[ORCID][1,2,*]
[1]Depts. of Otolaryngology, Neuroscience & Physiology, and the Neuroscience Institute, NYU Grossman School of Medicine
[2]Lead Contact
[*]Correspondence: schoppik@gmail.com

Fish actively control posture in the pitch axis (nose-up/nose-down) to counter instability and regulate their elevation in the
water column. To test the hypothesis that environmental cues shape strategies fish use to control posture, we leveraged
a serendipitous finding: larval zebrafish (*Danio rerio*) sink mildly after acute loss of lateral line hair cells. Using long-
term (48 h) recordings of unrestrained swimming, we discovered that sinking larvae compensated differently depending
on light conditions. In the dark, they swim more frequently with an increased nose-up posture. In contrast, larvae in the
light do not swim more frequently, but do climb more often. Finally, after lateral line regeneration, larvae returned to
normal buoyancy and swam comparably to control siblings. We conclude that larvae can switch postural control strategies
depending on the availability of visual information. Our findings complement and extend morphological and kinematic
analyses of locomotion. More broadly, by quantifying the variation in strategies our work speaks to the evolutionary
substrate for different balance behaviors.

INTRODUCTION

Actinopterygian (ray-finned) fishes adjust both posture and density to regulate their depth, allowing them to occupy a remarkable
range of elevations. Postural control is dynamic (Webb, 2002), and fishes will reorient relative to gravity and their environment
(Meyer et al., 1976a; 1977). While most fishes maintain a dorsal-up posture (Webb, 2006), notable exceptions speak to behavioral
flexibility. For example, *Syndodontus nigriventris* (Meyer et al., 1976b; Willoughby, 1976) swim ventral-up, *Aeoliscus punctulatus*
(Fish and Holzman, 2019) swim vertically with their heads pointed downward, and *Pleuronectiformes* swim on their side (Platt,
1973). Moreover, reorienting posture relative to gravity permits navigation in depth (Zhu et al., 2024), complementing morpholog-
ical specializations that decrease density (Alexander, 1990; Pelster, 1997). Key among these specializations is the swim bladder, a
gas-filled organ used by most teleosts, especially at larval stages, to countermand sinking by reducing density (Pelster, 2009; Steen,
1970). The remarkable diversity in behavior and habitat reflects selective pressures acting, at least in part, on postural control
strategies.

Understanding variability in postural behavior requires perturbations. For example, swim bladder inflation can be manipulated by
removing gas, attaching weights to fish, or changing external hydrostatic pressure (Fänge, 1966; Gee and Holst, 1992; Lindsey
et al., 2011; Steen, 1970). *Oncorhynchus nerka* adopt nose-up postures and increase pectoral fin movements as external hydrostatic
pressure increases (Harvey, 1963; Harvey and Bothern, 1972). Comparably, larval zebrafish (*Danio rerio*) change static posture and
swim frequency to compensate for oil-filled swim bladders (increased density) (Ehrlich and Schoppik, 2017a) or when swimming
in 1.5% glycerol (increased buoyancy) (Ehrlich and Schoppik, 2017b); neither manipulation is compatible with long-term imaging.
Taken together, the work establishes that fish change posture when challenged, but the strategies they use are underexplored.

Previous work establishes the larval zebrafish as an excellent model to investigate posture, and suggests a possible route to challenge
buoyancy. First, larvae swim in short translational bouts and, due to their low Reynolds number, do not glide (McHenry and Lauder,
2005). The passive period between bouts allows direct measurements of the effects of physical forces that challenge posture (e.g.
gravity) (Ehrlich and Schoppik, 2017a). Recent work measuring larval posture and locomotion from the side (Zhu et al., 2023)
reveals that larvae use a combination of trunk rotations and pectoral fin-generated lift (Ehrlich and Schoppik, 2019) to navigate in
depth (Zhu et al., 2024). Next, larvae inflate their swim bladders early in development to reduce density (Goolish and Okutake,
1999). Further, they sense flow using a set of superficial hair cell sensors known as the lateral line; these hair cells can be killed
pharmacologically and can regenerate (Davis et al., 2020). Early perturbations (pharmacological or genetic) to the lateral line
interfere with normal swim bladder inflation (Lin et al., 2023; Venuto et al., 2023), suggesting a means to vary buoyancy.

We hypothesized that, if their buoyancy changed, larval zebrafish would respond by changing posture and locomotion, and that the
nature of these changes would depend on the availability of visual information. We first observed that after loss of the lateral line,
larval zebrafish sank slightly between swim bouts; we could thus challenge fish in the light and the dark. We discovered that larvae
move more frequently in the dark, but climbed more often in the light; these changes could partially recover elevation. Finally,
behavior in larvae that had regenerated their lateral line was indistinguishable from age-matched control siblings. Our results reveal
that, when challenged, larval zebrafish adjust posture to maintain elevation with strategies that vary with the availability of visual
information.

MATERIALS AND METHODS

Animals

All procedures involving larval zebrafish (*Danio rerio*) were approved by the New York University Grossman School of Medicine Institutional Animal Care & Use Committee (IACUC). Zebrafish larvae were raised at 28.5°C on a standard light-dark cycle (14 hours light, 10 hours dark) at a density of 20–50 larvae in 25–40 mL of E3 medium before 6 days post-fertilization (dpf). During experiments, larvae older than 5 dpf were fed cultured rotifers (Reed Mariculture) daily for 30 minutes.

Zebrafish lines

Behavioral experiments were performed on the Schoppik lab's wild-type background (AB/WIK/TU mix), originally characterized in Zhu et al. (2023). Imaging experiments used the *Tg(myo6b:actb1-EGFP)* transgenic line to label lateral line hair cells.

Experiments

Behavior and Copper Treatment

Freely moving larvae were measured using the previously described Scalable Apparatus for Measuring Posture and Locomotion (SAMPL) (Zhu et al., 2023). In brief, 6–8 larvae were placed into custom vertical chambers (25 mL volume, 50 mm x 50 mm x 10 mm) filled with E3. Chambers were placed between an infrared light source (940 nm) and a camera fit with an infrared filter (Edmund Optics). Recordings were performed from the side in a light-tight enclosure with programmable white light illumination inside the box. The field of view (400 mm^2) was sampled from the lower middle of the chamber at a rate of 166 frames per second.

Experiments began at 6 dpf with 24 hours of recording. At 7 dpf, larvae were either treated with 10 µM copper sulfate (CuSO$_4$; Acros Organics 197722500) in E3 or untreated (transferred to E3 alone) for 90 minutes. Untreated fish were transferred and handled identically to copper-treated larvae. Larvae were then washed in E3 and returned to chambers for 24 hours of post-treatment recording. Copper sulfate treatment was repeated after 24 hours to avoid hair cell regeneration (Thomas et al., 2014). Larvae were fed for 30 minutes each day; behavior was not measured during feeding. A total of 5 experimental repeats using sibling controls (i.e., 5 clutches) were completed in the dark (N=114 total larvae per condition), and 8 in the light (N=150 total larvae per condition).

Extensive description of SAMPL analyses and a full Python-based analysis suite are freely available at Zhu et al. (2023). In short, swim bouts were extracted from captured epochs and aligned to peak speed. Bouts were defined as lasting 250 ms before to 200 ms after peak speed. Navigational categories were classified by bout trajectory at peak speed. Climb bouts had trajectories greater than 0.5°, dive bouts were less than -0.5°, and trajectories between -0.5° and 0.5° was categorized as "flat." The same criteria were applied to larvae in each lighting condition and treatment group. Only daytime activity (9:00–23:00) was analyzed to minimize the impact of circadian cues.

To evaluate larval ability to maintain elevation, chambers were filled with 6 mL of 2% agarose (Invitrogen 16500-500) so that the floor of the arena was in the field of view. Larvae were split into four groups: dark control (N=36), dark copper-treated (N=31), light control (N=57), and light copper (N=57) as described above. The number of larvae resting on agarose was counted every 30 minutes for 4 hours (8 observations total) across 5 experimental repeats in the dark and 6 repeats in the light.

Longitudinal Experiments

Effects of regenerated hair cells were examined over two periods of time: 7–9 dpf (early) and 14–16 dpf (late). Behavior recordings and copper treatments were conducted in the same manner as described above. In addition to an untreated sibling control group (N=85), one group of larvae was treated with copper sulfate in the early phase (N=94) and another group of larvae was treated during the late phase (N=64). Between behavioral recordings, larvae were raised in 450 mL of dense rotifer solution in 2 L tanks.

Imaging

One hour after copper treatment, larvae were washed three times for 5 minutes with E3. Larvae were next anesthetized in MS-222 (Sigma-Aldrich E10521) then mounted dorsally in 2% low-melting point agarose in E3 for imaging. The SO2 neuromast was identified on control and copper-treated larvae using a confocal microscope (Zeiss LSM800) equipped with a 20x water-immersion objective (1.0 NA). All images were analyzed using Fiji/ImageJ (Schindelin et al., 2012).

Statistical Analysis

Sample sizes were determined with respect to extensive prior characterization of wild-type variability (Zhu et al., 2023). An experimental repeat consists of all experiments run on a single clutch of fish (i.e. siblings). As most distributions of raw kinematic data were not normally distributed, central tendency and variance are reported as the median and interquartile range, and differences in medians between two groups were analyzed with Mann-Whitney U tests. Consequentially, identical p-values appear multiple times due to identical ranking across experimental repeats.

Fitting steering and righting gains requires a large number of bouts (Zhu et al., 2023); these were calculated on data aggregated across repeats, compared using unpaired t-tests. Effect size was estimated using Cohen's d (Lötsch and Ultsch, 2020). Finally, two-way ANOVA was used to evaluate differences across multiple conditions (light/dark, treated/untreated) and one-way ANOVA was used to assess longitudinal data, with Tukey's HSD post-hoc comparison to adjust for multiple comparisons. Statistical significance was set at $\alpha = 0.05$.

RESULTS

Larvae compensate for postural instability after lateral line ablation with increased movement frequency and nose-up
posture in the dark

We used a videographic assay called the Scalable Apparatus for Measuring Posture and Locomotion, or SAMPL, to measure pitch
axis posture (nose-up/nose-down) and locomotion as fish navigate in depth. In the SAMPL apparatus, larvae swim freely in a
vertical chamber. Behavior is sampled in a specific field of view (400 mm^2) located in the center of the lower half of the chamber,
away from the floor and walls (Figure 1A). Larval zebrafish swim in short epochs comprised of multiple short bursts of translation,
or "bouts" (Figure 1B). During the inter-bout interval (Figure 1C), larvae are largely passive, allowing us to measure postural
instability. To change elevation, larvae climb or dive by adjusting bout kinematics and/or posture (Figure 1D).

We selectively ablated lateral line hair cells by exposing larvae that were 7 days post-fertilization (dpf) to ototoxic levels (10 µM) of
copper sulfate (CuSO$_4$, or "copper") for 90 minutes every 24 hours (Figure 2A) per established protocols (Hernández et al., 2006;
Newton et al., 2023). Treatment produced rapid and complete loss of exposed lateral line hair cells (Figure 2B) while preserving
inner ear vestibular hair cells that are protected by the otic capsule. Using SAMPL, we monitored behavior of copper-treated larvae
and control siblings for 24 hours before and 48 hours after copper exposure in complete darkness. Before treatment, all larvae
showed comparable behavior (Table 1).

Copper-treated larvae sank considerably more than untreated siblings. Vertical displacement during the inter-bout interval was
greater in treated larvae than untreated siblings (-0.32 [0.65] mm vs. 0.02 [0.62] mm; p=0.0473, Figures 2C and 2D, Table 2).
Similarly, copper-treated larvae sank in the vertical axis far more quickly than untreated siblings (-0.27 [0.45] mm s^{-1} vs. -0.01
[0.30] mm s^{-1}; p=0.0183, Figures 2E and 2F). Sinking remained comparable across the duration of the experiment (Table 3), and
can be seen during individual inter-bout intervals (Supplementary Video 1). Finally, in the inter-bout interval, treated larvae also
rotated nose-down more quickly compared to untreated siblings (-6.59 [7.00]° s^{-1} vs. -2.70 [3.71]° s^{-1}; p=0.0060, Figures 2K
and 2L).

In response, copper-treated larvae swam more frequently and adopted a more nose-up pitch than untreated siblings. Their inter-bout
interval duration was significantly shorter than untreated siblings (1.42 [1.15] s vs. 1.78 [2.29] s; p=0.0300, Figures 2G and 2H).
Treated fish demonstrated a more nose-up posture than untreated siblings (12.88 [16.80]° vs. 8.75 [17.74]°; p=0.0473, Figures 2I
and 2J). However, across experiments we observed fewer bouts in treated siblings (14699 vs. 34041, or 43%), suggesting that
changes to posture and locomotion might not be completely restorative.

The sinking, increased bout frequency, increased posture, and increased angular velocity are similar to — but considerably milder
than — behaviors observed in larvae that inflate their swim bladders with oil instead of air (Ehrlich and Schoppik, 2017a;b).
We infer that after loss of the lateral line, larvae adjust their posture and swim frequency to partially compensate for decreased
buoyancy.

Treated larvae rely on postural changes in the dark without altering trajectory

To investigate how lateral line loss might influence bout kinematics, we examined the statistics of swim bouts after copper treatment.
Swim speed was comparable between treatment conditions (copper: 15.37 [10.36] mm s^{-1} vs. control: 11.48 [6.74] mm s^{-1};
p=0.0718). The average bout trajectory was also similar (copper: 13.53 [27.02]° vs. 9.77 [29.33]°; p=0.1481). We conclude that
gross locomotor capacity is comparable between treated and untreated siblings.

Only climbing would countermand sinking associated with copper treatment. Using posture at the beginning of the bout, we
separated bouts into climbs and dives based on their bout trajectories (Figure 3A). In the dark, the fraction of climb bouts was not
significantly different (p=0.0718) between copper-treated (75 [7]%) and control siblings (66 [4]%). When climbing, we observed
comparable depth changes between groups (copper: 0.53 [0.83] mm vs. control: 0.37 [0.57] mm; p=0.0718, Figure 3B). In contrast,
treated larvae moved significantly further in depth during dives compared to controls, consistent with decreased buoyancy (-0.21
[0.34] mm vs. -0.16 [0.23] mm; p=0.0473, Figure 3B).

During climbing and diving bouts, larvae perform two distinct angular rotations using their trunk for different purposes: to change
trajectory in the accelerative phase (steering), and to restore posture during the decelerative phase (righting). These rotations can
be parameterized as the steering and righting gains (Ehrlich and Schoppik, 2017b; Zhu et al., 2023). Steering gain is defined as the
slope of the best-fit line between posture and trajectory evaluated at the time of peak speed (Zhu et al., 2023). If the trajectory of a
given bout can be explained by the posture at the time of peak speed, the steering gain would be 1. Copper-treated larvae exhibited
greater steering gain than untreated siblings (0.69 [0.02] vs. 0.74 [0.02]; p=0.0367, Figure 3C). The gain of the righting rotation is
defined as the slope of the best-fit line between posture at the beginning of the bout and posture change from peak speed to 100 ms
after peak speed. A righting gain of 1 would indicate that the rotation perfectly restored posture. The righting gain did not differ
between copper-treated and sibling larvae (0.18 [0.01] vs. 0.18 [0.02]; p=0.6281, Figure 3D).

Larval zebrafish use a second effector, their pectoral fins, to generate lift when climbing (Ehrlich and Schoppik, 2019; Zhu et al.,
2023). Fin-related lift can be dissociated from steering-related changes using the difference between the predicted trajectory based

on posture and the observed trajectory ("attack angle," Figure 3E). Copper-treated larvae generated less lift, seen as lower attack
angles during climbs, than sibling controls (1.14 [6.59]° vs. 3.76 [6.92]°; p=0.0300, Figure 3F). Given that larvae do not engage
their fins during dives, as expected, dive bout attack angles were comparable between groups (copper: -6.27 [6.15]° vs. control:
-5.25 [5.37]°; p=0.1050).

We conclude that in addition to moving more frequently and adopting a more nose-up posture, copper-treated larvae engage their
trunks more when they climb. However, the average trajectory of bouts is not steeper, they do not climb more frequently, and per
bout, their vertical displacement is comparable.

Treated larvae climb more in the light

We hypothesized that visual input would influence how fish respond to lateral line hair cell loss. We therefore repeated our
experiments in a standard light-dark cycle (LD) to complement data gathered in the dark (DD; Figure 4A). All measurements and
comparisons across illumination conditions used data gathered during circadian day (Tables 2, 5 and 6). As expected (Prober et al.,
2006), both copper-treated and untreated larvae moved more frequently in the light (Figure 4B). The inter-bout interval duration in
the light was shorter for both treatment conditions (copper LD: 0.63 [0.50] s vs. DD: 1.42 [1.52] s; p=0.0010 — untreated LD: 0.69
[0.54] s vs. DD: 1.78 [2.29] s; p=0.0010). During recorded swimming epochs in the light, we observed no difference in movement
frequency between copper-treated and untreated siblings (1.58 [2.00] s^{-1} vs. 1.44 [1.85] s^{-1}; p=0.9000). Across experiments, we
observed fewer total bouts from copper-treated fish than controls in the light (127078 vs. 182534, or 69%), though this difference
was smaller than the 43% observed in the dark.

We next compared linear acceleration during the inter-bout interval to determine if the physical challenges that followed copper
treatment were comparable in the light and the dark. Instability after lateral line ablation was comparable in both the light and dark
during the inter-bout interval with a main effect of both light (p=0.0216) and treatment (p=1.5e-5) but no interaction (p=0.9176).
Copper-treated larvae showed greater acceleration in the downward direction than untreated siblings in both the light and the dark
(LD copper: -0.16 [0.08] mm s^{-2} vs. LD untreated -0.04 [0.03] mm s^{-2}; p=0.0017 — DD copper: -0.24 [0.03] mm s^{-2} vs. DD
untreated: -0.09 [0.03] mm s^{-2}; p=0.0099, Figure 4C). Crucially, downward acceleration did not change between light and dark for
either treatment (copper LD vs. DD: p=0.2900 — untreated LD vs. DD: p=0.3608). As linear acceleration was comparable in both
light and dark regardless of treatment, we conclude that the changes after copper treatment reflect true changes to density.

Given the comparable physical challenges to maintaining elevation (i.e. increased downward acceleration), we next asked if
compensatory strategies were similar after copper-treatment in the light and dark (Tables 5 and 6). Unlike in the dark, copper-
treated larvae in the light climbed with steeper trajectories than untreated siblings (LD copper: 4.28 [21.82]° vs. LD untreated:
-1.29 [18.31]°; p=0.0135, Figure 4D). Copper-treated larvae climbed more frequently (60 [13]%) than control larvae (46 [5]%;
p=0.0013). In the light, copper-treated larvae relied less on trunk steering to climb, instead engaging their fins to produce lift. Fin
engagement (attack angles) during climbs was comparable between copper-treated and untreated siblings in the light (LD control:
1.99 [4.33]° vs. LD copper: 1.71 [4.72]°; p=0.9000, Figure 4E). In the light, steering gain (trunk use) was unchanged between
copper-treated larvae than untreated siblings (control: 0.75 [0.03] vs. copper: 0.72 [0.02]; p=0.2755, Figure 4F), as was righting
gain (control: 0.09 [0.01] vs. copper: 0.10 [0.01]; p=0.7969, Figure 4G). However, similar to copper-treated larvae in the dark,
those in the light adopted a more nose-up posture than untreated siblings (4.96 [16.71]° vs. -0.37 [14.84]°; p=0.0037, Figure 4H).

Just as in the dark, copper-treated larvae in the light changed their behavior by increasing posture. However, unlike fish in the
dark, copper-treated larvae did not move more, nor did they change bout kinematics (attack angle and steering gain). Instead,
they climbed more frequently, and with steeper trajectories than untreated siblings. We conclude that ambient light can change
compensatory strategies after lateral line hair cell loss.

Ambient light improves elevation efficiency

In both the dark and the light, we observed fewer total bouts from copper-treated larvae compared to untreated siblings (Table 4).
We hypothesized that the decrease in observed bouts was due to ineffective elevation control. If so, copper-treated larvae would sit
at the floor of the chamber outside of the normal field of view of the camera. We adjusted the arena to observe the chamber floor
and observed larvae position in dark and light conditions (Figure 5A). There were main effects of both light condition (Figure 5B;
p=3.8e-10) and treatment (p=1.4e-27), as well as an interaction between the two (p=5.8e-4). Specifically, copper-treated larvae
were positioned at the bottom of the arena more often than controls in the dark (12.5 [12.3]% vs. 54.1 [50.6]% ; p=0.001) and in
the light (4.8 [12.5]% vs. 33.3 [13.8]% ; p=0.001). Fewer copper-treated larvae were observed on the floor in the light relative to
the dark. Importantly, ambient light improved elevation in copper-treated larvae (p=0.001), but not sibling controls (p=0.1188).

Given the downward acceleration, fewer bouts observed, and number of larvae observed at the arena floor, we conclude that lateral
line ablation compromises the ability of larvae to maintain elevation. The physical challenges faced by copper-treated larvae were
comparable between the light and the dark. We therefore propose that the strategies adopted in the light are more effective at
compensating for decreased buoyancy.

Larvae recover postural stability after lateral line regeneration

We hypothesized that buoyancy would be restored following regeneration of lateral line hair cells. Larvae were treated with copper
either in an early phase (7–9 dpf) or late phase (14–16 dpf) and compared to untreated control siblings (Figure 6A). Copper

treatment killed lateral line hair cells similarly at both time points (Figure 2B and Figure 6B), but early-treated fish had regenerated
their hair cells by 14 dpf (Figure 6B). Similar to exposure at 7 dpf, late-treated larvae sank faster than controls (-0.23 [0.46] mm
s^{-1}; p=0.0013) and early-treated animals (p=0.0012). In the early phase, larvae presented similarly to data presented in Figures 2
and 3. Here, we describe data collected in the late phase only.

Larval buoyancy returns to control levels after hair cells regenerate. Vertical velocity between bouts was similar for early-treated
larvae and controls (-0.03 [0.24] mm s^{-1} vs. -0.04 [0.31] mm s^{-1}; p=0.9000, Figure 6C). Other properties of the inter-bout interval
also recovered after lateral line regeneration. The inter-bout interval duration of early-treated larvae returned to control values (2.92
[3.49] s vs. 2.50 [3.69] s; p=0.7858, Figure 6D). Early-treated animals also held similar postures to untreated siblings (7.08 [18.09]°
vs. 11.25 [21.51]°; p=0.5569). We conclude that that 1) lateral line regeneration resolves the positional challenges imposed by
copper treatment, and 2) lateral line loss at 14 dpf similarly affects buoyancy.

Because early-treated larvae sank less and improved stability after hair cell regeneration, we asked whether their compensatory
climbing behaviors persisted. Early-treated and control larvae exhibited similar attack angles (10.68 [6.21]° vs. 9.77 [6.80]°;
p=0.1860; Figure 6E), which were comparable to late-treated animals (7.55 [6.11]°; p=0.0639). Larvae across all three groups
performed similar fractions of climbs and dives (control: 71[7]%; early: 68 [2]%; late: 71 [20]%; p=0.6314), consistent with
previous observations in the dark. However, unlike compensatory actions at 7 dpf, late-treated larvae swam faster than sibling
controls (13.07 [9.91] mm s^{-1} v. 10.46 [6.56] mm s^{-1}; p=0.0330; Figure 6F) and early-treated larvae (9.88 [5.89] mm s^{-1}; p=0.0119),
while control and early-treated larvae displayed similar speeds (p=0.3964).

Early-treated larvae swam comparably to controls after hair cell regeneration. While late-treated larvae sank like early controls,
the major change to their behavior was an increase in swim speed. We conclude that hair cell regeneration is sufficient to recover
buoyancy, and that stabilizing strategies can change with age.

DISCUSSION

We hypothesized that postural control would shift to incorporate multisensory information, and investigated the ways larval ze-
brafish maintain elevation when buoyancy is challenged. Lateral line loss causes fish to become denser and sink, which triggers
changes to behavior that vary in the light and the dark. Changes to behavior were only partially effective, but differentially so:
treated fish were more likely to rest on the bottom of the arena, but less so in the light. Finally, sinking and compensatory changes
recovered after peripheral regeneration, demonstrating transient effects of ablation. We conclude that fish have multiple strategies
for maintaining elevation, and selectively engage them depending on available sensory information. The behavioral flexibility we
observe could act as a substrate for selective pressures that gave rise to the vast array of postures and preferred elevations among
fish.

We observed that larvae sank after copper treatment. The small increase in downward acceleration we observed was stable across
clutches of fish and illumination. While establishing the physiological mechanism is beyond the scope of this paper, the most likely
explanation for the changes we observe is that acute loss of the lateral line causes larvae to slightly deflate their swim bladders
(Dumbarton et al., 2010). Early lateral line manipulations can induce hyperbuoyancy (Lin et al., 2023; Venuto et al., 2023); our
results expand this link between the lateral line and swim bladder volume. Importantly, near-neutral buoyancy is re-established
after lateral line regeneration. We conclude that the inferred decrease in swim bladder volume is reversible.

Illumination-dependent findings reflect limits of behavior fish can produce when challenged with lateral line loss. First, challenged
larvae swim more frequently than control siblings in the dark, with less fin engagement (lower attack angles). Neither swim
frequency nor fin engagement changed in the light, presumably because in the light, both are at their ceiling. In the light larvae
climbed more frequently with greater average trajectories. A similar ceiling for climb fraction and trajectory likely exists in the
dark. While we restricted our analysis to comparable points in time, it would be interesting to explore these ceilings in the context
of circadian day and night, particularly given diel-related elevation changes in fish (Helfman, 1986).

The strategies we observed were more effective in the light than in the dark. Evidence comes from two sources: first, we observed
greater number of observed bouts (which can only happen if fish are off the bottom of the arena) in the light; second, a greater
fraction of fish were observed on the bottom of the arena in the dark. The change to buoyancy, measured by downward acceleration
after treatment, was comparable in the light and the dark. While there was a significant change in vertical displacement per climb
bout in the light, the effect size was quite small. Instead, we propose that the improvement in the light reflects an increased
fraction of climbs that, together, allows for elevation changes. Larval zebrafish sequence climbs and dives to change elevation
when navigating in depth (Zhu et al., 2024). We propose that a similar mechanism explains improved control of elevation we
observed in the light.

We propose that the changes we observe when larvae are challenged might reflect differential engagement of neural circuits respon-
sible for postural set point, swim frequency, and fin control. Two likely, but not mutually exclusive, substrates for these behaviors
are the cerebellum and the interstitial nucleus of Cajal, also called the nucleus of the medial longitudinal fasciculus (INC/nMLF).
First, the cerebellum is a hub of sensory integration (Ishikawa et al., 2015), modulating vestibular function (Auer et al., 2023;
Manzoni, 2007) and motor activity (Cullen, 2023; Manzoni, 2007). The cerebellum has been reported to modulate fin-body coor-
dination (Auer et al., 2023; Ehrlich and Schoppik, 2019), a key difference in the strategies observed. Next, the INC/nMLF is likely
engaged during different compensatory tactics given its activity during various swim modes (Sankrithi and O'Malley, 2010) and

involvement in posture and navigation (Sugioka et al., 2023; Zhu et al., 2024). The INC/nMLF is responsive to light, head taps (Sankrithi and O'Malley, 2010), body tilts (Sugioka et al., 2023), and horizontal visual motion (Naumann et al., 2016). Notably, visual cues during sinking might induce an optomotor response, similar to reported optokinetic responses (Bianco et al., 2012; Dehmelt et al., 2021; Matsuda and Kubo, 2021) that could engage compensatory behaviors. Loss-of-function and functional experiments comparing neural responses in the light and dark will speak directly to the circuits responsible for strategy selection and implementation to maintain elevation.

Dealing with gravity is a biomechanical challenge for all animals. Without a substrate, maintaining elevation becomes vital for aquatic and airborne creatures. We discovered that larval zebrafish become denser and sink after acute lateral line loss, enabling comparisons of compensatory strategies with and without ambient light. Larvae correct for destabilization by climbing using a combination of postural and kinematic adjustments. As hypothesized, these mechanisms diverged when visual information was available into distinct strategies, with different efficacy. Our work reveals separate behaviors to accomplish the same purpose — maintaining elevation — framing future investigation into how different sensory inputs and locomotor actions regulate position. Different behavioral strategies can serve as a substrate for natural selection (Brown and Wilson, 1956). Understanding how animals change behavior when challenged extends existing studies into morphological (Drucker, 2002) and locomotor specializations (Fish and Lauder, 2006; Tytell et al., 2010) well-suited to different environmental niches.

ACKNOWLEDGMENTS

Research was supported by the National Institute on Deafness and Communication Disorders of the National Institutes of Health under award numbers R01DC017489, the National Institute of Neurological Disorders and Stroke under award number R61NS125280, by the Leon Levy Foundation (YZ), the Rainwater Charitable Foundation (YZ) and by the Irma T. Hirschl/Monique Weill-Caulier Trust (DS). The authors would like to thank Hannah Gelnaw for assistance with fish care, Lavinia Sheets and lab for sharing the *Tg(myo6b:actb1-EGFP)* line, Dena Goldblatt and other members of the Schoppik and Nagel labs for their valuable feedback and discussions.

AUTHOR CONTRIBUTIONS

Conceptualization: YZ; Methodology: YZ, SD; Investigation: SD, YZ; Visualization: SD; Writing: DS, SD; Editing: SD, DS; Funding Acquisition: DS, YZ; Supervision: DS.

AUTHOR COMPETING INTERESTS

The authors have no competing interests to declare.

REFERENCES

R. M. Alexander. Size, speed and buoyancy adaptations in aquatic animals. *American Zoologist*, 30(1):189–196, Feb. 1990. ISSN 0003-1569. doi: 10.1093/icb/30.1.189. URL http://dx.doi.org/10.1093/icb/30.1.189.

F. Auer, K. Nardone, K. Matsuda, M. Hibi, and D. Schoppik. Purkinje cells control posture in larval zebrafish (*Danio rerio*). *bioRxiv*, Sept. 2023. doi: 10.1101/2023.09.12.557469. URL https://doi.org/10.1101/2023.09.12.557469.

I. H. Bianco, L.-H. Ma, D. Schoppik, D. N. Robson, M. B. Orger, J. C. Beck, J. M. Li, A. F. Schier, F. Engert, and R. Baker. The tangential nucleus controls a gravito-inertial vestibulo-ocular reflex. *Current Biology*, 22(14):1285–1295, Jul 2012. doi: 10.1016/j.cub.2012.05.026. URL http://dx.doi.org/10.1016/j.cub.2012.05.026.

W. L. Brown and E. O. Wilson. Character displacement. *Systematic Zoology*, 5(2):49, June 1956. ISSN 0039-7989. doi: 10.2307/2411924. URL http://dx.doi.org/10.2307/2411924.

K. E. Cullen. Internal models of self-motion: neural computations by the vestibular cerebellum. *Trends in Neurosciences*, 46(11):986–1002, Nov. 2023. ISSN 0166-2236. doi: 10.1016/j.tins.2023.08.009. URL http://dx.doi.org/10.1016/j.tins.2023.08.009.

S. N. Davis, P. Wu, E. D. Camci, J. A. Simon, E. W. Rubel, and D. W. Raible. Chloroquine kills hair cells in zebrafish lateral line and murine cochlear cultures: Implications for ototoxicity. *Hearing Research*, 395:108019, Sept. 2020. ISSN 0378-5955. doi: 10.1016/j.heares.2020.108019. URL http://dx.doi.org/10.1016/j.heares.2020.108019.

F. A. Dehmelt, R. Meier, J. Hinz, T. Yoshimatsu, C. A. Simacek, R. Huang, K. Wang, T. Baden, and A. B. Arrenberg. Spherical arena reveals optokinetic response tuning to stimulus location, size, and frequency across entire visual field of larval zebrafish. *eLife*, 10, June 2021. ISSN 2050-084X. doi: 10.7554/elife.63355. URL http://dx.doi.org/10.7554/eLife.63355.

E. G. Drucker. Wake dynamics and locomotor function in fishes: Interpreting evolutionary patterns in pectoral fin design. *Integrative and Comparative Biology*, 42(5):997–1008, Nov. 2002. ISSN 1557-7023. doi: 10.1093/icb/42.5.997. URL http://dx.doi.org/10.1093/icb/42.5.997.

T. C. Dumbarton, M. Stoyek, R. P. Croll, and F. M. Smith. Adrenergic control of swimbladder deflation in the zebrafish (danio rerio). *Journal of Experimental Biology*, 213(14):2536–2546, July 2010. ISSN 0022-0949. doi: 10.1242/jeb.039792. URL http://dx.doi.org/10.1242/jeb.039792.

D. E. Ehrlich and D. Schoppik. Control of movement initiation underlies the development of balance. *Current Biology*, 27(3):334–344, Feb. 2017a. ISSN 0960-9822. doi: 10.1016/j.cub.2016.12.003. URL http://dx.doi.org/10.1016/j.cub.2016.12.003.

D. E. Ehrlich and D. Schoppik. A novel mechanism for volitional locomotion in larval zebrafish. *bioRxiv*, Sept. 2017b. doi: 10.1101/189191. URL http://dx.doi.org/10.1101/189191.

D. E. Ehrlich and D. Schoppik. A primal role for the vestibular sense in the development of coordinated locomotion. *eLife*, 8, Oct. 2019. doi: 10.7554/elife.45839. URL https://doi.org/10.7554/elife.45839.

R. Fänge. Physiology of the swimbladder. *Physiological Reviews*, 46(2):299–322, Apr. 1966. ISSN 1522-1210. doi: 10.1152/physrev.1966.46.2.299. URL http://dx.doi.org/10.1152/physrev.1966.46.2.299.

F. Fish and G. Lauder. Passive and active flow control by swimming fishes and mammals. *Annual Review of Fluid Mechanics*, 38(1):193–224, Jan. 2006. ISSN 1545-4479. doi: 10.1146/annurev.fluid.38.050304.092201. URL http://dx.doi.org/10.1146/annurev.fluid.38.050304.092201.

F. E. Fish and R. Holzman. Swimming turned on its head: Stability and maneuverability of the shrimpfish (aeoliscus punctulatus). *Integrative Organismal Biology*, 1(1), Jan. 2019. ISSN 2517-4843. doi: 10.1093/iob/obz025. URL http://dx.doi.org/10.1093/iob/obz025.

J. H. Gee and H. M. Holst. Buoyancy regulation by the sticklebacks culaea inconstans and pungitius pungitius in response to different salinities and water densities. *Canadian Journal of Zoology*, 70(8):1590–1594, Aug. 1992. ISSN 1480-3283. doi: 10.1139/z92-219. URL http://dx.doi.org/10.1139/z92-219.

E. M. Goolish and K. Okutake. Lack of gas bladder inflation by the larvae of zebrafish in the absence of an air-water interface. *Journal of Fish Biology*, 55(5): 1054–1063, Nov. 1999. ISSN 1095-8649. doi: 10.1111/j.1095-8649.1999.tb00740.x. URL http://dx.doi.org/10.1111/j.1095-8649.1999.tb00740.x.

H. H. Harvey. *Pressure in the early life history of sockeye salmon*. University of British Columbia, 1963. doi: 10.14288/1.0105582. URL https://doi.library.ubc.ca/10.14288/1.0105582.

H. H. Harvey and C. R. Bothern. Compensatory swimming in the kokanee and sockeye salmon oncorhynchus nerka (walbaum). *Journal of Fish Biology*, 4 (2):237–247, Apr. 1972. ISSN 1095-8649. doi: 10.1111/j.1095-8649.1972.tb05670.x. URL http://dx.doi.org/10.1111/j.1095-8649.1972.tb05670.x.

G. S. Helfman. *Fish Behaviour by Day, Night and Twilight*, page 366–387. Springer US, 1986. ISBN 9781468482614. doi: 10.1007/978-1-4684-8261-4_14. URL http://dx.doi.org/10.1007/978-1-4684-8261-4_14.

P. P. Hernández, V. Moreno, F. A. Olivari, and M. L. Allende. Sub-lethal concentrations of waterborne copper are toxic to lateral line neuromasts in zebrafish (danio rerio). *Hearing Research*, 213(1–2):1–10, Mar. 2006. ISSN 0378-5955. doi: 10.1016/j.heares.2005.10.015. URL http://dx.doi.org/10.1016/j.heares.2005.10.015.

T. Ishikawa, M. Shimuta, and M. Häusser. Multimodal sensory integration in single cerebellar granule cells in vivo. *eLife*, 4, Dec. 2015. ISSN 2050-084X. doi: 10.7554/elife.12916. URL http://dx.doi.org/10.7554/eLife.12916.

A. Lin, E. Álvarez Salvado, N. Milicic, N. Pujara, and D. E. Ehrlich. Multisensory navigational strategies of hatchling fish for dispersal. *Current Biology*, 33(22): 4917–4925.e4, Nov. 2023. ISSN 0960-9822. doi: 10.1016/j.cub.2023.09.070. URL http://dx.doi.org/10.1016/j.cub.2023.09.070.

B. W. Lindsey, T. C. Dumbarton, S. J. Moorman, F. M. Smith, and R. P. Croll. Effects of simulated microgravity on the development of the swimbladder and buoyancy control in larval zebrafish (danio rerio). *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 315A(5):302–313, Mar. 2011. ISSN 1932-5231. doi: 10.1002/jez.677. URL http://dx.doi.org/10.1002/jez.677.

J. Lötsch and A. Ultsch. A non-parametric effect-size measure capturing changes in central tendency and data distribution shape. *PLOS ONE*, 15(9):e0239623, Sept. 2020. ISSN 1932-6203. doi: 10.1371/journal.pone.0239623. URL http://dx.doi.org/10.1371/journal.pone.0239623.

D. Manzoni. The cerebellum and sensorimotor coupling: Looking at the problem from the perspective of vestibular reflexes. *The Cerebellum*, 6(1), Mar. 2007. ISSN 1473-4230. doi: 10.1080/14734220601132135. URL http://dx.doi.org/10.1080/14734220601132135.

K. Matsuda and F. Kubo. Circuit organization underlying optic flow processing in zebrafish. *Frontiers in Neural Circuits*, 15, July 2021. ISSN 1662-5110. doi: 10.3389/fncir.2021.709048. URL http://dx.doi.org/10.3389/fncir.2021.709048.

M. J. McHenry and G. V. Lauder. The mechanical scaling of coasting in zebrafish (danio rerio). *Journal of Experimental Biology*, 208(12):2289–2301, June 2005. ISSN 0022-0949. doi: 10.1242/jeb.01642. URL http://dx.doi.org/10.1242/jeb.01642.

D. L. Meyer, W. Heiligenberg, and T. H. Bullock. The ventral substrate response: A new postural control mechanism in fishes. *Journal of Comparative Physiology A*, 109(1):59–68, 1976a. ISSN 1432-1351. doi: 10.1007/bf00663435. URL http://dx.doi.org/10.1007/BF00663435.

D. L. Meyer, C. Platt, and H. J. Distel. Postural control mechanisms in the upside-down catfish (synodontis nigriventris). *Journal of Comparative Physiology A*, 110(3):323–331, 1976b. ISSN 1432-1351. doi: 10.1007/bf00659148. URL http://dx.doi.org/10.1007/BF00659148.

D. L. Meyer, R. Becker, and W. Graf. The ventral substrate response of fishes: Comparative investigation of the vsr about the roll and the pitch axis. *Journal of Comparative Physiology A*, 117(2):209–217, 1977. ISSN 1432-1351. doi: 10.1007/bf00612787. URL http://dx.doi.org/10.1007/BF00612787.

E. A. Naumann, J. E. Fitzgerald, T. W. Dunn, J. Rihel, H. Sompolinsky, and F. Engert. From whole-brain data to functional circuit models: The zebrafish optomotor response. *Cell*, 167(4):947–960.e20, Nov. 2016. ISSN 0092-8674. doi: 10.1016/j.cell.2016.10.019. URL http://dx.doi.org/10.1016/j.cell.2016.10.019.

K. C. Newton, D. Kacev, S. R. O. Nilsson, A. L. Saettele, S. A. Golden, and L. Sheets. Lateral line ablation by ototoxic compounds results in distinct rheo-taxis profiles in larval zebrafish. *Communications Biology*, 6(1), Jan. 2023. doi: 10.1038/s42003-023-04449-2. URL https://doi.org/10.1038/s42003-023-04449-2.

B. Pelster. *5 Buoyancy At Depth*, page 195–237. Elsevier, 1997. ISBN 9780123504401. doi: 10.1016/s1546-5098(08)60230-7. URL http://dx.doi.org/10.1016/S1546-5098(08)60230-7.

B. Pelster. *Buoyancy Control in Aquatic Vertebrates*, page 65–98. Springer Berlin Heidelberg, 2009. ISBN 9783540939856. doi: 10.1007/978-3-540-93985-6_4. URL http://dx.doi.org/10.1007/978-3-540-93985-6_4.

C. Platt. Central control of postural orientation in flatfish. *Journal of Experimental Biology*, 59(2):491–521, Oct. 1973. ISSN 1477-9145. doi: 10.1242/jeb.59.2.491. URL http://dx.doi.org/10.1242/jeb.59.2.491.

D. A. Prober, J. Rihel, A. A. Onah, R.-J. Sung, and A. F. Schier. Hypocretin/orexin overexpression induces an insomnia-like phenotype in zebrafish. *The Journal of Neuroscience*, 26(51):13400–13410, Dec. 2006. ISSN 1529-2401. doi: 10.1523/jneurosci.4332-06.2006. URL http://dx.doi.org/10.1523/JNEUROSCI.4332-06.2006.

N. Sankrithi and D. O'Malley. Activation of a multisensory, multifunctional nucleus in the zebrafish midbrain during diverse locomotor behaviors. *Neuroscience*, 166(3):970–993, Mar. 2010. ISSN 0306-4522. doi: 10.1016/j.neuroscience.2010.01.003. URL http://dx.doi.org/10.1016/j.neuroscience.2010.01.003.

J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. Fiji: an open-source platform for biological-image analysis. *Nature Methods*, 9(7):676–682, June 2012. ISSN 1548-7105. doi: 10.1038/nmeth.2019. URL http://dx.doi.org/10.1038/nmeth.2019.

J. B. Steen. *10 The Swim Bladder as a Hydrostatic Organ*, page 413–443. Elsevier, 1970. ISBN 9780123504043. doi: 10.1016/s1546-5098(08)60135-1. URL http://dx.doi.org/10.1016/S1546-5098(08)60135-1.

T. Sugioka, M. Tanimoto, and S. ichi Higashijima. Biomechanics and neural circuits for vestibular-induced fine postural control in larval zebrafish. *Nature Communications*, 14(1), Mar. 2023. doi: 10.1038/s41467-023-36682-y. URL https://doi.org/10.1038/s41467-023-36682-y.

E. D. Thomas, I. A. Cruz, D. W. Hailey, and D. W. Raible. There and back again: development and regeneration of the zebrafish lateral line system. *WIREs Developmental Biology*, 4(1):1–16, Oct. 2014. ISSN 1759-7692. doi: 10.1002/wdev.160. URL http://dx.doi.org/10.1002/wdev.160.

E. D. Tytell, I. Borazjani, F. Sotiropoulos, T. V. Baker, E. J. Anderson, and G. V. Lauder. Disentangling the functional roles of morphology and motion in the swimming of fish. *Integrative and Comparative Biology*, 50(6):1140–1154, May 2010. ISSN 1557-7023. doi: 10.1093/icb/icq057. URL http://dx.doi.org/10.1093/icb/icq057.

A. Venuto, S. Thibodeau-Beganny, J. G. Trapani, and T. Erickson. A sensation for inflation: initial swim bladder inflation in larval zebrafish is mediated by the mechanosensory lateral line. *Journal of Experimental Biology*, 226(11), June 2023. doi: 10.1242/jeb.245635. URL https://doi.org/10.1242/jeb.245635.

P. W. Webb. Control of posture, depth, and swimming trajectories of fishes. *Integrative and Comparative Biology*, 42(1):94–101, 2002. doi: 10.1093/icb/42.1.94. URL https://doi.org/10.1093/icb/42.1.94.

P. W. Webb. Stability and maneuverability. *Fish physiology*, 23:281, 2006.

N. G. Willoughby. The buoyancy and orientation of the upside-down catfishes of the genus synodontis (pisces: Siluroidei). *Journal of Zoology*, 180(3):291–314, Nov. 1976. ISSN 1469-7998. doi: 10.1111/j.1469-7998.1976.tb04680.x. URL http://dx.doi.org/10.1111/j.1469-7998.1976.tb04680.x.

Y. Zhu, F. Auer, H. Gelnaw, S. N. Davis, K. R. Hamling, C. E. May, H. Ahamed, N. Ringstad, K. I. Nagel, and D. Schoppik. Sampl is a high-throughput solution to study unconstrained vertical behavior in small animals. *Cell Reports*, 42(6):112573, June 2023. ISSN 2211-1247. doi: 10.1016/j.celrep.2023.112573. URL http://dx.doi.org/10.1016/j.celrep.2023.112573.

Y. Zhu, H. Gelnaw, F. Auer, K. R. Hamling, D. E. Ehrlich, and D. Schoppik. A brainstem circuit for gravity-guided vertical navigation. *bioRxiv*, Mar. 2024. doi: 10.1101/2024.03.12.584680. URL http://dx.doi.org/10.1101/2024.03.12.584680.

Figure 1: Physical forces and kinematic parameters of larval locomotion in the vertical plane.
(A) Example field of view (4 cm^2) with a 6 dpf larva. Inset shows the pitch of the larva, defined as the angle between the long axis of the body and the horizon; the gravity vector acting at the center of mass; and the buoyancy vector acting at the center of volume.
(B) Representative sequence of swim bouts, plotted as speed (top) and pitch (bottom) as a function of time. Positive values for pitch are nose-up.
(C) Speed trace of two swim bouts illustrating the speed threshold (dashed line; 5 mm s^{-1}) that defines the active bout, as well as the passive inter-bout interval.
(D) Schematic of larvae at the beginning and end of a climb and dive bout, illustrating the vertical displacement.

Figure 2: Copper treatment ablates lateral line hair cells and disrupts stability between bouts.

(A) Experimental timeline indicating pre-treatment recording (grey), treatment periods (dark orange), and post-treatment recording (light orange). Full sample numbers and statistics in Table 2.

(B) Confocal images of a control (left) and a copper-treated (right) SO2 neuromast transgenically labelled with EGFP in 7 dpf larvae. Complete loss of hair cells follows copper treatment. Scale bar = 10 μm.

(C) Schematic (left) and distribution (right) of vertical displacement between bouts. Orange are copper-treated larvae, black untreated siblings.

(D) Median displacement in depth during the inter-bout interval is significantly lower in copper-treated larvae (p=0.0473). Grey circles represent experimental repeats.

(E) Schematic and distributions of vertical velocity during the inter-bout interval.

(F) Median vertical velocity between bouts is lower (i.e. downward) in copper-treated larvae (p=0.0183).

(G) Schematic (left) and distribution (right) of inter-bout interval duration.

(H) Median inter-bout interval duration is lower for copper-treated larvae (p=0.0300).

(I) Schematic (left) and distribution (right) of inter-bout interval pitch.

(J) Median inter-bout interval pitch is higher in copper-treated larvae (p=0.0473).

(K) Schematic (left) and distribution (right) of angular velocity during the inter-bout interval.

(L) Median inter-bout interval angular velocity is lower (i.e. nose-down rotation) in copper-treated larvae. (p=0.0060).

Mann-Whitney U test. P, probability.

Figure 3: Copper-treated larvae rely on trunk rotations to produce climb bouts in the dark.

(A) Schematic (left) and distribution (right) of displacement in depth during climbing and diving bouts.

(B) Median displacement in depth is unchanged for climbs (top; p=0.0718) but greater for copper-treated diving bouts (bottom; p=0.0473).

(C) Steering rotation gain is higher in copper-treated larvae (p=0.0367).

(D) Righting rotation gain does not change between treatment groups (p=0.6281).

(E) Schematic (left) and distribution (right) of attack angle during climbing and diving bouts.

(F) Median attack angle is lower in copper-treated larvae for climbs (top; p=0.0300) but not dives (bottom; p=0.1050).

See Table 4 for full breakdown of bouts. Mann-Whitney U test. P, probability; NS, not significant.

Figure 4: Copper-treated larvae are similarly unstable and adopt different posture and kinematic strategies in the light and in the dark

(A) Experimental timeline of each lighting condition (filled = dark, open = light) and treatment (grey = control, orange = copper). Treatment times indicated in dark orange, and dotted rectangles show time window during which behavior was compared. Full sample numbers and statistics in Tables 5 and 6.

(B) Inter-bout interval (IBI) durations are longer in the dark than in the light (control: p=0.0010, copper: p=0.0010). Copper-treatment shortens IBI duration in the dark (p=0.0033) but not the light (p=0.9000).

(C) Vertical acceleration during the IBI is lower (more downward) in copper-treated larvae in both the dark (p=0.0099) and the light (p=0.0017). Acceleration did not vary as a function of light for control (p=0.3608) or copper-treated larvae (p=0.2900).

(D) Median swim trajectory of copper-treated larvae is higher than untreated siblings in the light (p=0.0135) but not the dark (p=0.3277). Trajectories are lower (more horizontal) in the light (control: p=0.0010, copper: p=0.0010).

(E–H) Data are plotted as the difference between control and copper-treated larvae for each experimental repeat for both lighting conditions. Dotted lines at 0 indicate no difference between copper-treated and untreated siblings. P values are comparisons of the differences in the dark vs. light.

(E) Differences in attack angle for climb bouts are small in the light, unlike the dark (p=0.0001).

(F) Differences in steering rotation gain are higher in the dark than in the light (p=0.0017).

(G) Differences in righting rotation gain are unchanged (p=0.9807).

(H) Differences in IBI pitch are higher but not significantly different in the light (p=0.5123).

See Table 4 for full breakdown of bouts. B–D: Two-way ANOVA with Tukey's HSD. E–I: Mann-Whitney U test (multiple comparison p-values listed in Table 5). IBI, inter-bout interval; NS, not significant.

Figure 5: Copper-treated larvae are inefficient at maintaining elevation, but are better in the light

(A) Experimental timeline of each lighting condition (filled = dark, open = light) and treatment (grey = control, orange = copper). Treatment times indicated in dark orange, and dotted lines demonstrate observation times. Dark control (N=36), Dark copper (N=31), Light control (N=57), Light copper (N=57).

(B) Fraction of larvae resting on the chamber floor is greater after lateral line ablation in both the dark (control v. copper: p=0.001) and light (control v. copper: p=0.001). Larvae in the light were less likely to rest on the floor between copper-treated groups (p=0.001) but not controls (p=0.1188). Two-way ANOVA with Tukey's HSD.

Figure 6: Copper-treated larvae are similarly unstable and adopt different posture and kinematic strategies in the light and in the dark

(A) Experimental timeline of each condition in the early (7–9 dpf) and late phase (14–16 dpf). Color indicates treatment (grey = control, orange = acute copper ablation, green = regenerated hair cells). Treatment times indicated in dark orange, dotted rectangle shows time window during which behavior was compared. Full sample numbers and statistics in Table 7.

(B) Confocal imaging of SO2 neuromast hair cells at 14 dpf in representative control (left), early-treated (center), and late-treated larvae (right). Hair cells were regenerated to controls in early-treated larvae, and similarly ablated by copper in late-treated siblings. Scale bar = 10 μm.

(C) Vertical velocity during the IBI is similar between early-treated and control animals (p=0.9000), while late-treated larvae sink significantly (late v. control: p=0.0013; late v. early: p=0.0012).

(D) IBI duration is similar between all groups (p=0.1832).

(E) Attack angle during climb bouts is comparable between all groups (p=0.0671).

(F) Peak speed is similar between early-treated and control animals (p=0.3964), but increased in late-treated larvae (late v. control: p=0.0330; late v. early: p=0.0119).

See Table 4 for full breakdown of bouts. One-way ANOVA with Tukey's HSD post-hoc comparison. disp, displacement; IBI, inter-bout interval; NS, not significant.

Table 1: Pre-treatment results for dark-dark and light-dark datasets. Median and interquartile range (IQR) are listed for non-normal distributions of raw data. P-value calculated using the Mann-Whitney U test ($\alpha = 0.05$), and effect size calculated using Cohen's d. ^Steering and righting rotation gain p-value calculated using unpaired t-test (mean and standard deviation listed).

Parameter	Unit	Control	Copper	P-Value [Effect size]
Dark-Dark Dataset				
Inter-bout intervals				
Vertical displacement	mm	0.03 [0.59]	-0.02 [0.66]	0.5000 [0.10]
Vertical velocity	mm s^{-1}	-0.04 [0.33]	-0.07 [0.40]	0.4172 [0.10]
Duration	s	1.48 [2.15]	1.52 [2.24]	0.3002 [0.03]
Pitch	deg	14.39 [20.57]	15.81 [18.78]	0.2654 [0.09]
Angular velocity	° s^{-1}	-1.43 [4.23]	-2.09 [4.52]	**0.0473** [0.20]
Bouts				
Peak speed	mm s^{-1}	11.36 [6.14]	11.01 [6.75]	0.1481 [0.07]
Trajectory	deg	20.84 [32.90]	23.37 [30.31]	0.2017 [0.11]
Vertical displ (climb)	mm	0.54 [0.68]	0.56 [0.73]	0.5000 [0.04]
Vertical displ (dive)	mm	-0.17 [0.25]	-0.16 [0.23]	0.2017 [0.09]
Attack angle (climb)	deg	6.37 [6.94]	6.27 [7.66]	0.4172 [0.01]
Attack angle (dive)	deg	-5.45 [5.68]	-5.81 [5.92]	0.1481 [0.08]
Steering rotation gain^		0.68 [0.02]	0.66 [0.03]	0.3975 [0.56]
Righting rotation gain^		0.17 [0.01]	0.18 [0.01]	0.3027 [0.69]
Fraction of climbs	%	74.58 [0.97]	81.09 [6.77]	0.2017 [0.97]
Fraction of dives	%	24.09 [0.90]	18.14 [7.00]	0.2017 [0.94]
Light-Dark Dataset				
Inter-bout intervals				
Vertical displacement	mm	-0.08 [0.19]	-0.07 [0.19]	0.1350 [0.05]
Vertical velocity	mm s^{-1}	-0.15 [0.23]	-0.13 [0.21]	0.2817 [0.11]
Duration	s	0.62 [0.34]	0.65 [0.42]	0.2472 [0.13]
Pitch	deg	0.08 [15.03]	0.62 [15.59]	0.3964 [0.04]
Angular velocity	° s^{-1}	-2.20 [3.85]	-1.97 [3.53]	0.3566 [0.08]
Bouts				
Peak speed	mm s^{-1}	12.41 [7.07]	11.91 [6.53]	0.3182 [0.09]
Trajectory	deg	0.001 [3.31]	0.16 [4.04]	0.4790 [0.03]
Vertical displ (climb)	mm	0.11 [0.19]	0.11 [0.20]	0.4790 [0.01]
Vertical displ (dive)	mm	-0.11 [0.17]	-0.10 [0.25]	0.2474 [0.07]
Attack angle (climb)	deg	1.42 [3.46]	1.28 [3.39]	0.2474 [0.05]
Attack angle (dive)	deg	-0.98 [3.13]	-1.24 [3.20]	0.1860 [0.11]
Steering rotation gain^		0.73 [0.03]	0.74 [0.03]	0.5436 [0.31]
Righting rotation gain^		0.07 [0.01]	0.07 [0.008]	0.3120 [0.52]
Fraction of climbs	%	48.25 [10.42]	49.18 [12.23]	0.3964 [0.10]
Fraction of dives	%	48.76 [10.13]	47.91 [12.70]	0.3964 [0.09]

Table 2: Effects of lateral line ablation in the dark. Median [IQR] used for non-normal distributions of raw data. N = 114 control larvae, 114 copper-treated larvae. P-value calculated using Mann-Whitney U test (α = 0.05), and effect size calculated using Cohen's d. ^Steering and righting rotation gain p-value calculated using unpaired t-test (mean and standard deviation listed).

Parameter	Unit	Control	Copper	P-Value [Effect size]
Inter-bout Interval				
Vertical displacement	mm	0.02 [0.62]	-0.32 [0.65]	**0.0473** [0.73]
Vertical velocity	mm s^{-1}	-0.01 [0.30]	-0.27 [0.45]	**0.0183** [1.00]
Duration	s	1.78 [2.29]	1.42 [1.15]	**0.0300** [0.29]
Pitch	deg	8.75 [17.74]	12.88 [16.80]	**0.0473** [0.32]
Angular velocity	° s^{-1}	-2.70 [3.71]	-6.59 [7.00]	**0.0060** [1.06]
Bout				
Peak speed	mm s^{-1}	11.48 [6.74]	15.37 [10.36]	0.0718 [0.67]
Trajectory	deg	9.77 [29.33]	13.53 [27.02]	0.1481 [0.17]
Vertical displacement (climb)	mm	0.37 [0.57]	0.53 [0.83]	0.0718 [0.35]
Vertical displacement (dive)	mm	-0.16 [0.23]	-0.21 [0.34]	**0.0473** [0.25]
Attack angle (climb)	deg	3.76 [6.92]	1.14 [6.59]	**0.0300** [0.52]
Attack angle (dive)	deg	-5.25 [5.37]	-6.27 [6.15]	0.1050 [0.25]
Steering rotation gain^		0.69 [0.02]	0.74 [0.02]	**0.0367** [1.58]
Righting rotation gain^		0.18 [0.01]	0.18 [0.02]	0.6281 [0.31]
Fraction of climbs	%	66 [4]	75 [7]	0.0718 [1.59]
Fraction of dives	%	32 [4]	23 [7]	0.0718 [1.76]

Table 3: Inter-bout interval vertical velocity by time post-treatment. Early and late represent data collected in the first and second 5 hours after the treatment, respectively, for each 24-hour behavior session using Dark-Dark dataset. P-value calculated using Wilcoxon rank sum ($\alpha = 0.05$).

Treatment	Time Post-Treatment	Median [IQR]	P-Value
Control	Early	-0.06 [0.29]	0.14164
	Late	0.02 [0.45]	
Copper	Early	-0.3 [0.20]	0.59081
	Late	-0.29 [0.41]	

Table 4: Number of inter-bout intervals and bouts across datasets. Percentages reflect median fraction and [IQR] of total bouts for that group and time point. P-value calculated using Mann-Whitney U test with $\alpha = 0.05$ for dark and light datasets, and using one-way ANOVA for longitudinal data.

Dataset	Condition	IBIs	Climb bouts	Flat bouts	Dive bouts	Total bouts	Climb % P-Value
Dark-Dark	control	31228	22537 (66 [4]%)	681 (2 [0.4]%)	10823 (32 [4]%)	34041	0.0718
	copper	14407	10811 (75 [7]%)	258 (2 [0.5]%)	3630 (23 [7]%)	14699	
Light-Dark	control	174820	81680 (46 [5]%)	5394 (3 [0.4]%)	95460 (51 [5]%)	182534	**0.0013**
	copper	120012	75311 (60 [13]%)	3165 (3 [0.4]%)	48602 (38 [13]%)	127078	
Longitudinal– Early Phase	control	21526	16150 (69 [12]%)	544 (2 [0.9]%)	7138 (29 [11]%)	23832	0.4717
	copperEarly	18504	12205 (68 [8]%)	371 (2 [1]%)	5826 (30 [7]%)	18402	
	copperLate	30536	21072 (62 [9]%)	811 (3 [1]%)	11159 (34%)	33042	
Longitudinal– Late Phase	control	17020	12149 (71 [7]%)	202 (1 [0.2]%)	4831 (27 [7]%)	17182	0.6314
	copperEarly	15902	11270 (68 [2]%)	264 (1 [1]%)	5093 (31 [2]%)	16627	
	copperLate	19892	12242 (71 [20]%)	231 (1 [0.9]%)	5683 (27 [19]%)	18156	

Table 5: Effects of vision on swimming after lateral line loss. Median [IQR] used for non-normal distributions of raw data. N = 150 LD control, 150 LD copper. P-value calculated using two-way ANOVA and Tukey's HSD post-hoc comparison (α = 0.05), and effect size calculated using Cohen's d. IBI, inter-bout interval

Parameter	Unit	LD Control	LD Copper	Effect of Light	Effect of Treatment	Interaction Effect
IBI Duration	s	0.69 [0.54]	0.63 [0.50]	**8.4e-13**	**9.6e-3**	**9.5e-3**
Vertical acceleration	mm s^{-2}	-0.04 [0.03]	-0.16 [0.08]	**0.0216**	**1.5e-5**	0.9176
Bout trajectory	deg	-1.29 [18.31]	4.28 [21.82]	**7.4e-8**	**1.17e-3**	0.4744

Table 6: Effects of lateral line loss on swimming in the light. Median [IQR] used for non-normal distributions of raw data. N = 150 LD control, 150 LD copper. P-value calculated using Mann-Whitney U test ($\alpha = 0.05$), and effect size calculated using Cohen's d. ˆSteering and righting rotation gain p-value calculated using unpaired t-test (mean and standard deviation listed).

Parameter	Unit	Control	Copper	P-Value [Effect Size]
Inter-bout interval				
Vertical displacement	mm	-0.08 [0.20]	-0.09 [0.21]	0.2474[0.08]
Vertical velocity	mm s^{-1}	-0.12 [0.19]	-0.21 [0.28]	**0.0090** [0.52]
Duration	s	0.69 [0.54]	0.63 [0.50]	0.3180 [0.15]
Pitch	deg	-0.37 [14.84]	4.96 [16.71]	**0.0019** [0.46]
Angular velocity	° s^{-1}	-4.07 [4.14]	-4.56 [5.50]	0.1860 [0.13]
Bout				
Peak speed	mm s^{-1}	14.22 [7.66]	13.43 [8.82]	0.2817 [0.13]
Trajectory	deg	-1.29 [18.31]	4.28 [21.82]	**0.0013** [0.38]
Vertical displacement (climb)	mm	0.20 [0.25]	0.21 [0.29]	**0.0415** [0.09]
Vertical displacement (dive)	mm	-0.22 [0.21]	-0.21 [0.24]	0.4790 [0.02]
Attack angle (climb)	deg	1.99 [4.33]	1.71 [4.72]	0.2817 [0.06]
Attack angle (dive)	deg	-2.45 [4.03]	-4.26 [5.09]	**0.0067** [0.41]
Steering rotation gainˆ		0.75 [0.03]	0.72 [0.02]	0.0915 [0.90]
Righting rotation gainˆ		0.09 [0.01]	0.10 [0.01]	**0.2557** [0.59]
Fraction of climbs	%	46 [5]	60 [13]	**0.0013** [1.73]
Fraction of dives	%	51 [5]	38 [13]	**0.0013** [1.65]

Table 7: Effects of hair cell regeneration after lateral line ablation. Median [IQR] used for non-normal distributions of raw data. N=85 control larvae, 94 larvae treated with copper in the early phase (7–8 dpf), and 64 larvae treated with copper in the late phase (14–15 dpf). Data from late phase of recordings. P-value calculated using one-way ANOVA with Tukey's HSD post-hoc comparison ($\alpha = 0.05$; bold numbers indicate significance) and effect size calculated using Cohen's d and enclosed in []. ˆSteering and righting gain p-value calculated using unpaired t-test (mean and standard deviation listed).

Parameter	Unit	Control	Copper Early	Copper Late	One-Way ANOVA P-Value	Control v. Early Post-Hoc	Early v. Late Post-Hoc	Control v. Late Post-Hoc
Inter-bout Interval								
Vertical displacement	mm	-0.05 [0.79]	-0.04 [0.70]	-0.35 [0.89]	**0.0048**	0.9000[0.01]	**0.0122** [0.51]	**0.0095** [0.47]
Vertical velocity	mm s^{-1}	-0.04 [0.31]	-0.03 [0.24]	-0.23 [0.46]	**0.0004**	0.9000 [0.03]	**0.0012** [0.68]	**0.0013** [0.63]
Duration	s	2.50 [3.69]	2.92 [3.49]	1.86 [2.48]	0.1832	0.7858 [0.18]	0.1671 [0.53]	0.4424 [0.32]
Pitch	deg	11.25 [21.51]	7.08 [18.09]	9.57 [19.11]	0.5480	0.5569 [0.28]	0.9000 [0.18]	0.6640 [0.11]
Angular velocity	° s^{-1}	-2.63 [4.77]	-3.02 [4.57]	-5.60 [7.46]	**0.0237**	0.9000 [0.11]	**0.0476** [0.55]	**0.0386** [0.63]
Bout								
Peak speed	mm s^{-1}	10.46 [6.56]	9.88 [5.89]	13.07 [9.91]	**0.0180**	0.3964 [0.13]	**0.0119** [0.55]	**0.0330** [0.43]
Trajectory	deg	17.82 [38.01]	12.52 [35.13]	14.50 [37.83]	0.6417	0.4374 [0.19]	0.2474 [0.07]	0.2474 [0.11]
Vertical displacement (climb)	mm	1.20 [0.98]	1.09 [0.90]	1.52 [1.07]	0.5238	0.3182 [0.15]	0.0067 [0.57]	0.0090 [0.41]
Vertical displacement (dive)	mm	-0.19 [0.39]	-0.14 [0.34]	-0.25 [0.46]	0.2605	0.3964 [0.19]	0.0946 [0.40]	0.0639 [0.21]
Attack angle (climb)	deg	9.77 [6.80]	10.68 [6.21]	7.55 [6.11]	0.0671	0.1860 [0.18]	0.0119 [0.70]	0.0639 [0.46]
Attack angle (dive)	deg	-5.37 [6.00]	-4.33 [6.46]	-5.78 [6.19]	0.3155	0.1860 [0.22]	0.0517 [0.31]	0.4790 [0.09]
Steering rotation gainˆ		0.67 [0.03]	0.67 [0.06]	0.67 [0.07]	0.9937	0.8980 [0.06]	0.9592 [0.02]	0.9584 [0.02]
Righting rotation gainˆ		0.16 [0.03]	0.16 [0.02]	0.18 [0.01]	0.4476	0.9026 [0.06]	0.1924 [0.68]	0.3251 [0.50]
Fraction of Climbs	%	71 [7]	68 [2]	71 [20]	0.6314	0.0946 [0.08]	0.2474 [0.08]	0.4790 [0.007]
Fraction of Dives	%	27 [7]	31 [2]	27 [19]	0.6440	0.0781 [0.08]	0.2474 [0.08]	0.4790 [0.007]

SUPPLEMENTAL VIDEO

Video S1

Video S1. Real-time recording of control and copper-treated larvae after treatment. Notably, copper-treated larvae sink between bouts, while controls maintain their position in depth. Related to Figure 2.