Neuropilin-1 controls vascular permeability through juxtacrine regulation of endothelial adherens junctions

Sagnik Pal, Yangyang Su, Lena Claesson-Welsh, and Mark Richards[*]

Dept. of Immunology, Genetics and Pathology, Beijer and Science for Life Laboratories, Uppsala University, Uppsala, Sweden

*) Corresponding author at the address above or mark.richards@igp.uu.se

Key words: Neuropilin 1, vascular permeability, signalling, VE-cadherin, VEGF

Abstract

Neuropilin-1 (NRP1) regulates endothelial cell (EC) biology through modulating vascular endothelial growth factor receptor 2 (VEGFR2) signalling by presenting VEGFA. How NRP1 impacts VEGFA-mediated vascular hyperpermeability however is unresolved, being described as having a positive or passive function. Using EC-specific *Nrp1* knock-out mice, we discover that EC-expressed NRP1 exerts an organotypic role. In ear skin, VEGFA/VEGFR2-mediated vascular leakage increased following EC NRP1 knock-out, showing that NRP1 negatively regulates VEGFR2 signalling. Conversely, in back skin and trachea, EC NRP1 knock-out decreased vascular leakage. Accordingly, VE-cadherin phosphorylation increased in the ear skin but was suppressed in back skin of *Nrp1* iECKO mice. NRP1 has been shown to have the ability to act in a juxtacrine manner. Importantly, NRP1 was more abundant in perivascular cells of the ear skin than back skin. Global NRP1 knock-out suppressed VEGFA-induced vascular leakage in the ear skin, implicating perivascular NRP1 as a juxtacrine co-receptor of VEGFA in this compartment. Altogether, we demonstrate that perivascular NRP1 is an active participant in EC VEGFA/VEGFR2 signalling and acts as an organotypic modifier of EC biology.

Introduction

Neuropilin 1 (NRP1) is a multifunctional transmembrane protein that is expressed abundantly on the surface of a range cell types, where it binds class 3 semaphorins (SEMA3), heparan sulfate and vascular endothelial growth factors (VEGFs) (Goshima et al., 1999, Mamluk et al., 2002, Gu et al., 2002). VEGFA is the main activator of VEGF receptor 2 (VEGFR2) signalling in endothelial cells (ECs) and its binding to NRP1 promotes heterocomplex formation between NRP1 and VEGFR2 (Soker et al., 2002). Thus, NRP1 has been designated as a co-receptor of VEGFR2 and is known to modulate VEGFR2 signalling. The cytoplasmic domain of NRP1 contains a C-terminal SEA motif, a PDZ binding domain that mediates binding to synectin (GIPC1) and in-turn regulates the endocytic trafficking of both NRP1 and VEGFR2 (Cai and Reed, 1999, Naccache et al., 2006, Horowitz and Seerapu, 2012, Lanahan et al., 2010). Accordingly, removal of the NRP1 cytoplasmic tail delays VEGFR2 endocytosis following VEGFA binding, leading to enhanced surface retention and reduced phosphorylation of tyrosine (Y)1175 in VEGFR2 (Lanahan et al., 2013). NRP1 thus acts as an important modulator of VEGFR2 activation and its downstream signalling pathways. However, reports have also suggested that VEGFA transduces biological responses in ECs via NRP1, in a VEGFR2-independent manner (Roth et al., 2016).

Global NRP1 knockout in mice results in lethality at E10-E12.5 due to abnormal vessel sprouting in major organs and impaired yolk sac vascularization (Kawasaki et al., 1999). Moreover, NRP1 overexpression leads to excessive vessel growth and promotes leaky and haemorrhagic vessels (Kitsukawa et al., 1995). Endothelial-specific knockout of NRP1 in mouse embryos only show mild embryonic brain defects, while in the adult vasculature no gross abnormalities are evident (Lanahan et al., 2013). Several reports also link increased NRP1 expression in tumours to a poor prognosis for survival (Miao et al., 2000, Kawakami et al., 2002, Hong et al., 2007). Furthermore, in diseases such as age related macular degeneration, NRP1 has been linked with increased neovascularization and vessel hyperpermeability, likely due to enhanced signalling downstream of VEGFA (Raimondi et al., 2014, Fernández-Robredo et al., 2017).

VEGFA-induced vascular permeability is mediated through the Y949 and Y1173 phosphosites of VEGFR2. The phosphorylated Y949 presents a binding site for TSAd (T-cell specific adaptor), which in turn binds SFKs (Src family kinases) that phosphorylate and promote internalization of junctional proteins such as VE-cadherin (Matsumoto et al., 2005, Sun et al., 2012, Li et al., 2016, Jin et al., 2022). Concurrently, pY1173 phosphorylates PLCϒ, leading to Ca2+/Protein kinase C (PKC) activation of endothelial nitric oxide synthase (eNOS), contributing to Src activation and phosphorylation of VE-cadherin (Sjoberg et al., 2023). The role of NRP1 in VEGFA-mediated permeability has been controversial. Studies employing *in* vivo and *in vitro* models have found NRP1 to be a positive regulator of VEGFA-mediated vascular permeability, or to lack effect, dependent on tissue or

experimental setup (Fantin et al., 2017, Acevedo et al., 2008, Wang et al., 2015, Becker et al., 2005,

Pan et al., 2007, Cerani et al., 2013). Additionally, treatment of ECs with CendR peptides, which bind

and induce internalisation of NRP1, increases permeability in a NRP1-dependent, but VEGFR2-

independent manner (Roth et al., 2016). Therefore, the exact role of NRP1 in VEGFA-VEGFR2

mediated permeability remains unclear. NRP1 is a promising target in a number of pathologies where

vascular dysfunction is exacerbative. Understanding the relationship between NRP1 and VEGFA-

VEGFR2 signalling is thus of potential therapeutic benefit.

Here, we aimed to further uncover the role of NRP1 in VEGFA-mediated vascular permeability. Using

mice with global or EC-specific loss of NRP1 expression, we identify a tissue-specific role for

endothelial NRP1 in the modulation of VEGFA/VEGFR2 signalling regulating vascular leakage. We find

that endothelial-specific loss of NRP1 (*Nrp1* iECKO) can both increase and decrease VEGFA-mediated

vascular leakage, dependent on the vascular bed. NRP1 is widely expressed and is an important

constituent of peri-vascular cells in some, but not all, tissues and vessel subtypes. The consequence

of global loss of NRP1 expression demonstrates that peri-vascular NRP1 can modify VEGFA/VEGFR2-

induced vascular leakage and steer the tissue-specific role of endothelial NRP1 in barrier integrity.

Collectively, these data reveal that the relative expression levels of NRP1 between perivascular cells

and ECs acts as a tissue-specific modifier of EC VEGFA/VEGFR2 signalling upstream of vascular

leakage.

Results

<u>NRP1 regulates VEGFA-mediated permeability in an organotypic manner</u>

NRP1 has been studied extensively as a co-receptor of the VEGFA/VEGFR2 pathway, and its ability to

modulate VEGFR2 signalling in developmental and pathological angiogenesis has been clearly

established. However, results concerning the role of NRP1 in VEGFA-mediated vascular permeability

have remained contradictory (Fantin et al., 2017, Acevedo et al., 2008, Wang et al., 2015, Becker et

al., 2005, Pan et al., 2007, Cerani et al., 2013). *Nrp1*$^{fl/fl}$; *Cdh5*^{CreERT2} (*Nrp1* iECKO) mice were used to

study the EC-specific role of NRP1 in VEGFA-induced vascular leakage. Following tamoxifen

administration, a 75% reduction in NRP1 protein levels was achieved in lung tissue, chosen for

analysis due to the high EC content (Figure 1A and B). Furthermore, using immunofluorescent

stainings, reduction in NRP1 levels could be observed in the ear and back skin vascular beds with the

loss of EC NRP1 (Supp. figure 1A and B). Using a highly sensitive assay combining intravital

microscopy and atraumatic intradermal microinjection in the ear skin of *Nrp1* iECKO mice, VEGFA-

induced vascular leakage was significantly increased following the loss of NRP1 (Figure 1C and D)

(Honkura et al., 2018). Additionally, we measured qualitative aspects of vascular leakage including lag

period (time from stimulation to leakage onset) and degree of leakage (rate of dextran

extravasation). These analyses showed that ECs in *Nrp1* iECKO mice respond faster to stimulation

(Figure 1E) and exhibit more profuse leakage at each site (Figure 1F), indicative of increased

disruption of EC-EC junctions. These data suggest that NRP1 exerts a stabilizing role by suppressing

VEGFA-induced endothelial junction disruption in the ear dermis. This finding is in contrast to reports

showing NRP1 to be a positive regulator of VEGFA-mediated vascular leakage (Fantin et al., 2017,

Acevedo et al., 2008, Becker et al., 2005).

One possible explanation for these differences compared to previous reports is that the role of NRP1

may be organ dependent, in line with the growing insights into the distinct properties of ECs in

different vessel types and vascular beds (Augustin and Koh, 2017). To investigate a potential

organotypic role for NRP1 in EC biology, we studied the consequence of *Nrp1* knockout on the barrier

properties of ECs in different vascular beds.

Initially, selected organs, chosen based on prior experience of their different EC barrier properties

(Aird, 2007, Augustin and Koh, 2017, Richards et al., 2022), were investigated for their susceptibility

to VEGFA-induced vascular leakage via the systemic administration of fluorescent dextrans with or

without VEGFA. After 30 minutes, organs were collected and acute vascular leakage assessed

microscopically or following solvent-based extraction of dextran and fluorescence spectroscopy. All

organs analysed, except for the liver, showed vascular leakage after VEGFA stimulation (Supp. figure

2A - E). The role of endothelial NRP1 in these organs was further studied in *Nrp1* iECKO mice,

showing significantly decreased VEGFA-mediated vascular leakage in the trachea and back skin

(Figure 1G and H, Supp. figure 3). Kidney, skeletal muscle and heart in contrast did not show a

significant difference in VEGFA-mediated vascular leakage following the loss of EC-expressed NRP1

(Figure 1I and J).

These data collectively suggest that endothelial NRP1 has an organ-specific role in controlling VEGFA-

mediated vascular permeability. Importantly, endothelial NRP1 is a positive regulator of VEGFA

mediated permeability in the trachea and back skin but a negative regulator in the ear skin.

<u>Global inactivation of NRP1 reduces VEGFA-mediated vascular permeability</u>

Our results here, where NRP1 may play positive (trachea and back skin), negative (ear skin) and

passive (kidney, skeletal muscle and heart) role in VEGFA-induced vascular leakage, were collected

using EC-specific knockout mice. In previous studies, mouse models of both EC-specific deletion and a

globally expressed NRP1 C-terminal deletion mutant were employed (Fantin et al., 2017, Roth et al.,

2016). We thus set out to investigate whether global inactivation of NRP1 using an anti-NRP1

blocking antibody might differently modify the VEGFA-induced leakage response. Wild-type C57Bl/6

mice were locally treated in the ear dermis with an anti-NRP1 antibody via intradermal injection,

followed by intravital assessment of VEGFA-induced vascular permeability. Interestingly, NRP1

blockade reduced vascular leakage in the ear dermis when compared to isotype control (Supp. figure

4A). This is in contrast to the EC-specific loss of NRP1 in *Nrp1* iECKO mice, which resulted in enhanced

VEGFA-mediated vascular leakage (Figure 1C-F). We similarly investigated the effect of blocking NRP1

on the back skin and tracheal vasculatures. Treatment with anti-NRP1 antibody was found to reduce

VEGFA-induced leakage in both tissues following local and systemic administration of VEGFA

respectively (Supp. figure 4B and C).

To verify the findings using neutralizing anti-NRP1 antibodies, we crossed *Nrp1*$^{fl/fl}$ with *Actb*^{CreERT} mice

to generate global *Nrp1* knock-out mice (*Nrp1* iKO). Efficient reduction of NRP1 protein was

confirmed in lung lysates (Figure 2A and B). Complete loss of NRP1 could also be seen in the ear and

back skin of these mice using immunofluorescent staining (Supp. figure 1). Intravital imaging of the

ear dermis of *Nrp1* iKO mice showed that global loss of NRP1 reduced VEGFA-induced vascular

leakage, in contrast to the phenotype seen in *Nrp1* iECKO mouse ear skin and in agreement with

blocking antibody data (Figure 2C and D). Furthermore, global *Nrp1* knock-out resulted in a ~75%

reduction in VEGFA-mediated vascular leakage also in the back skin and tracheal vasculatures, similar

to that seen in *Nrp1* iECKO mice (Figure 2E and F).

These data illustrate the heterogenous nature of EC signalling regulation in different vascular beds,

and show that NRP1 modulates VEGFA signaling in an organotypic and cell-specific manner. Notably,

we demonstrate that VEGFA-induced leakage in the back skin and trachea is indistinguishable

between endothelial-specific and global *Nrp1* KO mice, whereas in the ear skin there is a contrasting

phenotype.

<u>NRP1 is heterogeneously expressed in perivascular cells</u>

NRP1 is known to form a heterocomplex with VEGFR2 upon VEGFA binding. Interestingly,

VEGFR2/NRP1/VEGFA can assemble as a juxtacrine *trans* complex, where NRP1 and VEGFR2 are

expressed on the surface of adjacent cells, as well as a *cis* complex, where NRP1 and VEGFR2 are

expressed on the same cell (Koch et al., 2014). Importantly, even though the kinetics of *trans*

VEGFR2/NRP1 complex formation is slow, these complexes are stable and produce a distinct

signalling output compared to the *cis* configuration (Koch et al., 2014). Thus, NRP1 presented in *cis* or

trans produces differential VEGFR2 signalling output upon VEGFA binding. Given the above findings

we reasoned that peri-endothelial distribution of NRP1 could be an important modifier of EC VEGFR2

signalling and possibly explain organotypic differences observed in *Nrp1* iECKO mice.

To investigate the pattern of NRP1 expression we employed a *Pdgfrβ* promoter-driven GFP (*Pdgfrβ-*

GFP) mouse to visualize PDGFRβ-positive pericytes. To visualize ECs, and the localization of NRP1 in

the ear skin and back skin from these mice, tissues were immunostained for CD31 and NRP1,

respectively. NRP1 expression could be seen in both endothelial and perivascular cells, which was

lost after global *Nrp1* deletion (Figure 3A and Supp. Figure 1). Both arteriolar (Figure 3B) and capillary

(Figure 3C) PDGFRβ-positive perivascular cells expressed NRP1. The relative level of perivascular/EC

expression of NRP1 was higher in the ear skin compared to back skin. In venules, the low expression

of NRP1 in perivascular cells in the back skin resulted in a 5-fold difference in the perivascular/EC

NRP1 ratio, when comparing the ear skin to back skin (Figure 3D).

These data thus show that NRP1 is expressed in perivascular cells of the ear skin and back skin. We

find however that the ratio of NRP1 expression between ECs and perivascular cells differs between

ear skin and back skin, and between different vessel types. In ear skin the perivascular/EC ratio is

higher compared to the back skin, which may support a higher ratio of *trans* NRP1/VEGFR2 relative

to cis complexes in the ear skin.

<u>NRP1 distribution modifies VEGFA-mediated signalling and vascular leakage</u>

NRP1 modifies VEGFR2 signalling by controlling its internalisation and intracellular trafficking (Bayliss

et al., 2020, Ballmer-Hofer et al., 2011). The presence of NRP1 in *trans* however, modifies this

dynamic and retains VEGFR2 at the cell surface for longer time periods, altering its signalling output

(Koch et al., 2014).

We thus wished to investigate whether perivascular NRP1 expression might interact with and alter

VEGFR2 signalling upstream of vascular leakage, and explain the above described organotypic effects

of EC NRP1 loss. For this purpose, *Nrp1* iECKO mice were crossed with homozygous *Vegfr2$^{Y949F/Y949F}$*

mice, to produce *Vegfr2$^{Y949F/Y949F}$;Nrp1* iECKO mice that are deficient in both EC NRP1 expression and

VEGFA-mediated vascular leakage ability. In these mice, loss of NRP1 protein was efficiently

established in lung tissue (Figure 4A and B). Analysis of VEGFA-mediated vascular leakage in the ear

dermis showed that the increased leakage, induced by the loss of EC NRP1, was abrogated by the loss

of the VEGFR2 phosphosite Y949, known to be required for activation of Src in response to VEGFA

(Figure 4C and D). These analyses suggest that, in the ear dermis, EC NRP1 negatively regulates

VEGFA-induced vascular leakage by modulation of VEGFR2 activation and phosphorylation.

We next sought to determine the relationship between NRP1 and VEGFR2 Y949 signalling in the

trachea and back skin vasculatures. Leakage in $Vegfr2^{Y949F/Y949F}$;$Nrp1$ iECKO mice was induced by

systemic VEGFA and fluorescent 2000 kDa dextran administration in both $Nrp1$ iECKO and

$Vegfr2^{Y949F/Y949F}$;$Nrp1$ iECKO mice. As we observed previously, loss of EC NRP1 reduced vascular

leakage in the trachea and back skin by 75%, an effect that was enhanced further by the VEGFR2

Y949F mutation (Figure 4E and F). Importantly, leakage was similarly inhibited in the $Vegfr2^{Y949F/Y949F}$

model as in the combined $Vegfr2^{Y949F/Y949F}$;$Nrp1$ iECKO mice in both the trachea and back skin. These

data show that NRP1 presented in *trans* by perivascular cells in the ear dermis can signal through

VEGFR2 to induce vascular leakage, and that VEGFR2 Y949 is vital for mediating NRP1's effects on

endothelial junctions.

<u>Perivascular NRP1 modifies VEGFA-induced signalling</u>

VEGFA/VEGFR2 signalling mediates phosphorylation of VE-Cadherin on Y685, required for vascular

permeability (Orsenigo et al., 2012, Wessel et al., 2014). We have also recently shown that PLCγ is an

important mediator of VEGFA-induced vascular leakage by allowing the production of nitric oxide

through eNOS, which in turn modified Src by tyrosine nitration, required for full Src activity (Sjoberg

et al., 2023). To clarify the impact of perivascular NRP1 on VEGFA-induced signalling we studied VE-

Cadherin and PLCγ phosphorylation in $Nrp1$ iECKO and $Nrp1$ iKO mice (Figure 5). Perivascular NRP1

expression in the ear dermis of $Nrp1$ iECKO mice was lost in $Nrp1$ iKO mice (Supp. figure 1). In

agreement with the enhanced vascular leakage in the ear dermis of $Nrp1$ iECKO mice, VE-Cadherin

phosphorylation at Y685 was enhanced following intradermal VEGFA injection (Figure 5A). No such

induction was seen in $Nrp1$ iKO mice (Figure 5B). In the back skin, VE-Cadherin pY685 levels were

induced by VEGFA in the control but not in the $Nrp1$ iECKO, nor the iKO models (Figure 5C and D), in

accordance with the permeability phenotype (see Figure 1). Similarly, PLCγ phosphorylation was

potentiated in the ear skin of $Nrp1$ iECKO but not $Nrp1$ iKO mice (Supp. figure 5A and B). Moreover,

in the back skin loss of either endothelial or global NRP1 led to a reduction in PLCγ phosphorylation

following intradermal VEGFA stimulation (Supp. figure 5C and D).

Taken together these data show that NRP1 is expressed by perivascular cells in a tissue-specific

manner and that perivascular NRP1 can be an important modulator of VEGFA-signalling. In the ear

dermis, NRP1 is expressed by both endothelial and perivascular cells, forming *cis* and *trans*

interactions with VEGFR2 respectively. Whilst the *cis* EC interaction predominates, loss of EC NRP1

promotes *trans* NRP1/VEGFR2 interaction and potentiates VEGFA-induced signalling regulating

vascular leakage (Figure 6). Meanwhile, due to a paucity of perivascular NRP1 in the trachea and back

skin vasculatures, loss of EC NRP1 leads to a loss of NRP1:VEGFR2 complexes, attenuating VEGFA-

regulated vascular leakage.

<u>Discussion</u>

Here we find that NRP1 expression by perivascular cells is an important modulator of

VEGFA signalling regulating endothelial junction stability. Our data suggest that NRP1 presented in a

juxtacrine manner can potentiate certain VEGFR2 signalling upstream of vascular leakage. NRP1 is

expressed by a number of cells adjacent to blood vessels, including PDGFRβ-positive pericytes.

However, the abundance of these cells and their level of NRP1 expression differ between vascular

beds. Loss of EC-specific NRP1 thus has a differential effect on VEGFA-mediated vascular leakage in

different vascular beds, dependent on the ability of ECs to form *tran*s NRP1/VEGFR2 complexes.

Numerous studies have implicated endothelial NRP1 as a positive regulator of

permeability (Fantin et al., 2017, Becker et al., 2005, Acevedo et al., 2008). However, multiple studies

argue against this, with NRP1 blockade or deletion having no effect on VEGFA-induced vascular

leakage in the skin and retina (Pan et al., 2007, Cerani et al., 2013). Here we find that the function of

endothelial NRP1 in VEGFA-mediated vascular permeability varies in different vascular beds. In

keeping with previous data, we find that loss of NRP1 reduces VEGFA-induced vascular leakage in the

back skin and trachea. Meanwhile, endothelial NRP1 appears to have no effect on VEGFA-induced

vascular leakage in kidney, skeletal muscle and heart. In the ear skin, however, loss of NRP1

surprisingly leads to an increase in VEGFA-induced vascular leakage. This appears to be through

enhanced VEGFR2 Y949 signalling, as mutation of this phosphosite normalises the enhanced leakage

seen after loss of NRP1. Previous publications have described NRP1 as being pivotal for VEGFR2-

induced Src activity via recruitment of Abl kinase (Fantin et al., 2017). How exactly VEGFA-VEGFR2

signals to Abl, and whether this requires the Y949 phosphosite, however is unknown. Further work is

required to understand exactly how NRP1 modifies VEGFR2 activity and how this differs between

different vascular beds.

As others have noted and we see here, NRP1 is expressed by perivascular cells,

including PDGFRβ positive pericytes (Bartlett et al., 2017, Wnuk et al., 2017). However, we find that

perivascular NRP1 expression and its ratio to EC expression differs between different vascular beds,

being higher in the ear skin than back skin. When the *trans* NRP1/VEGFR2 complex predominates, as

in the ear skin of *Nrp1* iECKO mice, our data indicate that VEGFA/VEGFR2 signalling impacting

junction stability is potentiated. This is in keeping with reports showing that NRP1 presented in *trans* can modify VEGFR2 signalling kinetics, resulting in prolonged PLCγ and ERK2 phosphorylation (Koch et al., 2014). However how such prolonged kinetics might affect the VEGFR2 signalling pathways upstream of vascular leakage has not been previously explored. In addition, juxtacrine NRP1 interactions have been shown to have important functions in cell types other than ECs. For example, NRP1 expressed on microglia is important for the trans-activation of PDGFRα on oligodendrocyte precursor cells, resulting in enhanced proliferation and myelin repair (Sherafat et al., 2021). Furthermore, NRP1 has been posited as a potential immune checkpoint inhibitor (Sarris et al., 2008). NRP1 is highly expressed by regulatory T cells, as well as antigen presenting dendritic cells, and the formation of *trans* NRP1 homodimers has been suggested to stabilise regulatory T cell function and promote immune tolerance. Finally, numerous studies have described the expression of NRP1 on tumour cells, which can modify EC behaviour and alter their angiogenic potential (Koch et al., 2014, Miao et al., 2000, Hu et al., 2007, Shahrabi-Farahani et al., 2016). This may partly be due to the ability of NRP1 to bind VEGFA and thus provide a pool of tumoural VEGFA towards which ECs migrate. In tumours, NRP1 can also form juxtacrine interactions with VEGFR2 on ECs upon VEFGA binding and modify intracellular EC signalling compared to NRP1 action in a *cis* manner (Koch et al., 2014, Morin et al., 2020, Morin et al., 2018).

One possible mechanism by which NRP1 presented in *trans* might alter VEGFA/VEGFR2 signalling, when compared to *cis*, is via modification of VEGFR2 trafficking. The adapter molecule synectin is an important intracellular binding partner of NRP1 and facilitates myosin IV-mediated endocytosis (Cai and Reed, 1999, Naccache et al., 2006). A recent study also found that NRP1 interacts with members of the ERM family of intracellular proteins, which act as intermediaries between the plasma membrane and actin cytoskeleton and that have also been associated with endocytosis and intracellular trafficking (Ponuwei, 2016, Gioelli et al., 2022). NRP1 may thus function not just as a co-receptor, but also as a linker to mediate the intracellular trafficking of cell surface receptors. This idea is supported by studies showing that NRP1 co-ordinates PDGFRα internalisation and its intracellular signalling, as well as the trafficking of integrin α5β1 during cell adhesion (McGowan and McCoy, 2021). Furthermore, it has recently been shown that NRP1 binds to VE-Cadherin and regulates its cell surface turnover, a role that has been proposed to at least partly explain its role in EC barrier integrity (Gioelli et al., 2022, Bosseboeuf et al., 2023). In response to VEGFA the cytoplasmic domain of NRP1, which binds synectin, has been found to be crucial in mediating NRP1's positive regulation of leakage (Fantin et al., 2017). Furthermore, mutant mice lacking the NRP1 cytoplasmic domain have impaired arteriogenesis due to delayed VEGFR2 endocytosis and resulting reduction in phosphorylation at Y1175 (Lanahan et al., 2013). Interestingly,

the C-terminal tail is also crucial for the stability of trans NRP1-VEGFR2 complexes (Koch et al., 2014).

A central function of NRP1 is thus to mediate VEGFR2 internalisation and trafficking following VEGFA

stimulation. From our data we hypothesise that NRP1 presented in *trans* delays VEGFR2

internalisation and potentiates VEGFR2 signalling upstream of vascular leakage. Indeed, a recent

study found that proper internalisation of VEGFR2 is required to shutdown VEGFR2 Y949 signalling

through increasing the stoichiometric ratio of cell surface phosphatases that dephosphorylate Y949

(Corti et al., 2022). Our data are compatible with the hypothesis that VEGFR2 held at the cell surface

via juxtracrine NRP1 interaction is capable of enhancing VEGFR2 Y949 signalling and resulting

disruption of EC junctions, but due to the rapid turnover of VEGFR2 phosphorylation, this is

technically very challenging to conclusively demonstrate. We hypothesize however that NRP1

expression within the perivascular microenvironment could be an important modifying factor of EC

signalling and function, and might play an important role in mural cell function during vascular

development and quiescence.

Combined, we identify a differential function of endothelial NRP1 in VEGFA-mediated

vascular leakage in different vascular beds. Different levels and patterning of perivascular NRP1

expression has the potential to modify VEGFR2 signalling and alter VEGFA-induced vascular

hyperpermeability. Further elucidation of NRP1 expression patterns in different tissues and how the

VEGFA/NRP1/VEGFR2 complex presented in *cis* or *trans* can differentially modify cellular signalling

will allow to better understand how NRP1 may serve in therapeutic applications in a distinct manner

in different tissues.

<u>Acknowledgements</u>

We thank the BioVis imaging facility, Uppsala University, for expert services. We appreciate expert

advice from Dr. Konstantin Gängel. This study was supported by the Swedish Research Council (2022-

00896), the Knut and Alice Wallenberg foundation (KAW 2020.0057 and KAW 2019.0276), Fondation

Leducq Transatlantic Network of Excellence Grant in Neurovascular Disease (17 CVD 03) to LCW. MR

was supported by Olle Engkvist (218-0057) SSMF (201912) and an EMBO long-term fellowship (ALTF

923–2016).

Author contributions

Conceptualization, M.R. and L.C.-W.; methodology, M.R. and S.P.; investigation, M.R., S.P., Y.S.; writing, M.R., S.P. and L.C.-W.; funding acquisition; M.R. and L.C.-W.; resources, L.C.-W.

Conflict of interest

M.R. is now an employee for AstraZeneca (Biopharmaceuticals R&D, Cambridge, UK). All of this work was performed at Dept. of Immunology, Genetics and Pathology, Beijer and Science for Life Laboratories, Uppsala University, Uppsala, Sweden. No funding or support was received from AstraZeneca. Other authors declared no conflict of interest.

Figure legends

Figure 1: NRP1 regulates VEGFA-mediated permeability in an organotypic manner

(A) Schematic illustration of experimental design to induce recombination in *Nrp1* iECKO mice.

(B) Western blot quantification of lung lysates from tamoxifen-treated control and *Nrp1* iECKO mice (n=4).

(C) Representative images showing leakage of 2000 kDa FITC-dextran (Pseudo-colour) in response to intradermal VEGFA injection in the ear dermis of control and *Nrp1* iECKO mice.

(D) Leakage sites per vessel length in response to intradermal VEGFA stimulation in the ear skin of control and *Nrp1* iECKO mice. n=6 mice, two or more acquisitions/mouse.

(E) Lag period between intradermal VEGFA injection and initiation of leakage in the ear skin of control and *Nrp1* iECKO mice. n≥3 mice, two or more acquisitions/mouse, three or more sites/acquisition.

(F) Quantification of 2000 kDa dextran extravasation over time in the ear skin of control and *Nrp1* iECKO mice following intradermal VEGFA stimulation. Black lines represent lines of best fit for the slope between leakage initiation and leakage termination. n≥3 mice, two or more acquisitions/mouse, three or more sites/acquisition.

(G-H) Leakage of fixable 2000 kDa FITC dextran in trachea (G) and back skin (H) after systemic administration of VEGFA in control and *Nrp1* iECKO mice. Left, representative images. Right, quantification of tracer leakage area / vessel area (n ≥ 8 mice, 2 or more fields of view/mouse).

(I-J) Leakage of 70 kDa (I) and 2000 kDa (J) dextran extracted from perfused tissues after systemic administration of VEGFA in kidney, skeletal muscle and heart in control and *Nrp1* iECKO mice (n ≥ 8 mice).

Error bars; mean ± SEM. Statistical significance: Two-tailed unpaired student's t-test and linear regression with ANCOVA. Scale bar: 100 μm

Figure 2: Global loss of NRP1 reduces VEGFA-mediated vascular permeability

(A) Schematic illustration of experimental design to induce recombination in *Nrp1* iKO mice.

(B) Western blot quantification of lung lysates from tamoxifen-treated control and *Nrp1* iKO mice (n ≥ 2 mice).

(C) Representative images showing leakage of 2000 kDa FITC dextran (Pseudo-colour) in response to intradermal VEGFA injection in the ear dermis of control and *Nrp1* iKO mice.

(D) Leakage sites per vessel length in response to intradermal VEGFA stimulation in the ear skin of control and *Nrp1* iKO mice. n=3 mice, two or more acquisitions/mouse.

(E-F) Leakage of fixable 2000 kDa FITC dextran in trachea (E) and back skin (F) after systemic administration of VEGFA in control and *Nrp1* iKO mice. Left, representative images. Right, quantification of tracer leakage area / vessel area (n ≥ 5 mice, 3 or more fields of view/mouse).

Error bars; mean ± SEM. Statistical significance: Two-tailed unpaired student's t-test. Scale bar: 100μm

Figure 3: Heterogeneous expression of perivascular NRP1

(A) Schematic illusatration of mouse strain and representation of NRP1 expression in endothelial, and perivascular cells using *Pdgfrβ*-GFP mice. White dashed line encompasses vascular area, space between white and yellow dashed lines represents perivascular area. Note perivascular, NRP1 expressing cells in between yellow and white lines.

(B-D) Images, left, and quantification, right, showing vascular and perivascular NRP1 expression in arterioles **(A)**, capillaries **(B)** and venules **(C)** of ear skin and back skin and its relative expression.

Error bars; mean ± SEM. Statistical significance: Two-tailed unpaired student's t-test. Scale bar: 50μm

Figure 4: NRP1 distribution modifies VEGFA-induced vascular permeability

(A) Schematic illustration of breeding scheme to obtain homozygous *Vegfr2* $^{Y949F/Y949F}$; *Nrp1* iECKO mice and experimental design to induce recombination.

(B) Western blot quantification of lung lysates from tamoxifen treated *Vegfr2* $^{Y949F/Y949F}$ and *Vegfr2* $^{Y949F/Y949F}$; *Nrp1* iECKO mice (n=3 mice).

(C) Representative images showing leakage in response to intradermal VEGFA injection in the ear dermis of *Vegfr2* $^{Y949F/Y949F}$ and *Vegfr2* $^{Y949F/Y949F}$; *Nrp1* iECKO mice.

(D) Leakage sites per vessel length in response to intradermal VEGFA stimulation in the ear skin of in control, *Nrp1* iECKO, *Vegfr2* $^{Y949F/Y949F}$ and *Vegfr2* $^{Y949F/Y949F}$; *Nrp1* iECKO mice, normalized to control (n ≥ 6 mice, two or more acquisitions/mouse.

(E-F) Leakage of 2000 kDa dextran in trachea (E) and back skin (F) of control, *Nrp1* iECKO, *Vegfr2* $^{Y949F/Y949F}$ and *Vegfr2* $^{Y949F/Y949F}$; *Nrp1* iECKO mice after systemic administration of VEGFA. Left, representative images. Right, quantification of tracer leakage area/vessel area (n ≥ 3 mice, 2 or more fields of view/mouse).

Error bars; mean ± SEM. Statistical significance: Two tailed unpaired student's t-test and Two-way ANOVA. Scale bar: 100μm

Figure 5: Perivascular NRP1 modifies VEGFA-induced signalling

(A-B) Phosphorylation of VE-Cadherin (VEC) Y685 in response to intradermal PBS or VEGFA injections in the ear dermis of control and *Nrp1* iECKO (A) or *Nrp1* iKO (B) mice. Left, representative images.

Right, quantification of phosphorylated VEC Y685 area per total VEC area normalized to PBS control
in the ear dermis of control and *Nrp1* iECKO or *Nrp1* iKO mice. n≥3 mice, two or more fields of
view/mouse.

(C-D) Phosphorylation of VE-Cadherin Y685 in response to intradermal PBS or VEGFA injections in the
back skin of control and *Nrp1* iECKO (C) or *Nrp1* iKO (D) mice. Left, representative images. Right,
quantification of phosphorylated VEC Y685 area per total VEC area normalized to PBS control in the
back skin of control and *Nrp1* iECKO or *Nrp1* iKO mice. n≥3 mice, two or more fields of view/mouse.

Error bars; mean ± SEM. Statistical significance: Two-way ANOVA. Scale bar: 50μm

Figure 6: Schematic model of NRP1's spatial effects on VEGFA-induced vascular permeability

With global loss of NRP1, VEGFA/VEGFR2 signalling regulating vascular leakage is suppressed and
thus leakage is reduced (left). In the presence of EC NRP1, a VEGFA/VEGFR2/NRP1 complex is formed
in *cis,* which has a rapid and transient effect on VEGFR2 downstream signalling (middle). Loss of EC
NRP1 but the presence of perivascular NRP1 leads to formation of a stable VEGFA/VEGFR2/NRP1
complex in *trans* and enhanced VEGFR2 signalling and increased vascular permeability.

Supplementary Figure 1. NRP1 expression in the ear dermis and back skin of adult mice

(A-B) Whole mount immunostainings showing NRP1 expression in ear dermis (A) and back skin (B) of
control, *Nrp1* iECKO and *Nrp1* iKO mice. Scale bar: 50μm

Supplementary Figure 2. Organotypic assessment of VEGFA-induced leakage

(A,B) Leakage of 10 kDa **(A)** and 70 kDa **(B)** dextran after systemic administration of only dextran
(control) and dextran plus VEGFA in liver, kidney, skeletal muscle and heart of wild-type C57Bl/6 mice
(n ≥ 3 mice).

(C-E) Leakage of 2000 kDa dextran after systemic administration of only dextran (control) and
dextran plus VEGFA in ear dermis **(C)**, back skin **(D)** and trachea **(E)** of wild-type C57Bl/6 mice (n ≥ 2
mice).

Error bars; mean ± SEM. Statistical significance used: Two-tailed unpaired student's t-test

Supplementary Figure 3. Loss of EC NRP1 reduces VEGFA mediated vascular permeability in the
back skin

Evans blue leakage with intradermal administration of VEGFA in the back skin of *Nrp1* iECKO mice.
Left, representative images. Right, quantification of Evans blue extravasation shown as VEGFA
induced leakage fold over PBS (n ≥ 3 mice).

Error bars; mean ± SEM. Statistical significance used: Two-tailed unpaired student's t-tes

Supplementary figure 4. Global inactivation of NRP1 reduces VEGFA mediated vascular
permeability

(A) VEGFA-induced leakage in the ear skin of C57Bl/6 mice treated intradermally with isotype control
or NRP1 blocking antibody. Left, representative images. Right, quantification of leakage sites per 100
μm of vessel length (n=3 mice, 3 acquisitions/mouse).

(B) Leakage of 2000 kDa dextran in trachea of C57Bl/6 mice treated systemically with isotype control
or NRP1 blocking antibody. Left, representative images. Right, quantification of tracer area/vessel
area (n=3 mice).

(C) PBS and VEGFA-induced leakage in the back skin of C57Bl/6 mice treated intradermally with isotype control or NRP1 blocking antibody. Left, representative images. Right, quantification of Evans blue leakage (n=3 mice).

Error bars; mean ± SEM. Statistical significance used: Two-tailed unpaired student's t-test. Scale bar: 100µm

Supplementary figure 5. Perivascular NRP1 modifies PLCγ phosphorylation

(A-B) Phosphorylation of PLCγ in response to intradermal PBS or VEGFA injections in the ear dermis of control and *Nrp1* iECKO (A) or *Nrp1* iKO (B) mice. Left, representative images. Right, quantification of phosphorylated PLCγ area per total VEC area normalized to PBS control in the ear dermis of control and *Nrp1* iECKO or *Nrp1* iKO mice. n≥3 mice, two or more fields of view/mouse.

(C-D) Phosphorylation of PLCγ in response to intradermal PBS or VEGFA injections in the back skin of control and *Nrp1* iECKO (C) or *Nrp1* iKO (D) mice. Left, representative images. Right, quantification of phosphorylated PLCγ area per total VEC area normalized to PBS control in the back skin of control and *Nrp1* iECKO or *Nrp1* iKO mice. n≥3 mice, two or more fields of view/mouse.

Error bars; mean ± SEM. Statistical significance: Two-way ANOVA. Scale bar: 50µm

Methods

Animals

In vivo animal experiments were carried out in accordance with the ethical permit provided by the Committee on the Ethics of Animal Experiments of the University of Uppsala (permit 6789/18). *Nrp1*$^{flox/flox}$ mice contain *loxP* sites flanking exon 2 and were obtained from The Jackson Laboratory (005247). This strain was crossed with Cdh5(PAC)-CreER^{T2} mice (kind gift from Dr. Ralf Adams, Max-Planck Institute Münster) or Tmem163Tg(ACTB-cre)2Mrt mice to generate endothelial-specific *Nrp1* iECKO mice and global *Nrp1* iKO mice respectively. Mice possessing a knock-in mutation in the VEGFR gene, *Kdr*, at the Y949 phosphosite have been described previously (Li et al., 2016). These mice were crossed with *Nrp1* iECKO mice to generate *Vegfr2* $^{Y949F/Y949F}$; *Nrp1* iECKO mice. *Pdgfrb*-GFP reporter mice (Tg(Pdgfrb-EGFP)jn169Gsat/Mmucd, Stock number 031796-UCD) have been described previously (Jung et al., 2018). Mice were maintained in ventilated cages with group housing (2–5 per cage) and access to water and feed ad libitum. Male and female mice age 8-18 weeks were used. Each experiment was conducted on tissue from at least three age-matched animals representing individual biological repeats. Sample size (number of acquired images / movies and number of mice) were chosen to ensure reproducibility and allow stringent statistical analysis. To induce Cre recombinase-mediated gene recombination tamoxifen (SigmaAldrich, T5648) formulated in peanut oil was injected intraperitoneally (80 mg/kg) for 5 consecutive days. The mice were allowed to rest for 4 days before experiments were conducted.

<u>Intravital vascular leakage assay</u>

Intravital imaging of the mouse ear with intradermal injection has been described previously

(Honkura et al., 2018). Briefly, following systemic administration of 2000 kDa FITC (SigmaAldrich,

FD2000S) or TRITC Dextran (ThermoFischer Scientific, D7139) by tail-vein injection, mice were

sedated by <u>intraperitoneal injection</u> of Ketamine-Xylazine (120 mg/kg Ketamine, 10 mg/kg Xylazine)

and the ear secured to a solid support. Mice were maintained at a body temperature of 37 °C for the

entire experiment, maximum 90 min. Time-lapse imaging was performed using single-photon

microscopy (Leica SP8). For intradermal EC stimulation, a volume of approximately 0.1 µl VEGFA

(Peprotech, 450-32), concentration 100 ng/µl, was injected using a sub-micrometer capillary needle.

10 kDa TRITC Dextran (ThermoFischer Scientific, D1817) was used as a tracer. Leakage sites were

identified in time-lapse imaging as defined sites of concentrated dextran in the extravascular space.

<u>Miles' assay</u>

Mice were injected intraperitoneally with pyrilamine maleate salt (4mg/kg, Sigma, P5514) diluted in

0.9% saline, to inhibit histamine release, 30 min prior to tail vein injection of Evans blue (100 µl, 1%

Evans blue). Evans blue was allowed to circulate for 10 min before intradermal injection in the back

skin of recombinant mouse VEGFA165 (Peprotech, 450-32) (80 ng in 25 µl) or PBS. The skin was

excised 30 minutes subsequent to VEGFA injection, placed in formamide over night at 55°C and

absorbance at 620nm was measured using a spectrophotometer and normalized to tissue weight.

<u>Systemic VEGFA-induced permeability</u>

To assess EC permeability, all tissues were cleaned of excess cartilage, fat and connective tissue.

Skeletal muscle was assessed using the tibialis anterior and liver was assessed using the left lateral

lobe. To analyse liver, kidney, skeletal muscle and heart permeability, dextran (30 mg/kg) with or

without VEGFA (160 µg/kg) were injected systemically through the tail vein. Thirty minutes later mice

were anesthetized with Ketamine/Xylazine before intracardiac perfusion with Dulbecco's phosphate

buffered saline (DPBS). Tissues were dissected, washed in DPBS and incubated in formamide for 48

hr at 55 °C. Dextran fluorescence was then measured using a spectrophotometer and normalized to

tissue weight. To assess VEGFA-induced leakage microscopically, mixtures of fixable dextran (FITC,

Tdb labs; TRITC, ThermoFischer Scientific, D7139) (30 mg/kg) with VEGFA (160 µg/kg) were injected

systemically through the tail vein. Thirty minutes later mice were anesthetized with

Ketamine/Xylazine before intracardiac perfusion with DPBS followed by 1% paraformaldehyde.

Tissues were then immersed in 1% paraformaldehyde for 2 hr before proceeding with

immunohistochemistry. For leakage quantification at least three large tile scan areas ($\geq$1 mm^2) were

captured for each mouse<u>.</u>

<u>Intradermal VEGFA injection in the ear and back skin</u>

Mice were anesthetised using isofluorane and injected with PBS or VEGFA (Peprotech, 450-32) (20ug

in 5 ul for the ear dermis and 80ug in 20ul for the back skin) followed by perfusion using 1% PFA. The

ear and back skin samples were excised and further fixed for 1 hour in 1% PFA. Afterwards, the tissue

samples were processed for immunostaining.

<u>NRP1 blockade</u>

Anti-NRP1 (R&D systems, MAB59941) or isotype control (R&D Systems, MAB006) were administered

intradermally in the ear skin (1 µg) or back skin (5 µg), or systemically (2 mg/kg). 24 hours later

VEGFA-mediated vascular leakage was assessed using the intravital vascular leakage assay, Miles'

assay or through systemic VEGFA-induced permeability.

<u>Immunohistochemistry</u>

Tissues were fixed through intracardiac perfusion with 1% paraformaldehyde (PFA) followed by

immersion in 1% PFA for 2 hr at room temperature. For whole mount preparation of the ear skin,

back skin and trachea removal of excess cartilage, fat and connective tissue tissues was carried out.

Tissues were blocked overnight at 4 °C in Tris-buffered saline (TBS), 5% (w/v) Bovine Serum Albumin

(BSA), 0.2% Triton X-100. Samples were incubated overnight with primary antibody in blocking

solution, followed by washing in TBS, 0.2% Triton X-100 and incubation with appropriate secondary

antibody for 2 hr at room temperature in blocking buffer before washing and mounting in

fluorescent mounting medium (DAKO). Images were acquired using a Leica SP8 confocal microscope.

Commercial antibodies used were: rat anti-CD31 (BD Biosciences, 553370), goat anti-CD31 (R&D

Systems, AF3628), goat anti-NRP1 (R&D Systems, AF566), chicken anti-GFP (Abcam, Ab13970), rabbit

anti-phospho-PLCγ1 (Tyr783) (Cell signaling technology, 2821s), rabbit anti-PLCγ1 (Cell signaling

technology, 2822s), and goat anti–VE-cadherin (R&D systems, AF1002). Secondary antibodies against

rat (ThermoFischer Scientific; Alexa 488, A21208 and Alexa 594, A21209), rabbit (ThermoFischer

Scientific; Alexa 488, A21206 and Alexa 568, A10042), goat (ImmunoResearch Laboratories, Alexa

647, 705-605-147), chicken (ImmunoResearch Laboratories, Alexa 488, 703-545-155) were used.

pVEC Y685 antibody was prepared by immunizing rabbits with phospho-peptides of the

corresponding region in mouse VE-cadherin (New England Peptide) and further verified using

immunostaining in Cdh5$^{Y685F/Y685F}$ mice. Primary and secondary antibodies were prepared at a dilution

of 1:100 and 1:400, respectively unless otherwise stated.

<u>Western blot analysis</u>

Lungs from mice were removed and snap frozen. Protein was obtained by mechanical dissociation in RIPA buffer supplemented with 50 nM Na3VO4, Phosphatase inhibitor cocktail (Roche 04906837001) and Protease inhibitor cocktail (Roche, 04693116001). LDS sample buffer (Invitrogen, NP0007) and Sample Reducing Agent (Invitrogen, NP0009) were added to the samples and heated to 70 °C for 10 min. Proteins were separated on Nu Page 4–12% Bis-Tris Gel (Invitrogen) in MOPS SDS Running buffer (Invitrogen, NP0001), transferred to PVDF membrane (Thermo scientific, 88518) in NuPAGE transfer buffer (Novex, NP006), 10% methanol and subsequently blocked with 5% BSA in Tris-buffered saline with Tween 20 (TBST) for 60 min. The immunoblots were analysed using primary antibodies incubated overnight at 4 °C and secondary antibodies linked to horseradish peroxidase (HRP) (Cytiva) incubated for 1 hr at room temperature. After each step filters were washed four times with TBST. HRP signals were visualized by enhanced chemiluminescence (ECL) (Cytiva) (1:25000) and imaged with Chemidoc (Bio-Rad).

Primary antibodies targeting GAPDH (Chemicon, MONOCLONAL ANTIBODY374), NRP1 (R&D Systems, AF566) were used at a dilution of 1:1,000.

<u>Image quantification</u>

For confocal images, macromolecular leakage was quantified by measurement of tracer area following image thresholding. Tracer area was then normalized to vessel area. For analysis of vascular and peri-vascular NRP1 expression, masks were generated of the CD31 vascular area and the peri-vascular area 3µm around the vessel (Figure 3A). NRP1 area following thresholding was quantified within, and normalised to these areas. Phosphorylation values were quantified by measurement of phosphorylated vessel area following image thresholding. Phosphorylation area was then normalized to vessel area. Threshold values were kept constant within experiments.

Analysis of leakage from intravital movies has been described previously (Richards et al., 2021). Briefly, leakage sites were identified in time-lapse imaging as defined sites of concentrated dextran in the extravascular space. To quantify these their numbers were normalized to vessel length. To assess lag period the time of the appearance of these sites following injection of stimulus was quantified. To assess the extent of barrier disruption the rate of dextran extravasation specifically at these sites was quantified over time.

All measurements were done with Fiji processing package of Image J2 software.

<u>Statistical analysis</u>

Data are expressed as mean ± SEM. The statistical tests used were the Students't test and two-way

ANOVA. P-values given are from independent samples analysed by two-tailed paired t tests or

multiple comparisons obtained from two-way ANOVA. Rate of leakage was compared using linear

regression and ANCOVA. All statistical analyses were conducted using GraphPad Prism. A p-value

<0.05 was considered statistically significant and significances indicated as $p<0.05$ (*), $p<0.01$ (**),

and $p<0.001$ (***). For animal experiments, no statistical methods were used to predetermine

sample size. The investigators were blinded to allocation during experiment and outcome

assessment.

<u>References</u>

ACEVEDO, L. M., BARILLAS, S., WEIS, S. M., GÖTHERT, J. R. & CHERESH, D. A. 2008. Semaphorin 3A
suppresses VEGF-mediated angiogenesis yet acts as a vascular permeability factor. *Blood,*
111**,** 2674-80.
AIRD, W. C. 2007. Phenotypic heterogeneity of the endothelium: I. Structure, function, and
mechanisms. *Circ Res,* 100**,** 158-73.
AUGUSTIN, H. G. & KOH, G. Y. 2017. Organotypic vasculature: From descriptive heterogeneity to
functional pathophysiology. *Science,* 357.
BALLMER-HOFER, K., ANDERSSON, A. E., RATCLIFFE, L. E. & BERGER, P. 2011. Neuropilin-1 promotes
VEGFR-2 trafficking through Rab11 vesicles thereby specifying signal output. *Blood,* 118**,** 816-
26.
BARTLETT, C. S., SCOTT, R. P., CAROTA, I. A., WNUK, M. L., KANWAR, Y. S., MINER, J. H. & QUAGGIN,
S. E. 2017. Glomerular mesangial cell recruitment and function require the co-receptor
neuropilin-1. *Am J Physiol Renal Physiol,* 313**,** F1232-F1242.
BAYLISS, A. L., SUNDARARAMAN, A., GRANET, C. & MELLOR, H. 2020. Raftlin is recruited by
neuropilin-1 to the activated VEGFR2 complex to control proangiogenic signaling.
Angiogenesis, 23**,** 371-383.
BECKER, P. M., WALTENBERGER, J., YACHECHKO, R., MIRZAPOIAZOVA, T., SHAM, J. S., LEE, C. G.,
ELIAS, J. A. & VERIN, A. D. 2005. Neuropilin-1 regulates vascular endothelial growth factor-
mediated endothelial permeability. *Circ Res,* 96**,** 1257-65.
BOSSEBOEUF, E., CHIKH, A., CHAKER, A. B., MITCHELL, T. P., VIGNARAJA, D., RAJENDRAKUMAR, R.,
KHAMBATA, R. S., NIGHTINGALE, T. D., MASON, J. C., RANDI, A. M., AHLUWALIA, A. &
RAIMONDI, C. 2023. Neuropilin-1 interacts with VE-cadherin and TGFBR2 to stabilize
adherens junctions and prevent activation of endothelium under flow. *Sci Signal,* 16**,**
eabo4863.
CAI, H. & REED, R. R. 1999. Cloning and characterization of neuropilin-1-interacting protein: a PSD-
95/Dlg/ZO-1 domain-containing protein that interacts with the cytoplasmic domain of
neuropilin-1. *J Neurosci,* 19**,** 6519-27.
CERANI, A., TETREAULT, N., MENARD, C., LAPALME, E., PATEL, C., SITARAS, N., BEAUDOIN, F.,
LEBOEUF, D., DE GUIRE, V., BINET, F., DEJDA, A., REZENDE, F. A., MILOUDI, K. & SAPIEHA, P.
2013. Neuron-derived semaphorin 3A is an early inducer of vascular permeability in diabetic
retinopathy via neuropilin-1. *Cell Metab,* 18**,** 505-18.
CORTI, F., RISTORI, E., RIVERA-MOLINA, F., TOOMRE, D., ZHANG, J., MIHAILOVIC, J., ZHUANG, Z. W. &
SIMONS, M. 2022. Syndecan-2 selectively regulates VEGF-induced vascular permeability.
Nature Cardiovascular Research, 1**,** 518-528.

FANTIN, A., LAMPROPOULOU, A., SENATORE, V., BRASH, J. T., PRAHST, C., LANGE, C. A., LIYANAGE, S. E., RAIMONDI, C., BAINBRIDGE, J. W., AUGUSTIN, H. G. & RUHRBERG, C. 2017. VEGF165-induced vascular permeability requires NRP1 for ABL-mediated SRC family kinase activation. *J Exp Med,* 214**,** 1049-1064.

FERNÁNDEZ-ROBREDO, P., SELVAM, S., POWNER, M. B., SIM, D. A. & FRUTTIGER, M. 2017. Neuropilin 1 Involvement in Choroidal and Retinal Neovascularisation. *PLoS One,* 12**,** e0169865.

GIOELLI, N., NEILSON, L. J., WEI, N., VILLARI, G., CHEN, W., KUHLE, B., EHLING, M., MAIONE, F., WILLOX, S., BRUNDU, S., AVANZATO, D., KOULOURAS, G., MAZZONE, M., GIRAUDO, E., YANG, X. L., VALDEMBRI, D., ZANIVAN, S. & SERINI, G. 2022. Neuropilin 1 and its inhibitory ligand mini-tryptophanyl-tRNA synthetase inversely regulate VE-cadherin turnover and vascular permeability. *Nat Commun,* 13**,** 4188.

GOSHIMA, Y., HORI, H., SASAKI, Y., YANG, T., KAGOSHIMA-MAEZONO, M., LI, C., TAKENAKA, T., NAKAMURA, F., TAKAHASHI, T., STRITTMATTER, S. M., MISU, Y. & KAWAKAMI, T. 1999. Growth cone neuropilin-1 mediates collapsin-1/Sema III facilitation of antero- and retrograde axoplasmic transport. *J Neurobiol,* 39**,** 579-89.

GU, C., LIMBERG, B. J., WHITAKER, G. B., PERMAN, B., LEAHY, D. J., ROSENBAUM, J. S., GINTY, D. D. & KOLODKIN, A. L. 2002. Characterization of neuropilin-1 structural features that confer binding to semaphorin 3A and vascular endothelial growth factor 165. *J Biol Chem,* 277**,** 18069-76.

HONG, T. M., CHEN, Y. L., WU, Y. Y., YUAN, A., CHAO, Y. C., CHUNG, Y. C., WU, M. H., YANG, S. C., PAN, S. H., SHIH, J. Y., CHAN, W. K. & YANG, P. C. 2007. Targeting neuropilin 1 as an antitumor strategy in lung cancer. *Clin Cancer Res,* 13**,** 4759-68.

HONKURA, N., RICHARDS, M., LAVINA, B., SAINZ-JASPEADO, M., BETSHOLTZ, C. & CLAESSON-WELSH, L. 2018. Intravital imaging-based analysis tools for vessel identification and assessment of concurrent dynamic vascular events. *Nat Commun,* 9**,** 2746.

HOROWITZ, A. & SEERAPU, H. R. 2012. Regulation of VEGF signaling by membrane traffic. *Cell Signal,* 24**,** 1810-20.

HU, B., GUO, P., BAR-JOSEPH, I., IMANISHI, Y., JARZYNKA, M. J., BOGLER, O., MIKKELSEN, T., HIROSE, T., NISHIKAWA, R. & CHENG, S. Y. 2007. Neuropilin-1 promotes human glioma progression through potentiating the activity of the HGF/SF autocrine pathway. *Oncogene,* 26**,** 5577-86.

JIN, Y., DING, Y., RICHARDS, M., KAAKINEN, M., GIESE, W., BAUMANN, E., SZYMBORSKA, A., ROSA, A., NORDLING, S., SCHIMMEL, L., AKMERIÇ, E. B., PENA, A., NWADOZI, E., JAMALPOUR, M., HOLSTEIN, K., SÁINZ-JASPEADO, M., BERNABEU, M. O., WELSH, M., GORDON, E., FRANCO, C. A., VESTWEBER, D., EKLUND, L., GERHARDT, H. & CLAESSON-WELSH, L. 2022. Tyrosine-protein kinase Yes controls endothelial junctional plasticity and barrier integrity by regulating VE-cadherin phosphorylation and endocytosis. *Nature Cardiovascular Research,* 1**,** 1156-1173.

JUNG, B., ARNOLD, T. D., RASCHPERGER, E., GAENGEL, K. & BETSHOLTZ, C. 2018. Visualization of vascular mural cells in developing brain using genetically labeled transgenic reporter mice. *J Cereb Blood Flow Metab,* 38**,** 456-468.

KAWAKAMI, T., TOKUNAGA, T., HATANAKA, H., KIJIMA, H., YAMAZAKI, H., ABE, Y., OSAMURA, Y., INOUE, H., UEYAMA, Y. & NAKAMURA, M. 2002. Neuropilin 1 and neuropilin 2 co-expression is significantly correlated with increased vascularity and poor prognosis in nonsmall cell lung carcinoma. *Cancer,* 95**,** 2196-201.

KAWASAKI, T., KITSUKAWA, T., BEKKU, Y., MATSUDA, Y., SANBO, M., YAGI, T. & FUJISAWA, H. 1999. A requirement for neuropilin-1 in embryonic vessel formation. *Development,* 126**,** 4895-902.

KITSUKAWA, T., SHIMONO, A., KAWAKAMI, A., KONDOH, H. & FUJISAWA, H. 1995. Overexpression of a membrane protein, neuropilin, in chimeric mice causes anomalies in the cardiovascular system, nervous system and limbs. *Development,* 121**,** 4309-18.

KOCH, S., VAN MEETEREN, L. A., MORIN, E., TESTINI, C., WESTRÖM, S., BJÖRKELUND, H., LE JAN, S., ADLER, J., BERGER, P. & CLAESSON-WELSH, L. 2014. NRP1 presented in trans to the

667   endothelium arrests VEGFR2 endocytosis, preventing angiogenic signaling and tumor
668   initiation. *Dev Cell,* 28**,** 633-46.

669 LANAHAN, A., ZHANG, X., FANTIN, A., ZHUANG, Z., RIVERA-MOLINA, F., SPEICHINGER, K., PRAHST, C.,
670   ZHANG, J., WANG, Y., DAVIS, G., TOOMRE, D., RUHRBERG, C. & SIMONS, M. 2013. The
671   neuropilin 1 cytoplasmic domain is required for VEGF-A-dependent arteriogenesis. *Dev Cell,*
672   25**,** 156-68.

673 LANAHAN, A. A., HERMANS, K., CLAES, F., KERLEY-HAMILTON, J. S., ZHUANG, Z. W., GIORDANO, F. J.,
674   CARMELIET, P. & SIMONS, M. 2010. VEGF receptor 2 endocytic trafficking regulates arterial
675   morphogenesis. *Dev Cell,* 18**,** 713-24.

676 LI, X., PADHAN, N., SJÖSTRÖM, E. O., ROCHE, F. P., TESTINI, C., HONKURA, N., SÁINZ-JASPEADO, M.,
677   GORDON, E., BENTLEY, K., PHILIPPIDES, A., TOLMACHEV, V., DEJANA, E., STAN, R. V.,
678   VESTWEBER, D., BALLMER-HOFER, K., BETSHOLTZ, C., PIETRAS, K., JANSSON, L. & CLAESSON-
679   WELSH, L. 2016. VEGFR2 pY949 signalling regulates adherens junction integrity and
680   metastatic spread. *Nat Commun,* 7**,** 11017.

681 MAMLUK, R., GECHTMAN, Z., KUTCHER, M. E., GASIUNAS, N., GALLAGHER, J. & KLAGSBRUN, M.
682   2002. Neuropilin-1 binds vascular endothelial growth factor 165, placenta growth factor-2,
683   and heparin via its b1b2 domain. *J Biol Chem,* 277**,** 24818-25.

684 MATSUMOTO, T., BOHMAN, S., DIXELIUS, J., BERGE, T., DIMBERG, A., MAGNUSSON, P., WANG, L.,
685   WIKNER, C., QI, J. H., WERNSTEDT, C., WU, J., BRUHEIM, S., MUGISHIMA, H.,
686   MUKHOPADHYAY, D., SPURKLAND, A. & CLAESSON-WELSH, L. 2005. VEGF receptor-2 Y951
687   signaling and a role for the adapter molecule TSAd in tumor angiogenesis. *Embo j,* 24, 2342-
688   53.

689 MCGOWAN, S. E. & MCCOY, D. M. 2021. Neuropilin-1 directs PDGFRalpha-entry into lung fibroblasts
690   and signaling from very early endosomes. *Am J Physiol Lung Cell Mol Physiol,* 320, L179-L192.

691 MIAO, H. Q., LEE, P., LIN, H., SOKER, S. & KLAGSBRUN, M. 2000. Neuropilin-1 expression by tumor
692   cells promotes tumor angiogenesis and progression. *FASEB J,* 14**,** 2532-9.

693 MORIN, E., LINDSKOG, C., JOHANSSON, M., EGEVAD, L., SANDSTRÖM, P., HARMENBERG, U.,
694   CLAESSON-WELSH, L. & SJÖBERG, E. 2020. Perivascular Neuropilin-1 expression is an
695   independent marker of improved survival in renal cell carcinoma. *The Journal of Pathology,*
696   250**,** 387-396.

697 MORIN, E., SJÖBERG, E., TJOMSLAND, V., TESTINI, C., LINDSKOG, C., FRANKLIN, O., SUND, M.,
698   ÖHLUND, D., KIFLEMARIAM, S., SJÖBLOM, T. & CLAESSON-WELSH, L. 2018. VEGF receptor-
699   2/neuropilin 1 trans-complex formation between endothelial and tumor cells is an
700   independent predictor of pancreatic cancer survival. *The Journal of Pathology,* 246**,** 311-322.

701 NACCACHE, S. N., HASSON, T. & HOROWITZ, A. 2006. Binding of internalized receptors to the PDZ
702   domain of GIPC/synectin recruits myosin VI to endocytic vesicles. *Proc Natl Acad Sci U S A,*
703   103**,** 12735-40.

704 ORSENIGO, F., GIAMPIETRO, C., FERRARI, A., CORADA, M., GALAUP, A., SIGISMUND, S., RISTAGNO,
705   G., MADDALUNO, L., KOH, G. Y., FRANCO, D., KURTCUOGLU, V., POULIKAKOS, D., BALUK, P.,
706   MCDONALD, D., GRAZIA LAMPUGNANI, M. & DEJANA, E. 2012. Phosphorylation of VE-
707   cadherin is modulated by haemodynamic forces and contributes to the regulation of vascular
708   permeability in vivo. *Nat Commun,* 3**,** 1208.

709 PAN, Q., CHANTHERY, Y., LIANG, W. C., STAWICKI, S., MAK, J., RATHORE, N., TONG, R. K., KOWALSKI,
710   J., YEE, S. F., PACHECO, G., ROSS, S., CHENG, Z., LE COUTER, J., PLOWMAN, G., PEALE, F.,
711   KOCH, A. W., WU, Y., BAGRI, A., TESSIER-LAVIGNE, M. & WATTS, R. J. 2007. Blocking
712   neuropilin-1 function has an additive effect with anti-VEGF to inhibit tumor growth. *Cancer*
713   *Cell,* 11**,** 53-67.

714 PONUWEI, G. A. 2016. A glimpse of the ERM proteins. *J Biomed Sci,* 23**,** 35.

715 RAIMONDI, C., FANTIN, A., LAMPROPOULOU, A., DENTI, L., CHIKH, A. & RUHRBERG, C. 2014. Imatinib
716   inhibits VEGF-independent angiogenesis by targeting neuropilin 1-dependent ABL1 activation
717   in endothelial cells. *J Exp Med,* 211**,** 1167-83.

RICHARDS, M., NWADOZI, E., PAL, S., MARTINSSON, P., KAAKINEN, M., GLOGER, M., SJÖBERG, E.,
KOLTOWSKA, K., BETSHOLTZ, C., EKLUND, L., NORDLING, S. & CLAESSON-WELSH, L. 2022.
Claudin5 protects the peripheral endothelial barrier in an organ and vessel-type-specific
manner. *Elife,* 11.
RICHARDS, M., PAL, S., SJÖBERG, E., MARTINSSON, P., VENKATRAMAN, L. & CLAESSON-WELSH, L.
2021. Intra-vessel heterogeneity establishes enhanced sites of macromolecular leakage
downstream of laminin α5. *Cell Rep,* 35**,** 109268.
ROTH, L., PRAHST, C., RUCKDESCHEL, T., SAVANT, S., WESTRÖM, S., FANTIN, A., RIEDEL, M.,
HÉROULT, M., RUHRBERG, C. & AUGUSTIN, H. G. 2016. Neuropilin-1 mediates vascular
permeability independently of vascular endothelial growth factor receptor-2 activation. *Sci*
Signal, 9**,** ra42.
SARRIS, M., ANDERSEN, K. G., RANDOW, F., MAYR, L. & BETZ, A. G. 2008. Neuropilin-1 expression on
regulatory T cells enhances their interactions with dendritic cells during antigen recognition.
Immunity, 28**,** 402-13.
SHAHRABI-FARAHANI, S., GALLOTTINI, M., MARTINS, F., LI, E., MUDGE, D. R., NAKAYAMA, H., HIDA,
K., PANIGRAHY, D., D'AMORE, P. A. & BIELENBERG, D. R. 2016. Neuropilin 1 Receptor Is Up-
Regulated in Dysplastic Epithelium and Oral Squamous Cell Carcinoma. *Am J Pathol,* 186**,**
1055-64.
SHERAFAT, A., PFEIFFER, F., REISS, A. M., WOOD, W. M. & NISHIYAMA, A. 2021. Microglial neuropilin-
1 promotes oligodendrocyte expansion during development and remyelination by trans-
activating platelet-derived growth factor receptor. *Nat Commun,* 12**,** 2265.
SJOBERG, E., MELSSEN, M., RICHARDS, M., DING, Y., CHANOCA, C., CHEN, D., NWADOZI, E., PAL, S.,
LOVE, D. T., NINCHOJI, T., SHIBUYA, M., SIMONS, M., DIMBERG, A. & CLAESSON-WELSH, L.
2023. Endothelial VEGFR2-PLCgamma signaling regulates vascular permeability and
antitumor immunity through eNOS/Src. *J Clin Invest,* 133.
SOKER, S., MIAO, H. Q., NOMI, M., TAKASHIMA, S. & KLAGSBRUN, M. 2002. VEGF165 mediates
formation of complexes containing VEGFR-2 and neuropilin-1 that enhance VEGF165-
receptor binding. *J Cell Biochem,* 85**,** 357-68.
SUN, Z., LI, X., MASSENA, S., KUTSCHERA, S., PADHAN, N., GUALANDI, L., SUNDVOLD-GJERSTAD, V.,
GUSTAFSSON, K., CHOY, W. W., ZANG, G., QUACH, M., JANSSON, L., PHILLIPSON, M., ABID,
M. R., SPURKLAND, A. & CLAESSON-WELSH, L. 2012. VEGFR2 induces c-Src signaling and
vascular permeability in vivo via the adaptor protein TSAd. *J Exp Med,* 209**,** 1363-77.
WANG, J., WANG, S., LI, M., WU, D., LIU, F., YANG, R., JI, S., JI, A. & LI, Y. 2015. The Neuropilin-1
Inhibitor, ATWLPPR Peptide, Prevents Experimental Diabetes-Induced Retinal Injury by
Preserving Vascular Integrity and Decreasing Oxidative Stress. *PLoS One,* 10**,** e0142571.
WESSEL, F., WINDERLICH, M., HOLM, M., FRYE, M., RIVERA-GALDOS, R., VOCKEL, M., LINNEPE, R., IPE,
U., STADTMANN, A., ZARBOCK, A., NOTTEBAUM, A. F. & VESTWEBER, D. 2014. Leukocyte
extravasation and vascular permeability are each controlled in vivo by different tyrosine
residues of VE-cadherin. *Nat Immunol,* 15**,** 223-30.
WNUK, M., ANDEREGG, M. A., GRABER, W. A., BUERGY, R., FUSTER, D. G. & DJONOV, V. 2017.
Neuropilin1 regulates glomerular function and basement membrane composition through
pericytes in the mouse kidney. *Kidney Int,* 91**,** 868-879.

Figure 1

Figure 2

A Tamoxifen IP (80mg/kg)
d1 d2 d3 d4 d5
4 days post treatment
VEGFA
8-16 weeks
Nrp1^fl/fl; Actb^CreERT2 mice

B Lung lysates
Control *Nrp1* iKO
NRP1 140 kDa
GAPDH 36 kDa
NRP1/GAPDH
Control *Nrp1* iKO
**

C Ear dermis
t=0s t=200s
2000 kDa dextran
Control
Nrp1 iKO

D Leakage sites / 100um
Control *Nrp1* iKO
Venule Capillary
*

E Control *Nrp1* iKO
CD31
2000 kDa dextran
Trachea
Dextran area / Vessel area
Control *Nrp1* iKO
**

F Control *Nrp1* iKO
CD31
2000 kDa dextran
Back skin
Dextran area / Vessel area
Control *Nrp1* iKO
*

Figure 3

Figure 4

A

Nrp1 iECKO **X** *Vegfr2*^{Y949F/Y949F} → *Vegfr2*^{Y949F/Y949F}; *Nrp1* iECKO 8-16 weeks

Tamoxifen IP (80mg/kg)

4 days post treatment

d1 d2 d3 d4 d5 → VEGFA

B

Lung lysates

Vegfr2^{Y949F/Y949F}   *Vegfr2*^{Y949F/Y949F}; *Nrp1* iECKO

NRP1 — 140 kDa

GAPDH — 36 kDa

NRP1 / GAPDH   ****

Vegfr2^{Y949F/Y949F}   *Vegfr2*^{Y949F/Y949F}; *Nrp1* iECKO

C Ear dermis

t = 0s   t = 200s

2000 kDa dextran

Vegfr2^{Y949F/Y949F}

Vegfr2^{Y949F/Y949F}; *Nrp1* iECKO

D

Number of leakage sites normalized to control

Venule

- Control
- ▲ *Nrp1* iECKO
- ○ *Vegfr2*^{Y949F/Y949F}
- △ *Vegfr2*^{Y949F/Y949F}; *Nrp1* iECKO

** ** *

E

Vegfr2^{+/+}   *Vegfr2*^{Y949F/Y949F}

Control   *Nrp1* iECKO   Control   *Nrp1* iECKO

Trachea

CD31
2000 kDa dextran

Dextran area/Vessel area

*** ** ***

p=0.06

Trachea

- Control
- ▲ *Nrp1* iECKO
- ○ *Vegfr2*^{Y949F/Y949F}
- △ *Vegfr2*^{Y949F/Y949F}; *Nrp1* iECKO

F

Vegfr2^{+/+}   *Vegfr2*^{Y949F/Y949F}

Control   *Nrp1* iECKO   Control   *Nrp1* iECKO

Back skin

CD31
2000 kDa dextran

Dextran area/Vessel area

*** *** ***

Back skin

- Control
- ▲ *Nrp1* iECKO
- ○ *Vegfr2*^{Y949F/Y949F}
- △ *Vegfr2*^{Y949F/Y949F}; *Nrp1* iECKO

Figure 5

Figure 6

Supp fig 1

A

NRP1 expression in ear dermis

B

NRP1 expression in back skin

Supp fig 2

A

10 kDa

B

70 kDa

C

2000 kDa dextran

D

2000 kDa dextran

E

2000 kDa dextran

Supp fig 3

Supp fig 4

A

VEGFA stimulation in the ear dermis

B

VEGFA stimulation in the trachea

C

VEGFA stimulation in the back skin

Supp fig 5