Local Inhibitory Dynamics Underpin Temporal Integration and Functional Segregation between Barrels and Septa in the Mouse Barrel Cortex

Ali Özgür Argunşah[1,2], Tevye Jason Stachniak[1,2,3], Jenq-Wei Yang[1,2], Linbi Cai[1,2], Alexander van der Bourg[1,2], Rahel Kastli[1,2,4], Theofanis Karayannis[1,2,5,*]

[1]Laboratory of Neural Circuit Assembly, Brain Research Institute, University of Zurich Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.
[2]Neuroscience Center Zurich, University of Zurich and ETH Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.
[3]Division of Biomedical Sciences, Faculty of Medicine, Memorial University of Newfoundland, St. John's, A1B 3V6, Canada.
[4]Department of Stem Cell & Regenerative Biology, Harvard University, Cambridge, MA, USA.
[5]University Research Priority Program (URPP), Adaptive Brain Circuits in Development and Learning, University of Zurich, Zurich, Switzerland.
*Correspondence: karayannis@hifo.uzh.ch

**Abstract**

Just as separated digits and repeated sampling enhance somatosensation in humans, mice sense objects through multiple segregated whiskers through successive contacts. Individual whisker identity is maintained through the somatotopic organization of the Whisker→Brainstem→Thalamus→Cortex axis, culminating in distinct cortical domains: barrels and the surrounding septa. By performing simultaneous recordings using in-vivo electrophysiology in wild-type (WT) mice, we identify a progressive divergence in spiking activity between these domains upon repeated behaviorally relevant (10Hz) single- and multi-whisker stimulation. While the spiking activity ratio of multi- to single-whisker stimulation remains stable in barrels, it increases progressively in septa, suggesting inhibitory cell recruitment. Using genetic fate-mapping and tissue clearing, we indeed reveal that SST+ and VIP+ interneurons exhibit distinct laminar and regional distributions in barrel and septa domains. Further, calcium imaging of SST+ and VIP+ interneurons

shows that while both neuron types respond to single-whisker stimulus, SST+
interneurons preferentially engage more in 10Hz multi-whisker stimulation, indicating their
critical role in progressive stimulus preference. Genetic removal of Elfn1, which regulates
the incoming excitatory synaptic dynamics onto SST+ interneurons, leads to the loss of
the progressive increase in septal spiking activity upon multi-whisker stimulation,
resembling the stable response pattern observed in barrels. The importance of the loss
of functional segregation of barrels, versus septa is revealed by cumulative temporal
decoding analysis, supporting the notion that SST+ interneuron-mediated inhibition
contributes to temporal encoding and stimulus integration. Finally, viral tracing combined
with whole brain clearing and imaging reveals that barrel and septa domains project
differentially to secondary somatosensory (S2) and motor (M1) cortices. These distinct
projection patterns suggest that differential inhibitory processing in barrels and septa may
contribute to functionally specialized downstream signaling. Together, our findings
indicate that the progressive engagement of SST+ interneurons, mediated by Elfn1-
dependent synaptic facilitation, underlies the preferential integration of multi-whisker
stimuli in septa. This local inhibitory mechanism likely contributes to the functional
segregation of barrel and septa domains and their distinct cortical projections, shaping
how sensory information is processed and relayed to higher-order brain regions.
50

51 **Keywords**
52 Barrel Cortex, Elfn1 Knockout, Somatosensory Whisker Coding, Cortical Domains,
53 Temporal Dynamics, Synaptic Integration, Sensory Projections, Network Segregation
54

**Introduction**
Although the somatosensory whisker system is one of the principal means by which mice
sense their environment and navigate the world, these tasks are carried out by a relatively
small number of whiskers. This small number of vibrissae is represented in the whisker
somatosensory cortex as barrel "islands" (columns) separated and surrounded by the
septal "sea" (compartments), reminiscent of the whisker pad pattern on a rodent's face,
where sparse hair follicles are separated by spaces between them. In contrast, other
sensory systems, such as the auditory and the visual ones, have densely packed

anatomical receptor configurations and cortical representations. Although all sensory
cortices are composed of canonical microcircuits (Douglas and Martin, 2004) that include
similar populations of excitatory and inhibitory neurons, maintaining segregated individual
whisker information may require anatomical specializations, the barrel and septa
domains. These domains may therefore create a distinct spatiotemporal stimulus
representation and information coding in the barrel cortex, different from the other sensory
modalities. In the adult murine barrel cortex, thalamocortical inputs coming from the
ventro-postero-medial (VPM) and postero-medial (PoM) nuclei segregate into the barrel
and septa domains, respectively (Alloway, 2008; Bureau et al., 2006; Kim and Ebner,
1999; Staiger and Petersen, 2021). Further, in mice (Audette et al., 2018; Sato and
Svoboda, 2010; Shepherd and Svoboda, 2005) and rats (Alloway, 2008; Chakrabarti and
Alloway, 2006; Melzer et al., 2006b) it was shown that information processing and flow
within and from these domains target anatomically distinct sensory and motor areas. It
has been suggested that the primary whisker somatosensory cortex (wS1) sends higher
dimensional information needed for object recognition to the secondary somatosensory
cortex (wS2), while the projections to the primary motor cortex (M1) carry less complex
information that does not require segregated streams of information coming from
individual whiskers (Alloway, 2008; Brecht and Sakmann, 2002; Cai et al., 2022; Sato
and Svoboda, 2010). This divergence in information processing is also found within the
barrel- and septa-related circuits. While barrel circuits are more involved in processing
spatiotemporal whisker-object interactions, the septal circuits are more sensitive to the
frequency of whisker movements, which suggests the use of temporal versus rate coding
of information in barrel and septa, respectively (Melzer et al., 2006a, 2006b). Hence the
existence of the septal column separating the barrels may provide mice with the means
to separate these two interconnected yet very distinct information-processing routes
along the temporal domain, and to make sense of their somatosensory environment
through repeated sampling via cloistered information streams. Recent studies in mice
have shown that barrel and septa compartments are not only anatomically segregated in
terms of cell-type specific thalamocortical projections and differential expression of
different genetic markers (Young et al., 2023) but also in their tuning upon single- and
multi- whisker stimulation (Wang et al. 2022).

In both rats and mice, the operational identity of barrels and septa has often been
attributed to the parallel bottom-up thalamocortical pathways of VPM and PoM. Here, by
performing *in-vivo* silicon probe recordings simultaneously in the barrel and septa
domains upon repeated single- and multi- whisker stimulation in wS1 of mice, we first
show and characterize the temporal divergence between these domains. Utilizing genetic
fate-mapping, passive tissue clearing, and light-sheet microscopy, we reveal that cortical
somatostatin (SST+) and vasoactive intestinal peptide (VIP+) expressing interneurons
show a layer-dependent differential distribution in the barrel versus septa columns.
Through two-photon calcium imaging of these two populations, we show a differential
engagement of SST+ cells upon multi- vs. single-whisker stimulation. By altering the
short-term synaptic dynamics of incoming excitation onto these interneurons through the
removal of the trans-synaptic protein Elfn1 (Elfn1 KO), we find that the column-specific
contrast ratio between multi- over single- whisker response to a repetitive stimulus is
diminished. Through a progressive decoding analysis, we find that in contrast to control,
whisker stimulation-evoked Elfn1 KO responses in different domains cannot be decoded
efficiently. Finally, using *in-vivo* retrograde viral-based tracing, we show a layer-specific
projection preference between these domains and downstream regions wS2 and M1.
Hence, in addition to the anatomical specialization through thalamocortical pathways, we
reveal a key contribution of local lateral cortical inhibition, provided by SST+ interneurons,
in setting up the functional segregation of barrel and septa domains, and subsequently,
their downstream targets.
115
116   **Results**
117
**Barrel and Septa columns display differentially adaptive spiking upon repeated**
**single- and multi- whisker stimulation**
To characterize the temporal stimulus-response profiles of barrel and septal domains
simultaneously, we performed *in-vivo* multi-shank (8x8) silicon probe recordings under
urethane-induced light anesthesia from mouse somatosensory whisker barrel cortex
(wS1), after the age of active whisking onset, postnatal day (P)20-30. Whisker stimulation
was performed on either just one principal whisker (single-whisker stimulus or SWS) or
via both the principal whisker and most of the macro vibrissae together (multi-whisker

stimulus or MWS; see methods) (Figure 1A). Single- or multi- whisker evoked activity within identified regions was recorded upon a 2s-long stimulation at 10Hz, similar to the average frequency at which mice actively whisk (Carvell and Simons, 1990; Grant et al., 2012). Each mouse received both stimulation paradigms. Current source density (CSD) analysis was used to assess the location of each silicon probe shank, which was later verified by post-hoc histology. The recorded activity was assigned to the stimulated principal barrel column (B), the adjacent septal column (S), or the adjacent unstimulated neighboring barrel column (N) according to SWS paradigm (Figure 1B and Supp Figure 1A to D). An example probe assignment can be seen in Supp Figure 1E. While this distinctive experimental paradigm is relatively low throughput, it has allowed us to investigate all three of these domains simultaneously. The septal compartments in mouse barrel cortex are quite narrow (Bureau et al., 2006; Sato et al., 2007) and there is the possibility of rotational misalignment during the insertion of silicon probes into the brain (Supp. Figure 1F). For these reasons we restricted our analysis of *in-vivo* responses upon SWS and MWS between barrel and septa responses to upper layers and L4, for which we have higher confidence about the anatomical domain allocation after our histological (Supp. Figure 1B) and CSD analysis (Supp. Figure 1C and D). In line with this domain allocation, t-distributed stochastic neighbor embedding (t-SNE) analysis of the multi-unit activity (MUA) recorded across the entire train of stimuli in the different domains revealed that the spiking responses between the principal barrel and the neighboring one form two separate clusters, while the responses recorded in the septa lay in between (Figure 1C; see methods).

It has previously been shown that electrophysiological responses upon whisking-based object detection saturate rapidly after a single whisker touch, while object localization may require multiple repeated whisker touches (O'Connor et al., 2013; Pammer et al., 2013). Correspondingly, when a single whisker is stimulated repeatedly, the response to the first pulse is principally bottom-up thalamic-driven responses, while the later pulses in the train are expected to also gradually engage cortico-thalamo-cortical and cortico-cortical loops. We therefore first tested whether the leading pulses of the stimulus train evoked differential responses among the principal barrel, the adjacent septa, and the neighboring barrel upon SWS and MWS. For easier visualization and full appreciation of the data, we

present our findings both in classical time series firing-rate plot and stimulation pulse
aligned stacked images (Supp. Figure 2A and B, 200ms baseline followed by 2000ms
evoked response to 20 pulses at 10Hz reshaped to 20 stacked 100ms-long evoked
response image, left to right). Earlier studies that separately recorded whisker-evoked
responses from either barrel or septa domains in rat reported that the responses of barrel
neurons to single principal whisker stimulation are much higher than those evoked in the
septal neurons (Armstrong-James and Fox, 1987; Brecht and Sakmann, 2002). The
analysis of the data we obtained from the simultaneous recordings at these domains in
the mouse do not reveal such differences in the first two stimulus responses in L4 upon
single whisker stimulation (SWS) (Figure 1D; statistics on the bottom; white represents
no significance). However, response differences emerged from the 3$^{rd}$ pulse onward upon
SWS (Figure 1D, bottom). Similar differences also emerged between stimulated principal
and unstimulated neighboring barrel columns upon SWS (starting from 4$^{th}$ pulse,
Figure1D, bottom, Barrel-vs-Neighbor (BN) row). Interestingly, an inverted pattern of
significance appeared between septa and adjacent unstimulated neighboring barrels
(Figure 1D, bottom, Septa-vs-Neighbor (SN) row). When multiple whiskers (MWS) were
activated simultaneously, like SWS, a divergence between barrel and septa domain
activity also occurred in Layer 4 from the 2$^{nd}$ pulse onward (Figure 1E). Finally, when we
calculated the response ratio upon MW over SW stimulation, we saw that the septal
MW/SW ratio diverged from barrel MW/SW ratio over the course of repeated stimulation
(Figure 1F), indicating a role of septal neurons in the progressive processing MW
information.
179

**Barrel and Septa columns display differential SST+ and VIP+ neuron densities**
It is known that barrel and septa domains receive different thalamic projections, from VPM
and PoM respectively, that might affect stimulus response properties (Ahissar et al., 2001;
Chmielowska et al., 1989; Koralek et al., 1988; Sosnik et al., 2001). Our
electrophysiological experiments show a significant divergence of responses over time
upon both SWS and MWS in L4 between barrels (principal and neighboring) and adjacent
septa, with minimal initial difference. We hypothesized that this late divergence might be
driven by differences in the local circuitry, and specifically inhibitory cells. We therefore

next assessed the spatial distribution of two inhibitory cell populations that could provide

distinct regulation of cortical activity in the temporal domain: SST+ and VIP+ cells.

Earlier research examining the three largest inhibitory neuron populations in the mouse

barrel cortex showed that, while parvalbumin positive (PV+) cells do not show any

differential density preference between barrel and septa columns, SST+ and VIP+ cells

have higher density in the septa in L4 (Almási et al., 2019). We attempted to verify the

reported differential SST+ and VIP+ neuronal distributions using an alternative imaging

and counting strategy. To label the two neuronal populations, we crossed the SST-Cre

and VIP-Cre lines with a tdTomato reporter mouse line (Ai14). To comprehensively

quantify the density of SST+ and VIP+ cells in the barrel and septa domains in 3D, we

utilized a passive CLARITY-based tissue clearing protocol, followed by light-sheet

microscopy using a custom-built light-sheet microscope (Vladimirov et al., 2023; Voigt et

al., 2019) (Figure 2A). The individual barrels in wS1 were reliably detected using auto-

fluorescence from the tissue acquired with a 488nm laser. tdTomato positive SST+ or

VIP+ neurons were detected by a 561nm laser, allowing us to accurately localize and

count cells in barrel and septal domains (Figure 2B and C). By adjusting the angle of each

barrel column in 3D (using Imaris, RRID:SCR_007370), we precisely identified the

column  through L2/3, L4 and L5, using the barrel borders for every plane (XY, YZ, XZ)

in L4 (Figure 2B, also see Methods). Once the columns were accurately determined, we

counted the number of tdTomato positive neurons in the barrels and septa and

corresponding volumes for density estimation. Despite their opposite distributions and

neuronal density changes along the depth of wS1 (Figure 2B and C), we found that both

SST+ (Figure 2D) and VIP+ (Figure 2E) neuronal densities were higher in L4 septa

compared to barrels, but we detected no difference in L2/3 and L5.

These results confirmed that SST+ and VIP+ interneurons have higher densities in septa

compared to barrels in L4 and suggest these interneurons may play a role in diverging

responses between barrels and septa upon SWS.

**SST+ neurons are more strongly activated by multi- versus single-whisker**

**stimulation**

Having identified spatial distribution patterns of SST+ or VIP+ interneurons that could
contribute to the single- vs multi- whisker response scaling observed in Figure 1F; we
sought to assess these interneurons' response profiles upon similar whisker stimulation
paradigms. Therefore, we performed similar SWS and MWS experiments using acute in-
vivo two-photon calcium (Ca2+) imaging under light anesthesia after weaning (P21–41)
(Figure 3A and B). We injected OGB-1, the membrane-permeable AM-ester form of the
calcium indicator, into wS1 of animals expressing tdTomato in either VIP+ or SST+ INs
(VIPCre-Ai14 and SSTCre-Ai14 lines, respectively) to localize VIP+ and SST+ activity
(Figure 3A). Two-photon Ca2+ imaging was performed upon stimulation of either the
principal C2 whisker at 10Hz for 2s (SWS) or most of the macro vibrissae including C2
(MWS) (Figure 3B). We found that both SST+ and VIP+ interneurons increased their firing
upon SWS (Figure 3C and D) as has been reported previously (Kastli et al., 2020), but
only SST+ interneurons changed their firing significantly between single- and multi-
whisker stimulation at 10-Hz (Figure 3D, SW vs MW comparison). Although these optical
recordings were performed in layer 2/3 (L2/3) rather than layer 4 (L4) due to depth
limitations, the similarity of the peak MW/SW ratios of L2/3 populations in putative barrel
and septa columns (c.f. Figure 1F and Supp. Figure 3) provides support for the validity of
using L2/3 data to infer respective L4 cell activity. This result suggests that differential
activation of SST+ interneurons is more likely to be involved in the domain specific
temporal ratio differences between barrels and septa.
238

**Progressive multi- over single-whisker response divergence in septal populations**
**is abolished in Elfn1 KO mice**
SST+ interneurons in the cortex are known to show distinct short-term synaptic plasticity,
particularly strong facilitation of excitatory inputs, which enables them to regulate the
temporal dynamics of cortical circuits. This facilitation allows SST+ cells to progressively
enhance their inhibitory influence in response to repeated stimulation, a property critical
for shaping network activity. A key regulator of this plasticity is the synaptic protein Elfn1,
which mediates short-term synaptic facilitation of excitation on SST+ interneurons
(Stachniak et al., 2021, 2019; Tomioka et al., 2014).

Having identified SST+ cells as potential regulators of temporal MW/SW ratio divergence,
we investigated how the absence of Elfn1 alters sensory-driven activity in vivo, potentially
shaping septal function in MW/SW response scaling. Using the same whisker stimulation
paradigms and domain-assignment strategy as in WT animals (Figure 1A and B; and
Supp. Figure 1), we recorded from barrel and septa domains in Elfn1 KO mice under both
single-whisker stimulation (SWS) and multi-whisker stimulation (MWS) conditions.
Unsupervised t-SNE analysis revealed that the domain-specific clustering of responses
seen in WT (Figure 1C) mice was disrupted in Elfn1 KO mice, with barrel and septa
responses intermingling (Figure 4A). This convergence of response profiles was further
clear in stimulus-aligned stacked images, where the emergent differences between
barrels and septa under SWS were largely abolished in the KO (Figure 4B). The only
remaining distinction was between directly stimulated barrels and neighboring barrels,
which persisted in the KO (Figure 4B). Under MWS conditions, the temporal divergence
between barrel and septa domains observed in WT Layer 4 was diminished in Elfn1 KO
mice (Figure 4C). Additionally, the progressive increase in the MW/SW response ratio
seen in septal neurons of WT mice was no longer seen in the KO (Figure 4D).
These findings suggest that in WT animals, activity spillover from principal barrels is
normally constrained by the progressive engagement of SST+ interneurons, driven by
Elfn1-dependent facilitation at their excitatory synapses. In the absence of Elfn1, this local
inhibitory mechanism is disrupted, leading to a loss of the distinct temporal response
divergence between barrel and septa domains.
269
**Barrel and Septa response decoding identity is lost in Elfn1 KO mice**
Building on our findings that Elfn1-dependent facilitation in SST+ interneurons is critical
for maintaining barrel-septa response divergence, we sought to quantify how sensory
information accumulates and is integrated over time in these domains—and how this
process is altered in Elfn1 KO mice.
To investigate this, we performed an accumulative temporal decoding analysis using
neural responses to 20 pulses of either single-whisker stimulation (SWS) or multi-whisker
stimulation (MWS) at 10 Hz for 2 seconds. We segmented the response window into three
time ranges to identify the temporal segments driving divergence between domains: the

full response window (1-95ms), the first part (1-50ms) which includes the first peak, and the late part (51-95ms), which includes the second firing peal (Figure 5A). A one-versus-all error-correcting output code (ECOC) classifier, using a GentleBoost ensemble of decision trees (Allwein et al., 2001; Friedman et al., 2000), was trained on firing profiles from WT barrel, septa, and neighboring barrel domains (3-class classification for SWS) or barrel and septa domains (2-class classification for MWS). We conducted 10-fold cross-validation within WT and KO datasets (WT CrossVal and KO CrossVal) and tested a WT-trained classifier on Elfn1 KO responses (WT Train - KO Test) to evaluate decoding performance across accumulated pulses (see Methods).

In WT mice, decoding accuracy increased progressively with the number of pulses upon SWS (Figure 5B, red) reflecting accumulated segregation of sensory information over time. This SWS WT CrossVal trend was more pronounced in the first response window (1-50ms) (Figure 5C vs 5D, red), indicating that domain-specific information accumulates predominantly in first temporal segments in repeated single whisker sampling. Cross-validation within the WT dataset (WT CrossVal) showed higher accuracy compared to the WT-trained classifier tested on KO responses (WT Train - KO Test) and KO CrossVal at this first segment, but not in the second (Figure 5B-D). Elfn1 KO dataset showed higher cross validation performance at the second part of the response profiles (Figure 5D) indicating a temporal information delay under the reduced progressive inhibition. While decoding curves started at similar initial levels for SWS data (Figures 5B-D, first pulse number), initial MW decoding accuracies were remarkably different (Figures 5E-G). In WT mice, while the decoding accuracy increased progressively as in SWS (Figure 5E, red), the difference between accumulated $20^{th}$ and the starting ($1^{st}$) accuracies were lower than KO CrossVal results (Figures 5E-G, red vs blue) which was predominantly due to the first part of the signal (Figure 5F vs 5G).

These results highlight the critical role of Elfn1 in facilitating the temporal integration of sensory inputs through its effects on SST+ interneurons. In WT mice, Elfn1-dependent short-term facilitation enables SST+ interneurons to progressively enhance their inhibitory influence, maintaining the functional segregation between barrel and septa domains. In Elfn1 KO mice, the loss of this facilitation mechanism disrupts the temporal dynamics necessary for accumulating domain-specific information, leading to a failure in

310 segregating sensory streams. This finding reinforces the importance of local inhibitory
311 circuits, particularly those involving SST+ interneurons, in shaping the temporal
312 processing of sensory information in the barrel cortex.
313

**wS1 Barrel Columns and Septal Domains have different wS2 and M1 Layer-**
**dependent projection preferences**
We identified that temporal MW/SW response divergence in septal populations is
abolished in Elfn1 KO mice. Given that barrel and septal compartments differentially
process sensory input over time, this raises the question of how these distinct streams of
information are relayed to downstream regions and whether they contribute to separate
functional pathways. Following up from this observation, we finally aimed to explore if the
projection targets of these two wS1 domains are distinct, in a similar manner to what has
been revealed for visual cortex domain projections (Meier et al., 2021). Previously, it has
been shown that cortical pyramidal neurons in wS1 exhibit distinct long-range projection
patterns and sensory tuning properties depending on their projection target (S2 or M1).
S2-projecting pyramidal neurons tend to have narrower receptive fields, are more likely
to be tuned to the columnar whisker and carry more precise sensory information about
whisker deflections. In contrast, M1-projecting pyramidal neurons are more broadly tuned,
integrating information from multiple whiskers, which aligns with their role in sensorimotor
integration and behavior (Sato and Svoboda, 2010; Yamashita et al., 2013). This data
could be suggestive of distinct localization of the different projection neurons in the
different wS1 domains. By utilizing *in vivo* retrograde AAV injections in two of the main
target cortical regions of wS1, we aimed to characterize the projections from wS1 to wS2
and M1.
We injected a retrograde AAV virus expressing GFP in wS2 or M1 in separate adult mice
and dissected the brains 4 weeks after the injection (Figure 6A). We utilized again the
passive CLARITY-based tissue clearing and imaged the whole brain using mesoSPIM.
The auto-fluorescence of the barrels allowed for labeling of barrel columns in 3D (Figure
6B). With the same strategy as our previous analysis on interneuron population
distributions (Figure 2), we counted the number of neurons in wS1 barrel and septa
domains across the cortical layers (Figure 6C) and measured the corresponding domain

volumes to estimate projection densities as a proxy for connectivity levels between S1-S2 and S1-M1.

As an internal control and in line with expectations, fewer projections arose from layer 4 barrels to either wS2 or M1 than from other cortical layers (Figure 6C and D). However, septa cells between mid-L4 and upper-L5B sent more wS2 projections (Figure 6D, left, turquois), whereas more neurons in septa from L3 to L4 project to M1, compared to the projection cells in barrels (Figure 6D, right, magenta). Additionally, a secondary comparative analysis showed that barrels in L3 host significantly more wS2-projecting-neurons (Figure 6E, left, purple), while L5A septal domains host significantly more M1-projecting neurons (Figure 6E, right, magenta). This differential targeting suggests that the temporal whisker-evoked processing specialization of barrel and septa domains may shape how distinct streams of sensory information are conveyed to higher-order cortical areas, potentially influencing perceptual processing in wS2 and sensorimotor integration in M1.

**Discussion**

In the absence of visual stimuli, humans explore objects through tactile interactions, where the initial touch begins the process of information accumulation, with subsequent touches refining the identification and characterization of the object. Rodents rely on a similar strategy during active whisking, where sequential whisker contacts allow for the progressive refinement of object recognition and spatial mapping (Armstrong-James and Fox, 1987; Kheradpezhouh et al., 2017; Petersen, 2007). This process involves parallel ascending pathways that transmit peripheral touch sensations to the cortex, with the primary somatosensory cortex mapping whisker spatial patterns directly onto cortical barrels, and in between spaces, septa (Van Der Loos and Woolsey, 1973; Woolsey et al., 1975).

Our study reveals that the functional segregation between barrel and septa domains in the mouse whisker primary somatosensory cortex (wS1) is not only anatomically defined by distinct thalamocortical projections but is also dynamically shaped by local inhibitory circuits, particularly those involving somatostatin-positive (SST+) interneurons. Through a combination of functional and anatomical techniques, we show that the temporal

divergence in spiking activity between barrel and septa domains is mediated by the short-
term synaptic facilitation properties of SST+ interneurons and are critically dependent on
the synaptic protein Elfn1. Furthermore, using a decoder we show that this temporal
divergence may affect the integration of sensory information during repeated whisker
stimulation, which may be  subsequently routed to differential downstream cortical areas,
such as the secondary somatosensory cortex (wS2) and primary motor cortex (M1).
Among the inhibitory neuron classes, SST+ interneurons are known for their high
spontaneous activity and powerful regulation of local neural networks through dense
feedback connections onto nearby pyramidal neurons (Urban-Ciecko and Barth, 2016).
In contrast, VIP+ interneurons influence network function through a disinhibitory circuit,
inhibiting SST+ (and to a lesser extent PV+) interneurons upon activation, which in turn
reduces inhibition on pyramidal cells (Karnani et al., 2016; Lee et al., 2013; Pi et al., 2013).
Thus SST+ interneurons may contribute to barrel-septal identity using surround inhibition,
while being regulated themselves by VIP+ interneuron activity under certain behaviorally
relevant sensory-motor tasks. Our analysis of SST+ cell distribution, shows a higher
density in the septa in L4, compared to barrels. Interestingly, layer 4 SST+ interneuron
axons remain within layer 4 (Xu et al., 2013) in the somatosensory cortex, unlike SST+ of
layers 2/3 and 5/6 which are Martinotti type and target layer 1. In addition, while Martinotti
cells receive facilitating synapses from pyramidal cells and form inhibitory synapses onto
dendrites of neighboring pyramidal cells (Silberberg and Markram, 2007), non-Martinotti
SST cells predominantly inhibit fast spiking cells in layer 4 (Ma et al., 2006; Xu et al.,
2013). During naturalistic whisking, SST+ and PV+ cells show divergent activation
patterns, with FS cells showing transient excitation due to rapid synaptic depression,
while SST+ cells show the opposite (Tan et al., 2008). It is currently unclear though under
which behavioral circumstances SST+ cells of L4 would engage in inhibition of excitatory
cells versus inhibition of PV+ cells. Our data suggest that under our experimental
conditions, a bottom-up activation paradigm, SST+ cells overall engage in the progressive
inhibition of the local L4 circuit, rather than disinhibition. To assess whether temporal
divergence in spiking carries meaningful sensory information, we conducted a detailed
accumulative temporal decoding analysis. We find that in wild-type mice, decoding
accuracy for SWS increased progressively with each additional pulse, particularly in the

initial response window (1-50ms), indicating that domain-specific information
accumulates primarily in early temporal segments during repeated whisker sampling. This
progressive accumulation of decodable information was severely disrupted in Elfn1 KO
mice, with cross-validation showing significantly reduced performance compared to wild-
type, especially in early response segments. Interestingly, Elfn1 KO mice showed higher
cross-validation performance in the later response segment (51-95ms), suggesting a
temporal delay in information processing due to reduced progressive inhibition. Under
multi-whisker stimulation, wild-type and knockout mice showed markedly different initial
decoding accuracies, with the difference between accumulated and initial accuracies
being lower in wild-type than in Elfn1 KO mice. Our anatomical connectivity mapping
would suggest that disruption in temporal information processing in the two domains is
crucial for maintaining the functional segregation of parallel processing streams to wS2
and M1.
The differential projection patterns of barrel and septa domains to wS2 and M1 further
suggest that the temporal processing capabilities of these domains influence how sensory
information is integrated with motor output. For example, the projection of septa domains
to M1 may facilitate the rapid translation of sensory information into motor commands,
while the projection of barrel domains to wS2 may support more detailed sensory
discrimination. The disruption of the informational integrity of these projections in Elfn1
KO mice could therefore have significant implications for sensorimotor integration and
behavior.
It is important to note that while our study focuses on the mouse barrel cortex, there are
significant differences between rodent species, such as rats and mice, in terms of cortical
organization and whisker system function. Most research on whisker stimulation had
traditionally focused on rats (Estebanez et al., 2016; Petersen, 2007; Roy et al., 2011;
Tanke et al., 2018). The existence of septa in the mouse barrel cortex has been debated,
with some suggesting that the distinction between barrel and septal circuits is less
pronounced in mice compared to rats (Bureau et al., 2006; Sato et al., 2007). For
example, the mouse barrel cortex has fewer neurons per column and thinner septa
compared to rats, which may influence the degree of functional segregation between
barrel and septa domains. While many animals have whiskers, only a subset of animals

434 have visible cortical barrels. For instance, cats have whiskers and brainstem barrelettes
435 but lack cortical barrels (Nomura et al., 1986; Rice, 1985; Rice et al., 1985). The presence
436 of barrels in the cortex seems to correlate with volitional whisking behavior, underlining
437 the importance of one-to-one mapping for processing individual whisker information. Our
438 findings show that septa are crucial in separating these parallel whisker streams by
439 modulating the lateral cortico-cortical information flow during repetitive whisker touches.
440 This modulation is significantly influenced by the short-term facilitation properties of
441 synaptic inputs onto SST+ interneurons, facilitated by the Elfn1 gene, alongside distinct
442 bottom-up inputs to each domain. The loss of temporal segregation in Elfn1 KO mice
443 could hinder object localization, which relies on multi-whisker integration in septa, while
444 sparing simpler detection tasks mediated by barrels.

445 In summary, our study demonstrates that there exist a functional segregation of barrel
446 and septa domains in the mouse whisker somatosensory cortex, and that is dynamically
447 shaped by the temporal dynamics of SST+ interneuron activity, imparted by the synaptic
448 protein Elfn1. This temporal divergence is essential for the integration of sensory
449 information over repeated stimulation and for the differential projection of information to
450 downstream cortical areas. These findings highlight the importance of local inhibitory
451 circuits in shaping cortical processing and offer new insights into mechanisms underlying
452 sensory perception and behavior.

454 **Acknowledgments**
455 We thank members of the Karayannis lab for suggestions and comments on the
456 manuscript. Authors also thank George Kanatouris for helping with animal handling and
457 breeding.

459 **Funding**
460 This work was supported by Swiss National Science Foundation Grant S-41260-01-01
461 (T.K) and European Research Council Grant 679175 (T.K).

463 **Author Contributions**

A.Ö.A. curated the data, analyzed in-vivo silicon probe recordings and the two-photon
calcium imaging data, performed the decoder analysis and wrote the manuscript.
T.J.S. conceptualized the study, maintained Elfn1 KO line, performed earlier
electrophysiological characterizations and contributed to the drafting of the
manuscript, J.W.Y. performed silicon probe recordings, AvdB collected two-photon
calcium imaging data, L.C. performed light-sheet data analysis, R.K. performed tissue
clearing and light-sheet image acquisition, T.K. conceptualized and supervised the study
and wrote the manuscript.
472

473 **Declaration of Interest**
474 The authors declare no competing interests.
475

476 **Methods**
477

478 RESOURCE AVAILABILITY
479 Lead Contact
480 Further information and requests for resources and reagents should be directed to the
481 Lead Contact, T.K. (karayannis@hifo.uzh.ch)
482

483 Materials Availability
484 This study did not generate new unique reagents.
485

486 Data and Code Availability
487 The datasets and analysis routines are available from the first author on reasonable
488 request.
489

**Mice**
All the animal experiments followed the guidelines of the Veterinary Office of Switzerland
and were approved by the Cantonal Veterinary Office Zurich and the University of Zurich.
husbandry with a 12-h reverse dark-light cycle (7 a.m. to 7 p.m. dark) at 24 °C and variable
humidity. Adult (5- to 10-week-old) male C57BL6J wide type mice were used for

retrograde tracing. Animal lines used in this study are VIP-IRES-Cre

(Vip[tm1(cre)Zjh]/J)(Taniguchi et al., 2011), SST-IRES-Cre (Sst[tm2.1(cre)Zjh]/J) (Taniguchi et al.,

2011), Ai14 (B6;129S6-Gt(ROSA)26Sor[tm14(CAG-tdTomato)Hze]/J)(Madisen et al., 2010) and

Elfn1 KO (Elfn1tm1(KOMP)Vlcg). Both male and female mice were used in all

experiments.

**In-Vivo Silicon Probe Recording**

We used 4 Elfn1 KO and 4 WT littermates at the age of P20 to P30 for the multi-electrode

silicon probe recordings. Mice were under urethane induced anesthesia (1.5g/1kg)

throughout the experiment. A heating pad was used to maintain the mouse's body

temperature at 37°C. The depth of anesthesia was checked with breathing speed and

paw reflexes throughout the experiment. The skull of the right hemisphere was exposed

by removing the skin on top, and a metallic head holder was implanted on the skull with

cyanoacrylate glue and dental cement. A 26G needle was used to open a ~2mm x 2mm

cranial window which exposed the S1 barrel field (wS1). Extreme care was taken not to

cause damage or surface bleeding during surgery.

wS1 neural activities were recorded with an 8-shank-64-channel silicon probe. Each of

the 8 shanks has 8 recording sites (100μm apart). The distance between each shank is

200μm (NeuroNexus Technologies, Ann Arbor, MI, USA). The silicon probe was labeled

with DiI (1,1′-dioctadecyl-3,3,3′,3′-tetramethylindocarbocyanine, Molecular Probes,

Eugene, OR, USA) and inserted perpendicularly into the barrel cortex. A silver wire was

placed into the cerebellum as a ground electrode. All data were acquired at 20 kHz and

stored with MC Rack (RRID:SCR_014955) software (Multi Channel systems). The total

duration of multi-electrode recordings varied between 3 to 5 hours. After each experiment,

the animal was deeply anesthetized by ketamine (120 mg/kg, ketamine, 50 mg/mL,

HamelnPharma, Hameln Germany) and perfused through the aorta with ringer solution.

The brain was kept in 4% PFA. Tangential sections (200-μm thick) were prepared for

vesicular glutamate transporter 2 (vGlut2, Synaptic Systems, 135404; 1:1000)

immunohistochemistry. By combining the DiI and vGlut2 staining, the insertion position of

the 8 shank probes were identified in the barrel cortex.

**Whisker Stimulation**

Either a single principal whisker (SW, single whisker) or multiple whiskers (MW, multi whisker) were stimulated 1mm from the snout in rostral-to caudal direction (about 1mm displacement) using a stainless-steel rod (1 mm diameter) connected to a miniature solenoid actuator. The movement of the tip of the stimulator bar was measured precisely using a laser micrometer (MX series, Metralight, CA, USA) with a 2500 Hz sampling rate. The stimulus takes 26ms to reach the maximal 1 mm whisker displacement, with a total duration of 60ms until it reaches baseline (Yang et al., 2017). Both SW and MW stimulation was performed at 10Hz for 2s and each stimulation repeated for 20 times (20 Trials). The inter-trial interval was 30s. In every experiment, multiple whiskers were singly stimulated to increase the probability of finding a proximal barrel-septa and neighboring barrel triplets. We first performed multi-whisker stimulation and then individual single whisker stimulations were performed at the end. Individual whisker stimulation was performed by attaching a pipet tip to the stimulator, which was then placed in contact with the principal whisker under a dissecting scope. MW stimulation was performed by contacting multiple whiskers with 1cm-wide masking tape attached to the pipet tip.

**In vivo two-photon calcium imaging**

Neuronal ensembles in superficial layers of the principal whisker barrel field mapped by intrinsic signal imaging were bolus-loaded with the AM ester form of Oregon Green BAPTA-1 by pressure injection (OGB-1; 1 mM solution in $Ca^{2+}$-free Ringer's solution; 2-min injection at 150–200 µm depth) as described previously (Stosiek et al., 2003). The craniotomy was then filled with agarose (type III-A, 1% in Ringer's solution; Sigma) and covered with an immobilized glass plate. Two-photon $Ca^{2+}$ imaging was performed with a Scientifica HyperScope two-photon laser scanning microscope one hour after bolus loading using a Ti:sapphire laser system at 900 nm excitation (Coherent Chameleon; ~120 femtosecond laser pulses). Two-channel fluorescence images of 256 × 128 pixels at 11.25 Hz (HyperScope galvo-mode) were collected with a 16x water-immersion objective lens (Nikon, NA 0.8). In all, 3–5 separate spots have been imaged per animal. Per imaging spot, 10 trials of 20s-long evoked activity recorded for each of single- and

multi- whisker stimulation paradigms. Data acquisition was controlled by ScanImage

(Pologruto et al., 2003).

**Analysis of calcium imaging data**

$Ca^{2+}$ imaging data is analyzed using custom MATLAB scripts. First, fluorescence image

time-series for a given region were concatenated. The concatenated data was then

aligned using a Fourier domain cross-correlation based subpixel registration algorithm

(Guizar-Sicairos et al., 2008) to correct for translational drift (registered on red tdTomato

channel and transferred to OGB-1 channel). Average intensity projections of the imaging

data were used as reference images to manually annotate regions of interest (ROIs)

corresponding to individual neurons. Neurons with somata partly out-of-focus were not

included. $Ca^{2+}$ signals were expressed as the mean pixel value of the relative

fluorescence change $\Delta F/F = (F-F_0)/F_0$ in each given ROI. $F_0$ was calculated as the bottom

5% of the fluorescence trace. Neuropil patches surrounding each neuron is defined by all

pixels not assigned to a neuronal soma or astrocyte of the corresponding neuron ROI

annotation (Peron et al., 2015) (A disk shaped region around the neuron of interest with

excluding any intersecting neighboring neuronal ROI). Neuropil correction is performed

as $F_{corrected} = F_{neuron} - \alpha * F_{neuropil}$. α is estimated for each imaging spot separately using the

formula $F_{blood\_vessel}/F_{surrounding\_neuropil}$ (Kerlin et al., 2010). For each stimulus, the evoked

responses of 10 trials were analyzed and the response magnitude expressed as the mean

of the evoked $\Delta F/F$ integral (%·s; integral of the first 2s response starting at stimulus

onset).

**Analysis of Silicon Probe Data**

Extracellular silicon probe data were analyzed using a custom-made Matlab script

(MATLAB (RRID:SCR_001622), Version: 2024a, MathWorks, MA, USA). The raw data

signal was band-pass filtered (0.8-5 kHz), and the multi-unit activity (MUA) was extracted

with the threshold of 7.5 times the standard deviation (SD) of baseline (Yang et al., 2017).

The current source density (CSD) map was used to identify laminar probe contact

location. The earliest CSD sink was identified as layer 4, followed by L2-3 (Reyes-Puerta

et al., 2015; Van Der Bourg et al., 2017). MUA was smoothed using a Gaussian kernel (0

mean, 5ms sigma).

**Decoder Analysis**

Recordings from the principal barrel, adjacent septa, and a neighboring unstimulated

barrel were organized into three matrices for wild-type (WT) animals: a 280 × 95 × 20

matrix for the stimulated barrel (14 Barrels, 95ms, 20 pulses), a 180 × 95 × 20 matrix for

the septa (9 Septa, 95ms, 20 pulses), and a 360 × 95 × 20 matrix for the neighboring

barrel (18 Neighboring barrels, 95ms, 20 pulses). Each matrix's dimensions correspond

to the number of recording channels, time points (95ms epochs), and 20 individual

stimulation trials, respectively. Although each trial originally spanned 2000ms (20 pulses

at 10Hz), the analysis focused on the first 95ms post-stimulation (95ms instead of

100ms). To examine the impact of different temporal windows on classification

performance, three versions of the analysis were performed by varying the time-range

parameter, 1 to 95ms for the full epoch, 1 to 50ms for the first part, and 51 to 95ms for

the second part.

For each stimulation trial, data was accumulated pulse-by-pulse. In each iteration

(ranging from 1 to 20), the corresponding segments from the three matrices were

reshaped into two-dimensional feature matrices by concatenating the selected time points

across the accumulated pulses. Labels were then assigned to the reshaped data ('B' for

the barrel, 'S' for septa, and 'N' for the neighboring barrel).

We employed a classification approach utilizing error-correcting output codes (ECOC)

models with a gentle adaptive boosting (GentleBoost) strategy to train classifiers. This

method is particularly effective in handling imbalanced data and unequal misclassification

costs. Like Logit Boost, each weak learner in the ensemble fitted a regression model to

response values, $y_n \in \{-1, +1\}$. The mean-squared error is computed as:

612
$$N \sum_{n=0}^{d} d(t)_n \big(y_n - h_t(x_n)\big)^2$$

where $d(t)_n$ represents observation weights at step t (summing up to 1) and $h_t(x_n)$

denotes predictions from the regression model $h_t$ fitted to response values $y_n$. As the

strength of individual learners diminishes, the weighted mean-squared error converges

towards 1.

Three distinct evaluations were performed. WT-to-KO Prediction, the ECOC classifier

trained on WT data (compiled from the three regions) was used to predict labels from

corresponding recordings in knockout (KO) mice. Within-Group Cross-Validation (WT), A

cross-validation procedure was applied to the WT dataset using MATLAB's *crossval*

function to assess classifier performance internally. Within-Group Cross-Validation (KO),

Similarly, an ECOC classifier was trained and cross-validated on the KO dataset.

For each evaluation, confusion matrices were generated and summary accuracy statistics

computed. Accuracy was tracked as a function of the number of accumulated pulses

(from 1 to 20). The accuracy value represents the average per-class accuracy. In other

words, for each class the accuracy is defined as: $Accuracy_{class}$ = (TP + TN) / Total

samples, where for that class, TP (True Positives) is the number of correctly predicted

samples (the corresponding diagonal element of the confusion matrix), and TN (True

Negatives) is the sum of all correct predictions for samples not in that class (i.e., all the

entries outside the row and column of that class). The function first computes the

confusion matrix C where rows represent actual classes and columns represent predicted

classes. It then calculates the total number of samples as the sum of all entries in C, and

it determines the number of classes. The overall accuracy reported is computed via the

formula:

Accuracy = (2*trace(C)+2*Total)/((Number of Classes)*Total)

Here, trace(C) is the sum of the diagonal elements, which means it is the sum of the true

positives for all classes, and *Total* is the sum of all entries in the confusion matrix.

Although this formula may look a bit unusual, it is designed to yield the average of the

per-class accuracies. In practice, for each class, one would compute (TP + TN) / Total

and then average these values over the classes that reflects how well the classifier

performed across all classes. To characterize the evolution of classification accuracy with

increasing pulse number, non-linear exponential functions of the form:

$$f(x) = a - b * \exp(-cx)$$

were fitted to the accuracy data using a non-linear least squares approach (implemented

via the *fminspleas* function). Separate fits were obtained for the WT-to-KO prediction, WT

cross-validation, and KO cross-validation accuracy curves. The fitted curves and raw
accuracy data were then plotted. This integrated analysis pipeline allowed for systematic
comparison of classification performance across spatial locations and conditions and an
assessment of the temporal dynamics underlying the neural responses to single-whisker
stimulation. The same strategy has been used for multi-whisker decoding but between
only Barrel and Septa.
652

**Retrograde tracing**
Mice aged from P10-37 were anesthetized by isoflurane and a small craniotomy was
made above the injection site (M1 or S2).  200nl retrograde AAV virus, ssAAV-retro/2-
CAG-EGFP-WPRE-SV40p(A) (physical titer: $5X10^{12}$ vg/ml) was injected 200-500μm
deep in the M1 or S2. Virus was purchased from Zurich Viral Vector Facility (Cat#: V24-
retro).
659

**Tissue Processing for Passive Clearing and Imaging**
The method used for hydrogel-based tissue clearing is explained in detail elsewhere
(Chung and Deisseroth, 2013). Briefly,  Mice age P29-70 (4 weeks after retrograde AAV
virus injection) or SST-Ai14 and VIP-Ai14 mice age from P14-77 were transcardially
perfused using 1x PBS and Hydrogel solution (1% PFA, 4% Acrylamide, 0.05% Bis). The
collected brains were post-fixed for 48 h in a Hydrogel solution (1% PFA, 4% Acrylamide,
0.05% Bis). Afterward, the Hydrogel polymerization was induced at 37 °C. Following the
polymerization, the brains were immersed in 40 mL of 8% SDS and kept shaking at room
temperature until the tissue was cleared sufficiently (15-90 days depending on the
animal's age). Finally, after 2–4 washes in PBS, the brains were put into a self-made
refractive index matching solution (RIMS). They were left to equilibrate in 5 mL of RIMS
for at least 4 days at RT (Room Temperature) before being imaged. After clearing, brains
were attached to a small weight and loaded into a quartz cuvette, then submerged in
RIMS, and imaged using a home-built mesoscale selective plane illumination microscope
(mesoSPIM, Voigt et al., 2019). The microscope consists of a dual-sided excitation path
using a fiber-coupled multiline laser combiner (405, 488, 515, 561, 594, 647 nm, Omicron
SOLE-6) and a detection path comprising an Olympus MVX-10 zoom macroscope with a

1× objective (Olympus MVPLAPO 1x), a filter wheel (Ludl 96A350), and a scientific CMOS

(sCMOS) camera (Hamamatsu Orca Flash 4.0 V3). For imaging tdTomato and eGFP a

594 nm excitation with a 594 long-pass filter (594 LP Edge Basic, AHF) and 488 nm &

520/35 (BrightLine HC, AHF) were used, respectively. The excitation paths also contain

galvo scanners (GCM-2280-1500, Citizen Chiba) for light-sheet generation and reduction

of streaking artifacts due to absorption of the light-sheet. In addition, the beam waist is

scanning using electrically tunable lenses (ETL, Optotune EL-16-40-5D-TC-L)

synchronized with the rolling shutter of the sCMOS camera. This axially scanned light-

sheet mode (ASLM) leads to a uniform axial resolution across the field-of-view of 5–10 µm

(depending on zoom and wavelength). The magnification 0.8X (pixel size: 8.23 µm) was

used for the view of the whole brain and 2X (pixel size: 3.26 µm) was used for the view

of barrel cortex.  Further technical details of the mesoSPIM are described elsewhere

(Vladimirov et al., 2023; Voigt et al., 2019).

**Analysis of cleared brain imaging data**

Barrel cortex was imaged in 3D with 2X objective. The images were processed through

Fiji and Imaris to visualize the 3D structure of the barrel cortex. XY-, YZ-, XZ- projections

were visualized in Imaris, which allows us to adjust the correct angle for each barrel

column to be vertical against XY-plane. This method allowed us identify barrel and septa

in layers 2-3 and 5 and layer 4. In addition, the depth information is also precisely acquired

for each barrel column. SST+ or VIP+ neuron numbers, S2-projected and M1-projected

S1 neuron numbers were counted in icy. Max projection for each 50 ums along barrel

columns was acquired and neurons on each max projection image were counted

automatically by icy (RRID:SCR_010587) with the same threshold for all the brains. Each

barrel was drawn according to the barrel map from each brain by max projection. The

area of septa was calculated by subtracting the whole selected area from all the selected

barrel areas.

**t-SNE Analysis**

To visualize the segregation of neural responses across the different cortical domains,

we applied t-distributed Stochastic Neighbor Embedding (t-SNE) to the averaged

datasets. t-SNE was used to reduce the high-dimensional data into a 2D space for
visualization. The parameters for t-SNE were set to use the *barneshut* algorithm for
efficiency, with a perplexity of 9 and exaggeration of 3 to enhance cluster separation. The
resulting t-SNE coordinates were plotted using MATLAB's *gscatter* function to distinguish
between the different domains (Barrel, Septa, Neighboring). This analysis allowed us to
visually inspect the clustering of neural responses from different cortical domains.

**Statistical Analysis of the Multi-Unit Activity**
To evaluate the statistical significance of differences in multi-unit activity (MUA)
responses between barrel (B), septa (S), and neighboring barrel (N) domains across 20
pulses of whisker stimulation, we performed a pulse-by-pulse comparison using the non-
parametric Mann-Whitney U test (also known as the Wilcoxon rank-sum test). This
analysis was conducted on the averaged MUA data recorded from Layer 4 of the whisker
somatosensory cortex (wS1) in wild-type (WT) and Elfn1 knockout (KO) mice during
single-whisker stimulation (SWS) and multi-whisker stimulation (MWS) paradigms, as
detailed above.  For each of the 20 stimulation pulses (1 to 20), the MUA data for each
domain were reshaped and averaged across trials. The Mann-Whitney U test was then
applied pairwise to assess differences between domains: barrel vs. septa (B vs S), barrel
vs. neighboring barrel (B vs N), and septa vs. neighboring barrel (S vs N). The resulting
p-values were stored in a 3×20 matrix, where rows correspond to the pairwise
comparisons and columns represent the pulse number.  To visualize the statistical
outcomes, a heatmap was generated using MATLAB (Version 2024a, MathWorks, MA,
USA). A custom discrete colormap was created by adapting the hot colormap, which
originally provides a gradient of 10 colors ranging from black (low values) to white (high
values). We selected specific indices from this gradient (10, 8, 6, 4, 2, 1) to define a 10-
level discrete colormap, assigning the darkest shade (black) to p-values ≤ 0.05 (indicating
statistical significance) and progressively lighter shades to higher p-values, with the
lightest shade (white) representing p-values > 0.05. The colormap was inverted to ensure
that significant p-values (≤ 0.05) appeared as light shades against a dark background,
enhancing visual contrast.  The heatmap was plotted with p-values scaled between 0 and
0.05, using the custom colormap.

 **Statistical Analysis**

 Data are represented as mean ± SEM unless stated otherwise. Statistical comparisons

 have been done using a one-tailed Mann-Whitney-U test for non-paired data and one-

 tailed Wilcoxon signed-rank test for paired comparisons (the statistical significance

 threshold was set to $p < 0.05$; in the Figures, different degrees of evidence against the null

 hypothesis are indicated by asterisks ($p < 0.05$: *; $p < 0.01$: **; $p < 0.001$: ***). Cell densities

 in Figures 2 and 6 were compared using repeated measures ANOVA. All tests were

 conducted using custom codes in Matlab.

 **Figure Legends**

 **Figure 1.** Temporal divergences emerge between barrel and septa domains upon

 repeated whisker stimulation in Layer 4 of mouse wS1. (A) Either principal whiskers

 (SWS, or single whisker) or most of the principal whiskers (MWS, or multi whisker) were

 stimulated for 2s at 10Hz (20 pulses) for 20 times (20 trials). (B) Acute simultaneous *in-*

 *vivo* silicon probe (8x8) recordings were performed from barrel and septa domains after

 the start of active whisking (>P21) (C) t-SNE analysis of the average firing dynamics from

 three domains after averaging 20 single trials. (D) Left: Average multi-unit firing profiles

 recorded at Layer 4 of barrel (black), unstimulated neighbor (blue) and septa (pink)

 columns upon 2s-long 10Hz repeated single whisker stimulation. Middle: Matrix

 representation of the data. Right: Pair-wise statistical comparisons of first 50ms of the

 responses (yellow line represents the cut off) of three conditions for all pulses (Mann-

 Whitney-U-Test). (E) Same with E but response profiles upon MW stimulation. (F) Top:

 Temporal ratio dynamics (average AUC MW responses divided by average AUC SW

 responses) for barrel and septa. Bottom: statistical comparison of average AUC ratios of

 all pulses (1-20), first six (1-6) or last 14 pulses (7-20), respectively. (p values are color

 coded in D, stars represents *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$ in E and F.)

 **Figure 2.** SST+ and VIP+ neuron densities differ at Barrel and Septa Domains in Layer

 4. (A) A passive CLARITY-based tissue clearing protocol was performed on the collected

 brains which were subsequently imaged in their entirety using a custom-built light-sheet

 microscope (mesoSPIM). 561nm wavelength was used to image tdTomato, 488nm

wavelength was used to image autofluorescence. (B) 3D reconstruction of cleared SST-
Ai14 (left) and VIP-Ai14 (right) barrel cortex. X-Y, X-Z and Y-Z views are presented
focusing with a single barrel max projection. The brain was rotated to achieve the
orientation that the barrel columns are perpendicular to the X-Y view. The scale bars are
set for 500μm. (C) Max projection examples of SST (red) and VIP (blue) neurons in wS1
L2-3, L4 and L5. Quantified barrels (yellow lines) and septa (green lines) are circled out
which are perpendicular to X-Y (top-down) view. The scale bar is set for 200μm. (D) SST+
neuron density distribution over laminae for barrel and septal domains (N=4 mice, n=54
Barrels and surrounding septa per barrel). Normalized cell densities plotted as a function
of depth. 0μm represents the top of layer 4. SST+ interneuron density is significantly
higher in septa (p=0.0015, F=30.64, repeated-measures ANOVA). (E) VIP+ neuron
density distribution over laminae for barrel and septal domains. (N=4 mice, n=53 Barrels
and surrounding septa per barrel). Normalized cell densities plotted as a function of depth.
0μm represents the top of layer 4. VIP+ interneuron density is also significantly higher in
septa (p=0.021, F=9.63, repeated-measures ANOVA).

**Figure 3**. Single- vs multi- whisker responses of SST+ and VIP+ Interneurons (A)
Interneurons are labeled with tdTomato using reporter mouse lines and Ca$^{2+}$ imaging has
been performed after bulk loading of OGB-1. scale bar 35μm. (B) Schematic
representation of the single- and multi- whisker stimulation protocol upon 10Hz
stimulation. (C) $\Delta F/F$ traces of evoked activity of SST (on top) and VIP (on bottom)
interneurons ($N$ = 3 animals per group, VIP P21+: 138 cells, SST P21+: 51 cells). (D) Area
under the curve (AUC) of the first 2s of the $\Delta F/F$ traces of all SST and VIP cells upon
single- (SW: 2s-long principal whisker evoked activity) and multi- whisker (MW: 2s-long
multi whisker stimulation evoked activity) stimulation (Wilcoxon signed-rank test, Base:
2s-long baseline activity).

**Figure 4.** Loss of Elfn1 abolishes barrel-septa response divergences upon single-whisker
stimulation. (A) t-SNE analysis of the average firing dynamics from three domains. (B)
Left: Average multi-unit firing profiles recorded at Layer 4 of barrel (gray), unstimulated
neighbor (light blue) and septa (purple) columns (11-barrel columns, 7 septal columns,

11 unstimulated neighbors from N=4 mice) upon 2s-long 10Hz repeated single whisker

stimulation. Middle: Matrix representation of the same data by representing each

stimulation pulse response on one row. Bottom: Pair-wise statistical comparisons of three

conditions for all pulses (Mann-Whitney-U-Test). (C) Same with but response profiles

upon MW stimulation. (D) Top: Ratio of MW/SW dynamics for barrel and septa. Bottom:

statistical comparison of average ratios of 20 average of all pulses, first six or last 14

pulses, respectively. (p values are color coded in B, stars represents *: p<0.05; **: p<0.01;

***: p<0.001 in C and D.)

**Figure 5.** Accumulative decoding analysis shows an alteration of columnar domain

identity in Elfn1 KO animals. (A) For the decoding analysis, the entire pulse response,

only the first 50ms or only the last 45ms has been used. (B) Decoder analysis of the full

pulse response profiles in L4 upon SWS. (C) Decoder analysis of the first part of the pulse

response (1-50ms) profiles in L4 upon SWS.  (D) Decoder analysis of the second part of

the pulse response (51-95ms) profiles in L4 upon SWS. (E) Decoder analysis of the full

pulse response profiles in L4 upon MWS. (F) Decoder analysis of the first part of the pulse

response (1-50ms) profiles in L4 upon MWS (G) Decoder analysis of the second part of

the pulse response (51-95ms) profiles in L4 upon MWS.

**Figure 6.** wS1 barrel and septa columns differentially project to wS2 and M1 (A) The

schematics of the experimental protocol. The retro-AAV-CAG-GFP was injected in either

S2, or M1 at P20-30 and the brains were dissected 4 weeks after the virus injection

following the tissue clearing and whole brain imaging. (B) 3D reconstruction of the cleared

brain with S2(Left panels) or M1(right panels) injection. X-Y, X-Z and Y-Z views are

presented focus with a single barrel max projection. The brain was rotated to achieve the

orientation that the septa is perpendicular to the X-Y view (top-down). Scale bars in whole

brain view are set for 2000um and the scale bars in X-Y, X-Z and Y-Z views are set for

200um. (C) Max projection examples of S2-projected wS1 neurons and M1-projected

wS1 neurons in L2-3, L4 and L5 in barrel vs septa. Quantified barrels (yellow lines) and

septa (blue lines) are circled out which are perpendicular to X-Y (top-down) view. Scale

bars are set for 200um. (D) Left: Normalized cell density profiles of S2 projection neurons

at barrel and septa columns. 0μm represents the top of layer 4. L4 and L5a septa columns

send significantly more projections to S2 than barrel column (N=4 mice) (p=0.0139,

F=11.79, repeated-measures ANOVA). Right: Normalized cell density profiles of M1

projections neurons at barrel and septa columns. 0μm represents the top of layer 4. L3

and L4 septa columns send significantly more projections to M1 (N=6 mice) (p=0.0032,

F=14.82, repeated measures ANOVA). (E) Left: Normalized cell density profiles of S2

and M1 projection neurons at barrel columns. 0μm represents the top of L4. L3 barrel

columns send significantly more projections to S2 (N=4 mice with S2 injections, N=6 mice

with M1 injections) (p=0.0293, F=7.02, repeated-measures ANOVA). Right: Normalized

cell density profiles of S2 and M1 projections neurons at septa columns. 0μm represents

the top of L4. L5a septal columns send significantly more projections to M1 N=4 mice with

S2 injections, N=6 mice with M1 injections) (p=0.0402, F=5.98, repeated measures

ANOVA).

**Supplementary Figure 1.** An example of assignment of barrel, septa and Neighbor

electrodes using histology and current source density (CSD) analysis. (A) Shank

electrode map of 8x8 silicon probes. (B) Probe locations were labeled (DiI) after each

experiment using histology (vGlut2 staining). Matching shank locations are indicated

using yellow numbers. Blue dots represent the insertion sites. The red and white text

depicts corresponding barrel names (C) All principal whiskers are stimulated one-by-one.

C1 to C5 barrel responses upon single whisker stimulation are shown from top to bottom.

(D) A closer look on C1, C2 and C3 whisker stimulations and corresponding CSD at

shanks 5,6,7 and 8. (E), Barrel (B), adjacent septa (S) and the adjacent neighboring barrel

(N) were assigned according to the probe insertion sites and L4 barrel autofluorescence

in combination with CSD responses. For example, while shanks 6 and 8 show strong

response upon single whisker stimulation (and hence assigned as barrels), shank 7 does

not show the expected response (hence assigned as septa). Only the triplets of

electrodes were used in the analysis in which principal barrel (B), adjacent septa (S) and

the adjacent neighboring barrel (N) were captured by the probe insertion sites for any

experiment (Left). 14-barrel columns, 9 septal columns, 18 unstimulated neighboring

barrels from N=4 mice. (F) An ideal schematic representation (on the left). Realistically,

we cannot always be sure if an entire shank is consistently going through a barrel or
septal column due to the curvature of the cortex (on the right) for above and below of
layer4.

**Supplementary Figure 2.** Representation of the conversion of the two-second-long
single trial recordings into 20x100ms matrices. (A) Schematic representation. (B) An
example data. The last 5ms of each pulse have been discarded in further analysis.

**Supplementary Figure 3.** (A) Left: Average multi-unit firing profiles recorded at Layer 2/3
of barrel (black), unstimulated neighbor (blue) and septa (pink) columns upon 2s-long
10Hz repeated single whisker stimulation. Right: Another representation of the same data
but converted into an image by representing each stimulation pulse response on one row.
(B) Left: Average multi-unit firing profiles recorded at Layer 2/3 of barrel (black),
unstimulated neighbor (blue) and septa (pink) columns upon 2s-long 10Hz repeated multi
whisker stimulation. Right: Another representation of the same data but converted into an
image by representing each stimulation pulse response on one row. (C) Left: Temporal
ratio dynamics (average AUC MW responses divided by average AUC SW responses)
for barrel and septa for layer2/3. Right: statistical comparison of average AUC ratios of
all pulses (1-20), first six (1-6) or last 14 pulses (7-20), respectively (stars represents *:
$p<0.05$; **: $p<0.01$; ***: $p<0.001$ in F and G. Wilcoxon rank sum test).

**References**

Ahissar E, Sosnik R, Bagdasarian K, Haidarliu S. 2001. Temporal frequency of whisker
movement. II. Laminar organization of cortical representations. *J Neurophysiol* **86**.
doi:10.1152/jn.2001.86.1.354
Alloway KD. 2008. Information processing streams in rodent barrel cortex: The differential
functions of barrel and septal circuits. *Cerebral Cortex*. doi:10.1093/cercor/bhm138
Allwein EL, Schapire RE, Singer Y. 2001. Reducing multiclass to binary: A unifying
approach for margin classifiers. *Journal of Machine Learning Research* **1**.
Almási Z, Dávid C, Witte M, Staiger JF. 2019. Distribution patterns of three molecularly
defined classes of gabaergic neurons across columnar compartments in mouse
barrel cortex. *Front Neuroanat* **13**. doi:10.3389/fnana.2019.00045
Armstrong-James M, Fox K. 1987. Spatiotemporal convergence and divergence in the
rat S1 "Barrel" cortex. *Journal of Comparative Neurology* **263**.
doi:10.1002/cne.902630209

Audette NJ, Urban-Ciecko J, Matsushita M, Barth AL. 2018. POm Thalamocortical Input
Drives Layer-Specific Microcircuits in Somatosensory Cortex. *Cereb Cortex* **28**.
doi:10.1093/cercor/bhx044
Brecht M, Sakmann B. 2002. Dynamic representation of whisker deflection by synaptic
potentials in spiny stellate and pyramidal cells in the barrels and septa of layer 4 rat
somatosensory cortex. *Journal of Physiology*. doi:10.1113/jphysiol.2002.018465
Bureau I, Von Paul F Saint, Svoboda K. 2006. Interdigitated paralemniscal and lemniscal
pathways in the mouse barrel cortex. *PLoS Biol* **4**. doi:10.1371/journal.pbio.0040382
Cai L, Yang JW, Wang CF, Chou SJ, Luhmann HJ, Karayannis T. 2022. Identification of
a Developmental Switch in Information Transfer between Whisker S1 and S2 Cortex
in Mice. *Journal of Neuroscience* **42**. doi:10.1523/JNEUROSCI.2246-21.2022
Carvell GE, Simons DJ. 1990. Biometric analyses of vibrissal tactile discrimination in the
rat. *Journal of Neuroscience* **10**. doi:10.1523/jneurosci.10-08-02638.1990
Chakrabarti S, Alloway KD. 2006. Differential origin of projections from SI barrel cortex to
the whisker representations in SII and MI. *Journal of Comparative Neurology* **498**.
doi:10.1002/cne.21052
Chmielowska J, Carvell GE, Simons DJ. 1989. Spatial organization of thalamocortical and
corticothalamic projection systems in the rat SmI barrel cortex. *Journal of
Comparative Neurology* **285**. doi:10.1002/cne.902850304
Chung K, Deisseroth K. 2013. CLARITY for mapping the nervous system. *Nat Methods*
**10**. doi:10.1038/nmeth.2481
Douglas RJ, Martin KAC. 2004. Neuronal circuits of the neocortex. *Annu Rev Neurosci*.
doi:10.1146/annurev.neuro.27.070203.144152
Estebanez L, Bertherat J, Shulz DE, Bourdieu L, Léger JF. 2016. A radial map of multi-
whisker correlation selectivity in the rat barrel cortex. *Nat Commun* **7**.
doi:10.1038/ncomms13528
Friedman J, Hastie T, Tibshirani R. 2000. Additive logistic regression: A statistical view of
boosting. *Ann Stat*. doi:10.1214/aos/1016218223
Grant RA, Mitchinson B, Prescott TJ. 2012. The development of whisker control in rats in
relation to locomotion. *Dev Psychobiol* **54**. doi:10.1002/dev.20591
Guizar-Sicairos M, Thurman ST, Fienup JR. 2008. Efficient subpixel image registration
algorithms. *Opt Lett* **33**. doi:10.1364/ol.33.000156
Karnani MMM, Jackson J, Ayzenshtat I, Tucciarone J, Manoocheri K, Snider WGG, Yuste
R. 2016. Cooperative Subnetworks of Molecularly Similar Interneurons in Mouse
Neocortex. *Neuron* **90**. doi:10.1016/j.neuron.2016.02.037
Kastli R, Vighagen R, van der Bourg A, Argunsah AÖ, Iqbal A, Voigt FF, Kirschenbaum
D, Aguzzi A, Helmchen F, Karayannis T. 2020. Developmental divergence of sensory
stimulus representation in cortical interneurons. *Nat Commun* **11**.
doi:10.1038/s41467-020-19427-z
Kerlin AM, Andermann ML, Berezovskii VK, Reid RC. 2010. Broadly Tuned Response
Properties of Diverse Inhibitory Neuron Subtypes in Mouse Visual Cortex. *Neuron*
**67**. doi:10.1016/j.neuron.2010.08.002
Kheradpezhouh E, Adibi M, Arabzadeh E. 2017. Response dynamics of rat barrel cortex
neurons to repeated sensory stimulation. *Sci Rep* **7**. doi:10.1038/s41598-017-11477-
6

Kim U, Ebner FF. 1999. Barrels and septa: Separate circuits in rat barrel field cortex.
*Journal of Comparative Neurology* **408**. doi:10.1002/(SICI)1096-
9861(19990614)408:4<489::AID-CNE4>3.0.CO;2-E
Koralek KA, Jensen KF, Killackey HP. 1988. Evidence for two complementary patterns of
thalamic input to the rat somatosensory cortex. *Brain Res* **463**. doi:10.1016/0006-
8993(88)90408-8
Lee S, Kruglikov I, Huang ZJ, Fishell G, Rudy B. 2013. A disinhibitory circuit mediates
motor integration in the somatosensory cortex. *Nat Neurosci* **16**.
doi:10.1038/nn.3544
Ma Y, Hu H, Berrebi AS, Mathers PH, Agmon A. 2006. Distinct subtypes of somatostatin-
containing neocortical interneurons revealed in transgenic mice. *Journal of
Neuroscience* **26**. doi:10.1523/JNEUROSCI.0661-06.2006
Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD,
Hawrylycz MJ, Jones AR, Lein ES, Zeng H. 2010. A robust and high-throughput Cre
reporting and characterization system for the whole mouse brain. *Nat Neurosci*
**13**:133–140. doi:10.1038/nn.2467
Meier AM, Wang Q, Ji W, Ganachaud J, Burkhalter A. 2021. Modular network between
postrhinal visual cortex, amygdala, and entorhinal cortex. *Journal of Neuroscience*
**41**. doi:10.1523/JNEUROSCI.2185-20.2021
Melzer P, Champney GC, Maguire MJ, Ebner FF. 2006a. Rate code and temporal code
for frequency of whisker stimulation in rat primary and secondary somatic sensory
cortex. *Exp Brain Res* **172**:370–386. doi:10.1007/s00221-005-0334-1
Melzer P, Sachdev RNS, Jenkinson N, Ebner FF. 2006b. Stimulus frequency processing
in awake rat barrel cortex. *Journal of Neuroscience* **26**.
doi:10.1523/JNEUROSCI.2620-06.2006
Nomura S, Itoh K, Sugimoto T, Yasui Y, Kamiya H, Mizuno N. 1986. Mystacial vibrissae
representation within the trigeminal sensory nuclei of the cat. *Journal of Comparative
Neurology* **253**. doi:10.1002/cne.902530110
O'Connor DH, Hires SA, Guo Z V., Li N, Yu J, Sun QQ, Huber D, Svoboda K. 2013.
Neural coding during active somatosensation revealed using illusory touch. *Nat
Neurosci* **16**. doi:10.1038/nn.3419
Pammer L, O'Connor DH, Hires SA, Clack NG, Huber D, Myers EW, Svoboda K. 2013.
The mechanical variables underlying object localization along the axis of the whisker.
*Journal of Neuroscience* **33**. doi:10.1523/JNEUROSCI.4316-12.2013
Peron S, Chen TW, Svoboda K. 2015. Comprehensive imaging of cortical networks. *Curr
Opin Neurobiol*. doi:10.1016/j.conb.2015.03.016
Petersen CCH. 2007. The functional organization of the barrel cortex. *Neuron*.
doi:10.1016/j.neuron.2007.09.017
Pi HJ, Hangya B, Kvitsiani D, Sanders JI, Huang ZJ, Kepecs A. 2013. Cortical
interneurons that specialize in disinhibitory control. *Nature* **503**.
doi:10.1038/nature12676
Pologruto TA, Sabatini BL, Svoboda K. 2003. ScanImage: Flexible software for operating
laser scanning microscopes. *Biomed Eng Online* **2**. doi:10.1186/1475-925X-2-13
Reyes-Puerta V, Sun JJ, Kim S, Kilb W, Luhmann HJ. 2015. Laminar and Columnar
Structure of Sensory-Evoked Multineuronal Spike Sequences in Adult Rat Barrel
Cortex in Vivo. *Cerebral Cortex* **25**. doi:10.1093/cercor/bhu007

Rice FL. 1985. An attempt to find vibrissa-related barrels in the primary somatosensory
cortex of the cat. *Neurosci Lett* **53**. doi:10.1016/0304-3940(85)90180-6
Rice FL, Gomez C, Barstow C, Burnet A, Sands P. 1985. AComparative analysis of the
development of the primary somatosensory cortex: Interspecies similarities during
barrel and laminar development. *Journal of Comparative Neurology* **236**.
doi:10.1002/cne.902360405
Roy NC, Bessaih T, Contreras D. 2011. Comprehensive mapping of whisker-evoked
responses reveals broad, sharply tuned thalamocortical input to layer 4 of barrel
cortex. *J Neurophysiol* **105**. doi:10.1152/jn.00939.2010
Sato TR, Gray NW, Mainen ZF, Svoboda K. 2007. The functional microarchitecture of the
mouse barrel cortex. *PLoS Biol* **5**. doi:10.1371/journal.pbio.0050189
Sato TR, Svoboda K. 2010. The functional properties of barrel cortex neurons projecting
to the primary motor cortex. *Journal of Neuroscience* **30**.
doi:10.1523/JNEUROSCI.3774-09.2010
Shepherd GMG, Svoboda K. 2005. Laminar and columnar organization of ascending
excitatory projections to layer 2/3 pyramidal neurons in rat barrel cortex. *Journal of
Neuroscience* **25**. doi:10.1523/JNEUROSCI.1173-05.2005
Silberberg G, Markram H. 2007. Disynaptic Inhibition between Neocortical Pyramidal
Cells Mediated by Martinotti Cells. *Neuron* **53**. doi:10.1016/j.neuron.2007.02.012
Sosnik R, Haidarliu S, Ahissar E. 2001. Temporal frequency of whisker movement. I.
Representations in brain stem and thalamus. *J Neurophysiol* **86**.
doi:10.1152/jn.2001.86.1.339
Stachniak TJ, Kastli R, Hanley O, Argunsah AÖ, Van Der Valk EGT, Kanatouris G,
Karayannis T. 2021. Postmitotic prox1 expression controls the final specification of
cortical VIP interneuron subtypes. *Journal of Neuroscience* **41**.
doi:10.1523/JNEUROSCI.1021-21.2021
Stachniak TJ, Sylwestrak EL, Scheiffele P, Hall BJ, Ghosh A. 2019. Elfn1-induced
constitutive activation of mglur7 determines frequency-dependent recruitment of
somatostatin interneurons. *Journal of Neuroscience* **39**.
doi:10.1523/JNEUROSCI.2276-18.2019
Staiger JF, Petersen CCH. 2021. Neuronal circuits in barrel cortex for whisker sensory
perception. *Physiol Rev* **101**. doi:10.1152/physrev.00019.2019
Stosiek C, Garaschuk O, Holthoff K, Konnerth A. 2003. In vivo two-photon calcium
imaging of neuronal networks. *Proc Natl Acad Sci U S A* **100**.
doi:10.1073/pnas.1232232100
Tan Z, Hu H, Huang ZJ, Agmon A. 2008. Robust but delayed thalamocortical activation
of dendritic-targeting inhibitory interneurons. *Proc Natl Acad Sci U S A* **105**.
doi:10.1073/pnas.0710628105
Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, Kvitsani D, Fu Y, Lu J, Lin Y, Miyoshi
G, Shima Y, Fishell G, Nelson SB, Huang ZJ. 2011. A Resource of Cre Driver Lines
for Genetic Targeting of GABAergic Neurons in Cerebral Cortex. *Neuron* **71**:995–
1013. doi:10.1016/j.neuron.2011.07.026
Tanke N, Borst JGG, Houweling AR. 2018. Single-cell stimulation in barrel cortex
influences psychophysical detection performance. *Journal of Neuroscience* **38**.
doi:10.1523/JNEUROSCI.2155-17.2018

Tomioka NH, Yasuda H, Miyamoto H, Hatayama M, Morimura N, Matsumoto Y, Suzuki
T, Odagawa M, Odaka YS, Iwayama Y, Won Um J, Ko J, Inoue Y, Kaneko S, Hirose
S, Yamada K, Yoshikawa T, Yamakawa K, Aruga J. 2014. Elfn1 recruits presynaptic
mGluR7 in trans and its loss results in seizures. *Nat Commun* **5**.
doi:10.1038/ncomms5501
Urban-Ciecko J, Barth AL. 2016. Somatostatin-expressing neurons in cortical networks.
*Nat Rev Neurosci* **17**. doi:10.1038/nrn.2016.53
Van Der Bourg A, Yang JW, Reyes-Puerta V, Laurenczy B, Wieckhorst M, Stüttgen MC,
Luhmann HJ, Helmchen F. 2017. Layer-Specific Refinement of Sensory Coding in
Developing Mouse Barrel Cortex. *Cerebral Cortex* **27**. doi:10.1093/cercor/bhw280
Van Der Loos H, Woolsey TA. 1973. Somatosensory cortex: Structural alterations
following early injury to sense organs. *Science (1979)* **179**.
doi:10.1126/science.179.4071.395
Vladimirov N, Voigt F, Naert T, Araujo GR, Cai R, Reuss AM, Zhao S, Schmid P,
Hildebrand S, Schaettin M, Groos D, Mateos JM, Bethge P, Yamamoto T, Aerne V,
Roebroeck A, Ertürk A, Aguzzi A, Ziegler U, Stoeckli ET, Baudis L, Lienkamp SS,
Helmchen F. 2023. The Benchtop mesoSPIM: a next-generation open-source light-
sheet microscope for large cleared samples. *bioRxiv*.
Voigt FF, Kirschenbaum D, Platonova E, Pagès S, Campbell RAA, Kastli R, Schaettin M,
Egolf L, van der Bourg A, Bethge P, Haenraets K, Frézel N, Topilko T, Perin P, Hillier
D, Hildebrand S, Schueth A, Roebroeck A, Roska B, Stoeckli ET, Pizzala R, Renier
N, Zeilhofer HU, Karayannis T, Ziegler U, Batti L, Holtmaat A, Lüscher C, Aguzzi A,
Helmchen F. 2019. The mesoSPIM initiative: open-source light-sheet microscopes
for imaging cleared tissue. *Nat Methods* **16**. doi:10.1038/s41592-019-0554-0
Wang HC, LeMessurier AM, Feldman DE. 2022. Tuning instability of non-columnar
neurons in the salt-and-pepper whisker map in somatosensory cortex. *Nat Commun*
**13**. doi:10.1038/s41467-022-34261-1
Woolsey TA, Welker C, Schwartz RH. 1975. Comparative anatomical studies of the Sml
face cortex with special reference to the occurrence of "barrels" in layer IV. *Journal*
*of Comparative Neurology* **164**. doi:10.1002/cne.901640107
Xu H, Jeong HY, Tremblay R, Rudy B. 2013. Neocortical Somatostatin-Expressing
GABAergic Interneurons Disinhibit the Thalamorecipient Layer 4. *Neuron* **77**.
doi:10.1016/j.neuron.2012.11.004
Yamashita T, Pala A, Pedrido L, Kremer Y, Welker E, Petersen CCH. 2013. Membrane
potential dynamics of neocortical projection neurons driving target-specific signals.
*Neuron* **80**. doi:10.1016/j.neuron.2013.10.059
Yang JW, Prouvot PH, Reyes-Puerta V, Stüttgen MC, Stroh A, Luhmann HJ. 2017.
Optogenetic modulation of a minor fraction of parvalbumin-positive interneurons
specifically affects spatiotemporal dynamics of spontaneous and sensory-evoked
activity in mouse somatosensory cortex in vivo. *Cerebral Cortex* **27**.
doi:10.1093/cercor/bhx261
Young TR, Yamamoto M, Kikuchi SS, Yoshida AC, Abe T, Inoue K, Johansen JP, Benucci
A, Yoshimura Y, Shimogori T. 2023. Thalamocortical control of cell-type specificity
drives circuits for processing whisker-related information in mouse barrel cortex. *Nat*
*Commun* **14**. doi:10.1038/s41467-023-41749-x
1079

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Figure 1

Supplementary Figure 2

**A** Layer$_{2/3}$ Single Whisker Stimulation

Barrel

Septa

2s

**B** Layer$_{2/3}$ Multi Whisker Stimulation

Barrel

Septa

2s

Pulse Number

Time (ms)

**C** Layer$_{2/3}$ Multi/Single

Barrel WT
Septa WT

AUC$_{MW}$/AUC$_{SW}$

Pulse Number

**1-20**

****

**1-6**

*

**7-20**

***

AUC(MW/SW)%

B    S

Supplementary Figure 3