Full Title:

Progressive engagement of SST+ interneurons via Elfn1 regulates the barrel-septa response identity in the somatosensory cortex of mice

Short Title:

Progressive engagement of SST+ neurons regulate barrel-septa identity

Authors:

Ali Özgür Argunşah[1,2], Tevye Jason Stachniak[1,2,3], Jenq-Wei Yang[1,2], Linbi Cai[1,2], George Kanatouris[1,2], Theofanis Karayannis[1,2,*]

[1]Laboratory of Neural Circuit Assembly, Brain Research Institute, University of Zurich Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.
[2]Neuroscience Center Zurich, University of Zurich and ETH Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.
[3]Division of Biomedical Sciences, Faculty of Medicine, Memorial University of Newfoundland, St. John's, A1B 3V6, Canada.

[*]**Correspondence**: karayannis@hifo.uzh.ch

Abstract

The vibrissae system of rodents, akin to human hands and fingers, provides somatosensory information coming from individual whiskers for object exploration and recognition. Just as separated digits enhance somatosensation in humans, the ability of mice to sense objects through multiple whiskers in segregated streams is crucial. The segregation begins at the level of the whiskers and is maintained through their precise somatotopic organization in the Brainstem→ Thalamus→ Cortex axis, culminating in the so-called barrels and the in-between "spaces" called septa. Here, by performing *in-vivo*

silicon probe recordings simultaneously in the barrel and septa domains in mice upon
repeated 10Hz single and multi-whisker stimulation, we identify and characterize a
temporal divergence in the spiking activity between these domains. Further, through
genetic fate-mapping, we reveal that cortical SST+ and VIP+ inhibitory neurons show a
layer-dependent differential preference in septa versus barrel domains. Utilizing a genetic
manipulation that affects the temporal facilitation dynamics onto only these two inhibitory
cell classes, we largely abolish the temporal response divergence between the two
cortical domains. Finally, using *in-vivo* viral tracing, whole-brain clearing and imaging, we
show a differential barrel and septa projection pattern to cortical regions S2 and M1. We
hence reveal that local temporally engaging cortical inhibition provided by SST+ neurons
contribute to the functional segregation of barrel and septa domains and potentially their
downstream targets.
44

Introduction
Although the somatosensory whisker system is one of the principal means by which mice
sense their environment and navigate the world, the number of whiskers is relatively low.
This relatively small number of vibrissae is represented in the whisker somatosensory
cortex as barrel "islands" (columns) separated by the septal "sea" (compartments),
reminiscent of the whisker pad pattern, with sparse hair follicles separated with significant
spaces between them. In contrast, other sensory systems such as the auditory and the
visual have densely packed anatomical receptor configurations and cortical
representations. Although all sensory cortices are composed of canonical microcircuits[1]
that include similar populations of excitatory and inhibitory neurons, maintaining
separation of individual whisker information may require anatomical specializations found
only in barrel cortex, such as the barrel and septa domains. These domains may therefore
create a distinct spatiotemporal stimulus representation and information coding in the
barrel cortex, compared to other sensory modalities.Such cortical columns have also
been shown to exist in other sensory cortical areas organized in specific sensory-receptor
to cortical-column mapping[2,3]. Cortical columns are thought to be separate processing
units acting as transfer modules between converging sensory stimuli and cognitive
faculties and/or motor output. In the barrel cortex, thalamocortical inputs coming from the

Ventro-Postero medial (VPM) and Postero-medial (PoM) nuclei segregate into the barrel and septa domains, respectively[4–7]. Further, information processing and flow within and from these domains target anatomically distinct sensory and motor areas[8–11]. It has been suggested that while the primary whisker somatosensory cortex (wS1) sends high dimensional information needed for object recognition to the secondary somatosensory cortex (wS2), the projections to the primary motor cortex (M1) carry less complex information unrelated to information coming from individual whiskers[6,10,12,13]. This divergence is also found in the barrel- and septa-related circuits. While barrel circuits are more involved in processing spatiotemporal whisker-object interactions, the septal circuits are more sensitive to the frequency of whisker movements, which suggests the use of temporal versus rate coding of information in barrel and septa, respectively[14,15]. Hence the existence of the septal region separating the barrels may provide the means that mice possess to separate these two interconnected yet very distinct information-processing routes along the temporal domain, in order to make sense of their somatosensory environment. This operational identity of barrels and septa has often been attributed to the parallel bottom-up thalamocortical pathways of the the VPM and PoM. Here, by performing *in-vivo* simultaneous silicon probe recordings in the barrel and septa domains in mice, we first characterize the temporal divergence between these domains. Through genetic labeling and passive tissue clearing, we reveal that cortical SST+ and VIP+ neurons show a layer-dependent differential distribution in the barrel versus septa domains. By altering the short-term synaptic dynamics of incoming excitation onto these interneurons upon removal of the synaptic protein *Elfn1*, we find that the domain-specific divergence of responses to a repetitive single-whisker stimulus is degraded. Finally, utilizing *in-vivo* retrograde viral-based tracing, we show a layer-specific projection preference between these domains and downstream regions wS2 and M1. Hence, in addition to the thalamocortical pathway-specific inputs, we reveal a key contribution of local lateral cortical inhibition provided by SST+ neurons in setting up the functional segregation of barrel and septa domains and subsequently the receiving downstream regions.

Results

Barrel and Septa columns display differentially adaptive spiking to repeated single whisker stimulation

To characterize the temporal stimulus response profiles of barrel and septal domains simultaneously, we performed *in-vivo* silicon probe recordings under urethane-induced light-anesthesia from mouse somatosensory whisker barrel cortex (wS1), after the onset age of whisking, postnatal day (P)20-30. Activity within identified barrels was recorded upon a 2s-long stimulation at 10Hz. Whisker stimulation was performed on either just one principal whisker (single-whisker stimulus or SWS, Figure 1A, 20 repetitions per principal whisker) or via both the principal whisker and most of the macro vibrissae together (multi-whisker stimulus or MWS, Supp. Figure 1A). Every mouse received both stimulation paradigms and post-hoc histology was used to assess the location of the multi-shank silicon probes, in order to assign the recorded activity to the stimulated principal barrel, the adjacent septa, or the adjacent unstimulated neighboring barrel (in the case of SWS) (Figure 1B and Supp Figure 1B). While this distinctive experimental paradigm is relatively low throughput, it has allowed us to investigate all these domains simultaneously.

It has previously been shown that electrophysiological responses upon whisking-based object detection saturate rapidly after a single whisker touch, whereas object localization may require multiple whisker touches[16,17], Correspondingly, when a single whisker is stimulated repeatedly, the response to the first pulse represents principally bottom-up thalamic-driven responses, whereas the later pulses in the train are expected to also gradually engage cortico-thalamo-cortical and cortico-cortical loops. We therefore first tested whether the initial pulses of the stimulus train (the first three out of total 20 pulses in 2s) evoked differential responses among the principal barrel, the adjacent septa, and the neighboring barrel upon SWS and MWS.

Earlier studies in which whisker-evoked responses were recorded either from barrel or from septa domains in separate experiments reported that the responses of barrel neurons to single principal whisker stimulation are much higher than those evoked in the

septal neurons[12,18]. The analysis of the data we obtained from the simultaneous
recordings at these domains, do not reveal such differences in first two pulse responses
among these columns in any of the laminae recorded upon single whisker stimulation
(SWS) (Figure 1D,G and J, First and Second Pulses). However response differences
emerged starting from the 3rd pulse upon SWS for L2/3 (Figure 1C,D and E, Third Pulse),
L4 (Figure 1F,G and H, Third Pulse) and L5 (Figure 1I,J and K, Third Pulse), and starting
from the 2nd pulse upon MWS in L4 (Supp. Figure 1E and F, Second Pulse), but not in
L2/3 or L5 (Supp. Figure 1C,D,G,H and I). While such differences also emerged between
stimulated principal and unstimulated neighboring barrels in L2/3 and L4 (starting at
different pulses; for L2/3: 7th pulse, for L4: 4th pulse, Figure1E and H, Barrel-vs-
Neighbour (BN) column), L5 did not show any significant difference (Figure 1K, BN
column). Interestingly, an inverted pattern of significant changes emerged between septa
and adjacent unstimulated neighboring barrels (Figure 1E, H and K, SN column). When
multiple whiskers were activated simultaneously, a divergence between barrel and septa
domains occured almost exclusively in Layer 4 (Supp. Figure 1I).
140
141 **SST+ and VIP+ neuron densities within Barrel and Septa Domains Diverge in Layer**
142 **4**
143
It is known that barrel and septa domains receive different thalamic projections, from VPM
and PoM respectively[19–22]. Our electrophysiological experiments show a significant
divergence of responses over time upon both SWS and MWS in L4 between barrels
(principal and neighboring) and adjacent septa that could be driven by this distinct
thalamocortical innervation or by differences in local circuitry. We therefore next assessed
the spatial distribution of local inhibitory cells in these domains, which could indicate their
active role in domain-specific activitation. Specifically, we made a comprehensive
analysis of the barrel versus septa localization of two inhibitory cell populations that
provide distinct regulation of cortical activity also in the temporal aspect, somatostatin
positive (SST+) and vasointestinal peptite positive (VIP+) cells.
154

Earlier research examining three inhibitory neuron populations in mouse barrel cortex
showed that, while parvalbumin positive (PV+) cells do not show any differential density
preference between barrel and septa columns, SST+ and VIP+ cells are significantly
denser in the septa than in the barrels in L4[23]. Using a different strategy, we aimed to
test the reported differential SST+ and VIP+ neuronal distributions. To label the two
neuronal populations, we crossed the SST-Cre and VIP-Cre lines with a tdTomato
reporter mouse line (Ai14). To comprehensively quantify the density of SST+ and VIP+
cells in the barrel and septa domains in 3D we utilized a passive CLARITY-based tissue
clearing protocol, followed by light-sheet microscopy using a custom-built light-sheet
microscope (mesoSPIM)[24,25]. The auto-fluorescence from the tissue was acquired with
488nm laser, which allows for the reliable detection of individual barrels in wS1, whereas
the tdTomato positive SST+ or VIP+ neurons were detected by a 594nm laser, which
permits us to localize and count cells in these two distinct domains accurately (Figure 2A
and Supp. Figure 2). By adjusting the angle of each barrel column (in 3D using Imaris),
we precisely identified the column of the barrel and septa in L2-3, L4 and L5, using the
barrel borders for every plane (XY,YZ,XZ) in L4 (see Methods). Once the columns were
accurately determined, we counted the number of tdTomato positive neurons in the
barrels and septa and corresponding volumes for density estimation. We found that
despite the known opposite distribution change along the depth of wS1 , both SST+ and
VIP+ neuronal densities were higher in L4 septa compared to barrels, but we detected no
difference in L2/3 and L5 (Figure 2B and C).
176
177 Based on the similarity of initial responses of all columns upon SWS and the subsequent
178 divergence emerging over time (Figure 1C-K), as well as the confirmation that SST+ and
179 VIP+ cell have higher densities in septa compared to barrel in L4 (Figure 2B and C), we
180 wanted to identify potential temporal adaptation mechanisms present in SST+ or VIP+
181 interneurons that could cause such physiological differences.
182
Out of these two interneuron classes, SST+ interneurons tend to be more spontaneously
active and supply powerful regulation to the local neuronal activity through dense
feedback connections to nearby pyramidal neurons[26] In contrast, VIP+ interneurons

have been reported to exert their influence on network function via a dis-inhibitory circuit
such that when driven by local, long-range, or neuromodulatory inputs, they inhibit SST+
interneurons (and to a lesser extent PV+ interneurons), resulting in reduced inhibition of
pyramidal neurons[27–29].
190
191 **SST+ and VIP+ interneurons exclusively express Elfn1 and the former contribute**
192 **stronger to the temporal regulation**
193
To explore potential genetic mechanisms that would be common to both SST+ and VIP+
inhibitory populations, which could contribute to the sensory adaptation observed, we
performed gene expression analysis using the Allen Institute database[30] (Figure 3A).
Subsequently, we also explored gene expression differences amongst all neuronal
populations in three different databases to confirm identified candidates[31] (Figure 3B).
To find gene with expression profiles that match closest to SST+ and VIP+ cells and
furthest from excitatory cells (Exc) and parvalbumin positive (PV+) cell , we defined two
distance metrics, combined correlation (CC) and combined p-value (Cp) (see Methods).
This approach favors genes that are significantly correlated with the SST and VIP genes
and uncorrelated with Exc and PV+ cell gene markers . In the Allen database, the Exc
neuron marker included everything that expressed the Emx1 gene, whereas VIP, SST
and PV genes were used for the respective neuronal populations. Although Emx1 is a
developmental gene that may also label non-neuronal cell types, since we only
considered neuronal expression, it does not pose a problem for our analysis.
208
We characterized five genes that have the highest CC and lowest Cp as potential
contributors to the temporal adaptation profiles present in SST+ or VIP+ interneurons
(Figure 3B and Supp. Figure 3A and B). Out of the five candidate genes that were
common between the two SST+ and VIP+ populations of neurons, *Elfn1* is known to be
involved in short-term plasticity and has been shown to affect the short-term excitatory
input dynamics onto SST+ neurons[32].
215

Having identified *Elfn1* as the sole synaptic gene that is expressed in SST+ and VIP+ with

known short-term plasticity effects, we went on to directly characterize the synaptic

facilitation levels of excitatory inputs onto SST+ and VIP+ interneurons in WT and *Elfn1*

knock-out (KO) mice. We made acute slices from SST-Cre X tdTomato and VIP-Cre X

tdTomato (Ai14) mice. Upon local neuropil stimulation and whole-cell recording from

SST+ neurons in L2/3, we found a strong short-term facilitation of excitatory transmission

in response to 10Hz neuropil stimulation (Figure 3C), which is in line with previous

work[32,33]. When performing the same type of recordings from VIP+ cells, we found that

genetic deletion of *Elfn1* also reduced the facilitation ratio at VIP interneurons (Multipolar,

posthoc characterization). However, at 10 Hz, WT VIP+ multipolar cells displayed minimal

facilitation, hence the loss of *Elfn1* produced synaptic depression instead (Figure 3D).

Although 10 Hz represents the same frequency used for the whisker stimulation, closer

examination of whisker evoked spiking, trace by trace, indicated that spiking consistency

is low, thus comparisons between in vitro and in vivo data should be made cautiously.

Nonetheless, the substantial reduction in facilitation for SST+ neurons, compared to

multipolar VIP+ neurons, which comprise only a minority of the VIP+ population (the

majority being bipolar, calretinin positive VIP+ neurons, which are unaffected in *Elfn1KO*

mice[33]) suggest that any effect that the removal of *Elfn1* would have on the domain-

dependent effects we observe in sensory-driven activity *in vivo,* would be dominated by

its impact on SST+ neurons, and not on VIP+ ones.

Barrel-Septa response differences disappear in *Elfn1KO* mice

Having characterized physiological and anatomical differences between barrel and septa

domains and having found a potential responsible gene for the physiological divergence

between these anatomical domains, we next explored how the lack of *Elfn1* affects the

sensory-driven activity *in vivo*.

Simultaneous recordings in barrel and septa domains in *Elfn1KO mice* upon SWS (Figure

4) and MWS (Supp. Figure 4) in fact revealed a lack of stimulus-dependent emerging

differences between these domains found in WT controls upon SWS. While the

differences between barrels, septa and unstimulated neighbors upon SWS were largely
diminished in L2/3 (Figure 4A,B and C) and L5 (Figure 4G,H and I), the observed
differences in L4 between barrels and septa were not as strong between barrels and
unstimulated neighbors (Figure 4D,E and F). Interestingly, response differences between
barrel and septa domains upon MWS did not differ compared to WT controls, apart from
the fact that the emergence of dissimilarities started at the 5^{th} pulse as opposed to 2^{nd} in
WT (Supp. Figure 4G and 2I, respectively). This finding suggests that activity spillover
from principal barrels is normally constrained by the delayed inhibition provided by the
presence of *Elfn1* in excitatory synapses onto SST+ neurons, which have been shown to
supply late recurrent inhibition in the cortex and hippocampus[32,34].
257
258   **Barrels and Septa response identity is lost in *Elfn1KO* mice**
259
It has been shown that the septal circuits are more sensitive to the frequency of whisker
movements, which exemplifies the use of temporal vs rate coding of information in barrel
and septa, respectively[14,15]. Hence it is conceivable that the existence of the septal
regions separating the barrels provides a means to allow mice to separate these two
interconnected yet very distinct information-processing routes along the temporal domain.
In order to assess the information content of each of the two anatomical domains in WT
animals and how it may change when activity between them converges in the *KO* animals,
we trained a one-versus-all error-correcting output code (ECOC) classifier[35] with a
GentleBoost[36] ensemble of decision trees using WT firing profile data from three
domains (3-class classification). First, we performed 10-fold cross validation over the WT
dataset (Figure 5A and B, left columns) and later we tested the WT-data-trained classifier
with *Elfn1KO* firing profiles to compare the classifier performance.
272
273   The results show that in L2/3 and L4, barrel columns (Stimulated barrels and unstimulated
274   neighbors) we get higher classification accuracy compared to septa for SWS data (Figure
275   5A, left column, L2/3 and L4), while L5 receiver operating characteristic (ROC) curves
276   almost completely overlap for all conditions (Figure 5A, left columns, L5). Interestingly,
277   while decoding accuracy for unstimulated neighbors was not drastically altered upon *KO-*

testing (Figure 5A, right column, all layers, turquoise), barrel and septa performances dropped significantly (Figure 5A, right column, gray and purple). This might not be surprising given that barrel, septa and neighbor response profiles are closer in Elfn1KO compared to WT (Supp. Figure 5A). We also saw that MWS-based decoders perform worse both for WT-crossvalidation and *KO* testing (Figure 5B) which is most likely due to the fact that barrel vs septa responses in WT animals are more similar to one another, while this is not the case for *Elfn1KO* animals (S5B Fig). However, most interestingly, layer 5 decoding performance completely drops to chance levels for the *KO*-test case (Figure 5B, L5, WT-Train, KO-Test).

It is quite interesting to observe that while response profiles of the tested domains got closer in *Elfn1KO* versus WT upon SWS (Supp. Figure 5A), the opposite happened upon MWS (Supp. Figure 5B).

wS1 Barrel Columns and Septal Domains have different wS2 and M1 Layer-dependent projection preferences

We identified that barrel and septa respond differently in WT mice and that this difference disappears in the *Elfn1KO* animals upon SWS and amplified upon MWS. Following up from this observation, we finally aimed to explore if the projection targets of these two wS1 domains are distinct, in a similar manner to what has been revealed for visual cortex domain projections[37]. By utilizing *in vivo* retrograde AAV injections in two of the main target cortical regions of wS1, we characterized the projections from wS1 to wS2 and M1.

We injected a retrograde AAV virus expressing GFP in wS2 or M1 in different mice (N=4 for S2 and N=6 for M1) and 4 weeks after the injection the brains were processed using passive tissue clearing and imaged using a custom-built light-sheet microscope (mesoSPIM) (Figure 6A). The auto-fluorescence of the barrels allowed for labeling of barrel columns in 3D, similar to our analysis on interneuron population distributions (Fig 2A and S2 Fig). We counted the number of neurons in wS1 barrel and septa domains

and measured the corresponding domain volumes to estimate projection densities as a

proxy for connectivity levels between S1-S2 and S1-M1.

Our analysis showed that while barrels host significantly more wS2-projection cells in L2/3

(Figure 6B), septa domains host more M1-projecting neurons (Figure 6C). Overall our

findings suggest that barrels verus septa domains do not only process distinct sensory-

driven information regulated by local SST+ cell inhibition, but also transmit this information

in a biased manner to downstream regions wS2 and M1.

Discussion

In the absence of visual stimuli, humans aim to identify an object by handling, with the

first touch initiating the information accumulation process, and every subsequent touch

leading to the full identification and characterization of the object, through evidence

accumulation(Armstrong-James & Fox, 1987; Kheradpezhouh et al., 2017; Petersen,

2007).As is known, this process includes parrarel ascending pathways that bring the

peripheral touch events through central pathways to the cortex in a somatotopic manner.

Similarly, mice actively use their whiskers in place of arms and hands, again

somatotopically mapping the whisker spatial pattern information one-to-one to the

characterisitc cortical barrels.[40,41].

Although a large number of other species have whiskers, only a subset of animals with

whiskers display a barrel pattern. For example, cats have whiskers and barrel-like

structures in the brainstem nuclei (barrelettes) but have no barrels visible in their

cortex[42–44]. It is the volitional whisking behavior that seems to correlate with an animal

having barrels in their wS1 or not. The fact that animals whisk to identify surrounding

structures and only have cortical barrels when they whisk, is suggestive of the importance

of the one to one mapping in individual whisker information processing in the cortex. Here

we find that septa play an important role in separating these parrarel whisker streams by

regulating the lateral cortico-cortical information flow generated upon repetitive whisker

touch. This process is tightly controlled by the short-term facilitation properties of

incoming input onto SST+ interneurons which is enabled by the expression of the Elfn1
gene.

It has previously been shown that, in contrast to L2/3 and L5/6, where SST+ cells are
layer 1 targeting Martinotti type, the axons of SST+ INs in layer 4 of somatosensory cortex
largely remain within layer 4[45]. In addition, while Martinotti cells receive facilitating
synapses from PCs and form inhibitory synapses onto dendrites of neighboring PCs[46],
non-Martinotti SST cells predominantly inhibit L4 FS cells in Layer 4[45,47]. Regarding
the inputs onto the two L4 populations, SST+ and PV+ inhibitory cells are also activated
in a divergent manner during thalamocortical activation at naturalistic whisking
frequencies, with FS interneurons transiently excited due to the rapid depression of their
thalamocortical inputs, whereas SST+ cells the opposite[48].

Somatosensation requires the accumulation of spatial and temporal information through
repeated coordinated movement of corresponding sensory organs and these organs are
often consisted of separate units such as hands and fingers or snout and whiskers.
Although it has been speculated that the information relayed to and from barrels may
differ from that of the septa[6,9,38], it has been unclear how this segregation is
accomplished. The one-to-one topographic mapping of whiskers, thalamic barreloids, and
cortical barrels favors an *a priori* assumption of bottom-up segregation of cortical domain
identity. In contrast, our simultaneous recordings in the barrel and septa domains
demonstrate that their sensory responses to the first whisker deflection do not significantly
differ (Figure 1C-K). This would suggest that the first touch serves as an initiating signal
in both domains, and it is only with repetitive activation of individual whiskers that their
responses strongly diverge. Our work reveals that in addition to distinct feedforward
thalamocortical projections, the cortical distribution of SST+ and VIP+ neurons (Figure 2)
and the short-term synaptic dynamics onto SST+ interneurons (Figure 3) also contribute
to this divergence and help shape cortical response identity. In our comparative gene
analysis between septa-enriched and non-enriched cell types, we identified new gene
candidates, including a long non-coding RNA of unknown function, for further study as
potential mediators of differential response dynamics in wS1. Amongst the handful of

other genes found in our bioinformatics screen and based on previous work[32–34,49]*,*

Elfn1 was a strong candidate for mediating a stimulius-dependent domain diversgence.

Indeed we found that its removal abolishes the functional temporal divergence between

barrel and septa domains (Figure 4). Thus, in addition to thalamo-cortical feedforward

projections, we find that intracortical activity, shaped by SST+ inhibitory cells contributes

to cortical domain response identity.

To further study if the spiking divergence in the temporal domain carries important

sensory information, we trained an ECOC-based classifier to decode sensory stimuli

among these separate yet interconnected domains. While the temporal profiles recorded

at barrel domains upon SWS were successfully classified by the algorithm, the septal

data was not (Figure 5A). These findings are consistent with previous suggestions that

high dimensional temporal information (e.g., relating to object identity or location) is

transmitted from wS1 to wS2 via the barrels, whereas low dimensional sensory

information (e.g., object detection) is sent from wS1 to wM1 via septa[6,10,12,14,15].

Indeed, our retrograde labeling is consistent with this interpretation, with projection cells

from wS1 barrels showing a bias to wS2, whereas projections from wS1 septa being

biased to M1.When we instead examine multi-whisker stimulation, our sensory decoding

from the classifier fails (Figure 5B), suggesting an impaired transmission of high

dimensional data, at least in our experimental paradigm. Interestingly, the deletion of

Elfn1 decayed the ability of SWS to support sensory decoding which we attribute to the

corresponding loss of SST+ interneuron contributions to frequency segregation. In

addition, we saw a loss of differentiation between SW and MW stimuli and between barrel

and septal regions (Figure 5A and Supp. Figure 5A and B).

Thus, in addition to any contributions from distinct thalamocortical input pathways, our

findings reveal an important role for short-term plasticity driving cortical SST+ cell-

mediated delayed inhibition in diversifying the response properties between individual

barrels and septal regions, creating contrast in somatosensory information processing.

Acknowledgments

We thank members of the Karayannis lab for suggestions and comments on the manuscript. Authors also thank Rahel Kaestli for help with tissue clearing and light-sheet image acquisition.

Funding

This work was supported by Swiss National Science Foundation Grant S-41260-01-01 (T.K) and European Research Council Grant 679175 (T.K).

Author Contributions

Conceptualization: Ali Özgür Argunşah, Tevye Jason Stachniak, Jenq-Wei Yang, Theofanis Karayannis.

Data curation: Ali Özgür Argunşah, Tevye Jason Stachniak, Jenq-Wei Yang, Linbi Cai.

Formal analysis: Ali Özgür Argunşah, Tevye Jason Stachniak.

Funding acquisition: Theofanis Karayannis.

Investigation: Ali Özgür Argunşah, Tevye Jason Stachniak, Linbi Cai.

Methodology: Ali Özgür Argunşah, Tevye Jason Stachniak, Jenq-Wei Yang, Linbi Cai, George Kanatouris.

Project administration: Theofanis Karayannis.

Software: Ali Özgür Argunşah.

Supervision: Theofanis Karayannis.

Validation: Ali Özgür Argunşah, Tevye Jason Stachniak.

Visualization: Ali Özgür Argunşah, Tevye Jason Stachniak, Linbi Cai.

Mice: Tevye Jason Stachniak, Linbi Cai, George Kanatouris.

Immunohistochemistry: Tevye Jason Stachniak, Linbi Cai, George Kanatouris.

Writing – original draft: Ali Özgür Argunşah

Writing – review & editing: Ali Özgür Argunşah, Tevye Jason Stachniak, Theofanis Karayannis.

Declaration of Interest

The authors declare no competing interests.

Methods

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Theofanis Karayannis (karayannis@hifo.uzh.ch)

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

The datasets and analysis routines are available from the corresponding author on reasonable request.

Mice

All the animal experiments followed the guidelines of the Veterinary Office of Switzerland and were approved by the Cantonal Veterinary Office Zurich and the University of Zurich. husbandry with a 12-h reverse dark-light cycle (7 a.m. to 7 p.m. dark) at 24 °C and variable humidity. Adult (5- to 10-week-old) male C57BL6J wide type mice were used for retrograde tracing. Animal lines used in this study are VIP-IRES-Cre (Vip$^{tm1(cre)Zjh}$/J)[50], SST-IRES-Cre (Sst$^{tm2.1(cre)Zjh}$/J) [50], Ai14 (B6;129S6-Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J)[51] and HTB ((Gt(ROSA)26Sortm1(CAG-neo,-HTB)Fhg)[52], and Elfn1KO (Elfn1tm1(KOMP)Vlcg). Both male and female mice were used in cortical brain slice physiology and in vivo physiology experiments.

In-Vitro Electrophysiology

Whole-cell patch-clamp electrophysiological recordings were performed on labelled SST neurons located in neocortical layers 2/3 and 4 of barrel cortex (L2/3, L4, approximately bregma -0.5 to -2.0 mm) in acute brain slices prepared from postnatal day 7- 22 (P7–P22) male and female mice. Coronal brain slices from barrel cortex were prepared in cold

artificial cerebrospinal fluid (aCSF) containing (in mM): 128 NaCl, 26 NaHCO3, 10 D-glucose, 3 KCl, 1 MgCl2, 2 CaCl2 and 1.25 NaH2PO4, aerated with 95% O_2 / 5% CO_2. Acute slices were perfused at a rate of 2–3 ml/min with oxygenated ACSF at room temperature. Patch electrodes were made from borosilicate glass (Harvard Apparatus) and had a resistance of 2-4 MΩ. The intracellular solution had (in mM): 125 K gluconate, 2 KCl, 10 HEPES, 10 phosphocreatine, 4 MgCl2, 1 EGTA, 0.1 CaCl2, 4 ATP, 0.4 GTP, pH 7.35, 290 mOsm. Experiments were performed in voltage-clamp mode using the Axopatch 200B amplifier (Molecular Devices). Visually guided patch of fluorescent labelled cells was performed on a Zeiss Axioscope using a Retiga Electro camera (01-ELECTRO-M-14-C-OC, Teledyne Scientific Imaging). Access resistance was monitored to ensure the stability of recording conditions. Recordings with access resistance >40 MΩ, or whole cell capacitance <4 pF were excluded. No compensation was made for access resistance and no correction was made for the junction potential between the pipette and the ACSF. Following a baseline stabilization period (2-3 min), evoked synaptic currents recorded in 2 min (12 sweeps) at Vh= -70 mV were averaged and analyzed using Clampfit10 (Molecular Devices). Five electrical stimuli from a Digitimer isolated stimulator (DS2A Mk.II) were delivered at 50 Hz through a monopolar glass pipette (2-4 MΩ) positioned in L2/3, close to the soma of the recorded cells. The stimulating electrode was placed typically 100 - 250 μm from the recorded cell, parallel to the pial surface. A similar distance was maintained in L4 recordings, with the stimulating electrode parallel to the L2/3-L4 boundary. For the L2/3 to L4 recordings, the simulating electrode was then moved to lower L3 within the same column, again approximately 100 - 250 μm from the recorded cell but offset slightly to avoid directly stimulating the recorded cell's axon. Stimulation intensity and duration were adjusted to produce stable evoked EPSC amplitudes.

In-Vivo Silicon Probe Recording

We used 4 Elfn1KO and 4 WT littermates at the age of P20 to P30 for the multi-electrode recordings. Mice were anesthetized by urethane (1.5g/1kg) throughout the whole experiment. A heating pad was used to maintain the mouse's body temperature at 37°C. The depth of anesthesia was checked with breathing speed and paw reflexes throughout

the experiment. The skull of the right hemisphere was exposed by removing the skin on

top, and a metallic head holder was implanted on the skull with cyanoacrylate glue and

dental cement. A 20G needle was used to open a ~3mm x 3mm cranial window which

exposed the S1 barrel field (wS1). Extreme care was taken not to cause damage or

surface bleeding during surgery.

wS1 neural activities were recorded with an 8-shank-64-channel silicon probe. Each of

the 8 shanks has 8 recording sites (100 µm apart). The distance between each shank is

200 µm (NeuroNexus Technologies, Ann Arbor, MI, USA). The silicon probe was labeled

with DiI (1,1′-dioctadecyl-3,3,3′,3′-tetramethylindocarbocyanine, Molecular Probes,

Eugene, OR, USA) and inserted perpendicularly into the barrel cortex. A silver wire was

placed into the cerebellum as a ground electrode. All data were acquired at 20 kHz and

stored with MC_RACK software (Multi Channel systems). The total duration of multi-

electrode recordings varied between 3 h and 5 h. After each experiment, the animal was

deeply anesthetized by ketamine (120 mg/kg, ketamine, 50 mg/mL, HamelnPharma,

Hameln Germany) and perfused through the aorta with ringer solution. The brain was

kept in 4% PFA. Tangential sections (200-µm thick) were prepared for cytochrome-

oxidase (CO) histochemistry. By combining the DiI and CO staining, the insertion position

of the 8 shank probes were identified in the barrel cortex. Only the shanks located within

the identified individual barrels were used for the data analysis.

Whisker Stimulation

A single whisker was stimulated 1mm from the snout in rostral-to caudal direction (about

1mm displacement) using a stainless-steel rod (1 mm diameter) connected to a miniature

solenoid actuator. The movement of the tip of the stimulator bar was measured precisely

using a laser micrometer (MX series, Metralight, CA, USA) with a 2500 Hz sampling rate.

The stimulus takes 26 ms to reach the maximal 1 mm whisker displacement, with a total

duration of 60 ms until it reaches baseline (Yang et al., 2017). The whisker was stimulated

at 10Hz for 2s. Each principal whisker in case of SWS and multiple whiskers at once in

the case MWS was stimulated 20 times (20 trials).

Analysis of Silicon Probe Data

Extracellular silicon probe data were analyzed using a custom-made Matlab script (Matlab 2019a, Mathworks, MA, USA). The raw data signal was band-pass filtered (0.8-5 kHz) and the multi-unit activity (MUA) was extracted with the threshold of 7.5 times the standard deviation (SD) of baseline. The current source density (CSD) map was used to identify L2-3 and L4. The earliest CSD sink was identified as layer 4, followed by L2-3 (Reyes-Puerta et al., 2015; van der Bourg et al., 2017). MUA was smoothed using a Gaussian kernel (0 mean, 5ms sigma).

Decoder Analysis

We employed a classification approach utilizing error-correcting output codes (ECOC) models with a gentle adaptive boosting (GentleBoost) strategy to train classifiers. This method is particularly effective in handling imbalanced data and unequal misclassification costs.

Like Logit Boost, each weak learner in the ensemble fitted a regression model to response values, $y_n \in \{-1, +1\}$. The mean-squared error, stored in the FitInfo property of the ensemble object by fitcensemble, is computed as:

$$N \sum_{n=0}^{d} d(t)_n (y_n - h_t(x_n))^2$$

where, $d(t)_n$ represents observation weights at step t (summing up to 1), and $h_t(x_n)$ denotes predictions from the regression model h_t fitted to response values y_n. As the strength of individual learners diminishes, the weighted mean-squared error converges towards 1.

The training of classifiers utilized wild-type data. In the case of single whisker (SW) data, three classes were considered: barrel, septa, and neighbors. For multi-whisker (MW) data, two classes were defined: barrel and septa. Subsequently, Elfn1 knockout (KO) SW or MW data was introduced into the trained model to obtain predictions, respectively.

Retrograde tracing and tissue processing

Mice were anesthetized by isoflurane and a small craniotomy was made above the injection site (M1 or S2). 200nl AAV24-retro GFP was injected in the M1 or S2. 4 weeks after injection, the animals were transcardially perfused using 1X PBS and 4% PFA. Following that, the whole brains were used for hydrogel-based tissue clearing by putting in a hydrogel solution (1% PFA, 4% Acrylamide, 0.05% Bis) for 48h before the hydrogel polymerization was induced at 37 C. Afterwards, the brains were put in 40ml of 8% SDS and kept shaking at room temperature until the tissue was cleared sufficiently (around 2 months). Then the brains were washed 3 times in PBS and put into a self-made refractive index matching solution (RIMS) for the last clearing step. They were left to equilibrate in 5ml of RIMS for at least 4 days at room temperature before being imaged.

Imaging of injected brain

After clearing, brains were imagined by a home-built mesoscale selective plane illumination microscope (mesoSPIM). The detailed steps were described elsewhere (Voigt, F.F et al, 2019). The images were acquired by laser 488nm & 561nm with 0.8X magnification (pixel size: 8.23um) for the view of the whole brain and 2X magnification (pixel size: 3.26um) for the view of the ipsilateral barrel cortex. 488nm laser was to acquire the fluorescence of virus while 561nm was to acquire an auto-fluorescence background to visualize barrels in the barrel cortex.

Tissue Processing for Passive Clearing and Imaging

The method used for hydrogel-based tissue clearing is explained in detail elsewhere [53]. Briefly, the animals were transcardially perfused using 1x PBS and Hydrogel solution (1% PFA, 4% Acrylamide, 0.05% Bis). The collected brains were post-fixed for 48 h in a Hydrogel solution (1% PFA, 4% Acrylamide, 0.05% Bis). Afterward, the Hydrogel polymerization was induced at 37 °C. Following the polymerization, the brains were immersed in 40 mL of 8% SDS and kept shaking at room temperature until the tissue was cleared sufficiently (10–40 days depending on the animal's age). Finally, after 2–4 washes in PBS, the brains were put into a self-made refractive index matching solution (RIMS)

[54]. They were left to equilibrate in 5 mL of RIMS for at least 4 days at RT (Room
Temperature) before being imaged. After clearing, brains were attached to a small weight
and loaded into a quartz cuvette, then submerged in RIMS, and imaged using a home-
built mesoscale selective plane illumination microscope (mesoSPIM)[24]. The
microscope consists of a dual-sided excitation path using a fiber-coupled multiline laser
combiner (405, 488, 515, 561, 594, 647 nm, Omicron SOLE-6) and a detection path
comprising an Olympus MVX-10 zoom macroscope with a 1× objective (Olympus
MVPLAPO 1x), a filter wheel (Ludl 96A350), and a scientific CMOS (sCMOS) camera
(Hamamatsu Orca Flash 4.0 V3). For imaging tdTomato and eGFP a 594 nm excitation
with a 594 long-pass filter (594 LP Edge Basic, AHF) and 488 nm & 520/35 (BrightLine
HC, AHF) were used respectively. The excitation paths also contain galvo scanners
(GCM-2280-1500, Citizen Chiba) for light-sheet generation and reduction of streaking
artifacts due to absorption of the light-sheet. In addition, the beam waist is scanning using
electrically tunable lenses (ETL, Optotune EL-16-40-5D-TC-L) synchronized with the
rolling shutter of the sCMOS camera. This axially scanned light-sheet mode (ASLM) leads
to a uniform axial resolution across the field-of-view of 5–10 μm (depending on zoom and
wavelength). The field of views ranged from 10.79 mm at ×1 or x0.8 magnification (Pixel
size: 5.27 μm) for overview datasets. Further technical details of the mesoSPIM are
described elsewhere[24,25]. The images generated with the mesoSPIM were
preprocessed using Fiji and Imaris software to generate the images of the barrel cortex.
For the quantification of neurons, the collected RAW files were converted to HDF5 format,
and each z-plane was median filtered (window size 2 pixels) and convolved with a disc
kernel (radius 1.5, convolution window size 10). The convolution was performed on the
log-transformed z-plane and the convolution result was thresholded (threshold 0.3). On
the convolved, thresholded z-planes local maxima were detected and then consolidated
into cell detections in 3D space by matching and grouping detections on adjacent slices
that are less than 5 pixels apart. Grouped detections were discarded if they spanned less
than 4 adjacent slices. For the remaining grouped detections 3D cell positions were
defined by the detection at median z location. 3D consolidated detections were filtered
anatomically and morphologically using a semi-automated procedure to exclude non

cortical detections and detections not corresponding to neurons (e.g., blood vessels).
Analysis was performed manually using Imaris
618

Analysis of cleared brain imaging data
Barrel cortex was imaged in 3D with 2X objective. The images were processed through
Fiji and Imaris to visualize the 3D structure of the barrel cortex. XY-, YZ-, XZ- projections
were visualized in Imaris, which allows us to adjust the correct angle for each barrel
column to be vertical against XY-plane. This method allowed us identify barrel and septa
in layers 2-3 and 5 and layer 4. In addition, the depth information is also precisely acquired
for each barrel column. SST+ or VIP+ neuron numbers, S2-projected and M1-projected
S1 neuron numbers were counted in icy. Max projection for each 50 ums along barrel
columns was acquired and neurons on each max projection image were counted
automatically by icy with the same threshold for all the brains. Each barrel was drawn
according to the barrel map from each brain by max projection. The area of septa was
calculated by subtracting the whole selected area from all the selected barrel areas.
631

Gene expression analysis
Two metrics were computed, Expression of all genes correlated with SST, VIP, PV and
EMX genes and compared with ttest and plotted in 2D. Since we wanted high correlation
and strong similarity with VIP and SST and the opposite for PV and Emx1, we used a
combined similarity and correlation scores using the following formulas, $Cp =$
$pSST*pVIP*(1-pPV)*(1-pEMX)$, where pSST is the p value came out of a ttest between
all genes and SST etc. $CC = cSST*cVIP*(1-cPV)*(1-cEMX)$ where cSST is the correlation
between SST gene and all genes.
640

Statistical Analysis
Data are represented as mean ± SEM unless stated otherwise. Statistical comparisons
have been done using a one-tailed Mann-Whitney-U test. In the case of the Gaussian
cluster analysis, a two-tailed t-test was applied on the epsilon distance between starter
cells and pre-synaptic partners to check the statistical significance threshold was set to
$p<0.05$; in the Figures, different degrees of evidence against the null hypothesis are

indicated by asterisks ($p<0.05$: *; $p<0.01$: **; $p<0.001$: ***). Cell densities in Figure 2 and

6 were compared using repeated measures ANOVA. All tests were conducted using

custom codes in Matlab.

Figures Legends

Figure 1. Layer specific temporal divergences emerge in barrel and septa domains

upon repeated single-whisker stimulation. (A) Acute *in-vivo* silicon probe (8x8)

recordings were performed after the start of active whisking (>P21). Principal whiskers

were stimulated one by one for 2s at 10Hz. (B) Left: Probe locations were labeled after

each experiment using histology. Right: Only the triplets of electrodes were used in the

analysis in which principal barrel (B), adjacent septa (S) and the adjacent neighboring

barrel (N) were captured by the probe insertion sites for any experiment. 14 barrel

columns, 9 septal columns, 18 unstimulated neighboring barrels from N=4 mice. (C)

Analysis for Layer 2/3. Left: Average firing profiles of barrel (black), unstimulated neighbor

(blue) and septa (pink) columns upon 2s-long 10Hz repeated whisker stimulation. Middle:

Average firing rates per stimulating pulse. Right: Average firing rates for 1^{st}, 2^{nd}, 3^{rd}, and

20^{th} pulses zoomed in. (D) Pair-wise statistical comparisons of three conditions for 1^{st},

2^{nd}, 3^{rd}, and 20^{th} pulses (Mann-Whitney-U-Test). (E) Pair-wise statistical comparisons of

three conditions for all pulses (Mann-Whitney-U-Test). (F-G-H) Same with C-D-E but for

Layer 4. (I-J-K) Same with C-D-E but for Layer 5. (*: $p<0.05$; **: $p<0.01$;***:$p<0.001$; Dark

shading is a function of p value and white stands for $p>0.05$.)

Figure 2. SST+ and VIP+ neuron densities differ at Barrel and Septa Domains in

Layer 4. (A) A passive CLARITY-based tissue clearing protocol was performed on the

collected brains which were subsequently imaged in their entirety using a custom-built

light-sheet microscope (mesoSPIM). 594nm wavelength was used to image tdTomato,

488nm wavelength was used to image autofluorescence. (B) SST+ neuron density

distribution over laminae for barrel and septal domains (N=4 mice, n=54 Barrels and

surrounding septa per barrel). Normalized cell densities plotted as a function of depth.

0μm represents the top of layer 4. SST+ interneuron density is significantly higher in septa

(p=0.0015, F=30.64, repeated-measures ANOVA). (C) VIP+ neuron density distribution

over laminae for barrel and septal domains. (N=4 mice, n= 53 Barrels and surrounding

septa per barrel). Normalized cell densities plotted as a function of depth. 0µm represents

the top of layer 4. VIP+ interneuron density is also significantly higher in septa (p=0.0.21,

F=9.63, repeated-measures ANOVA)

Figure 3. Elfn1 is predominantly expressed in SST+ and VIP+ Interneurons and

responsible for the short-term plasticity related temporal engagement. (A) Scatter

plot showing genes exclusively expressed by SST+ and VIP+ neurons only. X-axis is the

combined correlation which is calculated as follows: cSST.*cVIP.*(1-cPV).*(1-cEXC),

where cX represents the correlation between the test sequence and the neuronal type of

interest X. This formula favors genes that are highly expressed in SST and VIP cells and

not expressed in Exc and PV cells. The higher the combined correlation, the better.

Similarly, Y-Axis is the combined p-value which is calculated as follows pSST.*pVIP.*(1-

pPV).*(1-pEXC), where pX represents the p-value (two-sample t-test) between the test

sequence and the neuronal type of interest X. This formula favors genes that significantly

similar with SST and VIP cells and different from Exc and PV cells. Lower the combined

p-value, better. (B) Elfn1 gene expression data across three RNA-seq datasets containing

cortical excitatory neurons and inhibitory interneuron subtypes visualized as boxplots of

RPKM expression levels. Boxes show median and interquartile ranges, and whiskers

show full data without outliers. (C) Excitatory synaptic events evoked by repetitive local

electrical stimulation produce strongly facilitating EPSCs onto SST interneurons. At 10

Hz, this is principally due to a suppression of initial release probability. For wildtype SST

interneurons EPSCs facilitate rapidly at 10 Hz (p=0.003, 3×10^{-7}, 3×10^{-11} and 3×10^{-15}

for 2^{nd} 3^{rd} 4^{th} and 5^{th} stimuli vs 1^{st} stimuli, respectively). Elfn1 knockout removes this

facilitation (P = 1, 1, 1, 1 vs 1^{st}). Responses in KO neurons facilitate less than in WT

neurons (P= 0.0001, 6×10^{-7} for 4^{th} and 5^{th} stimuli ANOVA with Bonferroni post hoc). (D)

Elfn1 KO also alters ESPC facilitation ratio onto multipolar VIP interneurons at 10 Hz.

Wildtype VIP interneurons do not display substantial facilitation at 10 Hz (p=1, 1, 1 and 1

for 2^{nd} 3^{rd} 4^{th} and 5^{th} stimuli vs 1^{st} stimuli, respectively). Elfn1 knockout induces synaptic

depression (p=1,6×10^{-4}, 5×10^{-6}, 1×10^{-6} vs 1^{st}). Responses in KO neurons depress more

than in WT neurons (p=0.03, 0.0004, 0.001 for 3rd, 4th, and 5th stimuli ANOVA with

Bonferroni post hoc). *p<0.05, **p<0.01, ***p<0.001

Figure 4. Loss of Elfn1 abolishes barrel-septa response divergences upon single-

whisker stimulation. (A) Analysis for Layer 2/3. Left: Average firing profiles of barrel

(gray), unstimulated neighbor (turquois) and septa (purple) columns upon 2s-long 10Hz

repeated whisker stimulation. Middle: Average firing rates per stimulating pulse (11 barrel

columns, 7 septal columns, 11 unstimulated neighbors from N=4 mice.). Right: Average

firing rates for 1st, 2nd, 3rd, and 20th pulses. (B) Pair-wise statistical comparisons of three

conditions for four pulses. (C) Pair-wise statistical comparisons of three conditions for all

pulses. (D-E-F) Same with A-B-C but for Layer 4. (G-H-I) Same with A-B-C but for Layer

5. (*: p<0.05; **: p<0.01;***:p<0.001; Two-sided Mann-Whitney U Test. White represents

p>0.05.)

Figure 5. Decoding analysis shows an alteration of columnar domain identity in

__Elfn1KO__ **animals.** (A) Decoder analysis based on the single whisker stimulation

response profiles. Left column represents the receiver operating characteristic (ROC)

curve of the classification performance of the decoder trained and cross-validated using

WT data. Right column represents the ROC curve of the classification performance after

training with WT and testing with KO data. (B) Decoder analysis based on the multi

whisker stimulation response profiles. Left column represents the receiver operating

characteristic (ROC) curve of the classification performance of the decoder trained and

cross-validated using multi-whisker WT data. Right column represents the ROC curve of

the classification performance after training with WT and testing with KO data.

Figure 6. wS1 barrel and septa columns differentially project to wS2 and M1 (A) The

representation of the experimental protocol and timelines. The retro-AAV-CAG-GFP was

injected in either S2, or M1 at P20-30 and the brains were dissected 4 weeks after the

virus injection following the tissue clearing and whole brain imaging. (B) Normalized cell

density profiles of S2 and M1 projection neurons at barrel columns. 0µm represents the

top of layer 4. Layer 2/3-barrel columns send significantly more projections to S2

(p=0.0293, F=7.02, repeated-measures ANOVA). (C) Normalized cell density profiles of
S2 and M1 projections neurons at septa columns. 0µm represents the top of layer 4.
Layer 5 septal columns send significantly more projections to M1 (p=0.0402, F=5.98,
repeated measures ANOVA).
744
745 **Supplementary Figure Legends**
746
Supplementary Figure 1. Layer 4 presents the highest level of temporal
divergences between barrel and septa domains upon repeated multi-whisker
stimulation. (A) Acute *in-vivo* silicon probe (8x8) recordings were performed after the
start of active whisking (>P21). Most of the principal whiskers were stimulated for 2s at
10Hz. (B) Left: Probe locations were labeled after each experiment using histology. Right:
Only the triplets of electrodes were used in the analysis in which principal barrel (B),
adjacent septa (S) and the adjacent neighboring barrel (N) were captured by the probe
insertion sites for single whisker stimulation experiments. Only barrels and septa were
used for multi whisker analysis. 14 barrel columns, 9 septal columns from n=4 mice. (C)
Analysis for Layer 2/3. Left: Average firing profiles of barrel (black) and septa (pink)
columns upon 2s-long 10Hz repeated whisker stimulation. Middle: Average firing rates
per stimulating pulse. Right: Average firing rates for 1^{st}, 2^{nd}, 3^{rd}, and 20^{th} pulses. (D) Pair-
wise statistical comparisons of three conditions for four pulses. (E-F) Same with C-D but
for Layer 4. (G-H) Same with C-D but for Layer 5. (I) Barrel-Septa comparison for Layer
2/3,4 and 5 (*: p<0.05; **: p<0.01;***:p<0.001; Two-sided Mann-Whitney U Test. White
represents p>0.05.)
763
764 **Supplementary Figure 2. SST+ and VIP+ neuron densities differ at Barrel and Septa**
765 **Domains in Layer 4.** Left: Example SST-tdTomato brain virtual slice on left. Right Top:
766 Inner ring represents barrel borders; outer ring represents septal borders. Right Bottom:
767 Same as top but without green channel.
768
769 **Supplementary Figure 3. Elfn1 is solely expressed in SST+ and VIP+ Interneurons**
770 **and responsible for the short-term plasticity related temporal engagement.** (A) Each

bar shows the differential expression of Elfn1 in a comparison within one of the datasets. Fold-Changes are relative to the other 3 groups of neuron subtypes within the same dataset. (B) Gene expression data for other four candidates across three RNA-seq datasets containing cortical excitatory neurons and inhibitory interneuron subtypes. Each dot represents a single sample.

Supplementary Figure 4. Loss of Elfn1 does not alter barrel-septa response profiles upon multi-whisker stimulation. (A) Analysis for Layer 2/3. Left: Average firing profiles of barrel (gray) and septa (purple) columns upon 2s-long 10Hz repeated whisker stimulation. Middle: Average firing rates per stimulating pulse (11 barrel columns, 7 septal columns from n=4 mice.)). Right: Average firing rates for 1^{st}, 2^{nd}, 3^{rd}, and 20^{th} pulses. (B) Pair-wise statistical comparisons of three conditions for four pulses. (C-D) Same with A-B but for Layer 4. (E-F) Same with A-B but for Layer 5. (G) Barrel-Septa comparison for Layer 2/3,4 and 5 (*: $p<0.05$; **: $p<0.01$;***:$p<0.001$; Two-sided Mann-Whitney U Test. White represents $p>0.05$.)

Supplementary Figure 5. Visual presentation of WT vs Elfn1KO response profiles. (A) Visual comparison of mean firing rates of WT vs KO SWS responses for all three conditions and layers. (B) Visual comparison of mean firing rates of WT vs KO MWS responses for both conditions and all layers.

References

1. Douglas RJ, Martin KAC. Neuronal circuits of the neocortex. Annual Review of Neuroscience. 2004. doi:10.1146/annurev.neuro.27.070203.144152
2. Horton JG, Adams DL. The cortical column: A structure without a function. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005. doi:10.1098/rstb.2005.1623
3. Harris KD, Mrsic-Flogel TD. Cortical connectivity and sensory coding. Nature. 2013. doi:10.1038/nature12654
4. Kim U, Ebner FF. Barrels and septa: Separate circuits in rat barrel field cortex. Journal of Comparative Neurology. 1999;408. doi:10.1002/(SICI)1096-9861(19990614)408:4<489::AID-CNE4>3.0.CO;2-E

804   5.   Bureau I, Von Paul F Saint, Svoboda K. Interdigitated paralemniscal and lemniscal
805   pathways in the mouse barrel cortex. PLoS Biol. 2006;4.
806   doi:10.1371/journal.pbio.0040382
807   6.   Alloway KD. Information processing streams in rodent barrel cortex: The differential
808   functions of barrel and septal circuits. Cerebral Cortex. 2008. doi:10.1093/cercor/bhm138
809   7.   Staiger JF, Petersen CCH. Neuronal circuits in barrel cortex for whisker sensory
810   perception. Physiol Rev. 2021;101. doi:10.1152/physrev.00019.2019
811   8.   Shepherd GMG, Svoboda K. Laminar and columnar organization of ascending excitatory
812   projections to layer 2/3 pyramidal neurons in rat barrel cortex. Journal of Neuroscience.
813   2005;25. doi:10.1523/JNEUROSCI.1173-05.2005
814   9.   Chakrabarti S, Alloway KD. Differential origin of projections from SI barrel cortex to the
815   whisker representations in SII and MI. Journal of Comparative Neurology. 2006;498.
816   doi:10.1002/cne.21052
817   10.   Sato TR, Svoboda K. The functional properties of barrel cortex neurons projecting to the
818   primary motor cortex. Journal of Neuroscience. 2010;30. doi:10.1523/JNEUROSCI.3774-
819   09.2010
820   11.   Audette NJ, Urban-Ciecko J, Matsushita M, Barth AL. POm Thalamocortical Input Drives
821   Layer-Specific Microcircuits in Somatosensory Cortex. Cereb Cortex. 2018;28.
822   doi:10.1093/cercor/bhx044
823   12.   Brecht M, Sakmann B. Dynamic representation of whisker deflection by synaptic
824   potentials in spiny stellate and pyramidal cells in the barrels and septa of layer 4 rat
825   somatosensory cortex. Journal of Physiology. 2002. doi:10.1113/jphysiol.2002.018465
826   13.   Cai L, Yang JW, Wang CF, Chou SJ, Luhmann HJ, Karayannis T. Identification of a
827   Developmental Switch in Information Transfer between Whisker S1 and S2 Cortex in
828   Mice. Journal of Neuroscience. 2022;42. doi:10.1523/JNEUROSCI.2246-21.2022
829   14.   Melzer P, Champney GC, Maguire MJ, Ebner FF. Rate code and temporal code for
830   frequency of whisker stimulation in rat primary and secondary somatic sensory cortex.
831   Exp Brain Res. 2006;172: 370–386. doi:10.1007/s00221-005-0334-1
832   15.   Melzer P, Sachdev RNS, Jenkinson N, Ebner FF. Stimulus frequency processing in awake
833   rat barrel cortex. Journal of Neuroscience. 2006;26. doi:10.1523/JNEUROSCI.2620-
834   06.2006
835   16.   O'Connor DH, Hires SA, Guo Z V., Li N, Yu J, Sun QQ, et al. Neural coding during
836   active somatosensation revealed using illusory touch. Nat Neurosci. 2013;16.
837   doi:10.1038/nn.3419
838   17.   Pammer L, O'Connor DH, Hires SA, Clack NG, Huber D, Myers EW, et al. The
839   mechanical variables underlying object localization along the axis of the whisker. Journal
840   of Neuroscience. 2013;33. doi:10.1523/JNEUROSCI.4316-12.2013
841   18.   Armstrong-James M, Fox K. Spatiotemporal convergence and divergence in the rat S1
842   "Barrel" cortex. Journal of Comparative Neurology. 1987;263.
843   doi:10.1002/cne.902630209
844   19.   Koralek KA, Jensen KF, Killackey HP. Evidence for two complementary patterns of
845   thalamic input to the rat somatosensory cortex. Brain Res. 1988;463. doi:10.1016/0006-
846   8993(88)90408-8
847   20.   Chmielowska J, Carvell GE, Simons DJ. Spatial organization of thalamocortical and
848   corticothalamic projection systems in the rat SmI barrel cortex. Journal of Comparative
849   Neurology. 1989;285. doi:10.1002/cne.902850304

21. Ahissar E, Sosnik R, Bagdasarian K, Haidarliu S. Temporal frequency of whisker
movement. II. Laminar organization of cortical representations. J Neurophysiol. 2001;86.
doi:10.1152/jn.2001.86.1.354
22. Sosnik R, Haidarliu S, Ahissar E. Temporal frequency of whisker movement. I.
Representations in brain stem and thalamus. J Neurophysiol. 2001;86.
doi:10.1152/jn.2001.86.1.339
23. Almási Z, Dávid C, Witte M, Staiger JF. Distribution patterns of three molecularly
defined classes of gabaergic neurons across columnar compartments in mouse barrel
cortex. Front Neuroanat. 2019;13. doi:10.3389/fnana.2019.00045
24. Voigt FF, Kirschenbaum D, Platonova E, Pagès S, Campbell RAA, Kastli R, et al. The
mesoSPIM initiative: open-source light-sheet microscopes for imaging cleared tissue. Nat
Methods. 2019;16. doi:10.1038/s41592-019-0554-0
25. Vladimirov N, Voigt F, Naert T, Araujo GR, Cai R, Reuss AM, et al. The Benchtop
mesoSPIM: a next-generation open-source light-sheet microscope for large cleared
samples. bioRxiv. 2023.
26. Urban-Ciecko J, Barth AL. Somatostatin-expressing neurons in cortical networks. Nat
Rev Neurosci. 2016;17. doi:10.1038/nrn.2016.53
27. Lee S, Kruglikov I, Huang ZJ, Fishell G, Rudy B. A disinhibitory circuit mediates motor
integration in the somatosensory cortex. Nat Neurosci. 2013;16. doi:10.1038/nn.3544
28. Pi HJ, Hangya B, Kvitsiani D, Sanders JI, Huang ZJ, Kepecs A. Cortical interneurons that
specialize in disinhibitory control. Nature. 2013;503. doi:10.1038/nature12676
29. Karnani MMM, Jackson J, Ayzenshtat I, Tucciarone J, Manoocheri K, Snider WGG, et al.
Cooperative Subnetworks of Molecularly Similar Interneurons in Mouse Neocortex.
Neuron. 2016;90. doi:10.1016/j.neuron.2016.02.037
30. Yao Z, van Velthoven CTJ, Nguyen TN, Goldy J, Sedeno-Cortes AE, Baftizadeh F, et al.
A taxonomy of transcriptomic cell types across the isocortex and hippocampal formation.
Cell. 2021;184. doi:10.1016/j.cell.2021.04.021
31. Huntley MA, Srinivasan K, Friedman BA, Wang TM, Yee AX, Wang Y, et al. Genome-
wide analysis of differential gene expression and splicing in excitatory neurons and
interneuron subtypes. Journal of Neuroscience. 2020;40. doi:10.1523/JNEUROSCI.1615-
19.2019
32. Stachniak TJ, Sylwestrak EL, Scheiffele P, Hall BJ, Ghosh A. Elfn1-induced constitutive
activation of mglur7 determines frequency-dependent recruitment of somatostatin
interneurons. Journal of Neuroscience. 2019;39. doi:10.1523/JNEUROSCI.2276-18.2019
33. Stachniak TJ, Kastli R, Hanley O, Argunsah AÖ, Van Der Valk EGT, Kanatouris G, et al.
Postmitotic prox1 expression controls the final specification of cortical VIP interneuron
subtypes. Journal of Neuroscience. 2021;41. doi:10.1523/JNEUROSCI.1021-21.2021
34. Tomioka NH, Yasuda H, Miyamoto H, Hatayama M, Morimura N, Matsumoto Y, et al.
Elfn1 recruits presynaptic mGluR7 in trans and its loss results in seizures. Nat Commun.
2014;5. doi:10.1038/ncomms5501
35. Allwein EL, Schapire RE, Singer Y. Reducing multiclass to binary: A unifying approach
for margin classifiers. Journal of Machine Learning Research. 2001;1.
36. Friedman J, Hastie T, Tibshirani R. Additive logistic regression: A statistical view of
boosting. Annals of Statistics. 2000. doi:10.1214/aos/1016218223

37.   Meier AM, Wang Q, Ji W, Ganachaud J, Burkhalter A. Modular network between
895   postrhinal visual cortex, amygdala, and entorhinal cortex. Journal of Neuroscience.
896   2021;41. doi:10.1523/JNEUROSCI.2185-20.2021
897   38.   Petersen CCH. The functional organization of the barrel cortex. Neuron. 2007.
898   doi:10.1016/j.neuron.2007.09.017
899   39.   Kheradpezhouh E, Adibi M, Arabzadeh E. Response dynamics of rat barrel cortex
900   neurons to repeated sensory stimulation. Sci Rep. 2017;7. doi:10.1038/s41598-017-11477-
901   6
902   40.   Van Der Loos H, Woolsey TA. Somatosensory cortex: Structural alterations following
903   early injury to sense organs. Science (1979). 1973;179. doi:10.1126/science.179.4071.395
904   41.   Woolsey TA, Welker C, Schwartz RH. Comparative anatomical studies of the Sml face
905   cortex with special reference to the occurrence of "barrels" in layer IV. Journal of
906   Comparative Neurology. 1975;164. doi:10.1002/cne.901640107
907   42.   Rice FL, Gomez C, Barstow C, Burnet A, Sands P. AComparative analysis of the
908   development of the primary somatosensory cortex: Interspecies similarities during barrel
909   and laminar development. Journal of Comparative Neurology. 1985;236.
910   doi:10.1002/cne.902360405
911   43.   Rice FL. An attempt to find vibrissa-related barrels in the primary somatosensory cortex
912   of the cat. Neurosci Lett. 1985;53. doi:10.1016/0304-3940(85)90180-6
913   44.   Nomura S, Itoh K, Sugimoto T, Yasui Y, Kamiya H, Mizuno N. Mystacial vibrissae
914   representation within the trigeminal sensory nuclei of the cat. Journal of Comparative
915   Neurology. 1986;253. doi:10.1002/cne.902530110
916   45.   Xu H, Jeong HY, Tremblay R, Rudy B. Neocortical Somatostatin-Expressing GABAergic
917   Interneurons Disinhibit the Thalamorecipient Layer 4. Neuron. 2013;77.
918   doi:10.1016/j.neuron.2012.11.004
919   46.   Silberberg G, Markram H. Disynaptic Inhibition between Neocortical Pyramidal Cells
920   Mediated by Martinotti Cells. Neuron. 2007;53. doi:10.1016/j.neuron.2007.02.012
921   47.   Ma Y, Hu H, Berrebi AS, Mathers PH, Agmon A. Distinct subtypes of somatostatin-
922   containing neocortical interneurons revealed in transgenic mice. Journal of Neuroscience.
923   2006;26. doi:10.1523/JNEUROSCI.0661-06.2006
924   48.   Tan Z, Hu H, Huang ZJ, Agmon A. Robust but delayed thalamocortical activation of
925   dendritic-targeting inhibitory interneurons. Proc Natl Acad Sci U S A. 2008;105.
926   doi:10.1073/pnas.0710628105
927   49.   Stachniak TJ, Argunsah AÖ, Yang J-W, Cai L, Karayannis T. Presynaptic kainate
928   receptors onto somatostatin interneurons are recruited by activity throughout development
929   and contribute to cortical sensory adaptation. Journal of Neuroscience. 2023;43.
930   doi:10.1523/JNEUROSCI.1461-22.2023
931   50.   Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, et al. A Resource of Cre Driver
932   Lines for Genetic Targeting of GABAergic Neurons in Cerebral Cortex. Neuron. 2011;71:
933   995–1013. doi:10.1016/j.neuron.2011.07.026
934   51.   Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, et al. A robust and
935   high-throughput Cre reporting and characterization system for the whole mouse brain. Nat
936   Neurosci. 2010;13: 133–140. doi:10.1038/nn.2467
937   52.   Li Y, Stam FJ, Aimone JB, Goulding M, Callaway EM, Gage FH. Molecular layer
938   perforant path-associated cells contribute to feed-forward inhibition in the adult dentate

gyrus. - Supporting Information. Proc Natl Acad Sci U S A. 2013;110: 9106–11.
doi:10.1073/pnas.1306912110
53.   Chung K, Deisseroth K. CLARITY for mapping the nervous system. Nat Methods.
2013;10. doi:10.1038/nmeth.2481
54.   Yang B, Treweek JB, Kulkarni RP, Deverman BE, Chen CK, Lubeck E, et al. Single-cell
phenotyping within transparent intact tissue through whole-body clearing. Cell. 2014;158.
doi:10.1016/j.cell.2014.07.017
946

947   Figures

Figure 1, Argunsah et al., 2024

Figure 2, Argunsah et al., 2024

200 µm

Supplementary Figure 2, Argunsah et al., 2024

Figure 3, Argunsah et al., 2024

Supplementary Figure 3, Argunsah et al., 2024

Figure 4, Argunsah et al., 2024

Supplementary Figure 4, Argunsah et al., 2024

Figure 5, Argunsah et al., 2024

Supplementary Figure 5, Argunsah et al., 2024

Figure 6, Argunsah et al., 2024