GTDrift: A resource for exploring the interplay between genetic drift, genomic and transcriptomic characteristics in eukaryotes

Florian Bénitière[1,2] , **Laurent Duret**[1] , **Anamaria Necsulea**[1]

[1]Laboratoire de Biométrie et Biologie Évolutive, Université Lyon 1, UMR CNRS 5558, Villeurbanne, France.
[2]Laboratoire d'Écologie des Hydrosystèmes Naturels et Anthropisés, Université Lyon 1, UMR CNRS 5023, Villeurbanne, France.
Correspondence: florian.benitiere@univ-lyon1.fr

January 23, 2024

Abstract

We present GTDrift, a comprehensive data resource that enables explorations of genomic and transcriptomic characteristics alongside proxies of the intensity of genetic drift in individual species. This resource encompasses data for 1,507 eukaryotic species, including 1,414 animals and 93 green plants, and is organized in three components. The first two components contain approximations of the effective population size, which serve as indicators of the extent of random genetic drift within each species. In the first component, we meticulously investigated public databases to assemble data on life history traits such as longevity, adult body length and body mass for a set of 969 species. The second component includes estimations of the ratio between the rate of non-synonymous substitutions and the rate of synonymous substitutions (dN/dS) in protein-coding sequences for 1,324 species. This ratio provides an estimate of the efficiency of natural selection in purging deleterious substitutions. The third component encompasses various genomic and transcriptomic characteristics. With this component, we aim to facilitate comparative transcriptomics analyses across species, by providing easy-to-use processed data for more than 16,000 RNA-seq samples across 491 species. These data include intron-centered alternative splicing frequencies, gene expression levels and sequencing depth statistics for each species, obtained with a homogeneous analysis protocol. To enable cross-species comparisons, we provide orthology predictions for conserved single-copy genes based on BUSCO gene sets. To illustrate the possible uses of this database, we identify the most frequently used introns for each gene and we assess how the sequencing depth available for each species affects our power to identify major and minor splice variants.

***K*eywords** Dataset · Random genetic drift · Effective population size · N_{e} · Life history traits · Alternative splicing · dN/dS · Splice variants · Genomes · Transcriptomic

Introduction

Genetic drift refers to stochastic fluctuations in allele frequencies within a population across successive generations. These fluctuations arise due to the inherently random sampling of individuals that reproduce and pass on their alleles to subsequent generations (Wright, 1929; Graur *et al.*, 2000). Population genetics principles state that the ability of natural selection to promote beneficial mutations or eliminate deleterious mutations depends on the intensity of selection (s) relative to the power of genetic drift (defined by the effective population size, N_e): if the selection coefficient is sufficiently weak relative to drift ($|N_e s| < 1$), alleles behave as if they are effectively neutral (Kimura *et al.*, 1963; Ohta, 1973). Thus, random drift sets an upper limit on the efficiency of selection. This limit is called the "drift barrier" (Lynch, 2007, 2010). Genomes that are subject to intense genetic drift are expected to be less well-optimized compared to those experiencing lower genetic drift. Michael Lynch proposed that variation in the ability to purge slightly deleterious mutations (*i.e.* variation in N_e) can account for differences in genome characteristics among species (Lynch and Conery, 2003). This hypothesis has been empirically validated for multiple genome characteristics and phylogenetic clades. For example, it was shown that the genomes of crustacean species with low N_e values are larger than those of their sister species (Lefébure *et al.*, 2017). Moreover, species with large N_e tend to have a lower mutation rate than species with low N_e, illustrating the notion that natural selection acts to improve replication fidelity, within the constraints defined by random genetic drift (Lynch *et al.*, 2016).

We recently examined the variations in transcriptome complexity across animal species in light of the "drift barrier" hypothesis (Bénitière *et al.*, 2023). In multicellular eukaryotes, the vast majority of genes give rise to multiple isoforms through alternative splicing (Chen *et al.*, 2014). This phenomenon has attracted a great deal of interest since its discovery almost 50 years ago (Berget *et al.*, 1977). Alternative splicing is commonly hypothesized to be adaptive, because it can increase the number of biological functions that are encoded in each genome. Indeed, numerous instances of alternative splicing patterns with beneficial effects have been identified (Mudge *et al.*, 2011; Barbosa-Morais *et al.*, 2012; Merkin *et al.*, 2012; Reyes *et al.*, 2013; Verta and Jacobs, 2022; Singh and Ahi, 2022; Wright *et al.*, 2022). However, these examples represent only a small fraction of all splice variants that are now known, especially given the substantial detection power brought by next-generation RNA sequencing (RNA-seq) techniques. Many of the splice variants that can now be detected with RNA-seq are present at very low frequencies (Gonzàlez-Porta *et al.*, 2013; Tress *et al.*, 2017) and are poorly conserved during evolution (Barbosa-Morais *et al.*, 2012; Merkin *et al.*, 2012). It was thus hypothesized that they may be the result of errors of the splicing machinery, rather than functional isoforms (Pickrell *et al.*, 2010; Gout *et al.*, 2013; Xu and Zhang, 2014; Saudemont *et al.*, 2017; Xu and Zhang, 2018; Liu and Zhang, 2018b,a; Xu *et al.*, 2019; Xu and Zhang, 2020; Zhang and Xu, 2022). Notably, according to the "drift barrier" hypothesis, one may hypothesize that if alternative splicing (AS) primarily serves functional roles, the rate of alternative splicing should increase with N_e. Conversely, if AS predominantly involves deleterious processes, its rate should decline with increasing N_e. We applied this reasoning in our previous work (Bénitière *et al.*, 2023), which led us to deduce that AS is predominantly non-functional.

This methodology for exploring the impact of N_e on biological processes holds potential for broader applications. For example, one could examine the functional importance of alternative polyadenylation sites (Xu and Zhang,

2018). Such investigations demand cross-species comparative transcriptomics analyses, a task facilitated by
the abundant availability of publicly accessible RNA-seq data. Yet, analysis of transcriptome sequencing data
is resource-intensive in terms of time, energy, and computational power. To facilitate future analyses, we
provide a comprehensive database that streamlines the process by offering pre-processed data. This dataset
includes proxies for effective population size, sets of orthologous single-copy genes, gene expression levels,
and intron-centered alternative splicing frequencies, along with phylogenetic trees to control for phylogenetic
inertia. These resources have been compiled for 16,000 RNA-seq samples spanning 1,507 multicellular
eukaryotic species.

This database, that we name GTDrift, complements other public transcriptomic data resources, such as
Bgee (Bastian *et al.*, 2021), which provides gene expression levels for 52 species (Version 15.0.1), but not
alternative splicing frequencies. Other databases do provide alternative splicing frequencies. For example,
MeDAS (Li *et al.*, 2021) provides AS data for 18 metazoan species, and MetazExp (Liu *et al.*, 2022) provides
data for 72 metazoan species. This latter resource is substantial, including data for $\sim 53,000$ RNA-seq
samples. However, this database favors insects (53 species, with $\sim 26\,000$ RNA-seq samples for *Drosophila*
melanogaster) and does not include any representative of the vertebrate clade, for which more computational
resources are required because of their large genomes. Our database encompasses a broader phylogenetic
distribution of species (Fig. 1), with 93 green plant species, 561 invertebrates and 853 vertebrates. Moreover,
while other public databases such as MetazExp are aimed at biologists want to analyze alternative splicing
patterns in a gene-by-gene manner through web queries, in GTDrift we provide all data in flat files, which
enable downstream computational analyses. GTDrift is thus mainly aimed at users with some computational
skills. Nevertheless, we have created a user-friendly Shiny app to facilitate exploration of the database and
species-specific data downloads (Chang *et al.*, 2023).

In GTDrift, we used assemblies and annotations data collected from The National Center for Biotechnology
Information (NCBI) (Sayers *et al.*, 2022), as well as publicly available RNA-seq data to investigate alternative
splicing patterns and gene expression profiles. We computed summary statistics across all analyzed RNA-seq
samples for each species, which enabled us to determine whether the available sequencing depth is sufficient
for the study of alternative splicing. To ensure comparability across species, we annotated Benchmarking
Universal Single Copy Orthologs (BUSCO) (Waterhouse *et al.*, 2018) genes in all species and provide
phylogenetic trees to control for phylogenetic inertia.

We believe that this tremendous amount of information should be shared with the scientific community,
because it provides the means to investigate the impact of genetic drift on genome and transcriptome
architecture, on a broad phylogenetic scale.

Methods

Species selection

95 The first criterion for species inclusion in GTDrift is the availability of a genome assembly and annotation in
96 the NCBI database (NCBI Resource Coordinators, 2018; Sayers *et al.*, 2022), as well as the availability of

Clades

- Embryophyta N=93
- Diptera N=96
- Lepidoptera N=54
- Coleoptera N=30
- Hymenoptera N=94
- Other Insects N=47
- Nematoda N=58
- Teleostei N=188
- Mammalia N=213
- Aves N=372
- Other Vertebrates N=80
- Other Metazoans N=182

Figure 1: **Phylogenetic distribution of the species included in the GTDrift database.** The phylogeny was retrieved from TimeTree (Kumar *et al.*, 2022). Not all species studied are present (N=1,221).

RNA-seq data in the Short Read Archive (Leinonen *et al.*, 2011). We included 1,507 multicellular eukaryotic species. This collection encompasses 1,414 animal species as well as 93 species of green plants (Fig. 1). Our Snakemake pipeline can be applied to any species for which genome sequence, genome annotation and RNA-seq data are available, which will enable us to further expand GTDrift in the future(Mölder *et al.*, 2021).

Collecting life history traits

We queried several databases to acquire three specific life history traits, namely: maximum longevity, body mass, and body length. These traits were previously identified as suitable proxies for estimating the effective population size (Romiguier *et al.*, 2014; Waples, 2016; Figuet *et al.*, 2016; Galtier, 2016; Weyna and Romiguier,

2020). For eusocial species, which live in colonies and have both reproductive and non-breeding individuals, we gather data on the queen of the colony. For solitary species, we collected data for females if available; otherwise, males were considered.

We employed several distinct methodologies to screen the databases. We initially used a manual approach to search across various sources of information, including scientific papers and databases.

We manually searched for information on life history traits from four prominent databases, which encompass diverse taxonomic groups. The Animal Ageing and Longevity Database (AnAge) (Tacutu *et al.*, 2013), is renowned for its comprehensive collection of vertebrates, particularly mammals. The Encyclopedia of Life (EOL) (Wilson, 2003; Parr *et al.*, 2014) encompasses a wide spectrum of species, prominently featuring invertebrates. The Animal Diversity Web (ADW) (Myers *et al.*, 2023), is a particularly rich resource for invertebrates. The FishBase (Froese and Pauly, 2023) predominantly houses data on teleostei species. While AnAge furnishes extensive information regarding body mass and lifespan, it is lacking data pertaining to body length (Fig. 2A,B,C). Furthermore, as previously noted, certain databases are tailored to specific clades. For instance, in comparison to EOL and ADW, AnAge contains relatively fewer records for invertebrates (Fig. 2D,E,F).

We then made efforts to automate the manual search procedures. The primary automated procedure involved the development of a bash script, which utilized the Latin nomenclature of the species to navigate the textual content within the research pages of the 4 databases listed above. The bash script was designed to extract sentences, words, and numerical data in proximity to keywords such as "longevity", "mass", and "weight", serving as indicators of relevant information. Its output was then reformatted through an R script. While this approach proved effective for databases like AnAge, EOL, and FishBase, its applicability to the ADW database was limited due to the manner in which information is embedded within textual paragraphs. Consequently, we employed an alternative method for the ADW database, involving Machine Learning and Natural Language Processing Question-Answering techniques. We obtained a trained model named "tinyroberta-squad2" from huggingface.co (noa, 2023). This model was used to answer questions related to specific attributes, such as 'what is the body length ?'; 'what is the body mass ?'; 'what is the longevity ?'. Each question retrieved a pool of 100 potential answers derived from the database's textual content, ranked by their predictive scores provided by the model.

We implemented an iterative selection process to identify the highest predicted answer containing relevant units and numeric values. To avoid redundancy, the selected answer was then removed from the text, and the process was repeated up to 10 times. The entire procedure was implemented in a Python script. We processed the script's output to restructure the obtained results.

Discrepancies between the manual approach and the other two methodologies were further re-investigated manually and corrected as needed after a further re-reading of the text. As a result, the curated dataset that we share reflects our highest level of confidence.

In total, our data collection effort resulted in the acquisition of life history traits for 969 metazoan species.

Figure 2: **Representation of life history traits retrieved from diverse data sources.** Depiction of data origins
for lifespan (**A**), body length (**B**), and body mass (**C**). Additionally, distribution of species and their respective clades
with at least one recorded life history trait in ADW (**D**), EOL (**E**), and AnAge (**F**).

Acquisition of the reference genome sequence and annotations

Using the sra-tools software, we performed an automated identification of the reference genome for each
species. Subsequently, we downloaded the annotation data in GFF format, the nucleotide coding sequences in
FASTA format, and the peptide sequences in FASTA format from the NCBI database (Sayers *et al.*, 2022).

dN/dS **pipeline**

We developed a pipeline to estimate the rate of non-synonymous substitutions divided by the synonymous
substitutions rate (dN/dS), representing the frequency at which non-synonymous changes occur relative to
synonymous ones. Since non-synonymous substitutions are commonly perceived as errors, dN/dS serves as a
measure of the rate of erroneous substitutions per neutral substitution. This ratio is directly dependant of N_e
as it is jointly determined by the distribution of selection coefficient of new mutations (s) and the magnitude
of genetic drift as defined by N_e (Yang and Nielsen, 1998; Nielsen and Yang, 2003). The transcriptome-wide
dN/dS is expected to rise over prolonged periods of small N_e due to the increasing number of slightly
deleterious mutations reaching fixation (Ohta, 1992; Galtier, 2016).

155 Estimating the dN/dS necessitates the annotation of genes shared across all species, their evolutionary
156 history depicted by a phylogenetic tree, and finally a comparative analysis of site evolution to derive the
157 dN/dS ratio.

BUSCO genes identification

159 We used the BUSCO v.3.1.0 software to identify single-copy orthologous genes within three datasets selected
160 from OrthoDB v9 (Zdobnov *et al.*, 2017): eukaryota (N=303 genes), embryophyta (N=1,440 genes) and
161 metazoan (N=978 genes) sourced from BUSCOv3 (Waterhouse *et al.*, 2018; Seppey *et al.*, 2019; Manni *et al.*,
162 2021). The search was performed against the longest annotated protein sequences *per* gene within each
163 genome.

Phylogenetic tree reconstruction

Due to the considerable time and resource demands associated with phylogenetic inference for large numbers
of species, we employed a strategy in which the analysis was partitioned by clades. On initial releases of the
database, which did not encompass all current species, we performed 3 comparable and independent analyses
that rely on the three BUSCO datasets, corresponding to the following lineages: eukaryota, embryophyta and
metazoa. For each BUSCO dataset, we selected a subset of species that matched the lineage of interest from
the available database records at the time of analysis. All of these selected species underwent transcriptomic
analyses (see Transcriptomic analyses). We then collected the longest corresponding proteins identified in each
species for each BUSCO gene family. We removed proteins for which the amino acid sequence provided with
the annotations did not perfectly correspond to the translation of the corresponding coding sequences. We
then aligned the resulting sets of protein-coding sequences for each BUSCO gene, using the codon alignment
option in PRANK v.170427 (Löytynoja and Goldman, 2008). We translated the codon alignments into
protein alignments using the R package seqinr (Charif and Lobry, 2007).

A filter was applied to retain only genes for which enough species have been detected (85% of the analyzed
species), reducing the eukaryota set to 126 genes (embryophyta N=387 genes, metazoa N=731 genes). Then,
species were removed from the analysis if they had less than 80% of the studied genes, reducing the number
of studied species from 336 to 279 for the eukaryota BUSCO dataset (embryophyta 93 to 80 species, metazoa
293 to 257 species).

To infer the phylogenetic tree rapidly, we sub-sampled the resulting multiple alignments, selecting alignments
with the highest number of species (eukaryota N=25 genes, embryophyta N=77 genes, metazoa N=146 genes).
We then concatenated these alignments and kept sites that were aligned in most of the analyzed species (see
information provided in the supplementary archive for more details). The final alignment for the eukaryota
BUSCO dataset included 279 taxa (embryophyta N=80 species, metazoa N=257 species) taxa and 600 sites
(embryophyta N=700 sites, metazoa N=3,000 sites). We used RAxML-NG (Kozlov *et al.*, 2019), to infer
the species phylogeny on these final alignments. RAxML was set to perform one model *per* gene with a
fixed empirical substitution matrix (LG), empirical amino acid frequencies from alignment (F) and 8 discrete
GAMMA categories (G8). These parameters were specified in a partition file with one line *per* BUSCO gene

multiple alignment. The analysis generated at least 10 starting trees. The best-scoring topology was kept as the final ML tree and 10 bootstrap replicates have been generated.

The phylogenetic trees were rooted using as a reference source the TimeTree phylogeny, which synthesizes data from numerous published studies, despite its incomplete representation of all species (Kumar *et al.*, 2022).

To encompass a broader spectrum of the species included in our latest database release, the one published here, we also reconstructed phylogenetic trees *per* clade. To do this, we divided the full set of metazoan species in 9 groups (Hymenoptera, Diptera grouped with Lepidoptera under the superorder Mecopterida, Nematoda, other insects, Aves, Mammalia, Teleostei, other vertebrates, and finally other invertebrates). We used as a basis for the analysis 73 highly prevalent metazoan BUSCO genes among the 731 genes preselected in the metazoa analysis. We applied the protocol described above to each individual clade. The resulting clade-specific trees were merged using outgroup species as a reference point to construct the complete metazoan phylogenetic tree.

dN/dS computation

We computed dN/dS ratios for BUSCO gene families that were present in at least 85 percent of the species under investigation. We conducted four independent analyses. We first analyzed each of the three BUSCO gene sets: eukaryota (N=126 genes), embryophyta (N=387 genes), metazoa (N=731 genes). We also performed an analysis '*per* clade', as explained above for the phylogenetic tree reconstruction, using the same 731 genes preselected in the metazoa analysis. Codon alignments obtained using PRANK (Löytynoja and Goldman, 2008) served as the basis for this estimation. To manage the computational memory demands during the substitution rate estimation step, we segmented the sequence alignments into clusters. Following the approach recommended by Bolívar *et al.* (2019), these clusters were defined based on the average GC3 content across species, in order to group genes with similar parameters. We then concatenated the alignments within each group, obtaining alignments that were 200 kb long on average. This process yielded 13 groups for eukaryota (15 for embryophyta and 73 for metazoa). We used bio++ v.3.0.0 libraries (Dutheil and Boussau, 2008; Guéguen *et al.*, 2013; Bolívar *et al.*, 2019) to estimate the dN/dS on each branch of the phylogenetic tree, for each concatenated alignment.

In a first step, we used an homogeneous codon model implemented in bppml to infer the most likely branch lengths, codon frequencies at the root, and substitution model parameters. We used YN98 (F3X4) (Yang and Nielsen, 1998) substitution model, which allows for different nucleotide content dynamics across codon positions. In a second step, we used the MapNH substitution mapping method to count synonymous and non-synonymous substitutions (Dutheil *et al.*, 2012; Guéguen and Duret, 2018). We defined dN as the total number of non-synonymous substitutions divided by the total number of non-synonymous mutational opportunities, both summed across concatenated alignments, for each branch of the phylogenetic tree. Likewise, we defined dS as the total number of synonymous substitutions divided by the total number of synonymous mutational opportunities, both summed across concatenated alignments. The *per*-species dN/dS corresponds to the

227 ratio between dN and dS, on the terminal branches of the phylogenetic tree. We also provide the dN and dS
228 values for each branch within the phylogenetic trees.

229 For the 'per clade' approach, the results pertaining to distinct clades were combined in a single table.

Transcriptomic analyses

231 We developed a pipeline facilitating the detection of alternative splicing events within genes. This process
232 entails the selection of RNA-seq data, subsequent alignment to the reference genome, and the identification
233 of splicing events through the recognition of introns. Utilizing the aligned transcriptomic data, we computed
234 gene expression levels across each sample.

Selection of the RNA-seq samples

To extract RNA-seq data, we queried the Short Read Archive (SRA) database for samples where the library
source was 'TRANSCRIPTOMIC' and the library strategy was 'RNA-seq'.

For perfect comparability of transcriptome data among species, we would need to have the same representation
of individual tissues, developmental stages *etc.* for each species, with data generated with the same protocol
by the same person. However, such data exist only for limited sets of species (*e.g.*, Cardoso-Moreira *et al.*
(2019)). Here, we decided not to filter the RNA-seq samples on criteria pertaining to sample origin or
experimental protocols, mainly because the relevant information is not always provided in sufficient detail
in the SRA database (Leinonen *et al.*, 2011). Moreover, depending on the clade, the biological sample of
origin can vary from "whole body" in insects, to specific tissues or cell types in mammals. Thus, perfectly
comparable sample collections are difficult to obtain across such a broad phylogenetic scale.

Rather than filtering samples on these criteria prior to inclusion in the database, in GTDrift we provide users
with the information collected from SRA for all RNA-seq samples. This information includes the library type,
the date of extraction and the name of the laboratory that performed the experiment (see Description of the
data available in GTDrift).

After evaluating the amount of RNA-seq data that is needed to evaluate global alternative splicing patterns
for each species (see below), we decided to include a maximum of 50 RNA-seq samples *per* species in GTDrift.
We included more than 50 samples for 150 species (43 embryophyta, 107 metazoa), for which we performed
more detailed analyses, considering various tissues or developmental stages.

In the current version of GTDrift, the RNA-seq dataset encompasses a total of 491 distinct species, including
92 plants and 399 animals. (Fig. 3A).

Indexing genomes and aligning RNA-seq data

257 The RNA-seq alignment phase represents the most time-consuming stage in the pipeline (Fig. 4), and can
258 extend up to one week when utilizing 16 cores for each RNA-seq dataset, particularly for larger genomes such
259 as those of mammals.

Figure 3: **Species with transcriptomic data and alternative splicing estimation.** (*cf* Fig. 2A Bénitière *et al.* (2023)) **A**: Taxonomic distribution of the species for which transcriptomic data was included in GTDrift. **B**: Definition of the variables used to compute the relative splicing frequency of a focal intron, compared to splice variants with a common alternative splice boundary (RAS) or compared to the unspliced form (RANS): N_s: number of spliced reads corresponding to the precise excision of the focal intron; N_a: number of reads corresponding to alternative splice variants relative to this intron (*i.e.* sharing only one of the two intron boundaries); N_u: number of unspliced reads, co-linear with the genomic sequence. **C**: Definitions of the main variables used in this study. The definition of the variables corresponds to the one provided in Bénitière *et al.* (2023).

For this step, HISAT2 version 2.1.0 was employed to align RNA-seq reads to the respective reference
genomes (Kim *et al.*, 2019). To enhance the sensitivity of splice junction detection, we constructed genome
indexes incorporating annotated intron and exon coordinates along with genome sequences. The maximum
permitted intron length was set at 2,000,000 base pairs. The processed and compressed files generated during
this procedure can amass a size exceeding 20 terabytes.

We extracted intron coordinates from the HISAT2 alignments, utilizing a custom Perl script that scanned
for CIGAR strings containing "N" characters, which indicate skipped regions in the reference sequence. For
intron identification and quantification, we exclusively utilized uniquely mapped reads with a maximum
mismatch fraction of 0.02. In the context of new intron identification, we imposed a minimum anchor length
(*i.e.*, part of the read that spans each of the two exons flanking a given intron) of 8 base pairs. We then
quantified intron splicing frequencies by including aligned reads with a minimum anchor length of 5 base
pairs. We retained predicted introns exhibiting GT-AG, GC-AG, or AT-AC splice signals and determined the
intron strand based on the splice signal.

Introns were assigned to genes if at least one of their boundaries was within 1 base pair of annotated exon
coordinates, combined across all isoforms for each gene. Intron assignments were limited to those that could
be unambiguously associated with a single gene. Notably, we differentiated between annotated introns,
present in the reference genome annotations, and unannotated introns, identified through RNA-seq data and
assigned to previously annotated genes.

We identified introns situated within protein-coding regions. To do this, for each protein-coding gene, we
extracted annotated start and stop codon positions across all annotated isoforms. The minimum start codon
and maximum end codon positions were identified, and introns located upstream or downstream of these
extreme coordinates were considered as interrupting untranslated regions.

Alternative splicing variables

For each intron, we recorded two key variables: N_s representing the number of reads corresponding to the
precise removal of the intron (referred to as spliced reads), and N_a representing the count of reads supporting
alternative splicing events (*i.e.* spliced variants sharing only one of the two boundaries of the focal intron).
Additionally, we denoted N_u as the count of unspliced reads that align linearly with the genomic sequence and
span at least 10 base pairs on both sides of an exon-intron junction. These definitions are visually clarified in
(Fig. 3B,C). Subsequently, we introduced the relative measurement of the target intron's abundance compared
to introns with a single alternative splice boundary (RAS $= \frac{N_s}{N_s + N_a}$), as well as relative to unspliced reads
(RANS $= \frac{N_s}{N_s + \frac{N_u}{2}}$).

To compute these ratios, we required at least 10 reads in their denominators. Thus, we computed the RAS
only when $(N_s + N_a) \geq 10$, and the RANS only when $(N_s + \frac{N_u}{2}) \geq 10$. We divided N_u by 2 because unspliced
reads that span the two intron boundaries likely refer to the same intron retention event. If these conditions
were not met, the resulting values were designated as unavailable (NA). These ratios were computed utilizing
data from all available RNA-seq samples, unless explicitly specified (*e.g.* in sub-sampling analyses). Based on
these ratios, we divided introns into three categories: major-isoform introns, defined as those introns that
have RANS > 0.5 and RAS > 0.5 (these likely correspond to the introns of major isoforms (Gonzàlez-Porta
et al., 2013; Tress *et al.*, 2017; Bénitière *et al.*, 2023)); minor-isoform introns, defined as those introns that
have RANS ≤ 0.5 or RAS ≤ 0.5 (these introns are detected in a minority of transcripts); unclassified introns,
which do not satisfy the above conditions.

Gene expression estimation

Gene expression levels were computed using Cufflinks version 2.2.1 (Trapnell *et al.*, 2010; Roberts *et al.*, 2011), utilizing the read alignments obtained with HISAT2 for each individual RNA-seq sample. We thus evaluated gene expression levels with the Fragment *Per* Kilobase of exon *per* Million mapped reads (FPKM) method. To determine the representative expression level of each gene, the mean FPKM was calculated across all samples, taking into consideration the sequencing depth of each sample, called 'weighted FPKM'. We used this measure to evaluate the relationship between alternative splicing rates and gene expression levels, within each species.

Estimation of the sequencing depth

We determined for each gene the union of all annotated exon coordinates (termed here exon blocks). Using bedtools v2.25.0 (Quinlan and Hall, 2010), we assessed the read coverage at each position of the exon blocks. The average exonic *per*-base read coverage was subsequently computed for each gene. The sequencing depth of a given sample was evaluated through the median *per*-base read coverage across BUSCO (Benchmarking Universal Single-Copy Orthologs) genes.

Data visualisation using a Shiny app

A Shiny app available at https://lbbe-shiny.univ-lyon1.fr/ShinyApp-GTDrift/ was deployed to allow users to visualize and compare the summarized data (Chang *et al.*, 2023). Most of the graphics shown in this paper are directly reproducible from the app. In this app, users can also visualize intra-species variables, for example comparing introns or gene characteristics. Furthermore, a specific tab is dedicated to the investigation of gene structure in relation to the splicing attributes found in the underlying database. Users can also visualize the phylogenetic tree and employ these trees for conducting Phylogenetic Generalized Least Square regression analyses.

The app is organized in several panels or "tabs" in the web page.

The tab 'Inter-species graphics' facilitates the comparison of genome characteristics across different species through graphical representation. Additionally, users have the option to upload their own data in a tab-separated text format, where each species is represented in a separate row, with the variables of interest organized in columns. An example of such a tabular dataset can be found in the repository.

The 'Inter-species Axis' tab explains the variables available in the 'Inter-species graphics' tab.

The 'Intra-species graphics' tab permits the exploration of characteristics within a species, focusing on introns or on genes. Furthermore, users have the ability to retrieve metadata related to BUSCO annotation, gene expression, or intron splicing events (see Methods).

The 'Intra-species Axis' tab describes the variables featured in the 'Intra-species graphics' tab.

Within the 'Gene structure' tab, users can delve into the introns detected in RNA-seq alignments for a specific gene. These introns are color-coded based on various criteria, including their location within the CDS or outside of it, as well as whether they are classified as major or minor-isoform introns (see Methods).

336 The 'Phylogenetic tree' tab facilitates the examination of phylogenetic trees used for conducting Phylogenetic
337 Generalized Least Squares regression within the 'Inter-species graphics' tab.

Data and code availability

339 All processed data that we generated and used in this study, as well as the scripts that we used to analyze
340 the data and to generate the figures, are available at the following Zenodo DOI: https://doi.org/10.5281/
341 zenodo.10022493. The database is provided on Zenodo with the DOI: https://doi.org/10.5281/zenodo.
342 10017653. Finally, the Shiny app is available at: https://lbbe-shiny.univ-lyon1.fr/ShinyApp-GTDrift/
343 and on Zenodo with the DOI: https://doi.org/10.5281/zenodo.10022520.

Results

Description of the data available in GTDrift

In GTDrift, we provide a manageable number of compressed data tables for each species processed via our
pipeline (Fig. 4). Tables are stored in tab-delimited text format, which makes them easy to access for users
with experience in bioinformatics. They are user-friendly because of the simplicity of their contents. To access
these tables, users can visit the Zenodo DOI: https://doi.org/10.5281/zenodo.10017653 and select their
desired data type. The data can also be easily explored through a web application written in Shiny at
https://lbbe-shiny.univ-lyon1.fr/ShinyApp-GTDrift/. Data exploration is thus easily accessible even
for users who do not have a background in bioinformatics.

Our database is centered around transcriptomics data. At the time of publication, the database contained
over 15,935 RNA-seq samples distributed over 491 embryophytes and metazoans (Fig. 1), providing gene
expression and alternative splicing events data. Additionally, we have enriched the database with annotations
for orthologous single-copy genes (BUSCO genes) and proxies of effective population size, including the
molecular evolutionary rate dN/dS and life history traits such as longevity, body mass, body length. We
used similar types of data in our recent publication exploring the relationship between random genetic drift
and alternative splicing patterns (Bénitière *et al.*, 2023). However, here we provide considerably more data,
for 1,507 species compared to 53 in this publication.

Below, we provide information on the data types that are currently available in GTDrift.

Life history traits

363 The table labeled 'life_history_traits.tab' comprises values pertaining to three distinct traits (longevity, body
364 mass, and body weight), for 969 species. This table includes bibliographic references which attribute these
365 values to each species. The species are defined by their scientific names and by the corresponding NCBI
366 taxonomy identifier (taxID).

Figure 4: **Description of the bioinformatic analysis pipeline.** (Adapted from Supplementary Fig. 11 Bénitière *et al.* (2023)) First, we retrieved genomic sequences and annotations from the NCBI Genomes database. We aligned RNA-seq reads on the corresponding reference genomes with HISAT2. We used these alignments to estimate various variables related to splicing patterns (see Fig. 2), to compute the AS rate, and to estimate gene expression using Cufflinks. To compute the dN/dS ratios, we first identified BUSCO genes with BUSCOv3 and aligned their coding sequences (CDS) using PRANK (codon model). We reconstructed a phylogenetic tree using RAxML-NG. Using bio++, we estimated dN/dS along the phylogenetic tree on concatenated alignments. This pipeline was previously used in Bénitière *et al.* (2023).

Protein-coding sequence evolution features

We provide estimates of the representative dN/dS ratio for most species (N=1,324 species after filtering for a sufficient number of annotated orthologous genes). The data are available in the directory 'dNdS'.

We provide the phylogenetic tree of the studied species, with the dN/dS ratios as branch lengths, in the Newick file format. We provide this data separately for the four approaches used to estimate the ratios dN/dS, using the eukaryota, embryophyta or metazoa BUSCO gene sets, or a different gene set for each clade (Methods). In addition, we provide a table comprising the dN and dS values for each terminal branch of the phylogenetic tree, along with the species scientific name and NCBI taxonomy ID, for each of the four approaches.

Gene expression

In the 'Transcriptomic' directory, each species is represented by a dedicated table named 'by_gene_analysis.tab.gz'. This table contains annotated gene coordinates, the mean and median FPKM (Fragments *Per* Kilobase of exon *per* Million mapped reads) across samples. Additionally, the table includes information about RNA-seq read coverage for exonic regions for each gene, including the total read coverage across samples. The individual gene expression data for each RNA-seq experiment can be accessed within the 'RUN' directory. The data are provided in a separate directory for each SRA accession number. The file 'by_gene_db.tab.gz' containing the exon coverage and the FPKM measured for each gene corresponding in line to the previous file 'by_gene_analysis.tab.gz'.

Alternative splicing data

For each species, we provide a summarized table named 'by_intron_analysis.tab.gz', containing for each intron the cumulative counts of spliced reads (N_s), the number of reads supporting alternative splicing of this introns (N_a), and the number of unspliced reads overlapping with this intron (N_u) detected through RNA-seq analysis (Methods). This table contains data combined across all analyzed RNA-seq samples. Detailed information for individual RNA-seq experiments can be found within the 'RUN' directory, in the file 'by_intron_db.tab.gz'. In these files, introns are listed in the same order as in the file 'by_intron_analysis.tab.gz'.

RNA-seq sample description

In the file named 'SRAruninfo.tab', we provide information extracted from the SRA database, for each RNA-seq sample. Depending on the sample, this information can include the library source, the tissue from which the sample is derived, the sex of the sampled individual, the lab that conducted the analysis, the methods used to prepare the library, *etc*.

BUSCO gene identification

In the directory 'BUSCO_annotations', we provide the correspondence between NCBI gene identifiers and BUSCO gene identifiers, determined for three distinct BUSCO datasets: eukaryota, metazoa, and embryophyta.

Data quality validation

Acquiring life history traits

To facilitate the acquisition of life history traits, we have devised and shared a pipeline that uses an automatic screening technique complemented by a Machine Learning method.

To assess the effectiveness of the automatic screening technique that we used to extract life history traits from various databases, we conducted a comparative analysis, contrasting it with the manual methodology. We also compared it to the Machine Learning (ML) approach for the ADW database. The screening procedure yielded accurate information with varying false positive rates depending on the source database, as follows: AnAge (98.9% accuracy; 0% false positive), fishbase (100%; 0.2%), EOL (94.5%; 0.18%), and ADW (88%;

Figure 5: N_e **proxies.** **A**: Relationship between body length (cm, log scale) and longevity (days, log scale) of the organism. Each dot represents one species (colored by clade). **B**: Relationship between body length (cm, log scale) and the body weight (kg, log scale). **A,B**: Pagel's *lambda* model is used to take into account the phylogenetic structure of the data in a regression model.

5.4%). These results highlight the utility of our screening pipeline for identifying three key life history traits
across AnAge, EOL, ADW, and fishbase databases.

For the ADW database, the ML approach exhibited a slight advantage over the screening method, and
its results did not completely align with those obtained through the screening approach. Specifically, for
life history traits, the ML approach correctly retrieved 89.7% of the results obtained through the manual
approach, while introducing a 9.2% false positive rate.

When combining both the ML approach and the screening process, we achieved a 95% accuracy rate in
identifying positive cases. However, a 7% error rate persisted in this merged approach.

In GTDrift, we provide data corresponding to a synthesis of the three methodologies including only manually-
checked values (Methods).

Estimating the intensity of random drift

As expected, a positive correlation is observed in Fig. 5A,B between the different life history traits, used as
indirect predictors of the effective population size (N_e) (Romiguier *et al.*, 2014; Waples, 2016; Figuet *et al.*,
2016; Galtier, 2016; Weyna and Romiguier, 2020).

When examining the dN/dS ratio across distinct time scales and using various BUSCO datasets, we
consistently observe comparable dN/dS ratios at terminal branches. This uniformity across a range of
methodological approaches highlights their concordance (Fig. 6A,B).

Furthermore, the observed dN/dS ratio are significantly correlated with proxies derived from life history
traits (Fig. 7A,B) as reported in the literature (Romiguier *et al.*, 2014; Figuet *et al.*, 2016).

Figure 6: **Reproducibility of the** dN/dS **ratio. A**: Relation between the dN/dS ratio on terminal branches of the phylogenetic tree of the metazoa set compared to the ones measured in the *per* clades set. **B**: Relation between the dN/dS ratio on terminal branches of the phylogenetic tree of the eukaryota set compared to the ones measured in the embryophyta and the metazoa set. **A,B**: LM stands for Linear regression Model and Pagel's *lambda* model is used to take into account the phylogenetic structure of the data in a regression model.

Quality of genome annotations

To assess gene expression levels and alternative splicing patterns, the quality of genome annotations is of paramount importance. We evaluated genome annotation quality by examining the presence of BUSCO genes. We note that the results depend on the BUSCO dataset that is used as a starting point. When using the BUSCO dataset designed for eukaryota, which comprises 303 genes, we have effectively identified nearly all single-copy orthologous genes, and this feature exhibits a high degree of homogeneity across different species (Fig. 8). However, the aves clade demonstrates a deficiency in the number of BUSCO genes compared to the anticipated count based on BUSCO expectations. This is expected given the known genome incompleteness problem for this clade, due to the presence of GC-rich chromosomes (Li *et al.*, 2022).

Because the eukaryota BUSCO gene set is limited, we also performed gene identification for the metazoa and embryophyta BUSCO datasets, leading to substantially larger collections of genes. Specifically, we detected 978 BUSCO genes for the metazoa dataset and 1,440 genes for the embryophyta dataset.

Spliced introns classification

A significant body of literature has consistently reported that the majority of genes typically exhibit one predominant isoform (Gonzàlez-Porta *et al.*, 2013; Tress *et al.*, 2017). This isoform is commonly termed "major isoform". Here, we aimed to assess the influence of sequencing depth on the identification of major-isoform introns, that is, those introns that belong to major isoforms (see Alternative splicing variables). Employing the model organism *Drosophila melanogaster*, we randomly selected between 1 and 20 RNA-seq samples. For each subset of samples, we computed the median read coverage across the exons of BUSCO genes, providing a standardized measure of transcriptome sequencing depth that can be compared across different species. Additionally, we tallied the count of introns falling into various categories (major-isoform introns,

Figure 7: **Interplay between N_e proxies.** Correlation between the dN/dS ratio on terminal branches of the phylogenetic tree of the *per* clade set and life history traits: longevity (days, log scale) (**A**), body weight (kg, log scale) (**B**), body length (cm, log scale) (**C**). **A,B,C**: Pagel's *lambda* model is used to take into account the phylogenetic structure of the data in a regression model.

minor-isoform introns or unclassified introns - see Methods) for each subset of samples. This entire process was repeated 10 times (Fig. 9A).

As expected, we observed that the number of major-isoform introns that could be identified increased with greater sequencing depth until it reached a threshold of 200 read coverage *per* base (Fig. 9A). Beyond this threshold, no additional major-isoform introns are discernible. Simultaneously, the count of unclassified introns decreased to nearly zero, indicating that introns newly detected above the 200-read coverage threshold predominantly consisted of minor-isoform introns that shared a boundary with a major intron. Indeed, the count of minor-isoform introns continued to rise steadily beyond this point.

We then assessed the proportion of annotated introns that fall within the categories defined above. Our results reveal that the majority of species exhibit well-annotated major-isoform introns, indicating the accuracy of the intron annotation (Fig. 9B). Additionally, as sequencing depth increases, we observed a decreasing fraction of annotated minor-isoform introns. This trend is consistent with expectations, given that higher sequencing depth expands the pool of rare variants and potential spontaneous errors that may not have been previously observed. It is important to note that there appears to be no inherent limit to this phenomenon, as the intricacies of alternative splicing machinery can give rise to unpredictable errors (Bénitière *et al.*, 2023).

Figure 8: **BUSCO genes annotation.** Proportion of BUSCO genes, from the BUSCO gene set eukaryota (N=303 genes), identified in each species.

Discussion

GTDrift is a comprehensive data resource facilitating investigations of genomic and transcriptomic characteristics alongside indicators of genetic drift intensity for distinct species. Notably, this resource offers information on life history traits, including longevity, adult body length, and body mass, for a curated set of 969 species. Additionally, it provides estimates of the ratio between the rate of non-synonymous substitutions over synonymous substitutions (dN/dS) for 1,324 species.

For individual species, intron-centered alternative splicing frequencies, gene expression levels, and sequencing depth statistics have been systematically quantified and shared, encompassing more than 15,935 RNA-seq samples across 491 species. To enable cross-species comparisons, orthology predictions for conserved single-copy genes are provided, based on BUSCO gene sets, encompassing a total of 1,507 eukaryotic species, including 1,414 animals and 93 green plants, along with phylogenetic trees to account for phylogenetic inertia.

The number of species per data type varies due to different limitations: availability of life history traits data; completeness of gene annotations for dN/dS calculation; computational resources and availability of RNA-seq samples for transcriptomic analysis (Fig. 4).

These pre-processed data streamlines the work for those interested in investigating the impact of drift on biological processes across a wide range of species. All data are provided in flat files, which enable downstream computational analyses and render GTDrift mainly aimed at users with some computational

Figure 9: **Sequencing depth impact on intron classification. A**: Number of major (RANS > 0.5 and RAS > 0.5), minor (RANS $\geq$ 5% or RAS $\geq$ 5%) and unclassified introns for *Drosophila melanogaster*. The sequencing depth is measured by taking the median *per*-base read coverage across BUSCO genes from eukaryota gene set. **B**: *Per* species major-isoform introns, minor-isoform introns and undetermined introns (N_s) $\geq$ 10) annotated proportion and sequencing depth measured by taking the median *per*-base read coverage eukaryota BUSCO genes.

482   skills. Nonetheless, to enhance accessibility, we have developed a user-friendly Shiny app that facilitates
483   database exploration and allows for species-specific data downloads (available at `https://lbbe-shiny.univ-`
484   `lyon1.fr/ShinyApp-GTDrift/`).

Cautionary considerations in utilizing N_e proxies

Users should bear in mind that the scientific community has yet to establish the most adequate proxies for effective population size. A prominent hypothesis suggests that these proxies are associated with the number of individuals (N). Indeed, species with greater longevity and larger body mass tend to be less abundant within their ecological niche due to resource (mass) and spatial (length, mass) requirements (Damuth, 1981; Nee *et al.*, 1991; White *et al.*, 2007). Therefore, variations in life history traits should correspond to variations in the number of individuals (N), which subsequently impact N_e.

When using the dN/dS ratio as a proxy for N_e, rather than focusing on correlations with the population census, we evaluated the efficiency of natural selection to purge deleterious mutations. This efficiency can be represented as the product of N_e and 's', which denotes the selection coefficient. The extent to which a well-estimated dN/dS ratio can be considered as a proxy for N_e remains a subject of debate. Notably, when the rate of synonymous substitutions (dS) exceeds 1, it indicates a point of saturation where multiple substitutions occur *per* site, rendering dS susceptible to considerable noise due to the challenge of accurately identifying the number of substitutions at given sites. In such cases, the dN component can often still be reliably determined. Given that non-synonymous substitutions have a lower rate compared to synonymous ones, dN reaches a saturation point at a later stage.

Moreover, when the evolutionary time frame is relatively short, characterized by small dS values, the variants under examination are primarily attributed to polymorphism rather than fixed substitutions. In such cases, we are not effectively measuring substitution rates. Consequently, the discussion also revolves around determining a divergence threshold, above which we could assume that dS and dN predominantly represent substitutions, with minimal influence from polymorphism. In this perspective, the expanding polymorphism data could potentially serve as a means to distinguish between polymorphism and substitutions, offering a more efficient approach to investigate dN/dS (Mugal *et al.*, 2014).

Overall, we found that the various N_e proxies were significantly correlated, even when accounting for the underlying phylogenetic structure. Thus, our dataset, which encompasses information on dN and dS across all branches of the phylogenetic trees, holds the potential to estimate the long-term effective population size (N_e) and its interaction with life history traits over time.

Comparing transcriptomic data

In our study, we have identified BUSCO genes for the eukaryota, metazoa, or embryophyta BUSCO reference gene sets. To ensure meaningful comparisons between species with a sufficient number of detected BUSCO genes, we evaluated the median RNA-seq coverage of these BUSCO genes. As demonstrated in Data quality validation, the median *per*-base read exonic RNA-seq coverage of BUSCO genes is a good indicator of the power to detect alternative splicing patterns. We believe that, for the inclusion of additional species, an examination of the RNA-seq read coverage on BUSCO genes is needed to ensure that we could identify major-isoform introns and analyze alternative splicing patterns.

Additionally, it is essential to assess the completeness of the genome and of the annotation, which can be estimated based on the number of identified BUSCO genes. Some species may have a limited number of well-annotated BUSCO genes, or global gene duplications may result in the presence of two copies of a BUSCO gene, which no longer qualifies as a single copy gene.

Our RNA-seq description table offers users access to information collected from the Sequence Read Archive (SRA) for the RNA-seq datasets under study. This table enables users to filter and select RNA-seq data that align with their specific research needs. Users can tailor their selection based on factors such as sex, tissue, or protocol. Depending on the research question that is asked, it may be important to extract and analyse RNA-seq samples that were generated for the same biological conditions. We provide this information so that GTDrift users are able to filter the data as needed.

To facilitate cross-species comparisons, especially in the context of alternative splicing and gene expression, users can make use of BUSCO gene sets, which should exhibit consistent expression patterns, functionality, and evolutionary constraints across diverse species. However, users should thoroughly validate this assumption and proceed with vigilance.

Conclusion

In conclusion, we are confident that the GTDrift database can be a valuable resource for studies aiming to investigate the relationship between the intensity of genetic drift, genomic and transcriptomic characteristics.

Acknowledgements

We thank Loïc Guille for his contribution to an initial pilot study, Tristan Lefébure for insightful discussions and Laurent Guéguen for his help on dN/dS analyses. Computational analyses were performed using the computing facilities of the CC LBBE/PRABI and the Core Cluster of the Institut Français de Bioinformatique (IFB) (ANR-11-INBS-0013). Silhouette images of taxonomic Families originate from PhyloPic developed and maintained by Mike Keesey available at https://www.phylopic.org/.

Author contributions statement

F.B. conceived the pipeline and conducted the analyses. F.B. and A.N. drafted the manuscript. All authors reviewed the manuscript.

Funding

This work was funded by the French National Research Agency (ANR-20-CE02-0008-01 "NeGA" and ANR-17-CE12-0019-01 "LncEvoSys").

Competing interests

The authors declare no conflicts of interest.

References

2023. Hugging Face – The AI community building the future.

Barbosa-Morais, N. L., Irimia, M., Pan, Q., Xiong, H. Y., Gueroussov, S., Lee, L. J., Slobodeniuc, V., Kutter, C., Watt, S., Colak, R., Kim, T., Misquitta-Ali, C. M., Wilson, M. D., Kim, P. M., Odom, D. T., Frey, B. J., and Blencowe, B. J. 2012. The evolutionary landscape of alternative splicing in vertebrate species. *Science (New York, N.Y.)*, 338(6114): 1587–1593.

Bastian, F. B., Roux, J., Niknejad, A., Comte, A., Fonseca Costa, S., de Farias, T. M., Moretti, S., Parmentier, G., de Laval, V. R., Rosikiewicz, M., Wollbrett, J., Echchiki, A., Escoriza, A., Gharib, W. H., Gonzales-Porta, M., Jarosz, Y., Laurenczy, B., Moret, P., Person, E., Roelli, P., Sanjeev, K., Seppey, M., and Robinson-Rechavi, M. 2021. The Bgee suite: integrated curated expression atlas and comparative transcriptomics in animals. *Nucleic Acids Research*, 49(D1): D831–D847.

Berget, S. M., Moore, C., and Sharp, P. A. 1977. Spliced segments at the 5' terminus of adenovirus 2 late mRNA. *Proceedings of the National Academy of Sciences*, 74(8): 3171–3175. Publisher: Proceedings of the National Academy of Sciences.

Bolívar, P., Guéguen, L., Duret, L., Ellegren, H., and Mugal, C. F. 2019. GC-biased gene conversion conceals the prediction of the nearly neutral theory in avian genomes. *Genome Biology*, 20(1): 5.

Bénitière, F., Necsulea, A., and Duret, L. 2023. Random genetic drift sets an upper limit on mRNA splicing accuracy in metazoans. Pages: 2022.12.09.519597 Section: New Results.

Cardoso-Moreira, M., Halbert, J., Valloton, D., Velten, B., Chen, C., Shao, Y., Liechti, A., Ascenção, K., Rummel, C., Ovchinnikova, S., Mazin, P. V., Xenarios, I., Harshman, K., Mort, M., Cooper, D. N., Sandi, C., Soares, M. J., Ferreira, P. G., Afonso, S., Carneiro, M., Turner, J. M. A., VandeBerg, J. L., Fallahshahroudi, A., Jensen, P., Behr, R., Lisgo, S., Lindsay, S., Khaitovich, P., Huber, W., Baker, J., Anders, S., Zhang, Y. E., and Kaessmann, H. 2019. Gene expression across mammalian organ development. *Nature*, 571(7766): 505–509.

Chang, W., Cheng, J., Allaire, J. J., Sievert, C., Schloerke, B., Xie, Y., Allen, J., McPherson, J., Dipert, A., and Borges, B. 2023. *shiny: Web Application Framework for R*.

Charif, D. and Lobry, J. R. 2007. SeqinR 1.0-2: A Contributed Package to the R Project for Statistical Computing Devoted to Biological Sequences Retrieval and Analysis. In U. Bastolla, M. Porto, H. E. Roman, and M. Vendruscolo, editors, *Structural Approaches to Sequence Evolution: Molecules, Networks, Populations*, Biological and Medical Physics, Biomedical Engineering, pages 207–232. Springer, Berlin, Heidelberg.

Chen, L., Bush, S. J., Tovar-Corona, J. M., Castillo-Morales, A., and Urrutia, A. O. 2014. Correcting for Differential Transcript Coverage Reveals a Strong Relationship between Alternative Splicing and Organism Complexity. *Molecular Biology and Evolution*, 31(6): 1402–1413.

Damuth, J. 1981. Population density and body size in mammals. *Nature*, 290(5808): 699–700.

Dutheil, J. and Boussau, B. 2008. Non-homogeneous models of sequence evolution in the Bio++ suite of libraries and programs. *BMC Evolutionary Biology*, 8(1): 255.

Dutheil, J. Y., Galtier, N., Romiguier, J., Douzery, E. J. P., Ranwez, V., and Boussau, B. 2012. Efficient selection of branch-specific models of sequence evolution. *Molecular Biology and Evolution*, 29(7): 1861–1874.

Figuet, E., Nabholz, B., Bonneau, M., Mas Carrio, E., Nadachowska-Brzyska, K., Ellegren, H., and Galtier, N. 2016. Life History Traits, Protein Evolution, and the Nearly Neutral Theory in Amniotes. *Molecular Biology and Evolution*, 33(6): 1517–1527.

Froese, R. and Pauly, D. 2023. FishBase. Published: World Wide Web electronic publication.

Galtier, N. 2016. Adaptive Protein Evolution in Animals and the Effective Population Size Hypothesis. *PLOS Genetics*, 12(1): e1005774.

Gonzàlez-Porta, M., Frankish, A., Rung, J., Harrow, J., and Brazma, A. 2013. Transcriptome analysis of human tissues and cell lines reveals one dominant transcript per gene. *Genome Biology*, 14(7): 1–11. Number: 7 Publisher: BioMed Central.

Gout, J.-F., Thomas, W. K., Smith, Z., Okamoto, K., and Lynch, M. 2013. Large-scale detection of in vivo transcription errors. *Proceedings of the National Academy of Sciences*, 110(46): 18584–18589. Publisher: Proceedings of the National Academy of Sciences.

Graur, D., Li, W.-H. L., Graur, D., and Li, W.-H. L. 2000. *Fundamentals of Molecular Evolution*. Oxford University Press, Oxford, New York, second edition, second edition edition.

Guéguen, L. and Duret, L. 2018. Unbiased Estimate of Synonymous and Nonsynonymous Substitution Rates with Nonstationary Base Composition. *Molecular Biology and Evolution*, 35(3): 734–742.

Guéguen, L., Gaillard, S., Boussau, B., Gouy, M., Groussin, M., Rochette, N. C., Bigot, T., Fournier, D., Pouyet, F., Cahais, V., Bernard, A., Scornavacca, C., Nabholz, B., Haudry, A., Dachary, L., Galtier, N., Belkhir, K., and Dutheil, J. Y. 2013. Bio++: efficient extensible libraries and tools for computational molecular evolution. *Molecular Biology and Evolution*, 30(8): 1745–1750.

Kim, D., Paggi, J. M., Park, C., Bennett, C., and Salzberg, S. L. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, 37(8): 907–915. Number: 8 Publisher: Nature Publishing Group.

Kimura, M., Maruyama, T., and Crow, J. F. 1963. The Mutation Load in Small Populations. *Genetics*, 48(10): 1303–1312.

Kozlov, A. M., Darriba, D., Flouri, T., Morel, B., and Stamatakis, A. 2019. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*, 35(21): 4453–4455.

Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M., Stecher, G., and Hedges, S. B. 2022. TimeTree 5: An Expanded Resource for Species Divergence Times. *Molecular Biology and Evolution*, 39(8): msac174.

Lefébure, T., Morvan, C., Malard, F., François, C., Konecny-Dupré, L., Guéguen, L., Weiss-Gayet, M., Seguin-Orlando, A., Ermini, L., Sarkissian, C. D., Charrier, N. P., Eme, D., Mermillod-Blondin, F., Duret, L., Vieira, C., Orlando, L., and Douady, C. J. 2017. Less effective selection leads to larger genomes. *Genome Research*, 27(6): 1016–1028.

Leinonen, R., Sugawara, H., and Shumway, M. 2011. The Sequence Read Archive. *Nucleic Acids Research*, 39(Database issue): D19–D21.

Li, M., Sun, C., Xu, N., Bian, P., Tian, X., Wang, X., Wang, Y., Jia, X., Heller, R., Wang, M., Wang, F., Dai, X., Luo, R., Guo, Y., Wang, X., Yang, P., Hu, D., Liu, Z., Fu, W., Zhang, S., Li, X., Wen, C., Lan, F., Siddiki, A. Z., Suwannapoom, C., Zhao, X., Nie, Q., Hu, X., Jiang, Y., and Yang, N. 2022. De Novo Assembly of 20 Chicken Genomes Reveals the Undetectable Phenomenon for Thousands of Core Genes on Microchromosomes and Subtelomeric Regions. *Molecular Biology and Evolution*, 39(4): msac066.

Li, Z., Zhang, Y., Bush, S. J., Tang, C., Chen, L., Zhang, D., Urrutia, A. O., Lin, J.-w., and Chen, L. 2021. MeDAS: a Metazoan Developmental Alternative Splicing database. *Nucleic Acids Research*, 49(D1): D144–D150.

Liu, J., Yin, F., Lang, K., Jie, W., Tan, S., Duan, R., Huang, S., and Huang, W. 2022. MetazExp: a database for gene expression and alternative splicing profiles and their analyses based on 53 615 public RNA-seq samples in 72 metazoan species. *Nucleic Acids Research*, 50(D1): D1046–D1054.

Liu, Z. and Zhang, J. 2018a. Human C-to-U Coding RNA Editing Is Largely Nonadaptive. *Molecular Biology and Evolution*, 35(4): 963–969.

Liu, Z. and Zhang, J. 2018b. Most m6A RNA Modifications in Protein-Coding Regions Are Evolutionarily Unconserved and Likely Nonfunctional. *Molecular Biology and Evolution*, 35(3): 666–675.

Lynch, M. 2007. The frailty of adaptive hypotheses for the origins of organismal complexity. *Proceedings of the National Academy of Sciences*, 104(suppl_1): 8597–8604. Publisher: Proceedings of the National Academy of Sciences.

Lynch, M. 2010. Evolution of the mutation rate. *Trends in genetics : TIG*, 26(8): 345–352.

Lynch, M. and Conery, J. S. 2003. The origins of genome complexity. *Science (New York, N.Y.)*, 302(5649): 1401–1404.

Lynch, M., Ackerman, M. S., Gout, J.-F., Long, H., Sung, W., Thomas, W. K., and Foster, P. L. 2016. Genetic drift, selection and the evolution of the mutation rate. *Nature Reviews Genetics*, 17(11): 704–714. Number: 11 Publisher: Nature Publishing Group.

Löytynoja, A. and Goldman, N. 2008. Phylogeny-Aware Gap Placement Prevents Errors in Sequence Alignment and Evolutionary Analysis. *Science*, 320(5883): 1632–1635. Publisher: American Association for the Advancement of Science.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., and Zdobnov, E. M. 2021. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution*, 38(10): 4647–4654.

Merkin, J., Russell, C., Chen, P., and Burge, C. B. 2012. Evolutionary dynamics of gene and isoform regulation in Mammalian tissues. *Science (New York, N.Y.)*, 338(6114): 1593–1599.

Mudge, J. M., Frankish, A., Fernandez-Banet, J., Alioto, T., Derrien, T., Howald, C., Reymond, A., Guigó, R., Hubbard, T., and Harrow, J. 2011. The Origins, Evolution, and Functional Potential of Alternative Splicing in Vertebrates. *Molecular Biology and Evolution*, 28(10): 2949–2959.

Mugal, C. F., Wolf, J. B., and Kaj, I. 2014. Why Time Matters: Codon Evolution and the Temporal Dynamics of dN/dS. *Molecular Biology and Evolution*, 31(1): 212–231.

Myers, P., Espinosa, R., Parr, C. S., Jones, T., Hammond, G. S., and Dewey, T. A. 2023. The Animal Diversity Web (online). Published: Accessed at https://animaldiversity.org.

Mölder, F., Jablonski, K. P., Letcher, B., Hall, M. B., Tomkins-Tinch, C. H., Sochat, V., Forster, J., Lee, S., Twardziok, S. O., Kanitz, A., Wilm, A., Holtgrewe, M., Rahmann, S., Nahnsen, S., and Köster, J. 2021. Sustainable data analysis with Snakemake. Technical Report 10:33, F1000Research. Type: article.

NCBI Resource Coordinators 2018. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Research*, 46(D1): D8–D13.

Nee, S., Read, A. F., Greenwood, J. J. D., and Harvey, P. H. 1991. The relationship between abundance and body size in British birds. *Nature*, 351(6324): 312–313. Number: 6324 Publisher: Nature Publishing Group.

Nielsen, R. and Yang, Z. 2003. Estimating the Distribution of Selection Coefficients from Phylogenetic Data with Applications to Mitochondrial and Viral DNA. *Molecular Biology and Evolution*, 20(8): 1231–1239.

Ohta, T. 1973. Slightly Deleterious Mutant Substitutions in Evolution. *Nature*, 246(5428): 96–98. Number: 5428 Publisher: Nature Publishing Group.

Ohta, T. 1992. The Nearly Neutral Theory of Molecular Evolution. *Annual Review of Ecology and Systematics*, 23: 263–286. Publisher: Annual Reviews.

Parr, C. S., Wilson, N., Leary, P., Schulz, K., Lans, K., Walley, L., Hammock, J., Goddard, A., Rice, J., Studer, M., Holmes, J., Corrigan, R., and Jr 2014. The Encyclopedia of Life v2: Providing Global Access to Knowledge About Life on Earth. *Biodiversity Data Journal*, 2: e1079. Publisher: Pensoft Publishers.

Pickrell, J. K., Pai, A. A., Gilad, Y., and Pritchard, J. K. 2010. Noisy Splicing Drives mRNA Isoform Diversity in Human Cells. *PLOS Genetics*, 6(12): e1001236. Publisher: Public Library of Science.

Quinlan, A. R. and Hall, I. M. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics (Oxford, England)*, 26(6): 841–842.

Reyes, A., Anders, S., Weatheritt, R. J., Gibson, T. J., Steinmetz, L. M., and Huber, W. 2013. Drift and conservation of differential exon usage across tissues in primate species. *Proceedings of the National Academy of Sciences*, 110(38): 15377–15382. Publisher: Proceedings of the National Academy of Sciences.

Roberts, A., Pimentel, H., Trapnell, C., and Pachter, L. 2011. Identification of novel transcripts in annotated genomes using RNA-Seq. *Bioinformatics*, 27(17): 2325–2329.

Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., Chiari, Y., Dernat, R., Duret, L., Faivre, N., Loire, E., Lourenco, J. M., Nabholz, B., Roux, C., Tsagkogeorga, G., Weber, A. A.-T., Weinert, L. A., Belkhir, K., Bierne, N., Glémin, S., and Galtier, N. 2014. Comparative population genomics in animals uncovers the determinants of genetic diversity. *Nature*, 515(7526): 261–263.

Saudemont, B., Popa, A., Parmley, J. L., Rocher, V., Blugeon, C., Necsulea, A., Meyer, E., and Duret, L. 2017. The fitness cost of mis-splicing is the main determinant of alternative splicing patterns. *Genome Biology*, 18.

Sayers, E. W., Bolton, E. E., Brister, J. R., Canese, K., Chan, J., Comeau, D. C., Connor, R., Funk, K., Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z., Thibaud-Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., Tse, T., Wang, J., Williams, R., Trawick, B. W., Pruitt, K. D., and Sherry, S. T. 2022. Database resources of the national center for biotechnology information. *Nucleic Acids Research*, 50(D1): D20–D26.

Seppey, M., Manni, M., and Zdobnov, E. M. 2019. BUSCO: Assessing Genome Assembly and Annotation Completeness. *Methods in Molecular Biology (Clifton, N.J.)*, 1962: 227–245.

Singh, P. and Ahi, E. P. 2022. The importance of alternative splicing in adaptive evolution. *Molecular Ecology*, 31(7): 1928–1938. Publisher: John Wiley & Sons, Ltd.

Tacutu, R., Craig, T., Budovsky, A., Wuttke, D., Lehmann, G., Taranukha, D., Costa, J., Fraifeld, V. E., and de Magalhães, J. P. 2013. Human Ageing Genomic Resources: integrated databases and tools for the biology and genetics of ageing. *Nucleic Acids Research*, 41(Database issue): D1027–1033.

Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., Salzberg, S. L., Wold, B. J., and Pachter, L. 2010. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nature Biotechnology*, 28(5): 511–515. Number: 5 Publisher: Nature Publishing Group.

Tress, M. L., Abascal, F., and Valencia, A. 2017. Alternative Splicing May Not Be the Key to Proteome Complexity. *Trends in Biochemical Sciences*, 42(2): 98–110.

Verta, J.-P. and Jacobs, A. 2022. The role of alternative splicing in adaptation and evolution. *Trends in Ecology & Evolution*, 37(4): 299–308.

Waples, R. S. 2016. Life-history traits and effective population size in species with overlapping generations revisited: the importance of adult mortality. *Heredity*, 117(4): 241–250.

Waterhouse, R. M., Seppey, M., Simão, F. A., Manni, M., Ioannidis, P., Klioutchnikov, G., Kriventseva, E. V., and Zdobnov, E. M. 2018. BUSCO Applications from Quality Assessments to Gene Prediction and Phylogenomics. *Molecular Biology and Evolution*, 35(3): 543–548.

Weyna, A. and Romiguier, J. 2020. Relaxation of purifying selection suggests low effective population size in eusocial Hymenoptera and solitary pollinating bees. *bioRxiv*, page 2020.04.14.038893. Publisher: Cold Spring Harbor Laboratory Section: New Results.

White, E. P., Ernest, S. K. M., Kerkhoff, A. J., and Enquist, B. J. 2007. Relationships between body size and abundance in ecology. *Trends in Ecology & Evolution*, 22(6): 323–330.

Wilson, E. O. 2003. The encyclopedia of life. *Trends in Ecology & Evolution*, 18(2): 77–80.

Wright, C. J., Smith, C. W. J., and Jiggins, C. D. 2022. Alternative splicing as a source of phenotypic diversity. *Nature Reviews Genetics*, 23(11): 697–710. Number: 11 Publisher: Nature Publishing Group.

Wright, S. 1929. The Evolution of Dominance. *The American Naturalist*, 63(689): 556–561. Publisher: [University of Chicago Press, American Society of Naturalists].

Xu, C. and Zhang, J. 2018. Alternative polyadenylation of mammalian transcripts is generally deleterious, not adaptive. *Cell systems*, 6(6): 734–742.e4.

Xu, C. and Zhang, J. 2020. A different perspective on alternative cleavage and polyadenylation. *Nature Reviews Genetics*, 21(1): 63–63. Number: 1 Publisher: Nature Publishing Group.

Xu, C., Park, J.-K., and Zhang, J. 2019. Evidence that alternative transcriptional initiation is largely nonadaptive. *PLoS Biology*, 17(3): e3000197.

Xu, G. and Zhang, J. 2014. Human coding RNA editing is generally nonadaptive. *Proceedings of the National Academy of Sciences*, 111(10): 3769–3774. Publisher: Proceedings of the National Academy of Sciences.

Yang, Z. and Nielsen, R. 1998. Synonymous and nonsynonymous rate variation in nuclear genes of mammals. *Journal of Molecular Evolution*, 46(4): 409–418.

Zdobnov, E. M., Tegenfeldt, F., Kuznetsov, D., Waterhouse, R. M., Simão, F. A., Ioannidis, P., Seppey, M., Loetscher, A., and Kriventseva, E. V. 2017. OrthoDB v9.1: cataloging evolutionary and functional annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs. *Nucleic Acids Research*, 45(D1): D744–D749.

Zhang, J. and Xu, C. 2022. Gene product diversity: adaptive or not? *Trends in Genetics*, 38(11): 1112–1122.