Fundamental constraints on the evolution of vertebrate life histories

George C. Brooks[1]*, Josef C. Uyeda[2], Nicholas Bone[2], Hailey M. Conrad[1], Christopher G.
Mull[3], and Holly K. Kindsvater[1]

[1]Department of Fish and Wildlife Conservation, Virginia Tech, Blacksburg, Virginia, USA

[2]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, USA

[3]Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada

*corresponding author: boa10gb@vt.edu, ORCID - 0000-0001-9006-6465

Author contact information

George Brooks: boa10gb@vt.edu

Josef Uyeda: juyeda@vt.edu

Nicholas Bone: nicbone@vt.edu

Hailey Conrad: haileyconrad@vt.edu

Christopher Mull: creeas@gmail.com

Holly Kindsvater: hkindsvater@vt.edu

*keywords:-*body size, ecological opportunity, lifetime mortality, parity mode, viviparity

ABSTRACT

Vertebrate life histories evolve in response to selection imposed by abiotic and biotic

environmental conditions while being limited by genetic, developmental, physiological,

demographic, and phylogenetic processes that constrain adaptation. Despite the well-

recognized shifts in selective pressures accompanying transitions among environments,

identifying the conditions driving innovation and the consequences for life-history

evolution remain an outstanding question. Here, we compare the traits of aquatic and

terrestrial vertebrates to infer shifts in demographic and evolutionary constraints that

explain differences in life-history optimization. Specifically, our results emphasize the

reduced potential for life-history diversification on land, especially that of reproductive

strategies. Moreover, our study reveals differences between the evolution of viviparity in

the two realms. Transitions from egg laying to live birth represents a major shift across

life-history space for aquatic organisms, whereas terrestrial egg-laying organisms evolve

live birth without drastic changes in life-history strategy. Whilst trade-offs in the

allocation of resources place fundamental constraints on the way life histories can vary,

ecological setting influences the position of species within the viable phenotypic space

available for adaptive evolution.

INTRODUCTION

The evolution of diversity of vertebrate life histories has been fuelled by key transitions among water and land, and the evolution of innovative traits or reproductive strategies that allow species to survive in novel environments. The consequences of such innovations can have cascading effects on available niche space (i.e., arising from physiological constraints), life histories, and macro-evolutionary patterns. For example, internal fertilization is a necessary but insufficient innovation to explain both the colonization of terrestrial environments, and the repeated transitions to bearing live young (viviparity) [1,2,3,4]. Viviparity is intricately connected to parental care, egg size, and fertilization mode [4,5,6,7]. These characteristics, in turn, can limit evolutionary lability or allow species to capitalize on novel ecological opportunities [8,9,10,11]. The evolution of viviparity, for example, enables vertebrates to overcome some abiotic constraints, such fact that eggs without shells will desiccate on land, but will require increased investment in each offspring, shifting the fundamental trade-off between offspring size and number. These processes can affect the parameter space of life history traits that are viable, which underly demographic constraints, ultimately affecting the course of evolution. We can infer that the trait-space occupied by species persisting over evolutionary timescales emerges from these combined processes. We use this perspective to understand how transitions in reproductive mode have shaped the evolution of vertebrate diversity. While links between live-birth and adaptive radiation have been documented in fish [8], mammals [12], and reptiles [13], the consequences of the evolution of viviparity for life-history traits and demography in aquatic and terrestrial environments are not fully understood.

Life history traits related to growth, reproduction, and survival determine an individual's

maximum size, fitness, and pace-of-life and provide a fundamental link between population

demography and growth as well as evolutionary dynamics [14,15,16,17]. Life-history

diversity evolves according to the biotic stressors (e.g., predation, competition),

characteristics of the environment, and the genetic and developmental context of each

lineage [14,18,19]. The selection pressures of each environment dictate which life-history

strategies persist over evolutionary time and render certain strategies impossible (e.g.,

extremely large body mass on land, broadcast spawning on land) [20,21], while phylogenetic

inertia contributes by determining which strategies can be attained via evolution.

Transitions between aquatic and terrestrial habitats have led to radiations into unexploited

niches in both habitats. These transitions require key adaptations in development and

reproduction that foster success in the new context. Alongside the preeminent colonization

of land that marks the origin of tetrapods, transitions to an entirely terrestrial life history

have evolved independently in multiple amphibian clades, accompanied by the evolution of

direct development, in which juveniles complete metamorphosis prior to hatching on land

[7]. Reversions back to the water have occurred in most vertebrate clades [22].

The fundamental differences between species in water versus on land reflect the

consequences of the unique physical and ecological features of life in water for growth,

survival, and reproduction [23,24,25,26]. Aquatic environments prevent desiccation, buffer

thermal ranges relative to the extreme temperatures possible on land and offer structural

support against the pull of gravity [23,24,25]. Additionally, aquatic food webs differ in their

complexity and the spatial distribution of resources [27,28,29]. External fertilization is

common among aquatic species, most of which release small eggs that disperse in the water

column [23,25]. However, the repeated evolution of internal fertilization has made the

transition to viviparity possible in some ancestrally egg-laying lineages of elasmobranchs

(sharks and rays), teleosts (ray finned fishes), and amphibians [30]. A key innovation in

vertebrate diversification on land was the evolution of amniotic egg, which requires internal

fertilization, and prevents desiccation of eggs in air. Several amniotic vertebrates have

subsequently evolved partially or fully aquatic life histories but must lay and care for their

eggs on land. A few amniotic lineages, such as some sea snakes and aquatic mammals, have

escaped this constraint by evolving viviparity [30]. While the sequence of parity mode

evolution remains a topic of debate in squamate reptiles [30,31], this uncertainty mostly

concerns the choice of theoretical and statistical model, and a more comprehensive

framework with multiple lines of evidence from physiology, morphology, and

developmental biology supports the general sequence of egg-laying to live-bearing with

potential for reversals [32]. Egg-laying amniotes, such as some reptiles and all birds, can

approximate viviparity by way of producing large eggs and providing extensive parental

care, [7,33], but are prohibited from a fully aquatic existence.

Despite the well-recognized shifts in selective pressures accompanying transitions among

environments, the conditions driving developmental innovation and its consequences for

life history diversity among vertebrates have never been comprehensively tested. One

challenge is the extensive macroevolutionary scale that is necessary to identify the

constraints imposed by transitions among habitats or reproductive mode. At this scale, the

traits that taxa use to modulate their life history may not be comparable across clades (e.g.,

parity mode may be a good proxy for parental investment in many squamates, but not in

amphibians). This means that finding traits that allow quantitative comparisons of across

clades can be challenging [34,35]. Here, we compared extant vertebrates, including

cartilaginous fishes, bony fishes, amphibians, reptiles, birds, and mammals to demonstrate

that the requirements of life on land, or the ecological release experienced when colonizing

a novel environment, exert a predictable shift in demographic and evolutionary constraints,

with important consequences for the evolution of biodiversity. To accomplish this, we

define a novel trait space that enables comparisons of the complex, interrelated sets of

trade-offs underlying all life histories [35,36]. Our use of broadly available trait data

facilitates the generation of hypotheses and tests of these hypotheses with

macroevolutionary data. Specifically, we use life-history traits that serve as proxies for

mortality of juveniles and adults [37,38,39] to model the diversification of vertebrates,

including during the vertebrate invasion of land, reversions back to the water, and after the

evolution of live birth.

We identify a three-way trade-off on the bounds of the relationship between juvenile

mortality, adult mortality, and body size (Supplementary Fig. 1-3). Using this trade-off, we

propose an alternative phylogenetic framework for studying constraints on life history

evolution in vertebrates. Given the 450-million-year history of vertebrate evolution, we

assume that enough time has passed for the species in our dataset to fully explore the

bounds of life-history space, i.e., areas of trait space where species are absent represent

non-viable life-history strategies as opposed to trait space that species have yet to evolve

into [40,41]. We reformulate the bounded state space of life history as a "Ternary plot" –

equilateral triangles that define a space based on trade-offs in investment among these

interacting variables [34,42]. Focusing on traits that are fundamental to fitness and placing

each species in subdivided locations of this common space allows us to explore transitions

among locations and quantify which direction of transitions are more common over

evolutionary time. With this approach, we can go beyond previous research grouping

species by traits in ternary space to formally quantify whether and how the evolutionary

consequences of transitions from oviparity to viviparity differ among aquatic and terrestrial

environments.

RESULTS

Life-history patterns in aquatic and terrestrial environments - We used phylogenetic

comparative tests to identify constraints on the value of traits in life-history space. By

focusing on defining these constraints, as opposed to variation in the rates of change, we

uncovered fundamental shifts in life history across vertebrates due to transitions between

aquatic and terrestrial life histories. Larger vertebrates have lower mortality, as expected,

but for aquatic and terrestrial organisms this is achieved through different means (Fig. 1).

Terrestrial vertebrates have low juvenile mortality regardless of body size but exhibit a

steep negative relationship between body size and adult mortality ($\beta = -0.04, \beta = -0.58$

respectively, $p < 0.001$, Fig. 1). In contrast, species with aquatic offspring show a large

reduction in juvenile mortality with increasing body size, and low adult mortality across taxa

with a weak negative relationship with body size ($\beta = -0.40, \beta = -0.17$ respectively, $p <$

0.001, Fig. 1). When taken together, aquatic and terrestrial species exhibit highly

concordant patterns of lifetime mortality with body size ($\beta = -0.54, \beta = -0.57$

respectively, $p = 0.29$, Fig. 1). Body size and environment explain over half of the variation

($R^2_{adj} = 0.57$) in juvenile mortality of vertebrates, and over one third of the variation in

adult ($R^2_{adj} = 0.41$) and lifetime mortality ($R^2_{adj} = 0.37$).

These analyses additionally highlighted that key trade-offs among life history traits depend

on the scale of comparisons [32,43,44]. For example, we found that non-phylogenetic

estimates of the covariance between body size, adult mortality, and juvenile mortality are

all strongly negative (Supplementary Table 1). While under most scenarios phylogenetic

estimates are also negative, these estimates are considerably weaker than the non-

phylogenetic ones (except size and adult mortality, which was roughly of similar in

magnitude in the phylogenetic estimates, Supplementary Table 2). Furthermore, after

accounting for the interaction between size and terrestrial habit, we found size and juvenile

mortality to have positive evolutionary correlations in aquatic taxa, despite having strong

negative correlations in the non-phylogenetic analyses (Supplementary Table 1). Differences

between the non-phylogenetic and phylogenetic covariances suggest a contrast between

the trade-offs operating in the evolutionary dynamics between closely related species (i.e.

the covariance of changes in among diverging lineages) in life history traits with high

phylogenetic signal (Supplementary Table 3) and the overall bounds on the relationship (the

overall covariance among variables across species). This is apparent by visual inspection of

the relationships between variables (Fig. 1 and Supplementary Fig. 3-5). This difference

emerges because while evolutionary changes may appear positively or negatively correlated

in the short term, the full space of life history variation in vertebrates is bounded by

fundamental trade-offs that become visible in the non-phylogenetic regressions (and in

examination of the life history space ordinated across species).

Parity mode evolution in aquatic and terrestrial environments - When modelling life-

history evolution in ternary space, we found differences in the evolutionary consequences

of transitions to viviparity in each environment. Overall, aquatic species can occupy virtually

every corner of life-history space, whereas terrestrial species have strong selection pressure

to evolve toward reduced adult and juvenile mortality and larger body size (Fig. 2). In both

environments, while oviparity does not appear to exert strong constraints on life-history

strategies, transitions to viviparity tend to be associated with reduced adult and juvenile

mortality and larger body sizes (Fig. 3). This shift is explained by movement to a common

location within the life-history triangle, despite different starting points within oviparous

ancestors, and is generally driven by the shifts to viviparity in aquatic species including

cartilaginous fishes (sharks and rays), bony fishes, and amphibians (Fig. 3, 4, and 6). The

strength of this pattern varies among these clades; for example, transitions to live-bearing in

sharks and rays are associated with a larger increase in body size than in bony fishes [45].

For terrestrial species, viviparity is associated with a much smaller shift in life-history space

(Fig. 5, 6). In reptiles, which have the greatest number of origins of viviparity in terrestrial

vertebrates, life history traits vary only minimally between oviparous and viviparous species.

DISCUSSION

Here we present a novel framework for studying life history evolution that quantifies how

the reproductive strategies of vertebrates have been shaped by contrasting fitness

landscapes of aquatic and terrestrial environments. Specifically, transitions from egg laying

to live birth have led to major shifts in life-history trait-space for aquatic organisms. By

contrast, the life histories of oviparous and viviparous species in terrestrial environments

are largely similar. Our analysis is based on a novel trait-based approach to classifying

species based on three components of fitness (body size, age of maturity, and lifetime

fecundity) which can inform comparative studies that seek to explain life-history evolution

across a broad range of taxa. Our approach builds on previous research placing species in

trait space to identify trade-offs shaping life history strategies, most notably the

Equilibrium-Periodic-Opportunistic (E-P-O) categories [34,46]. Our method differs from prior

work in that it relies on lifetime fecundity instead of offspring size as a proxy for juvenile

mortality, and we use maturation age as a proxy of adult mortality. In our framework,

lifetime mortality (the summed mortality of juvenile and adults) shows a remarkably

consistent relationship with body size (Fig. 1), as expected, justifying our use of these proxy

traits. Elsewhere, our approach has been used to identify Periodic and Opportunistic life

histories, and to divide Equilibrium species into small-bodied species that have relatively

few, large offspring (Precocial species), and large-bodied species, which may produce low

numbers of large offspring but also live a long time (Survivor species) [16,37]. This

distinction captures an important range of life history variation in mammals, despite their

low numbers of offspring when compared to amphibians and fishes. Our focal traits are also

closely connected to a species' placement on the slow-fast life history continuum [19,47].

Our approach requires less knowledge of age-specific vital rates to place species in ternary

space. Ultimately, our three metrics simplify life-history space over macroevolutionary time

and highlight long-term constraints on the evolution of vertebrate diversity.

Overall, the selection pressures that favour live birth, as well as the benefits conferred, are

different in the water versus on land. In aquatic environments, the evolution of viviparity

requires the precursory evolution of internal fertilization, which is a prerequisite of life on

land [3,48,49,50]. The origins and advantages of different modes of fertilization are not fully

known; female choice, male opportunity to manipulate female investment, sperm storage,

increased control over fertilization timing, and reduced egg mortality have all been

suggested drivers of the evolution of internal fertilization [51,52]. The idiosyncrasies of

development and early life history of aquatic species that have evolved along with

externally fertilized eggs are likely a barrier to the evolution of viviparity for most aquatic

vertebrates. Aquatic vertebrates with external fertilization produce many tiny offspring,

which is hypothesized to maximize fitness when prey encounter is stochastic and survival is

independent of density [53,54]. By contrast, viviparity in aquatic species with internal

fertilization acts to increase juvenile survival, presumably through reduced predation of

offspring held *in utero*. Terrestrial vertebrates have internal fertilization and exhibit life-

history strategies that effectively minimize juvenile mortality regardless of whether they

have eggs or bear live young. For example, terrestrial egg deposition requires larger eggs to

prevent desiccation. Larger eggs require a longer period to develop, which in turn favours

extended parental care. Juvenile animals on land are not bathed in a nutrient broth as they

are in aquatic systems, thus precluding the evolution of particularly small offspring. Evolving

viviparity on land does not appreciably reduce juvenile mortality, but instead could act to

maintain high levels of juvenile survival across a wider range of environmental conditions.

Differences in life-history diversity and evolution in water versus on land carry important

implications for the adaptive capacity of species to environmental change [55,56]. Firstly,

terrestrial environments constrain the diversification of life-history strategies. As a result,

terrestrial vertebrates will be more limited than aquatic species in their capacity to adapt to

global change. Secondly, when considering the relative difference in vulnerability to

extinction between egg-laying and live-bearing taxa, environmental considerations are

paramount. Terrestrial live bearers have similar life-history traits to their egg-laying

counterparts, and thus will show comparable response to anthropogenic threats, all else

being equal. In contrast, the large shift in life-history strategies seen as viviparity evolves

from oviparity in aquatic species suggests their response to threats may be starkly different.

For cartilaginous fish, the evolution of live birth represents a distinct shift to a 'slower' life

history strategy, increasing generation time and reducing their adaptive capacity to rapid

environmental change. For bony fish, however, live birth appears to have evolved --when

possible-- to buffer juvenile stages from highly stochastic, ephemeral environments, and is

often accompanied by small body size and 'faster' life histories. The adaptive capacity of

aquatic livebearers with short generation times in the face of anthropogenic threats may be

greater than oviparous species.

Strong linear relationships (Fig. 1) and a triangular distribution of life-history strategies

(Supplementary Fig. 3) indicate that long term macroevolutionary patterns are highly

directional and suggest that three axes are sufficient to define the vertebrate trait space.

Phylogenetically corrected analyses showed less clearly defined trade-offs between the

three axes (Supplementary Fig. 4-6), suggesting that evolution over the short term is not

constrained by classic life-history trade-offs between growth, survival, and reproduction

[44,57], exemplified by the ubiquitous positive relationship between body size and clutch

size within clades [58]. Individual clades may have different constraints, but they are all

limited to this common space. In other words, a ternary plot is naturally bounded and

inherently accounts for the trade-offs in resource allocation that every living organism must

make [36]. Indeed, by defining the common life-history space of all vertebrates we provide a

useful way to summarize the limits of evolution within a clade. Although the overarching

constraint is the inability for species to evolve into 'Darwinian demons' which do not face

trade-offs between growth, reproduction, and survival, here we illustrate how life-history

space is filled by different clades in diverse ways. Only by understanding the

macroevolutionary trade-offs that define trait space does the challenge of comparing such a

disparate array of organisms become tractable; disparities in how species move through

space can be used to answer questions about which traits are key levers controlling

diversification.

Fundamentally, placing species in a common life-history framework helps us to understand

the rules of life. Once all species are mapped in trait space, one can model

macroevolutionary dynamics and explore covariance between key life-history traits. For

example, we show that the evolution of viviparity in aquatic and terrestrial environments

proceeds through different paths and at different rates. As an initial first step, we tested

models where transitions happened in these traits independently of their location in state

space. However, an interesting alternative model is one where each location in life history

space has its own transition rate. Future elaborations of our approach could treat the

evolution of viviparity as a cause, effect, or interaction between other factors. Additionally,

our approach could be used to understand how differences in food web structure in aquatic

and terrestrial environments shapes life-history strategies. Productivity, the patchiness of

resources, and the distribution of predators in space will have profound implications for

both a species' transitions in life-history space and the position of life-history optima.

Lastly, a potentially fruitful application of this work is to species' invasions, for which it is

possible to predict shifts in trait space following the introduction of a non-native species. If

species are typically on the edge of replacement, they will likely not be able invest in one of

the trait axes (larger body size, higher survival, increased reproductive output) without

some kind of ecological release that temporarily relaxes existing constraints. The

colonization of land and the ability of viviparous species to exploit cold climates both

exemplify the interplay between life-history evolution and ecological opportunity. These

examples demonstrate how unifying classic life history theory with trait-based approaches

and contemporary computational techniques can be leveraged to make progress on

previously intractable questions and generate novel insights into the forces that speed up or

slow evolutionary innovation.

METHODS

Trait-based framework - We collated published data on species' body size, age at maturity,

and lifetime fecundity, allowing us to place species on common life-history axes, despite

differences in their ecology. We compared distribution of body sizes, age at maturity, and

lifetime fecundity across vertebrate classes (Supplementary Fig. 1). We related this

combination of traits to live-bearing (viviparous) and egg-laying (oviparous) reproductive

strategies (Supplementary Fig. 2), and the occupancy of aquatic and terrestrial

environments (Supplementary Fig. 3). Several vertebrate groups (e.g., amphibians, sea

turtles) occupy multiple environments during their lifetimes, but for simplicity we classified

species as aquatic or terrestrial based on the initial environment a species inhabits (egg

deposition site for oviparous species, juvenile environment for viviparous species). Thus,

most amphibians are classified as aquatic despite extended terrestrial phases and sea

turtles are classified as terrestrial owing to their use of nesting beaches. The choice to focus

on the environment of early-life stages stems from our interest in the evolution of

reproductive strategies (e.g., parity mode) and the observation that fecundity exhibits the

greatest variability and disparity across aquatic and terrestrial realms (Supplementary Fig. 1-

3). Whilst a model that includes both juvenile and adult environments would be more

informative, it is computationally infeasible to include all possible life cycle transitions in the

state-space approach presented here.

Like other trait-based analyses of trade-offs and patterns in life histories [19,21,34], our trait

proxies were derived from life-history theory [14,15]. However, our approach differs in that

it uses only three traits to represent vital rates that are fundamental to both individual

fitness and population demography, allowing us to maximize the number of taxa that we

included in our analysis. Furthermore, unlike prior studies, we did not control for body size

when comparing birth and death rates, instead addressing explicitly its changes after

evolutionary transitions in habitat or reproductive mode. We relied on the principle that

persistence of populations and species over evolutionary timescales must all fulfil the same

requirement, namely that individual females, on average, must replace themselves and their

mate [18,35,36]. Lifetime production of offspring is expected to be positively correlated

with juvenile mortality and negatively correlated with per-offspring parental investment

[35,37]. Similarly, age at maturity is strongly linked to adult mortality, whereby species that

mature later have higher survival and longer expected lifespans [38,39]. Organisms with

longer lifespans therefore must, on average, replace themselves at a slower rate than those

with shorter lifespans; any other scenario would result in the unrealistic outcomes of

exponential population growth or extinction. Both adult and juvenile mortality are also

strongly related to body size, with larger animals often exhibiting larger clutch sizes and

delayed maturity [40,41]. Fundamentally, body size represents investment that must be

made in the growth and metabolic maintenance of body tissues outside of direct investment

in reproduction.

Trait and phylogenetic data – We assembled a phylogeny for vertebrates on an order-level

backbone obtained using median divergence times from the timetree of life [59]. A

supertree was made by grafting published trees for each clade. We combined trees for fish

[60], mammals [61,62], birds [63], squamate reptiles [64], amphibians [65], turtles

(testudines) [66], crocodilians [67], and chondrichthyans [68]. As the full chondrichthyan

phylogeny was constructed with a birth-death polytomy resolver, we chose to base our

analyses on the molecular backbone phylogeny calibrated with 10 fossil calibration dates

comprising 610 species from all major orders. The phylogeny includes both oviparous and

viviparous lineages and provides good coverage for the order Carcharhiniformes, which is

the largest order of sharks and exhibits the greatest diversity of reproductive modes in this

group.

Life history traits were compiled from publicly available databases that cover the major

vertebrate clades (Table 1). A link to the complete dataset for all vertebrate clades used in

this paper is provided in the data availability statement, and graphically displayed in the

supplementary material (Supplementary Fig. 7). Data for bony fish (Actinopterygii) were

downloaded from FishBase using the rfishbase package (www.fishbase.org) [69] and data

for viviparous species was supplemented with the primary literature. Data for sharks and

rays (Chondrichthyans) were taken from the Sharkipedia data repository

(www.sharkipedia.org) [70]. Amphibian trait data were taken from AmphiBio [71]. Data for

squamate reptiles were taken from [72] and supplemented with estimates of longevity

sourced from the literature. Data for birds and mammals were obtained from [73]. Manual

searches of the literature were conducted and bolstered our dataset for an additional 17

species of bony fish, 137 snakes, 137 amphibians, 45 turtles and 9 crocodilians. Owing to the

number of species included in the analysis, we had to assume that the information obtained

from previously published datasets were reliable but retrospectively checked outliers from

visual inspection of the model output to confirm their accuracy.

Body size measurements are reported as snout-vent length (SVL) for all amphibians, reptiles,

and mammals. Body sizes reported for bony fish, sharks, skates, and chimaeras represent

total length, and measurements for rays are disc (wing) width. When multiple

measurements were available for a species, mean values were used for analysis. If sex-

specific estimates were available, we used numbers reported for females. Most trait

databases did not include mass, so to convert lengths to mass we used available female

length-weight (L-W) or width-weight conversion relationships (see supporting data for

sources of L-W relationships). When sex and species-specific L-W relationships were

unavailable, we estimated reasonable L-W relationships from the same genus or family.

Reproductive mode was characterized as either egg-laying or live-bearing, and fertilization

mode was coded as internal or external. Species were classified as either aquatic or

terrestrial; given our interest in reproductive modes, species with biphasic life cycles were

coded based on the location of egg deposition (e.g., most amphibians are coded as aquatic

despite having terrestrial adults). Given our emphasis on the evolution of reproductive

strategies, we only considered juvenile environments in the present analysis.

Proxies for mortality can be inferred from the life-history traits that are most

sensitive to changes in mortality schedules [14]. For example, increased predation or

harvest typically selects for earlier maturity [74,75]. Therefore, we use the inverse of age at

maturity to estimate adult mortality, such that:

$$M_a = 1/(1 + A_{mat})$$

381   where A_{mat} is the age at maturity and M_a is adult mortality [37]. In a similar fashion,

382   lifetime fecundity is indicative of mortality experienced in early life-stages [14,15].

383   Calculating lifetime fecundity requires information on lifespan, age at maturity, breeding

384   frequency (e.g. annual, biennial) and clutch size. We logged this quantity such that lifetime

385   egg production is equal to:

$$M_j = ln((A_{max} - A_{mat}) \times (F * b))$$

where A_{max} is longevity, A_{mat} is maturity, F is fecundity, and b represents the number of

breeding bouts per year. The distribution of all three life-history traits, split between aquatic

and terrestrial species, with egg-laying and live-bearing reproductive modes is presented in

Supplementary Fig. 1 and summarized in Supplementary Table 4. To produce a rough

measure of lifetime mortality for each species, we scaled and summed our proxies of

juvenile mortality and adult mortality.

$$M_{total} = scaled(M_j) + scaled(M_a)$$

where our proxies of juvenile and adult mortality are first scaled before they are summed.

Statistical Analyses:-We estimated the covariances among traits using both phylogenetic

and non-phylogenetic estimates using multivariate Brownian Motion models of trait

evolution. The assumption of multivariate Brownian Motion is likely inappropriate, and

therefore we also used an alternative phylogenetic modelling approach (Supplementary

Table 2). Nevertheless, estimates of phylogenetic and non-phylogenetic correlation matrices

among traits are a meaningful summary statistic of the covariances among traits across the

observable evolutionary process. Therefore, we used a Pagel's λ model and the R package

phylopars [76] to accommodate for partially missing data, with analyses run multiple times

with species missing 0, 1, or 2 trait values excluded. Analyses were performed once with a

fully phylogenetic model (λ = 1) and again in a non-phylogenetic model (λ = 0) to facilitate

comparisons of the phylogenetic and non-phylogenetic estimates with missing data. For

each trait individually, we additionally report estimated phylogenetic signal metrics of

Pagel's λ and Blomberg's K for all taxa with complete data using the function *phylosig* in the

R package *phytools* [77].

To determine the relationship between maximum body size and mortality estimates, we fit

three separate linear regression models with juvenile mortality, adult mortality, and lifetime

mortality as response variables. In each model body size, habitat (aquatic or terrestrial), and

their interaction were included as predictors. As we were interested in the overall bounds

on life-history space, rather than clade-specific relationships, our primary analyses did not

correct for phylogeny – the uncorrected formulation is likely more informative for our

research question as it emphasizes the bounds of the trait space rather than the covariances

of changes among lineages [32,43,44]. We then performed the same regressions accounting

for phylogeny using the function *phylopars.lm* [76]. We assumed phylogenetically correlated

residuals evolving under an Ornstein-Uhlenbeck process (Supplementary Fig. 5). For each

model we obtained average effect sizes, and their associated standard errors and p-values.

Model fit was evaluated by comparing predicted and observed mortality values and

estimating R^2 using the *rr2* package [78].

Ternary plots are often used for "phases" or regions of interactions that have a distinct

shared feature (e.g. 'solid', 'liquid' or 'gas' equilibria in a Gibbs triangle). These features

make them useful for projecting relationships between life history traits (e.g., Mims et al.

2010). We created ternary plots for our proxies for body size, adult mortality, and juvenile

mortality. Allocation to each of these trait axes for each species was taken as their relative

location across the full span of vertebrate life history trait space, with each of these ranges

standardized to a new axis on a 0-100 scale $[100 \times (x - x_{min})/(x_{max} - x_{min})]$. We then

discretized this space into 16 equal triangles (4 bins for each trait) representing the possible

states that each species in our dataset could occupy. This defined life-history space enables

us to employ a discrete-state evolutionary model. By doubling the state space and allowing

transitions between "layers" to be defined by binary characters, we can compare how such

binary factors influence life history evolution across vertebrates. For example, we defined a

32-state continuous-time Markov chain with two ternary plot spaces corresponding to

aquatic and terrestrial life history (Supplementary Fig. 8).

We then estimated transitions between aquatic and terrestrial environments, and between

regions of the life history space assuming that movement within this space is only allowed

between discrete bins that neighbor each other with a shared edge. Transitions in the binary

character move species to corresponding bins in the replicated ternary space for the

alternative binary state (Supplementary Fig. 8). Since a 32-by-32 transition matrix would

have an exceptionally large number of parameters under most models, we simplified

movement within each ternary space by defining four parameters governing the dynamics

of trait change. These parameters are defined by the directional trends toward each of the

three vertices of space, as well as an overall parameter governing whether species are

attracted or repelled to the center of the triangle. Each layer of the life history space is

defined by four unique parameters plus two parameters for asymmetric transitions in the

binary trait, for a total of 10 parameters.

We conducted a second set of analyses that accommodate missing data in our three life

history proxies by coding their location within the state space with ambiguous coding, which

can be modeled using continuous-time Markov Chains. When modeled in this way, the

posterior probabilities of species with missing data are computed jointly and informed by

both their observed life history values, as well as the life history values of related species

(see supplementary material). All analyses were performed in R using the *fit_mk* function in

the castor package modified to account for ternary space [79](R Core Team 2023).

Data availability

The dataset used in this study can be accessed at https://doi.org/10.5281/zenodo.11457349 [80]

Code availability

The code generated in this study can be accessed at https://doi.org/10.5281/zenodo.11457349 [80]

Acknowledgements

We would like to thank Jacob Berman and Ty Stephenson for help compiling trait data. We would like to thank the Department of Fish and Wildlife Conservation and the Department of Biological Sciences at Virginia Tech for financial and logistical support.

Author contributions

GCB and HKK conceived the idea. GCB, HMC, and CM compiled and proofed the data. JU designed the analyses. GCB, JU, and NB performed the analyses and confirmed the reproducibility of the code and results. GCB, JU, and HKK wrote the manuscript. All authors revised and approved the final manuscript.

Competing Interests Statement

The authors declare no competing interests.

Additional Information

Supplementary Information is available for this paper.

REFERENCES

1. Blackburn, D.G., 1992. Convergent evolution of viviparity, matrotrophy, and specializations for fetal nutrition in reptiles and other vertebrates. American Zoologist, 32(2), pp.313-321.
2. Wourms, J. P., & Lombardi, J. (1992). Reflections on the evolution of piscine viviparity. American Zoologist, 32(2), 276-293.
3. Kalinka, A.T., 2015. How did viviparity originate and evolve? Of conflict, co-option, and cryptic choice. BioEssays, 37(7), pp.721-731.
4. Vági, B., Marsh, D., Katona, G., Végvári, Z., Freckleton, R.P., Liker, A. and Székely, T., 2022. The evolution of parental care in salamanders. Scientific Reports, 12(1), p.16655.
5. Gross, M. R., and R. C. Sargent. 1985. The Evolution of Male and Female Parental Care in Fishes. American Zoologist, 25:807–22.
6. Farmer CG. 2000. Parental Care: The key to understanding endothermy and other convergent features in birds and mammals. Am Nat. 155(3):326-334. doi: 10.1086/303323.
7. Vági, B., Végvári, Z., Liker, A., Freckleton, R.P. and Székely, T., 2019. Parental care and the evolution of terrestriality in frogs. Proceedings of the Royal Society B, 286(1900), p.20182737.

8. Helmstetter, A.J., Papadopulos, A.S., Igea, J., Van Dooren, T.J., Leroi, A.M. and
Savolainen, V., 2016. Viviparity stimulates diversification in an order of fish. Nature
communications, 7(1), p.11271.
9. Reznick, D.N., Travis, J., Pollux, B.J. and Furness, A.I., 2021. Reproductive mode and
conflict shape the evolution of male attributes and rate of speciation in the fish
family Poeciliidae. Frontiers in Ecology and Evolution, 9, p.639751.
10. Miller, A.H., Stroud, J.T. and Losos, J.B., 2023. The ecology and evolution of key
innovations. Trends in Ecology & Evolution, 38(2), pp.122-131.
11. Nuño de la Rosa, L., 2023. Agency in Reproduction. Evolution & Development.
25:418–429.
12. Blackburn, D.G., 1999. Viviparity and oviparity: evolution and reproductive
strategies. Encyclopedia of reproduction, 4, pp.994-1003.
13. Recknagel, H., Kamenos, N.A. and Elmer, K.R., 2021. Evolutionary origins of viviparity
consistent with palaeoclimate and lineage diversification. Journal of evolutionary
biology, 34(7), pp.1167-1176.
14. Stearns, Stephen C., 1992. The Evolution of Life Histories. Oxford University Press,
London.
15. Roff, D. ed., 1993. Evolution of life histories: theory and analysis. Springer Science &
Business Media.
16. Kindsvater, H.K., Mangel, M., Reynolds, J.D. and Dulvy, N.K., 2016. Ten principles
from evolutionary ecology essential for effective marine conservation. Ecology and
Evolution, 6(7), pp.2125-2138.
17. White, C.R., Alton, L.A., Bywater, C.L., Lombardi, E.J. and Marshall, D.J., 2022.
Metabolic scaling is the product of life-history optimization. Science, 377(6608),
pp.834-839.
18. Partridge, L. and Harvey, P.H., 1988. The ecological context of life history evolution.
Science, 241(4872), pp.1449-1455.
19. Healy, K., Ezard, T.H.G., Jones, O.R., Salguero-Gómez, R., Buckley, Y.M., 2019., Animal
Life History is Shaped by the Pace of Life and the Distribution of Age-Specific
Mortality and Reproduction. Nature Ecology & Evolution, 3(8), 1217-1224.
20. Southwood, T.R.E., 1988. Tactics, strategies and templets. Oikos, pp.3-18.
21. Brown, J.H., Hall, C.A. and Sibly, R.M., 2018. Equal fitness paradigm explained by a
trade-off between generation time and energy production rate. Nature Ecology &
Evolution, 2(2), pp.262-268.
22. Colbert, E.H., 1955. Evolution of the vertebrates. A history of the backboned animals
through time. Evolution of the vertebrates. A history of the backboned animals
through time.
23. Strathmann, R. R. (1990). Why life histories evolve differently in the sea. American
Zoologist, 30(1), 197–207.
24. Dawson, M. N., & Hamner, W. M. (2008). A biophysical perspective on dispersal and
the geography of evolution in marine and terrestrial systems. Journal of the Royal
Society, Interface/the Royal Society, 5(19), 135–150.
25. Burgess, S.C., Baskett, M.L., Grosberg, R.K., Morgan, S.G. and Strathmann, R.R., 2016.
When is dispersal for dispersal? Unifying marine and terrestrial perspectives.
Biological Reviews, 91(3), pp.867-882.
26. Capdevila, P., Beger, M., Blomberg, S.P., Hereu, B., Linares, C. and Salguero-Gómez,
R., 2020. Longevity, body dimension and reproductive mode drive differences in

aquatic versus terrestrial life-history strategies. Functional Ecology, 34(8), pp.1613-
1625.

27. Chase, J.M., 2000. Are there real differences among aquatic and terrestrial food
        webs? Trends in ecology & Evolution, 15(10), pp.408-412.

28. Weston, N.H., Vanni, M.J., Yang, L.H. 2008. Comparing Resource Pulses in Aquatic
        and Terrestrial Systems. Ecology, 89(3), 647-659.

29. Steele, J.H., Brink, K.H. and Scott, B.E., 2019. Comparison of marine and terrestrial
        ecosystems: suggestions of an evolutionary perspective influenced by environmental
        variation. ICES Journal of Marine Science, 76(1), pp.50-59.

30. Blackburn, D.G., 2015. Evolution of vertebrate viviparity and specializations for fetal
        nutrition: a quantitative and qualitative analysis. Journal of Morphology, 276(8),
        pp.961-990.

31. Pyron, R.A. and Burbrink, F.T., 2014. Early origin of viviparity and multiple reversions
        to oviparity in squamate reptiles. Ecology letters, 17(1), pp.13-21.

32. Uyeda, J.C., Zenil-Ferguson, R. and Pennell, M.W., 2018. Rethinking phylogenetic
        comparative methods. Systematic Biology, 67(6), pp.1091-1109.

33. Martin, K. L., and Carter, A. L. 2013. Brave new propagules: terrestrial embryos in
        anamniotic eggs. Integrative and Comparative Biology 53:233–247.

34. Winemiller, K.O. and Rose, K.A., 1992. Patterns of life-history diversification in North
        American fishes: implications for population regulation. Canadian Journal of
        Fisheries and aquatic sciences, 49(10), pp.2196-2218.

35. Gadgil, M. and Bossert, W.H., 1970. Life historical consequences of natural selection.
        The American Naturalist, 104(935), pp.1-24.

36. Leimar, O., 2002. Evolutionary change and Darwinian demons. Selection, 2(1-2),
        pp.65-72.

37. Kindsvater, H.K., Dulvy, N.K., Horswill, C., Juan-Jordá, M.J., Mangel, M. and
        Matthiopoulos, J., 2018. Overcoming the data crisis in biodiversity conservation.
        Trends in Ecology & Evolution, 33(9), pp.676-688.

38. Charnov, E.L., 1990. On evolution of age of maturity and the adult lifespan. Journal of
        Evolutionary Biology, 3(1-2), pp.139-144.

39. Charnov, E.L. and Berrigan, D., 1990. Dimensionless numbers and life history
        evolution: age of maturity versus the adult lifespan. Evolutionary Ecology, 4(3),
        pp.273-275.

40. Darwin, C. R. 1874. The descent of man, and selection in relation to sex. 2d ed.
        Appleton, New York.

41. Williams, G. C. 1966. Adaptation and natural selection. Princeton University Press,
        Princeton, N.J.

42. Grime, J.P., 1974. Vegetation classification by reference to strategies. Nature,
        250(5461), pp.26-31.

43. Simpson, E.H., 1951. The interpretation of interaction in contingency tables. Journal
        of the Royal Statistical Society: Series B (Methodological), 13(2), pp.238-241.

44. Reznick, D., Nunney, L. and Tessier, A., 2000. Big houses, big cars, superfleas and the
        costs of reproduction. Trends in ecology & evolution, 15(10), pp.421-425.

45. Goodwin, N.B., Dulvy, N.K. and Reynolds, J.D., 2002. Life-history correlates of the
        evolution of live bearing in fishes. Philosophical Transactions of the Royal Society of
        London. Series B: Biological Sciences, 357(1419), pp.259-267.

46. Winemiller, K.O., 1989. Patterns of variation in life history among South American
fishes in seasonal environments. Oecologia, 81, pp.225-241.

47. Salguero-Gómez, R., Jones, O.R., Jongejans, E., Blomberg, S.P., Hodgson, D.J., Mbeau-
Ache, C., Zuidema, P.A., De Kroon, H. and Buckley, Y.M., 2016. Fast–slow continuum
and reproductive strategies structure plant life-history variation worldwide.
Proceedings of the National Academy of Sciences, 113(1), pp.230-235.

48. Gross, M.R. and Shine, R., 1981. Parental care and mode of fertilization in
ectothermic vertebrates. Evolution, pp.775-793.

49. Beck, C.W., 1998. Mode of fertilization and parental care in anurans. Animal
Behaviour, 55(2), pp.439-449.

50. Benun Sutton, F. and Wilson, A.B., 2019. Where are all the moms? External
fertilization predicts the rise of male parental care in bony fishes. Evolution, 73(12),
pp.2451-2460.

51. Orr, T.J. and Brennan, P.L., 2015. Sperm storage: distinguishing selective processes
and evaluating criteria. Trends in Ecology & Evolution, 30(5), pp.261-272.

52. Lyons, K., Kacev, D. and Mull, C.G., 2021. An inconvenient tooth: Evaluating female
choice in multiple paternity using an evolutionarily and ecologically important
vertebrate clade. Molecular Ecology, 30(7), pp.1574-1593.

53. Winemiller, K.O. and Rose, K.A., 1993. Why do most fish produce so many tiny
offspring?. The American Naturalist, 142(4), pp.585-603.

54. Olsson, K.H., Gislason, H. and Andersen, K.H., 2016. Differences in density-
dependence drive dual offspring size strategies in fish. Journal of Theoretical Biology,
407, pp.118-127.

55. Forster, J., Hirst, A.G. and Atkinson, D., 2012. Warming-induced reductions in body
size are greater in aquatic than terrestrial species. Proceedings of the National
Academy of Sciences, 109(47), pp.19310-19314.

56. Kuparinen, A. and Festa-Bianchet, M., 2017. Harvest-induced evolution: insights
from aquatic and terrestrial systems. Philosophical Transactions of the Royal Society
B: Biological Sciences, 372(1712), p.20160036.

57. Chang, C.C., Moiron, M., Sánchez-Tójar, A., Niemelä, P.T. and Laskowski, K.L., 2023.
What is the meta-analytic evidence for life-history trade-offs at the genetic level?
Ecology Letters.

58. Shine, R., 1988a. The evolution of large body size in females: a critique of Darwin's"
fecundity advantage" model. The American Naturalist, 131(1), pp.124-131.

59. Hedges, S.B., Dudley, J. and Kumar, S., 2006. TimeTree: a public knowledge-base of
divergence times among organisms. Bioinformatics, 22(23), pp.2971-2972.

60. Rabosky, D.L., Chang, J., Cowman, P.F., Sallan, L., Friedman, M., Kaschner, K., Garilao,
C., Near, T.J., Coll, M. and Alfaro, M.E., 2018. An inverse latitudinal gradient in
speciation rate for marine fishes. Nature, 559(7714), pp.392-395.

61. Zhou, C.F., Wu, S., Martin, T. and Luo, Z.X., 2013. A Jurassic mammaliaform and the
earliest mammalian evolutionary adaptations. Nature, 500(7461), pp.163-167.

62. Upham, N.S., Esselstyn, J.A. and Jetz, W., 2019. Inferring the mammal tree: species-
level sets of phylogenies for questions in ecology, evolution, and conservation. PLoS
biology, 17(12), p.e3000494.

63. Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K. and Mooers, A.O., 2012. The global
diversity of birds in space and time. Nature, 491(7424), pp.444-448.

64. Tonini, J.F.R., Beard, K.H., Ferreira, R.B., Jetz, W. and Pyron, R.A., 2016. Fully-sampled phylogenies of squamates reveal evolutionary patterns in threat status. Biological Conservation, 204, pp.23-31.

65. Jetz, W. and Pyron, R.A., 2018. The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. Nature ecology & evolution, 2(5), pp.850-858.

66. Thomson, S.A., 2021. Turtles of the world: annotated checklist and atlas of taxonomy, synonymy, distribution, and conservation status. Phyllomedusa: Journal of Herpetology, 20(2), pp.225-228.

67. Hedges S.B., Marin J., Suleski M., Paymer M., Kumar S. 2015. "Tree of life reveals clock-like speciation and diversification." Mol. Biol. Evol. 32:835–845.

68. Stein, R.W., Mull, C.G., Kuhn, T.S., Aschliman, N.C., Davidson, L.N., Joy, J.B., Smith, G.J., Dulvy, N.K. and Mooers, A.O., 2018. Global priorities for conserving the evolutionary history of sharks, rays and chimaeras. Nature ecology & evolution, 2(2), pp.288-298.

69. Froese, R., 1992. 1992 progress report on FishBase. ICLARM Contribution No. 852.

70. Mull, C. G., N. Pacoureau, S. A. Pardo, L. S. Ruiz, E. García-Rodríguez, B. Finucci, M. Haack et al. "Sharkipedia: a curated open access database of shark and ray life history traits and abundance time-series." Scientific Data 9, no. 1 (2022): 559.

71. Oliveira, B.F., São-Pedro, V.A., Santos-Barrera, G., Penone, C. and Costa, G.C., 2017. AmphiBIO, a global database for amphibian ecological traits. Scientific data, 4(1), pp.1-7.

72. Meiri, S., 2018. Traits of lizards of the world: Variation around a successful evolutionary design. Global ecology and biogeography, 27(10), pp.1168-1172.

73. Myhrvold, N.P., Baldridge, E., Chan, B., Sivam, D., Freeman, D.L. and Ernest, S.M., 2015. An amniote life-history database to perform comparative analyses with birds, mammals, and reptiles: Ecological Archives E096-269. Ecology, 96(11), pp.3109-3109.

74. Abrams, P.A. and Rowe, L., 1996. The effects of predation on the age and size of maturity of prey. Evolution, 50(3), pp.1052-1061.

75. Sharpe, D.M. and Hendry, A.P., 2009. Life history change in commercially exploited fish stocks: an analysis of trends across studies. Evolutionary applications, 2(3), pp.260-275.

76. Goolsby EW, Ane C, Bruggeman J. 2017. Rphylopars: Fast Multivariate Phylogenetic Comparative Methods for Missing Data and Within-Species Variation. Methods in Ecology & Evolution. 2017. 8:22-27.

77. Revell, L.J. 2024. phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and other things). PeerJ, 12, p.e16505.

78. Ives, A.R. and Li, D., 2018. rr2: An R package to calculate R2s for regression models. Journal of Open Source Software, 3(30), p.1028.

79. R Core Team. 2024. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

80. Brooks, G. C., Uyeda J. C., H. Conrad, N. Bone, C. Mull, and H. K. Kindsvater. 2024. Code for: Fundamental constraints on the evolution of vertebrate life histories v1.0. https://doi.org/10.5281/zenodo.11457349

81. Smith (2017). Ternary: An R Package for Creating Ternary Plots. Comprehensive R
Archive Network. doi: 10.5281/zenodo.1068996

Table 1. Number of species for each of the major vertebrate clades for which partial
and complete life-history data was available. Numbers in parentheses show the
proportion of total diversity represented within each clade. Also shown is the
proportion of species (with complete data) that are viviparous or have terrestrial egg
deposition.

Clade	Partial Data	Complete Data	Viviparity	Terrestrial Eggs
Chondrichthyes	571 (46%)	115 (19%)	81%	0%
Actinopterygii	9441 (29%)	117 (0.3%)	15%	0%
Amphibia	5250 (61%)	260 (4%)	3%	6%
Reptilia	4516 (39%)	213 (3%)	14%	100%
Aves	8690 (80%)	922 (8%)	0%	100%
Mammalia	4480 (70%)	1465 (30%)	99%	97%

Figure 1: The relationship between body size and a) juvenile mortality, b) adult mortality, and c) lifetime mortality for aquatic (n = 577) and terrestrial (n = 2345) vertebrates.
Aquatic and terrestrial species differ in their relationship between juvenile mortality and body size ($GLM, p < 0.001$), and in their relationship between adult mortality and body size ($GLM, p < 0.001$). When taken together however, aquatic and terrestrial species show no difference in the scaling of lifetime mortality with body size ($GLM, p = 0.29$). Lines represent predicted trends, and shaded areas reflect 95% confidence intervals.

Aquatic   **Terrestrial**

701   **Figure 2: Movement through life-history space for aquatic and terrestrial vertebrates.**
702   Plots were generated from 2922 species for which complete life-history data was available.
703   The colours represent the number of taxa represented in each segment, with darker colours
704   representing more species, and lighter colours representing fewer species. The arrows
705   indicate the direction and relative strength of selection through life-history space.

Oviparity   **Viviparity**

Figure 3: Movement through life-history space for egg-laying and live-bearing vertebrates.
Plots were generated from 2922 species for which complete life-history data was available.
The colours represent the number of taxa represented in each segment, with darker colours
representing more species, and lighter colours representing fewer species. The arrows
indicate the direction and relative strength of selection through life-history space.

Figure 4. Phylogeny of cartilaginous fishes (sharks, rays, and chimaeras) and the relative position in life-history space for egg-laying (top right) and live-bearing (bottom right) taxa. Ternary plots show the counts of species (n = 561) in each segment with overlaid contours indicating the density of species in life-history space, with species with complete data indicated as points. Densities were estimated using the two-dimensional kernel density estimation procedure from the *Ternary* package [81]. The branches of the phylogeny are coloured using stochastic mapping to match the cells in the ternary plots, with lighter and darker colors indicating oviparous and viviparous species, respectively. Silhouettes were obtained from PhyloPic (https://www.phylopic.org/) under a creative commons license.

Figure 5. Phylogeny of reptiles and the relative position in life-history space for egg-laying (top right) and live-bearing (bottom right) taxa. Ternary plots show the counts of species (n = 3031) in each segment with overlaid contours indicating the density of species in life-history space, with species with complete data indicated as points. Densities were estimated using the two-dimensional kernel density estimation procedure from the *Ternary* package [81]. The branches of the phylogeny are coloured using stochastic mapping to match the cells of the ternary plots, with lighter and darker colors indicating oviparous and viviparous species, respectively. Silhouettes were obtained from PhyloPic (https://www.phylopic.org/) under a creative commons license.

**Figure 6. Comparison of egg-laying and live-bearing species of bony fish (n = 3799),
amphibians (n = 4520), mammals (n = 4480), and birds (n = 8690).** Ternary plots show the
counts of species in each segment with overlaid contours indicating the density of species in
life-history space, with species with complete data indicated as points. Densities were
estimated using the two-dimensional kernel density estimation procedure from the *Ternary*
package [81]. The branches of the phylogeny are coloured using stochastic mapping to
match the cells in the ternary plots, with lighter and darker colors indicating oviparous and
viviparous species, respectively. Silhouettes were obtained from PhyloPic
(https://www.phylopic.org/) under a creative commons license.