Fundamental constraints on the evolution of vertebrate life histories

George C. Brooks[1]*, Josef C. Uyeda[2]*, Nicholas Bone[2], Hailey M. Conrad[1], Christopher G.

Mull[3], and Holly K. Kindsvater[1]

[1]Department of Fish and Wildlife Conservation, Virginia Tech, Blacksburg, Virginia, USA

[2]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, USA

[3]Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada

*authors contributed equally to this work.

corresponding author: George Brooks, boa10gb@vt.edu

Author contact information

George Brooks: boa10gb@vt.edu

Josef Uyeda: juyeda@vt.edu

Nicholas Bone: nicbone@vt.edu

Hailey Conrad: haileyconrad@vt.edu

Christopher Mull: creeas@gmail.com

Holly Kindsvater: hkindsvater@vt.edu

keywords:-body size, ecological opportunity, lifetime mortality, parity mode, viviparity

ABSTRACT

The evolution of vertebrate life histories has been fuelled by key transitions among water
and land. Life history diversity evolves according to the interplay of predation on juveniles
and adults, the physical properties of the environment, and the developmental context of
each lineage. The selection pressures of each environment dictate which life-history
strategies persist over evolutionary time and render certain strategies impossible. Despite
the well-recognized shifts in selective pressures accompanying transitions among
environments, the conditions driving innovation and its consequences for life-history
evolution have never been comprehensively tested. Here, we show that the contrasting
fitness landscapes operating in aquatic and terrestrial environments result in predictable
shifts in demographic and evolutionary constraints that explain differences in vertebrate
life histories. Specifically, our results emphasize the reduced potential for life-history
diversification on land, especially that of reproductive strategies. Moreover, our study
reveals differences between the evolution of viviparity in the two environments.
Transitions from egg laying to live birth represents a major shift across life-history space
for aquatic organisms, whereas terrestrial egg-laying organisms evolve live birth without
drastic changes in life-history strategy. Whilst trade-offs in the allocation of resources
place fundamental constraints on the way life histories can vary, ecological setting defines
the macroevolutionary space that species can explore.

INTRODUCTION

The evolution of diversity of vertebrate life histories has been fuelled by key transitions among water and land, as well as a myriad of life history traits that act to modulate investment in development and reproduction. For example, the evolution of internal fertilization is a necessary but insufficient innovation to explain repeated transitions to bearing live young (viviparity) in aquatic and terrestrial environments (Blackburn 1992, Wourms and Lombardi 1992, Kalinka 2015, Vági et al. 2022). The consequences of innovations in reproduction can have cascading effects on available niche space, life histories, and macro-evolutionary patterns. Viviparity is intricately connected to parental care, egg size, and fertilization mode (Gross and Sargent 1985, Farmer 2000, Vági et al. 2019, 2022). These characteristics, in turn, can limit evolutionary lability or allow species to capitalize on novel ecological opportunities (Helmstetter et al. 2016, Reznick et al. 2021, Miller et al. 2023, Nuño de la Rosa 2023). Transitions in reproductive mode are key to the evolution of vertebrate diversity, but the consequences for life-history traits and demography could differ in aquatic and terrestrial environments.

Life history traits related to growth, reproduction, and survival determine an individual's maximum size, fitness, and pace-of-life (Stearns 1992; Roff 1992; Kindsvater et al. 2016, White et al. 2022) and provide a fundamental link between population demography and growth as well as evolutionary dynamics. Life history diversity evolves according to the interplay of predation on juveniles and adults, the physical properties of the environment, and the developmental context of each lineage (Partridge and Harvey 1988, Stearns 1992, Healy et al. 2019). The selection pressures of each environment dictate which life-history strategies persist over evolutionary time and render certain strategies impossible

(Southwood 1988), while phylogenetic inertia contributes by determining which strategies

can be attained via evolution. Transitions between aquatic and terrestrial habitats have led

to radiations into unexploited niches in both habitats. These transitions require key

adaptations in development and reproduction that foster success in the new context.

Alongside the preeminent colonization of land that marks the origin of tetrapods, transitions

to an entirely terrestrial life history have evolved independently in multiple amphibian

clades, accompanied by the evolution of direct development, in which juveniles complete

metamorphosis prior to hatching on land. Reversions back to the water have occurred in

most vertebrate clades (Colbert 1955).

The fundamental differences between species in water versus on land reflect the

consequences of the unique physical and ecological features of life in water for growth,

reproduction, and dispersal (Strathmann 1990, Dawson and Hamner 2008, Burgess et al.

2016, Capdevila et al. 2020). Aquatic environments prevent desiccation, buffer thermal

ranges relative to the extreme temperature possible on land and offer structural support

against the pull of gravity (Strathmann 1990, Dawson and Hamner 2008, Burgess et al.

2016). Additionally, aquatic food webs are more complex, and resources tend to be less

patchy in space and time (though exceptions are common; Chase 2000, Weston et al. 2008,

Steele et al. 2019). External fertilization is common among aquatic species, most of which

release small eggs that disperse in the water column (Strathmann 1990, Burgess et al. 2016).

However, the repeated evolution of internal fertilization has made the transition to

viviparity possible in some ancestrally egg-laying lineages of elasmobranchs (sharks and

rays), teleosts (ray finned fishes), and amphibians. A key innovation in vertebrate

diversification on land was the evolution of amniotic egg, which requires internal

fertilization, and prevents desiccation of eggs in air. Several amniotic vertebrates have

subsequently evolved partially or fully aquatic life histories but must lay and care for their

eggs on land. A few amniotic lineages, such as some sea snakes and aquatic mammals, have

escaped this constraint by evolving viviparity. Other amniotic clades including some reptiles

and all birds instead produce large eggs that are protected via an eggshell, extensive

parental care, or both (Martin and Carter 2013, Vági et al. 2019).

Despite the well-recognized shifts in selective pressures accompanying transitions among

environments, the conditions driving developmental innovation and its consequences for

life history diversity and evolutionary processes have never been comprehensively tested.

One challenge is that the phylogenetic scales that hold the necessary variation for

comprehensively identifying the constraints to life history variation also carry considerable

variation in the traits that taxa use to modulate their life history (e.g. parity mode may be a

good proxy for parental investment in many squamates, but not in amphibians). This means

that finding traits that allow quantitative comparisons of across clades is challenging. Here,

we used extant vertebrates, including cartilaginous fishes, bony fishes, amphibians, reptiles,

birds, and mammals to demonstrate that the constraints imposed by life on land, or indeed

the evolutionary release experienced when entering the water, exert a predictable shift in

demographic and evolutionary constraints, with important consequences for the evolution

of biodiversity. To accomplish this study, we define a novel trait space that enables

comparisons of the complex, interrelated sets of trade-offs underlying all life histories with

broadly available trait data, facilitating the generation of hypotheses and guiding tests of

these hypotheses with macroevolutionary data. Specifically, we used proxies for life history

traits where precise information is lacking to understand life history evolution throughout

the diversification of vertebrates. This approach improves our understanding of the

vertebrate invasion of land, reversions back to the water, and the evolution of live birth.

METHODS

Trait and phylogenetic data:-A phylogeny for vertebrates was assembled on a order-level

backbone obtained using median divergence times from the timetree of life (Hedges et al.

2006). A supertree was made by grafting published trees for each clade (Figure S2). We

combined trees for fish (Rabosky et al. 2018), mammals (Zhou et al. 2013, Upham et al.

2019), birds (Jetz et al. 2012), squamates (Tonini et al. 2016), amphibians (Jetz & Pyron

2018), testudines (Thomson et al. 2021), crocodylians (Hedges et al. 2015), and

chondrichthyans (sharks, skates, rays, and chimaeras; hereafter sharks, Stein et al. 2018). As

the full shark phylogeny was constructed with a birth-death polytomy resolver, we chose to

base our analyses on the molecular backbone phylogeny calibrated with 10 fossil calibration

dates comprising 610 species from all major orders. The phylogeny includes both oviparous

and viviparous lineages and provides good coverage for the order Carcharhiniformes which

exhibits the greatest diversity of reproductive modes.

Life history traits were compiled from publicly available databases that cover the major

vertebrate clades. The complete dataset for all vertebrate clades used in this paper is

provided as supplementary material, and summarized in Table 1 (Table 1, Appendix 1). Data

for bony fish (Actinopterygii) were downloaded from FishBase using the rfishbase package

(www.fishbase.org; Froese 1992). Data for sharks and rays (Chondrichthyans) were taken

from the Sharkipedia data repository (www.sharkipedia.org; Mull et al. 2022). Amphibian

trait data were taken from AmphiBio (Oliveira et al. 2017). Data for squamate reptiles were

taken from Meiri et al. (2018) and supplemented with estimates of longevity sourced from

the literature. Data for birds and mammals were obtained from Myhrvold et al. (2015). We

performed manual searches of the literature to bolster our dataset. We were able to source

data for an additional 17 species of bony fish, 137 snakes, 137 amphibians, 45 turtles

(Testudines) and 9 crocodilians. Owing to the number of species included in the analysis we

had to assume that the information obtained from previously published datasets were

reliable but performed post-hoc checks of influential data points as to their accuracy.

Table 1. Number of species for each of the major vertebrate clades for which life-history data was available. Numbers in parentheses are percentages of the total number of species included in each group.

Clade	Total	Internal Fertilization	Viviparity	Terrestrial Eggs	Terrestrial Adults
Chondrichthyes	115 (19%)	100%	81%	0%	0%
Actinopterygii	117 (0.05%)	17%	15%	0%	0%
Amphibia	260 (4%)	33%	3%	6%	95%
Reptilia	213 (3%)	100%	14%	100%	64%
Aves	356 (4%)	100%	0%	100%	100%
Mammalia	1465 (30%)	100%	99%	97%	94%

Body size measurements are snout-vent length (SVL) for all amphibians, reptiles, and

mammals. Body sizes reported for bony fish, sharks, skates, and chimaeras represent total

length, and measurements for rays reflect disc width. When multiple measurements were

available for a species, mean values were used for analysis. If sex-specific estimates were

available, we used numbers reported for females. Most of the trait databases did not

include mass, so to convert lengths to mass we used available female length-weight (L-W) or

width-weight conversion relationships. When sex and species-specific L-W relationships

were unavailable we estimated reasonable L-W relationships from the same genus or

family. Reproductive mode was characterized as either egg-laying or live-bearing, and

fertilization mode was coded as internal or external. Species were classified as either

154 aquatic or terrestrial, and to account for biphasic life-cycles juveniles and adults were

155 classified separately (e.g., most amphibians are coded as aquatic juveniles and terrestrial

156 adults). In total, we compiled a data set including… (Figure S3, Table 1).

157 Our proxy for adult mortality only requires an estimate of age at maturity, such that:

158
$$M_a \;=\; ln(1/A_{mat})$$

159 where A_{mat} is the age at maturity and M_a is adult mortality (Kindsvater et al. 2018).

160 Calculating lifetime fecundity requires information on lifespan, age at first reproduction,

161 breeding frequency (e.g. annual, biennial) and clutch size, such that lifetime egg production

162 is equal to:

163
$$M_j \;=\; ln((A_{max} \;-\; A_{mat}) \;*\; (F \;*\; b))$$

164 where A_{max} is longevity, A_{mat} is maturity, F is fecundity, and b represents the number of

165 breeding bouts per year. The distribution of all three life-history traits, split between aquatic

166 and terrestrial species, with egg-laying and live-bearing reproductive modes is presented in

167 Figure S1. To produce a crude measure of lifetime mortality for each species, we scaled and

168 summed our proxies of juvenile mortality and adult mortality.

169
$$M_{total} \;=\; scaled(M_j) \;+\; scaled(M_a)$$

where our proxies of juvenile and adult mortality are first scaled before they are summed.

Collating data on body size, age at maturity, and lifetime reproductive output allowed us to

place species on common life-history axes, despite differences in ecology and traits

underlying these values. We compared the distribution of body sizes, age at first

reproduction, and lifetime fecundity across species, and related them to occupancy of

aquatic and terrestrial environments in different life stages (Figure S1). We also considered

whether species have internal or external fertilization and whether they are live-bearing

(viviparous) or egg-laying (oviparous). Our proxies were derived based on life-history theory

and the principle that persistence of populations and species over evolutionary timescales

must all fulfil the same requirement, namely that individual females, on average, must

replace themselves and their mate (Gadgil and Bossert 1970, Partridge and Harvey 1988,

Leimar 2002). Expected lifetime production of offspring is an indirect measure of juvenile

mortality and per offspring parental investment (Gadgil and Bossert 1970; Kindsvater et al.

2018). Similarly, age at first reproduction is strongly linked to adult mortality, whereby

species that mature later have higher survival and longer expected lifespans (Charnov 1990,

Charnov and Berrigan 1990). Organisms with longer lifespans therefore must, on average,

replace themselves at a slower rate than those with shorter lifespans; any other scenario

would result in the unrealistic outcomes of exponential growth or extinction. However, both

adult and juvenile mortality are also strongly related to body size, with larger animals often

exhibiting larger clutch sizes and delayed maturity (Darwin 1874, Williams 1966).

Fundamentally, body size represents investment that must be made in the growth and

metabolic maintenance of body tissues outside of direct investment in reproduction.

However, the consequences of transitions from water to land on species' survival and

growth are poorly understood.

Statistical Analyses:-We estimated the covariances among traits using both phylogenetic

and non-phylogenetic estimates using multivariate Brownian Motion models of trait

evolution. The assumption of multivariate Brownian Motion is likely inappropriate, and

therefore we used an alternative phylogenetic modeling approach (see below).

Nevertheless, estimates of phylogenetic and non-phylogenetic correlation matrices among

traits is a meaningful summary statistic of the covariances among traits across the

observable evolutionary process. Therefore, we used a Pagel's λ model and the R package

phylopars (Goolsby et al. 2017) to accommodate for partially missing data, with analyses run

multiple times with species missing 0, 1, or 2 trait values excluded. Analyses were

performed once with a fully phylogenetic model (λ = 1) and again in a non-phylogenetic

model (λ = 0) to facilitate comparisons of the phylogenetic and non-phylogenetic estimates

with missing data.

To determine the relationship between maximum body size and mortality estimates, we fit

three separate linear regression models with juvenile mortality, adult mortality, and lifetime

mortality as response variables. In each model body size, habitat (aquatic or terrestrial), and

their interaction were included as predictors. As we were interested in the overall bounds

on life-history space, rather than clade-specific relationships, our primary analyses did not

correct for phylogeny – the uncorrected formulation is likely more informative for our

research question as it emphasizes the bounds of the trait space rather than the covariances

of changes among lineages (Simpson 1951, Reznick et al. 2000, Uyeda et al. 2018). We then

performed the same regressions corrected for phylogeny using the function *phylopars.lm*

(Goolsby et al. 2017). We assumed phylogenetically correlated residuals evolving under an

Ornstein-Uhlenbeck process. To quantify taxonomic heterogeneity in the relationship

between body size and mortality, we performed the comparative analyses separately for

every major clade (Figure S4). For each model we obtained average effect sizes, and their

associated standard errors and p-values. Model fit was evaluated by comparing predicted

and observed mortality values and estimating R^2 using the *rr2* package (Ives and Li 2018).

Ternary plots are often used for "phases" or regions of interactions that have a distinct

shared feature (e.g. 'solid', 'liquid' or 'gas' equilibria in a Gibbs triangle). These features

make them useful for projecting relationships between life history traits (e.g., Mims et al.

2010). We created ternary plots for our proxies for body size, adult mortality, and juvenile

mortality. Allocation to each of these trait axes for each species was taken as their relative

location across the full span of vertebrate life history trait space, with each of these ranges

standardized to a new axis on a 0-100 scale. We then discretized this space into 16 equal

triangles (4 bins for each trait) representing the possible states that each species in our

dataset could occupy. This defined life-history space enables us to employ a discrete-state

evolutionary model. By doubling the state space and allowing transitions between "layers"

defined by binary characters, we can compare how such binary factors influence life history

evolution across vertebrates. For example, we defined a 32-state continuous-time Markov

chain with two ternary plot spaces corresponding to aquatic and terrestrial life history

(Figure 1).

We then estimated transitions between aquatic and terrestrial environments, and between

regions of the life history space assuming that movement within this space is only allowed

between discrete bins that neighbor each other with a shared edge. Transitions in the binary

character move species to corresponding bins in the replicated ternary space for the

alternative binary state (Figure 1). Since a 32 by 32 transition matrix would have an

exceptionally large number of parameters under most models, we simplified movement

within each ternary space by defining four parameters governing the dynamics of trait

change. These parameters are defined by the directional trends toward each of the three

vertices of the space, as well as an overall parameter governing whether species are

attracted or repelled to the center of the triangle (see Supplement). Each layer of the life

history space is defined by four unique parameters plus two parameters for asymmetric

transitions in the binary trait, for a total of 10 parameters.

We conducted a second set of analyses that accommodate missing data in our three life

history proxies by coding their location within the state space with ambiguous coding, which

can be modeled using continuous-time Markov Chains. When modeled in this way, the

posterior probabilities of species with missing data are computed jointly and is informed by

both their observed life history values, as well as the life history values of related species

(see supplementary material). All analyses were performed in R using the *fit_mk* function in

the castor package modified to account for ternary space (R Core Team 2023).

RESULTS

We identified a three-way trade-off on the bounds of the relationship between offspring

mortality, adult mortality, and body size (Figure S1), suggesting an alternative phylogenetic

framework for studying constraints on life history evolution in vertebrates. Assuming that

over the course of vertebrate evolution, the bounds on vertebrate life history extremes

have largely been reached by the species in our dataset, we reformulated the bounded state

space of life history as a "Ternary plot" – equilateral triangles that define a space based on

trade-offs in investment among these interacting variables (Grime 1974). Placing each

species in subdivided locations of this common space then allowed us to explore transitions

among locations and quantify which direction of transitions are more common over

evolutionary time. With this approach, we address whether the evolutionary significance of

transitions from oviparity to viviparity differ among aquatic and terrestrial environments.

Life-history patterns in aquatic and terrestrial environments:-We used phylogenetic

comparative tests to identify constraints on the value of traits in life-history space. By

focusing on defining these constraints and the "arena" for life history evolution, as opposed

to variation in the rates of change, we uncovered fundamental shifts in life history across

vertebrates due to transitions between aquatic and terrestrial life histories. Larger

vertebrates have lower mortality, as expected, but for aquatic and terrestrial organisms this

is achieved through different means (Figure 2). Terrestrial vertebrates have low juvenile

mortality regardless of body size but exhibit a steep negative relationship between body size

and adult mortality ($\beta = -0.04$, $\beta = -0.58$ respectively, $p < 0.001$, Figure 2A, 2B). In

contrast, species with aquatic offspring show a large reduction in juvenile mortality with

increasing body size, and low adult mortality across taxa with a weak negative relationship

with body size ($\beta = -0.40$, $\beta = -0.17$ respectively, $p < 0.001$, Figure 2A, 2B). When taken

together however, aquatic and terrestrial species exhibit highly concordant patterns of

lifetime mortality with body size ($\beta = -0.54$, $\beta = -0.57$ respectively, $p = 0.29$, Figure

2C). Body size and environment explain over a half of the variation ($R^2_{adj} = 0.57$) in juvenile

mortality of vertebrates, and over a third of the variation in adult ($R^2_{adj} = 0.41$) and

lifetime mortality ($R^2_{adj} = 0.37$).

These analyses additionally highlighted key trade-offs among life history traits depend on

the scale of comparisons (Simpson 1951, Reznick et al. 2000, Uyeda et al. 2018). For

example, while non-phylogenetic estimates of the covariance between body size, adult

mortality, and juvenile mortality are all strongly negative, phylogenetic estimates vary

between negative and positive for most traits (except size and adult mortality, which were

negatively related in both types of analyses). We found size and both juvenile mortality or

adult mortality have a positive evolutionary correlation, but strong negative correlations in

the non-phylogenetic analyses. This disparity in the direction of trade-offs suggests a

contrast between the evolutionary dynamics between closely related species and the

overall bounds on the relationship, which is likewise apparent by visual inspection of the

relationships between variables (Figure 2, S4). This difference emerges because while

evolutionary changes may appear positively or negatively correlated in the short term, the

full space of life history variables in vertebrates is bounded by the fundamental trade-offs

that become visible in the non-phylogenetic regressions (and overall examining of the life

history space ordinated across species).

Parity mode evolution in aquatic and terrestrial environments:-When modelling life-history

evolution in ternary space, we found fundamental differences in the evolutionary

consequences of transitions to viviparity in each environment. Overall, aquatic species can

occupy virtually any space in life-history space, whereas terrestrial species have strong

selection pressure to evolve toward reduced adult and juvenile mortality and larger body

size (Figure 3). In both environments, while oviparity does not appear to exert strong

constraints on life-history strategies, transitions to viviparity are associated with reduced

adult and juvenile mortality and larger body size (Figure 4). This shift is explained by

movement to a common location within the life-history triangle, despite different starting

points within oviparous ancestors. However, for aquatic species, transitions from oviparity

to viviparity are typically accompanied by reductions in adult and juvenile mortality, and

larger body sizes (Figure 5, 7). By contrast, for terrestrial species, viviparity is associated with

a much smaller shift in life-history space (Figure 6, 7). In sum, we infer that while aquatic

environments permit the evolution of more diverse life histories, the evolution of viviparity

constrains life history traits more for aquatic species than terrestrial environments.

DISCUSSION

Here we present a novel framework for studying life history evolution that quantifies how

developmental and reproductive strategies of aquatic and terrestrial vertebrates have been

shaped by contrasting fitness landscapes throughout their radiation. Specifically, transitions

from egg laying to live birth represents a major shift across life-history space for aquatic

organisms, whereas terrestrial egg-laying organisms are situated adjacent to viviparous

strategists and evolve live birth in response to environmental pressures more as a means to

remain in that region of life-history space. Our analysis is based on a unique trait-based

approach to classifying species based on three metrics, or components of fitness, which can

inform comparative studies that seek to explain life history evolution across a broad range

of taxa. Ultimately, our three metrics simplify life history space over macroevolutionary time

and highlight the long-term constraints on vertebrate diversity.

Overall, the selection pressures that favour live birth, as well as the benefits conferred, are

different in the water versus on land. In aquatic environments the evolution of viviparity

necessitates the precursory evolution of internal fertilization (Gross and Shine 1981, Beck

1998, Kalinka 2015, Benun Sutton and Wilson 2019). The origins and advantages of internal

fertilization are unclear; female choice, male opportunity to manipulate female investment,

sperm storage, increased control over fertilization timing, and reduced egg mortality have

all been suggested as potential benefits (Orr and Brennan 2015, Lyons et al. 2021). Viviparity

in aquatic species acts to increase juvenile survival. Terrestrial organisms, in contrast, have

already evolved the necessary preadaptation for live birth; internal fertilization is a

prerequisite of life on land. Moreover, terrestrial vertebrates have already evolved life-

history strategies that effectively minimize juvenile mortality. Terrestrial egg deposition

requires larger eggs to prevent desiccation, which prolongs developmental times, which in

turn favours extended parental care. Animals on land are not bathed in a nutrient broth as

they are in aquatic systems, thus precluding particularly small offspring. As such, viviparity

on land does not appreciably reduce juvenile mortality, but instead acts more to maintain

high levels of juvenile survival across a wider range of environmental conditions.

Differences in life-history diversity and evolution in water versus on land carry important

implications for the adaptive capacity of species to environmental change (Forster et al.

2012, Kuparinen and Festa-Bianchet 2017). Firstly, terrestrial environments permit a greatly

reduced diversity of life-history strategies. As a result, terrestrial vertebrates will be more

vulnerable to global change. Secondly, when considering the relative difference in

vulnerability between egg-laying and live-bearing taxa, environmental considerations are

paramount. Terrestrial live bearers have similar life-history traits to their egg-laying

counterparts, and thus will show comparable response to anthropogenic threats, all else

being equal. In contrast, the large shift in life-history strategies seen as viviparity evolves

from oviparity in aquatic species suggests their response to threats may be starkly different.

For cartilaginous fish, the evolution of live birth represents a distinct shift to a 'slower' life

history strategy, reducing their adaptive capacity to rapid environmental change (though

this may be mediated by an increased capacity for dispersal). For bony fish, however, live

birth appears to have evolved to keep pace with highly stochastic, ephemeral environments.

As such, their adaptive capacity in the face of anthropogenic threats may be greater than

oviparous species.

Strong linear relationships (Figure 2) and a triangular distribution of strategies (Figure S1)

indicate that long term macroevolutionary patterns are highly directional and that three

axes are sufficient to define the vertebrate trait space. Phylogenetically corrected analyses

showed less clearly defined trade-offs between the three axes (Figure S4), suggesting that

evolution over the short term is not constrained by classic life-history trade-offs between

growth, survival, and reproduction (Reznick et al. 2000, Chang et al. 2023). The ubiquitous

positive relationship between body size and clutch size within clades serves to highlight this

notion (Shine 1988a). Individual clades may have different constraints, but they are all

limited to a common space. Indeed, by defining the common life-history space of all

vertebrates we provide a useful way to summarize the limits of evolution within a clade.

Although the overarching constraint is the inability for species to evolve into 'Darwinian

demons' which do not face trade-offs between growth, reproduction, and survival, here we

show that life-history space is filled by different clades in different ways and at different

rates. Only by understanding the macroevolutionary trade-offs that define trait space does

the challenge of comparing such a diaspora of organisms become tractable; disparities in

how species move through space can be used to answer questions about which traits are

key levers controlling diversification.

Fundamentally, placing species in a common life-history framework helps us to understand

the rules of life. Once all species are mapped in trait space, one can model

macroevolutionary dynamics and explore covariance between key life-history traits. For

example, we show that the evolution of viviparity in aquatic and terrestrial environments

precedes through different paths and at different rates. Our framework could also be used

to understand how differences in food web structure in aquatic and terrestrial

environments shapes life-history strategies. Productivity, the patchiness of resources, and

the distribution of predators will have profound implications for both a species' movement

through life-history space and the position of life-history optima. Lastly, a potentially

fruitful application of this work are species invasions – in our framework, it will be possible

to predict shifts in trait-space following the introduction of a non-native species. If species

are typically on the edge of replacement, they will likely not be able invest in one of the trait

axes (larger body size, higher survival, increased reproductive output) without some kind of

ecological release that temporarily relaxes constraints in trait space. For example, the

colonization of land, and the ability of viviparous species to exploit cold climates exemplify

the interplay between life-history evolution and ecological opportunity. Further research

unifying classic life history theory with trait-based approaches and contemporary

computational techniques can be leveraged to make progress on previously intractable

questions and generate novel insights on the forces that speed up or slow evolutionary

innovation.

Data and code availability statement

The dataset and code generated during the current study are available in the Github

repository, https://github.com/geobro1992/VTparity

Acknowledgements

We would like to thank Jacob Berman and Ty Stevenson for help compiling trait data. We

would like to thank the Department of Fish and Wildlife Conservation and the Department

of Biological Sciences at Virginia Tech for financial and logistical support.

Author contributions

GB and HK conceived the idea. GB, HC, and CM compiled and proofed the data. JU designed

the analyses. GB, JU, and NB performed the analyses and confirmed the reproducibility of

the code and results. GB and HK wrote the manuscript. All authors revised and approved the

final manuscript.

Competing Interests Statement

The authors declare no competing interests.

REFERENCES

Beck, C.W., 1998. Mode of fertilization and parental care in anurans. Animal Behaviour,

55(2), pp.439-449.

Benoit-Bird, K., 2023. Resource patchiness as a resolution to the food paradox in the sea.

The American Naturalist.

Benun Sutton, F. and Wilson, A.B., 2019. Where are all the moms? External fertilization

predicts the rise of male parental care in bony fishes. Evolution, 73(12), pp.2451-2460.

Blackburn, D.G., 1992. Convergent evolution of viviparity, matrotrophy, and specializations

for fetal nutrition in reptiles and other vertebrates. American Zoologist, 32(2), pp.313-321.

Burgess, S.C., Baskett, M.L., Grosberg, R.K., Morgan, S.G. and Strathmann, R.R., 2016. When

is dispersal for dispersal? Unifying marine and terrestrial perspectives. Biological Reviews,

91(3), pp.867-882.

Capdevila, P., Beger, M., Blomberg, S.P., Hereu, B., Linares, C. and Salguero-Gómez, R., 2020.

Longevity, body dimension and reproductive mode drive differences in aquatic versus

terrestrial life-history strategies. Functional Ecology, 34(8), pp.1613-1625.

Chang, C.C., Moiron, M., Sánchez-Tójar, A., Niemelä, P.T. and Laskowski, K.L., 2023. What is

the meta-analytic evidence for life-history trade-offs at the genetic level? Ecology Letters.

Charnov, E.L., 1990. On evolution of age of maturity and the adult lifespan. Journal of

Evolutionary Biology, 3(1-2), pp.139-144.

Charnov, E.L. and Berrigan, D., 1990. Dimensionless numbers and life history evolution: age

of maturity versus the adult lifespan. Evolutionary Ecology, 4(3), pp.273-275.

Chase, J.M., 2000. Are there real differences among aquatic and terrestrial food webs?

Trends in ecology & Evolution, 15(10), pp.408-412.

Clavel J, Escarguel G, Merceron G (2015). "mvMORPH: an R package for fitting multivariate

evolutionary models to morphometric data." Methods in Ecology and Evolution, 6, 1311-

1319.

Colbert, E.H., 1955. Evolution of the vertebrates. A history of the backboned animals

through time. Evolution of the vertebrates. A history of the backboned animals through

time.

Darwin, C. R. 1874. The descent of man, and selection in relation to sex. 2d ed. Appleton,

New York.

Dawson, M. N., & Hamner, W. M. (2008). A biophysical perspective on dispersal and the

geography of evolution in marine and terrestrial systems. Journal of the Royal Society,

Interface/the Royal Society, 5(19), 135–150.

Duarte, C.M. and Alcaraz, M., 1989. To produce many small or few large eggs: a size-

independent reproductive tactic of fish. Oecologia, 80, pp.401-404.

Farmer CG. 2000. Parental Care: The key to understanding endothermy and other

convergent features in birds and mammals. Am Nat. 155(3):326-334. doi: 10.1086/303323.

Forster, J., Hirst, A.G. and Atkinson, D., 2012. Warming-induced reductions in body size are

greater in aquatic than terrestrial species. Proceedings of the National Academy of Sciences,

109(47), pp.19310-19314.

Froese, R., 1992. 1992 progress report on FishBase. ICLARM Contribution No. 852.

Gadgil, M. and Bossert, W.H., 1970. Life historical consequences of natural selection. The

American Naturalist, 104(935), pp.1-24.

Goolsby EW, Ane C, Bruggeman J. 2017. Rphylopars: Fast Multivariate Phylogenetic

Comparative Methods for Missing Data and Within-Species Variation. Methods in Ecology &

Evolution. 2017. 8:22-27.

Grime, J.P., 1974. Vegetation classification by reference to strategies. Nature, 250(5461),

pp.26-31.

Gross, M.R. and Shine, R., 1981. Parental care and mode of fertilization in ectothermic

vertebrates. Evolution, pp.775-793.

Gross, Mart R., and R. Craig Sargent. 1985. The Evolution of Male and Female Parental Care

in Fishes. American Zoologist, 25:807–22.

Healy, K., Ezard, T.H.G., Jones, O.R., Salguero-Gómez, R., Buckley, Y.M., 2019., Animal Life

History is Shaped by the Pace of Life and the Distribution of Age-Specific Mortality and

Reproduction. Nature Ecology & Evolution, 3(8), 1217-1224.

Hedges, S.B., Dudley, J. and Kumar, S., 2006. TimeTree: a public knowledge-base of

divergence times among organisms. Bioinformatics, 22(23), pp.2971-2972.

Hedges S.B., Marin J., Suleski M., Paymer M., Kumar S. 2015. "Tree of life reveals clock-like

speciation and diversification." Mol. Biol. Evol. 32:835–845.

Helmstetter, A.J., Papadopulos, A.S., Igea, J., Van Dooren, T.J., Leroi, A.M. and Savolainen,

V., 2016. Viviparity stimulates diversification in an order of fish. Nature communications,

7(1), p.11271.

Ho, Lam si Tung, and Cécile Ané. 2014. "A Linear-Time Algorithm for Gaussian and Non-

Gaussian Trait Evolution Models." *Systematic Biology* 63 (3): 397–408.

Ho, L.S.T., Ane, C., Lachlan, R., Tarpinian, K., Feldman, R., Yu, Q., van der Bijl, W., Maspons,

J., Vos, R. and Ho, M.L.S.T., 2016. Package 'phylolm'.

Ives, A.R. and Li, D., 2018. rr2: An R package to calculate R2s for regression models. J. Open

Source Softw., 3(30), p.1028.

Kalinka, A.T., 2015. How did viviparity originate and evolve? Of conflict, co-option, and

cryptic choice. BioEssays, 37(7), pp.721-731.

Kindsvater, H.K., Mull, C.G., Pardo, S.A., Pacoureau, N. and Dulvy, N.K., 2020. Sharkipedia:

Elasmobranch Traits and Trends Global Database.

Kuparinen, A. and Festa-Bianchet, M., 2017. Harvest-induced evolution: insights from

aquatic and terrestrial systems. Philosophical Transactions of the Royal Society B: Biological

Sciences, 372(1712), p.20160036.

Leimar, O., 2002. Evolutionary change and Darwinian demons. Selection, 2(1-2), pp.65-72.

Lyons, K., Kacev, D. and Mull, C.G., 2021. An inconvenient tooth: Evaluating female choice in

multiple paternity using an evolutionarily and ecologically important vertebrate clade.

Molecular Ecology, 30(7), pp.1574-1593.

Martin, K. L., and Carter, A. L. 2013. Brave new propagules: terrestrial embryos in

anamniotic eggs. Integrative and Comparative Biology 53:233–247.

Miller, A.H., Stroud, J.T. and Losos, J.B., 2023. The ecology and evolution of key innovations.

Trends in Ecology & Evolution, 38(2), pp.122-131.

Mims, M.C., Olden, J.D., Shattuck, Z.R. and Poff, N.L., 2010. Life history trait diversity of

native freshwater fishes in North America. Ecology of Freshwater Fish, 19(3), pp.390-400.

Mull, C. G., N. Pacoureau, S. A. Pardo, L. S. Ruiz, E. García-Rodríguez, B. Finucci, M. Haack et

al. "Sharkipedia: a curated open access database of shark and ray life history traits and

abundance time-series." Scientific Data 9, no. 1 (2022): 559.

Myhrvold, N.P., Baldridge, E., Chan, B., Sivam, D., Freeman, D.L. and Ernest, S.M., 2015. An

amniote life-history database to perform comparative analyses with birds, mammals, and

reptiles: Ecological Archives E096-269. Ecology, 96(11), pp.3109-3109.

Nuño de la Rosa, L., 2023. Agency in Reproduction. Evolution & Development. 25:418–429.

Oliveira, B.F., São-Pedro, V.A., Santos-Barrera, G., Penone, C. and Costa, G.C., 2017.

AmphiBIO, a global database for amphibian ecological traits. Scientific data, 4(1), pp.1-7.

Orr, T.J. and Brennan, P.L., 2015. Sperm storage: distinguishing selective processes and

evaluating criteria. Trends in Ecology & Evolution, 30(5), pp.261-272.

Partridge, L. and Harvey, P.H., 1988. The ecological context of life history evolution. Science,

241(4872), pp.1449-1455.

Reznick, D., Nunney, L. and Tessier, A., 2000. Big houses, big cars, superfleas and the costs of

reproduction. Trends in ecology & evolution, 15(10), pp.421-425.

Reznick, D.N., Travis, J., Pollux, B.J. and Furness, A.I., 2021. Reproductive mode and conflict

shape the evolution of male attributes and rate of speciation in the fish family Poeciliidae.

Frontiers in Ecology and Evolution, 9, p.639751.

Sargent, R.C., Taylor, P.D. and Gross, M.R., 1987. Parental care and the evolution of egg size

in fishes. The American Naturalist, 129(1), pp.32-46.

Shine, R., 1988a. The evolution of large body size in females: a critique of Darwin's"

fecundity advantage" model. The American Naturalist, 131(1), pp.124-131.

Shine, R., 1988b. Constraints on reproductive investment: a comparison between aquatic

and terrestrial snakes. Evolution, 42(1), pp.17-27.

Simpson, E.H., 1951. The interpretation of interaction in contingency tables. Journal of the

Royal Statistical Society: Series B (Methodological), 13(2), pp.238-241.

Stearns, Stephen C., 1992. The Evolution of Life Histories. Oxford University Press, London.

Steele, J.H., Brink, K.H. and Scott, B.E., 2019. Comparison of marine and terrestrial

ecosystems: suggestions of an evolutionary perspective influenced by environmental

variation. ICES Journal of Marine Science, 76(1), pp.50-59.

Stein, R.W., Mull, C.G., Kuhn, T.S., Aschliman, N.C., Davidson, L.N., Joy, J.B., Smith, G.J.,

Dulvy, N.K. and Mooers, A.O., 2018. Global priorities for conserving the evolutionary history

of sharks, rays and chimaeras. Nature ecology & evolution, 2(2), pp.288-298.

Strathmann, R. R. (1990). Why life histories evolve differently in the sea. American Zoologist,

30(1), 197–207.

Vági, B., Végvári, Z., Liker, A., Freckleton, R.P. and Székely, T., 2019. Parental care and the

evolution of terrestriality in frogs. Proceedings of the Royal Society B, 286(1900),

p.20182737.

Vági, B., Marsh, D., Katona, G., Végvári, Z., Freckleton, R.P., Liker, A. and Székely, T., 2022.

The evolution of parental care in salamanders. Scientific Reports, 12(1), p.16655.

Weston, N.H., Vanni, M.J., Yang, L.H. 2008. Comparing Resource Pulses in Aquatic and

Terrestrial Systems. Ecology, 89(3), 647-659.

Williams, G. C. 1966. Adaptation and natural selection. Princeton University Press,

Princeton, N.J.

Wourms, J.P., 1981. Viviparity: the maternal-fetal relationship in fishes. American Zoologist,

21(2), pp.473-515.

Wourms, J. P., & Lombardi, J. (1992). Reflections on the evolution of piscine viviparity.

American Zoologist, 32(2), 276-293.

Figure 1. A conceptual diagram of the discrete-space evolutionary model in ternary space.

In this modelling framework, species can evolve to neighbouring patches within the

defined life-history space or can transition between states. The two state spaces

(triangles) can represent egg-laying and live-bearing reproductive modes or aquatic and

terrestrial environments.

Figure 2: The relationship between body size and a) juvenile mortality, b) adult mortality, and c) lifetime mortality for aquatic and terrestrial vertebrates. Offspring mortality in terrestrial environments is hyper-constrained, and variation in life-histories is chiefly explained by changes in adult mortality. Regression lines account for phylogenetic relatedness.

Figure 3: Movement through life-history space for aquatic (left) and terrestrial (right)

vertebrates. Plots were generated from 2922 species for which life-history data was

available. The colours represent the number of taxa represented in each segment, with

darker colours representing fewer species, and lighter colours representing more species.

The arrows indicate the direction and relative strength of selection through life-history

space.

Figure 4: Movement through life-history space for egg-laying (left) and live-bearing (right) vertebrates. Plots were generated from 2922 species for which life-history data was available. The colours represent the number of taxa represented in each segment, with darker colours representing fewer species, and lighter colours representing more species. The arrows indicate the direction and relative strength of selection through life-history space.

Figure 5. Phylogeny (left) of cartilaginous fishes (sharks, rays, and chimaeras) and the

relative position in life-history space for egg-laying (top right) and live-bearing (bottom

right) taxa. Ternary plots show the counts of species in each segment with overlaid

contours indicating the density of species in life-history space. The branches of the

phylogeny are coloured to match the ternary plots.

Figure 6. Phylogeny (left) of reptiles and the relative position in life-history space for egg-laying (top right) and live-bearing (bottom right) taxa. Ternary plots show the counts of species in each segment with overlaid contours indicating the density of species in life-history space. The branches of the phylogeny are coloured to match the ternary plots.

Figure 7. Comparison of egg-laying (left) and live-bearing (right) species of bony fish (top row), amphibians (second row), mammals (third row), and birds (bottom row). Ternary plots show the counts of species in each segment with overlaid contours indicating the density of species in life-history space.