Embryonic origins of forebrain oligodendrocytes revisited by combinatorial

genetic fate mapping

Yuqi Cai[1#], Zhirong Zhao[1#], Mingyue Shi[1], Mingfang Zheng[1], Ling Gong[1], Miao He[1*]

[1] Institutes of Brain Science, State Key Laboratory of Medical Neurobiology and MOE Frontiers
Center for Brain Science, Department of Neurobiology, Zhongshan Hospital, Fudan University,
Shanghai 200032, China.

[#]These authors contributed equally to this work.

[*]Corresponding author hem@fudan.edu.cn

Abstract

Multiple embryonic origins give rise to forebrain oligodendrocytes (OLs), yet controversies

and uncertainty exist regarding their differential contributions. We established intersectional

and subtractional strategies to genetically fate map OLs produced by medial ganglionic

eminence/preoptic area (MGE/POA), lateral/caudal ganglionic eminences (LGE/CGE) and

dorsal pallium. We found that, contrary to the canonical view, LGE/CGE-derived OLs make

minimum contributions to the neocortex and corpus callosum, but dominate piriform cortex

and anterior commissure. Additionally, MGE/POA-derived OLs, instead of being entirely

eliminated, make small but sustained contribution to cortex with a distribution pattern

distinctive from those derived from the dorsal origin. Our study provides a revised and more

comprehensive view of cortical and white matter OL origins, and established valuable new

tools and strategies for future OL studies.

Introduction

Oligodendrocytes (OLs) are an important class of macroglia responsible for producing the myelin sheaths that insulate and protect neuronal axons. Forebrain OLs arise from multiple embryonic origins. Previous fate mapping study using $Nkx2.1^{Cre}$[1], $Gsh2^{Cre}$[2] and $Emx1^{Cre}$[3] reported consecutive and competing waves of OLs derived from medial ganglionic eminence/preoptic area (MGE/POA), lateral/caudal ganglionic eminences (LGE/CGE) and dorsal pallium[2]. The first-wave of OLs generated by MGE/POA ($_{MP}$OLs) were believed to be eliminated postnatally, while those from the second and third waves ($_{LC}$OLs and dOLs) survive and populate the cortex and corpus callosum at comparable proportions. Several other studies provided both supporting and contradicting evidence to this model[4-9]. Moreover, $Gsh2$ was recently found to be expressed in dorsal progenitors[10], casting doubt on the interpretation of lineage tracing data from $Gsh2^{Cre}$.

In this study, we generated new genetic tools and combinatorial fate mapping strategies which allow direct visualization and comparison among OLs derived from different origins. We found that neocortical OLs are primarily composed of dOLs, rather than similar proportions of $_{LC}$OLs and dOLs. In contrast, $_{LC}$OLs and dOLs made comparable contributions to piriform cortex. We also found that although $_{MP}$OLs only make a small contribution, they do persist in the cortex beyond adulthood with a unique spatial pattern distinct from that of the dOLs. In the two major white matter commissure tracts, dOLs are the vast majority in corpus callosum but make little contribution to anterior commissure, while $_{LC}$OLs behaved the opposite. These findings significantly revised the classical view and provided a new and more comprehensive picture of cortical and white matter OL origins.

Results and Discussion

To unambiguously track OLs from different embryonic origins, we first generated a knock-in

driver, $Opalin^{P2A-Flpo-T2A-tTA2}$ (Figure 1), orthogonal to Cre drivers that label dorsal or ventral

progenitors ($Progenitor^{Cre}$). *Opalin* (also known as *Tmem10*) encodes oligodendrocytic

myelin paranodal and inner loop protein that are specifically expressed in differentiated

oligodendrocytes[11-14]. In $Opalin^{P2A-Flpo-T2A-tTA2}$, *Flpo* and *tTA2* were inserted before the

STOP codon and linked by self-cleavage peptide P2A and T2A (Figure 1A-C), allowing

co-transcription and translation with *Opalin*. Flp-mediated recombination by this driver

(hereinafter referred to as $Opalin^{Flp}$ for simplicity) enables highly specific, efficient and

irreversible OL labeling, while the tTA2 component offers the flexibility for labeling in

tunable densities (Figure 1D-F).

Next, we established two types of genetic combinatorial fate mapping strategies to directly

visualize OLs from different embryonic origins (Figure 2): (1) combining $Opalin^{Flp}$ and

$Progenitor^{Cre}$ with intersectional reporters *Ai65* to label OLs derived from Cre+ progenitor

domain by RFP (Figure 2A); (2) combining $Opalin^{Flp}$ and $Progenitor^{Cre}$ with *RC::FLTG*[15]

to simultaneously label OLs derived from Cre+ progenitors by GFP and OLs derived from the

complementing Cre- progenitors by RFP (Figure 2B). The first approach allowed us to track

dOLs and $_{MP}$OLs (Figure 2C, E). The second approach empowered us to observe and compare

OLs generated from dorsal and ventral origins, or those from Gsh2+ and Gsh2- progenitors,

in the same brain (Supplementary Figure 1). Importantly, the subtraction power enabled us to

target OLs derived from LGE/CGE progenitors that express neither *Emx1* nor *Nkx2.1* (Figure

2D). In addition, these strategies greatly facilitated the identification of OLs derived from

specific origin which exist at relatively low density in certain regions.

Deploying these strategies, we assessed the differential contributions of dOLs, $_{LC}$OLs and

$_{MP}$OLs by analyzing RFP+ cells in the following mice: *Opalin^{Flp}::Emx1^{Cre}::Ai65* (Figure 2C),

Opalin^{Flp}::Emx1^{Cre}::Nkx2.1^{Cre}::RC::FLTG (Figure 2D) and *Opalin^{Flp}::Nkx2.1^{Cre}::Ai65* (Figure

2E). We observed two significant differences from the traditional model in the neocortex. The

first major deviation is that, instead of comparable contributions by dOLs and $_{LC}$OLs, the vast

majority of neocortical OLs were dOLs but not $_{LC}$OLs. The density of dOLs is significantly

higher than that of $_{LC}$OLs in the motor cortex (Mo) (504.5±25.1 vs. 5.3±0.7; P<0.001; n=3)

and somatosensory cortex (SS) (518.0±22.6 vs.21.6±2.3; P<0.001; n=3) (Figure 2C, D, F).

Co-staining with aspartoacylase (ASPA)[16], a marker for mature OL, also revealed a very

high neocortical OL labeling rate in *Opalin^{Flp}::Emx1^{Cre}::Ai65* (RPF+ASPA+/ASPA+: 94.6±

2.0% in Mo and 90.5±2.2% in SS; n=3).Considering the possibility of incomplete

recombination in combinatorial reporters, and the relatively low Cre activity in the dorsal

MGE of *Nkx2.1^{Cre}*[1], the genuine contribution of $_{LC}$OLs to the neocortex could be even lesser

than our current observation. Therefore, the large quantity of neocortical OLs labeled by

Gsh2^{Cre} in previous study[2] or by GFP in *Opalin^{Flp}::Gsh2^{Cre}::RC::FLTG* (Supplementary

Figure 1B) most likely were predominantly dOLs generated by Gsh2+ dorsal progenitors[10],

rather than bona fide $_{LC}$OLs.

The second major deviation is that cortical $_{MP}$OLs are not completely depleted postnatally,

instead, they make a small but continued contribution with a unique spatial distribution

pattern (Figure 2D and Supplementary Figure 2). $_{MP}$OLs display a clear rostrocaudal density

decline (Supplementary Figure 2A-B) and a laminar preference towards layer 4 (L4) in SS

(Supplementary Figure 2C-E). In contrast, the distribution of dOLs does not vary significantly

across the rostrocaudal axis but exhibits increased density towards deeper layers

(Supplementary Figure 2A-E). Importantly, we have observed cortical $_{MP}$OLs in mice as old

as one year (Supplementary Figure 2F), well beyond the age analyzed in previous reports[2, 7,

8], suggesting a persisted contribution.

We then turned our attention to the lateral three-layer archicortex, piriform cortex (Pir).

Different from the neocortex, Pir contains comparable density of dOLs and $_{LC}$OLs (114.1±

12.5 vs. 171.7±23.6; P=0.058; n=3) (Figure 2F). $_{MP}$OLs make the lowest contribution at a

density similar to SS(16.4±2.4 vs.15.8±1.4; P=0.966; n=4) (Figure 2E).

These combinatorial models also grant us the opportunity to revisit the differential

contributions of dOLs, $_{LC}$OLs and $_{MP}$OLs to the two commissural white matter tracts, corpus

callosum (cc) and anterior commissure (ac), which contain higher density of OLs than the

grey matter (Figure 3). We found that, similar to the neocortex, cc is mainly populated by

dOLs and supplemented by very low density of $_{LC}$OLs and $_{MP}$OLs(Figure 3A-D). Interestingly,

$_{LC}$OLs and $_{MP}$OLs seem to show preferential distribution in the lateral and medial regions of cc

(cc-l and cc-m), respectively (Figure 3B-D, boxed regions in right panels). Different from cc,

ac is mainly populated by $_{LC}$OLs (2454.6±254.1, n=3) and $_{MP}$OLs (1146.3±101.6; n=4) and

supplemented by very low density of dOLs (85.2±34.5; n=3) (Figure 3D).

To substantiate the above results, we further breed *Opalin^{Flp}::Emx1^{Cre}::Nkx2.1^{Cre}::Ai65* to label

dOLs together with $_{MP}$OLs by RFP and co-stained them with ASPA (Supplementary Figure 3).

RFP-ASPA+ cells were difficult to find in Mo, SS and cc, but were more easily observed in

Pir   and ac, consistent with the respective low and high $_{LC}$OL densities in these regions.

In summary, our findings significantly revised the canonical model of forebrain OL origins

(Figure 4A), and provided a new and more comprehensive view (Figure 4B). We

demonstrated that neocortical OLs are mainly derived from dorsal origin with small but

lasting contribution from the ventral origin (Figure 2, Supplementary Figure 1A). Our data

showed that LGE/CGE makes little contribution to neocortex and cc, but makes major

contribution to piriform cortex and ac. This finding is supported by another report[17] in

which in utero electroporation failed to label LGE-derived cortical OLs. Taken together, these

results suggest that the lack of $_{LC}$OL in neocortex is less likely caused by competitive

postnatal elimination, but more likely due to limited production and/or allocation. We further

discovered that MGE/POA makes a small but persistent contribution to the neocortex with a

distinct distribution pattern featured by a rostral-high to caudal-low gradient and a preference

towards L4 in SS. Whether their enduring existence and highly biased localization has

functional implications awaits future exploration. In addition, we found that the cc showed a

similar OL composition as the neocortex, but the Pir and the ac each exhibited distinct OL

compositions in term of their embryonic origins. $_{LC}$OLs are the major contributor to both

regions, while dOLs and $_{MP}$OLs mainly contribute to Pir and ac, respectively.

In addition to the new framework of forebrain OL origins (Figure 4), we also generated a new

driver (Figure 1) and established multiple combinatorial genetic models (Figure 2) for

efficient tracking and direct visualization of OLs from different embryonic origins without

interference from other cells types. These tools set up a firm foundation and will provide

reliable experimental access for future inquiries on the development and function of diverse

OLs in healthy and disease brains[18], especially to uncover the relationship between

developmental origins and the functional and molecular heterogeneity of OLs.

Material and Methods

Mice

All mouse studies were carried out in strict accordance with the guidelines of the Institutional

Animal Care and Use Committee of Fudan University. All husbandry and experimental

procedures were reviewed and approved by the same committee. All applicable institutional

and/or national guidelines for the care and use of animals were followed. The following

transgenic mouse lines were used in this study: $Nkx2.1^{Cre}$ (Jax 008661)[1],$Gsh2^{Cre}$(Jax

025806)[2], $Emx1^{Cre}$ (Jax 005628)[3], $Ai65$ (Jax 021875)[19], $RC::FLTG$ (Jax 026932)[15].

The tTA2-dependent tdTomato reporter (TRE-RFP) was derived from $Ai62$ (Jax 022731) [19],

by removing LoxP-STOP-LoxP with $E2a$-Cre (Jax 003724). The Flp dependent H2B-GFP

reporter (HG-FRT) was derived from HG-$dual$ (Jax 028581) via removal of loxP flanking

STOP cassette by CMV-Cre[20]. The $Opalin^{P2A\text{-}Flpo\text{-}T2A\text{-}tTA2}$ allele was generated by targeted

insertion of the T2A-Flpo-P2A-tTA2 sequence immediately before the STOP codon of the

endogenous $Opalin$ gene using homologous recombination. Gene targeting vector was

generated using PCR-based cloning approach as described before[20]. More specifically, a

4.7 kb 5' homology arm, a loxP flanking Neo positive selection cassette, a

T2A-FlpO-P2A-tTA2 cassette and a 2.7 kb 3' homology arm were cloned into a building

vector containing the DTA negative selection cassette to generate the targeting vector.

Targeting vector was linearized and transfected into a C57/black6 ES cell line. ES clones that

survived through negative and positive selections were first screened by genomic PCR, then

confirmed by Southern blotting using appropriate DIG-dUTP labelled probes. One positive

ES cell clone was used for blastocyst injection to obtain male chimera mice carrying the

modified allele following standard procedures. Chimera males were bred with C57BL/6J

females to confirm germline transmission by genomic PCR. The Neo selection cassette was

self-excised during spermatogenesis of F0 chimeras. Heterozygous F1 siblings were bred with

one another to establish the colony. Targeting vector construction, ES cell transfections and

screening, blastocyst injections and chimera breeding were performed by Biocytogen Co,

China.

Genomic PCR

Genomic DNA was prepared from mouse tails. Tissue was lysed by incubation in tail lysis

buffer (Viagen, 102-T) with 0.1 mg/ml proteinase K (Diamond, A100706) overnight at 55°C

followed by 45 min at 90°C in an air bath to inactivate proteinase K. The lysate was cleared

by centrifugation at maximum speed (21,130 G) for 15 min in a table-top centrifuge.

Supernatant containing genomic DNA was used as the PCR template for amplifying DNA

products. The following primers were used:

Opalin-F: 5'- GGCCTATGTTTGATTTCCAGCACTG -3'

Opalin-R: 5'- AGCACTTATGACTGCTGAGCCGTTC -3'

Immunohistochemistry and microscopy

Mice were anaesthetized by intraperitoneal injection of 1.5% sodium pentobarbital (0.09 mg/g

body weight) and then intracardially perfused with saline followed by 4% paraformaldehyde

in 0.1 M PB. Following post fixation at 4°C for 24 hours, brain samples were sectioned at 30

μm using a vibratome (Leica VT1000S), or transferred into 30% sucrose in 0.1M PB for

cryoprotection, OCT embedded, and sectioned using a cryostat (Leica CM1950). For CC1

immunostaining, antigen retrieval was performed prior to blocking by boiling for 3 min in 10

mM citrate buffer (pH6.0). Sections were blocked in PBS containing 0.05% Triton and 5%

normal donkey serum and then incubated with the following primary antibodies in the

blocking solution at 4 °C overnight: RFP (goat polyclonal antibody, 1:2000, SICGEN

AB0081-200; rabbit polyclonal antibody, 1:2000, Rockland 600-401-379), GFP (chicken

polyclonal antibody, 1:1000, Aves Labs, GFP-1020), MBP (rat polyclonal antibody, 1:500,

AbD Serotec, MCA409S), CC1 (rabbit polyclonal antibody, 1:500, Oasis Biofarm,

OB-PRB070, mouse polyclonal antibody, 1:300, Millipore, OP80), ASPA (rat polyclonal

antibody, 1:200, Oasis Biofarm, OB-PRT005), Sox9(rabbit polyclonal antibody, 1:2000,

Chemicon, AB5535), NeuN (mouse monoclonal antibody, 1:500, Millipore, MAB377).

Sections were then incubated with appropriate Alexa fluor dye-conjugated IgG secondary

antibodies (1:500, Thermofisher Scientific) or CF dye-conjugated IgG secondary antibodies

(1:250, Sigma) in blocking solution and mounted in Aqua-mount (Southern Biotech, 0100-01).

Sections were counterstained with DAPI. Sections were imaged with confocal microscopy

(Olympus FV3000), fluorescence microscopy (Nikon Eclipse Ni; Olympus VS120; Olympus

VS200) and fluorescent stereoscope (Nikon SMZ25). Anatomical regions in the postnatal

brain were identified according to the *Paxinos "The Mouse Brain" Atlas* and the *Allen*

Reference Atlas, and their areas were measured in ImageJ, whenever applicable. One out of

every four sections within the range of selection from each brain and at least three brains per

genotype were analyzed. To quantify density and co-localization, cells were identified and

counted in Image J or QuPath.

Statistical analysis

GraphPad Prism version 8.0.1 was used for statistical calculations. No statistical methods

were used to predetermine sample sizes, but our sample sizes are similar to those reported in

previous publications. Data collection and analysis were performed blind to the conditions of

the experiments whenever possible. No animals or data points were excluded from the

analysis. Normalcy was assessed using Shapiro-Wilk test. Equal variances were assessed

using F test or Bartlett's test. Statistical significance was tested using two-tailed unpaired

t-test, Welch's t-test, one-way ANOVA and two-way ANOVA followed by Tukey's or

Bonferroni post hoc test, wherever appropriate. Data are presented as mean $\pm$ standard error

of the mean (S.E.M.). $P < 0.05$ was considered significant. Significance is marked as $*P <$

0.05; $**P < 0.01$ and $***P < 0.001$.

Acknowledgements. We thank Dr. Yilin Tai for helpful discussion and Dr. Min Jiang from

core facility of IOBS, Fudan University, for technical support on imaging experiments. This

study was supported by funds from the National Science and Technology Innovation 2030

Major Projects of China (STI2030-Major Projects-2022ZD0206500), National Natural

Science Foundation of China (32171087, 32200918, 32371145).

Author contributions.

Conceptualization: MH

Methodology: YC, MH

Investigation: YC, ZZ, MS

Visualization: YC, MH, LG

Supervision: MH, LG

Writing—original draft: MH, YC, LG

Writing—Review and editing: MH, YC, ZZ, MZ

Declaration of interests. The authors declare no competing interests.

Reference

1.   Xu, Q., M. Tam, and S.A. Anderson, *Fate mapping Nkx2.1-lineage cells in the mouse*
telencephalon. J Comp Neurol, 2008. **506**(1): p. 16-29.

2.   Kessaris, N., et al., *Competing waves of oligodendrocytes in the forebrain and postnatal*
elimination of an embryonic lineage. Nat Neurosci, 2006. **9**(2): p. 173-9.

3.   Gorski, J.A., et al., *Cortical excitatory neurons and glia, but not GABAergic neurons, are*
produced in the Emx1-expressing lineage. The Journal of neuroscience : the official journal of
the Society for Neuroscience, 2002. **22**(15): p. 6309-14.

4.   Nakahira, E., et al., *Direct evidence that ventral forebrain cells migrate to the cortex and*
contribute to the generation of cortical myelinating oligodendrocytes. Dev Biol, 2006. **291**(1):
p. 123-31.

5.   Tsoa, R.W., et al., *Spatiotemporally different origins of NG2 progenitors produce cortical*
interneurons versus glia in the mammalian forebrain. Proc Natl Acad Sci U S A, 2014.
111(20): p. 7444-9.

6.   Naruse, M., et al., *The Dorsoventral Boundary of the Germinal Zone is a Specialized Niche*
for the Generation of Cortical Oligodendrocytes during a Restricted Temporal Window. Cereb
Cortex, 2016. **26**(6): p. 2800-2810.

7.   Orduz, D., et al., *Developmental cell death regulates lineage-related*
interneuron-oligodendroglia functional clusters and oligodendrocyte homeostasis. Nat
Commun, 2019. **10**(1): p. 4249.

8.   Liu, R., et al., *In Vivo Clonal Analysis Reveals Development Heterogeneity of Oligodendrocyte*
Precursor Cells Derived from Distinct Germinal Zones. Adv Sci (Weinh), 2021. **8**(20): p.
e2102274.

9.   Shen, Z., et al., *Distinct progenitor behavior underlying neocortical gliogenesis related to*
tumorigenesis. Cell Rep, 2021. **34**(11): p. 108853.

10.   Zhang, Y., et al., *Cortical Neural Stem Cell Lineage Progression Is Regulated by Extrinsic*
Signaling Molecule Sonic Hedgehog. Cell Rep, 2020. **30**(13): p. 4490-4504 e4.

11.   Kippert, A., et al., *Identification of Tmem10/Opalin as a novel marker for oligodendrocytes*
using gene expression profiling. BMC Neurosci, 2008. **9**: p. 40.

12.   Yoshikawa, F., et al., *Opalin, a transmembrane sialylglycoprotein located in the central*
nervous system myelin paranodal loop membrane. J Biol Chem, 2008. **283**(30): p. 20830-40.

13.   Jiang, W., et al., *Identification of Tmem10 as a novel late-stage oligodendrocytes marker for*
detecting hypomyelination. Int J Biol Sci, 2013. **10**(1): p. 33-42.

14.   Marques, S., et al., *Oligodendrocyte heterogeneity in the mouse juvenile and adult central*
nervous system. Science, 2016. **352**(6291): p. 1326-1329.

15.   Plummer, N.W., et al., *Expanding the power of recombinase-based labeling to uncover*
cellular diversity. Development, 2015. **142**(24): p. 4385-93.

16.   Huang, H., et al., *Immunological Markers for Central Nervous System Glia.* Neurosci Bull,
261   2023. **39**(3): p. 379-392.
262   17.   Li, J., et al., *The Lateral Ganglionic Eminence Does Not Generate Cortical Oligodendrocytes.*
263   bioRxiv, 2023: p. 2023.12.01.569674.
264   18.   Gong, L., et al., *Emerging strategies for the genetic dissection of gene functions, cell types,*
265   *and neural circuits in the mammalian brain.* Mol Psychiatry, 2022. **27**(1): p. 422-435.
266   19.   Madisen, L., et al., *Transgenic mice for intersectional targeting of neural sensors and*
267   *effectors with high specificity and performance.* Neuron, 2015. **85**(5): p. 942-58.
268   20.   He, M., et al., *Strategies and Tools for Combinatorial Targeting of GABAergic Neurons in*
269   *Mouse Cerebral Cortex.* Neuron, 2016. **91**(6): p. 1228-1243.

Figures

Figure 1. A new driver mouse for efficient and specific OL labeling

(A) Scheme for generating the *Opalin*$^{P2A-Flp-T2A-tTA2}$ allele. (B) Southern blot confirmation of
correctly targeted ES clone. (C) Genomic PCR to genotype F1 offspring. (D) OL labeling by
Flp. (E) OL labeling by tTA2. High magnification images of the boxed region showing
co-localization of RFP with MBP staining, which further demonstrated the myelination ability
of labeled OLs. (F) Quantification of marker staining. Left panel quantified the labeling
specificity. Both systems are highly specific, as shown by the complete co-localization with
OL marker (CC1) and lack of co-staining by the neuronal marker (NeuN) or astrocyte marker
(Sox9). Right panel quantified the labeling efficiency. Close to complete OL labeling was
achieved by Flp-dependent H2B-GFP reporter in all analyzed regions, while sparser labeling
with variable regional density was achieved by tTA2-dependent tdTomato reporter driven by
TRE promoter. Scale bar: 50 μm in low magnification images, 5μm in high magnification
images. Quantification: n=3. Dots represent data from individual mice.

Figure 2. Combinatorial fate mapping of dOLs, $_{MP}$OLs and $_{LC}$OLs

(A) Strategy for intersectional labeling. Flp-AND-Cre labels OLs from Cre-expressing progenitors with RFP. (B) Strategy for simultaneous intersectional and subtractional labeling of OLs derived from complementary origins. Flp-NOT-Cre labels OLs from non-Cre-expressing progenitors with RFP, while Flp-AND-Cre labels OLs from Cre-expressing progenitors with GFP. (C-E) Intersectional labeling of dOLs in *Opalin^{Flp}*::*Emx1^{Cre}*::*Ai65*, subtractional labeling of $_{LC}$OLs in *Opalin^{Flp}*::*Emx1^{Cre}*::*Nkx2.1^{Cre}*::*RC::FLTG*, and intersectional labeling of $_{MP}$OLs in *Opalin^{Flp}*::*Nkx2.1^{Cre}*::*Ai65* and their cortical distribution. Schematics, representative images and quantification of RFP+ cell density in motor cortex (Mo), somatosensory cortex (SS) and piriform cortex (Pir) were displayed. (F) Differential contribution of OLs from the three embryonic origins to Mo, SS and Pir.

Scale bar: 1mm in low magnification images, 500 μm in high magnification images of the boxed area. Quantification: n=3 for dOLs and $_{LC}$OLs; n=4 for $_{MP}$OLs. Dots represent data from individual mice. **P < 0.01, ***P < 0.001. Error bar: S.E.M.

Figure 3. dOLs, _MP_OLs and _LC_OLs differentially contribute to the two major white matter tracts

(A-C) Schematics and representative images of dOLs, _MP_OLs and _LC_OLs in the two major commissure white matter tracts: corpus callosum(cc) and anterior commissure (ac). _MP_OLs and _LC_OLs preferentially reside in the medial and lateral cc (cc-m and cc-l), respectively. (D) Quantification of dOL, _MP_OL and _LC_OL densities in cc-m, cc-l and ac. Scale bar: 1mm in low magnification images, 500μm in cc and ac, 100 μm in high magnification images of the boxed area (cc-m and cc-l) in cc. Quantification: n=3 for dOLs and _LC_OLs; n=4 for _MP_OLs. Dots represent data from individual mice. **P < 0.01, ***P < 0.001. Error bar: S.E.M.

Figure 4. The classical and revised model of forebrain OL origins

(A) In the classical model[2], OLs derived from MGE/POA (orange) were largely eliminated postnatally (thin dashed line), while those from LGE/CGE (blue) and dorsal origin (purple) survive at similar proportions (thick solid line). Therefore, neocortex (NCx) and corpus callosum (cc) contain comparable density of $_{LC}$OLs (blue dots) and dOLs (purple dots) and are devoid of $_{MP}$OLs (orange dots). (B) In the new model, NCx and cc mainly contains dOLs with very low contribution from the ventral origins. $_{LC}$OLs mainly contribute to Pir and ac. $_{MP}$OLs makes a small but sustained contribution to NCx, with a strong laminar preference towards layer 4 in somatosensory cortex (SS). In addition, dOLs and $_{MP}$OLs also make substantial contributions to Pir and ac, respectively. Grey dots indicate OLs in unanalyzed regions.

Supplementary information

Supplementary Figure 1. Simultaneous differential labeling of OLs derived from
complementary embryonic origins

(A) Coronal sections showing GFP+ OLs from dorsal origin (dOLs) and RFP+ OLs from
ventral origin (vOLs) in $Opalin^{Flp}$::$Emx1^{Cre}$::RC::$FLTG$. (B) Coronal sections showing
GFP+ OLs derived from Gsh2+ progenitors ($_{Gsh2+}$OL) and RFP+ OLs derived from Gsh2-
progenitors ($_{Gsh2-}$OL) in $Opalin^{Flp}$::$Gsh2^{Cre}$::RC::$FLTG$. Scale bar: 1mm.

Supplementary Figure 2. The distribution pattern of neocortical dOLs and $_{MP}$OLs

(A-B) Rostrocaudal distribution of neocortical dOLs and $_{MP}$OLs. (A) Within the range of
analysis, RFP+ OLs in neocortex of every fourth section were quantified. (B) Slices were
grouped into seven bins numbered rostrocaudally. Density of dOLs showed no significant
change across bins, while $_{MP}$OLs exhibited lower density in more caudal regions. (C-E)
Laminar distribution of dOLs and $_{MP}$OLs in somatosensory cortex (SS). (C) Representative
images of SS from P56 mice. (D) Distribution of dOLs and $_{MP}$OLs across 6 layers. More
dOLs reside in deeper layers, while $_{MP}$OLs are highly enriched in L4. (E) dOLs and $_{MP}$OLs
exhibited significant differences in their relative laminar distributions. (F) Representative
image of SS from 1 year old *Opalin^{Flp}::Nkx2.1^{Cre}::Ai65* mouse. Scale bar: 200 μm. Dots
represent data from individual mice. Error bar: S.E.M. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Supplementary Figure 3. Intersectional labeling of OLs derived from both dorsal origin and MGE/POA

(A) Coronal sections showing both dOLs and $_{MP}$OLs labeled by RFP in *Opalin^{Flp}::Emx1^{Cre}::Nkx2.1^{Cre}::Ai65.* (B) Higher magnification images showing ASPA+ RFP+ dOLs/$_{MP}$OLs (closed arrow heads) and ASPA+RFP- putative $_{LC}$OLs (open arrow heads).The latter cells were difficult to find in neocortical regions such as motor cortex (Mo) and somatosensory cortex (SS), and corpus callosum (cc), but were frequently encountered in piriform cortex (Pir) and anterior commissure (ac). Scale bar: 1mm in low magnification images, 200μm in Mo, SS, cc and ac, 10 μm in high magnification images of boxed area (all shared the same scale).