Single-cell RNA sequencing of iPSC-derived brain organoids reveals

Treponema pallidum infection inhibiting neurodevelopment

Qiu-Yan Xu[1,†], Yong-Jing Wang[1,2,†], Yun He[1,†], Xin-Qi Zheng[1], Man-Li Tong[1,3,*], Yu Lin[1,3,*], and Tian-Ci Yang[1,3,*]

[1] Centre of Clinical Laboratory, Zhongshan Hospital of Xiamen University, School of Medicine, Xiamen University, Xiamen, 361004, China.

[2] Guangyuan Hospital of Traditional Chinese Medicine, Guangyuan, 628000, China.

[3] Institute of Infectious Disease, School of Medicine, Xiamen University, Xiamen, 361004, China.

[†] These authors have contributed equally to this work.

* For correspondence: yangtianci@xmu.edu.cn (T.-C. Y.);

linyuxmuzsyy0620@xmu.edu.cn (Y. L.); tongmanli@xmu.edu.cn (M.-L. T.)

Abstract

Congenital syphilis is a vertically transmitted bacterial infection caused by *Treponema pallidum*, often causing multidomain neurodevelopmental disabilities. However, little is known about the pathogenesis of this disease. Brain organoids platform derived from the induced pluripotent stem cell (iPSC) is employed and exposed to *T. pallidum* infection for modelling congenital neurodevelopmental impairment. Single-cell RNA sequencing is used for identifying the subpopulations of differentially expressed genes and cellular heterogeneity and reconstructing differentiation trajectories following *T. pallidum* infection. The results reveal that *T. pallidum* infection influences the formation of neural rosette structures, reduces the cell number of the neural progenitor cell subcluster 1B (subNPC1B) and hindbrain neurons, and affects the neurodevelopment of the brain organoid. Moreover, it is speculated that *T. pallidum* inhibit the hindbrain neuron cell number through the suppression of subNPC1B subgroup in the organoids and inhibits transcription factor 3 activity in the subNPC1B-hindbrain neuronal axis. This is the first report on the inhibited effects of *T. pallidum* on the neurodevelopment of the iPSC-derived brain organoid model. It is concluded that *T. pallidum* could inhibit the differentiation of subNPC1B in brain organoids, thereby reducing the differentiation from subNPC1B to hindbrain neurons, and ultimately affecting the development and maturation of hindbrain neurons.

Keywords: Congenital syphilis, *Treponema pallidum*, Brain organoids, Single-cell RNA sequencing, Neurodevelopment.

Introduction

Congenital infections, which refer to maternal infections with vertical transmission to the fetus, pose a significant global public health issue and are associated with adverse pregnancy outcomes, neonatal diseases, long-term neurodevelopmental consequences, and increased healthcare expenditures(Grosse et al., 2021). Notably, an expanding roster of congenital infections, such as cytomegalovirus, Zika virus, syphilis, rubella, lymphocytic choriomeningitis virus, and toxoplasmosis, have been implicated in neurodevelopmental impairments(Fortin and Mulkey, 2023). With the increasing number of syphilis cases in recent years in many countries, there has been a growing focus on the prenatal exposure of the fetus to maternal *Treponema pallidum* and its consequential effects on the developing brain and subsequent neurodevelopmental outcomes(David et al., 2022; Medoro and Sánchez, 2021). Tonni et al. utilized a transabdominal neuroscan of pregnant woman with primary syphilis and observed hyperechogenicity involving both choroid plexuses of the lateral ventricles and the falx cerebri(Tonni et al., 2022). Moreover, the fetal brain contour exhibited a smooth appearance devoid of discernible gyri and sulci, indicating a significant influence of *T. pallidum* on fetal brain development and subsequent neurodevelopmental outcomes(Tonni et al., 2022). However, the precise mechanisms underlying the impact of *T. pallidum* on fetal neurodevelopment remain constrained.

Self-renewing and pluripotency properties of human pluripotent stem cells have greatly facilitated the understanding of the human nervous system development and pathogenesis of various neurological diseases(Mertens et al., 2016; Ramani et al., 2020). Human brain organoids derived from induced pluripotent stem cells (iPSCs), which more closely resemble the developing human brain than previous approaches(Lancaster et al., 2013), provide unprecedented opportunities for studying viral infections and their impact on human neurodevelopment(Kim et al., 2019; Krenn et al., 2021). The development of organoids requires robust strategies to assess the cellular composition of organoids and their complex interactions and neuronal networks. Single-cell RNA sequencing (scRNA-seq), a high-resolution approach that

fits the above requirement, can deconstruct organoid cell composition, help to study

the molecular underpinnings of cellular heterogeneity, and reconstruct differentiation

trajectories in these complex developing tissues(Kanton et al., 2019; Velasco et al.,

2019). Several laboratories using scRNA-seq for the direct comparison of cultured

and primary cortical structures *in vitro* and *in vivo*, respectively, demonstrated great

similarity in the gene expression profiles(Camp et al., 2015; Pollen et al., 2019).

However, understanding and interpreting gene expression and regulatory patterns

from single-cell genomic data on organoids in the field of syphilis remains a major

challenge. Here, we constructed the iPSC-derived brain organoids and infected them

with *T. pallidum*, and evaluated the neurodevelopment of the brain organoids based on

the scRNA-seq to model and analyse the pathogenesis of neurodevelopmental

disorders of congenital syphilis. This is the first study to evaluate the effects of *T.*

pallidum on the neural development of an iPSC-derived brain organoid model,

thereby laying the foundation for further research elucidating the pathogenesis of

congenital syphilis.

Results

***T. pallidum* infection inhibited brain organoid neurodevelopment**

To study the neurodevelopment of *T. pallidum*-infected brain organoids instead of

patient-derived brain tissue, we first generated iPSC-derived brain organoids for

nearly 2 months as previously described(Madhavan et al., 2018; Setoh et al., 2019)

(Figure 1A, Supplement file- figure supplement 1). Immature brain organoids on day

25, which coincides with the gradual differentiation and maturation of brain organoid,

were utilised for *T. pallidum* infection. Brain organoid growth was tracked over 30

days post-infection to monitor organoid growth and development. At day 20

post-infection, the size on average of the *T. pallidum*-infected organoids was

significantly smaller than that of the control ($P<0.01$) (N=15 organoids from 3

separate bioreactors per group), and the edges of the organoids were uneven, similar

to signs of disintegration; while the control group grew well (Figure 1B). To address

whether *T. pallidum*-infected organoid modulates the expression of gene markers for

three germ layers in neurodevelopment, i.e., endoderm, mesoderm and ectoderm,

quantitative real-time PCR (qRT-PCR) was performed (N=15 organoids from 3

separate bioreactors per group). The qRT-PCR analysis revealed that the expressions

of endodermal marker Ve-cad, KDR, and SOX17 were significantly upregulated in

the *T. pallidum* group ($P < 0.05$). In contrast, expressions of the ectodermal genes

MAP2, Nestin, and SOX2 were significantly decreased ($P < 0.05$) (Figure 1C). This

finding was confirmed by immunofluorescent staining against TUBB3 and SOX2,

which are expressed in the central nervous system from the earliest developmental

stages throughout vertebrate evolution(Mercurio et al., 2019). *T. pallidum* infection

resulted in brain organisational changes in neural rosette-like structures resembling

the proliferative regions of the human ventricular zone, and loss of structural integrity

(Figure 1D) (N=15 organoids from 3 separate bioreactors per group). These results

suggest that *T. pallidum* infection influenced the formation of neural rosette structures

and affected the neurodevelopment of the brain organoid.

scRNA-seq revealed that *T. pallidum* infection inhibited the differentiation of

neural progenitor cell subcluster 1B in the brain organoid

Individual genetic markers did not address the cellular diversity of the human brain,

where closely related cell classes can be identified by combining genetic signatures,

while single-cell RNA sequencing allows for systematic analysis of many genes. To

better understand the effects of *T. pallidum* infection on organoid development,

especially cell type specification changes, scRNA-seq was performed to molecularly

profile 21,504 single cells isolated from 12 organoids grown in 3 separate bioreactors

on day 55 (after *T. pallidum* treatment for 30 days) (Figure 2A). After quality control,

a mass of data from single cells of brain organoids from the *T. pallidum* (10,138 cells)

and control groups (11,366 cells) were obtained. All single-cell transcriptomes

underwent unsupervised clustering using *t*-distributed stochastic neighbour

embedding (tSNE), yielding a total of 19 clusters (Figure 2B), which were then

assigned to 13 cell types based on the known marker-gene expression enrichments

(Figure 2C, Supplement file- figure supplement 2). Thereafter, we compared the

regional specification in organoids between two groups, and analysed them at the

single-cell level. The proportions of neural progenitor cell (NPC)1, NPC2 and neuron

population were 12.5%, 1.72%, and 9.1%, respectively, following *T. pallidum*

infection ($P < 0.05$), while the proportion of the neural ectoderm cell, neural epithelial

cell, and undefined cells in the *T. pallidum* group were higher than that in the control

group ($P < 0.05$) (Figure 2D). To identify the NPC subclusters that could be

particularly vulnerable to *T. pallidum* infection, we analysed the expression of

candidate genes across the NPC cell populations on brain organoids following *T.*

pallidum infection. Heatmaps revealed that the key transcription factors of neural

differentiation, such as HOTAIRM1, HOXB2, and HOXA2 were downregulated

following *T. pallidum*-infection in the NPC1 cluster and genes regarded as neural

repairing markers, such as EN2, IGFBP1 and SPRY1 were upregulated following *T.*

pallidum infection (Figure 2E). However, no change in these genes was found in the

NPC2 cluster between the two groups (Supplement file- figure supplement 3),

illustrating that *T. pallidum* directly inhibited the development of the NPC1 cluster.

To probe the effect of *T. pallidum* in the NPC1 cluster development of brain

organoids, we first performed a whole-transcriptome analysis of the NPC1 cluster to

discriminate any changes in the transcriptional level between the *T. pallidum* (1,290

cells) and control groups (2,114 cells) (Figure 2F). Then, Gene ontology (GO) term

analysis of the upregulated and downregulated genes were identified, respectively.

The upregulated genes revealed that the biological processes were mainly those

related to metabolic process and blood vessel genesis, and the molecular function was

closely associated with pathogenic infection. Moreover, the biological processes

mainly enriched those processes associated with neural development and

neurofunctional zoning, and the molecular function enriched genes promoting neural

differentiation with downregulated expression (Figure 2G).

To dissect the specific cell type(s) affected by *T. pallidum* in the NPC1 population,

we re-clustered and subdivided the NPC1 population into five subclusters (subNPC1

A–E) based on the high expression of general NPC1 subcluster markers (Figure 2H).

The proportions of subclusters between the two groups were analysed by uniform

manifold approximation and projection (UMAP) and clustering of the NPC1

population. Comparative analysis of the NPC1 subclusters between the two groups

demonstrated that *T. pallidum* downregulated the number of subNPC1B population (*P*

< 0.05); however, no significant difference was observed in the other subclusters. The

highly expressed genes in the subNPC1B population were related to the HOX family,

which is considered positively regulated for regional division and functional

formation during brain development (Figure 2I). Overall, these results indicated that *T.*

pallidum reduced the cell number of subNPC1B subclusters and affected brain

organoid development.

scRNA-seq revealed that *T. pallidum* infection inhibited neuronal differentiation

in the brain organoid

To infer inhibition in the neuron cluster by *T. pallidum*, differential genes were

grouped for volcano plot analysis (2292 cells) (Figure 3A). In the neuron population,

compared with those in the control group, the upregulated and downregulated genes

in the *T. pallidum* group were evaluated by GO analysis; the main functions of these

upregulated genes were enriched in the items related to neuronal apoptosis and

endocytosis, while the downregulated genes were enriched for cerebrum functional

partition and energy metabolism in neuronal related functions, including neuron fate

commitment, brain segmentation, neural tube development, etc (Figure 3B). These

results indicated that the altered genes following *T. pallidum* infection were closely

linked to the neuronal dysfunctional and cerebral partition. To examine the abundance

of target gene transcripts based on scRNA-seq data, eight neural-specific highly

expressed genes were selected and validated. The results demonstrated that the early

neural-specific transcription factors HOXA5, HOXC4, HOXC5, and NEUROG1; the

positive regulatory factors during neural maturation PCP4, HOTRAIRM1 and GNG5;

and the key RNA binding protein for nerve system segmentation HNRNPH1 were

significantly downregulated following *T. pallidum* infection (*P* <0.01–0.001) (Figure

3C). We subsequently validated the genes by performing qRT-PCR, and found that the

HOXA5, HOXC5, and HOXA4 genes of *T. pallidum* group were significantly lower

than that of the control group (P <0.05–0.01) (Figure 3D) (N=15 organoids from 3

separate bioreactors per group). Immunostaining of day 55 brain organoids of neuron

marker-MAP2 revealed that the neural rosette-like structure was damaged following *T.*

pallidum infection (Figure 3E) (N=15 organoids from 3 separate bioreactors per

group).

Following the confirmation that CD71 and CD49b are expressed in pluripotent

stem cells, neural crest, neural stem cells but not in neurons(Menon et al., 2019), the

identified difference was expended to develop novel cell selection strategies. We

isolated CD71⁻/CD49b⁻ subpopulations from day 55 brain organoids (N=21

organoids from 4 separate bioreactors per group) differentiation using

fluorescence-activated cell sorting (FACS), which revealed the proportion of

CD71⁻/CD49b⁻ neurons in the *T. pallidum* group was significantly lower than that in

the control group (P < 0.01), indicating that *T. pallidum* inhibited the normal

differentiation of brain organoids into mature neurons (Figure 3F). The phase contrast

images of 2-day cultured post- CD71⁻/CD49b⁻ subpopulation sorting demonstrated

that the neurite of CD71⁻/CD49b⁻ neuron populations were sparse and mostly

fragmented in the *T. pallidum* group, and the immunocytochemistry of neuronal

markers-MAP2 increased perinuclear immunofluorescence were affected by *T.*

pallidum (Figure 3G). These results indicate that *T. pallidum* inhibited the

differentiation of brain organoid neurons.

scRNA-seq revealed that *T. pallidum* infection inhibited the differentiation of the

hindbrain neurons of the brain organoid

Neuron segmentation is a key step for neuronal digital reconstruction, which is

essential for exploring brain circuits and understanding brain function(Li and Shen,

2022; Sugino et al., 2019). Distinct neuronal cell types acquire and maintain their

identity by expressing different genes(Nelson et al., 2006). To clarify the region that

was specifically inhibited by *T. pallidum*, UMAP analysis revealed the production of

both unique and overlapping cell types between the two groups (Figure 4A). Based on

the cerebral cortex-specific markers in each segmentation, the neuron cluster was

grouped into four subclusters: hindbrain neuron, choroid plexus, noncerebral neuron,

and cortical neuron (Figure 4B). Thereafter, the cell number proportions of four

subclusters within the two groups were calculated; the proportion of hindbrain neuron

subclusters decreased to 14.23% following *T. pallidum* infection, while the

proportions of the other three subclusters showed no obvious changes between the

two groups (Figure 4C). The neural-specific highly expressed genes of four

subclusters were selected and validated, and the results depicted that the hindbrain

neuronal markers MAGEH1, MEIS3 and USP47 were significantly decreased

following *T. pallidum* infection ($P<0.05$–0.01), whereas no significant alterations

were detected in the other three subclusters between the two groups ($P > 0.05$) (Figure

4D). Furthermore, immunostaining for hindbrain neuron specific markers-SHOX2

and HOXA5 in the cryosections from day 55 brain organoid revealed that the

fluorescence intensity of SHOX2 and HOXA5 decreased following *T. pallidum*

infection (Figure 4E) (N=15 organoids from 3 separate bioreactors per group).

Overall, these results suggest that *T. pallidum* inhibited the differentiation of

hindbrain neurons in the brain organoids.

scRNA-seq revealed the mechanism of *T. pallidum* affecting neurodevelopment in

the brain organoids

To study the differentiation of subNPC1B into hindbrain neuron cluster and the

corresponding gene expression, we selected HES4 and NEUROGD1 to construct a

new trajectory corresponding to two distinct cell fates. The root of the trajectory was

mainly populated by HES4 for subNPC1B subcluster, while the termini of the tree

was populated by NEUROGD1 for the hindbrain neuron subcluster (Figure 5A).

Thereafter, we assessed the expression of the genes-regulated trajectory of the

subNPC1B-hindbrain neuron. ASPM, CCNB1, TPX2, UBE2C, TOP2A, MKI67,

CENPF, PTTG1 and HMGN2 expression, which were the top genes in subNPC1B,

were markedly reduced from the root to the termini. SNCG, S100A10, NEUROD1,

RGS10, STMN2, PPP1R17, BASP1, NEFL, CALM1, DCX, TMSB10, TUBB3,

TMSB4X, STMN4, C11orf96, TUBA1A, RTN1, SYT4, and SST expression, which

were the top genes in the hindbrain neuron subcluster, were lowly expressed at the early stage of the pseudotime axis, and gradually increased until the end of the pseudotime axis (Figure 5B). These results further confirmed that the differentiation direction is from subNPC1B to the hindbrain neuron.

To determine whether *T. pallidum* inhibited the differentiation of hindbrain neurons by disturbing subNPC1B-hindbrain neuron differentiation axis or by directly affecting the hindbrain neuron, subNPC1B and hindbrain neuron markers were displayed using dot shapes (Figure 5C). The subNPC1B specific markers-LRRC75A-AS1, ZFAS1, SEC11A, GAS5, ID1, and NDUFB5 were decreased following *T. pallidum* infection; the inhibitors markers- BNIP3, PGK1 and IFITM3 during the NPC differentiation were increased; the early NPC markers- POU3F3 and INSIG2 also increased in the *T. pallidum* group; however, there was no difference in the expression of the hindbrain neuronal markers- NEFM, NEFL, ENO2, B4ALT5, SPTAN1, and TTR in either the *T. pallidum* group or control group, indicating that the decreased number of hindbrain neurons was attributed to the disturbance of the subNPC1B- hindbrain neuron differentiation axis.

To uncover the mechanism underlying the subNPC1B-hindbrain neuron differentiation process, GO analysis of the total differential genes in subNPC1B demonstrated that the GO terms were mainly enriched in transcription regulation (Figure 5D), suggesting that the transcriptional regulatory network mediated by the transcription factors of subNPC1B could play a major role during the inhibition of the subNPC1B-hindbrain neuron differentiation process by *T. pallidum*. Next, the single-cell regulatory network inference and clustering (SCENIC) analysis for subNPC1B subcluster was performed to assess the differences in the transcriptional activity of the transcription factors between the two groups, and found that transcription factor 3 (TCF3) was specifically enriched in subNPC1B of the *T. pallidum* group (Figure 5E), speculating that TCF3 is the key transcription factor for the inhibition of subNPC1B-hindbrain neuron differentiation caused by *T. pallidum*. Furthermore, we explored TCF3 function which could be extended to known TCF3-specific downstream factors and pathway activities (Figure 5F), Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis

subsequently performed on the hindbrain neuron subcluster (Figure 5G), and found

that they were in the common notch signaling pathway. Therefore, we speculate that

TCF3 and notch signalling pathway could play important roles in the inhibition of

subNPC1B-hindbrain neuronal differentiation axis by *T. pallidum*, warranting further

study to confirm these findings.

Discussion

Infants born to mothers with *T. pallidum*-infection and who had a positive syphilis

serologic test during pregnancy have a high risk for neurodevelopmental

disorders(Auriti et al., 2022; Verghese et al., 2018), with varied clinical manifestations,

such as bulging fontanelle, seizures, cranial nerve palsies, etc(Mattei et al., 2012;

Woods, 2009). Owing to ethical and practical challenges, guinea pig and rabbit animal

models have been widely employed to study the pathogenesis of congenital

syphilis(Wicher and Wicher, 2001); however, neither neuroinvasive nor congenital

syphilis has been well explained(Tantalo et al., 2005). In the study, we evaluated

iPSC-derived brain organoids for modelling to reveal the pathogenesis of

neurodevelopmental disorders associated with congenital syphilis. The results showed

that *T. pallidum* infection inhibited brain organoid neurodevelopment. This conclusion

is supported by the results showing brain organoids infected with *T. pallidum*

demonstrated significant decrease in the overall organoid size and interfered with the

formation of neural rosette-like structure of brain organoids. Combined with

scRNA-seq analysis, it clarified that *T. pallidum* inhibited the differentiation of

hindbrain neurons through the suppression of subNPC1B in brain organoids.

Furthermore, we found that TCF3 may be a key transcription factor associated with

the inhibition of subNPC1B-hindbrain neuron axis by *T. pallidum*.

As brain organoids reflect developmental stages during pregnancy and the early

postnatal, they are currently the most relevant tools for research on

neurodevelopmental disorders(Andrews and Kriegstein, 2022), helping to discover

novel biological information on the early stages of neurodevelopment and disease

progression, as well as the ability to manipulate these processes *in vitro* (Trujillo and

Muotri, 2018). Tonni et al. reported that a routine third-trimester scan of pregnant

woman with primary syphilis, which revealed fetal growth restriction associated with

absent fetal movement, a pathologic neuroscan characterized by cortical calcifications,

and a transabdominal neuroscan was able to detect hyperechogenicity involving both

choroid plexuses of the lateral ventricles as well as the falx cerebri, with smooth

without recognizable gyri and sulci (Tonni et al., 2022). The imaging features and the

neurodevelopmental outcome in the fetal brain were verified by organoids infected

with *T. pallidum* inducing a significant decrease in the overall organoid size and

interfering with the formation of neural rosette-like structure of brain organoids in our

study. Previous studies have shown that Zika virus-induced cellular death and reduced

size of cortical organoids are associated with the clinical microcephaly phenotype

(Nowakowski et al., 2016; Qian et al., 2016; Watanabe et al., 2017). Whether *T.*

pallidum induced the decrease in overall organoid size and disruption of neural

rosette-like structure in the brain organoids is related to the clinical microcephaly

phenotype required further confirmation in the infection animals.

Uterine environmental pollution taking place in early brain development could

have long-lasting consequences for adult brain functioning(Meyer et al., 2007).

Several studies demonstrated that the development of the central nervous system was

blocked at an immature stage following Zika virus infection(Hughes et al., 2016; Li et

al., 2016; Zhang et al., 2019), preferentially targeting and hindering the growth of

neural precursor cells in the monolayer and cortical brain organoid culture

systems(Zhang et al., 2019). Notably, similar results were observed in our study: the

proportion of neuroectodermal cells, neuroepithelial cells, neural stem cells, and other

cells in the early stages of neurodevelopment of *T. pallidum*-infected organoids were

significantly higher than that the control group, suggesting that organoids infected

with *T. pallidum* retained neural precursor cells in the developmental stage and

inhibited their differentiation in brain organoids towards maturity. Besides, NPCs

appear to be the most susceptible cells of the central nervous system to Zika virus

infection(Rubio-Hernández et al., 2023). Zika virus can directly infect the NPC and

disrupt neural development, resulting in reduced organoid growth and overall size, thereby recapitulating microcephaly observed in foetuses and infants exposed to the virus(Watanabe et al., 2017). In this study, the proportion of NPC and neuron population decreased following *T. pallidum* infection, suggesting that *T. pallidum* inhibited the differentiation of immature cells into NPCs in the brain organoids, resulting in the number of NPC (especially subNPC1B), and subsequently led to a decrease in the number of neurons. Moreover, 40% of children with congenital syphilis have similar mental disorders to those in congenital Zika virus infection(Carrier and Haughton, 2019).

The vertebrate hindbrain contains a complex network of dedicated neural circuits that play an essential role in controlling many physiological processes and behaviours. The hindbrain also contains a wide variety of inter- and relay neurons, including a network of reticulospinal neurons that regulate alertness, sleep, posture, and fine-grained locomotor activities(Krumlauf and Wilkinson, 2021; Valakh et al., 2023). Our scRNA-seq revealed that the number of hindbrain neurons significantly decreased following *T. pallidum* infection, which supports the fact that *T. pallidum* infection causes systemic paresis, progressive dementia, and other nervous system related symptoms. HOX family genes are key factors regulating early hindbrain development and segmentation(Frohman et al., 1990; McNulty et al., 2005). SHOX2 is required for vestibular statoacoustic neuronal development with nerve fibres mostly terminating in the pons and medulla oblongata (hindbrain regions)(Laureano et al., 2022). Immunofluorescence staining of HOXA5 and SHOX2 markers demonstrated that the fluorescence intensity of SHOX2 and HOXA5 decreased following *T. pallidum* infection, thereby revealing that *T. pallidum* inhibited the development and maturation of hindbrain neurons on organoids, which was again confirmed by the scRNA-seq analysis results. Moreover, our scRNA-seq analysis revealed that *T. pallidum* inhibited the hindbrain neuron cell number through the suppression of subNPC1B population in organoids, and found that TCF3 and notch signalling pathway may play important roles in the inhibition of subNPC1B-hindbrain neuron differentiation axis by *T. pallidum*. However, in-depth exploration of the mechanism

of *T. pallidum*-inhibited hindbrain neurogenesis is warranted for further understanding
the pathogenesis and early intervention of congenital neurodevelopmental
impairment.

Nevertheless, certain limitations remain to be addressed and could likely be the
topic of future studies. First, although several recent protocols have made use of
growth factors to promote further neuronal maturation and survival, the organoid
culture scheme needs to be further improved owing to the lower percentage of mature
neurons and the challenge of cell necrosis within the organoids at this stage in day 55
organoids. Second, only changes in the appearance, morphology, and cell type of the
organoids infected with *T. pallidum* were observed in this study, while research on
physiological function was still blank. Third, further experiments are warranted to
confirm that *T. pallidum* restricts the neurodevelopment of organoids by inhibiting the
subNPC1B-hindbrain neuron differentiation axis, which is limited to bioinformatics
analysis. Finally, future animal experiments could allow a more comprehensive
understanding of the mechanisms discovered in organoid studies.

In summary, even though the current organoid systems demonstrated limitations
and require additional optimisation for better use in disease modelling, they have
opened up important novel avenues for related research on congenital
neurodevelopmental impairment. Based on the bioinformatics analysis of scRNA-seq,
we found that *T. pallidum* could inhibit the differentiation of subNPC1B in brain
organoids, thereby reducing the differentiation from subNPC1B to hindbrain neurons,
and ultimately affecting the development and maturation of hindbrain neurons during
pregnancy. The TCF3 and notch signalling pathway could play important roles in this
process, which needs further experimental verification. This is the first report to study
the effects of *T. pallidum* on the neural development of an iPSC-derived brain
organoid model, which could serve as a critical benchmark for studying the
mechanism of congenital neurodevelopmental impairment in the future.

Materials and methods

Preparation of *T. pallidum*

The *T. pallidum* Nichols strain was kindly provided by Lorenzo Giacani, PhD (University of Washington, Seattle, WA, USA) and propagated in male New Zealand rabbits as described previously(Tong et al., 2017). *T. pallidum* was mixed with the organoids maturation stage medium and thereafter used to treat the organoids.

iPSC culture and brain organoids generation.

iPSC was purchased from the American Type Culture Collection, which were derived from fibroblasts that were reprogrammed by the ectopic expression of four specific genes (OCT4, SOX2, KLF4, and MYC)(Yang et al., 2022). Brain organoids were generated from iPSC using established protocols as previously described(Lin et al., 2018; Madhavan et al., 2018), but slightly modified. Briefly, iPSC colonies were cultured in a 6-well plate coated with Matrigel (Corning, 354277) in the mTeSR™ medium (STEMCELL Technologies，85851) until the cells reached 90–100% confluence and dissociated into single cells with Accutase (Gibco, A1110501). Thereafter, the iPSCs were transferred to a 6-well plate with Ultra-Low Attachment surface polystyrene (Corning, REF3471) in StemFlex™ DMEM/F12 Medium (Gibco, A33493) containing with 20% KnockOut Serum (Gibco, 10828028), 1×NEAA (Gibco, 11140-050), 1×GlutaMAX-1 (Gibco, 35050-061), 1×β-Mercaptoethanol (Invitrogen, 21985-023), 5 mM Dorsomorphin (Sigma, P5499), 10 mM SB-431542 (Sigma, S4317), and 1% Penicillin-Streptomycin (Gibco, 15140-0122); 10 μM Rock inhibitor (MCE, Y-27632) was added to the medium after the first 3 days and then the medium was half changed every other day. At day 7, the old medium was removed and replaced with a differentiation medium consisting of Neurobasal-A-Medium (Gibco, 10888-022) containing 1×B27 (Gibco, 12587010), 1×GlutaMAX-1, 20ng/mL FGF2 (R&D Systems, 233-FB), 10 ng/mL EGF (R&D Systems, 236-EG), 1×β-Mercaptoethanol, and 1% Penicillin-Streptomycin. On day 25, the differentiation medium was exchanged with a maturation medium consisting of Neurobasal-A-Medium, 20ng/mL BDNF (R&D Systems, 248-BDB), 20 ng/mL GDNF (R&D Systems, 212-GD), 20 ng/mL NT-3 (R&D Systems, 267-N3), 10 ng/mL EGF, and 1% Penicillin-Streptomycin. The medium was half changed every other day

until 55 days.

Immunohistochemistry

For immunohistochemistry of the brain organoids, organoids were processed at day

55 as previously described(Qian et al., 2016). Briefly, whole organoids were fixed in

4% paraformaldehyde for 45–60 min with slow shaking at 25 ℃. The organoids were

washed three times with phosphate buffer solution (PBS) and then incubated in 30%

sucrose solution overnight at 4 ℃. The organoids were embedded in plastic

cryomolds (Tissue Tek) and filled with O.C.T cryoembedding medium (Sakura, 4583).

The embedded tissue was frozen on dry ice and then stored at –80 ℃ until cryostat

sectioning. Frozen organoid tissue was sliced into 15-μm sections using a microtome

cryostat (MICROM International GmbH, HM550) at −15 to −20 ℃, and collected

using superfrost Ultra Plus slides. For immunostaining, the organoid sections were

quickly washed with PBS to eliminate any residual O.C.T, and then blocked with

blocking permeabilising solution (0.3% Triton-X, 0.2% Tween-20, 0.1% bull serum

albumin and 10% normal goat serum in PBS) for 1 h at room temperature. Thereafter,

sections were incubated overnight at 4 ℃ in an antibody diluent (1.1% Triton-X and 2%

normal goat serum in PBS) containing primary antibodies anti-TUBB3 (mouse,

1:1000, Sigma-Aldrich, T8660; rabbit, 1:300, Abcam, ab18207), anti-SOX2 (rabbit,

1:300, Abcam, ab92494), anti-MAP2 (rabbit, 1:200, Abcam, ab32454), anti-SHOX2

(rabbit, 1:100, Affinit, DF14735), and anti-HOXA5 (mouse, 1:100, Santa cruz,

sc-365784). On the following day, sections were rinsed three times in PBST (PBS

containing 0.05% Tween 20) before incubation for 1 h at room temperature in

secondary antibody solution, which contained Alexa Fluor 488-conjugated anti-mouse

antibody (goat, 1:1000, Thermo Fisher, A11001) and Alexa Fluor 647-conjugated

anti-rabbit antibody (goat, 1:1000, Thermo Fisher, A-21244). Finally, sections were

washed with PBST and stained with VECTASHIELD® Antifade Mounting Medium

with DAPI (Vector laboratories, H1200). Stained organoid cryosections were imaged

using a confocal laser scanning microscope (Zeiss, LSM 700).

For immunohistochemistry of neurons, following culture on Millicell glass

coverslips, neurons obtained by flow sorting were washed with PBS three times and

fixed with 4% paraformaldehyde for 15 min. After fixation, the neurons were washed

with PBS and permeabilised with blocking solution containing 10% normal goat

serum, 2% bull serum albumin, and 0.05% Triton X-100 in PBS for 1 h at room

temperature. Then neurons were incubated with anti-MAP2 (rabbit, 1:200, Abcam,

ab32454) overnight at 4 °C and Alexa Fluor 647-conjugated anti-rabbit antibody (goat,

1:1000, Thermo Fisher, A-21244) for 1 h at room temperature, and washed three

times with PBS before mounting on the glass slide.

Flow cytometric screening and candidate validation

Neuron cells obtained from brain organoids were screened for cell surface molecule

expression using an ultra-high-speed flow cytometry sorter (Beckman, MoFlo Astrios

EQS) as previously reported(Menon et al., 2019). Organoids were dissociated into

single cell, and then resuspended in PBS with 5% KnockOut Serum and stained using

CD71-PE (1:100, Bioscience, 12-0719-42) and CD49b-APC (1:100, Bioscience,

17-0500-42). Antibodies were incubated for 30 min at 4 °C and the cells were

subsequently washed twice in PBS with 5% KnockOut Serum. For flow cytometric

analysis, single cell suspensions were dispensed for sorting and then culture for

subsequent observation and validation. Data analyses were performed using FlowJo™

version 10.7.1 (Becton, Dickinson and Company).

RNA isolation and qRT-PCR

For each *T. pallidum* or control group, the organoids were collected at day 55 in 2 mL

RNAse-free tubes and chilled on ice throughout the procedure. Brain organoids were

homogenised in Trizol (Invitrogen, 1596026) and processed according to the

manufacturer's instructions. cDNA synthesis was performed using a TransScript

All-in-One First-Strand cDNA Synthesis SuperMix for qPCR kit (Invitrogen,

AT341-01) according to the manufacturer protocols. qPCR quantified cDNA via

SYBR$^®$ Premix Ex Taq™ II (Takara, RR82LR). Each reaction was run in triplicate

and analysed following the $\Delta\Delta$Ct value using glyceraldehyde-3-phosphate

dehydrogenase (GAPDH) as a reference gene. Data were presented as expression

level ($2^{-\Delta\Delta Ct}$) relative to GAPDH. The primer pairs are listed in Supplementary Table.

Preparation of single-cell suspensions and scRNA-seq of brain organoids

For preparation of single-cell suspensions, the organoids were harvested and

dissociated into single cells using Accutase. Single cell was resuspended with cold

DMEM/F-12 medium (Gibco, C11330500BT), and counted by Trypan Blue to

determine the live cell concentration. Living cell rate was preferably above 90%, and

then diluted for an appropriate concentration to obtain approximately 6000 cells per

lane of a 10×microfluidic chip device; 30000–50000 cells were needed to generate the

scRNA-seq. Single cell sequencing experiments were conducted using the

10×Genomics Chromium following the manufacturer's instructions by GENE

DENOVO, Guangzhou, China. A detailed description of the scRNA-seq protocol is

available in the experiment flow of the Supplementary Protocol.

scRNA-seq data analysis

10X Genomics Cell Ranger software (version 3.1.0) was employed to convert raw

BCL files to FASTQ files, alignment, and counts quantification. These FASTQ files

were aligned to the human reference genome and transcriptome

(GCF_000001405.38_GRCh38.p12). The cell by gene matrices for each sample were

individually imported to Seurat (v3.1.1)(Butler et al., 2018) for downstream analysis.

Cluster markers were interpreted and assigned cluster identity by using known

literature cell type annotations identifiable details, and the organoids were divided

into 13 cell populations using the tSNE algorithm in the R language. Subclustering

within the NPC and neuron populations was performed by selecting and clustering the

cells from the NPC and neuronal clusters, respectively, and repeating the clustering

procedure with batch effect correction. The resulting distance matrices of NPC and

neuron clusters were used as input for generating UMAP embeddings. After

subclustering, the NPC was identified and subclustered into NPC1 and NPC2; NPC1

were ulteriorly subclustered and generated into subNPC1A, subNPC1B, subNPC1C,

subNPC1D, and subNPC1E. Neuron populations were identified and subclustered to

generate four clusters: hindbrain neuron, choroid plexus, noncerebral neuron, and

cortical neuron.

Differentially expressed genes were identified using the SingleR R

package(Satija et al., 2015) and cut-off of |log fold-change| > 0.25 (Bonferroni

adjusted *p*-value < 0.05). Thereafter, the violin plot of differentially expressed genes

was displayed using the 'ggplot2' R package, and the heatmap of differentially

expressed genes was plotted using the 'heatmap' R package(Saliba et al., 2016).

In order to identify disturbed biological functions and understand the importance

of differentially expressed genes following *T. pallidum* infection on organoids, GO

classification was performed, which included the following categories: biological

process and molecular functions. GO functional enrichment analysis was conducted

for identifying differentially expressed genes using Database for Annotation,

Visualization and Integrated Discovery(Sherman et al., 2022; Zhang et al., 2016) with

default parameters. KEGG pathway enrichment analysis identified significantly

enriched metabolic pathways or signal transduction pathways in the hindbrain neuron

clusters. The distribution of expression of each cluster was demonstrated by using a

dot plot.

To estimate RNA velocity, spliced and unspliced transcripts were enumerated

using the scVelo (v0.2.4)(Bergen et al., 2020). We performed pseudotemporal

trajectory inference analysis on the subNPC1B and hindbrain neurons of human brain

organoids utilising Monocle2 using the DDR-Tree and default parameter.

SCENIC was used to analyse the transcription factors for identifying gene

regulatory networks as previously described(Aibar et al., 2017; Lin et al., 2021a).

Briefly, we predicted the gene regulatory networks and transcription factors using

SCENIC (pySCENIC, v.11.2), which consisted of the following three steps:

establishment of co-expression modules by Random Forest, identification of direct

relationship using motif analysis via RcisTarget, and calculation of the regulon

activity score with Area Under the recovery Curve(Lin et al., 2021b). The TCF3 was

identified from the subNPC1B subsets in the organoids from single-cell transcripts.

Data visualization and statistical analysis

All the statistical analyses and data visualization were performed under the R statistical computing environment (v3.6.3). Heatmaps of gene expression were produced using pheatmap (v1.0.12). Visualisation of the three-dimensional data embeddings were performed using the rgl package (v0.100.54). All other data visualisations were produced using functionalities provided in Seurat and the ggplot2 package (v3.3.0). The Chi-square test was used to compare the percentage differences among different groups. One-way analysis of variance with Tukey's multiple comparisons tests were used for multiple group comparisons. A paired t-test was used to perform the statistical analysis of the differences in the mRNA expression levels and mean fluorescence intensity in Prism program (v5.0). Data are presented as mean $\pm$ standard error of mean. $P < 0.05$ was considered statistically significant.

Acknowledgements

This work was funded by the National Natural Science Foundation of China (grant number 82272370 and 81973104 to T.-C. Y., and grant number 82001600 to Y. L.), Science and Technology Program of Xiamen City, Fujian Province, China (grant number 3502Z20224036 to T.-C. Y.), the Natural Science Foundation of Fujian Province, China (grant number 2021J02055 to T.-C. Y., and grant number 2020J05292 to Y. L., grant number 2021J01073 to M.-L. T.). The funders played no role in the study design, data collection, or analyses, the decision to publish, or manuscript preparation.

Author contributions

Q.-Y. X., Conceptualization, Data curation, Software, Formal analysis, Validation, Investigation, Visualization, Methodology, Writing – original draft, Writing – review and editing; Y.-J. W., Data curation, Software, Methodology; Y. H., Investigation, Visualization, Methodology; X.-Q. Z., Data curation, Software, Methodology; M.-L. T., Resources, Formal analysis, Supervision, Funding acquisition, Validation; Y. L., Conceptualization, Data curation, Software, Formal analysis, Validation, Investigation, Funding acquisition, Visualization; T.-C. Y., Conceptualization, Funding acquisition,

559   Writing – original draft, Project administration, Writing – review and editing

Competing interests

561   The authors declare that no competing interests exist.

Data and materials availability:

The raw scRNA-sequence data reported in this paper have been deposited in the

Genome Sequence Archive (Genomics, Proteomics & Bioinformatics 2021) in

National Genomics Data Center (Nucleic Acids Res 2022), China National Center for

Bioinformation / Beijing Institute of Genomics, Chinese Academy of Sciences

(GSA-Human: HRA004501)   that   are   publicly   accessible   at

https://ngdc.cncb.ac.cn/gsa-human. All data needed to evaluate the conclusions in the

paper are present in the paper and/or the Supplementary Materials.

References

Aibar, S., González-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H.,
Hulselmans, G., Rambow, F., Marine, J.C., Geurts, P., Aerts, J*., et al.* (2017).
SCENIC: single-cell regulatory network inference and clustering. Nat Methods
14, 1083-1086.
Andrews, M.G., and Kriegstein, A.R. (2022). Challenges of Organoid Research.
Annu Rev Neurosci *45*, 23-39.
Auriti, C., Bucci, S., De Rose, D.U., Coltella, L., Santisi, A., Martini, L., Maddaloni,
C., Bersani, I., Lozzi, S., Campi, F*., et al.* (2022). Maternal-Fetal Infections
(Cytomegalovirus, Toxoplasma, Syphilis): Short-Term and Long-Term
Neurodevelopmental Outcomes in Children Infected and Uninfected at Birth.
Pathogens *11*.
Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing
RNA velocity to transient cell states through dynamical modeling. Nat
Biotechnol *38*, 1408-1414.
Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and
species. Nat Biotechnol *36*, 411-420.
Camp, J.G., Badsha, F., Florio, M., Kanton, S., Gerber, T., Wilsch-Bräuninger, M.,
Lewitus, E., Sykes, A., Hevers, W., Lancaster, M*., et al.* (2015). Human cerebral
organoids recapitulate gene expression programs of fetal neocortex development.
Proc Natl Acad Sci U S A *112*, 15672-15677.
Carrier, J., and Haughton, V. (2019). Congenital Syphilis: A Challenging Case for

593 NICU Clinicians. Neonatal Netw *38*, 170-177.

David, M., Hcini, N., Mandelbrot, L., Sibiude, J., and Picone, O. (2022). Fetal and neonatal abnormalities due to congenital syphilis: A literature review. Prenat Diagn *42*, 643-655.

Fortin, O., and Mulkey, S.B. (2023). Neurodevelopmental outcomes in congenital and perinatal infections. Curr Opin Infect Dis *36*, 405-413.

Frohman, M.A., Boyle, M., and Martin, G.R. (1990). Isolation of the mouse Hox-2.9 gene; analysis of embryonic expression suggests that positional information along the anterior-posterior axis is specified by mesoderm. Development *110*, 589-607.

Grosse, S.D., Dollard, S.C., and Ortega-Sanchez, I.R. (2021). Economic assessments of the burden of congenital cytomegalovirus infection and the cost-effectiveness of prevention strategies. Semin Perinatol *45*, 151393.

Hughes, B.W., Addanki, K.C., Sriskanda, A.N., McLean, E., and Bagasra, O. (2016). Infectivity of Immature Neurons to Zika Virus: A Link to Congenital Zika Syndrome. EBioMedicine *10*, 65-70.

Kanton, S., Boyle, M.J., He, Z., Santel, M., Weigert, A., Sanchís-Calleja, F., Guijarro, P., Sidow, L., Fleck, J.S., Han, D*., et al.* (2019). Organoid single-cell genomic atlas uncovers human-specific features of brain development. Nature *574*, 418-422.

Kim, J., Koo, B.K., and Yoon, K.J. (2019). Modeling Host-Virus Interactions in Viral Infectious Diseases Using Stem-Cell-Derived Systems and CRISPR/Cas9 Technology. Viruses *11*.

Krenn, V., Bosone, C., Burkard, T.R., Spanier, J., Kalinke, U., Calistri, A., Salata, C., Rilo Christoff, R., Pestana Garcez, P., Mirazimi, A*., et al.* (2021). Organoid modeling of Zika and herpes simplex virus 1 infections reveals virus-specific responses leading to microcephaly. Cell Stem Cell *28*, 1362-1379.e1367.

Krumlauf, R., and Wilkinson, D.G. (2021). Segmentation and patterning of the vertebrate hindbrain. Development *148*.

Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E., Homfray, T., Penninger, J.M., Jackson, A.P., and Knoblich, J.A. (2013). Cerebral organoids model human brain development and microcephaly. Nature *501*, 373-379.

Laureano, A.S., Flaherty, K., Hinman, A.M., Jadali, A., Nakamura, T., Higashijima, S.I., Sabaawy, H.E., and Kwan, K.Y. (2022). shox2 is required for vestibular statoacoustic neuron development. Biol Open *11*.

Li, C., Xu, D., Ye, Q., Hong, S., Jiang, Y., Liu, X., Zhang, N., Shi, L., Qin, C.F., and Xu, Z. (2016). Zika Virus Disrupts Neural Progenitor Development and Leads to Microcephaly in Mice. Cell Stem Cell *19*, 672.

Li, Q., and Shen, L. (2022). Neuron segmentation using 3D wavelet integrated encoder-decoder network. Bioinformatics *38*, 809-817.

Lin, W., Wang, Y., Chen, Y., Wang, Q., Gu, Z., and Zhu, Y. (2021a). Role of Calcium Signaling Pathway-Related Gene Regulatory Networks in Ischemic Stroke Based on Multiple WGCNA and Single-Cell Analysis. Oxid Med Cell Longev *2021*,

637   8060477.

638 **Lin, W.W.,** Xu, L.T., Chen, Y.S., Go, K., Sun, C., and Zhu, Y.J. (2021b). Single-Cell
639   Transcriptomics-Based Study of Transcriptional Regulatory Features in the
640   Mouse Brain Vasculature. Biomed Res Int *2021*, 7643209.

641 **Lin, Y.T.,** Seo, J., Gao, F., Feldman, H.M., Wen, H.L., Penney, J., Cam, H.P.,
642   Gjoneska, E., Raja, W.K., Cheng, J*., et al.* (2018). APOE4 Causes Widespread
643   Molecular and Cellular Alterations Associated with Alzheimer's Disease
644   Phenotypes in Human iPSC-Derived Brain Cell Types. Neuron *98*,
645   1141-1154.e1147.

646 **Madhavan, M.,** Nevin, Z.S., Shick, H.E., Garrison, E., Clarkson-Paredes, C., Karl,
647   M., Clayton, B.L.L., Factor, D.C., Allan, K.C., Barbar, L*., et al.* (2018).
648   Induction of myelinating oligodendrocytes in human cortical spheroids. Nat
649   Methods *15*, 700-706.

650 **Mattei, P.L.**, Beachkofsky, T.M., Gilson, R.T., and Wisco, O.J. (2012). Syphilis: a
651   reemerging infection. Am Fam Physician *86*, 433-440.

652 **McNulty, C.L.,** Peres, J.N., Bardine, N., van den Akker, W.M., and Durston, A.J.
653   (2005). Knockdown of the complete Hox paralogous group 1 leads to dramatic
654   hindbrain and neural crest defects. Development *132*, 2861-2871.

655 **Medoro, A.K.,** and Sánchez, P.J. (2021). Syphilis in Neonates and Infants. Clin
656   Perinatol *48*, 293-309.

657 **Menon, V.,** Thomas, R., Elgueta, C., Horl, M., Osborn, T., Hallett, P.J., Bartos, M.,
658   Isacson, O., and Pruszak, J. (2019). Comprehensive Cell Surface Antigen
659   Analysis Identifies Transferrin Receptor Protein-1 (CD71) as a Negative
660   Selection Marker for Human Neuronal Cells. Stem Cells *37*, 1293-1306.

661 **Mercurio, S.,** Serra, L., and Nicolis, S.K. (2019). More than just Stem Cells:
662   Functional Roles of the Transcription Factor Sox2 in Differentiated Glia and
663   Neurons. Int J Mol Sci *20*.

664 **Mertens, J.,** Marchetto, M.C., Bardy, C., and Gage, F.H. (2016). Evaluating cell
665   reprogramming, differentiation and conversion technologies in neuroscience. Nat
666   Rev Neurosci *17*, 424-437.

667 **Meyer, U.,** Yee, B.K., and Feldon, J. (2007). The neurodevelopmental impact of
668   prenatal infections at different times of pregnancy: the earlier the worse?
669   Neuroscientist *13*, 241-256.

670 **Nelson, S.B.,** Hempel, C., and Sugino, K. (2006). Probing the transcriptome of
671   neuronal cell types. Curr Opin Neurobiol *16*, 571-576.

672 **Nowakowski, T.J.,** Pollen, A.A., Di Lullo, E., Sandoval-Espinosa, C., Bershteyn, M.,
673   and Kriegstein, A.R. (2016). Expression Analysis Highlights AXL as a Candidate
674   Zika Virus Entry Receptor in Neural Stem Cells. Cell Stem Cell *18*, 591-596.

675 **Pollen, A.A.,** Bhaduri, A., Andrews, M.G., Nowakowski, T.J., Meyerson, O.S.,
676   Mostajo-Radji, M.A., Di Lullo, E., Alvarado, B., Bedolli, M., Dougherty, M.L*.,*
677   *et al.* (2019). Establishing Cerebral Organoids as Models of Human-Specific
678   Brain Evolution. Cell *176*, 743-756.e717.

679 **Qian, X.,** Nguyen, H.N., Song, M.M., Hadiono, C., Ogden, S.C., Hammack, C., Yao,
680   B., Hamersky, G.R., Jacob, F., Zhong, C*., et al.* (2016). Brain-Region-Specific

Organoids Using Mini-bioreactors for Modeling ZIKV Exposure. Cell *165*, 1238-1254.

Ramani, A., Müller, L., Ostermann, P.N., Gabriel, E., Abida-Islam, P., Müller-Schiffmann, A., Mariappan, A., Goureau, O., Gruell, H., Walker, A.*, et al.* (2020). SARS-CoV-2 targets neurons of 3D human brain organoids. Embo j *39*, e106230.

Rubio-Hernández, E.I., Comas-García, M., Coronado-Ipiña, M.A., Colunga-Saucedo, M., González Sánchez, H.M., and Castillo, C.G. (2023). Astrocytes derived from neural progenitor cells are susceptible to Zika virus infection. PLoS One *18*, e0283429.

Saliba, A.E., Li, L., Westermann, A.J., Appenzeller, S., Stapels, D.A., Schulte, L.N., Helaine, S., and Vogel, J. (2016). Single-cell RNA-seq ties macrophage polarization to growth rate of intracellular Salmonella. Nat Microbiol *2*, 16206.

Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial reconstruction of single-cell gene expression data. Nat Biotechnol *33*, 495-502.

Setoh, Y.X., Amarilla, A.A., Peng, N.Y.G., Griffiths, R.E., Carrera, J., Freney, M.E., Nakayama, E., Ogawa, S., Watterson, D., Modhiran, N.*, et al.* (2019). Determinants of Zika virus host tropism uncovered by deep mutational scanning. Nat Microbiol *4*, 876-887.

Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T., and Chang, W. (2022). DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). Nucleic Acids Res *50*, W216-w221.

Sugino, K., Clark, E., Schulmann, A., Shima, Y., Wang, L., Hunt, D.L., Hooks, B.M., Tränkner, D., Chandrashekar, J., Picard, S.*, et al.* (2019). Mapping the transcriptional diversity of genetically and anatomically defined cell populations in the mouse brain. Elife *8*.

Tantalo, L.C., Lukehart, S.A., and Marra, C.M. (2005). Treponema pallidum strain-specific differences in neuroinvasion and clinical phenotype in a rabbit model. J Infect Dis *191*, 75-80.

Tong, M.L., Zhao, Q., Liu, L.L., Zhu, X.Z., Gao, K., Zhang, H.L., Lin, L.R., Niu, J.J., Ji, Z.L., and Yang, T.C. (2017). Whole genome sequence of the Treponema pallidum subsp. pallidum strain Amoy: An Asian isolate highly similar to SS14. PLoS One *12*, e0182768.

Tonni, G., Grisolia, G., Pisello, M., Zampriolo, P., Fasolato, V., Sindico, P., Araújo Junior, E., and Bonasoni, M.P. (2022). Congenital Syphilis Presenting with Brain Abnormalities at Neuroscan: A Case Report and a Brief Literature Review. Microorganisms *10*.

Trujillo, C.A., and Muotri, A.R. (2018). Brain Organoids and the Study of Neurodevelopment. Trends Mol Med *24*, 982-990.

Valakh, V., Wise, D., Zhu, X.A., Sha, M., Fok, J., Van Hooser, S.D., Schectman, R., Cepeda, I., Kirk, R., O'Toole, S.M.*, et al.* (2023). A transcriptional constraint mechanism limits the homeostatic response to activity deprivation in mammalian neocortex. Elife *12*.

Velasco, S., Kedaigle, A.J., Simmons, S.K., Nash, A., Rocha, M., Quadrato, G.,
Paulsen, B., Nguyen, L., Adiconis, X., Regev, A.*, et al.* (2019). Individual brain
organoids reproducibly form cell diversity of the human cerebral cortex. Nature
570, 523-527.
Verghese, V.P., Hendson, L., Singh, A., Guenette, T., Gratrix, J., and Robinson, J.L.
(2018). Early Childhood Neurodevelopmental Outcomes in Infants Exposed to
Infectious Syphilis In Utero. Pediatr Infect Dis J *37*, 576-579.
Watanabe, M., Buth, J.E., Vishlaghi, N., de la Torre-Ubieta, L., Taxidis, J., Khakh,
B.S., Coppola, G., Pearson, C.A., Yamauchi, K., Gong, D.*, et al.* (2017).
Self-Organized Cerebral Organoids with Human-Specific Features Predict
Effective Drugs to Combat Zika Virus Infection. Cell Rep *21*, 517-532.
Wicher, V., and Wicher, K. (2001). Pathogenesis of maternal-fetal syphilis revisited.
Clin Infect Dis *33*, 354-363.
Woods, C.R. (2009). Congenital syphilis-persisting pestilence. Pediatr Infect Dis J *28*,
536-537.
Yang, S.G., Wang, X.W., Qian, C., and Zhou, F.Q. (2022). Reprogramming neurons
for regeneration: The fountain of youth. Prog Neurobiol *214*, 102284.
Zhang, T., Lin, Y., Liu, J., Zhang, Z.G., Fu, W., Guo, L.Y., Pan, L., Kong, X., Zhang,
M.K., Lu, Y.H.*, et al.* (2016). Rbm24 Regulates Alternative Splicing Switch in
Embryonic Stem Cell Cardiac Lineage Differentiation. Stem Cells *34*,
1776-1789.
Zhang, W., Tan, Y.W., Yam, W.K., Tu, H., Qiu, L., Tan, E.K., Chu, J.J.H., and Zeng,
L. (2019). In utero infection of Zika virus leads to abnormal central nervous
system development in mice. Sci Rep *9*, 7298.

Figures

Figure 1. *T. pallidum* infection inhibited brain organoid neurodevelopment. (**A**) Schematic view of the methods for generating brain organoid from iPSCs. (**B**) Representative bright field images of individual brain organoids treated with *T. pallidum* over time. (**C**) Changes of the transcription levels of genes related to the three germ layers following *T. pallidum* infection in the brain organoids. (**D**) Immunofluorescent assay of the effect of *T. pallidum* infection on the brain organoid. A paired *t*-test was used for evaluating the statistical differences between the two groups. (*: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$, NS: not significant).

Figure 2. scRNA-seq revealed that *T. pallidum* infection inhibited the differentiation of neural progenitor cell subcluster 1B in the brain organoid. (**A**) Schematic overview of the 10×Genomics scRNA-seq procedure and analysis. (**B**) tSNE plot of single cells identified a total of 19 clusters. (**C**) tSNE plot of single cell depicting the separation into the 13 cell populations. (**D**) Distinguishing the tSNE plot of cell types and

proportion analysis of 13 cell populations within the two groups. (**E**) Expression profile of key genes related to NPC1 and NPC2 cell populations between the two groups. Powder blue: control group; Pink: *T. pallidum* group; Pale green: NPC1; Yellow: NPC2; Red indicates high gene expression; Green indicates low gene expression. (**F**) Volcano plot of differential genes (*T. pallidum* vs control) in the NPC1 cluster. (**G**) GO analysis of the upregulated genes (left) and downregulated genes (right) in the NPC1 cluster following *T. pallidum* infection. (**H**) UMAP plot of distinguished NPC1 subclusters (left) and proportion analysis of each subcluster within two groups (right). Chi-square test was used to compare the changes in the proportion of the five small subgroups among the different groups. (**I**) Specificity marker expression of NPC1 subcluster.

Figure 3. *T. pallidum* infection inhibited the differentiation of neurones in the brain organoid. (**A**) Volcano Plots of differentially expressed genes related to neurons between the two groups. (**B**) Representative GO terms of downregulated and upregulated enrichment genes related to the neurons between two groups. (**C**)Violin diagram of genes related to early neuronal development. (**D**) The effect of *T. pallidum* infection on the transcription level of genes related to the nervous system in the brain organoids. (**E**) The frozen section of brain organoids by MAP2 fluorescence staining. (**F**) Flow sorting of brain organoids and statistical analysis of the two groups using a paired *t*-test. (**G**) Cell morphology and validation of CD71⁻/CD49b⁻ neuron populations. A paired *t*-test was used for evaluating the statistical differences between the two groups. (*: $P < 0.05$, **: $P < 0.01$, ***：$P < 0.001$, NS: not significant).

Figure 4. scRNA-seq revealed that *T. pallidum* infection inhibited the differentiation of the hindbrain neurons in the brain organoid. (**A**) UMAP embedding plots comparing the distribution of the brain organoids within the two groups. (**B**) UMAP embedding of cell subset in each region of the neurons calibrated by marker genes. (**C**) Subcluster cell number as a percentage of the neuron cluster cell number. The Chi-square test was used to compare the differences in the percentages among different cell subsets. (**D**) Violin diagram of the relevant marker genes in each neuronal region. (**E**) Immunohistochemistry of the brain organoids validating the expression of hindbrain neuron markers, including HOXA5 and SHOX2. A paired *t*-test was used to compare the differences between the two groups (*: $P < 0.05$, **: $P < 0.01$, NS: not significant).

Figure 5. scRNA-seq revealed the mechanism of *T. pallidum* affecting neurodevelopment in the brain organoids. (**A**) Left panel: temporal analysis of subNPC1B-hindbrain neuron by Monocle 2, with subNPC1B subgroup in blue and hindbrain neuron subgroup in magenta. Right: HES4 localises to subNPC1B and NEUROGD1 localises to the hindbrain neuron. (**B**) Heatmap of differentially expressed transcripts along pseudotime. Cells are ordered by cell type and pseudotime. (**C**) Bubble difference diagram of subNPC1B and hindbrain neuron marker genes before and after *T. pallidum* infection. (**D**) GO analysis of the total number of

810 differential genes in the subNPC1B subgroup. (**E**) The differential expression of

811 transcription factors between the *T. pallidum* and control group by SCENIC analysis.

812 (**F**) Extended analysis of TCF3. (**G**) KEGG enrichment pathway analysis of hindbrain

813 neuron subgroup.