Repeated horizontal acquisition of lagriamide-producing symbionts in Lagriinae beetles

Short title: Repeated beetle symbiont acquisition

Siddharth Uppal*[1], Samantha C. Waterworth*[1,2], Alina Nick (anick@ice.mpg.de)[3], Heiko Vogel (hvogel@ice.mpg.de)[3], Laura V. Flórez[4], Martin Kaltenpoth[3†], Jason C. Kwan[1‡]

*These authors contributed equally to this work

†To whom correspondence should be addressed. Address: Hans-Knöll-Straße 8, 07745, Jena, Germany. Email: kaltenpoth@ice.mpg.de

‡To whom correspondence should be addressed. Address: 777 Highland Ave., Madison, WI 53705. Email: jason.kwan@wisc.edu

[1]Division of Pharmaceutical Sciences, School of Pharmacy, University of Wisconsin-Madison, Madison, USA.

[2]Current address: National Cancer Institute, Frederick, Maryland, USA

[3]Max Planck Institute for Chemical Ecology, Jena, Germany

[4]Department of Plant and Environmental Science, University of Copenhagen, Copenhagen, Denmark

Study funding information: NSF DBI-1845890 (S.U., S.C.W., J.C.K.), Max Planck Society (A.N., H.V., M.K.), ERC CoG 819585 "SYMBeetle" (M.K.).

Abstract

Microbial symbionts associate with multicellular organisms on a continuum from facultative associations to mutual codependency. In some of the oldest intracellular symbioses there is exclusive vertical symbiont transmission, and co-diversification of symbiotic partners over millions of years. Such symbionts often undergo genome reduction due to low effective population sizes, frequent population bottlenecks, and reduced purifying selection. Here, we describe multiple independent acquisition events of closely related defensive symbionts followed by genome erosion in a group of Lagriinae beetles. Previous work in *Lagria villosa* revealed the dominant genome-eroded symbiont of the genus *Burkholderia* produces the antifungal compound lagriamide and protects the beetle's eggs and larvae from antagonistic fungi. Here, we use metagenomics to assemble 11 additional genomes of lagriamide-producing symbionts from seven different host species within Lagriinae from five countries, to unravel the evolutionary history of this symbiotic relationship. In each host species, we detected one dominant genome-eroded *Burkholderia* symbiont encoding the lagriamide biosynthetic gene cluster (BGC). Surprisingly, however, we did not find evidence for host-symbiont co-diversification, or for a monophyly of the lagriamide-producing symbionts. Instead, our analyses support at least four independent acquisition events of lagriamide-encoding symbionts and subsequent genome erosion in each of these lineages. By contrast, a clade of plant-associated relatives retained large genomes but secondarily lost the lagriamide BGC. In conclusion, our results reveal a dynamic evolutionary history with multiple independent symbiont acquisitions characterized by high degree of specificity. They highlight the importance of the specialized metabolite lagriamide for the establishment and maintenance of this defensive symbiosis.

Keywords: lagriamide, *Burkholderia*, symbiosis, symbiont replacement, biosynthetic gene cluster, metagenomics, Lagriinae, chemical defence, secondary metabolism

Introduction

Eukaryotes have been associated with prokaryotic microbes since the initial endosymbiotic events that led to the acquisition of mitochondria and chloroplasts [1]. These organelles represent the presumed endpoint of ancient symbioses with α-proteobacteria and cyanobacteria, respectively, that over time led to a progressive shrinkage of the symbionts' genomes and eventual transfer of genes from symbionts to host [1]. Although organelle acquisition appears to be a rare event [2], other more recent symbioses appear to be on a similar evolutionary trajectory of profound genome reduction and absolute dependence on host cells. For example, the acquisition of the intracellular symbiont *Buchnera aphidicola* in the common ancestor of aphids allowed them to diversify as sap-feeding insects since the symbiont synthesises several essential amino acids not found in plant sap, as evidenced by a rapid basal radiation of aphid species [3] and strict co-evolution of aphids and *Buchnera* [4]. *B. aphidicola* has been vertically transmitted for at least 200 million years [4] and has a profoundly reduced chromosome, about 11% of the size of *Escherichia coli* [5]. Through comparison of various symbionts, a model of genome reduction has emerged whereby host-restriction initially weakens purifying selection on formerly essential genes, through both host-provided metabolites and symbiont population structure, with low effective population sizes and isolation within individual hosts [6]. When symbionts are vertically transmitted, population bottlenecks occur during every transmission event, causing the fixation of deleterious mutations within the population [6]. These factors combine to first cause an increase in pseudogenes in the genome [6] and then deletion of those pseudogenes due to a known deletion-bias within bacteria [7]. The most reduced genomes lose even central functions such as DNA repair pathways [6], which leads to an increased rate of evolution and further gene loss, as well as increased AT-bias in many cases [8, 9]. In the cases of symbionts living inside host cells, it is likely that this process is exacerbated due to a lack of opportunity or ability to horizontally acquire functional genes.

However, genome reduction is also known to occur without genetic isolation. For instance, free-
living bacteria living in nutrient-poor environments such as *Prochlorococcus* spp. are thought to
have reduced genomes as a consequence of selection pressure to streamline their metabolism
[10], potentially explained through the Black Queen hypothesis [11], which posits that selection
drives pathways to be lost when they are produced by another species in the ecosystem as
"public goods". There are also genome-reduced symbionts which seemingly are not genetically
isolated. *Burkholderia* symbionts that reside extracellularly in leaf nodules in plants are mainly
transmitted vertically because the symbiosis is mutually co-dependent [12], although horizontal
transfer may have occurred rarely between plants, the soil microbiota, and insects [13]. This
suggests a lack of genetic isolation, and indeed there is evidence of repeated horizontal
transfers of biosynthetic genes for defensive molecules among leaf nodule symbionts of
Rubiacaeae plants [13]. Such systems may provide an opportunity to study the evolutionary
pressures that lead to the process of genome reduction, and the mechanisms of symbiosis that
underlie it.

The dichotomy of vertical versus horizontal transfer of symbionts may be one determinant of
genome reduction. A relatively clear-cut example is the two symbionts of the tunicate
Lissoclinum patella, the extracellular cyanobacterium *Prochloron didemni* [14] and the
intracellular "*Candidatus* Endolissoclinum faulkneri" [15]. The former is capable of horizontal
transmission, which is reflected in its almost clonal genome amongst very divergent hosts and a
lack of genome reduction [14], while the latter is vertically transmitted, as evidenced by its co-
divergence with its hosts across cryptic speciations, and profound genome reduction [15, 16].
However, the mode of transmission also exists on a continuum from strict vertical to strict
horizontal, with mixtures of vertical and horizontal transmission in between [17]. For instance,
the Tsetse fly symbiont *Sodalis glossinidius* shows some signs of genome-reduction such as
rampant pseudogenes, but remains culturable in the lab, meaning that horizontal transmission

cannot be excluded [18]. Likewise, symbionts long thought to be exclusively vertically

transmitted, such as the bryozoan symbiont "*Ca.* Endobugula sertula", which is packaged with

the hosts' larvae, show no signs of genome reduction [19], indicating that there is no compelling

reason why it should not be able to transmit horizontally between hosts. Indeed, "*Ca.* E. sertula"

has been found in genetically divergent but proximal bryozoan individuals, suggesting horizontal

transmission [20].

The *Lagria* and *Ecnolagria* beetles belong to subfamily Lagriinae within the family

Tenebrionidae (order Coleoptera). The *Lagria villosa* beetle, a known soybean pest [21], is a

source of lagriamide, an antifungal polyketide [21]. The compound is produced via a *trans*-AT

polyketide synthase (PKS)-non-ribosomal peptide synthetase (NRPS) hybrid biosynthetic gene

cluster (BGC), termed *lga,* which due to a nucleotide signature (kmer frequency) distinct from

the chromosome is predicted to have been horizontally acquired [21]. The *lga* BGC is encoded

by a *Burkholderia* symbiont (*Burkholderia* sp. LvStB) that is present in glandular structures

associated with the ovipositor of female beetles and secreted on its eggs as they are laid [21].

The symbiont has been shown to have a defensive role against fungi in the egg [21] and larval

stages [22]. Previously, we showed that the genome of *Burkholderia* sp. LvStB is reduced and

has lost several essential genes including some genes involved in the DNA repair pathways and

primary metabolism [23]. The genome has a low coding density, and a high number of

pseudogenes and transposases, indicative of a reduced genome [23]. These characteristics are

consistent with host restriction and vertical transmission of LvStB. However, there is evidence

that *Burkholderia* symbionts from *L. villosa* can be transferred to plant tissues and survive for

several days and that bacteria can be acquired by the beetle from the plant and soil

environment [24].

As some Lagriinae beetles harbor symbionts in special structures that likely evolved between 55

and 82 million years ago based on fossil evidence, positioned to deposit symbionts on the eggs

[25, 26], we hypothesized that lagriamide-producing *Burkholderia* symbionts might have co-

evolved with their hosts in a manner similar to other vertically-transmitted insect symbionts.

However, the possibility for transmission of the symbionts to and from plants, and the

accessibility of the symbionts' habitat on the surface of eggs and within adult females suggested

that horizontal symbiont acquisition may be possible. As the beetles harbor complex

microbiomes with multiple related *Burkholderia* strains as well as other bacteria [22, 24], both

genome-reduced and not, an alternative hypothesis is that the lagriamide BGC has been

repeatedly horizontally transferred among environmental strains and symbionts. Moreover,

partnerships in defensive symbionts are usually more dynamic as compared to intracellular

nutritional symbionts [27]. It is also possible that the lagriamide-producing strain is restricted to

L. villosa, and that different Lagriinae species have symbionts with different BGCs, as this would

allow the association to react much more flexibly to changes in antagonist communities. To

clarify this evolutionary picture, we analyzed the metagenomes of 12 beetle samples, spanning

seven species belonging to the genera *Lagria* and *Ecnolagria* across five different countries

(four continents) (**Table 1**). We recovered the metagenome-assembled genomes (MAGs) of

several different *Burkholderia* bacteria and confirmed the presence of the lagriamide BGC in

each specimen. We also report a complete genome of the genome-reduced, lagriamide-

producing *Burkholderia* sp. LvStB symbiont, obtained through long-read Nanopore sequencing.

We compared the phylogeny of the recovered *Burkholderia* MAGs, the lagriamide BGCs, and

the host beetles to determine whether co-cladogenesis occurred in this system, and to further

explore the evolutionary relationships in the symbiosis. The results indicate that the lagriamide

BGC was likely only acquired once in the common ancestor of beetle-associated *Burkholderia*

symbionts, and subsequently lost in the majority of the descendent free-living strains.

Remarkably, as all the lagriamide-bearing symbionts are genome-reduced but do not form a

monophyletic clade and do not correspond to host phylogeny, they likely represent multiple

acquisition events, followed by independent genome-reduction processes. The common factor

of lagriamide production might be one of the reasons for selection by and dependency on hosts.

This would suggest that a single group of natural products caused several independent

symbioses to be established over evolutionary time.

Results and Discussion

Beetle phylogeny

We sequenced and assembled the metagenomes of the collected *Lagria* and *Ecnolagria* beetle

populations (**Table 1**), and beetle mitogenomes (see **Table S1** for mitogenome statistics) were

extracted and annotated to infer host beetle phylogeny (**Fig. 1**). In line with previous studies,

mitogenomes belonging to the tenebrionid subfamilies Lagriinae, Blaptinae, Pimeliinae,

Stenochiinae, and Alleculinae were found to be monophyletic whereas Diaperinae and

Tenebrioninae were found to be para- or polyphyletic [28–31]. Maximum likelihood analysis

using RAxML [32] (**Fig. SI 1**) and Bayesian analysis using MrBayes [33] (**Fig. 1**) gave similar

results.

All collected *Lagria* beetle mitogenomes clustered into four distinct subclades: All *L. hirta* beetle

mitogenomes were clustered in a single clade, alongside a closely related clade of *Lagria*

species (*L. rufipennis* and *L. okinawana*) from Japan. The *L. atripes* and *L. grenieri* beetles

formed another clade more distantly related to the *L. hirta* and Japanese *Lagria* species. Finally,

the *L. villosa* and *Ecnolagria* sp. beetles formed a fourth clade along with *Chrysolagria* sp.

(JX412760), distinct from the other *Lagria* beetles. A small distinction was noted here, wherein

the Bayesian phylogeny (**Fig. 1**) suggested that the *Cerogria* beetles belonged to the clade with

L. villosa and *Ecnolagria* species, whereas the Maximum-likelihood phylogeny showed the

Cerogria to be in a clade with all other *Lagria* beetles (**Fig. SI 1**). However*,* in both cases the

branch support values are too low to make any definite conclusions. Publicly available

sequences of *L. hirta* (OX375806) clustered with collected *L. hirta* samples from Rhineland

Palatinate, Germany (LhHG), and *L. rufipennis* (MW802588) clustered with the two *L. rufipennis*

(Lruf1 and Lruf2) samples.

Recovery of Lagriamide BGCs

A complete, or mostly complete, *lga* BGC was found, using antiSMASH v7 [34, 35], in eleven of

the twelve samples, with the exception of *L. rufipennis 1* where only small fragments of the *lga*

BGC could be recovered. The BGC recovered from *L. rufipennis 2* was found over two contigs

and could not be manually joined following inspection of the assembly graph. The missing data

for this region spans from approximately halfway through the *lgaB* gene to approximately

halfway through the *lgaC* gene (**Fig. 2A**).

Analysis of representative BGCs revealed two differences in gene organization of the *lga* BGC

across the different Lagriinae beetle species (**Fig. 2A**). The first difference observed regarded

the *lgaC* gene. The *lgaC* gene from the BGCs recovered from the *L. hirta* G and *L. hirta* SB

samples appeared to be split into two, denoted *lgaC1* and *lgaC2* for clarity. Alignment of the

lgaC gene from the three *L. hirta* samples revealed that there was a perfect alignment of the

nucleotide sequences save for a 37-bp deletion in the BGCs from *L. hirta* G and *L. hirta* SB,

which introduced a frameshift (**Fig. SI 2A**). This frameshift consequently introduced a premature

stop codon which split the *lgaC* gene into two ORFs in *L. hirta* G and *L. hirta* SB (**Fig. SI 2B**).

The second difference we observed was in the *lga* BGC from the LvStB genome extracted from

the 2023 *L. villosa* metagenome (Lv23). One split in *lgaB* and two splits in *lgaC* were seen.

However, the assembly of the *L. villosa* 2023 metagenome was based solely on long-read data,

which is error-prone [36], and the splits may not be a true reflection of the BGC in this sample.

Normally Sanger sequencing would be the solution to validate these questionable regions but

unfortunately, there was no remaining DNA after the long read sequencing runs for this

particular sample. For this reason, we left the BGC with the splits but were cautious not to over-

interpret the apparent breaks in the genes in this BGC.

We then considered the domain organisation within the *lga* BGC genes (**Fig. 2B**). The domain

organisation is largely congruent across the *lga* BGCs recovered from the metagenomes. We

did note, however, an additional annotated "DHt" domain in *lgaB*, which is defined as

"Dehydratase domain variant more commonly found in *trans*-AT PKS clusters", in the *lga* BGCs

from all *L. hirta* samples and the *L. rufipennis 2* sample. Similarly, we detected an additional

carrier protein domain (phosphopantetheine acyl carrier protein group) near the N-terminus of

the *lgaC* protein in the BGCs from the *L. grenieri* and *L. okinawana* samples. In all cases, close

inspection of the primary sequence of these additional dehydratase and carrier domains

revealed mutations in the sequences that would likely render the encoded domain non-

functional (see Supplementary Material for full details).

Finally, as with the originally described *lga* BGC recovered from the *L. villosa* 2019 sample [21],

we found mutations in the catalytic or conserved motifs of *lgaG* DH2, *lgaG* KS6, *lgaB* KR3 and

lgaC KS5 domains, that we believe may render these domains inactive (see Supplementary

Material for full details). As a result, the domain architecture of all representative *lga* BGCs from

all samples appear functionally identical.

Together, the conservation of the *lga* BGC in at least seven different species of Lagriinae

beetles, across four geographically distant countries, implies that the production of lagriamide is

an important factor for the host beetle and that the *lga* BGC is under strong selective pressure.

The presence of additional domains in the *lga* BGC in several samples, even though they are

likely inactive, is intriguing as it suggests that these domains may have previously been present

in all *lga* BGCs but may have decayed over time and were lost. The reason as to why these

domains were selected against would be speculative at best and inspection of all lagriamide-like

compounds produced in the different beetle populations would need to be assessed to truly infer

differences that the domain architecture may have on the resulting chemistry. Conserved

production of other bioactive compounds has been observed, such as pederin, across

Staphylinidae beetle species (*Paederus* and *Paederidus* genera) [37], which are host to a

Pseudomonas symbiont that produces pederin [38].

The two systems have several parallels: both pederin and lagriamide are produced by a *trans*-

AT PKS NRPS hybrid BGC in *Pseudomonas* bacterium [37], where the compound is

concentrated in the female oviposition organs, coated onto the eggs and serves to protect

juveniles [39]. Further, both pederin and lagriamide are the sole insect-associated compounds

in suites of compounds otherwise associated with marine invertebrates. Groups of pederin

analogs, such as the onnamides, mycalamides, psymberins, and theopederins have been

isolated from a variety of marine sponges [40–44] and ascidians [45], while bistramide, the most

structurally similar compound to lagriamide, was isolated from an ascidian [46]. The question

remains, however, as to what the fundamental link is between these terrestrial and marine

systems.

Complete genome of the *lga*-carrying LvStB symbiont

Long-read sequencing of the *L. villosa* 2023 (Lv23) metagenome allowed us to assemble a

complete genome of a *lga*-carrying *Burkholderia* strain (referred to as LvStB_2023 from

hereon). LvStB_2023 was found to have a 2.5 Mbp long genome with a GC percentage of

58.63%. It has 2 circular chromosomes - chromosome 1 is 1.89 Mbp, chromosome 2 is 0.55

Mpb in size, and there is a plasmid 59.77 kbp long. The genome is estimated to be 97.1%

complete (98.8% with "specific" mode) and 0.02% contaminated as per CheckM2 [47] and thus

is a high-quality MAG according to the MIMAG standards [48]. Assembly graph analysis of

LvStB_2023 verified that we have the complete sequence of two circular chromosomes and a

plasmid. However, the CheckM2 estimate did not reflect a fully complete genome, at 98.8%,

and we believe that this small discrepancy in predicted completeness may be a result of

ongoing genome reduction [49].

LvStB_2023 has a coding density of 78% and 59.1% with and without pseudogenes

respectively. A large percentage (43.87%) of the ORFs in LvStB_2023 were identified as

pseudogenes (1613 out of 3676), the highest of any *lga*-carrying *Burkholderia* symbiont.

However, this estimate may be artificially high as pseudogenes are identified purely based on

their length relative to their closest BLASTP match and these counts are derived from an

assembly generated from only long-read data which can be prone to errors [50–52], particularly

homopolymeric runs. However, coding density and frequency of pseudogenes is not very

different from LvStB MAGs assembled from short-read data (see **Table S2** for complete

genome characteristics of recovered MAGs). Having multiple chromosomes is a common

phenomenon in *Burkholderia* [53, 54]. Generally in multi-chromosome bacteria, the majority of

the genes for essential functions are located on one larger or primary chromosome, while the

smaller or secondary chromosome has much fewer essential genes and mostly carries genes

for niche specific functions [55]. In the case of LvStB_2023, chromosome 1 appears to be the

primary chromosome as it is much larger in size, and has 77 out of 84 core genes (including

multiple copies) (**Fig 3A**). Functional analysis revealed chromosome 1 to have the highest

number of genes for all essential COG categories (**Fig 3B**), including categories L (replication,

recombination and repair), J (Translation, ribosomal structure and biogenesis), M (Cell

wall/membrane/envelope biogenesis) and H (Coenzyme transport and metabolism).

The *lga* BGC is on chromosome 2 (0.55 Mbp long) and can be distinguished by the continuous

block of coding sequences on the reverse strand in **Fig. 3A**. A detailed view of the *lga*-carrying

chromosome can be seen in **Fig. SI 3,** where the GC content of the *lga* BGC is significantly

higher than the rest of the genome, providing further evidence for its horizontal acquisition.

Chromosome 1, chromosome 2, and the plasmid have 44.75%, 37.59%, and 42.34% of their

coding capacity taken up by pseudogenes, respectively. The similar abundance of pseudogenes

in each of the contigs indicates that the whole genome is undergoing reduction simultaneously.

The chromosome with *lga* (chromosome 2) has the smallest percentage of pseudogenes, which

may be a reflection of the required conservation of the *lga* BGC in combination with the

presence of large genes in *lga*.

Diversity of beetle-associated *Burkholderia* symbionts

Recovery and analysis of metagenome-assembled genomes

Following assembly, the 12 beetle metagenomes were binned, and the resultant bins were

manually refined. A total of 77 MAGs were recovered from all samples, of which 24 MAGs were

of high quality, 30 of medium quality, and 23 of low quality (**Table S2**) in accordance with

published MIMAG standards [48]. Only medium and high-quality MAGs were used for

downstream analysis, with the exception of one low-quality bin carrying the *lga* BGC (LhHG_2).

Confirmation that the *lga* BGC had been assigned to the correct MAG has been covered in

detail in the Supplementary Material. Genome erosion, such as that already observed for the

lga-carrying symbiont *Burkholderia* sp. LvStB [23], can skew the completeness metric. To

determine if a lower quality MAG was incomplete or genome-reduced, we also considered

several other metrics, including core gene presence, number of pseudogenes [23], and coding

density (**Table S2**), and concluded that this particular MAG (LhHG_2) was likely both reduced

and incomplete.

For each beetle population, a single MAG belonging to the genus *Burkholderia* with a single

copy of the *lga* BGC was identified. Previous studies on the lagriamide-carrying symbiont strain

B. gladioli LvStB [21, 23, 56], showed that this strain was significantly more abundant than all

other bacteria associated with *L. villosa*, and had a reduced genome. Consistent with this, all

newly recovered MAGs that included *lga* BGCs were the most abundant MAGs in each sample,

had reduced genomes with an abundance of pseudogenes and transposases, and had lower

coding densities relative to other *B. gladioli* genomes (**Table S2**). In standing with previous

studies of *Lagria* beetles, where both reduced and non-reduced *B. gladioli* genomes were

recovered, additional *B. gladioli* MAGs (Latri_2, LhHG_3, and LhSB_5) were recovered that did

not carry the lagriamide BGC and showed no evidence of genome erosion. We also recovered

three small *B. gladioli* MAGs (Lgren_7, Lv19_6_18, Lv20_2) and one small *Burkholderia* MAG

(Lv19_6_14), as well as MAGs classified as *B. lata* (Lv19_4_0) and *B. arboris* (Lv20_1).

Average nucleotide identity (ANI) analysis of *B. gladioli* MAGs carrying the *lga* BGC showed that

MAGs from different beetle species and/or different locations were likely different bacterial

species due to shared ANI values less than 95% [57]. However, previous studies have

suggested that ANI alone is not a sufficient metric for species delineation and that the aligned

fraction (AF) must also be taken into account [57–60]. Following recent cutoffs adopted for

species delineation [58], we opted to use AF ≥ 60% along with ANI ≥ 95% as a cutoff for

species assignment. Subsequently, we found that the *Burkholderia* MAGs carrying the

lagriamide BGC appeared to be split into at least five novel species (**Table S3**).

Phylogenetic analysis of recovered metagenome-assembled genomes

In order to elucidate the evolutionary history of the association between Lagriinae beetles and *Burkholderia* symbionts, we reconstructed phylogenies of the *Burkholderia* symbionts and free-living relatives based on shared single-copy genes. A priori, we hypothesized that the *lga*-encoding, genome-eroded symbionts would form a monophyletic clade Showing co-diversification with the hosts, given that such patterns have been previously described across many ancient and co-evolved symbioses.

The phylogeny of the beetle-associated *Burkholderia* symbionts, relative to other *Burkholderia* species, was inferred using 126 single-copy hierarchical orthogroups (HOGs) (non-pseudogenes) present in more than 90% of the genomes (RAxML phylogeny is represented in **Fig. 4** and **Fig. SI 4** while **Fig. SI 5** represents the Bayesian approach). *Burkholderia* symbionts without the *lga* BGC were broadly present across the phylogeny containing *B. gladioli, B. lata,* and *B. arboris* strains. By contrast, and consistent with our expectation, symbionts of different host species carrying the *lga* BGC were closely related. Surprisingly, however, these genome-eroded, *lga*-encoding symbionts did not form a monophyletic clade. Because the tree indicates that the common ancestor of the genome-reduced *lga*-encoding symbionts also gave rise to a lineage of non genome-reduced descendents, this result indicates a non-reduced free living common ancestor and subsequent multiple independent acquisition events by Lagriinae beetles. To test for the robustness of our phylogenetic analysis, we repeated the analysis using single-copy HOGs present in 95%, 80%, 70%, and 60% (**Fig. SI 6**) of the genomes, as well as after removing any putative horizontally transferred genes (**Fig. SI 7 and 8**). Other than minor discrepancies in the terminal nodes, we obtained highly similar phylogenetic trees, supporting the lack of monophyly of the *lga* BGC carrying *Burkholderia* symbionts. Thus, all our analyses

support a phylogeny that contains a clade of mostly free-living (plus some beetle-associated

symbionts with non-eroded genomes) *Burkholderia* that groups within the *lga* BGC-containing

Lagriinae symbionts. Concerning the evolutionary history of the symbiosis, this leaves us with

two alternative scenarios: (i) an ancestral association of the whole clade of bacteria with beetles

and a certain degree of genome erosion on the deep branches, and a subsequent reversal to a

free-living stage of the clade containing many plant-associated *B. gladioli* strains; or (ii) at least

four independent transitions from a free-living (or plant-associated) to a symbiotic lifestyle, each

of which followed by genome erosion.

To unravel which of these scenarios is more likely, we analyzed shared HOGs between different

Burkholderia sp., after removing any pseudogenes from MAGs. We observed substantially

higher conservation of orthogroups between the potentially free-living *Burkholderia* sp. than

among the *lga*-containing symbionts, with the free-living strains sharing a large core genome

(**Fig. 4**). If the shared ancestor of all *lga*-encoding symbionts and the free-living strains would

have been tightly associated with beetles and experienced some degree of genome erosion

(scenario i), this observation would postulate a substantial increase in the genome size of the

bacteria after the reversal to the free-living/plant-associated lifestyle and before the clade split

into the different taxa. While theoretically possible, this scenario seems highly unlikely, because

acquisition of a large number of genes would have to have happened quickly and early in order

for extant strains in this clade to have such a degree of gene overlap. Instead, it appears much

more plausible that there were at least four independent transitions to a symbiotic lifestyle, each

of which was followed by genome erosion. This is consistent with the observation that the

genomes of the eroded strains retain distinct sets of genes, many of which represent subsets of

the free-living strains' core genomes (**Fig. SI 9**), since gene loss from independant host-

restriction events would be expected to be largely stochastic. This distinct set of genes can

however, also be due to symbiont replacement events followed by genome reduction.

Furthermore, the lack of synteny observed in the genes flanking the *lga* BGC (**Fig. SI 10**) is

indicative of either genomic rearrangement or independent acquisition of the *lga* BGC. Both of

these further support the independent acquisitions of symbionts followed by genome erosion. It

is important to note that this conclusion is based on the current data and may change as we

obtain more samples and long-read metagenomes that allow for synteny analyses across the

entire genome.

Consistent with the scenario of multiple independent transitions to a symbiotic lifestyle, the

phylogeny of the *lga* BGC-carrying *Burkholderia* symbionts was found to be largely incongruent

with the beetle phylogeny (**Fig 5A**), except for the symbionts grouping together for individuals of

the same host species, i.e. *L. hirta* and *L. rufipennis*, respectively. The incongruence between

host and symbiont phylogenies suggests both multiple symbiont acquisition and possibly host

switching events that lead to symbiont replacements. Symbiont replacement has often been

reported in nutritional symbionts as a way for the hosts to replace a genetically degraded

symbiont with a more complete and effective one and to acquire new adaptations for expanding

into different niches [61]. *Burkholderia* symbionts related to *B. gladioli* in *Lagria* beetles have

been reported to evolve from plant-associated bacteria [26] capable of transfer from beetles to

plants with subsequent survival [24]. It is possible that the horizontal acquisition might occur in

the egg and larval stages, where the symbionts are localized on the surface (eggs) or in

cuticular invaginations (larvae and pupae) that remain connected to the external surface via a

small duct [62]. As the closely related *Burkholderia* strain LvStA can be acquired horizontally

from the environment [24], and there is evidence of free-living bacteria carrying lagriamide-like

BGCs [63], we propose that there are *lga*-carrying *Burkholderia* strains persevering in the

environment (e.g. in plants or soil) [24] that can be horizontally acquired by the beetle host.

As we previously observed that *lga* has distinct nucleotide composition to the Lv19 genome [21],

suggestive of a recent horizontal transfer, we sought to determine if it has been independently

transferred to the corresponding symbiont in different beetle hosts. Phylogenetic analysis of the

representative *lga* BGCs from all samples resulted in two possible topologies using GTRCAT-V

and GTRGAMMAI models (**Fig. 5B**). Both topologies included conserved clades. However, the

relative positions of the three clades are poorly supported (**Fig. 5B**, highlighted in blue),

resulting in the two alternative topologies. A Bayesian tree was also constructed (**Fig. SI 11**)

which is congruent with the GTRGAMMAI tree topology. The inconsistent topology likely stems

from limited resolution of the phylogeny affecting deep nodes in the trees. The GTRCAT-V

topology is perfectly congruent with the symbiont phylogeny based on genome-wide marker

genes, while the GTRGAMMAI topology shows one discrepancy at one of the deep nodes.

Thus, these analyses do not provide evidence for additional horizontal transfer events of the *lga*

cluster, so it is likely that there was a single acquisition of *lga* in the common ancestor of the

symbiont and *B. gladioli* clade, with subsequent loss in the free-living group. It appears that

lagriamide production was highly selected for in symbiotic settings and hence retained, whereas

it was lost in the most of the larger genomes (assumed to be free-living) where it was not

selected for. These findings indicate that the *lga* BGC is important in the symbiosis, either for

symbiont establishment (e.g. competition with other symbionts) and/or because lagriamide is an

effective host-defensive molecule. Furthemore, the fact that highly similar *Burkholderia* species

with *lga* were identified across different Lagriinae beetles indicates that symbiont acquisition is

highly selective.

Interestingly, lagriamide seems to be highly conserved, despite the dynamics of the system,

where multiple species of bacteria associate with each beetle host, and several *lga*-producing

Burkholderia have apparently been independently acquired. A dynamic association in defensive

symbioses has been previously hypothesized, to allow for rapid adaptation to a changing

community of antagonists, or to individual co-evolving pathogens [27]. We expected to see

changes in the defensive chemistry used in a symbiotic context, akin to the rapid evolution of

immune genes in animals [64–66]. However, despite the dynamic nature of many defensive

symbioses, with symbiont replacements on ecological or evolutionary timescales, several

examples of defensive symbioses highlight that the same defensive compounds can be used

over long evolutionary timescales. In case of beewolf wasps, *Streptomyces* symbionts have

been found to produce piericidin and streptochlorin for an estimated 68 million years [67, 68].

Both compounds are found in different beewolf species and across different geographic

locations. Similarly, as discussed above, pederin is produced across different species of

Paederus and *Paederidus* beetles by *Pseudomonas* symbionts [37, 38]. Similarly, we now

describe the production of lagriamide by a *Burkholderia* symbiont across several species of

Lagria and *Ecnolagria* beetles. Thus, even though these defensive symbioses are dynamic in

the acquisition and replacement of microbial partners, the chemistry seems to be conserved.

This suggests a limited diversity of chemical compounds that can be used for defense against

eukaryotic antagonists (predators or fungi) in a symbiotic context, which is supported by the

convergence on similar compounds in terrestrial and aquatic symbioses. It is possible that this

might be due to the harmful side effects of the bioactive molecule on the eukaryotic host,

analogous to the cytotoxic side-effects of antifungal pharmaceuticals on humans, resulting in

only limited diversity of such compounds.

To gain insights on the possible origin of the *lga* BGC, we performed an analysis of

pentanucleotide (5-mer) frequencies of the beetle-associated, *lga*-carrying symbionts and their

associated BGCs, along with the genomes of recently identified soil-borne *Paraburkholderia*

species that carry the lagriamide B (*lgb)* BGC, which is highly similar to the *lga* BGC [63].

Visualisation of 5-mer frequencies of the BGCs and the genomes revealed three clusters of

BGCs: The BGCs from the two soil-borne *Paraburkholderia* strains, the BGCs from the Brazilian

L. villosa-derived LvStB strains, and then a third cluster of all other *lga* BGCs (**Fig. SI 12**). A

similar pattern was observed for the nucleotide composition of the respective genomes wherein

LvStB and Lv20_9 form an isolated cluster, the two soil-borne *Paraburkholderia* form a second,

distant cluster, and all other *lga*-carrying *Burkholderia* strains and cultured *Lagria*-associated

genomes (LvStA and LhStG) form a third cluster. None of the BGCs share similar 5-mer

composition with their respective genomes, providing additional evidence for the horizontal

acquisition of the *lga* BGC.

We noted during the analysis of the COG annotated genes in LvStB_2023 that there appeared

to be a particularly high number of genes in the L category (replication, recombination and

repair) that were likely pseudogenes (**Fig 3B**). We assessed the percentage change of COG

annotated genes in all *lga*-carrying *Burkholderia* and found that this pseudogenization of genes

involved in DNA replication, recombination and repair was particularly high in all the Brazilian *L.*

villosa-derived LvStB strains, as well as the MAGs LhSB_1, LhG_1, Loki_2 and Lgren_6 (**Fig.**

SI 13). Two of the three LvStB strains also exhibited high pseudogenization of the genes

associated with cell motility (Category N). While COG annotation of genes does not provide a

robust picture, as not all genes are successfully annotated, the increased pseudogenization of

genes involved in DNA replication and repair may explain the divergence of the LvStB strains

observed in both the phylogenetic analysis and the related 5-mer analysis. In particular, LvStB

MAGs possessed highly truncated and psuedogenized *polA* genes, coding for DNA polymerase

I used in many DNA-repair pathways and chromosome replication [69], whereas other *lga*-

containing MAGs, except LhHG_2, had intact *polA* genes (**Fig. SI 14**). The loss of *polA* in the *L.*

villosa symbionts explains their accelerated sequence evolution in the genome as a whole and

also in the *lga* BGC compared to other *lga*-possessing symbionts (**Fig. SI 12**). The absence of

polA in LhHG_2 could be due to its poor quality, as it is only 46% complete and has only 47.6%

percentage of core genes.

Previous studies have highlighted how symbionts can be conserved across host-speciation
events and millions of years, leading to genome reduction in the symbiont [6, 16]. A
disadvantage of such an exclusive relationship is that the symbiont inevitably suffers from
increasing genome erosion that can result in reduced efficiency in providing benefits to the host
[70]. Consequently, many long-term obligate symbioses have experienced symbiont
replacement events that can provide an escape route for the insect host after its symbiont
entered the irreversible phase of degenerative genome reduction [71]. Such replacement is a
common phenomenon in Hemipteran symbionts [61]. In the present study, however, we are
suggesting that the repeated replacement of symbionts may have happened with very closely
related strains that carry the same biosynthetic gene cluster and hence likely provide the same
functional benefit to the host. One reason we are suggesting multiple acquisitions and
displacements may have happened is that all the *lga*-containing symbionts appear to be at
different stages of genome reduction, with very different genome sizes and gene complement,
perhaps indicating that they have been symbionts for different amounts of time. That in
combination with the apparent importance of *lga* specifically, the incongruence of symbiont and
host phylogeny, and the fact that none of the symbionts is profoundly genome-reduced,
suggests that although Lagriinae likely hosted *lga*-containing symbionts since the evolution of
special symbiont storage structures, the current symbionts are not direct descendents. The
replaced symbionts were likely genome-reduced to an extent that they were outcompeted by
incoming *lga*-bearing strains from the environment. The *lga* BGC-containing *Burkholderia*
strains were consistently the most abundant symbionts in the metagenomes across seven
different Lagriinae species, indicating that the *lga* BGC or an as yet unknown genomic feature
shared among the symbiont strains provides a key selective advantage in the beetles' symbiotic
organs. Possibly, lagriamide is uniquely suited to defend the symbionts' niche against
competitors and/or protect its host from antagonists. However, as lagriamide shows lower
antifungal activity than some secondary metabolites of related *Burkholderia* strains [21, 26, 72],

another intriguing possibility is that it only provides a moderate degree of defence but at the

same time exhibits less harmful side effects on the host than other antifungal compounds.

Further elucidating the relevance of lagriamide in establishing the symbiotic association with

beetles will not only provide valuable insights into the ecological and evolutionary dynamics of

defensive symbioses, but may also unravel the mechanisms ensuring specificity in symbiotic

alliances.

491 Methods

492 For additional details see Supplementary Material.

493 Insect collection

Specimens were collected between 2009 and 2019 in Spain, Germany, Brazil, Japan and

Australia in the locations listed in **Table S4**. Female adults were dissected either directly after

chilling for ca. 15 min at −20°C or preserved in 70% ethanol or acetone until dissected. The

accessory glands were removed and preserved in 70% ethanol at −80°C until further

processing. For species in which we suspected the presence of symbiont-harbouring

compartments within the ovipositor in addition to the glands, the ovipositor was also dissected

and preserved along with the accessory glands.

501 DNA isolation and metagenomic sequencing

Genomic DNA from the preserved organs was extracted per individual after removing the

fixative and homogenizing the tissue in liquid nitrogen. The MasterPure[TM] complete DNA and

RNA isolation Kit (Epicentre Technologies) was used as indicated by the manufacturer,

including an additional incubation step at 37°C for 30 min with 4 µL lysozyme (100 mg mL[-1])

before protein precipitation. The nucleic acids were re-suspended in Low TE buffer (1:10 dilution

of TE) and pooled by species. Metagenomic sequencing was carried out in two batches. The

first batch included the samples corresponding to *L. atripes*, *L. grenieri*, *L. hirta* G, *L. hirta* SB, *L.*

hirta HG, and *L. villosa* 2020. This first batch was sent for DNA library preparation using a

Nextera XT DNA Library Prep Kit (Illumina) and metagenomic sequencing on a NovaSeq 6000

platform, using a paired-end approach (2 x 150 bp) to a depth of 30 M reads (9 Gbp) by CeGaT

GmbH (Tübingen, Germany). Samples from the second batch including *L. rufipennis 1 and 2*, *L.*

okinawana, and *Ecnolagria sp.*, were sequenced using Illumina NextSeq 2000 (paired end 2 x

150bp) to a depth of 28 to 44 million reads at the Max Planck Genome Centre (Cologne,

Germany). The data from sample Lv19 corresponds to that described in Florez *et al.* [21].

Taxonomic assignment of individual specimens of *L. rufipennis* was first done morphologically

according to Jung and Kim [73] very similar to the sympatrically occurring females of *Lagria*

nigricollis, and the specimens used for sample Lruf2 were originally identified as *L. nigricollis*

(see also [26]). Due to the uncertainty associated with morphological identification, we therefore

additionally barcoded the specimens after their metagenomes had been sequenced (see

Supplementary information for detailed methods) and compared their cytochrome oxidase I

sequences to those of male specimens of *L. rufipennis* and *L. nigricollis* that can be more easily

distinguished based on their morphology. All 19 *L. rufipennis* and 10 *L. nigricollis* COI

sequences that we obtained turned out to be very similar and formed a sister group to the *L.*

rufipennis sequence available in NCBI (MW802588). However, the *L. nigricollis* sequences

formed a distinct subclade, with the exception of the sample that had been used for

metagenomics (Lruf2), which grouped within *L. rufipennis*. Hence, we reassigned Lruf2 to *L.*

rufipennis, resulting in two replicate metagenomes for this species. Unfortunately, the *L.*

nigricollis samples were males (which do not contain symbionts), preventing us from

sequencing a metagenome of this species.

For the *L. hirta* HG population, genomic DNA was extracted from a pool of 6 egg clutches (20-
30 eggs per clutch) using the Qiagen Genomic-tip 20/G Kit (Qiagen) following the instructions
from the manufacturer. The sample was split in half for DNA extraction and the resulting
genomic DNA was again combined. This sample and an aliquot of the *L. villosa* 2020 genomic
DNA sample underwent End-DNA repair and library preparation using the NEBNext Ultra II DNA
Library Prep Kit (New England Biolabs, Ipswich, USA) and the Ligation Sequencing Kit V14
(SQK-LSK114; Oxford Nanopore Technologies, Oxford, UK) followed by a clean-up step with
AMPure XP beads (Beckman Coulter). These were sequenced on a MinION platform (Oxford
Nanopore Technologies) as described below for long-read sequencing of the *L. villosa* 2023
sample.

Metagenomic sequence assembly and binning

Sample *L. villosa* 2019 represented data previously assembled and analyzed [21, 23].
Sequence data generated from *L. atripes*, *L. grenieri*, *L. hirta* G, *L. hirta* SB, *L. rufipennis 1*, *L.*
rufipennis 2, *L. okinawana*, and *Ecnolagria sp.*, consisted of only short-read Illumina sequence
data. Sequences were trimmed using Trimmomatic v0.39 [74] using TruSeq3-PE as reference,
and sequences shorter than 25 bp being discarded. The trimmed sequences were assembled using
SPAdes v.3.14.1 [75] and binned using Autometa [76]. Sequence data generated from *L. villosa*
2020 and *L. hirta* HG consisted of both short-read Illumina sequence data and long-read
Nanopore sequence data, as mentioned above. After trimming, reads were assembled with
SPAdes v.3.14.1 as a hybrid assembly with the nanopore flag enabled. Assembled contigs were
binned using Autometa [76]. The quality of all MAGs was assessed using CheckM2 v1.0.1 [47]
and each MAG was classified using GTDB-Tk v.2.3.2 against database release 214. Coverage
reported by SPAdes for each contig was used to calculate the MAG coverage, except for
LvStB_2023 where coverage was calculated by read aligned using minimap2 [77].

Long-read sequencing and assembly of the *L. villosa* 2023 metagenome

The symbiotic organs connected to the reproductive system of six female adults of *L. villosa* were dissected and gently homogenized to release bacterial symbionts. The residual host tissue was separated from the bacterial suspension, and both samples were frozen at -20°C. Later, both samples were thawed and centrifuged for 2 minutes at 3,000 rpm + 2 minutes at 5,000 rpm to pellet the tissue and bacteria, respectively. The supernatant was removed, and 20µL sterile 1x PBS was added to both samples. Genomic DNA was extracted using the Nanobind CBB Big DNA kit (Circulomics, Baltimore, USA) followed by enrichment for HMW (high molecular weight) DNA using the Short Read Eliminator kit XS (Circulomics). Isolated HMW DNA purity and concentrations were measured using Qubit (Thermo Fisher). End-DNA repair and library preparation was performed using the NEBNext Ultra II DNA Library Prep Kit (New England Biolabs, Ipswich, USA) and the Ligation Sequencing Kit V14 (SQK-LSK114; Oxford Nanopore Technologies, Oxford, UK) followed by a clean-up step with AMPure XP beads (Beckman Coulter). Sequencing was performed on a MinION platform (Oxford Nanopore Technologies) and MinION flow cells (vR10.4.1) with 100 ng of the library during a 72 h run.

After sequencing, super-high-accuracy base calling of the raw reads was performed with Guppy v6.3.8 (Oxford Nanopore Technologies) (dna_r10.4.1_400bps_sup.cfg model; split-read function enabled), resulting in a total of 9 Gb sequence data. The resulting reads were *de novo* assembled using Flye v2.9.1 [78, 79] with setting minimum overlap as 10kb and with the "--meta" option, followed by four rounds of polishing with Racon v1.3.3 (Vaser et al., 2017) starting from the Flye assembly with option (-m 8 -x -6 -g -8 -w 500). After each polishing round, reads were re-aligned to the resulting assembly with minimap2 v2.17 [77]. A final round of polishing was performed using Medaka v1.2.0 (https://github.com/nanoporetech/medaka) with the r941_min_high_g344 model using the MinION raw reads. After polishing, haplotype

redundancies and overlaps in the assembly based on read depth were purged using

Purge_Dups v1.2.6 [80]. The relative contig coverage, GC content and contig taxonomic

classification were scanned after each genome assembly using Blobtools and TaxonAnalysis to

enable the identification of potential microbial symbiont contigs. We subsequently performed

several rounds of Flye assemblies, using only subsets (e.g. 25%) of the complete MinION data

and/or read length size-cutoffs (5kb) to optimize symbiont genome assembly.

586   Phylogeny of beetles

Mitochondrial genomes (mitogenomes) were recovered from the Eukaryote kingdom bins from

each respective sample. Mitogenomes used by Wei *et al.* [28] in their phylogenetic tree were

selected for references and outgroups. Mitogenomes from all metagenomic datasets and

reference mitogenomes were annotated using the MITOS2 webserver [81] against the RefSe89

Metazoan database, using genetic code 5 (invertebrate mitochondrial). Amino acid sequences

of the 13 protein coding genes (PCGs) from each mitogenome were collected and aligned using

muscle v5.1 [82]. Nucleic acid sequences of the corresponding PCGs were aligned using

pal2nal v14 [83] using the -codontable 5 flag. Nucleic acid alignments were concatenated and a

partition file was generated using the pxcat command from the phyx package [84]. Phylogenetic

analysis was performed by partitioning each codon position for each gene. AICC model

predicted by ModelTest-NG v0.1.7 [85, 86] was used to construct the phylogenetic tree using

RAxML-NG v1.2.1 [87] with the parameters --all and --tree pars{25},rand{25}. Alignment file in

FASTA format was converted to nexus format using Geneious Prime 2023.2.1

(www.geneious.com). Bayesian analysis was performed by partitioning each codon position for

each genes using MrBayes v3.2 [33] with seed and swapseed equal to 42 and using the

following parameters lset applyto=(all) nst=6 rates=invgamma; and unlink statefreq=(all)

revmat=(all) shape=(all) pinvar=(all); using 10 million generations and sample frequency of 500.

The final average standard deviation of split frequencies (ASDSF) was 0.0042.

Burkholderia symbiont phylogeny

Prokka [88] was used to annotate the ORFs of the genomes/ MAGs. Pseudogenes were

removed from the MAGs and orthofinder v2.5.5 [89] was run on amino acid sequences of the

genomes/ MAGs. A custom script was used to extract the genes with single-copy HOGs that are

present in more than 95% (23 HOGs), 90% (126 HOGs), 80% (336 HOGs), 70% (656 HOGs)

and 60% (888 HOGs) of the genomes/ MAGs. Muscle v5.1 [82] was used to align the amino

acid sequences of the selected HOGs, followed by pal2nal v14 [83] to align the corresponding

nucleic acid sequences using -codontable 11. Subsequent steps were similar to those

performed while constructing beetle phylogeny. Bayesian analysis was performed using

MrBayes v3.2 [33] following the steps and parameters described in beetle mitogenome tree

construction. The final ASDSF was 0.0002.

Amino acid sequences of MAGs were blasted (diamond blastP) [90, 91] (with parameters -k 1 --

max-hsps 1 --outfmt 6 qseqid stitle pident evalue qlen slen) against a local copy of the NCBI nr

database, where previously identified *Burkholderia* sequences [21, 23] were removed. Genes

where the top blastP hits had percent identity less than 50% or those without "Burkholderia" in

the subject sequence title of the top hit were classified as putative horizontally transferred

genes. These genes along with any pseudogenes were removed from MAGs and orthofinder

was used to detect HOGs present in more than 95% (16 HOGs), 90% (98 HOGs), 80% (304

HOGs), 70% (632 HOGs) and 60% (884 HOGs) of the genomes/ MAGs. Subsequent steps

were similar to those mentioned in the above paragraph.

Lagriamide BGC phylogeny

Lagriamide BGC genes from *lgaA* to *lgaI* were extracted. Protein sequences were aligned with muscle v5.1 [82] followed by alignment of DNA sequences using pal2nal v14 [83] using -codontable 11. Pxcat command in the phyx package [84] was used to concatenate the DNA alignments and generate a partition file. Maximum likelihood tree was made using RAxML v8.2.12 (raxmlHPC-PTHREADS-SSE3) [32], with the parameters -f a -# 1000 -p 1989 -x 1989. For the GTRGAMMAI model each gene was partitioned for each codon position, while when using GTRCAT -V model partitioning was only performed per gene as it resulted in higher bootstrap values than partitioning for each codon position in each gene. Bayesian analysis was performed as described in the beetle phylogeny with the final ASDSF being 0.0001.

Data availability

The data associated with this study was deposited under BioProject accession no. PRJNA1054523. Metagenomic reads have been deposited in the Sequence Read Archive with accessions SRR27332963–SRR27332975. Representative *lga* BGC sequences have been submitted to Genbank with accession numbers PP034267–PP034279. All lagriamide BGC-carrying MAGs were deposited with the following accession numbers: Ecno_1, JAYFRU000000000; Latri_1, JAYFRV000000000; Lgren_6, JAYFRW000000000; LhG_1, JAYFRX000000000; LhHG_2, JAYFRY000000000; LhSB_1, JAYFRZ000000000; Lruf2_2, JAYFSA000000000; Loki_2, JAYFSB000000000; Lruf1_1, JAYFSC000000000; Lv20_9, JAYFSD000000000.

Acknowledgments

We would like to thank Jean Keller for providing python scripts that helped in creating the symbiont phylogeny. We are grateful to Rebekka Janke, Takema Fukatsu, Kiyoshi Ando, Kimio Masumoto, Juan José López and Rolf Beutel for support with insect collection and to Domenica Schnabelrauch for carrying out PCRs, cloning and Sanger sequencing.

Author Contributions

SU: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing.

SCW: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing.

AN: Data curation, Formal Analysis, Investigation, Methodology

HV: Data curation, Formal Analysis, Investigation, Methodology.

LVF: Conceptualization, Investigation, Writing – review & editing, Supervision, Funding acquisition.

MK: Formal Analysis, Methodology, Conceptualization, Investigation, Writing – review & editing, Supervision, Funding acquisition.

JCK: Conceptualization, Investigation, Writing – review & editing, Supervision, Funding acquisition

Funding

This work was supported by the U.S. National Science Foundation [DBI-1845890 to J.C.K., S.U., and S.C.W.], the Max Planck Society (to A.N., H.V., and M.K.), and the European

666   Research Council through an ERC Consolidator Grant (ERC CoG 819585 "SYMBeetle", to
667   M.K.).

Competing Interests

669   The Kwan lab offers their metagenomic binning pipeline Autometa on the paid bioinformatics
670   and computational platform BatchX in addition to distributing it through open source channels.

References

1.   Dyall SD, Brown MT, Johnson PJ. Ancient Invasions: From Endosymbionts to Organelles.
Science 2004; **304**: 253–257.
2.   Stephens TG, Gabr A, Calatrava V, Grossman AR, Bhattacharya D. Why is Primary
Endosymbiosis so Rare? *New Phytol* 2021; **231**: 1693–1699.
3.   Von Dohlen CD, Moran NA. Molecular Data Support a Rapid Radiation of Aphids in the
Cretaceous and Multiple Origins of Host Alternation. *Biol J Linn Soc Lond* 2000; **71**: 689–
717.
4.   Nováková E, Hypša V, Klein J, Foottit RG, von Dohlen CD, Moran NA. Reconstructing the
Phylogeny of Aphids (Hemiptera: Aphididae) Using DNA of the Obligate Symbiont
Buchnera aphidicola. *Mol Phylogenet Evol* 2013; **68**: 42–54.
5.   Chong RA, Park H, Moran NA. Genome Evolution of the Obligate Endosymbiont *Buchnera*
aphidicola. *Mol Biol Evol* 2019; **36**: 1481–1489.
6.   McCutcheon JP, Moran NA. Extreme Genome Reduction in Symbiotic Bacteria. *Nat Rev*
Microbiol 2011; **10**: 13–26.
7.   Mira A, Ochman H, Moran NA. Deletional Bias and the Evolution of Bacterial Genomes.
Trends Genet 2001; **17**: 589–596.

8.   Hershberg R, Petrov DA. Evidence that Mutation is Universally Biased Towards AT in

Bacteria. *PLoS Genet* 2010; **6**: e1001115.

9.   Dietel A-K, Merker H, Kaltenpoth M, Kost C. Selective Advantages Favour High Genomic

AT-Contents in Intracellular Elements. *PLoS Genet* 2019; **15**: e1007778.

10. Dufresne A, Garczarek L, Partensky F. Accelerated Evolution Associated with Genome

Reduction in a Free-Living Prokaryote. *Genome Biol* 2005; **6**: R14.

11. Morris JJ, Lenski RE, Zinser ER. The Black Queen Hypothesis: Evolution of Dependencies

Through Adaptive Gene Loss. *MBio* 2012; **3**: e00036–12.

12. Pinto-Carbó M, Gademann K, Eberl L, Carlier A. Leaf Nodule Symbiosis: Function and

Transmission of Obligate Bacterial Endophytes. *Curr Opin Plant Biol* 2018; **44**: 23–31.

13. Pinto-Carbó M, Sieber S, Dessein S, Wicker T, Verstraete B, Gademann K, et al. Evidence

of Horizontal Gene Transfer Between Obligate Leaf Nodule Symbionts. *ISME J* 2016; **10**:

2092–2105.

14. Donia MS, Fricke WF, Partensky F, Cox J, Elshahawi SI, White JR, et al. Complex

Microbiome Underlying Secondary and Primary Metabolism in the Tunicate-*Prochloron*

Symbiosis. *Proc Natl Acad Sci U S A* 2011; **108**: E1423–32.

15. Kwan JC, Donia MS, Han AW, Hirose E, Haygood MG, Schmidt EW. Genome Streamlining

and Chemical Defense in a Coral Reef Symbiosis. *Proc Natl Acad Sci U S A* 2012; **109**:

20655–20660.

16. Kwan JC, Schmidt EW. Bacterial Endosymbiosis in a Chordate Host: Long-Term Co-

Evolution and Conservation of Secondary Metabolism. *PLoS One* 2013; **8**: e80822.

17. Ebert D. The Epidemiology and Evolution of Symbionts with Mixed-Mode Transmission.

Annu Rev Ecol Evol Syst 2013; **44**: 623–643.

18. Toh H, Weiss BL, Perkin SAH, Yamashita A, Oshima K, Hattori M, et al. Massive Genome

Erosion and Functional Adaptations Provide Insights Into the Symbiotic Lifestyle of *Sodalis*

glossinidius in the Tsetse Host. *Genome Res* 2006; **16**: 149–156.

19. Miller IJ, Vanee N, Fong SS, Lim-Fong GE, Kwan JC. Lack of Overt Genome Reduction in

the Bryostatin-Producing Bryozoan Symbiont, 'Candidatus Endobugula sertula'. Appl

Environ Microbiol 2016; **82**: 6573–6583.

20. Linneman J, Paulus D, Lim-Fong G, Lopanik NB. Latitudinal Variation of a Defensive

Symbiosis in the Bugula neritina (Bryozoa) Sibling Species Complex. PLoS One 2014; **9**:

e108783.

21. Flórez LV, Scherlach K, Miller IJ, Rodrigues A, Kwan JC, Hertweck C, et al. An Antifungal

Polyketide Associated With Horizontally Acquired Genes Supports Symbiont-Mediated

Defense in Lagria villosa Beetles. Nat Commun 2018; **9**: 2478.

22. Janke RS, Kaftan F, Niehs SP, Scherlach K, Rodrigues A, Svatoš A, et al. Bacterial

Ectosymbionts in Cuticular Organs Chemically Protect a Beetle During Molting Stages.

ISME J 2022; **16**: 2691–2701.

23. Waterworth SC, Flórez LV, Rees ER, Hertweck C, Kaltenpoth M, Kwan JC. Horizontal

Gene Transfer to a Defensive Symbiont With a Reduced Genome in a Multipartite Beetle

Microbiome. MBio 2020; **11**: e02430–19.

24. Wierz JC, Gaube P, Klebsch D, Kaltenpoth M, Flórez LV. Transmission of Bacterial

Symbionts With and Without Genome Erosion Between a Beetle Host and the Plant

Environment. Front Microbiol 2021; **12**: 715601.

25. Jürgen Stammer H. Die Symbiose Der Lagriiden (Coleoptera). Z Morph u Okol Tiere 1929;

15: 1–34.

26. Flórez LV, Scherlach K, Gaube P, Ross C, Sitte E, Hermes C, et al. Antibiotic-Producing

Symbionts Dynamically Transition Between Plant Pathogenicity and Insect-Defensive

Mutualism. Nat Commun 2017; **8**: 15172.

27. Flórez LV, Biedermann PHW, Engl T, Kaltenpoth M. Defensive Symbioses of Animals With

Prokaryotic and Eukaryotic Microorganisms. Nat Prod Rep 2015; **32**: 904–936.

28. Wei Z, Shi A. The Complete Mitochondrial Genomes of Four Lagriine Species (Coleoptera,

Tenebrionidae) and Phylogenetic Relationships Within Tenebrionidae. *PeerJ* 2023; **11**:

e15483.

29. Wu C, Zhou Y, Tian T, Li T-J, Chen B. First Report of Complete Mitochondrial Genome in

the Subfamily Alleculinae and Mitochondrial Genome-Based Phylogenetics in

Tenebrionidae (Coleoptera: Tenebrionoidea). *Insect Sci* 2022; **29**: 1226–1238.

30. Ragionieri L, Zúñiga-Reinoso Á, Bläser M, Predel R. Phylogenomics of Darkling Beetles

(Coleoptera: Tenebrionidae) From the Atacama Desert. *PeerJ* 2023; **11**: e14848.

31. Kergoat GJ, Soldati L, Clamens A-L, Jourdan H, Jabbour-Zahab R, Genson G, et al. Higher

Level Molecular Phylogeny of Darkling Beetles (Coleoptera: Tenebrionidae). *Syst Entomol*

2014; **39**: 486–499.

32. Stamatakis A. RAxML Version 8: A Tool for Phylogenetic Analysis and Post-analysis of

Large Phylogenies. *Bioinformatics* 2014; **30**: 1312–1313.

33. Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, et al. MrBayes

3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model

Space. *Syst Biol* 2012; **61**: 539–542.

34. Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, et al. antiSMASH 5.0: Updates to

the Secondary Metabolite Genome Mining Pipeline. *Nucleic Acids Res* 2019; **47**: W81–

W87.

35. Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, et al. antiSMASH 7.0:

New and Improved Predictions for Detection, Regulation, Chemical Structures and

Visualisation. *Nucleic Acids Res* 2023; **51**: W46–W50.

36. Hu Y, Fang L, Nicholson C, Wang K. Implications of Error-Prone Long-Read Whole-

Genome Shotgun Sequencing on Characterizing Reference Microbiomes. *iScience* 2020;

23: 101223.

37. Piel J, Höfer I, Hui D. Evidence for a Symbiosis Island Involved in Horizontal Acquisition of

Pederin Biosynthetic Capabilities by the Bacterial Symbiont of *Paederus fuscipes* Beetles. *J*

Bacteriol 2004; **186**: 1280–1286.

38. Kador M, Horn MA, Dettner K. Novel Oligonucleotide Probes for *in Situ* Detection of

Pederin-Producing Endosymbionts of *Paederus riparius* Rove Beetles (Coleoptera:

Staphylinidae). *FEMS Microbiol Lett* 2011; **319**: 73–81.

39. Kellner RLL, Dettner K. Differential Efficacy of Toxic Pederin in Deterring Potential

Arthropod Predators of *Paederus* (Coleoptera: Staphylinidae) Offspring. *Oecologia* 1996;

107: 293–300.

40. Witczak ZJ, Bommareddy A, VanWert AL. Pederin, Psymberin and the Structurally Related

Mycalamides: Synthetic Aspects and Biological Activities. In: Kim S-K (ed). *Handbook of*

Anticancer Drugs from Marine Origin. 2015. Springer, Cham, pp 683–700.

41. Sakemi S, Ichiba T, Kohmoto S, Saucy G, Higa T. Isolation and Structure Elucidation of

Onnamide A, a New Bioactive Metabolite of a Marine Sponge, *Theonella* sp. *J Am Chem*

Soc 1988; **110**: 4851–4853.

42. Perry NB, Blunt JW, Munro MHG, Thompson AM. Antiviral and Antitumor Agents From a

New Zealand Sponge, Mycale sp. 2. Structures and Solution Conformations of

Mycalamides A and B. *J Org Chem* 1990; **55**: 223–227.

43. West LM, Northcote PT, Hood KA, Miller JH, Page MJ. Mycalamide D, a New Cytotoxic

Amide From the New Zealand Marine Sponge *Mycale* Species. *J Nat Prod* 2000; **63**: 707–

709.

44. Simpson JS, Garson MJ, Blunt JW, Munro MH, Hooper JN. Mycalamides C and D,

Cytotoxic Compounds From the Marine Sponge *Stylinos* n. Species. *J Nat Prod* 2000; **63**:

704–706.

45. Dyshlovoy SA, Fedorov SN, Kalinovsky AI, Shubina LK, Bokemeyer C, Stonik VA, et al.

Mycalamide A Shows Cytotoxic Properties and Prevents EGF-Induced Neoplastic

Transformation Through Inhibition of Nuclear Factors. *Mar Drugs* 2012; **10**: 1212–1224.

46. Gouiffes D, Juge M, Grimaud N, Welin L, Sauviat MP, Barbin Y, et al. Bistramide A, a New

Toxin From the Urochordata *Lissoclinum bistratum* Sluiter: Isolation and Preliminary

Characterization. *Toxicon* 1988; **26**: 1129–1136.

47. Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. CheckM2: A Rapid, Scalable and

Accurate Tool for Assessing Microbial Genome Quality Using Machine Learning. *Nat*

Methods 2023; **20**: 1203–1212.

48. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et al.

Minimum Information About a Single Amplified Genome (MISAG) and a Metagenome-

Assembled Genome (MIMAG) of Bacteria and Archaea. *Nat Biotechnol* 2017; **35**: 725–731.

49. Manzano-Marín A, Kvist S, Oceguera-Figueroa A. Evolution of an Alternative Genetic Code

in the *Providencia* Symbiont of the Hematophagous Leech *Haementeria acuecueyetzin*.

Genome Biol Evol 2023; **15**: evad164.

50. Tvedte ES, Gasser M, Sparklin BC, Michalski J, Hjelmen CE, Johnston JS, et al.

Comparison of Long-Read Sequencing Technologies in Interrogating Bacteria and Fly

Genomes. *G3: Genes Genomes Genet* 2021; **11**: jkab083.

51. Marx V. Method of the Year: Long-Read Sequencing. *Nat Methods* 2023; **20**: 6–11.

52. Dong X, Du MRM, Gouil Q, Tian L, Jabbari JS, Bowden R, et al. Benchmarking Long-Read

RNA-Sequencing Analysis Tools Using *In Silico* Mixtures. *Nat Methods* 2023; **20**: 1810–

1821.

53. Mahenthiralingam E, Urban TA, Goldberg JB. The Multifarious, Multireplicon *Burkholderia*

cepacia complex. *Nat Rev Microbiol* 2005; **3**: 144–156.

54. Lessie TG, Hendrickson W, Manning BD, Devereux R. Genomic Complexity and Plasticity

of *Burkholderia cepacia*. *FEMS Microbiol Lett* 1996; **144**: 117–128.

55. Egan ES, Fogel MA, Waldor MK. Divided Genomes: Negotiating the Cell Cycle in

Prokaryotes With Multiple Chromosomes. *Mol Microbiol* 2005; **56**: 1129–1138.

56. Flórez LV, Kaltenpoth M. Symbiont Dynamics and Strain Diversity in the Defensive

Mutualism Between *Lagria* Beetles and *Burkholderia*. *Environ Microbiol* 2017; **19**: 3674–

3688.

57. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. DNA-

DNA Hybridization Values and Their Relationship to Whole-Genome Sequence Similarities.

Int J Syst Evol Microbiol 2007; **57**: 81–91.

58. Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC, et

al. Microbial Species Delineation Using Whole Genome Sequences. *Nucleic Acids Res*

2015; **43**: 6761–6771.

59. Konstantinidis KT, Tiedje JM. Genomic Insights That Advance the Species Definition for

Prokaryotes. *Proc Natl Acad Sci U S A* 2005; **102**: 2567–2572.

60. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. GTDB: An

Ongoing Census of Bacterial and Archaeal Diversity Through a Phylogenetically

Consistent, Rank Normalized and Complete Genome-Based Taxonomy. *Nucleic Acids Res*

2022; **50**: D785–D794.

61. Sudakaran S, Kost C, Kaltenpoth M. Symbiont Acquisition and Replacement as a Source of

Ecological Innovation. *Trends Microbiol* 2017; **25**: 375–390.

62. Janke RS, Moog S, Weiss B, Kaltenpoth M, Flórez LV. Morphological Adaptation for

Ectosymbiont Maintenance and Transmission During Metamorphosis in *Lagria* Beetles.

Front Physiol 2022; **13**: 979200.

63. Fergusson CH, Saulog J, Paulo BS, Wilson DM, Liu DY, Morehouse NJ, et al. Discovery of

a Lagriamide Polyketide by Integrated Genome Mining, Isotopic Labeling, and Untargeted

Metabolomics. *Chem Sci* 2024; **15**: 8089–8096.

64. McTaggart SJ, Obbard DJ, Conlon C, Little TJ. Immune Genes Undergo More Adaptive

Evolution Than Non-Immune System Genes in *Daphnia pulex. BMC Evol Biol* 2012; **12**: 63.

65. Erler S, Lhomme P, Rasmont P, Lattorff HMG. Rapid Evolution of Antimicrobial Peptide

Genes in an Insect Host-Social Parasite System. *Infect Genet Evol* 2014; **23**: 129–137.

66. Behrman EL, Howick VM, Kapun M, Staubach F, Bergland AO, Petrov DA, et al. Rapid

Seasonal Evolution in Innate Immunity of Wild *Drosophila melanogaster*. *Proc Biol Sci*

2018; **285**: 20172599.

67. Engl T, Kroiss J, Kai M, Nechitaylo TY, Svatoš A, Kaltenpoth M. Evolutionary Stability of

Antibiotic Protection in a Defensive Symbiosis. *Proc Natl Acad Sci U S A* 2018; **115**:

E2020–E2029.

68. Kroiss J, Kaltenpoth M, Schneider B, Schwinger M-G, Hertweck C, Maddula RK, et al.

Symbiotic Streptomycetes Provide Antibiotic Combination Prophylaxis for Wasp Offspring.

Nat Chem Biol 2010; **6**: 261–263.

69. Patel PH, Suzuki M, Adman E, Shinkai A, Loeb LA. Prokaryotic DNA Polymerase I:

Evolution, Structure, and 'Base Flipping' Mechanism for Nucleotide Selection. *J Mol Biol*

2001; **308**: 823–837.

70. Bennett GM, Moran NA. Heritable Symbiosis: The Advantages and Perils of an

Evolutionary Rabbit Hole. *Proc Natl Acad Sci U S A* 2015; **112**: 10169–10176.

71. Michalik A, Bauer E, Szklarzewicz T, Kaltenpoth M. Nutrient Supplementation by Genome-

Eroded *Burkholderia* Symbionts of Scale Insects. *ISME J* 2023; **17**: 2221–2231.

72. Dose B, Niehs SP, Scherlach K, Shahda S, Flórez LV, Kaltenpoth M, et al. Biosynthesis of

Sinapigladioside, an Antifungal Isothiocyanate from *Burkholderia* Symbionts. *Chembiochem*

2021; **22**: 1920–1924.

73. Jung BH, Kim JI. Taxonomy of the Genus *Lagria* Fabricius (Coleoptera: Tenebrionidae:

Lagriinae) in Korea. *Korean J Appl Entomol* 2009; **48**: 275–280.

74. Bolger AM, Lohse M, Usadel B. Trimmomatic: A Flexible Trimmer for Illumina Sequence

Data. *Bioinformatics* 2014; **30**: 2114–2120.

75. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A

New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J*

Comput Biol 2012; **19**: 455–477.

76. Miller IJ, Rees ER, Ross J, Miller I, Baxa J, Lopera J, et al. Autometa: Automated

Extraction of Microbial Genomes From Individual Shotgun Metagenomes. *Nucleic Acids*

Res 2019; **47**: e57.

77. Li H. Minimap2: Pairwise Alignment for Nucleotide Sequences. *Bioinformatics* 2018; **34**:

3094–3100.

78. Vaser R, Sović I, Nagarajan N, Šikić M. Fast and Accurate *De Novo* Genome Assembly

From Long Uncorrected Reads. *Genome Res* 2017; **27**: 737–746.

79. Kolmogorov M, Bickhart DM, Behsaz B, Gurevich A, Rayko M, Shin SB, et al. metaFlye:

Scalable Long-Read Metagenome Assembly Using Repeat Graphs. *Nat Methods* 2020; **17**:

1103–1110.

80. Guan D, McCarthy SA, Wood J, Howe K, Wang Y, Durbin R. Identifying and Removing

Haplotypic Duplication in Primary Genome Assemblies. *Bioinformatics* 2020; **36**: 2896–

2898.

81. Donath A, Jühling F, Al-Arab M, Bernhart SH, Reinhardt F, Stadler PF, et al. Improved

Annotation of Protein-Coding Genes Boundaries in Metazoan Mitochondrial Genomes.

Nucleic Acids Res 2019; **47**: 10543–10552.

82. Edgar RC. Muscle5: High-Accuracy Alignment Ensembles Enable Unbiased Assessments

of Sequence Homology and Phylogeny. *Nat Commun* 2022; **13**: 6968.

83. Suyama M, Torrents D, Bork P. PAL2NAL: Robust Conversion of Protein Sequence

Alignments Into the Corresponding Codon Alignments. *Nucleic Acids Res* 2006; **34**: W609–

W612.

84. Brown JW, Walker JF, Smith SA. Phyx: Phylogenetic Tools for Unix. *Bioinformatics* 2017;

33: 1886–1888.

85. Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. ModelTest-NG: A New

and Scalable Tool for the Selection of DNA and Protein Evolutionary Models. *Mol Biol Evol*

2020; **37**: 291–294.

86. Flouri T, Izquierdo-Carrasco F, Darriba D, Aberer AJ, Nguyen L-T, Minh BQ, et al. The

Phylogenetic Likelihood Library. *Syst Biol* 2015; **64**: 356–362.

87. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: A Fast, Scalable and

User-Friendly Tool for Maximum Likelihood Phylogenetic Inference. *Bioinformatics* 2019;

35: 4453–4455.

88. Seemann T. Prokka: Rapid Prokaryotic Genome Annotation. *Bioinformatics* 2014; **30**:

2068–2069.

89. Emms DM, Kelly S. OrthoFinder: Phylogenetic Orthology Inference for Comparative

Genomics. *Genome Biol* 2019; **20**: 238.

90. Buchfink B, Xie C, Huson DH. Fast and Sensitive Protein Alignment Using DIAMOND. *Nat*

Methods 2015; **12**: 59–60.

91. Buchfink B, Reuter K, Drost H-G. Sensitive Protein Alignments at Tree-of-Life Scale Using

DIAMOND. *Nat Methods* 2021; **18**: 366–368.

Table 1. Metadata for different beetles collected for this study. *Lagria villosa* samples were

collected at three different time points, with one sample reported in a previous study (referred to

as Lv19 in this work) [21, 23].

Sample	Location	Statistics for *Burkholderia* symbiont MAG with *lga* BGC					
		MAG ID	Genome size (Mbp)	N_{50} (bp)	No. of contigs	Longest contig (bp)	Coverage
Lagria villosa 2019 (Lv19)*	São Paulo, Brazil	LvStB	2.07	8,138	294	99,474	1983.67
Lagria	São	Lv20_9	1.88	11,699	200	99,421	281.97

Sample	Location	Statistics for *Burkholderia* symbiont MAG with *lga* BGC					
		MAG ID	Genome size (Mbp)	N_{50} (bp)	No. of contigs	Longest contig (bp)	Coverage
villosa 2020 (Lv20)	Paulo, Brazil						
Lagria villosa 2023 (Lv23)	São Paulo, Brazil	LvStB_2023	2.50	1,892,292	3	1,892,292	2263.86
Lagria rufipennis (Lruf1)	Osaka and Ibaraki, Japan	Lruf1_1	1.93	5,267	368	29,589	636.05
Lagria rufipennis (Lruf2)	Tokushima, Osaka, and Kogashima, Japan	Lruf2_2	2.66	7,451	387	55,249	582.66
Lagria okinawana (Loki)	Okinawa, Japan	Loki_2	2.14	6,360	360	60,396	928.22

Sample	Location	Statistics for *Burkholderia* symbiont MAG with *lga* BGC					
		MAG ID	Genome size (Mbp)	N_{50} (bp)	No. of contigs	Longest contig (bp)	Coverage
Lagria hirta (LhSB)	Hessen, Germany	LhSB_1	1.17	5,434	212	88,724	261.74
Lagria hirta (LhHG)	Rhineland Palatinate, Germany	LhHG_2	1.14	6,598	170	127,478	868.51
Lagria hirta (LhG)	Galicia, Spain	LhG_1	2.70	8,034	376	91,830	557.11
Lagria grenieri (Lgren)	Huelva, Spain	Lgren_6	1.14	5,417	205	91,929	83.02
Lagria atripes (Latri)	Rhineland-Palatinate, Germany	Latri_1	2.20	7,903	280	93,629	29.63
Ecnolagria sp. (Ecno)	New South Wales,	Ecno_3	3.16	8,701	419	87,409	87.61

Sample	Location	Statistics for *Burkholderia* symbiont MAG with *lga* BGC					
		MAG ID	Genome size (Mbp)	N$_{50}$ (bp)	No. of contigs	Longest contig (bp)	Coverage
	Australia						

Figure 1. Beetle mitogenome phylogenetic tree using 13 mitochondrial protein coding genes

constructed using MrBayes [33]. Branch values represent posterior probabilities. Mitogenomes

recovered in this study are highlighted with bold lettering.

Figure 2. Analysis of representative *lga* BGCs extracted from eleven Lagriinae beetle

metagenomes. A) Comparison of representative *lga* BGC gene organization. Individual genes in

the *lga* BGCs are represented by arrows oriented in the predicted direction of transcription and

colored according to identity. Pairwise amino acid similarity between BGCs is indicated in the

shaded areas between genes. A scale bar is provided for gene size. Dashed lines indicate

fragments missing from the respective assemblies. B) Comparison of predicted enzyme domain

organization in the representative *lga* BGCs, where genes are ordered according to biosynthetic

order. Boxes around the domains indicate differences between the BGCs.

Figure 3. A) Circular representation of LvStB_2023 genome from the *L. villosa* 2023 sample.

Individual chromosomes are indicated by shades of purple. Coding sequences (CDS) which are

core genes or pseudogenes, as indicated by shades of green, while the rest are indicated in

shades of blue. B) Raw count of COG categories present on different contigs of the LvStB_2023

genome (with and without pseudogenes) from the *L. villosa* 2023 sample.

Figure 4. RAxML phylogenetic tree (left) and shared hierarchical orthogroups (HOGs) (non-

pseudogenes) between different *Burkholderia* genomes (matrix on the right). Each blue line

indicates a shared HOG. HOGs have been hierarchically clustered on the x-axis to improve

visualization. Values on nodes indicate bootstrap values. Genome size and coverage is

represented in brackets next to MAG ID. Outgroups include - *Paraburkholderia acidiphila*

(GCF_009789655.1), *Cupriavidus necator* (GCF_000219215.1), *Herbaspirillum seropedicae*

(GCF_001040945.1). Tree has been truncated for visualisation purposes, full tree can be found

in **Fig. SI 4** (RAxML phylogeny) and **Fig. SI 5** (bayesian phylogeny).

Figure 5. Congruence between phylogenies of beetle host, *Burkholdria* symbionts and *lga*

BGCs in all samples. A) Tanglegram between *lga*-carrying symbionts and beetle host

phylogeny. B) Tanglegram between *lga*-carrying symbionts (centre) and the *lga* BGC, as

inferred via two models GTRCAT (left) and GTRGAMMAI (right). Clades within the beetle host

mitogenome phylogenetic tree are highlighted in shades of grey. In all panels, the three

conserved clades are highlighted in purple, green and orange. Dashed brown lines represent

nodes that are unique between the respective phylogenies, except when the node is absent

from the other tree. Blue circles indicate clade nodes that are poorly supported.

Tree scale: 0.1

Non producer | **_lga_ producer**
Culturable | **Reference**

Outgroup +
other Burkholderia

100 — Burkholderia sp. Lgren_6 (1.14 Mbp, 83.02x) | Spain
100 — Burkholderia sp. Latri_1 (2.02 Mbp, 29.63x) | Germany
Burkholderia sp. Ecno_3 (3.16 Mbp, 87.61x) | Australia
100 — Burkholderia sp. Loki_2 (2.14 Mbp, 928.22x) | Japan
Burkholderia sp. LhG_1 (2.70 Mbp, 557.11x) | Spain
100 — Burkholderia sp. LhSB_1 (1.17 Mbp, 261.74x) | Germany
100 — Burkholderia sp. LhHG_2 (1.14 Mbp, 868.51x) | Germany
96 — Burkholderia sp. Lv20_9 (1.88 Mbp, 281.97x) | Brazil
100 — Burkholderia sp. Lv2023 (2.50 Mbp, 2263.86x) | Brazil
100 — Burkholderia sp. LvStB (2.07 Mbp, 1983.67x) | Brazil
100 — B.gladioli Lruf1_1 (1.93 Mbp, 636.05x) | Japan
100 — B.gladioli Lruf2_2 (2.66 Mbp, 582.66x) | Japan
100 — B.gladioli A1 (8.11 Mbp) | GCF_000633145.1
100 — B.gladioli Lgren_7 (1.79 Mbp, 5.88x) | Spain
100 — B.gladioli MSMB1756 (8.19 Mbp) | GCF_001527485.1
100 — B.gladioli CHAPALOTE (8.52 Mbp) | GCF_000757585.1
100 — B.gladioli BSR3 (9.05 Mbp) | GCF_000194745.1
100 — B.gladioli ATCC 10248 (8.89 Mbp) | GCF_000959725.1
B.gladioli NBRC 13700 (8.76 Mbp) | GCF_000739755.1
100 — B.gladioli LvStA (8.56 Mbp) | Brazil
100 — B.gladioli SN82F6 (8.45 Mbp) | GCF_000981885.1
100 — B.gladioli Latri_2 (7.58 Mbp, 15.72x) | Germany
100 — B.gladioli LhHG_3 (5.09 Mbp, 100.61x) | Germany
98 — B.gladioli LhSB_5 (7.71 Mbp, 7.05x) | Germany
98 — B.gladioli LhStG (8.40 Mbp) | Germany

Shared HOGs

A

Burkholderia Symbiont

P. acidiphila | GCF_009789655.1

Ecnolagria sp. (Ecno_3) | Australia
Lagria atripes (Latri_1) | Germany
Lagria grenieri (Lgren_6) | Spain

Other *B. gladioli*

Lagria rufipennis (Lruf2_2) | Japan
Lagria rufipennis (Lruf1_1) | Japan
Lagria villosa 2020 (Lv20_9) | Brazil

Lagria okinawana (Loki_2) | Japan
Lagria hirta (LhG_1) | Spain
Lagria hirta (LhSB_1) | Germany
Lagria hirta (LhHG_2) | Germany

2.0   1.5   1.0   0.5   0.0

Beetle host Mitogenome

Mylabris sp. MYL01

Ecnolagria sp. (Ecno) | Australia
Lagria villosa 2020 (Lv20) | Brazil
Lagria atripes (Latri) | Germany
Lagria grenieri (Lgren) | Spain

Lagria rufipennis (Lruf2) | Japan
Lagria rufipennis (Lruf1) | Japan

Lagria okinawana (Loki) | Japan
Lagria hirta (LhG) | Spain
Lagria hirta (LhSB) | Germany
Lagria hirta (LhHG) | Germany

0.0   0.5   1.0   1.5   2.0

B

BGC phylogeny (GTRCAT model)

P. acidicola

Ecno | Australia
Latri | Germany
Lgren | Spain

Lruf2 | Japan
Lv20 | Brazil

Loki | Japan
LhG | Spain
LhSB | Germany
LhHG | Germany

2.0   1.0   0.0

Burkholderia Symbiont

P. acidiphila

Ecno | Australia
Latri | Germany
Lgren | Spain

Other *B. gladioli*

Lruf2 | Japan
Lruf1 | Japan
Lv20 | Brazil

Loki | Japan
LhG | Spain
LhSB | Germany
LhHG | Germany

0.0   1.0   2.0

Burkholderia Symbiont

P. acidiphila

Ecno | Australia
Latri | Germany
Lgren | Spain

Other *B. gladioli*

Lruf2 | Japan
Lruf1 | Japan
Lv20 | Brazil

Loki | Japan
LhG | Spain
LhSB | Germany
LhHG | Germany

1.0   0.0

BGC phylogeny (GTRGAMMAI model)

P. acidicola

Ecno | Australia
Latri | Germany
Lgren | Spain

Lruf2 | Japan
Lv20 | Brazil

Loki | Japan
LhG | Spain
LhSB | Germany
LhHG | Germany

0.0   0.5   1.0   1.5   2.0