Repeated horizontal acquisition of

lagriamide-producing symbionts in

Lagriinae beetles

Short title: Selective beetle symbiont acquisition

Siddharth Uppal*[1], Samantha C. Waterworth*[1,2], Alina Nick (anick@ice.mpg.de)[3], Heiko Vogel

(hvogel@ice.mpg.de)[3], Laura V. Flórez[4], Martin Kaltenpoth[3], Jason C. Kwan[1†]

*These authors contributed equally to this work

[†]To whom correspondence should be addressed. Address: 777 Highland Ave., Madison, WI

53705. Email: jason.kwan@wisc.edu

[1]Division of Pharmaceutical Sciences, School of Pharmacy, University of Wisconsin-Madison,

Madison, USA.

[2]Current address: National Cancer Institute, Frederick, Maryland, USA

[3]Max Planck Institute for Chemical Ecology, Jena, Germany

[4]Department of Plant and Environmental Science, University of Copenhagen, Copenhagen,

Denmark

Study funding information: NSF DBI-1845890 (S.U., S.C.W., J.C.K.)

Abstract

Microbial symbionts associate with multicellular organisms on a continuum from facultative associations to mutual codependency. In the oldest symbioses there is exclusive vertical symbiont transmission, and extensive co-diversification of symbiotic partners over millions of years and speciation events. Such symbionts often undergo genome reduction due to low effective populations, frequent population bottlenecks and weakened purifying selection. We describe here a group of Lagriinae beetle symbionts, which produce defensive lagriamide molecules to protect beetle eggs and larvae from fungal infection. These beetles likely evolved specialised symbiont-storage structures between 55 and 82 million years ago. Previous work found, despite the lagriamide-producing bacterium not being genetically isolated and potentially surviving in the environment, it underwent genome reduction in *Lagria villosa* hosts. Here, we use shotgun metagenomics to assemble 11 additional lagriamide-producing symbionts from diverse hosts within Lagriinae from five countries, to clarify the evolution of this symbiotic relationship. Surprisingly, we did not find evidence of co-evolution between symbionts and hosts, and although the lagriamide gene cluster (*lga*) had been horizontally transferred to the *L. villosa* symbiont, it likely was not transferred between symbionts. Instead, *lga* was likely obtained only once in the common ancestor of all lagriamide-producing symbionts and subsequently lost in related free-living strains. Our results suggest that beetles acquired lagriamide-producing symbionts specifically multiple times independently, which was followed by genome erosion. We also show that these symbionts have been replaced in the beetle host multiple times. The system therefore offers a unique opportunity to study the process of genome reduction.

Keywords: lagriamide, *Burkholderia*, symbiosis, symbiont replacement, biosynthetic gene cluster, metagenomics, Lagriinae, chemical defence, secondary metabolism

Introduction

Eukaryotic organisms have been associated with prokaryotic microbes since the initial endosymbiotic events which led to the acquisition of mitochondria and chloroplasts [1]. These organelles represent the presumed endpoint of ancient symbioses with α-proteobacteria and cyanobacteria, respectively, that over time led to a progressive shrinkage of the symbiont's genome and eventual transfer of genes from symbiont to host [1]. Although organelle acquisition appears to be a rare event [2], other more recent symbioses appear to be on a similar evolutionary trajectory of profound genome reduction and absolute dependence on host cells. For example, the acquisition of the intracellular symbiont *Buchnera aphidicola* in the common ancestor of aphids allowed them to diversify as sap-feeding insects since the symbiont synthesises several essential amino acids not found in plant sap, as evidenced by a rapid basal radiation of aphid species [3] and strict co-evolution of aphids and *Buchnera* [4]. *B. aphidicola* has been vertically transmitted for at least 200 million years [4] and has a profoundly reduced chromosome, about 11% of the size of that of *Escherichia coli* [5]. Through comparison of various symbionts, a model of genome reduction has emerged whereby host-restriction initially weakens purifying selection on formerly essential genes, through both host-provided metabolites and symbiont population structure, with low effective populations isolated in individual hosts [6]. When symbionts are vertically transmitted, population bottlenecks occur every time a host procreates, causing the fixation of deleterious mutations within the population [6]. These factors combine to first cause a proliferation of non-functional pseudogenes in the genome [6] and then deletion of those pseudogenes due to a known deletion-bias within bacteria [7]. The most reduced genomes lose even central functions such as DNA repair pathways [6], which leads to an increased rate of evolution and further gene loss, as well as increased AT-bias in many cases [8, 9]. In the cases of symbionts living inside host cells, it is easy to suppose that this process is exacerbated due to a lack of opportunity or ability to obtain

functional genes through homologous recombination - i.e. a combination of genetic isolation and

disruption of homologous recombination apparatus.

However, genome reduction is also known to occur in the absence of genetic isolation. For

instance, free-living bacteria living in nutrient poor environments such as *Prochlorococcus* spp.

are thought to have reduced genomes as a consequence of selection pressure to streamline

their metabolism [10], potentially explained through the Black Queen hypothesis [11], which

posits that selection drives pathways to be lost when they are produced by another species in

the ecosystem as "public goods". There are also genome-reduced symbionts which seemingly

are not genetically isolated. *Burkholderia* symbionts that reside extracellularly in leaf nodules in

plants are mainly transmitted vertically because the symbiosis is mutually co-dependent [12],

although horizontal transfer may have occurred rarely between plants, the soil microbiota and

insects [13]. This suggests a lack of genetic isolation, and indeed there is evidence of repeated

horizontal transfers of biosynthetic genes for defensive molecules among leaf nodule symbionts

of *Rubiacaeae* plants [13]. Such systems may provide an opportunity to study the evolutionary

pressures that lead to the process of genome reduction, and the mechanisms of symbiosis that

underlie it.

The dichotomy of vertical versus horizontal transfer of symbionts may be one determinant of

genome reduction. A relatively clear-cut example are two symbionts of the tunicate *Lissoclinum*

patella, the extracellular cyanobacterium *Prochloron didemni* [14] and the intracellular

"*Candidatus* Endolissoclinum faulkneri" [15]. The former is capable of horizontal transfer, as

evidenced by its almost clonal genome amongst very divergent hosts and lack of genome

reduction [14], while the latter is vertically transmitted, as evidenced by its divergence with its

hosts across cryptic speciations, and profound genome reduction [15, 16]. However, the mode

of transmission also exists on a continuum from strict vertical to strict horizontal, with mixtures of

vertical and horizontal transmission in between. For instance, the Tsetse fly symbiont *Sodalis*

glossinidius shows some signs of genome-reduction such as rampant pseudogenes, but

remains culturable in the lab, meaning that horizontal transfer cannot be excluded [17].

Likewise, symbionts long thought to be exclusively vertically transmitted, such as the bryozoan

symbiont "*Ca.* Endobugula sertula", which is packaged with the hosts' larvae, show no signs of

genome reduction [18], meaning that there is no compelling reason why it should not be able to

transfer horizontally between hosts. Indeed, "*Ca.* E. sertula" has been found in genetically

divergent but proximal bryozoan individuals, suggesting horizontal transfer [19].

The *Lagria* and *Ecnolagria* beetles belong to subfamily Lagriinae within the family

Tenebrionidae (order Coleoptera). The *Lagria villosa* beetle, a known soybean pest [20], is a

source of lagriamide, an antifungal polyketide [20]. The compound is produced via a *trans*-AT

polyketide synthase (PKS)-non-ribosomal peptide synthetase (NRPS) hybrid biosynthetic gene

cluster (BGC), termed *lga,* which due to a nucleotide signature (kmer frequency) distinct from

the chromosome is predicted to have been horizontally acquired [20]. The *lga* BGC is encoded

by a *Burkholderia* symbiont (*Burkholderia* sp. LvStB) that is present in glandular structures

associated with the ovipositor of female beetles and secreted on its eggs as they are laid [20].

The symbiont has been shown to have a defensive role against fungi in the egg [20] and larval

stages [21]. Previously, we showed that the genome of *Burkholderia* sp. LvStB is reduced and

has lost a number of essential genes including some genes involved in the DNA repair

pathways and primary metabolism [22]. The genome has a low coding density, and a high

number of pseudogenes and transposases, indicative of a reduced genome [22]. These

characteristics are consistent with host restriction and vertical transmission of LvStB. However,

there is evidence that *Burkholderia* symbionts from *L. villosa* can be transferred to plant tissues

and persist for several days, and that bacteria can be acquired by the beetle from the plant and

soil environment [23].

As some Lagriinae beetles harbour symbionts in special structures that likely evolved between

55 and 82 million years ago based on fossil evidence, positioned to deposit symbionts on the

eggs [24, 25], we hypothesised that lagriamide-producing *Burkholderia* symbionts might have

co-evolved with their hosts in a manner similar to other vertically-transmitted insect symbionts.

However, the possibility for transmission of the symbionts to and from plants, and the

accessibility of the symbionts' habitat on the surface of eggs and within adult females suggested

that horizontal symbiont acquisition may be possible. As the beetles harbour complex

microbiomes with multiple related *Burkholderia* strains as well as other bacteria [21, 23], both

genome-reduced and not, an alternative hypothesis is that the lagriamide BGC has been

repeatedly horizontally transferred among environmental strains and symbionts. Moreover,

partnerships in defensive symbionts are usually more dynamic as compared to intracellular

nutritional symbionts [26]. It's also possible that the lagriamide-producing strain is restricted to *L.*

villosa, and that different Lagriinae species have symbionts with different BGCs, as this would

allow the association to react much more flexibly to changes in antagonist communities. To

clarify this evolutionary picture, we analysed the metagenome of 12 beetle populations,

spanning eight species belonging to genera *Lagria* and *Ecnolagria* across five different

countries (four continents) (**Table 1**). We recovered the metagenome-assembled genomes

(MAGs) of several different *Burkholderia* bacteria, and confirmed the presence of the lagriamide

BGC in each. We also report a complete genome of the genome-reduced, lagriamide-producing

Burkholderia sp. LvStB symbiont, obtained through long-read Nanopore sequencing. We

compared the phylogeny of the recovered *Burkholderia* MAGs, the lagriamide BGCs, and the

host beetles to determine whether co-cladogenesis occurred in this system, and to further

explore the evolutionary relationships in the symbiosis. The results indicate that the lagriamide

BGC was likely only acquired once in the common ancestor of beetle-associated *Burkholderia*

symbionts, and subsequently lost in the majority of the descendent free-living strains.

Remarkably, as all the lagriamide-bearing symbionts are genome-reduced but they do not form

a monophyletic clade and do not correspond to host phylogeny, they may represent multiple

acquisition events, followed by independent genome-reduction processes. The common factor

of lagriamide production might be one of the reasons for selection by and dependency on hosts.

This would suggest that a single group of natural products caused many independent

symbioses to be established over evolutionary time.

149　Results and Discussion

150　Beetle phylogeny

We sequenced and assembled the metagenomes of the collected *Lagria* and *Ecnolagria* beetle

populations (**Table 1**), and beetle mitogenomes were extracted and annotated to infer host

beetle phylogeny (**Fig. 1**). In line with previous studies, mitogenomes belonging to subfamilies

Lagriinae, Blaptinae, Pimeliinae, Stenochiinae and Alleculinae were found to be monophyletic

whereas Diaperinae and Tenebrioninae were found to be para- or polyphyletic [27–30].

Maximum likelihood analysis using RAxML [31] (**Fig. SI 1**) and Bayesian analysis using

MrBayes [32] (**Fig. 1**) gave the same results.

All collected *Lagria* beetle mitogenomes clustered into four distinct subclades: All *L. hirta* beetle

mitogenomes were clustered in a single clade, alongside a closely related clade of several

Lagria species (*L. rufipennis*, *L. okinawana* and *L. nigricollis*) from Japan. The *L. atripes* and *L.*

grenieri beetles formed another clade more distantly related to the *L. hirta* and Japanese *Lagria*

species. Finally, the *L. villosa* and *Ecnolagria* sp. beetles formed a fourth clade along with

Chrysolagria sp. (JX412760), distinct from the other *Lagria* beetles. Furthermore, while publicly

available sequences of *L. hirta* (OX375806) clustered with collected *L. hirta* samples from

Rhineland Palatinate, Germany (LhHG), *L. rufipennis* (MW802588) clustered outside the *L.*

rufipennis (Lruf), and *L. nigricollis* (Lnig) clade. This slightly distant clustering of the two *L.*

rufipennis mitogenomes could be due to the different geographical origins (China and Japan,

respectively). Unfortunately, however, the scarcity of taxonomic work and lack of molecular

barcodes on the Asian *Lagria* species in the BOLD database [33] preclude us from

unambiguously resolving this issue.

172 Recovery of Lagriamide BGCs

A complete, or mostly complete, *lga* BGC was found in eleven of the twelve samples, with the

exception of *L. rufipennis* where only small fragments of the *lga* BGC could be recovered. The

BGC recovered from *L. nigricollis* was found over two contigs and could not be manually joined

following inspection of the assembly graph. The missing data for this region spans from

approximately halfway through the *lgaB* gene to approximately halfway through the *lgaC* gene

(**Fig. 2A**). Inspection of the assembly graphs with focus on the *lga* BGC regions revealed

several possible variants of the BGC present in the symbionts associated with *L. atripes*, *L. hirta*

SB, *L. okinawana*, and *L. nigricollis* (See Supplementary Material). Close analysis of the

variants revealed that neither the gene nor domain organisation and predicted functionality

thereof was affected within the variants (i.e intra-variant organisation). We used read coverage

as a proxy for the most dominant BGC variant and the variant with the greatest coverage

support was chosen as the representative BGC for subsequent analyses. Full analysis

regarding BGC variants and selection of representative BGCs can be found in Supplementary

Material.

Analysis of representative BGCs revealed several differences in gene organisation of the *lga*

BGC across the different Lagriinae beetle species (**Fig. 2A**). The first difference we observed

was among the *lga* BGCs recovered from the *L. hirta* samples, wherein most *lga* BGCs

exhibited a split in *lgaC* (**Fig. 2A**). We initially had only short-read sequence data for the *L. hirta*

HG population, from which we recovered a *lga* BGC that exhibited a split in *lgaC*, as observed in

other *L. hirta*-derived *lga* BGCs (**Fig. 2A**). However, we later acquired additional long-read

sequence data for the *L. hirta* HG population and, following reassembly, were surprised to find

that the recovered *lga* BGC did not exhibit the previously observed break in *lgaC*. PCR

amplification of this region in the *lgaC* gene revealed that both complete and split variations of

lgaC were real and present in the *L. hirta* HG population as sequence variants. The coverage of

the *lga* BGC is the same as that of the predicted host genome, which would suggest that it is

present as a single copy. We therefore concluded that different BGC variants exist in the *L.*

hirta HG-associated *Burkholderia* population and may exist in the other Lagriinae populations.

Both read coverage and PCR amplification suggest that the variant with the complete (not split)

lgaC is more abundant.

The second difference we observed was in the *lga* BGC from the LvStB genome extracted from

the 2023 *L. villosa* metagenome (Lv23). One split in *lgaB* and two splits in *lgaC* were seen.

However, the assembly of the *L. villosa* 2023 metagenome was based solely on long-read data,

which is error-prone [34], and the splits may not be a true reflection of the BGC in this sample.

Normally Sanger sequencing would be the solution to validate these questionable regions but

unfortunately, there was no remaining DNA after the long read sequencing runs for this

particular sample. For this reason, we left the BGC with the splits but were cautious not to over-

interpret the apparent breaks in the genes in this BGC.

We then considered the domain organisation within the *lga* BGC genes (**Fig. 2B**). The domain

organisation is largely congruent across the *lga* BGCs recovered from the metagenomes. We

did note, however, an additional annotated "DHt" domain in *lgaB,* which is defined as

"Dehydratase domain variant more commonly found in *trans*-AT PKS clusters", in the *lga* BGCs

from all *L. hirta* samples and the *L. nigricollis* population. Similarly, we detected an additional

carrier protein domain (phosphopantetheine acyl carrier protein group) near the N-terminus of

the *lgaC* protein in the BGCs from the *L. grenieri* and *L. okinawana* samples. In all cases, close

inspection of the primary sequence of these additional dehydratase and carrier domains

revealed mutations in the sequences that would likely render the enzyme non-functional (see

Supplementary Material for full details).

Finally, as with the originally described *lga* BGC recovered from the *L. villosa* 2019 sample [20],

we found mutations in the catalytic or conserved motifs of *lgaG* DH2, *lgaG* KS6, *lgaB* KR3 and

lgaC KS5 domains, that we believe may render these domains inactive (see Supplementary

Material for full details). As a result, the domain architecture of all representative *lga* BGCs from

all samples are likely functionally identical.

Together, the conservation of the *lga* BGC in at least seven different species of Lagriinae

beetles, across four geographically distant countries, implies that the production of lagriamide is

an important factor for the host beetle and that the *lga* BGC is under strong selective pressure.

The presence of additional domains in the *lga* BGC in several samples, even though they are

likely inactive, is intriguing as it suggests that these domains may have previously been present

in all *lga* BGCs but may have decayed over time and were lost. The reason as to why these

domains were selected against would be speculative at best and inspection of all lagriamide-like

compounds produced in the different beetle populations would need to be assessed to truly infer

differences that the domain architecture may have on the resulting chemistry. Conserved

production of of other bioactive compounds has been observed, such as pederin, across

Staphylinidae beetle species (*Paederus* and *Paederidus* genera) [35], which are host to a

Pseudomonas symbiont that produces pederin [36]. Pederin is a toxic compound, produced via

a *trans*-AT PKS-NRPS hybrid BGC by the *Pseudomonas* symbiont that is coated onto the

beetle eggs and subsequently maintained by the larvae into adulthood and serves to protect

them against predation [37].

The two systems have several parallels: both pederin and lagriamide are produced by a *trans*-

AT PKS NRPS hybrid BGC in *Pseudomonas* bacterium [35], where the compound is

concentrated in the female oviposition organs, coated onto the eggs and serves to protect

juveniles [37]. Further, both pederin and lagriamide are the sole insect-associated compounds

in suites of compounds otherwise associated with marine invertebrates. Groups of pederin

analogs, such as the onnamides, mycalamides, psymberins, and theopederins have been

isolated from a variety of marine sponges [38–42] and ascidians [43], while bistramide, the most

structurally similar compound to lagriamide, was isolated from an ascidian [44]. Another

analogous system is that of beewolf wasps and *Streptomyces* bacteria which exhibit some

characteristics of genome erosion [45] and produce at least two classes of defensive

compounds: piericidins, and streptochlorins [46]. In this system, the *Streptomyces* bacteria are

extracellular symbionts stored in cuticular invaginations in the female wasp antennae and are

transmitted to the brood cells and subsequently transferred to the cocoon to protect the larvae

from fungal infection [46]. Piericidins are similarly produced by a *Streptomyces* harboured within

marine ascidian *Didemnum molle* [47, 48], and streptochlorins were originally isolated from a

marine-sediment borne *Streptomyces* species [49]. The question remains, however, as to what

the fundamental link is between these terrestrial and marine systems. What is clear, however, is

that the production of these compounds is crucial in the presence of predators, which provides

additional evidence that the production of lagriamide likely provides equally crucial defence for

the Lagriinae beetle eggs and larvae against fungal pathogens [20].

Complete genome of the *lga*-carrying LvStB symbiont

Long read sequencing of the *L. villosa* 2023 (Lv23) metagenome allowed us to assemble a complete genome of a *lga*-carrying *Burkholderia* strain (referred to as LvStB_2023 from hereon). LvStB_2023 was found to have a 2.5 Mbp long genome with a GC percentage of 58.63%. It has 2 circular chromosomes - chromosome 1 is 1.89 Mbp, chromosome 2 is 0.55 Mpb in size, and there is a plasmid 59.77 kbp long. The genome is estimated to be 97.1% complete (98.8% with "specific" mode) and 0.02% contaminated as per CheckM2 [50] and thus is a high quality MAG as per the MIMAG standards [51]. Assembly graph analysis of LvStB_2023 verified that we have the complete sequence of two circular chromosomes and a plasmid. However, the CheckM2 estimate did not reflect a fully complete genome, at 98.8%, and we believe that this small discrepancy in predicted completeness may be a result of ongoing genome reduction [52].

LvStB_2023 has a coding density of 78% and 59.1% with and without pseudogenes respectively. A large percentage (43.87%) of the ORFs in LvStB_2023 were identified as pseudogenes (1613 out of 3676), the highest of any *lga*-carrying *Burkholderia* symbiont. However, this estimate may be artificially high as pseudogenes are identified purely based on their length relative to their closest BLASTP match and these counts are derived from an assembly generated from only long-read data which can be prone to errors, particularly homopolymeric runs. However, coding density and frequency of pseudogenes is not a great deal different to LvStB MAGs assembled from short-read data (see Table S1 for complete genome characteristics of recovered MAGs). Having multiple chromosomes is a common phenomenon in *Burkholderia* [53, 54]. Generally in multi-chromosome bacteria, the majority of the genes for essential functions are located on one larger or primary chromosome. The smaller or the secondary chromosome has much fewer essential genes and mostly carries genes for niche specific functions [55]. In the case of LvStB_2023, chromosome 1 appears to be the

primary chromosome as it is much larger in size, and has 77 out of 84 core genes (including

multiple copies) (**Fig 3A**). Functional analysis revealed chromosome 1 to have the highest

number of genes for all essential COG categories (**Fig 3B**), including categories L (replication,

recombination and repair), J (Translation, ribosomal structure and biogenesis), M (Cell

wall/membrane/envelope biogenesis) and H (Coenzyme transport and metabolism).

The *lga* BGC is on chromosome 2 (0.55 Mbp long) and can be distinguished by the continuous

block of reverse coding sequences in **Fig 3A**. A detailed view of the *lga*-carrying chromosome

can be seen in **Fig. SI 2,** where the GC content of the *lga* BGC is significantly higher than the

rest of the genome, providing further evidence towards its horizontal acquisition. Chromosome

1, chromosome 2 and the plasmid have 44.75%, 37.59% and 42.34% of their coding capacity

taken up by pseudogenes, respectively. The similar abundance of pseudogenes in each of the

contigs indicates that the whole genome is undergoing reduction simultaneously. The

chromosome with *lga* (chromosome 2) has the smallest percentage of pseudogenes, which may

be a reflection of the required conservation of the *lga* BGC in combination with the presence of

large genes in *lga*.

302 Diversity of beetle-associated *Burkholderia* symbionts

303 Recovery and analysis of metagenome assembled genomes

Following assembly, the 12 beetle metagenomes were binned, and resultant bins manually

refined. A total of 77 MAGs were recovered from all samples, of which 24 MAGs were of high

quality, 30 of medium quality, and 23 of low quality (**Table S1**) in accordance with published

MIMAG standards [51]. Only medium and high quality MAGs were used for downstream

analysis, with the exception of one low quality bin carrying the *lga* BGC (LhHG_2). Confirmation

that the *lga* BGC had been assigned to the correct MAG has been covered in detail in the

Supplementary Material. Genome erosion, such as that already observed for the *lga*-carrying

symbiont *Burkholderia* sp. LvStB [22], can skew the completeness metric. To determine if a

lower quality MAG was incomplete or genome-reduced, we also considered several other

metrics, including core gene presence, number of pseudogenes [22], and coding density (**Table**

S1), and concluded that this particular MAG (LhHG_2) was likely both reduced and incomplete.

For each beetle population a single MAG belonging to genus *Burkholderia* with a single copy of

the *lga* BGC was identified, with the exception of the symbiont within *L. nigricollis* where the

possibility exists that the *lga* BGC is present in two copies (Please see Supplementary Material

for full details). Previous studies into the lagriamide-carrying symbiont strain *B. gladioli* LvStB

[20, 22, 56], showed that this strain was significantly more abundant than all other bacteria

associated with *L. villosa*, and had a reduced genome. Consistent with this, all newly recovered

MAGs that included *lga* BGCs were the most abundant MAGs in each sample and had reduced

genomes with an abundance of pseudogenes and transposases, and had lower coding

densities relative to other *B. gladioli* genomes (**Table S1**). In standing with previous studies of

Lagria beetles, where both reduced and non-reduced *B. gladioli* genomes were recovered,

additional *B. gladioli* MAGs (Latri_2, LhHG_3, and LhSB_5) were recovered that did not carry

the lagriamide BGC and showed no evidence of genome erosion. We also recovered three

small *B. gladioli* MAGs (Lgren_7, Lv19_6_18, Lv20_2) and one small *Burkholderia* MAG

(Lv19_6_14). We also recovered MAGs classified as *B. lata* (Lv19_4_0) and *B. arboris*

(Lv20_1).

Average nucleotide identity (ANI) analysis of *B. gladioli* MAGs carrying the *lga* BGC showed that

MAGs from different beetle species and/or different locations were likely different bacterial

species due to shared ANI values less than 95% [57]. Studies have suggested that ANI alone is

not a sufficient metric for species delineation and that the aligned fraction (AF) must also be

taken into account [57–60]. Following recent cutoffs adopted for species delineation [58], we

opted to use AF ≥ 60% along with ANI ≥ 95% as a cutoff for species assignment. Subsequently,

we found that the *Burkholderia* MAGs carrying the lagriamide BGC appeared to be split into at

least five novel species (**Table S2**).

Phylogenetic analysis of recovered metagenome assembled genomes

Phylogeny of the *lga*-carrying *Burkholderia* symbionts, relative to other *Burkholderia* species,

was inferred using hierarchical orthogroups identified using Orthofinder [61] (**Fig 4** and **Fig. SI**

3). *Burkholderia* symbionts with the *lga* BGC were closely related but did not form a

monophyletic clade. Additionally, we included the genome of a free-living *Paraburkholderia*

acidicola ATCC 31363 from soil which was recently found to carry a related lagriamide-like BGC

(*lgb*), that encodes a structurally similar compound [62] (**Fig SI 4**). This bacterium clustered

with other *Paraburkholderia* species.

Unexpectedly, the *Burkholderia* symbionts with a *lga* BGC did not form a monophyletic clade, as

would be expected in a long-term vertically-transmitted symbiont, and their common ancestor

existed before the common ancestor of many of the free-living *B. gladioli* strains. Given that it is

highly unlikely that genome-eroded symbionts could revert to free-living strains with large

genomes, the symbiont phylogeny is indicative of at least four independent transitions to a

symbiotic lifestyle in closely related bacteria with the *lga* BGC. Each of the transitions was

subsequently followed by genome erosion. The lack of synteny observed in the genes flanking

the *lga* BGC is indicative of either genomic rearrangement or independent acquisition of the *lga*

BGC, and further supports the independent acquisitions of symbionts followed by genome erosion (**Fig SI 5**).

The phylogeny of the *lga* BGC-carrying *Burkholderia* symbionts was found to be largely incongruent with the beetle phylogeny (**Fig 5A**). The only clades that showed possible co-cladogenesis with the beetle host were the clade of *L. hirta* and their associated BGCs with the clade of *L. nigricollis* and *L. rufipennis*, and their associated *lga* BGCs. This incongruence in the symbiont-host phylogeny was indicative to us of possible host jumping of the symbiont or on-going replacement of the symbiont due to sporadic loss from genomic degradation of the symbiont. Vertical transmission of symbionts is often linked with genome erosion including high numbers of pseudogenes, low coding density and loss of genes required for host-independent survival [6, 63]. Symbiont replacement has often been reported in nutritional symbionts as a way for the hosts to replace a genetically degraded symbiont with a more complete and effective one and to acquire new adaptations for expanding into different niches [64]. *Burkholderia* symbionts related to *B. gladioli* in *Lagria* beetles have been reported to evolve from plant associated bacteria [25] capable of transfer from beetles to plants with subsequent survival [23]. It is possible that the horizontal acquisition might occur in the egg and larval stages, where the symbionts are localised on the surface (eggs) or in cuticular invaginations (larvae and pupae) that remain connected to the external surface via a small duct [65]. As the closely related *Burkholderia* strain LvStA can be acquired horizontally from the environment [23], and there is evidence of free-living bacteria carrying lagriamide-like BGCs [62], we propose that there are *lga*-carrying *Burkholderia* strains persevering in the environment (e.g. in plants or soil) [23] that can be horizontally acquired by the beetle host.

As we previously observed that *lga* has distinct nucleotide composition to the Lv19 genome [20],

suggestive of a recent horizontal transfer, we sought to determine if it has been independently

transferred to the corresponding symbiont in different beetle hosts. Phylogenetic analysis of the

representative *lga* BGCs from all samples resulted in two possible topologies using GTRCAT-V

and GTRGAMMAI models (**Fig. 5B**). Both topologies included conserved clades. However, the

relative positions of the three clades are poorly supported (**Fig. 5B**, highlighted in blue),

resulting in the two alternative topologies. A Bayesian tree was also constructed (**Fig. SI 6**)

which is congruent with the GTRGAMMAI tree topology. The inconsistent topology likely stems

from limited resolution of the phylogeny affecting deep nodes in the trees, and we interpret the

totality of trees to be largely congruent with symbiont phylogeny without additional evidence of

more horizontal transfer events. It is likely that there was a single acquisition of *lga* in the

common ancestor of the symbiont and *B. gladioli* clade, with subsequent loss in the free-living

group. Comparison of both BGC phylogenetic trees with the beetle host phylogeny revealed that

both possibilities were incongruent with the beetle host phylogeny (**Fig. 5C**), as seen with the

symbionts. It appears that lagriamide production was highly selected for in symbiotic settings

and hence retained, whereas it was lost in the most of the larger genomes (assumed to be free-

living) where it was not selected for. These findings indicate that the *lga* BGC is important in the

symbiosis, either for symbiont establishment (e.g. competition with other symbionts) and/or

because lagriamide is an effective host-defensive molecule. Furthemore, the fact that highly

similar *Burkholderia* species with *lga* were identified across different Lagriinae beetles indicates

that symbiont acquisition is highly selective.

We performed an analysis of pentanucleotide (5-mer) frequency of the beetle-associated, *lga*-

carrying symbionts and their associated BGCs, along with the genomes of recently identified

soil-borne *Paraburkholderia* species that carry the lagriamide B (*lgb)* BGC, which is highly

similar to the *lga* BGC [62], to determine if there was any evidence that the *lga* BGC had been

originally acquired from an organism like the *Paraburkholderia*. Visualisation of 5-mer frequencies of the BGCs and the genomes revealed three clusters of BGCs: The BGCs from the two soil-borne *Paraburkholderia* strains, the BGCs from the Brazilian *L. villosa*-derived LvStB strains, and then a third cluster of all other *lga* BGCs (**Fig. SI 7**). A similar pattern was observed for the nucleotide composition of the respective genomes wherein LvStB and Lv20_9 form an isolated cluster, the two soil-borne *Paraburkholderia* form a second, distant cluster, and all other *lga*-carrying Burkholderia strains and cultured *Lagria*-associated genomes (LvStA and LhStG) form a third cluster. None of the BGCs share similar 5-mer composition with their respective genomes suggesting that all are likely horizontally acquired from an independent donor organism.

We noted in a previous analysis of the COG annotated genes that there appeared to be a particularly high number of genes in the L category (replication, recombination and repair) that were likely pseudogenes (**Fig 3B**). We assessed the percentage change of COG annotated genes in all *lga*-carrying *Burkholderia* and found that this pseudogenization of genes involved in DNA replication, recombination and repair was particularly high in all the Brazilian *L. villosa*-derived LvStB strains, as well as MAGs LhSB_1, LhG_1, Loki_2 and Lgren_6 (**Fig. SI 8**). Two of the three the LvStB strains also exhibited high pseudogenization of the genes associated with cell motility (Category N). While COG annotation of genes does not provide a robust picture, as not all genes are successfully annotated, the increased pseudogenization of genes involved in DNA replication and repair may explain the drift of the LvStB strains observed in both the phylogenetic analysis and the related 5-mer analysis. In particular, LvStB MAGs possessed highly truncated and psuedogenized *polA* genes, coding for DNA polymerase I used in many DNA-repair pathways and chromosome replication [66], whereas other *lga*-containing MAGs, except LhHG_2, had intact *polA* genes (**Fig. SI 9**). The loss of *polA* in the *L. villosa* symbionts explains their accelerated sequence drift in the genome as a whole and also in the *lga* BGC

compared to the *lga*-possessing symbionts (**Fig. SI 7**). The relative shortness of the LhHG_2

branch in the symbiont tree suggests that its *polA* gene might have degraded relatively recently.

The presence of the *lga* BGC in closely related *Burkholderia* symbionts and not in free living *B.*

gladioli indicates it is heavily selected for in the beetle symbiosis ecosystem and only specific

bacteria with the BGC are acquired by the beetles. It is possible that lagriamide might be

protective against natural enemies other than fungi. How and why it is highly selected for

despite the different geographical locations of these hosts is intriguing and suggests a number

of interesting questions, considering that the beetles studied here have access to many other

symbionts that can produce secondary metabolites, including antifungals [22], such as the

antifungal caryoynencin, which was isolated from the culturable beetle symbiont *B. gladioli*

LvStA.

Previous studies have highlighted how symbionts can be conserved across host-speciation

events and millions of years, leading to genome reduction in the symbiont [6, 16]. A

disadvantage of such an exclusive relationship is that the symbiont inevitably suffers from

increasingly severe genome erosion, and the end point of that process may be complete

dysfunction [67]. Our data concerning the Lagriinae beetles presented here suggests another

possibility, that if there is a reservoir of bacteria in the environment that harbour a pathway for a

specific secondary metabolite, they could serve as replacements for degraded symbionts. The

apparently repeated acquisitions and then genome reduction of symbionts suggests a unique

symbiosis specific to a molecule and not to a symbiont. If that hypothesis is correct, it would

suggest that lagriamide plays a role in host-symbiont communication or symbiont acquisition, in

addition to being a chemical defence. Perhaps the most intriguing aspect of this system is the

seemingly repeated independent genome reduction events in different symbiont clades, which

offers a chance to study which aspects of reductive evolution are deterministic and which are

stochastic. For instance, the almost exclusive loss of *polA* in LvStB but no other *lga*-containing

symbionts except for LhHG_2 suggests that it is a driver of increased sequence drift in that

clade. In addition, the selectivity of genome reduction in this group of hosts amongst a complex

microbiome suggests a role of the host in selecting for genome reduction, which warrants

further study.

Methods

458 For full details see Supplementary Material.

Insect collection

Specimens were collected between 2009 and 2019 in Spain, Germany, Brazil, Japan and

Australia in the locations listed in **Table S3**. Female adults were dissected either directly after

chilling for ca. 15 min at −20°C or preserved in 70% ethanol or acetone until dissected. The

accessory glands were removed and preserved in 70% ethanol at −80°C until further

processing. For species in which we suspected the presence of symbiont-harbouring

compartments within the ovipositor in addition to the glands, the ovipositor was also dissected

and preserved along with the accessory glands.

Phylogeny of beetles

Mitogenomes were recovered from the Eukaryote kingdom bins from each respective sample.

Mitogenomes used by Wei *et al.* [27] in their phylogenetic tree were selected for references and

outgroups. Mitochondrial genomes (mitogenomes) from all metagenomic datasets and

reference mitogenomes were annotated using the MITOS2 webserver [68] against the RefSe89

Metazoan database, using genetic code 5 (invertebrate mitochondrial). Amino acid sequences

of the 13 protein coding genes (PCGs) from each mitogenome were collected and aligned using

muscle v5.1 [69]. Nucleic acid sequences of the corresponding PCGs were aligned using

pal2nal v14 [70] using the -codontable 5 flag. Nucleic acid alignments were concatenated and a

partition file was generated using the pxcat command from the phyx package [71]. Phylogenetic

analysis was performed by partitioning each codon position for each gene. Maximum likelihood

tree was made using RAxML 8.2.12 (raxmlHPC-PTHREADS-SSE3) [31], with the parameters -f

a -m GTRGAMMAI -# 1000 -p 1989 -x 1989. Alignment file in FASTA format was converted to

nexus format using Geneious Prime 2023.2.1 (www.geneious.com). Bayesian analysis was

performed using MrBayes v3.2 [32] with parameters lset applyto=(all) nst=6 rates=invgamma;

and unlink statefreq=(all) revmat=(all) shape=(all) pinvar=(all); using 600,000 generations and

sample frequency of 1,000. Phylogenetic trees were visualised and modified in the Interactive

Tree of Life server [72].

485 Lagriamide BGC phylogeny

Lagriamide BGC genes from *lgaA* to *lgaI* were extracted from the representative BGCs. Protein

sequences were aligned with muscle v5.1 [69] followed by alignment of DNA sequences using

pal2nal v14 [70] using -codontable 11. Subsequent steps were similar to those performed while

constructing beetle phylogeny. For RAxML phylogeny using GTRCAT -V, partitioning was only

performed per gene as it resulted in higher bootstrap values than partitioning for each codon

position in each gene.

492 *Burkholderia* symbiont phylogeny

Prokka [73] was used to annotate the ORFs of the genomes. Orthofinder v2.5.4 [61] was run on

amino acid sequences of the genomes. A custom script was used to extract the genes with

hierarchical orthogroups (HOGs) that are present in more than 90% of the genomes and are

only present once per genome (149 HOGs were selected). Muscle v5.1 [69] was used to align

the amino acid sequences of the selected HOGs, followed by pal2nal v14 [70] to align the

corresponding nucleic acid sequences using -codontable 11. Subsequent steps were similar to

those performed while constructing beetle phylogeny.

For symbiont phylogeny with *P. acidicola* ATCC 31363, 152 HOGs (identified using Orthofinder)

present only once and in more than 90% of the genomes were used. Subsequent steps were

similar to the symbiont phylogeny generated above were followed and the final tree was

generated using RAxML 8.2.12 (raxmlHPC-PTHREADS-SSE3) [31], with the parameters -f a -m

GTRGAMMAI -# 1000 -p 1989 -x 1989.

505 Data availability

515 Acknowledgments

516   We would like to thank Jean Keller for providing python scripts that helped in creating the

517   symbiont phylogeny. We are grateful to Rebekka Janke, Takema Fukatsu, Kiyoshi Ando, Kimio

518   Masumoto, Juan José López and Rolf Beutel for support with insect collection and to Domenica

519   Schnabelrauch for carrying out PCRs, cloning and Sanger sequencing.

520 Author Contributions

SU: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology,

Visualization, Writing – original draft, Writing – review & editing. SCW: Conceptualization, Data

curation, Formal Analysis, Investigation, Methodology, Visualization, Writing – original draft,

Writing – review & editing. AN: Data curation, Formal Analysis, Investigation, Methodology

HV: Data curation, Formal Analysis, Investigation, Methodology. LVF: Conceptualization,

Investigation, Writing – review & editing, Supervision, Funding acquisition.

MK: Formal Analysis, Methodology, Conceptualization, Investigation, Writing – review & editing,

Supervision, Funding acquisition. JCK: Conceptualization, Investigation, Writing – review &

editing, Supervision, Funding acquisition

530 Funding

531   This work was supported by the U.S. National Science Foundation [DBI-1845890 to J.C.K.,

532   S.U., and S.C.W.].

533 Competing Interests

534   The Kwan lab is planning to offer their metagenomic binning pipeline Autometa on the paid

535   bioinformatics and computational platform BatchX in addition to distributing it through open

536   source channels.

References

1.   Dyall SD, Brown MT, Johnson PJ. Ancient Invasions: From Endosymbionts to Organelles.
Science 2004; **304**: 253–257.

2.   Stephens TG, Gabr A, Calatrava V, Grossman AR, Bhattacharya D. Why is Primary
Endosymbiosis so Rare? *New Phytol* 2021; **231**: 1693–1699.

3.   Von Dohlen CD, Moran NA. Molecular Data Support a Rapid Radiation of Aphids in the
Cretaceous and Multiple Origins of Host Alternation. *Biol J Linn Soc Lond* 2000; **71**: 689–
717.

4.   Nováková E, Hypša V, Klein J, Foottit RG, von Dohlen CD, Moran NA. Reconstructing the
Phylogeny of Aphids (Hemiptera: Aphididae) Using DNA of the Obligate Symbiont
Buchnera aphidicola. Mol Phylogenet Evol 2013; **68**: 42–54.

5.   Chong RA, Park H, Moran NA. Genome Evolution of the Obligate Endosymbiont *Buchnera*
aphidicola. Mol Biol Evol 2019; **36**: 1481–1489.

6.   McCutcheon JP, Moran NA. Extreme Genome Reduction in Symbiotic Bacteria. *Nat Rev*
Microbiol 2011; **10**: 13–26.

7.   Mira A, Ochman H, Moran NA. Deletional Bias and the Evolution of Bacterial Genomes.
Trends Genet 2001; **17**: 589–596.

8.   Hershberg R, Petrov DA. Evidence that Mutation is Universally Biased Towards AT in
Bacteria. *PLoS Genet* 2010; **6**: e1001115.

9.   Dietel A-K, Merker H, Kaltenpoth M, Kost C. Selective Advantages Favour High Genomic
AT-Contents in Intracellular Elements. *PLoS Genet* 2019; **15**: e1007778.

10. Dufresne A, Garczarek L, Partensky F. Accelerated Evolution Associated with Genome
Reduction in a Free-Living Prokaryote. *Genome Biol* 2005; **6**: R14.

11. Morris JJ, Lenski RE, Zinser ER. The Black Queen Hypothesis: Evolution of Dependencies

Through Adaptive Gene Loss. *MBio* 2012; **3**: e00036–12.

12. Pinto-Carbó M, Gademann K, Eberl L, Carlier A. Leaf Nodule Symbiosis: Function and

Transmission of Obligate Bacterial Endophytes. *Curr Opin Plant Biol* 2018; **44**: 23–31.

13. Pinto-Carbó M, Sieber S, Dessein S, Wicker T, Verstraete B, Gademann K, et al. Evidence

of Horizontal Gene Transfer Between Obligate Leaf Nodule Symbionts. *ISME J* 2016; **10**:

2092–2105.

14. Donia MS, Fricke WF, Partensky F, Cox J, Elshahawi SI, White JR, et al. Complex

Microbiome Underlying Secondary and Primary Metabolism in the Tunicate-*Prochloron*

Symbiosis. *Proc Natl Acad Sci U S A* 2011; **108**: E1423–32.

15. Kwan JC, Donia MS, Han AW, Hirose E, Haygood MG, Schmidt EW. Genome Streamlining

and Chemical Defense in a Coral Reef Symbiosis. *Proc Natl Acad Sci U S A* 2012; **109**:

20655–20660.

16. Kwan JC, Schmidt EW. Bacterial Endosymbiosis in a Chordate Host: Long-Term Co-

Evolution and Conservation of Secondary Metabolism. *PLoS One* 2013; **8**: e80822.

17. Toh H, Weiss BL, Perkin SAH, Yamashita A, Oshima K, Hattori M, et al. Massive Genome

Erosion and Functional Adaptations Provide Insights Into the Symbiotic Lifestyle of *Sodalis*

glossinidius in the Tsetse Host. *Genome Res* 2006; **16**: 149–156.

18. Miller IJ, Vanee N, Fong SS, Lim-Fong GE, Kwan JC. Lack of Overt Genome Reduction in

the Bryostatin-Producing Bryozoan Symbiont, '*Candidatus* Endobugula sertula'. *Appl*

Environ Microbiol 2016; **82**: 6573–6583.

19. Linneman J, Paulus D, Lim-Fong G, Lopanik NB. Latitudinal Variation of a Defensive

Symbiosis in the *Bugula neritina* (Bryozoa) Sibling Species Complex. *PLoS One* 2014; **9**:

e108783.

20. Flórez LV, Scherlach K, Miller IJ, Rodrigues A, Kwan JC, Hertweck C, et al. An Antifungal

Polyketide Associated With Horizontally Acquired Genes Supports Symbiont-Mediated

Defense in *Lagria villosa* Beetles. *Nat Commun* 2018; **9**: 2478.

21. Janke RS, Kaftan F, Niehs SP, Scherlach K, Rodrigues A, Svatoš A, et al. Bacterial

Ectosymbionts in Cuticular Organs Chemically Protect a Beetle During Molting Stages.

ISME J 2022; **16**: 2691–2701.

22. Waterworth SC, Flórez LV, Rees ER, Hertweck C, Kaltenpoth M, Kwan JC. Horizontal

Gene Transfer to a Defensive Symbiont With a Reduced Genome in a Multipartite Beetle

Microbiome. *MBio* 2020; **11**: e02430–19.

23. Wierz JC, Gaube P, Klebsch D, Kaltenpoth M, Flórez LV. Transmission of Bacterial

Symbionts With and Without Genome Erosion Between a Beetle Host and the Plant

Environment. *Front Microbiol* 2021; **12**: 715601.

24. Jürgen Stammer H. Die Symbiose Der Lagriiden (Coleoptera). *Z Morph u Okol Tiere* 1929;

15: 1–34.

25. Flórez LV, Scherlach K, Gaube P, Ross C, Sitte E, Hermes C, et al. Antibiotic-Producing

Symbionts Dynamically Transition Between Plant Pathogenicity and Insect-Defensive

Mutualism. *Nat Commun* 2017; **8**: 15172.

26. Flórez LV, Biedermann PHW, Engl T, Kaltenpoth M. Defensive Symbioses of Animals With

Prokaryotic and Eukaryotic Microorganisms. *Nat Prod Rep* 2015; **32**: 904–936.

27. Wei Z, Shi A. The Complete Mitochondrial Genomes of Four Lagriine Species (Coleoptera,

Tenebrionidae) and Phylogenetic Relationships Within Tenebrionidae. *PeerJ* 2023; **11**:

e15483.

28. Wu C, Zhou Y, Tian T, Li T-J, Chen B. First Report of Complete Mitochondrial Genome in

the Subfamily Alleculinae and Mitochondrial Genome-Based Phylogenetics in

Tenebrionidae (Coleoptera: Tenebrionoidea). *Insect Sci* 2022; **29**: 1226–1238.

29. Ragionieri L, Zúñiga-Reinoso Á, Bläser M, Predel R. Phylogenomics of Darkling Beetles

(Coleoptera: Tenebrionidae) From the Atacama Desert. *PeerJ* 2023; **11**: e14848.

30. Kergoat GJ, Soldati L, Clamens A-L, Jourdan H, Jabbour-Zahab R, Genson G, et al. Higher

Level Molecular Phylogeny of Darkling Beetles (Coleoptera: Tenebrionidae). *Syst Entomol*

2014; **39**: 486–499.

31.   Stamatakis A. RAxML Version 8: A Tool for Phylogenetic Analysis and Post-analysis of

Large Phylogenies. *Bioinformatics* 2014; **30**: 1312–1313.

32.   Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, et al. MrBayes

3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model

Space. *Syst Biol* 2012; **61**: 539–542.

33.   Ratnasingham S, Hebert PDN. BOLD : The Barcode of Life Data System

(www.barcodinglife.org). *Mol Ecol Notes* 2007; **7**: 355–364.

34.   Hu Y, Fang L, Nicholson C, Wang K. Implications of Error-Prone Long-Read Whole-

Genome Shotgun Sequencing on Characterizing Reference Microbiomes. *iScience* 2020;

23: 101223.

35.   Piel J, Höfer I, Hui D. Evidence for a Symbiosis Island Involved in Horizontal Acquisition of

Pederin Biosynthetic Capabilities by the Bacterial Symbiont of *Paederus fuscipes* Beetles. *J*

Bacteriol 2004; **186**: 1280–1286.

36.   Kador M, Horn MA, Dettner K. Novel Oligonucleotide Probes for *in Situ* Detection of

Pederin-Producing Endosymbionts of *Paederus riparius* Rove Beetles (Coleoptera:

Staphylinidae). *FEMS Microbiol Lett* 2011; **319**: 73–81.

37.   Kellner RLL, Dettner K. Differential Efficacy of Toxic Pederin in Deterring Potential

Arthropod Predators of *Paederus* (Coleoptera: Staphylinidae) Offspring. *Oecologia* 1996;

107: 293–300.

38.   Witczak ZJ, Bommareddy A, VanWert AL. Pederin, Psymberin and the Structurally Related

Mycalamides: Synthetic Aspects and Biological Activities. In: Kim S-K (ed). *Handbook of*

Anticancer Drugs from Marine Origin. 2015. Springer, Cham, pp 683–700.

39.   Sakemi S, Ichiba T, Kohmoto S, Saucy G, Higa T. Isolation and Structure Elucidation of

Onnamide A, a New Bioactive Metabolite of a Marine Sponge, *Theonella* sp. *J Am Chem*

Soc 1988; **110**: 4851–4853.

40. Perry NB, Blunt JW, Munro MHG, Thompson AM. Antiviral and Antitumor Agents From a

New Zealand Sponge, Mycale sp. 2. Structures and Solution Conformations of

Mycalamides A and B. *J Org Chem* 1990; **55**: 223–227.

41. West LM, Northcote PT, Hood KA, Miller JH, Page MJ. Mycalamide D, a New Cytotoxic

Amide From the New Zealand Marine Sponge *Mycale* Species. *J Nat Prod* 2000; **63**: 707–

709.

42. Simpson JS, Garson MJ, Blunt JW, Munro MH, Hooper JN. Mycalamides C and D,

Cytotoxic Compounds From the Marine Sponge *Stylinos* n. Species. *J Nat Prod* 2000; **63**:

704–706.

43. Dyshlovoy SA, Fedorov SN, Kalinovsky AI, Shubina LK, Bokemeyer C, Stonik VA, et al.

Mycalamide A Shows Cytotoxic Properties and Prevents EGF-Induced Neoplastic

Transformation Through Inhibition of Nuclear Factors. *Mar Drugs* 2012; **10**: 1212–1224.

44. Gouiffes D, Juge M, Grimaud N, Welin L, Sauviat MP, Barbin Y, et al. Bistramide A, a New

Toxin From the Urochordata *Lissoclinum bistratum* Sluiter: Isolation and Preliminary

Characterization. *Toxicon* 1988; **26**: 1129–1136.

45. Nechitaylo TY, Sandoval-Calderón M, Engl T, Wielsch N, Dunn DM, Goesmann A, et al.

Incipient Genome Erosion and Metabolic Streamlining for Antibiotic Production in a

Defensive Symbiont. *Proc Natl Acad Sci U S A* 2021; **118**: e2023047118.

46. Engl T, Kroiss J, Kai M, Nechitaylo TY, Svatoš A, Kaltenpoth M. Evolutionary Stability of

Antibiotic Protection in a Defensive Symbiosis. *Proc Natl Acad Sci U S A* 2018; **115**:

E2020–E2029.

47. Hayakawa Y, Shirasaki S, Shiba S, Kawasaki T, Matsuo Y, Adachi K, et al. Piericidins C_7

and C_8, New Cytotoxic Antibiotics Produced by a Marine *Streptomyces* sp. *J Antibiot* 2007;

60: 196–200.

48. Hayakawa Y, Shirasaki S, Kawasaki T, Matsuo Y, Adachi K, Shizuri Y. Structures of New

Cytotoxic Antibiotics, Piericidins C_7 and C_8. *J Antibiot* 2007; **60**: 201–203.

49. Watabe H. A New Antibiotic SF2583A, 4-chloro-5-(3'-indolyl)oxazole, Produced by

Streptomyces. Meiji Seika Kenkyu Nenpo 1988; **27**: 55–62.

50. Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. CheckM2: A Rapid, Scalable and

Accurate Tool for Assessing Microbial Genome Quality Using Machine Learning. *Nat*

Methods 2023; **20**: 1203–1212.

51. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et al.

Minimum Information About a Single Amplified Genome (MISAG) and a Metagenome-

Assembled Genome (MIMAG) of Bacteria and Archaea. *Nat Biotechnol* 2017; **35**: 725–731.

52. Manzano-Marín A, Kvist S, Oceguera-Figueroa A. Evolution of an Alternative Genetic Code

in the Providencia Symbiont of the Hematophagous Leech *Haementeria acuecueyetzin.*

Genome Biol Evol 2023; **15**: evad164.

53. Mahenthiralingam E, Urban TA, Goldberg JB. The Multifarious, Multireplicon *Burkholderia*

cepacia complex. *Nat Rev Microbiol* 2005; **3**: 144–156.

54. Lessie TG, Hendrickson W, Manning BD, Devereux R. Genomic Complexity and Plasticity

of *Burkholderia cepacia. FEMS Microbiol Lett* 1996; **144**: 117–128.

55. Egan ES, Fogel MA, Waldor MK. Divided Genomes: Negotiating the Cell Cycle in

Prokaryotes With Multiple Chromosomes. *Mol Microbiol* 2005; **56**: 1129–1138.

56. Flórez LV, Kaltenpoth M. Symbiont Dynamics and Strain Diversity in the Defensive

Mutualism Between *Lagria* Beetles and *Burkholderia. Environ Microbiol* 2017; **19**: 3674–

3688.

57. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. DNA-

DNA Hybridization Values and Their Relationship to Whole-Genome Sequence Similarities.

Int J Syst Evol Microbiol 2007; **57**: 81–91.

58. Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC, et

al. Microbial Species Delineation Using Whole Genome Sequences. *Nucleic Acids Res*

2015; **43**: 6761–6771.

691 59. Konstantinidis KT, Tiedje JM. Genomic Insights That Advance the Species Definition for
692   Prokaryotes. *Proc Natl Acad Sci U S A* 2005; **102**: 2567–2572.

693 60. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. GTDB: An
694   Ongoing Census of Bacterial and Archaeal Diversity Through a Phylogenetically
695   Consistent, Rank Normalized and Complete Genome-Based Taxonomy. *Nucleic Acids Res*
696   2022; **50**: D785–D794.

697 61. Emms DM, Kelly S. OrthoFinder: Phylogenetic Orthology Inference for Comparative
698   Genomics. *Genome Biol* 2019; **20**: 238.

699 62. Fergusson CH, Saulog J, Paulo BS, Wilson DM, Liu DY, Morehouse NJ, et al. Discovery of
700   the Polyketide Lagriamide B by Integrated Genome Mining, Isotopic Labeling, and
701   Untargeted Metabolomics. *ChemRxiv* 2023.

702 63. Lo W-S, Huang Y-Y, Kuo C-H. Winding Paths to Simplicity: Genome Evolution in
703   Facultative Insect Symbionts. *FEMS Microbiol Rev* 2016; **40**: 855–874.

704 64. Sudakaran S, Kost C, Kaltenpoth M. Symbiont Acquisition and Replacement as a Source of
705   Ecological Innovation. *Trends Microbiol* 2017; **25**: 375–390.

706 65. Janke RS, Moog S, Weiss B, Kaltenpoth M, Flórez LV. Morphological Adaptation for
707   Ectosymbiont Maintenance and Transmission During Metamorphosis in *Lagria* Beetles.
708   *Front Physiol* 2022; **13**: 979200.

709 66. Patel PH, Suzuki M, Adman E, Shinkai A, Loeb LA. Prokaryotic DNA Polymerase I:
710   Evolution, Structure, and 'Base Flipping' Mechanism for Nucleotide Selection. *J Mol Biol*
711   2001; **308**: 823–837.

712 67. Bennett GM, Moran NA. Heritable Symbiosis: The Advantages and Perils of an
713   Evolutionary Rabbit Hole. *Proc Natl Acad Sci U S A* 2015; **112**: 10169–10176.

714 68. Donath A, Jühling F, Al-Arab M, Bernhart SH, Reinhardt F, Stadler PF, et al. Improved
715   Annotation of Protein-Coding Genes Boundaries in Metazoan Mitochondrial Genomes.
716   *Nucleic Acids Res* 2019; **47**: 10543–10552.

69. Edgar RC. Muscle5: High-Accuracy Alignment Ensembles Enable Unbiased Assessments

of Sequence Homology and Phylogeny. *Nat Commun* 2022; **13**: 6968.

70. Suyama M, Torrents D, Bork P. PAL2NAL: Robust Conversion of Protein Sequence

Alignments Into the Corresponding Codon Alignments. *Nucleic Acids Res* 2006; **34**: W609–

W612.

71. Brown JW, Walker JF, Smith SA. Phyx: Phylogenetic Tools for Unix. *Bioinformatics* 2017;

33: 1886–1888.

72. Letunic I, Bork P. Interactive Tree of Life (iTOL) v5: An Online Tool for Phylogenetic Tree

Display and Annotation. *Nucleic Acids Res* 2021; **49**: W293–W296.

73. Seemann T. Prokka: Rapid Prokaryotic Genome Annotation. *Bioinformatics* 2014; **30**:

2068–2069.

Table 1. Metadata for different beetles collected for this study. *Lagria villosa samples were

collected three times at different time points. With one sample reported in a previous study

(referred to as Lv19 in this work) [20, 22].

Sample	Location	Abbreviation(s) used in the paper
Lagria villosa*	São Paulo, Brazil	Lv19, Lv20, Lv23
Lagria rufipennis	Osaka and Ibaraki, Japan	Lruf
Lagria okinawana	Okinawa, Japan	Loki
Lagria nigricollis	Tokushima, Osaka, and Kogashima, Japan	Lnig

Lagria hirta	Hessen, Germany	LhSB
Lagria hirta	Rhineland Palatinate, Germany	LhHG
Lagria hirta	Galicia, Spain	LhG
Lagria grenieri	Huelva, Spain	Lgren
Lagria atripes	Rhineland-Palatinate, Germany	Latri
Ecnolagria sp.	New South Wales, Australia	Ecno

Figure 1. Beetle mitogenome phylogenetic tree using 13 mitochondrial protein coding genes

constructed using MrBayes [32]. Branch values represent posterior probabilities. Mitogenomes

recovered in this study are highlighted with bold lettering.

Figure 2. Analysis of representative *lga* BGCs extracted from eleven Lagriinae beetle

metagenomes. A) Comparison of representative *lga* BGC gene organisation. Individual genes in

the *lga* BGCs are represented by arrows oriented in the predicted direction of transcription and

coloured according to identity. Pairwise amino acid similarity between BGCs is indicated in the

shaded areas between genes. A scale bar is provided for gene size. B) Comparison of predicted

enzyme domain organisation in the representative *lga* BGCs, where genes are ordered

according to biosynthetic order. Boxes around the domains indicate differences between the

BGCs.

Fig 3. A) Circular representation of LvStB_2023 genome. B) Raw count of COG categories

present on different contigs of LvStB_2023 (with and without pseudogenes).

Fig 4. Bayesian phylogenetic tree of Lagriinae beetle associated *Burkholderia* symbionts.

Values on nodes indicate posterior probabilities. Outgroups include - *Paraburkholderia*

acidiphila (GCF_009789655.1), *Cupriavidus necator* (GCF_000219215.1), *Herbaspirillum*

seropedicae (GCF_001040945.1).

Fig 5. Congruence between phylogenies of beetle host, *Burkholdria* symbionts and *lga* BGCs in

all samples. A) Tanglegram between *lga*-carrying symbionts and beetle host phylogeny. B)

Tanglegram between *lga*-carrying symbionts (centre) and the *lga* BGC, as inferred via two

models GTRCAT (left) and GTRGAMMAI (right). C) Tanglegram of the BGC phylogenies using

the GTRCAT (left) and GTRGAMMAI (right) models relative to the phylogeny of the host beetle

based on the respective mitogenome sequences (centre). Clades within the beetle host

mitogenome phylogenetic tree are highlighted in shades of grey. In all panels, the three

conserved clades are highlighted in purple, green and orange. Dashed brown lines represent

nodes that are unique between the respective phylogenies, except when the node is absent

from the other tree.

Tree scale: 1

Oedemera virescens | HQ232826	
Mylabris sp. | JX412732	
Asbolus verrucosus | NC_027256	
Machla sappho | MZ342778	
Philolithus aegrotus | MZ342784	
Pelecyphorus foveolatus | MZ342781	
Pelecyphorus contortus | MZ342780	
Stenomorpha consobrina | MZ342785	
Stenomorpha obovata | MZ342786	Pimeliinae
Adelium sp. | FJ613422	
Luprops yunnanus | ON303728	
Spinolyprops cribricollis | ON303729	
Anaedus unidentatus | ON303730	
Impressosora sp. | JX412754	
Eutrapela ruficollis | HQ232805	
Paratenetus tropicalis | JX412774	
Cerogria janthinipennis | ON303727	
Cerogria popularis | NC_061196	
Ecnolagria sp. (Ecno) | Australia	
Chrysolagria sp. | JX412760	
Lagria villosa (Lv20) | Brazil	
Lagria grenieri (Lgren) | Spain	
Lagria atripes (Latri) | Germany	
Lagria hirta (LhG) | Spain	
Lagria hirta (LhSB) | Germany	
Lagria hirta (LhGH) | Germany	
Lagria hirta | OX375806	
Lagria okinawana (Loki) | Japan	
Lagria rufipennis | MW802588	
Lagria rufipennis (Lruf) | Japan	
Lagria nigricollis (Lnig) | Japan	Lagriinae
Crypsis chinensis | ON303731	
Scaphidema metallicum | KX087341	Diaperinae
Tenebrio obscurus | MG739327	
Tenebrio molitor | NC_024633	
Tribolium castaneum | NC_003081	
Tribolium confusum | NC_026702	Tenebrioninae
Blaps rhynchoptera | MN267802	
Gonocephalum sp. | KU236386	
Opatrum sabulosum | MN745102	
Gonocephalum kochi | MW822744	Blaptinae
Platydema sp. | JX412842	
Alphitobius diaperinus | MT610905	
Promethis valgipes | MW201671	
Strongylium suspicax | JX412780	
Morphostenophanes yunnanus | MZ298928	
Morphostenophanes yunnanus | MW822745	Stenochiinae
Ulomoides dermestoides | KM046492	
Paramarygmus sp. | JX412808	
Amarygmini sp. | MH789725	
Zophobas atratus | MK140669	Tenebrioninae
Cteniopus sp. | KX087267	
Alleculinae sp. | JX412818	Alleculinae

B.arboris Lv20_1 (8.35 Mbp) | Brazil

B.arboris BCC0049 (8.28 Mbp) | GCF_902830425.1

B.lata 383 (8.67 Mbp) | GCF_000012945.1

B.lata Lv19_4_0 (8.81 Mbp) | Brazil

Burkholderia sp. Lv19_6_14 (1.69 Mbp) | Brazil

B. gladioli Lv19_6_18 (1.62 Mbp) | Brazil

B. gladioli Lv20_2 (2.04 Mbp) | Brazil

Burkholderia sp. Lgren_6 (1.14 Mbp) | Spain

Burkholderia sp. Latri_1 (2.03 Mbp) | Germany

Burkholderia sp. Ecno_3 (3.15 Mbp) | Australia

Burkholderia sp. Loki_2 (2.13 Mbp) | Japan

Burkholderia sp. LhG_1 (2.70 Mbp) | Spain

Burkholderia sp. LhHG_2 (1.14 Mbp) | Germany

Burkholderia sp. LhSB_1 (1.17 Mbp) | Germany

Burkholderia sp. Lv20_9 (1.88 Mbp) | Brazil

Burkholderia sp. LvStB_2023 (2.50 Mbp) | Brazil

Burkholderia sp. LvStB (2.07 Mbp) | Brazil

B. gladioli Lruf_1 (1.93 Mbp) | Japan

B. gladioli Lnig_2 (2.64 Mbp) | Japan

B. gladioli A1 (8.11 Mbp) | GCF_000633145.1

B. gladioli Lgren_7 (1.79 Mbp) | Spain

B. gladioli CHAPALOTE (8.52 Mbp) | GCF_000757585.1

B. gladioli MSMB1756 (8.19 Mbp) | GCF_001527485.1

B. gladioli BSR3 (9.05 Mbp) | GCF_000194745.1

B. gladioli NBRC 13700 (8.76 Mbp) | GCF_000739755.1

B. gladioli ATCC 10248 (8.89 Mbp) | GCF_000959725.1

B. gladioli SN82F6 (8.45 Mbp) | GCF_000981885.1

B. gladioli LvStA (8.56 Mbp) | Brazil

B. gladioli Latri_2 (7.58 Mbp) | Germany

B. gladioli LhStG (8.40 Mbp) | Germany

B. gladioli LhSB_5 (7.71 Mbp) | Germany

B. gladioli LhHG_3 (5.09 Mbp) | Germany

Outgroup

lga producer

Non producer

Culturable

Tree scale: 0.1

A

B

C