1	Full Title:	Genome-Wide Profiling of Soybean WRINKLED1 Transcription Factor
2		Binding Sites Provides Insight into Seed Storage Lipid Biosynthesis
3		
4	Authors:	Leonardo Jo[a,1,2,3], Julie M. Pelletier[a,1], Robert B. Goldberg[b,3], John J.
5		Harada[a,3]
6		
7	Affiliations:	[a]Department of Plant Biology, College of Biological Sciences, University
8		of California, Davis, CA 95616, USA, and [b]Department of Molecular,
9		Cell, and Developmental Biology, University of California, Los Angeles,
10		CA 90095
11		
12	Footnotes:	[1]These authors contributed equally to this work
13		[2]Current address, Experimental and Computational Plant Development,
14		Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands
15		[3]To whom correspondence should be addressed. Email: John J. Harada,
16		Department of Plant Biology, 605 Hutchison Drive, University of
17		California, Davis, CA 95616, USA, (530) 752-0673,
18		jjharada@ucdavis.edu; Leonardo Jo, Experimental and Computational
19		Plant Development, Utrecht University, Padualaan 8, 3584 CH Utrecht,
20		The Netherlands, l.jo@uu.nl; and Robert B. Goldberg, Department of
21		Molecular, Cell, and Developmental Biology, 4121 Terasaki Research
22		Building, University of California, Los Angeles, CA 90095, (310) 825-
23		9093, bobg@g.ucla.edu
24		

25	Preprint Server:	biorxiv.org,
26		https://www.biorxiv.org/content/10.1101/2024.01.23.576967v1
27		
28	Classification:	Major - Biological Sciences, Minor - Plant Biology
29		
30	Keywords:	fatty acids, triacylglycerol, embryo
31		

Abstract

Understanding the regulatory mechanisms controlling storage lipid accumulation will inform strategies to enhance seed oil quality and quantity in crop plants. The WRINKLED1 transcription factor (WRI1 TF) is a central regulator of lipid biosynthesis. We characterized the genome-wide binding profile of soybean (Gm)WRI1 and show that the TF directly regulates genes encoding numerous enzymes and proteins in the fatty acid and triacylglycerol biosynthetic pathways. GmWRI1 binds primarily to regions downstream of target gene transcription start sites. We showed that GmWRI1 bound regions are enriched for the canonical WRI1 DNA binding element, the AW Box (CNTNGNNNNNNNCG), and another DNA motif, the CNC Box (CNCCNCC). Functional assays showed that both DNA elements mediate transcriptional activation by GmWRI1. We also show that GmWRI1 works in concert with other TFs to establish a regulatory state that promotes fatty acid and triacylglycerol biosynthesis. In particular, comparison of genes targeted directly by GmWRI1 and by GmLEC1, a central regulator of the maturation phase of seed development, reveals that the two TFs act in a positive feedback subcircuit to control fatty acid and triacylglycerol biosynthesis. Together, our results provide new insights into the genetic circuitry in which GmWRI1 participates to regulate storage lipid accumulation during seed development.

Significance Statement

We report the genome-wide profiling of DNA sequences bound by and the genes directly-regulated by soybean WRINKLED1, a central regulator of storage lipid accumulation in oilseed plants. The information offers new insights into the mechanisms by which WRINKLED1 regulates genes encoding lipid biosynthetic enzymes and establishes a regulatory environment that promotes oil accumulation, and it may aid in the design of strategy to alter storage lipid accumulation in oilseeds.

Introduction

Oils derived from plants are used primarily for food-based applications, although they also have broad utility in industrial applications (1). Given that lipids are essential for human nutrition and that biofuels are of increasing importance as a sustainable energy source, the demand for plant oils is anticipated to increase. Because oilseeds account for approximately 55% of world vegetable oil production (2), developing crops with increased and/or modified seed oil content may be instrumental in meeting these demands. To achieve this goal, it is crucial that we understand the regulatory mechanisms that underlie seed storage lipid accumulation to inform the development of biotechnological tools that will improve the quality and quantity of seed oils.

Fatty acids (FAs) are stored primarily in the form of triacylglycerol (TAG) in seeds. As summarized in Fig. 1A, the FA biosynthetic pathway in seeds occurs in plastids (3, 4). The first committed and rate-limiting step in FA biosynthesis is catalyzed by the heterotetrameric acetyl-CoA carboxylase (ACCase) enzyme that produces malonyl-CoA. Subsequent steps are catalyzed by fatty acid synthase, a multisubunit complex consisting of 10 distinct enzymes activities, that catalyzes the transfer of malonyl-CoA to acyl carrier protein (ACP), the sequential elongation of the acyl-ACP chain, and the release of non-esterified C_{16} or C_{18} fatty acids. In addition to the FA biosynthetic enzymes, several other enzymes responsible for the elongation and desaturation of FAs account for the diversity of FAs found in the seeds (5).

TAG synthesis occurs in the endoplasmic reticulum through one of two pathways (6, 7). The acyl CoA-dependent pathway, known as the Kennedy pathway, catalyzes the sequential addition of fatty acid moieties to a glycerol-3-phosphate backbone to generate TAGs. In the alternative pathway, the acyl-CoA-independent pathway which predominates in soybean, fatty acids are first incorporated into the membrane lipid phosphatidylcholine prior to their incorporation into TAG. TAGs are stored largely in oil bodies that bud off from the endoplasmic reticulum and are surrounded by a phospholipid monolayer (8). Storage lipid and protein accumulation occurs primarily during the maturation phase of seed development (Fig. 1B) which follows the morphogenesis phase during which embryonic domains and tissues are established (9).

Thus, storage lipid accumulation in seeds requires the coordinated activity of many

diverse enzymes that operate in distinct cellular compartments. Oilseed crops are particularly

efficient in FA and TAG s biosynthesis, with storage lipids accounting for more than 60% of

seed dry mass (10). This massive accumulation of TAGs and the observation that FA

biosynthesis genes show similar patterns of expression that peaks during the onset of the

maturation program and declines during the later stages of maturation suggests that FA and TAG

biosynthesis is precisely regulated during seed development (11). Several reports indicate that

FA biosynthesis is regulated primarily at the transcriptional level (12-15).

The major regulator of seed storage lipid biosynthesis is the WRINKLED1 (WRI1)

transcription factor (TF), a member of the APETALA2 TF family (16). WRI1 has been shown

to be both necessary and sufficient for FA accumulation in many oilseed crops. For example,

Arabidopsis (*at*)*wri1* loss-of-function mutants exhibit an 80% reduction in seed oil content (17),

and ectopic expression of *WRI1* results in significant increases of FA-related gene expression

and, consequently, increased oil content in several plant species (16, 18-22).

WRI1 affects storage lipid accumulation by regulating the expression of genes involved

in FA biosynthesis. Experimental modulations of *WRI1* expression that alters seed oil levels

cause corresponding changes in the expression of FA biosynthesis genes, including those

encoding ACP, 3-ketoacyl-ACP synthases (KAS), and subunits of ACCase, suggesting that these

genes are activated transcriptionally by WRI1 (19, 23, 24). Consistent with this hypothesis, the

AW Box (CNTNGNNNNNNNCG), a DNA element bound by WRI1 *in vitro* (23), is enriched in

FA biosynthesis genes.

Despite its key regulatory role, a detailed, mechanistic understanding of how WRI1

regulates storage lipid accumulation in seeds is lacking. In particular, a comprehensive, genome-

wide description of all genes regulated by WRI1 has not been reported, limiting our

understanding of the exact mechanism by which WRI1 acts to regulate seed oil accumulation.

To address this limitation, we analyzed the genome-wide binding profile of soybean WRI1

(GmWRI1) during seed development and provide the first comprehensive description of WRI1

binding sites and the genes directly regulated by WRI1 in plants. Our results provide new

insights into the diverse mechanisms by which WRI1 acts as a central regulator of storage lipid

accumulation in seeds.

Results

GmWRI1 Binds and Regulates Genes Involved in Fatty Acid Biosynthesis

We identified genes that are directly regulated by GmWRI1-A (Glyma.08G227700) and

GmWRI1-B (Glyma.15G221600). Although there are several *WRI1* homologs in soybean,

GmWRI1-A and *GmWRI1-B* showed the highest degree of DNA sequence similarity with

AtWRI1, and other have demonstrated their ability to rescue the *atwri1* seed oil mutant

phenotype in Arabidopsis (25). As summarized in Fig. 1B, data from existing RNA-Sequence

(RNA-Seq) databases (GEO accession GSE99571) showed that both mRNAs are present at

high levels in embryos at the cotyledon (cot) and early maturation (em) stages and decline at

later stages of embryo development, preceding the onset of FA accumulation in soybean seeds.

Analysis of the spatial distribution of *GmWRI1* mRNAs in em stage seeds (GEO accession

GSE46096) showed that both accumulated predominantly in embryo subregions (Fig. 1C). The

temporal and spatial expression patterns of *GmWRI1-A* and *GmWRI1-B* are consistent with their

roles in controlling the FA and TAG biosynthetic programs in embryos.

141   To identify genes that are directly regulated by GmWRI1 during seed development, we

first defined genes that were bound by the TFs. We performed chromatin immunoprecipitation

followed by DNA sequencing (ChIP-Seq) experiments using em stage soybean embryos with

antibodies that recognize both GmWRI1-A and GmWRI1-B (see Experimental Procedures).

GmWRI1 bound genes were defined as those with reproducible GmWRI1 ChIP-Seq peaks in a

genomic region between 1 kb upstream of the gene transcription start site (TSS) to the

transcription termination site. We identified 11,203 bound genes (Dataset S1). We also

performed GmWRI1 ChIP-Seq experiments with cotyledon (cot)-stage embryos, because

GmWRI1 mRNA accumulation is high at that stage (Fig. 1B). However, the cot-stage results did

not pass ENCODE quality thresholds for ChIP-Seq data unlike experiments at the em stage

(26)(Dataset S2), as indicated by the smaller number of bound genes that were detected at the cot

stage compared with the em stage (SI Appendix Fig. S1A, Dataset S1). Nevertheless, virtually

all cot-stage GmWRI1 bound genes were also bound at the em stage (SI Appendix Fig. S1A).

Because most TF binding interactions do not represent transcriptional regulatory events

(27), we filtered GmWRI1 bound genes for those that are co-expressed with the TF. The

rationale is that a TF and the genes that it directly regulates are expressed in the same cell.

Given the *GmWRI1* mRNA spatial accumulation pattern, we defined co-expressed genes as those

whose mRNA levels were at least two-fold or higher in embryo cotyledon parenchymal tissues

compared to seed coat parenchyma and seed coat hilum tissues (FDR < 0.05, Harada–Goldberg

Soybean Seed Development LCM RNA-Seq Dataset, GEO accession GSE99109; Dataset S1).

As shown in Fig. 1D, we identified 2,155 genes that were both bound by GmWRI1 and co-

expressed with *GmWRI1*, a significant overlap ($P < 3.4e\text{-}58$), and we designated these as

GmWRI1 direct target (DT) genes.

We compared GmWRI1 DT genes with Arabidopsis genes regulated by AtWRI1 to

confirm that the soybean genes were WRI1 regulated. Because neither genes bound by AtWRI1

nor genes mis-regulated by *atwri1* mutations have been reported genome-wide, we identified

genes that were differentially expressed (DE) in *atwri1-1* mutant embryos at the linear-cotyledon

stage (Fig. 2A and S2). We found that 1,213 DE genes were downregulated (two-fold, FDR <

0.05) in *atwri1-1* compared with wild-type (WT) embryos, 848 of which correspond to annotated

soybean genes (SI Appendix Fig. S3, Dataset S3). A significant number of these genes, 165 (P <

5.8e-14), overlapped with Arabidopsis homologs of GmWRI1 DT genes (1,690, Fig. 2C). To

define Arabidopsis genes upregulated by ectopic *GmWRI1* expression, we transfected

Arabidopsis leaf protoplasts with either the *GmWRI1-A* gene driven by the *CaMV35S* promoter

(*p35S:GmWRI1*) or the *p35S:mCHERRY* gene (Fig. 2B). We identified 1,490 DE genes that

were upregulated (two-fold, FDR < 0.05) in *35S:GmWRI1* versus *35S:MCHERRY* cells, 1097 of

which correspond to annotated soybean genes (SI Appendix Fig. S3, Dataset S3). We found that

212 of these genes overlapped significantly with Arabidopsis homologs of GmWRI1 DT genes

(P < 2.0e-17, Fig. 2D). Together, these results suggest that GmWRI1 regulates GmWRI1 DT

genes. We designated the 401 genes (p < 1.7e-214) that are downregulated in *atwri1* mutant

embryos and upregulated by *35S:GmWRI1* overexpression in leaf cells as AtWRI1 directly

regulated (AtWRI DR) genes (Fig. 2E).

Two lines of evidence suggest that GmWRI1 DT genes are involved in FA and TAG

biosynthesis. First, GmWRI1 DT genes were overrepresented for Gene Ontology (GO) terms

involved with FA biosynthesis, such as fatty acid biosynthetic process and acetyl-CoA metabolic

process, acetyl-CoA carboxylase activity, and fatty acid synthase complex (Fig. 1F, Dataset S1).

Similarly, AtWRI DR genes were enriched for GO terms related to FA biosynthetic processes,

such as fatty acid biosynthetic process, acyl carrier activity, and acetyl-CoA carboxylase activity

(Fig. 2F, Dataset S3). GmWRI1 DT genes were also enriched for GO terms related to cell

proliferation, such as cell population proliferation and cytokinesis by cell plate formation,

opening the possibility that GmWRI1 may also regulate genes involved in processes other than

storage lipid accumulation. Second, Fig. 1A shows that the vast majority of enzymes involved in

FA and TAG biosynthesis are encoded by GmWRI1 DT genes, such as the ACCase subunits,

biotin carboxylase (CAC2) and biotin carboxyl carrier protein (BCCP2), and KASIII, and in

TAG assembly in the endoplasmic reticulum, such as phosphatidic acid phosphohydrolase

(PAH), diacylglycerol acyltransferase (DGAT), and phosphatidylcholine diacylglycerol

cholinephosphotransferase (PDCT). We note that two genes are bound by but not coexpressed

with GmWRI1. To determine the extent to which GmWRI1 regulates genes involved in FA and

TAG biosynthesis, we manually curated a list of FA and TAG (FAT) biosynthesis genes based

on the Plant Metabolic Network (PMN) (https://plantcyc.org/databases/soycyc/8.0) and used the

Arabidopsis Acyl-Lipid Metabolism (Aralip) (http://aralip.plantbiology.msu.edu/) to annotate

these genes (Dataset S4). Of the 826 soybean FAT genes that encode different steps in the FA

and TAG biosynthetic pathways, 111 were GmWRI1 DT genes ($P < 1.8$ e-30, Fig. 1E).

Together, our results confirm that GmWRI1 directly regulates genes required for storage lipid

biosynthesis.

WRI1 Primarily Binds Intragenic Regions Enriched for Specific DNA Motifs

To gain insight into the mechanisms by which GmWRI1 regulates FA and TAG biosynthetic

genes transcriptionally, we analyzed bound regions in GmWRI1 DT genes. Surprisingly, we

showed that the ChIP-Seq peaks which presumably represent GmWRI1 binding sites in cot and

em stage embryos were mostly located downstream of the TSS in intragenic regions (Fig. 3, A

and B, and S1b). This binding site profile differed from those of GmLEC1, GmAREB3,

GmbZIP67, and GmABI3 TFs in soybean embryos whose ChIP-Seq peaks were mostly located

in region upstream of the TSS (28, 29).

AW Box DNA motif is highly enriched in GmWRI1 binding sites. We asked if DNA motifs

were enriched in a 100 bp region surrounding the ChIP-Seq peaks, designated as GmWRI1

binding sites. One motif identified using *de novo* discovery analysis of GmWRI1 DT binding

sites was the AW box motif, a DNA sequence previously shown to be bound by AtWRI1 *in vitro*

(23)(Fig. 3C, Dataset S5). We showed that GmWRI1 DT and FAT GmWRI1 DT genes were

significantly enriched for the AW Box (Bonferroni-adjusted *P* values < 0.01), with 35.2% and

56.3% of binding sites, respectively, containing full AW Boxes (Fig. 3D). These results suggest

that the AW Box underlies the direct interaction of GmWRI1 with many, but not all, of its

binding sites.

Because most GmWRI1 binding sites were located downstream of the TSS, we

determined AW Box distribution in GmWRI1 DT and FAT GmWRI1 DT genes. AW Box

motifs were found predominantly in regions downstream of the TSS, consistent with GmWRI1

binding site locations (SI Appendix Fig. S4, A and B). We note, however, that AW Box

enrichment may be influenced by the higher GC content of regions downstream versus upstream

of the TSS. Nevertheless, AW Box distribution and enrichment in GmWRI1 binding sites is

consistent with a role for this DNA motif in specifying GmWRI1 occupancy in downstream

regions.

Additional DNA motifs are enriched in WRI1 binding sites. *De novo* discovery analyses

identified another DNA motif, CNCCNCC (designated CNC Box), that was enriched in the

binding sites of GmWRI1 DT and FAT GmWRI1 DT genes (Fig. 3, C and D, Dataset S5).

Although the CNC Box has not been shown to be bound by WRI1, its enrichment in GmWRI1

binding sites suggests its potential role in mediating GmWRI1 DT gene transcription. Consistent

with this hypothesis, a similar motif was previously shown to be enriched in the 5' UTR of FA-

related genes in numerous plant species (30).

A GAGA-rich DNA motif was also discovered *de novo* in GmWRI1 binding sites (Fig. 3,

B and C). This motif is similar to the BPC1 motif that is bound by the BASIC

PENTACYSTEINE6 (BPC6) transcription factor which is proposed to recruit the Polycomb

Repressive Complex (31). The BPC1 motif is found in the binding sites of many TFs that

regulate the maturation phase of seed development, such as GmLEC1, GmAREB3, GmbZIP67

and GmABI3 (28, 29). Together, these results suggest that transcriptional regulation of

GmWRI1 DT genes may involve DNA elements other than the well characterized AW Box.

Because FAs and TAGs are synthesized during the maturation phase, we asked whether

the CCAAT Box (CCAAT), G Box (CACGTG) and RY (CATGCA) DNA elements that are

known to be bound by the major maturation TF regulators, LEC1, bZIP67, ABI3, FUS3, and

LEC2, were enriched in GmWRI1 binding sites (28, 32). These motifs were not identified by *de*

novo discovery analysis, and only the G Box motif was significantly enriched in 3.3% and 5.0%

of GmWRI1 DT binding sites and FAT GmWRI1 DT binding sites (Fig. 3D and Dataset S5).

This result suggests that GmWRI1-mediated transcriptional control of FA biosynthesis involves

distinct cis-regulatory modules from those utilized by these other maturation TFs.

AW and CNC Box motifs are enriched in intragenic regions of AtWRI Directly-Regulated

genes. We analyzed motif enrichment in AtWRI DR genes to determine if the motifs identified

in GmWRI1 binding sites are conserved in Arabidopsis. AW Boxes and CNC Boxes were

detected downstream of the TSS in 70.3% and 59.6%, respectively, of AtWRI DR genes (Fig.

S5, Dataset S3). By contrast, the AW Boxes and CNC Boxes were detected in only 41% and

48.9%, respectively, of the downstream regions of 1000 sets of randomly matched genes

(Dataset S3). In regions upstream of the TSS, the AW Box was enriched in 33.2% of AtWRI1

DR genes as compared with 18.5% in background regions (Fig. S5, Dataset S3), but no

significant enrichment of the CNC Box was observed. These results are consistent with the

observation that the AW Box and CNC Box are overrepresented in binding sites downstream of

the TSS of genes regulated by WRI1, suggesting that these elements contribute to the

transcriptional regulation of these genes.

Functional Analysis of AW Box and CNC Box Motifs in GmWRI1 Direct Target Gene

Binding Regions

We conducted transient assays to determine whether enriched AW Boxes and CNC Boxes are required for GmWRI1-mediated transcription of the four FAT genes shown in Fig. 4A: *ACP4* (Glyma.15G098500), *BCCP2* (Glyma.13G0 57400), *CAC2* (Glyma.05G221100), and *ACP3* (Glyma.19G240100). We cloned WT versions of the genes' binding sites and mutant variants in which all AW Box (mAW), CNC Box (mCCNCC) or both (mAWmCNC) motifs were mutated (Fig. 4B, Dataset S6). Binding sites for the *ACP4*, *BCCP2,* and *CAC2* genes contained at least one full AW Box motif, whereas *ACP3* contained an abbreviated version of the AW Box (CNTNGNNNNNNCG) which was shown by others to be bound by AtWRI1 *in vitro* (33). With the exception of *BCCP2,* the binding sites all contained at least one CNC Box. Binding sites were fused upstream of a firefly *Luciferase* gene with a *35S* minimal promoter in a plasmid that also contained a *35S:Renilla Luciferase* gene as a normalization control (Fig. 4, C and D). Constructs were assayed either by transfecting the plasmids into soybean embryo protoplasts or by co-transfecting the plasmid into *Nicotiana benthamiana* (tobacco) leaf protoplasts along with another plasmid with a *35S:GmWRI1* or *35S:mCHERRY* gene (Fig. 4, C and D).

We showed that all WT GmWRI1 binding sites promoted reporter gene activity in soybean embryo protoplasts, and all except *ACP3* resulted in reporter gene activation in leaf protoplasts when co-transfected with *35S:GmWRI1* (Fig. 4, C and D). These results suggest that *cis*-regulatory elements within these binding sites are sufficient to recruit GmWRI1 and, perhaps, other TFs to activate gene transcription. We assayed binding sites with mutated DNA motifs to determine if the AW Box or CNC Box acted as *cis*-regulatory elements and obtained variable results. As shown in Fig. 4, C and D, mutations in the AW Box of *ACP4* and *BCCP2* binding sites severely reduced reporter gene activity in both embryo and leaf protoplasts. By contrast, mutations in the CNC Boxes of these binding sites did not reduce reporter gene activity. These results suggest that GmWRI1 function depends primarily on AW Box elements in *ACP4* and *BCCP2* genes. Mutations in the AW Boxes and CNC Boxes of *CAC2* and *ACP3* binding sites did not compromise their enhancer activity in soybean embryos. However, mutations in either motif of *CAC2* binding sites compromised reporter gene activation in leaf protoplasts (Fig. 4D). Because there is positional overlap between one of the two CNC Boxes and the AW Box in the

CAC2 binding site (Fig. 4B), it was necessary to establish that the negative effect of the CNC

Box mutations on enhancer activity did not result from alteration of the AW Box. We showed

that mutation of only the upstream CNC Box (mCNCv2) was sufficient to cause a reduction in

reporter gene activity in transactivation experiments with GmWRI1 in leaf protoplasts (Fig. 4D).

Furthermore, *CAC2* binding sites with mutations in both CNC Boxes and in the AW Box

(mAWmCNC) resulted in a greater reduction in reporter gene activity than those resulting from

mutations in either motif separately (Fig. 4D). This suggests that the two motifs act additively or

synergistically in transcriptional regulation.

The *ACP3* binding site did not display enhancer activity in leaf protoplasts, and AW Box

and CNC Box mutations in this binding site did not compromise its enhancer activity in embryo

protoplast (Fig. 4, C and D). These results open the possibility that GmWRI1 occupancy and

function might require other TFs. We created a series of 100 bp overlapping deletions in the 500

bp region surrounding the *ACP3* ChIP-Seq peak and assayed the effects of deleting sequential

100 bp segments in embryo protoplasts to identify regions with *cis*-regulatory elements (SI

Appendix Fig. S6, A and B). Although each deletion caused some reduction in reporter gene

activity compared to the complete region, deletions of regions containing the GmWRI1 ChIP-

Seq peak resulted in the greatest reduction in reporter gene activity (SI Appendix Fig. S6).

Because AW Box and CNC Box mutations in the *ACP3* binding site did not cause a reduction in

reporter gene activity (Fig. 4, C and D) we conclude that other DNA elements are required for

ACP3 enhancer activity.

Together, these results suggest that the AW Box and CNC Box motifs in GmWRI1

binding sites function as enhancer elements. Although the majority of these *cis*-regulatory

elements are located downstream of the TSS, they were able to promote GmWRI1 transcription

when placed upstream of the TSS.

GmWRI1 and GmLEC1 Control Fatty Acid-Related Genes through Distinct Binding Sites

In addition to FA and TAG biosynthesis genes, many GmWRI1 DT genes encode TFs (Dataset

S1), several of which are involved in controlling major embryonic processes in soybean and/or

Arabidopsis, such as LEC1, ABI3, FUS3, ABI4, AGL15, and WRI1 (32).

GmLEC1 is particularly interesting because it and AtLEC1 are major regulators of FA

and TAG biosynthesis in embryos, and GmLEC1 binding sites have been profiled genome-wide

(28, 29, 34). We previously showed that *GmWRI1-A* and *GmWRI1-B* are direct target genes of

GmLEC1 in soybean embryos (28), and we show here that *GmLEC1-1* and *GmLEC1-2* are

GmWRI1 DT genes (Dataset S1). These results suggest that GmWRI1 and GmLEC1 act in a

transcriptional positive feedback circuit in the control of FA and TAG biosynthesis in embryos.

To understand how the two TFs regulate FA and TAG biosynthesis programs, we

identified FAT GmWRI1 DT genes that are also bound by GmLEC1 in em-stage embryos (28).

Fig. 5A shows that 69 of 111 of FAT GmWRI1 DT genes were also bound by GmLEC1 (p

<1.5e-07). The GmWRI1 DT/GmLEC1 bound genes encode the vast majority of enzymes

required for FA and TAG biosynthesis, although some key steps in the pathways are targeted

uniquely by each TF (SI Appendix Fig. S7). Specifically, the KASIII enzyme (Fig. 1A) is

encoded by a GmWRI1 DT gene that is not bound by GmLEC1 (SI Appendix Fig. S7, Dataset

S3). By contrast, genes encoding fatty acyl-ACP thioesterases B (FATB) and glycerol-3-

phosphate acyltransferase (GPAT) are direct targets of GmLEC1 but not GmWRI1 (28). These

results suggest that although complementary, the two TFs serve some specific roles in FA and

TAG biosynthesis in embryos. Moreover, FAT mRNAs encoded by GmWRI1 DT/GmLEC1

bound genes were at higher levels than those encoded by GmWRI1 DT genes alone (Fig. 5B).

Our results suggest that GmWRI1 and GmLEC1 serve additive and/or synergistic functions in

FAT gene regulation in embryos.

We next evaluated the positional relationship between GmWRI1 and GmLEC1 binding

sites in FAT GmWRI1 DT genes. Fig. 5C shows that the median distance between GmWRI1

and GmLEC1 ChIP-Seq peak positions was 142 bp, ranging between 3 and 5,655 bp, suggesting

that these TFs bind adjacent but distinct sites in target genes. To address this hypothesis, we

analyzed DNA motif enrichment in a 50 bp window around ChIP-Seq peaks. As observed

previously, GmWRI1 binding sites were strongly enriched for the presence of AW Box and CNC

Box in the FAT genes targeted by GmWRI1 only and in the FAT GmWRI1 DT genes that are

also bound by GmLEC1 (Fig. 5D). GmLEC1 binding sites were enriched for G Box-type DNA

motifs (G Box and ACGTG) and the AW Box but not the CNC Box motifs (Fig. 5D). Although

GmLEC1 has not been shown to bind G Box-type DNA motifs directly, its occupancy in FAT

GmWRI1 DT genes is likely anchored by a bZIP type TF that binds G Box elements and

physically interacts with GmLEC1, as suggested previously (28, 35). Given the distance

between binding sites and the absence of evidence indicating that the two TFs interact

physically, GmWRI1 and GmLEC1 appear to act independently to promote the transcription of

FAT GmWRI1 DT genes.

Discussion

GmWRI1 is a Central Regulator of Storage Lipid Biosynthesis in Embryos

Fatty acid and TAG biosynthesis in embryos requires the activity of several enzymes located in

distinct cellular compartments. Expression of genes encoding these enzymes must be highly

coordinated to ensure the rapid and efficient incorporation and storage of carbon as FAs and

TAGs during seed development (Fig. 1A)(3, 4). Given the critical role played by WRI1 in

regulating FA biosynthesis genes and in controlling storage lipid accumulation in embryos (36),

we obtained the first genome-wide profile of WRI1 binding sites and of the genes directly

regulated by WRI1 in any plant.

Several lines of evidence indicate that GmWRI1 DT genes are directly regulated by

GmWRI1 and that GmWRI1 is a key regulator of storage lipid biosynthesis. First, GmWRI1 DT

genes correspond to FA and TAG related genes that are upregulated by ectopic GmWRI1

overexpression in soybean. Specifically, 24 of 55 genes upregulated by ectopic *GmWRI1*

expression in transgenic soybean seeds, 19 of which are FAT genes, are GmWRI1 DT genes

(25). Similarly, 7 of 18 genes, 6 of which are FAT genes, upregulated by ectopic *GmWRI1* in

soybean roots are GmWRI1 DT genes (20). Second, there is significant overlap between

GmWRI1 DT genes and homologous Arabidopsis genes that are downregulated in *atwri1*

mutants (Fig. 2). We note that a microarray analysis of ~3,500 Arabidopsis gene identified 25

downregulated genes in *atwri1* mutant when compared to WT seeds (37), 11 of which

overlapped with the 1,213 list of downregulated genes in *atwri1* mutant embryos identified here

399 (Dataset S5). This restricted overlap likely reflects the limited gene numbers assayed previously

400 and reduced sensitivity of the analysis resulting from the use of RNA from whole seeds rather

401 than embryos given that *AtWRI1* is primarily expressed embryonic tissues (38). Third,

402 Arabidopsis gene homologs of GmWRI1 DT genes overlapped significantly with those that are

403 upregulated upon ectopic overexpression of GmWRI1 in Arabidopsis leaf protoplasts (Fig. 2,

404 Dataset S3). Fourth, GmWRI1 DT genes are enriched for FAT genes and encode the vast

405 majority of enzymes involved in the FA and TAG biosynthetic pathways from the synthesis of

406 malonyl-CoA in plastids to the assembly of TAG in the endoplasmic reticulum (Fig. 1, A and E).

408   Others have reported that AtWRI1 regulates genes encoding glycolytic enzymes (17, 19,

409 37). Hexoses funneled through the glycolysis pathway account for much of the carbon used for

410 FA biosynthesis. Although glycolytic process was not an enriched GO term for GmWRI1 DT

411 genes, seven genes encoding plastid-localized glycolytic enzymes were identified as GmWRI DT

412 genes (Dataset S1). Together, these results indicate that GmWRI1 is a major regulator of storage

413 lipid biosynthesis.

415 **Biological Relevance of GmWRI1 Binding Preference for Intragenic Regions**

416 GmWRI1 binds preferentially with intragenic regions, consistent with the observation that AW

417 Box and CNC Box motifs are enriched primarily downstream of the TSS (Figs. 3A, S1B, and

418 S4). We demonstrated that intragenic binding was significant transcriptionally by showing that

419 GmWRI1 binding sites with AW and CNC motifs located primarily downstream of the TSS in

420 *BCCP2, ACP4*, and *CAC2* genes promote GmWRI1-mediated transcription when placed

421 upstream of a minimal promoter (Fig. 4, C and D). Together, these results establish that the

422 intragenic binding sites contain transcriptional enhancer elements.

424   Our finding that GmWRI1 DT genes possess intragenic binding sites is not unique. A

425 survey of Arabidopsis TF binding sites revealed that 13 of 27 TFs analyzed exhibited substantial

426 binding to intragenic sites (39), and an estimated 36% of enhancer elements in Drosophila genes

427 are located in intragenic regions (40). A mechanistic explanation of how intragenic binding

428 affects transcription comes from a recent study of the cucumber TENDRIL IDENTITY

429 (CsTEN), a TCP TF of the CYC/TB1 clade. CsTEN binds primarily to target gene intragenic

regions resulting in chromatin disassembly and transcription activation (41). CsTEN possesses

an amino-terminal intrinsically disordered region domain with histone acetyltransferase activity

that facilitates chromatin accessibility during target gene transcription. Interestingly, the

AtWRI1 TF also has an amino-terminal intrinsically disordered region domain (42), opening the

possibility that CsTEN and WRI1 share mechanistic similarities, although further experiments

are needed to demonstrate such associations. Nevertheless, our results provided new insights

into the mechanisms by which GmWRI1 functions in storage lipid biosynthesis in embryos.

GmWRI1 Function is Dependent on AW Boxes and CNC Boxes

Our results provide compelling evidence that the AW Box serves as an enhancer element that

mediates transcriptional regulation by WRI1. Mutations in AW Box elements in *BCCP2, ACP4,*

and *CAC2* binding sites compromised their enhancer activity in transactivation assays with

GmWRI1 in leaf protoplasts. Consistent with these findings, others have shown that AW Box

elements are enriched in WRI1 target genes in several plants, that WRI1 binds AW Box elements

in vitro, and that mutations in the AW Box compromised transcriptional activation by WRI1 (19,

20, 23, 25, 43-45).

A novel finding of our study is that DNA elements other than AW Boxes are involved in

GmWRI1 transcriptional activation. Our finding that AW Box mutations did not compromise

the enhancer activity of *CAC2* and *ACP3* binding sites in soybean embryo assays and of *ACP3*

binding sites in leaf protoplast assays indicates that other DNA *cis* elements are present in these

binding sites (Fig. 4, C and D). Consistent with this conclusion, only ~55% of binding sites for

FAT GmWRI1 DT genes contain full AW Boxes, implying that GmWRI1 may bind other DNA

elements or that other TFs bind GmWRI1 binding sites.

Mutational analysis of the *CAC2* binding site in leaf cells demonstrated conclusively that

the CNC Box in the GmWRI1 binding site is an enhancer element that promotes GmWRI1

transcriptional activity (Fig. 4D). The *CAC2* binding site also possesses AW Box enhancer

elements that promote transcription. Comparison of the effects of AW Box and CNC Box

mutations singly and in combination suggests that the two elements act additively or

synergistically to enhance GmWRI1-mediated transcription. Consistent with the identification

of the CNC Box as an enhancer element, the CNC Box is highly enriched in regions downstream

of the TSS in GmWRI1 DT and AtWRI1 DR genes (Fig. S4 and S5). Enrichment of the CNC

Box in GmWRI1 binding sites opens the possibility that the CNC Box may enhance GmWRI1

occupancy at these binding sites.

The mechanism by which the CNC Box enhances GmWRI1 transcriptional activity is not

known, but similarities between the CNC Box and known DNA motifs provide potential clues

about its function as an enhancer element. The CNC Box shares strong similarity with the

CTCCGCC DNA motif that acts as an intragenic, positive transcriptional enhancer element that

is bound by the CsTEN TCP-like TF (41). One model is that a TCP-like TF physically binds

with the CNC Box and enhances GmWRI1 occupancy and functions in an AW Box-independent

manner (Fig. 6). Recent studies showed that AtWRI1 interacts physically with the TCP TFs,

AtTCP4, AtTCP10 and AtTCP24, although these TFs appear to repress seed oil accumulation

(46). An alternate hypothesis is based on the fact that WRI1 is a member of the AP2/EREPBP

TF family (16, 44). A survey of the Arabidopsis cistrome revealed that several AP2/EREBP

domain TFs, such as ERF11, 15 and 105 and RAP212, bind DNA motifs similar to the CNC Box

(47). It is possible that GmWRI1 could potentially bind directly with the CNC Box, although

WRI1 has not been reported to bind physically with DNA motifs other than the AW Box.

Additional experiments are needed to define the role of CNC Box elements in controlling the

transcription of FAT genes regulated by GmWRI1 (Fig. 6).

GmWRI1 Acts with GmLEC1 and Other Transcription Factors to Establish a Regulatory

Environment that Promotes Storage Lipid Accumulation in Embryos

In addition to establishing that GmWRI1 directly regulates genes encoding the vast majority of

enzymes in the FA and TAG biosynthetic pathways, our results also indicate that GmWRI1 acts

to alter the regulatory state of soybean embryos to promote storage lipid accumulation. We

showed that GmWRI1 directly regulates several other well characterized TFs in seeds, including

GmLEC1, GmFUS3, and GmABI3, whose counterparts in Arabidopsis transcriptionally regulate

genes involved in storage lipid accumulation (34, 48). We also showed that GmWRI1 is a

GmWRI1 DT, indicating that it is a self-activating TF. Thus, GmWRI1 is a critical component

of a transcriptional regulatory network that promotes storage lipid accumulation and other

processes during seed development.

The relationship between GmWRI1 and GmLEC1 provides a particularly illuminating

example of the regulatory interplay that occurs between these TFs. Here, we showed that

GmLEC1 is directly regulated by GmWRI1, that overexpression of GmWRI1 in Arabidopsis leaf

protoplasts results in the upregulation of *AtLEC1*, and that *AtLEC1-LIKE* is downregulated in

atwri1 mutant embryos (Dataset S5). We showed previously that GmWRI1 is directly regulated

by GmLEC1, that overexpression of *AtLEC1* results in elevated expression of *AtWRI1*, and that

AtWRI1 is downregulated in *atlec1* mutant seeds (28, 29). These findings indicate that WRI1

and LEC1 act in a positive feedback subcircuit to control FA and TAG biosynthesis in embryos

through reciprocal regulation of the other TF's expression and through their regulation of FA and

TAG biosynthetic genes (Fig. 6). Positive feedback gene circuits act in gene regulatory networks

to trigger irreversible changes in transcriptional programs (49). A hallmark of the onset of the

seed maturation program is the *de novo* initiation of many biological programs that are silenced

during the morphogenesis phase of the embryo development, such as storage lipid and protein

accumulation (50). Initiation of the FA biosynthetic program in seeds requires a robust and

stable regulatory state to ensure proper expression of genes required for the quick and efficient

conversion of carbon into oil bodies. In this sense, a positive feedback subcircuit between

GmLEC1 and GmWRI1 is likely to be required for the rapid onset of the FA and TAG

accumulation programs during seed development.

GmWRI1 also appears to act in positive feedback circuits with GmABI3 and GmFUS3,

which are both GmWRI1 DT genes (Dataset S1). GmABI3 directly regulates *GmWRI1*, and

ectopic *GmABI3* expression activates *GmWRI1* (28, 51). Although GmFUS3 DT genes have not

been identified, AtFUS3 directly regulates *AtWRI1* in Arabidopsis (52). These findings

emphasize that GmWRI1 is an integral component of the transcriptional regulatory networks

controlling storage lipid accumulation.

Materials and Methods

Chromatin Immunoprecipitation and DNA Sequencing

Soybean cv Williams 82 (W82) plants were grown, and embryos were isolated from seeds staged

as described previously (29). GmWRI1 ChIP assays were conducted as described previously

(29), with minor modifications. After nuclei purification, nuclei were resuspended in shearing

buffer (10mM Tris pH8, 1mM EDTA, 0.1% SDS, 1mM PMSF), and chromatin was sheared to

~250 bp fragments using a Covaris E220 ultrasonicator. After shearing, Triton X-100 and NaCl

were added to chromatin to final concentrations of 1% and 150mM, respectively. Anti-

GmWRI1 (Glyma.08G227700 and Glyma.15G221600) antibodies for ChIP were raised in

rabbits against the peptide FSTQPGDHETESDVN, corresponding to the amino acids 352-366 of

GLYMA.08G227700 and amino acids 348-361 (FSTQPGDETESDVN) of Glyma.15G221600.

Antibody reactivity was assessed using Western blot analysis (Dataset S2). ChIP DNA library

preparation and DNA sequencing was done as described previously (29).

RNA-Sequencing Experiments with Arabidopsis Leaf Protoplasts Overexpressing

GmWRI1 **and with** *atwri1* **Mutant Embryos**

For *GmWRI1* overexpression assays with Arabidopsis leaf protoplasts, three-week-old

Arabidopsis thaliana Col-0 plants were grown as described (53). Arabidopsis leaf mesophyll

protoplasts were isolated and transfected as described previously (28). For each biological

replicate, 500,000 cells were transfected either with 10 μg of a plasmid carrying the *GmWRI1-A*

cDNA that was synthesized using primers listed in Dataset S6, Table S1, and driven by the

CaMV35S promoter (*p35S:GmWRI1*) or with a *35S:mCHERRY:NOSt* plasmid as a negative

control. Protoplasts were incubated for 16 hours at room temperature following transfection.

Homozygous *atwri1-1* and WT Col-0 seeds were plated on GM media and grown on soil

as described previously (53). Flowers from six-week-old plants were tagged at anthesis, and

embryos were dissected from seeds 8 days after pollination (25-30 embryos per biological

replicate, three biological replicates per genotype).

Total RNA was isolated using the RNAqueous™-Micro Total RNA Isolation Kit,

following manufacturer's instructions. RNA samples were then treated with DNAse for

digestion of DNA using the TURBO DNA-free™ Kit, following manufacturer's instructions.

RNA-Seq libraries were prepared with Tecan Ovation RNA-Seq System V2 using 50 ng of total

RNA following protocols as described by Kumar *et al.* (54), with NEXTflex ChIP-Seq Barcodes

(BioScientific). DNA was sequenced with a NovaSeq 6000 sequencer.

Cloning and Mutation of GmWRI1 Binding Sites

GmWRI1 binding sites (200 to 500 bp around the WRI1 ChIP-seq peak) were PCR amplified

from the soybean genomic DNA and cloned as described previously (28). Mutated derivatives

of binding sites and *ACP3* deletion series were synthesized by Twist Bioscience and inserted into

pDLUC15 using the same cloning strategy. Mutant derivatives of *ACP4* were created by

amplifying the WT *ACP4* binding site with primers containing the mutated sequences. Primers

and the full DNA sequences of WRI1 binding sites and mutant derivatives are listed in Dataset

S6, Table S1 and S2.

Transient Expression Assays

Transient assays of firefly and Renilla luciferase activity in protoplasts were performed as

described previously (28). All trials were done with three biological replicates, and one-way or

two-way ANOVA with Tukey's post hoc tests ($P < 0.05$) were used for the statistical analysis in

soybean embryo protoplasts and tobacco leaf protoplasts, respectively.

Data Analysis

GmWRI1 ChIP-Seq in cot and em stage soybean embryos and identification of GmWRI1 DT

genes. ChIP-Seq data were analyzed as described previously (28), with the exception that

MACS3 software (https://github.com/macs3-project/MACS) was used in place of MACS2.

Quality assessment of the ChIP-seq libraries were determined by evaluating the non-redundant

fraction (NRF) and strand cross-correlation (CC), following ENCODE guidelines for ChIP-Seq

data (26). NRF and CC values are summarized in Dataset S2

Because GmWRI1 binding sites were located primarily in regions downstream of the

TSS, we designated genes bound by GmWRI1 as those with a reproducible ChIP-Seq peak

within a 1 kb window upstream of its TSS or within the annotated gene body.

584    GmWRI1 DT genes are defined as those bound by GmWRI1 and coexpressed by

585  GmWRI1 (Dataset S1). Because *GmWRI1* is expressed primarily in embryonic cells (Fig. 1C),

586  we used the Harada-Goldberg Soybean Seed Development LCM RNA-Seq Dataset (GEO

587  accessions, GSE57606, GSE46096, and GSE99109) to identify genes that are two-fold

588  upregulated (FDR < 0.01) in embryo cotyledon abaxial or adaxial parenchyma versus seed coat

589  hilum and seed coat parenchyma subregions. We then defined WRI1 co-expressed genes as

590  those upregulated in at least two of the above comparisons (Dataset S1).

592  ***Identification of soybean FAT genes.*** To define FAT genes, we used the soybean Plant

593  Metabolic Network database, SoyCyc, and extracted the soybean genes associated with the

594  pathways listed in the Dataset S3. We also included genes annotated with the GO category for

595  fatty acid biosynthetic process (GO:0006633).

597  ***RNA-Seq data from Arabidopsis protoplasts and embryos.*** Sequenced reads were

598  demultiplexed and reads corresponding to rRNA sequences were removed. The resulting filtered

599  reads were mapped to Arabidopsis primary transcripts (TAIR10) using bowtie v0.12.7 with

600  parameters -v 2 -5 10 -3 4 -m 1 --best –strata. The EdgeR package (v3.10.5) was used to obtain

601  normalized expression values using the Trimmed Mean of M-values (TMM) method, and to

602  identify differentially expressed genes between the different genotypes (FDR < 0.05).

604  ***GO enrichment analysis.*** GO term enrichment analyses were done as described previously (28,

605  29). GO functional annotation was extracted from TAIR10 and Soybase databases and used for

606  Arabidopsis and soybean datasets, respectively. A q value threshold of 0.01 was used to identify

607  enriched GO categories.

609  ***DNA motif analysis.*** DNA motif *de novo* discovery analyses of binding sites in GmWRI1 DT

610  genes and FAT GmWRI1 DT genes was performed using the MEME-ChIP tool from the MEME

611  suite v5.3.3 as described previously (28), using a motif maximum length of 15 and a E-value

612  cutoff of 0.01. Discovered motifs were compared to the Arabidopsis DAP-Seq TF and the top-

613  ranked DNA motifs identified are listed in Dataset S4.

DNA motif enrichment analysis was performed as described (28), using the DNA motifs

AW-BOX (CNTNGNNNNNNNCG), CNC-Box (CNCCNCC), G-Box (CACGTG), RY

(CATGCA), and CCAAT-Box (CCAAT) to screen the GmWRI1 and GmLEC1 binding sites

and 500 bps windows around the TSS of WRI1 directly regulated genes in Arabidopsis using the

"known" function from HOMER (55).

Accession Numbers

Data are available at GEO, under the accessions: GSE152310 and GSE253104 (GmWRI1 ChIP-

seq at cot and em stages, respectively), GSE253109 (*atwri1* mutant embryo RNA Seq), and

GSE253107 (GmWRI1 overexpression RNA Seq).

Acknowledgments

We thank Makenzie Melby, Sana Mohammad, and Mary Irani for their technical assistance.

This work was supported by a grant from the National Science Foundation Plant Genome

Research Program to RBG and JJH.

References

1.   J. M. Dyer, S. Stymne, A. G. Green, A. S. Carlsson, High-value oils from plants. *Plant J.* **54**, 640-655 (2008).

2.   OECD-FAO, OECD-FAO Agricultural Outlook 2022-2031.

3.   P. D. Bates, S. Stymne, J. Ohlrogge, Biochemical pathways in seed oil synthesis. *Curr Opin Plant Biol* **16**, 358-364 (2013).

4.   S. Baud, Seeds as oil factories. *Plant Reprod* **31**, 213-235 (2018).

5.   J. L. Harwood, Recent advances in the biosynthesis of plant fatty acids. *Bba-Lipid Lipid Met* **1301**, 7-56 (1996).

6.   P. D. Bates, J. Browse, The significance of different diacylgycerol synthesis pathways on plant oil composition and bioengineering. *Front Plant Sci* **3**, 147 (2012).

7.   T. P. Durrett, C. Benning, J. Ohlrogge, Plant triacylglycerols as feedstocks for the production of biofuels. *Plant J.* **54**, 593-607 (2008).

8.   T. L. Shimada, M. Hayashi, I. Hara-Nishimura, Membrane Dynamics and Multiple Functions of Oil Bodies in Seeds and Leaves. *Plant Physiol* **176**, 199-207 (2018).

9.   S. A. Braybrook, J. J. Harada, LECs go crazy in embryo development. *Trends Plant Sci.* **13**, 624-630 (2008).

10. D. A. Levin, The Oil Content of Seeds: An Ecological Perspective. *The American naturalist* **108**, 193-206 (1974).

11. S. Baud, L. Lepiniec, Regulation of de novo fatty acid synthesis in maturing oilseeds of Arabidopsis. *Plant Physiol. Biochem.* **47**, 448-455 (2009).

12. S. Baud, I. A. Graham, A spatiotemporal analysis of enzymatic activities associated with carbon metabolism in wild-type and mutant embryos of Arabidopsis using in situ histochemistry. *Plant J.* **46**, 155-169 (2006).

13. S. Baud, L. Lepiniec, Physiological and developmental regulation of seed oil production. *Prog Lipid Res* **49**, 235-249 (2010).

14. P. O'Hara, A. R. Slabas, T. Fawcett, Fatty acid and lipid biosynthetic genes are expressed at constant molar ratios but different absolute levels during embryogenesis. *Plant Physiol* **129**, 310-320 (2002).

15. M. A. Troncoso-Ponce *et al.*, Comparative deep transcriptional profiling of four developing oilseeds. *Plant Journal* **68**, 1014-1027 (2011).

16. A. Cernac, C. Benning, WRINKLED1 encodes an AP2/EREB domain protein involved
in the control of storage compound biosynthesis in Arabidopsis. *Plant J* **40**, 575-585
(2004).

17. N. Focks, C. Benning, wrinkled1: A novel, low-seed-oil mutant of Arabidopsis with a
deficiency in the seed-specific regulation of carbohydrate metabolism. *Plant Physiol* **118**,
91-101 (1998).

18. D. An, M. C. Suh, Overexpression of Arabidopsis WRI1 enhanced seed mass and storage
oil content in Camelina sativa. *Plant Biotechnology Reports* **9**, 137-148 (2015).

19. S. Baud, S. Wuilleme, A. To, C. Rochat, L. Lepiniec, Role of WRINKLED1 in the
transcriptional regulation of glycolytic and fatty acid biosynthetic genes in Arabidopsis.
Plant J. **60**, 933-947 (2009).

20. L. Chen *et al.*, Soybean (Glycine max) WRINKLED1 transcription factor, GmWRI1a,
positively regulates seed oil accumulation. *Mol Genet Genomics* **293**, 401-415 (2018).

21. J. Liu *et al.*, Increasing seed mass and oil content in transgenic Arabidopsis by the
overexpression of wri1-like gene from Brassica napus. *Plant Physiol. Biochem.* **48**, 9-15
(2010).

22. B. Shen *et al.*, Expression of ZmLEC1 and ZmWRI1 Increases Seed Oil Production in
Maize. *Plant Physiology* **153**, 980-987 (2010).

23. K. Maeo *et al.*, An AP2-type transcription factor, WRINKLED1, of Arabidopsis thaliana
binds to the AW-box sequence conserved among proximal upstream regions of genes
involved in fatty acid synthesis. *Plant J.* **60**, 476-487 (2009).

24. A. To *et al.*, WRINKLED transcription factors orchestrate tissue-specific regulation of
fatty acid biosynthesis in Arabidopsis. *Plant Cell* **24**, 5007-5023 (2012).

25. B. Chen *et al.*, Multiple GmWRI1s are redundantly involved in seed filling and
nodulation by regulating plastidic glycolysis, lipid biosynthesis and hormone signalling
in soybean (Glycine max). *Plant Biotechnol. J.* **18**, 155-171 (2020).

26. S. G. Landt *et al.*, ChIP-seq guidelines and practices of the ENCODE and modENCODE
consortia. *Genome Res* **22**, 1813-1831 (2012).

27. P. J. Farnham, Insights from genomic profiling of transcription factors. *Nat Rev Genet* **10**,
605-616 (2009).

28.   L. Jo *et al.*, Combinatorial interactions of the LEC1 transcription factor specify diverse

694   developmental programs during soybean seed development. *Proc Natl Acad Sci U S A*

695   **117**, 1223-1232 (2020).

696   29.   J. M. Pelletier *et al.*, LEC1 sequentially regulates the transcription of genes involved in

697   diverse developmental processes during seed development. *Proc Natl Acad Sci U S A*

698   **114**, E6710-E6719 (2017).

699   30.   G. Bonaventure, J. B. Ohlrogge, Differential Regulation of mRNA Levels of Acyl Carrier

700   Protein Isoforms in Arabidopsis. *Plant Physiology* **128**, 223-235 (2002).

701   31.   A. Forderer, Y. Zhou, F. Turck, The age of multiplexity: recruitment and interactions of

702   Polycomb complexes in plants. *Curr Opin Plant Biol* **29**, 169-178 (2016).

703   32.   M. Alizadeh, R. Hoy, B. Lu, L. Song, Team effort: Combinatorial control of seed

704   maturation by transcription factors. *Curr Opin Plant Biol* **63**, 102091 (2021).

705   33.   Q. Kong *et al.*, The Arabidopsis WRINKLED1 transcription factor affects auxin

706   homeostasis in roots. *Journal of experimental botany* **68**, 4627-4634 (2017).

707   34.   J. Mu *et al.*, LEAFY COTYLEDON1 is a key regulator of fatty acid biosynthesis in

708   Arabidopsis. *Plant Physiol* **148**, 1042-1054 (2008).

709   35.   A. Yamamoto *et al.*, Arabidopsis NF-YB subunits LEC1 and LEC1-LIKE activate

710   transcription by interacting with seed-specific ABRE-binding factors. *Plant J.* **58**, 843-

711   856 (2009).

712   36.   Q. Kong, Y. Yang, L. Guo, L. Yuan, W. Ma, Molecular Basis of Plant Oil Biosynthesis:

713   Insights Gained From Studying the WRINKLED1 Transcription Factor. *Front Plant Sci*

714   **11**, 24 (2020).

715   37.   S. A. Ruuska, T. Girke, C. Benning, J. B. Ohlrogge, Contrapuntal networks of gene

716   expression during Arabidopsis seed filling. *Plant Cell* **14**, 1191-1206 (2002).

717   38.   M. F. Belmonte *et al.*, Comprehensive developmental profiles of gene activity in regions

718   and subregions of the Arabidopsis seed. *Proc Natl Acad Sci USA* **110**, E435-E444 (2013).

719   39.   K. S. Heyndrickx, J. Van de Velde, C. Wang, D. Weigel, K. Vandepoele, A functional

720   and evolutionary perspective on transcription factor binding in Arabidopsis thaliana.

721   *Plant Cell* **26**, 3894-3910 (2014).

722   40.   E. Z. Kvon *et al.*, Genome-scale functional characterization of Drosophila developmental

723   enhancers in vivo. *Nature* 10.1038/nature13395 (2014).

41.   X. Yang *et al.*, Regulation of plant architecture by a new histone acetyltransferase
targeting gene bodies. *Nat Plants* **6**, 809-822 (2020).

42.   W. Ma *et al.*, Deletion of a C-terminal intrinsically disordered region of WRINKLED1
affects its stability and enhances oil accumulation in Arabidopsis. *Plant J.* **83**, 864-874
(2015).

43.   C. Kuczynski, S. McCorkle, J. Keereetaweep, J. Shanklin, J. Schwender, An expanded
role for the transcription factor WRINKLED1 in the biosynthesis of triacylglycerols
during seed development. *Front Plant Sci* **13**, 955589 (2022).

44.   Z. Qiao *et al.*, Molecular basis of the key regulator WRINKLED1 in plant oil
biosynthesis. *Science Advances* **8** (2022).

45.   Q. Tan *et al.*, The molecular mechanism of WRINKLED1 transcription factor regulating
oil accumulation in developing seeds of castor bean. *Plant Divers* **45**, 469-478 (2023).

46.   Q. Kong *et al.*, TEOSINTE BRANCHED1/CYCLOIDEA/PROLIFERATING CELL
FACTOR4 Interacts with WRINKLED1 to Mediate Seed Oil Biosynthesis. *Plant Physiol*
184, 658-665 (2020).

47.   R. C. O'Malley *et al.*, Cistrome and Epicistrome Features Shape the Regulatory DNA
Landscape. *Cell* **165**, 1280-1292 (2016).

48.   Y. Yang *et al.*, Transcriptional regulation of oil biosynthesis in seed plants: Current
understanding, applications, and perspectives. *Plant Commun* **3**, 100328 (2022).

49.   W. Xiong, J. E. Ferrell, A positive-feedback-based bistable 'memory module'that
governs a cell fate decision. *Nature* **426**, 460-465 (2003).

50.   L. Jo, J. M. Pelletier, J. J. Harada, Central role of the LEAFY COTYLEDON1
transcription factor in seed development. *J Integr Plant Biol* **61**, 564-580 (2019).

51.   D. Zhang *et al.*, Plasticity and innovation of regulatory mechanisms underlying seed oil
content mediated by duplicated genes in the palaeopolyploid soybean. *Plant J.* **90**, 1120-
1133 (2017).

52.   F. Wang, S. E. Perry, Identification of direct targets of FUSCA3, a key regulator of
Arabidopsis seed development. *Plant Physiol* **161**, 1251-1264 (2013).

53.   M. A. L. West *et al.*, LEAFY COTYLEDON1 Is an Essential Regulator of Late
Embryogenesis and Cotyledon Identity in Arabidopsis. *Plant Cell* **6**, 1731-1745 (1994).

54.   R. Kumar *et al.*, A high-throughput method for Illumina RNA-Seq library preparation.

755   *Frontiers in Plant Science* **3** (2012).

756   55.   S. Heinz *et al.*, Simple combinations of lineage-determining transcription factors prime

757   cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576-

758   589 (2010).

759   56.   E. Collakova *et al.*, Metabolic and Transcriptional Reprogramming in Developing

760   Soybean (Glycine max) Embryos. *Metabolites* **3**, 347-372 (2013).

 Figure Legends

Figure 1. GmWRI1 acts as a central regulator of fatty acid and triacylglycerol biosynthesis

in soybean seeds. (A) Simplified schematic diagram of FA and TAG biosynthesis from Baud

and Lepiniec (11). Enzymes (labeled in red) encoded by genes that are bound by (teal) and/or

co-expressed with (grey) GmWRI1. (B) Temporal changes in relative storage fatty acid and

protein levels (top, adapted from Collakova *et al.* (56)) and of *GmWRI1-A* (*Glyma.08G227700*)

and *GmWRI1-B* (*Glyma.15G221600*) mRNA levels (bottom, from the Harada Embryo mRNA-

Seq Dataset, GEO Series GSE99571) during soybean embryo development. (C) mRNA levels of

GmWRI1-A and *GmWRI1-B* in seed subregions of the embryo axis, and cotyledons, endosperm,

and seed coat at the early maturation stage (GEO Series GSE46096). (D) GmWRI1 DT genes in

soybean embryos at the early maturation stage. Venn diagram shows the overlap of GmWRI1

bound and coexpressed genes. Statistical significance (hypergeometric distribution) of the

overlap between bound genes and co-expressed genes is indicated. (E) Venn diagram showing

the overlap of the GmWRI1 DT genes at the em stage and of FAT genes, with the statistical

significance of the overlap indicated (hypergeometric distribution). (F) Heatmap showing the q

value significance of GO term enrichment for GmWRI1 co-expressed, bound, and GmWRI1 DT

genes. The GO terms listed are the top 20 most enriched biological process GO terms for

GmWRI1 DT genes. Abbreviations: cotyledon (cot), early maturation (em), mid-maturation

(mm), late maturation (lm), seedling (sdlg).

Figure 2. Identification of WRI1 directly regulated genes in Arabidopsis. (A) Overview of

strategy used to identify Arabidopsis genes differentially expressed (A) in *atwri1-1* mutants

compared with WT embryos at 8 days after pollination and (B) in leaf protoplasts transfected

with *35S:GmWRI1* or *35S:mCHERRY* as a negative control. RNA-Seq experiments identified

differentially expressed mRNAs that differed at least two-fold (FDR < 0.05) when compared to

the control. (C) Venn diagram showing the overlap between genes downregulated in *atwri1*

mutant embryos and GmWRI1 DT genes. (D) Venn diagram showing the overlap between

genes upregulated in leaf protoplasts overexpressing GmWRI1 and GmWRI1 DT genes. (E)

Venn diagram showing the overlap between genes downregulated in *atwri1* mutant embryos and

upregulated in leaf protoplasts overexpressing *GmWRI1* to identify the AtWRI1 DR genes. (F)

Heatmap showing the q value significance of enriched GO terms for downregulated genes in

atwri1 mutant embryos, upregulated genes in the *35S:GmWRI1* leaf cells, and of the AtWRI1

DR genes. GO terms listed are the top 10 most enriched biological process terms (q value <

0.01).

Figure 3. Binding profile and enriched DNA motifs in GmWRI1 binding sites. (A) Position

of GmWRI1 ChIP-seq peak in reference to the TSS of GmWRI1 DT genes. (B) Pie chart

showing the fraction of GmWRI1 ChIP-Seq peaks that overlap distinct genomic features

(intergenic, 2 kb upstream of the TSS, 5' UTR, exons, introns, 3' UTR and 1 kb downstream of

the TTS). (C) Position weight matrices of DNA motifs discovered *de novo* in GmWRI1 binding

sites of GmWRI1 DT (top), and FAT GmWRI1 DT (bottom) genes, and their significance as

indicated by their associated E values. (D) DNA motif enrichment in the binding sites of

GmWRI1 DT (left) and FAT GmWRI1 DT (right) genes. Graphs show the percentage of bound

regions (green) and randomly selected, equivalently sized and spaced regions (gray) containing

the indicated DNA motifs. A Bonferroni-adjusted *P* value threshold of 0.01 was used to indicate

the statistical significance (*) of enrichment, while n.s. indicates no significant difference.

Figure 4. Functional analysis of GmWRI1 direct target gene binding sites. (A) Genome

browser view showing the position of GmWRI1 ChIP-Seq peaks for *BCCP2*

(Glyma.13G057400), *ACP4* (Glyma.15G098500), *CAC2* (Glyma.05G221100), and *ACP3*

(Glyma.19G240100). ChIP-Seq peaks (dashed lines) and TSS positions (tick mark) are

indicated. Peak distance from the TSS in bp is shown. (B) Diagrams of GmWRI1 binding sites

for genes in (a) used in the transient assay experiments. Yellow, pink and brown symbols,

respectively, indicate the positions of AW Box, CNC Box, and short AW Box motifs. (C) Effect

of WT GmWRI1 binding sites and of binding sites with mutations in all detectable AW Box

(mAW), CNC Box (mCNC) or both (mAWmCNC) DNA motifs on reporter gene activity in

soybean cotyledon embryo protoplasts. (D) Transactivation analysis of WT and mutant

GmWRI1 binding sites in tobacco leaf protoplasts transfected with *p35S:GmWRI1* (teal) or

p35S:mCHERRY (grey). For (C) and (D), average firefly to Renilla luciferase activity ratios

were multiplied by 100 and plotted with the respective standard errors. Lower-case letters

indicate the significance of comparisons based on ANOVA and post hoc Tukey test results.

Figure 5. GmWRI1 and GmLEC1 act independently to control storage lipid biosynthesis in

soybean embryos. (A) Bar graph shows the number of FAT GmWRI1 DT genes that are bound

or not bound by GmLEC1. The *P* value significance of the overlap is indicated (hypergeometric

distribution). (B) Box plots comparing the distribution of mRNA levels in em embryos (GEO

accession GSE99571) of FAT genes are not bound by either GmWRI1 and GmLEC1 (grey),

GmWRI1 DT genes that are not bound by GmLEC1 (teal), and GmWRI1 DT genes that are

bound by GmLEC1 (dark red). Asterisk indicates significant differences between groups

(Wilcoxon rank-sum test, p < 0.01). (C) GmWRI1 ChIP-Seq peak positions relative to LEC1

ChIP-Seq peaks (position 0) in FAT GmWRI1 DT genes that are also bound by GmLEC1. (D)

DNA motif enrichment in GmWRI1 DT binding sites of genes that are not bound or bound by

GmLEC1. Circle diameters depict the frequencies at which DNA motifs were identified in the

indicated regions, and the color intensity indicates the statistical significance of the enrichment

relative to the normal distribution of a population of randomly generated regions (Bonferroni-

adjusted *P* values).

Figure 6. **Model for the regulation of storage lipid accumulation in embryos.** WRI1 and

LEC1 act in a transcriptional positive feedback subcircuit to regulate FA and TAG biosynthesis

in the embryo. GmWRI1 regulates FA and TAG-related gene expression by binding with AW

Box and, possibly, CNC Box elements located downstream of the TSS. An alternative model is

that another TF, such as a TCP-like TF, binds the CNC Box, potentially by interacting with

WRI1. GmLEC1 binds FAT genes at sites that are distinct from GmWRI1 binding sites.

GmLEC1 interacts with bZIP TFs that bind with G Box DNA elements located most frequently

in regions upstream of the TSS. Dashed arrows indicate transcriptional activation.

Figure 1. GmWRI1 acts as a central regulator of fatty acid and triacylglycerol biosynthesis in soybean seeds.

(*A*) Simplified schematic diagram of FA and TAG biosynthesis from Baud and Lepiniec (11). Enzymes (labeled in red) encoded by genes that are bound by (teal) and/or co-expressed with (grey) GmWRI1. (*B*) Temporal changes in relative storage fatty acid and protein levels (top, adapted from Collakova *et al.* (56)) and of *GmWRI1-A* (*Glyma.08G227700*) and *GmWRI1-B* (*Glyma.15G221600*) mRNA levels (bottom, from the Harada Embryo mRNA-Seq Dataset, GEO Series GSE99571) during soybean embryo development. (*C*) mRNA levels of *GmWRI1-A* and *GmWRI1-B* in seed subregions of the embryo axis, and cotyledons, endosperm, and seed coat at the early maturation stage (GEO Series GSE46096). (*D*) GmWRI1 DT genes in soybean embryos at the early maturation stage. Venn diagram shows the overlap of GmWRI1 bound and coexpressed genes. Statistical significance (hypergeometric distribution) of the overlap between bound genes and co-expressed genes is indicated. (*E*) Venn diagram showing the overlap of the GmWRI1 DT genes at the em stage and of FAT genes, with the statistical significance of the overlap indicated (hypergeometric distribution). (*F*) Heatmap showing the q value significance of GO term enrichment for GmWRI1 co-expressed, bound, and GmWRI1 DT genes. The GO terms listed are the top 20 most enriched biological process GO terms for GmWRI1 DT genes. Abbreviations: cotyledon (cot), early maturation (em), mid-maturation (mm), late maturation (lm), seedling (sdlg).

Figure 2. Identification of WRI1 directly regulated genes in Arabidopsis.
(*A*) Overview of strategy used to identify Arabidopsis genes differentially expressed (*A*) in *atwri1-1* mutants compared with WT embryos at 8 days after pollination and (*B*) in leaf protoplasts transfected with *35S:GmWRI1* or *35S:mCHERRY* as a negative control. RNA-Seq experiments identified differentially expressed mRNAs that differed at least two-fold (FDR < 0.05) when compared to the control. (*C*) Venn diagram showing the overlap between genes downregulated in *atwri1* mutant embryos and GmWRI1 DT genes. (*D*) Venn diagram showing the overlap between genes upregulated in leaf protoplasts overexpressing *GmWRI1* and GmWRI1 DT genes. (*E*) Venn diagram showing the overlap between genes downregulated in *atwri1* mutant embryos and upregulated in leaf protoplasts overexpressing *GmWRI1* to identify the AtWRI1 DR genes. (*F*) Heatmap showing the q value significance of enriched GO terms for downregulated genes in *atwri1* mutant embryos, upregulated genes in the *35S:GmWRI1* leaf cells, and of the AtWRI1 DR genes. GO terms listed are the top 10 most enriched biological process terms (q value < 0.01).

Figure 3. Binding profile and enriched DNA motifs in GmWRI1 binding sites.
(*A*) Position of GmWRI1 ChIP-seq peak in reference to the TSS of GmWRI1 DT genes. (*B*) Pie chart showing the fraction of GmWRI1 ChIP-Seq peaks that overlap distinct genomic features (intergenic, 2 kb upstream of the TSS, 5' UTR, exons, introns, 3' UTR and 1 kb downstream of the TTS). (*C*) Position weight matrices of DNA motifs discovered *de novo* in GmWRI1 binding sites of GmWRI1 DT (top), and FAT GmWRI1 DT (bottom) genes, and their significance as indicated by their associated E values. (*D*) DNA motif enrichment in the binding sites of GmWRI1 DT (left) and FAT GmWRI1 DT (right) genes. Graphs show the percentage of bound regions (green) and randomly selected, equivalently sized and spaced regions (gray) containing the indicated DNA motifs. A Bonferroni-adjusted *P* value threshold of 0.01 was used to indicate the statistical significance (*) of enrichment, while n.s. indicates no significant difference.

Figure 4. Functional analysis of GmWRI1 direct target gene binding sites.
(*A*) Genome browser view showing the position of GmWRI1 ChIP-Seq peaks for *BCCP2* (Glyma.13G057400), *ACP4* (Glyma.15G098500), *CAC2* (Glyma.05G221100), and *ACP3* (Glyma.19G240100). ChIP-Seq peaks (dashed lines) and TSS positions (tick mark) are indicated. Peak distance from the TSS in bp is shown. (*B*) Diagrams of GmWRI1 binding sites for genes in (a) used in the transient assay experiments. Yellow, pink and brown symbols, respectively, indicate the positions of AW Box, CNC Box, and short AW Box motifs. (*C*) Effect of WT GmWRI1 binding sites and of binding sites with mutations in all detectable AW Box (mAW), CNC Box (mCNC) or both (mAWmCNC) DNA motifs on reporter gene activity in soybean cotyledon embryo protoplasts. (*D*) Transactivation analysis of WT and mutant GmWRI1 binding sites in tobacco leaf protoplasts transfected with *p35S:GmWRI1* (teal) or *p35S:mCHERRY* (grey). For (*C*) and (*D*), average firefly to Renilla luciferase activity ratios were multiplied by 100 and plotted with the respective standard errors. Lower-case letters indicate the significance of comparisons based on ANOVA and post hoc Tukey test results.

Figure 5. GmWRI1 and GmLEC1 act independently to control storage lipid biosynthesis in soybean embryos.
(*A*) Bar graph shows the number of FAT GmWRI1 DT genes that are bound or not bound by GmLEC1. The *P* value significance of the overlap is indicated (hypergeometric distribution). (*B*) Box plots comparing the distribution of mRNA levels in em embryos (GEO accession GSE99571) of FAT genes that are not bound by either GmWRI1 and GmLEC1 (grey), GmWRI1 DT genes that are not bound by GmLEC1 (teal), and GmWRI1 DT genes that are bound by GmLEC1 (dark red). Asterisk indicates significant differences between groups (Wilcoxon rank-sum test, p < 0.01). (*C*) GmWRI1 ChIP-Seq peak positions relative to LEC1 ChIP-Seq peaks (Position 0) in FAT GmWRI1 DT genes that are also bound by GmLEC1. (*D*) DNA motif enrichment in GmWRI1 DT binding sites of genes that are not bound or bound by GmLEC1. Circle diameters depict the frequencies at which DNA motifs were identified in the indicated regions, and the color intensity indicates the statistical significance of the enrichment relative to the normal distribution of a population of randomly generated regions (Bonferroni-adjusted *P* values).

Figure 6. Model for the regulation of storage lipid accumulation in embryos.
WRI1 and LEC1 act in a transcriptional positive feedback subcircuit to regulate FA and TAG biosynthesis in the embryo. GmWRI1 regulates FA and TAG-related gene expression by binding with AW Box and, possibly, CNC Box elements located downstream of the TSS. An alternative model is that another TF, such as a TCP-like TF, binds the CNC Box, potentially by interacting with WRI1. GmLEC1 binds FAT genes at sites that are distinct from GmWRI1 binding sites. GmLEC1 interacts with bZIP TFs that bind with G Box DNA elements located most frequently in regions upstream of the TSS. Dashed arrows indicate transcriptional activation.