The effect of dietary fish oil replacement by microalgae on the gilthead sea bream

midgut bacterial microbiota

Stefanos Katsoulis-Dimitriou[1], Eleni Nikouli[1], Elli Zafeiria Gkalogianni[1], Ioannis T.
Karapanagiotidis[1], Konstantinos A. Kormas[1,2*]

[1] Department of Agriculture Ichthyology & Aquatic Environment, University of Thessaly,
Volos, Magnesia, Greece

[2] Agricultural Development Institute, University Research and Innovation Centre "IASON",
Argonafton & Filellinon, 38221 Volos, Greece

* Corresponding author: kkormas@uth.gr

ABSTRACT

It is well known that the gut microbiome and its interaction with the host influence several important factors for fish health such as nutrition and metabolism. Diet is one of the main factors influencing the composition of the gut microbiome in reared fish. Microalgae, due to their high fatty acid content, appear to be a promising alternative for replacing fish oil in aquafeed. Thus, the aim of this work was to evaluate the effects of dietary microalgae blends as fish oil replacers on the midgut bacterial microbiota of the gilthead sea bream (*Sparus aurata*). The control diet (FO) contained only fish oil as source of lipids, EPA and DHA fatty acids, while three experimental diets were used where fish oil was replaced at 67% by one of the following microalgae biomass blends: *Microchloropsis gaditana* and *Isochrysis* sp. (*Tisochrysis lutea*) (MI), *Phaeodactylum tricornutum* and *Isochrysis* sp. (PI) and *Schizochytrium* sp. and *P. tricornutum* (SP). The midgut bacterial community composition of the experimental diets was altered compared to the control diet. There were 11 operational taxonomic units (OTUs) which were highly abundant in FO compared to the three experimental diets (FO,MI,SP) and two OTUs that were found in high abundance in both FO and the experimental diets in all comparisons (FO-MI, FO-PI, FO-SP). Most of the highly abundant OTUs in the experimental diets were unique to each experimental diet, with two OTUs being found in common between FO-MI and FO-PI. Additional evidence from the presumptive bacterial functional metabolic pathways suggested that the microalgae-based diets resulted in one over-expressed and one under-expressed pathway. The overexpressed pathway was related to the metabolism of fucose, a major cell wall exopolysaccharide of several microalgal species. Peptidoglycan biosynthesis was the under-expressed metabolic pathway. This suggests that a new gut microbiota profile was selected due to the microalgae inclusion in the provided diet. This study showed that, with the absence of mortality in fish, the gilthead sea bream gut microbiome can smoothly adapt its function according to the metabolic capacity of the dietary microalgae combinations that were used. The MI feed seems to promote several beneficial bacteria with potential probiotic abilities in the fish gut, belonging to the *Pseudoalteromonas, Pseudomonas, Bacillus* and *Rhodopseudomonas* genera.

Keywords: *Sparus aurata*, fish oil, aquaculture, microbiome, aquafeed

INTRODUCTION

       In reared fish, the composition of the gut microbiome is mainly determined by host

52   selection (Roeselers et al. 2011) and diet (Liu et al. 2016), while environmental and other

stochastic factors seem to be more important for wild fish (Kim et al. 2021, Kormas et al.

54   2023). The gut microbiome and its interactions with the host influence several important

factors for fish health such as nutrition and metabolism (Liu et al. 2021), maintaining the

56   development and maturation of the immune system (Koch et al. 2018), the nervous system

functioning and fish development (Phelps et al. 2017). Furthermore, commensal bacteria have

58   the potential to protect the fish against pathogens (Kelly & Salinas 2017) and when a disease

occurs there is a loss of gut microbiota diversity (Felix et al. 2022). Gut microbiota dysbiosis

60   is common in aquaculture species and is associated with changes in microbial structure, an

increase in pathogenic microorganisms and/or decreases in the abundance of beneficial taxa

62   (Xavier et al., 2023). For these reasons it is very important to investigate the changes that

may occur in the composition of the gut microbiota with changes in the fish's diet.

64   The main sources of proteins and lipids for carnivorous farmed fish, such as the

gilthead sea bream (*Sparus aurata)*, are fish meals and fish oils, respectively, which are

66   added in the aquafeeds in order to satisfy fish nutritional requirements and to provide high

nutritional quality to the final product (Siddik et al. 2023). Fish meals and fish oils originate

68   mainly from wild small pelagic fish species, making the aquaculture sector dependent on wild

fish stocks and, thus, reducing their sustainability (Péron et al. 2010). As a result, several

70   studies have focused on alternative sources of proteins and lipids to reduce the incorporation

of fish meals and fish oils in aquafeeds (Oliva-Teles et al. 2015, Naylor et al. 2021). Among

72   these alternatives, microalgae and oomycetes are suitable and sustainable feed ingredients.

Relevant studies showed that microalgae/oomycetes containing feeds have high digestibility,

74   positive effect on the growth rate of aquatic species, high quantities of important pigments

like carotenoids and phycobiliproteins, polysaccharides with potential antiviral and

76   antibacterial properties, potential immunostimulant and probiotic effects and low carbon

footprint for their production (Shah et al. 2017, Ahmad et al. 2022, Ma & Hu 2024).

78   *Microchloropsis gaditana* (Fawley, Jameson & Fawley 2015; formerly

Nannochloropsis gaditana) is a marine microalga of the class Eustigmatophyceae with high

80   content of eicosapentaenoic acid (EPA) (usually 36% of total fatty acids) (Zanella & Vianello

2020). *Phaeodactylum tricornutum* is a marine diatom of the class Bacillariophyceae, rich in

82   EPA (29% of total fatty acids) (Matos et al. 2016). The oomycete genus *Schizochytrium*

belongs to the Thraustochytriaceae family. Members of this genus are especially rich in

84   docosahexaenoic acid (DHA) (41% of total fatty acids) (Karapanagiotidis et al. 2022). The

marine microalga *Tisochrysis lutea* (here referred to as *Isochrysis* sp. due to its

86   reclassification (Bendif et al. 2013) also produces high levels of DHA (49% of total fatty

acids) (Ren et al. 2010).

88   Existing research has evaluated the use of *P. tricornutum* and *M. gaditana* in

alternative feeds for the gilthead sea bream. In particular, *P. tricornutum* when used in

90   experimental feeds, either alone or in combination with *Tetraselmis chui* and *Bacillus*

subtilis, has been linked with reduced gut microbial diversity in gilthead sea bream

92   (Cerezuela et. al 2012). However, the above combination along with dietary inulin and the

utilization of its b-glycans, have demonstrated immunostimulant effects, influencing

94   intestinal gene expression (Cerezuela et al. 2013; Reis et al. 2021; Teixeira et al. 2022). The

use of *M. gaditana* showed that gut microbiota composition and intestinal integrity remained

96   unaltered (Cerezo-Ortega et al. 2021) or an increase in intestinal microbiota richness without

affecting intestinal morphology and function (Jorge et al. 2019). When used in combination

98   with other species (*Chlorella* sp., *Arthrospira* sp., *Tisochrysis lutea*, *Scenedesmus*

almeriensis) it was shown to favour the presence of probiotic bacteria in the gut and reduce

100   the presence of pathogenic species (Garcia-Márquez et al. 2023; Garcia-Márquez et al. 2023).

To the best of our knowledge there are no studies about the effects of *Isochrysis* sp. and

102   *Schizochytrium* sp. on the gilthead sea bream gut microbiota.

        These microalgal species when combined in fish diets, appear suitable for replacing

104   the levels of EPA and DHA typically provided by fish oils. To our knowledge, there are no

studies evaluating the potential differences on the midgut microbial composition and function

106   of gilthead seabream when fed with a diet containing these microalgae species in

combination. Thus, the aim of this study was to investigate the potential effect on the midgut

108   microbial composition and function of gilthead seabream when fed a diet containing

combination of *Phaeodactylum tricornutum, Isochrysis* sp.*, Schizochytrium* sp. and

110   *Microchloropsis gaditana* inclusions. It is hypothesised that microalgal-based diets will

promote gut microbiota related to the metabolism of these microalgae.

MATERIALS AND METHODS

Sampling. For the needs of the present study, 40 *Sparus aurata* specimens weighing 28-37g were collected from a feeding trial (Gkalogianni et al. 2023) conducted by FELASA

accredited scientists at the experimental facilities of the Department of Agriculture, Ichthyology and Aquatic Environment registered for the use of experimental fish for

scientific purposes (EL-43BIO/exp-01). Fish were obtained from a commercial hatchery (8 g of initial mean weight) and reared for totally 80 days into glass tanks within a closed seawater

recirculation system (21.0±1.0 °C, pH at 8.0±0.2, salinity at 33±0.5 g/L, dissolved oxygen >6.5 mg/L, total ammonia nitrogen <0.1 mg/L. Fish were divided into 4 nutritional groups in

triplicate tanks with each group fed with a different diet. The four diets were isoproteic (48%), isolipidic (15.5%) and isoenergetic (21 Mj/Kg) and satisfied the EPA+DHA

requirements of the species. The control diet (FO) contained 80 g/Kg fish oil serving as a sole source of EPA and DHA, 40 g/Kg soybean oil and 250 g/Kg fishmeal with a EPA+DHA

level at 2.8 g/Kg . Three other diets were formulated replacing 50% of the dietary fish oil level of the control diet by a blend of microalgae biomasses serving as sources of EPA and

DHA: The PI diet contained 173 g/Kg *P. tricornutum* and 103 g/Kg *Isochrysis* sp. with EPA+DHA level at 2 g/Kg, the SP diet contained 39 g/Kg *Schizochytrium* sp. and 174 g/Kg

P. tricornutum with EPA+DHA level at 2.6 g/Kg, and MI diet contained 137 g/Kg *M. gaditana* and 103 g/Kg *Isochrysis* sp. with EPA+DHA level at 1.9 g/Kg. Since the

microalgae used in the experimental diets were in the form of dry biomass (and not extracted oils), their dietary inclusion contributed to the overall protein content. For this reason, the

percentage of inclusion of fishmeal was replaced accordingly (at 44% in PI diet, 27% in SP diet and 36% in MI diet). Diets were pelletized at 1.5 mm diameter according to

Karapanagiotidis et al. (2022). Fish promptly accepted the feeds and survival was high (99%) in all groups. At the end of the dietary trial, 10 individuals were randomly collected from

each dietary group, euthanized with a strong dose of the anesthetic 2-phenoxyethanol (450 mg l^{-1}, 15 min) and placed directly on ice. The body mass growth parameter L/W^3 of the

sampled fish based on weight (W) and length (L) ranged between 0.013±0.001 (FO) and 0.012±0.002 (MI) (Karapanagiotidis & Kormas unpubl. data). For the gut microbiota

analysis, the whole intestine of each individual was removed in aseptic conditions, with scissors and scalpel after sterilization with 70% ethanol before each dissection and the midgut

part was stored at -80 °C until the DNA extraction procedure.

     DNA extraction and sequencing. The DNA of seven gut samples of each feeding

148 group (FO, PI, SP, MI) was extracted using the Qiagen DNeasy PowerSoil Pro Kit (Qiagen, Hilden, Germany), according to the manufacturer's instructions. Bacterial communities were

150   analysed by 16S rRNA gene amplicon sequencing targeting the V3-V4 hypervariable region with the primer pair S-D-Bact-0341-b-S-17 and S-D-Bact-0785-a-A-21 with T_m at 55^{0}C

152   (Klindworth et al. 2013). PCR amplification and sequencing was performed at MRDNA Ltd.1 (Shallowater, TX, United States) facilities on a MiSeq using paired end reads (2×300

154   bp) following the manufacturer's guidelines.

        Data analysis. The 16S rRNA gene sequencing raw data were processed using the

MOTHUR MiSeq standard of protocol procedure (Schloss et al. 2013); MOTHUR is a stand-alone bioinformatics platform covering the entire procedure from raw sequencing data to

bacterial taxa 16S rRNA gene abundances. After trimming of the primer sequences 'screen.seqs' command was used with the parameters maxambig=0, minlength=331,

maxhomop=8. The 'split.abund' command (cutoff = 1) was used to exclude the rare sequences. The operational taxonomic units (OTUs) at 97% cutoff similarity level were

classified with the SILVA database release 138.1 (Quast et al. 2013, Yilmaz et al. 2014). The VSEARCH algorithm (Rognes et al. 2016) was used to detect and remove chimeric reads.

Sequences assigned as mitochondria or chloroplasts were removed from subsequent analysis. The 'sub.sample' command was used and the data were normalized to a depth of 2859 reads

per sample maintaining, thus, a sufficient number of reads per sample but at the sample time excluding only two samples with ≤2000 reads. Raw sequence reads have been deposited in

the Short Read Archive (https://www.ncbi.nlm.nih.gov/sra) under the BioProject accession number PRJNA1068122. Identification of the closest relatives of the most abundant OTUs

was performed using a Nucleotide Blast (http://blast.ncbi.nlm.nih.gov). The search was conducted with the following parameters: Standard databases, highly similar sequences

(Megablast) and closest relatives were considered with percentage identity higher than 90%. The statistical analysis and graphic illustrations were performed using PAleontological

STudies (PAST) software v.4.16 (Hammer et al. 2001). The input matrix for all statistical analyses was the OTUs table of each sequenced sample.

Presumptive functional analysis of midgut microbiota. The PICRUSt2 v2.5.1 (Phylogenetic Investigation of Communities by Reconstruction of Unobserved STates)

(Douglas et al. 2020) software was used to predict the microbiota functional profile (Langille et al. 2013). The predicted functions were searched in the MetaCyc database (Caspi et al.

2013), to be assigned to each metabolic pathway. Also, the KEGG (Kyoto Encyclopedia of Genes and Genomes) database (Kanehisa & Goto 2000, Kanehisa et al. 2023) was used to

identify whether the genera of the most important OTUs had all the enzymes for specific metabolic pathways of interest.

RESULTS

Sequencing data and OTUs. After the quality filtering of the amplicon sequencing
of the 16S rRNA gene V3–V4 region, a total of 319.103 reads remained. The number of
reads ranged from 1.631 to 44.124 reads per sample. A subsample of 2.859 reads per sample
was used for the normalization of the data. As a result, two samples (one from each of the FO
and PI treatments) with only 1.631 and 1.867 reads, respectively, were excluded from the rest
of the analyses. The total number of OTUs to which the reads were taxonomically assigned
was 519.

Alpha diversity. The Shannon_H and Simpson_1-D indices were calculated to assess
the alpha diversity of the gut microbiota of gilthead sea bream in the four groups (Table 1).
Also, Taxa_S was calculated for the estimation of the OTUs richness and Chao-1 to estimate
the diversity by calculating the potentially missed diversity using the presence of singletons
and doubletons (Table 1). Compared to the control (FO), Shannon_H and Simpson_1-D
values showed a slight increase in PI, while they decreased in MI and SP. Chao-1 index
values showed a decrease compared to FO in all three groups, while Taxa_S increased in PI
and SP and showed a slight decrease in MI. Despite the existing differences, the Bray-Curtis
PERMANOVA (permutational analysis of variance) performed with 9999 permutations
(Table S1) showed no statistically significant differences between the indices values of the
three experimental groups with FO. The only statistically significant difference was observed
in Taxa_S ($p=0.046$) and Shannon_H ($p=0.049$) between PI and MI.

Beta diversity. Non-metric multidimensional scaling analysis (nMDS) was used to
evaluate the beta diversity of the samples of the four feeding groups, based on a Bray-Curtis
distance matrix (Figure 1). Samples from the MI groups showed the highest variation. The PI
group was clearly separated from the samples of the FO and SP group. The samples of the
FO, MI and SP groups overlapped and there was overlap between the samples of the PI and
MI groups. Group PI was the only treatment with no overlap with FO. PERMANOVA
pairwise test (Table S2) was used to further analyze the differences between dietary groups.
Significant differences were observed between the PI and FO and between the PI and SP
groups.

The unique OTUs in FO, PI, SP and MI groups (Figure S1a) were 40 (7.7% of all
OTUs), 61 (11.8%), 148 (28.5%) and 56 (10.8%) respectively. In total, 11.2% of the OTUs

218   were found in all groups (FO, PI, SP, MI). Furthermore, there were 91 unique OTUs in FO

compared to PI (30%), 95 unique OTUs compared to MI (34.2%) and 76 unique OTUs

220   compared to SP (19.5%) (Figure S1b). Shared OTUs were in a higher proportion between FO

and MI (34.5%) with FO and PI being very close (33%) and, lastly, FO and SP (29%). SP had

222   the highest number of unique OTUs (199 OTUs - 51%) compared to FO, with MI and PI

having smaller contributions of unique OTUs (87 OTUs - 31.3% and 112 OTUs -37%,

224   respectively).

           Midgut bacterial taxonomic composition. Proteobacteria and Actinobacteriota were

226   within the three most abundant phyla in all dietary groups (Figure S2). In FO and MI, the

third most abundant phylum was Firmicutes, while in PI and SP the third most abundant

228   phylum was Verrucomicrobiota and Bacteroidota, respectively. In FO, MI and SP, despite

differences in either the abundance rank of the phyla (FO-MI) or the phyla presence (FO-SP),

230   the ratios of the phyla abundances were quite close (Figure S2). In contrast, in PI there was a

large increase in the relative abundance of Proteobacteria, since Proteobacteria were x2.1

232   more abundant than Actinobacteriota and x8.1 than Verrucomicrobiota. Actinobacteriota also

had a large difference in relative abundance with the third most abundant phylum,

234   Verrucomicrobiota (x3.8). Regarding other phyla, only in MI the Fusobacteriota occurred

with a relative abundance of 4.3% and in PI and SP the Firmicutes with relative abundance of

236   5.9% and 13.9%, respectively.

           In all feeding groups the Mycobacteriaceae was most abundant family in the FO

238   (27.6% of reads), MI (28.4% of reads) and SP (27.2% of reads) and second most abundant in

PI with relative abundance 22.0% of reads (Figure 2). In the PI treatment, the most abundant

240   family was the Rhodobacteraceae (31.7% of reads), which in MI and SP occurred in very low

abundances (2.9% and 1.3% of reads, respectively). These abundances were due to the most

242   dominant OTUs, which in FO, MI and SP was OTU001 (27.6%, 28.3%, 27.1% of reads

respectively) and OTU004 in PI with relative abundance 23.9% (Table S3). Nucleotide

244   BLAST search assigned OTU001 as *Mycolicibacterium aurum* and OTU004 as *Yoonia*

litorea. Compared to FO, in the other feeding groups the family Rhizobiaceae was not present

246   and the family Xanthomonadaceae was only present in MI (<4% relative abundance). The

family Vibrionaceae occurred in MI (8.3% of reads) and PI (12.3% of reads) but it was absent

248   in SP and FO. Treatment MI had the highest number of families contributing >4% relative

abundance (nine families), while SP had the lowest (five families).

250   **Most important OTUs**. In the dominant OTUs (cumulative relative dominance >

80% - Figure 3) three OTUs occurred in all feeding groups (OTU001, OTU003, OTU012).

OTU003 and OTU012 occurred in small proportions in PI (1.67% and 2.44%). From the Nucleotide BLAST search, the OTU003 closest relative was the uncultured *Staphylococcus*

sp. and OTU012 closest relative was *Cutibacterium acnes*. OTU004, which was the most dominant in PI, was among the most dominants of FO and MI, but with low relative

abundances (2.95% and 2.16% respectively). Within the dominant OTUs there were several unique ones that occurred in each feeding group.

The qualitative differences already observed in phyla and families were also observed at the level of dominant OTUs in each group. To explore this further, we compared key

OTUs between the FO and the three experimental groups. This comparison unveiled which OTUs occurred in high abundance and were common in both the FO and experimental feeds,

or exclusive to the FO (Figure 4, Table S4). 11 common OTUs were highly abundant in FO compared to MI, PI and SP. Furthermore, among the highly abundant OTUs, two of them

(OTU001 and OTU003) were observed in all treatments. Most of the highly abundant OTUs in MI, PI and SP were unique to each experimental feed, with only two OTUs (OTU006,

OTU027) being common to FO-MI and FO-PI.

Presumptive functional pathways. Compared to FO, one metabolic pathway was

considerably overexpressed (FUCCAT-PWY with values: LOG(tax.function.abundance PI/FO) = 1.988, LOG(tax.function.abundance MI/FO) = 1.988, LOG(tax.function.abundance

SP/FO) = 1.536) and one considerably underexpressed (PWY-6470 with values: LOG(tax.function.abundance PI/FO) = -1.511, LOG(tax.function.abundance MI/FO) = -

1.618, , LOG(tax.function.abundance SP/FO) = -1.509) in all three experimental feeding groups (Figure 5, Table S5). According to the MetaCyc database, the FUCCAT-PWY

pathway corresponds to L-fucose degradation, and the PWY-6470 pathway corresponds to peptidoglycan biosynthesis. Several common pathways were overexpressed in MI and PI

relatively to FO and the highest number of unique pathways were overexpressed in SP. From the KEGG database it was found that the genomes of most of the OTUs in each of the three

experimental feeds (see Fig. 4) contained all the genes encoding for the enzymes of the L-Fucose degradation pathway (Table S6).

DISCUSSION

In this study, it was hypothesized that the inclusion of microalgal-based diets in experimental reared gilthead sea bream would induce structural bacterial microbiota changes

adapted to these novel feed ingredients. In similar studies with the use of microalgae in the

      diet of gilthead sea bream, variable results have been reported. Cerezuela et al. (2012)

288   reported that the dietary inclusion of 100g/kg *Tetraselmis chui* or *P. tricornutum* decreased

      the intestinal microbial richness with a lower Shannon index value than the control. In the

290   current study, the Shannon index decreased in fish fed with the *Schizochytrium - P.

      tricornutum* (SP) diet but slightly increased in fish fed with the *P. tricornutum – Isochrysis*

292   diet. Nevertheless, these two diets had more OTUs (Taxa_S) than the control (FO). Jorge et

      al. (2019) using *M. gaditana* for fishmeal replacement reported a higher, but insignificant to

294   the control, intestinal richness and diversity of the microbiota. Using a microalgae blend of

      Tisochrysis lutea (member of the *Isochrysis* spp. group), *M. gaditana* and *Scenedesmus*

296   *almeriensis* at 5%, 15% and 25% dietary inclusion levels, Garcia-Marquez et al. (2023) found

      that the microalgae blend induced an increase in bacterial species diversity and a distinct shift

298   in microbiota fingerprinting as inclusion levels increased. In our study, compared to the

      control (FO), the diet containing *M. gaditana* (MI) showed lower alpha diversity indices, but

300   this was not statistically significant. Similar lack of statistically significance has been

      reported in the *Sparus aurata* gut microbiota after the inclusion of 5% hydrolysed *M.

302   gaditana* (Cerezo-Ortega et al. 2021).

            Studies with dietary inclusion of *Schizochytrium* sp. in Nile tilapia (Souza et al. 2020),

304   zebrafish (Shi et al. 2021) and rainbow trout (Lyons et al. 2016) also showed varying results.

      Specifically, in Nile tilapia a diet with 1.2% inclusion of *Schizochytrium sp.* led to higher

306   Inverse Simpson and Shannon values in the *Schizochytrium*-fed fish, but these were not

      significantly different to those in the control group. In zebrafish, the dietary inclusion at 0, 60

308   and 120g/kg of *Schizochytrium sp.* resulted in insignificant differences in Shannon index

      among the three groups, but in the beta diversity nMDS showed that both the *Schizochytrium

310   sp.* supplemented diets were separated from the control group (Shi et al. 2021). The diet

      containing Schizochytrium sp. in our study (SP) had lower values in Shannon-H and

312   Simpson_1-D indices than the control (FO), but these were not statistically significant and

      also there were no statistically significant differences in beta diversity compared to the

314   control. Furthermore, a 5% inclusion of *Schizochytrium limacinum* for fish oil replacement in

      the diet of rainbow trout resulted in a greater level of microbial diversity (Lyons et al. 2016; a

316   fact that was also observed in our results using the SP diet that had a higher OTUs (Taxa_S)

      index number than the control (FO).

318   Overall, the effect on gut microbiota diversity is primarily determined by the different

      species of microalgae, the fish as the host and the levels of inclusion of each microalga in the

diet. The literature as discussed above in combination with the results of our study shows the various changes that occur as the gut microbiome adapts to the different components added

by microalgae to the diet. However, differences in diversity alone cannot show the overall effects and investigation of changes at the level of bacterial phyla, families, and specific

OTUs that modulate the diversity is required. As it has been shown that the phycoshpere of cultivated microalgae used as feed might affect the fish gut microbiota in early

developmental stages (e.g. Nikouli et al. 2019), in the future the impact of the bacteria contained in the microalgae to be incorporated in the aquafeed should be investigated as well.

Most important OTUs. The dominant phyla found (Proteobacteria, Actinobacteriota, Bacteroidota) are common in the gilthead seabream gut microbiome (Kormas et al. 2014,

Nikouli et al. 2018, 2019, Cerezo-Ortega et al 2021) and also in other fish species gut microbiome (Llewellyn et al. 2016, Rosenau et al. 2021, Zhang et al. 2022). In PI, however,

there was an increased occurrence of Verrucomicrobiota. The Mycobacteriaceae family is the one that is highly abundant in all groups, due to OTU001, which belongs to the genus

Mycolicibacterium. From the available literature it seems that this genus is not common in the gilthead sea bream gut. It includes some species of pathogenic micro-organisms for

humans (*Mycobacterium tuberculosis, Mycobacterium leprae*) but also several species that inhabit a diverse range of environments including water bodies, soil, and metalworking fluids

(Gupta, Lo & Son 2018). *Mycobacterium* genus has also been found in the gut of freshwater species (Gallet et al. 2022). The closest relative, *M. aurum* has never been reported as a

pathogen, but its possible functions in the gut remain unknown. The Rhodobacteraceae family member OTU004 is the dominant in the PI treatment, closely related to *Yoonia litorea*

(Paracocacceae family). *Y. litorea* though has been recently classified in the *Roseobacter* clade within the Rhodobacteraceae family (Wirth & Whitman 2018). Later, it was proposed

the family Roseobacteraceae for members of the *Roseobacter* clade (Liang et al. 2021). Misclassifications have been a recurring problem between the Paracocacceae and

Roseobacteraceae families (Zhang et al. 2023), but *Yoonia* sp. isolates are phylogenetically placed in the Roseobacteraceae family (Feng & Xing 2023). Species from the *Roseobacter*

clade have been isolated from seawater marine sediments and biofilms, and are often associated with phytoplankton, macroalgae and marine animals (Wirth & Whitman 2018).

They have also been found in intestine of finfish and they seem to have probiotic effects (Ringø et al. 2022). Specifically, for the *Yoonia-Loktanella* group it has been speculated that

they offer defensive secondary metabolites to deep-sea sponges (Lo Giudice et al. 2024).

OTU003 was highly abundant in FO and MI, PI and SP and its closest relative is

354   *Staphylococcus* sp., while OTU012 was highly abundant in FO, MI and PI and closely related

to *Cutibacterium acnes*. *Staphylococcus* sp. is a common bacterial genus in healthy fish gut

356   microbiomes and some species may have probiotic effects (Zorriehzahra et al.2015, Borah et

al. 2016, El-Saadony et al. 2021). *Cutibacterium acnes* has been found in other fish species

358   and it has been suggested to have potential probiotic features (Nikouli et al. 2021). OTU006

and OTU027, that belong to the *Vibrio* genus, were in high abundance in MI and PI

360   compared to FO. *Vibrio* is a common bacterial genus found in fish gut with both pathogenic

and non-pathogenic species. Some *Vibrio* species can act as symbionts assisting in the

362   breakdown of dietary components, with strains found to produce amylase, cellulose and

chitinase among others (Egerton et al. 2018). From the rest top abundant OTUs, in MI

364   OTU013, OTU031 and OTU041 were closely related to *Pseudoalteromonas* sp.*,*

Pseudomonas sp. and *Bacillus* sp. respectively. Several species of these genera are known to

366   have probiotic effects in the fish gut (Mladineo et al. 2016, Zorriehzahra et al. 2015, Salam et

al. 2021). OTU030 was abundant in FO and MI and was related to *Rhodopseudomonas* sp., a

368   genus that is also reported to have probiotic effects in the fish gut (Koga et al. 2022). Thus,

from the point of structural changes of the bacterial communities, the MI experimental feed

370   seems to promote several beneficial bacterial species in the gut of gilthead sea bream.

L-fucose degradation was the only considerably overexpressed bacterial metabolic

372   pathway in the three experimental feeds in comparison to FO. Fucose is a deoxyhexose that is

found as a constituent of exopolysaccharides from microalgae (Ahmed et al. 2013, Liu et al.

374   2019, Wan et al. 2019). It is also commonly found in animal cell glycans, macroalgae,

cyanobacteria and microbial exopolysaccharides (Becker & Lowe 2003). Furthermore, search

376   of the KEGG database confirmed that most of the important bacterial genera found are

affiliated to genomes with all five enzymes to carry out the L-fucose degradation. This

378   suggests that the microalgal-based diets induced a clear gut bacterial response, at the

functional level, favoring the proliferation of bacteria which benefit from specific microalgal

380   compounds with potential benefits to the host, too. The degradation of fucose is a likely

beneficial-to-the-host service. For example, in humans fucose degraders are known to induce

382   the production of the beneficial short-chain fatty acids (Morrison et al. 2016, Høgsgaard et al.

2023, Fusco et al. 2023). The metabolic fate of the fucose degradation and its possible effect

384   on the sea bream's physiology remains to be specifically investigated in the future.

The only pathway that was considerably under-expressed in the three experimental

386   feeds compared to FO is peptidoglycan biosynthesis. The cell wall of some microalgae

species contains peptidoglycan (Agboola et al. 2019, Machado et al. 2022), but the species used in our research do not seem to contain it (Domozych et al. 2012, Le Costaouëc et al. 2017, Nadzir et al. 2023). Peptidoglycan forms around 90% of the dry weight of Gram-positive bacteria but only 10% of Gram-negative strains (Malanovic & Lohner 2016). The relative abundance of Gram-negative phyla and families like Vibrionaceae in PI and MI and Bacteroidota in SP is higher than FO. Consequently, the putative under-expression of peptidoglycan synthesis with the three diets compared to FO is likely representative of the higher presence of Gram- taxa. Finally, SP was the only treatment with unique overexpressed pathways along with the highest number of unique OTUs compared to FO which, however, only one of the treatment's unique OTUs are included in these overexpressed pathways (Table S7). This microalgal feed inclusion requires further investigation with in vitro experiments using specific isolates and growth media to test the functionality of these presumptive metabolic pathways.

In conclusion, this study showed that the inclusion of three microalgal-based diets induced both structural and functional changes in the sea bream's gut microbiome. The structural changes in each dietary treatment were considerable compared to the control diet. The putative under-expression of peptidoglycan synthesis with the 3 diets compared to FO was likely representative of the higher presence of Gram- strains. However, the total midgut bacterial communities which were favoured by the microalgal inclusion were inferred to be able to metabolise L-fucose, a major microalgal carbohydrate.

AKNOWLEDGMENTS

This work was financially supported by the English MSc Program "Host-Microbe Interactions of the University of Thessaly and co-financed by the European Union and Greek co-funded project entitled "Use of insect protein and microalgal oil for fishmeal and fish oil replacement in the diets of gilthead seabream (*Sparus aurata*) and European seabass (*Dicentrarchus labrax*) under national funds through the Operational Programme "Competitiveness, Entrepreneurship and Innovation-EPAnEK 2014-2020" (project code: MIS 5045804)).

420 **Table 1.** Alpha diversity metrics (Shannon H and Simpson 1-D), OTUs richness and Chao-1
values for each treatment. OTUs: operational taxonomic units.

Dietary treatment	Number of OTUs (Taxa_S)	Shannon H	Simpson 1-D	Chao-1
FO: Fish oil	53 ± 9.4	2.02 ± 0.570	0.75 ± 0.170	152.6 ± 85.38
PI: *Phaeodactylum + Isochrysis*	57 ± 14.2	2.07 ± 0.366	0.77 ± 0.081	115.7 ± 26.74
SP: *Schizochytrium + Phaeodactylum*	64 ± 63.3	1.80 ± 0.984	0.65 ± 0.265	123.2 ± 70.58
MI: *Microchloropsis + Isochrysis*	44 ± 8.0	1.41 ± 0.657	0.56 ± 0.254	104.3 ± 37.40

REFERENCES

Agboola, Jeleel O., Teuling, Emma, Wierenga, Peter A., Gruppen, Harry, and Schrama, Johan W., 'Cell Wall Disruption: An Effective Strategy to Improve the Nutritive Quality of Microalgae in African Catfish (*Clarias gariepinus*)', Aquaculture Nutrition, 25/4 (2019), 783–97

Ahmad, Ashfaq, W. Hassan, Shadi, and Banat, Fawzi, 'An Overview of Microalgae Biomass as a Sustainable Aquaculture Feed Ingredient: Food Security and Circular Economy', Bioengineered, 13/4 (2022), 9521–47

Ahmed, Mehboob, Moerdijk-Poortvliet, Tanja C.W., Wijnholds, Anita, Stal, Lucas J., and Hasnain, Shahida, 'Isolation, Characterization and Localization of Extracellular Polymeric Substances from the Cyanobacterium *Arthrospira platensis* Strain MMG-9', European Journal of Phycology, 49/2 (2014), 143–50

Becker, Daniel J., and Lowe, John B., 'Fucose: Biosynthesis and Biological Function in Mammals', Glycobiology, 13/7 (2003), 41R-53R

Bendif, E.M., Probert, I., Schroeder, D.C. and de Vargas, C. (2013) On the description of *Tisochrysis lutea* gen. nov. sp. nov. and *Isochrysis nuda* sp. nov. in the Isochrysidales, and the transfer of Dicrateria to the Prymnesiales (Haptophyta). *Journal of Applied Phycology* **25**, 1763-1776.

Borah, Debajit, Gogoi, Olee, Adhikari, Chanakya, and Kakoti, B. B., 'Isolation and Characterization of the New Indigenous *Staphylococcus* Sp. DBOCP06 as a Probiotic Bacterium from Traditionally Fermented Fish and Meat Products of Assam State', Egyptian Journal of Basic and Applied Sciences, 3/3 (2016), 232–40

Caspi, Ron, Billington, Richard, Ferrer, Luciana, Foerster, Hartmut, Fulcher, Carol A., Keseler, Ingrid M., et al., 'The MetaCyc Database of Metabolic Pathways and Enzymes and the BioCyc Collection of Pathway/Genome Databases', Nucleic Acids Research, 44/D1 (2016), D471-480

Cerezo-Ortega, I. M., Di Zeo-Sánchez, D. E., García-Márquez, J., Ruiz-Jarabo, I., Sáez-Casado, M. I., Balebona, M. C., et al., 'Microbiota Composition and Intestinal Integrity Remain Unaltered after the Inclusion of Hydrolysed *Nannochloropsis gaditana* in *Sparus aurata* Diet', Scientific Reports, 11/1 (2021), 18779

Cerezuela, Rebeca, Fumanal, Milena, Tapia-Paniagua, Silvana Teresa, Meseguer, José, Moriñigo, Miguel Angel, and Esteban, Ma Angeles, 'Histological Alterations and Microbial Ecology of the Intestine in Gilthead Seabream (*Sparus aurata* L.) Fed Dietary Probiotics and Microalgae', Cell and Tissue Research, 350/3 (2012), 477–89

Cerezuela, Rebeca, Meseguer, José, and Esteban, M. Ángeles, 'Effects of Dietary Inulin, Bacillus Subtilis and Microalgae on Intestinal Gene Expression in Gilthead Seabream (*Sparus aurata* L.)', Fish & Shellfish Immunology, 34/3 (2013), 843–48

Domozych, David S., Ciancia, Marina, Fangel, Jonatan U., Mikkelsen, Maria Dalgaard, Ulvskov, Peter, and Willats, William G. T., 'The Cell Walls of Green Algae: A Journey through Evolution and Diversity', Frontiers in Plant Science, 3 (2012), 82

Douglas, Gavin M., Maffei, Vincent J., Zaneveld, Jesse R., Yurgel, Svetlana N., Brown, James R., Taylor, Christopher M., et al., 'PICRUSt2 for Prediction of Metagenome Functions', Nature Biotechnology, 38/6 (2020), 685–88

Egerton, Sian, Culloty, Sarah, Whooley, Jason, Stanton, Catherine, and Ross, R. Paul, 'The Gut Microbiota of Marine Fish', Frontiers in Microbiology, 9 (2018) , 873

El-Saadony, Mohamed T., Alagawany, Mahmoud, Patra, Amlan K., Kar, Indrajit, Tiwari, Ruchi, Dawood, Mahmoud A. O., et al., 'The Functionality of Probiotics in Aquaculture: An Overview', Fish & Shellfish Immunology, 117 (2021), 36–52

Fawley, Marvin W., Jameson, Ian, and Fawley, Karen P., 'The Phylogeny of the Genus *Nannochloropsis* (Monodopsidaceae, Eustigmatophyceae), with Descriptions of *N. australis* sp. nov. and *Microchloropsis* gen. nov.', Phycologia, 54/5 (2015), 545–52

Feng, Xiaoyuan, and Xing, Peng, 'Genomics of *Yoonia* sp. Isolates (Family Roseobacteraceae) from Lake Zhangnai on the Tibetan Plateau', Microorganisms, 11/11 (2023), 2817

Fusco, William, Lorenzo, Manuel Bernabeu, Cintoni, Marco, Porcari, Serena, Rinninella, Emanuele, Kaitsas, Francesco, et al., 'Short-Chain Fatty-Acid-Producing Bacteria: Key Components of the Human Gut Microbiota', Nutrients, 15/9 (2023), 2211

Gallet, Alison, Yao, Eric Kouamé, Foucault, Pierre, Bernard, Cécile, Quiblier, Catherine, Humbert, Jean-François, et al., 'Fish Gut-Associated Bacterial Communities in a Tropical Lagoon (Aghien Lagoon, Ivory Coast)', Frontiers in Microbiology, 13 (2022), 963456

García-Márquez, Jorge, Rico, Rosa María, Acién, Francisco Gabriel, Mancera, Juan Miguel, Figueroa, Félix L., Vizcaíno, Antonio Jesús, et al., 'Dietary Effects of a Short-Term Administration of Microalgae Blend on Growth Performance, Tissue Fatty Acids, and Predominant Intestinal Microbiota in *Sparus aurata*', Microorganisms, 11/2 (2023), 463

Gkalogianni, Elli Zafeiria, Psofakis, Pier, Asimaki, Adamantia, Fountoulaki, Eleni, Henry, Morgane, and Karapanagiotidis, Ioannis T. (2023). *Microchloropsis gaditana, Schizochytrium* sp., *Phaeodactylum tricornutum*, and *Tisochrysis lutea* as n-3 PUFA sources in the diet of juvenile gilthead seabream (*Sparus aurata*). Proceedings of Aquaculture Europe 2023, September 18 - 21, Vienna, Austria.

Gupta, Radhey S., Lo, Brian, and Son, Jeen, 'Phylogenomics and Comparative Genomic Studies Robustly Support Division of the Genus *Mycobacterium* into an Emended Genus *Mycobacterium* and Four Novel Genera', Frontiers in Microbiology, 9 (2018), 00067

Hammer, Oyvind, Harper, David, and Ryan, Paul, 'PAST: Paleontological Statistics Software Package for Education and Data Analysis', *Palaeontologia Electronica*, 4 (2001), 1–9

Høgsgaard, Karina, Vidal, Natalia P., Marietou, Angeliki, Fiehn, Oliver Gam, Li, Qing, Bechtner, Julia, et al., 'Fucose Modifies Short Chain Fatty Acid and H2S Formation through Alterations of Microbial Cross-Feeding Activities', *FEMS Microbiology Ecology*, 99/10 (2023), fiad107

Jorge, Sara S., Enes, Paula, Serra, Cláudia R., Castro, Carolina, Iglesias, Paula, Oliva Teles, Aires, et al., 'Short-Term Supplementation of Gilthead Seabream (*Sparus aurata*) Diets with *Nannochloropsis gaditana* Modulates Intestinal Microbiota without Affecting Intestinal Morphology and Function', *Aquaculture Nutrition*, 25/6 (2019), 1388–98

Kanehisa, M., and Goto, S., 'KEGG: Kyoto Encyclopedia of Genes and Genomes', *Nucleic Acids Research*, 28/1 (2000), 27–30

Kanehisa, Minoru, Furumichi, Miho, Sato, Yoko, Kawashima, Masayuki, and Ishiguro-Watanabe, Mari, 'KEGG for Taxonomy-Based Analysis of Pathways and Genomes', *Nucleic Acids Research*, 51/D1 (2023), D587–92

Karapanagiotidis, I. T., Metsoviti, M. N., Gkalogianni, E. Z., Psofakis, P., Asimaki, A., Katsoulas, N., et al., 'The Effects of Replacing Fishmeal by *Chlorella vulgaris* and Fish Oil by *Schizochytrium* sp. and *Microchloropsis gaditana* Blend on Growth Performance, Feed Efficiency, Muscle Fatty Acid Composition and Liver Histology of Gilthead Seabream (*Sparus aurata*)', *Aquaculture*, 561 (2022), 738709

Kelly, Cecelia, and Salinas, Irene, 'Under Pressure: Interactions between Commensal Microbiota and the Teleost Immune System', *Frontiers in Immunology*, 8 (2017), 559

Kim, Pil Soo, Shin, Na-Ri, Lee, Jae-Bong, Kim, Min-Soo, Whon, Tae Woong, Hyun, Dong-Wook, et al., 'Host Habitat Is the Major Determinant of the Gut Microbiome of Fish', *Microbiome*, 9/1 (2021), 166

Klindworth, Anna, Pruesse, Elmar, Schweer, Timmy, Peplies, Jörg, Quast, Christian, Horn, Matthias, et al., 'Evaluation of General 16S Ribosomal RNA Gene PCR Primers for Classical and Next-Generation Sequencing-Based Diversity Studies', *Nucleic Acids Research*, 41/1 (2013), e1

Koch, Bjørn E. V., Yang, Shuxin, Lamers, Gerda, Stougaard, Jens, and Spaink, Herman P., 'Intestinal Microbiome Adjusts the Innate Immune Setpoint during Colonization through Negative Regulation of MyD88', *Nature Communications*, 9/1 (2018), 4099

Koga, Aoi, Yamasaki, Takumi, Hayashi, Shuhei, Yamamoto, Shinjiro, and Miyasaka, Hitoshi, 'Isolation of Purple Nonsulfur Bacteria from the Digestive Tract of Ayu (*Plecoglossus altivelis*)', *Bioscience, Biotechnology, and Biochemistry*, 86/3 (2022), 407–12

Kormas, Konstantinos A., Meziti, Alexandra, Mente, Eleni, and Frentzos, Athanasios, 'Dietary Differences Are Reflected on the Gut Prokaryotic Community Structure of Wild and Commercially Reared Sea Bream (*Sparus aurata*)', *MicrobiologyOpen*, 3/5 (2014), 718–28

Kormas, Konstantinos, Nikouli, Eleni, Kousteni, Vasiliki, and Damalas, Dimitrios, 'Midgut Bacterial Microbiota of 12 Fish Species from a Marine Protected Area in the Aegean Sea (Greece)', *Microbial Ecology*, 86/2 (2023), 1405–15

Kozich, James J., Westcott, Sarah L., Baxter, Nielson T., Highlander, Sarah K., and Schloss, Patrick D., 'Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform', *Applied and Environmental Microbiology*, 79/17 (2013), 5112–20

Langille, Morgan G. I., Zaneveld, Jesse, Caporaso, J. Gregory, McDonald, Daniel, Knights, Dan, Reyes, Joshua A., et al., 'Predictive Functional Profiling of Microbial Communities Using 16S rRNA Marker Gene Sequences', *Nature Biotechnology*, 31/9 (2013), 814–21

Le Costaouëc, Tinaïg, Unamunzaga, Carlos, Mantecon, Lalia, and Helbert, William, 'New Structural Insights into the Cell-Wall Polysaccharide of the Diatom *Phaeodactylum tricornutum*', *Algal Research*, 26 (2017), 172–79

Liang, Kevin Y. H., Orata, Fabini D., Boucher, Yann F., and Case, Rebecca J., 'Roseobacters in a Sea of Poly- and Paraphyly: Whole Genome-Based Taxonomy of the Family Rhodobacteraceae and the Proposal for the Split of the "Roseobacter Clade" Into a Novel Family, Roseobacteraceae Fam. Nov.', *Frontiers in Microbiology*, 12 (2021), 683109

Liu, Han, Guo, Xianwu, Gooneratne, Ravi, Lai, Ruifang, Zeng, Cong, Zhan, Fanbin, et al., 'The Gut Microbiome and Degradation Enzyme Activity of Wild Freshwater Fishes Influenced by Their Trophic Levels', *Scientific Reports*, 6/1 (2016), 24340

Liu, Qishun, Yao, Changhong, Sun, Yongxin, Chen, Wei, Tan, Haidong, Cao, Xupeng, et al., 'Production and Structural Characterization of a New Type of Polysaccharide from Nitrogen-Limited *Arthrospira platensis* Cultivated in Outdoor Industrial-Scale Open Raceway Ponds', *Biotechnology for Biofuels*, 12/1 (2019), 131

Liu, Yaqiu, Li, Xinhui, Li, Jie, and Chen, Weitao, 'The Gut Microbiome Composition and Degradation Enzymes Activity of Black Amur Bream (*Megalobrama terminalis*) in Response to Breeding Migratory Behavior', *Ecology and Evolution*, 11/10 (2021), 5150–63

Llewellyn, Martin S., McGinnity, Philip, Dionne, Melanie, Letourneau, Justine, Thonier, Florian, Carvalho, Gary R., et al., 'The Biogeography of the Atlantic Salmon (*Salmo salar*) Gut Microbiome', *The ISME Journal*, 10/5 (2016), 1280–84

Lo Giudice, Angelina, Papale, Maria, Azzaro, MMaurizio, Rizzo, Carmen, 'Prokaryotic diversity in the sponges *Mycale* (*Oxymycale*) *acerata* (Kirkpatrick, 1907) and *Dendrilla*
antarctica (Topsent, 1905) from two distant Antarctic marine areas: South Cove at Rothera Point (Adelaide Island, Western Antarctic Peninsula) and Thetys Bay (Terra
Nova Bay, Ross Sea), *Deep Sea Research Part II: Topical Studies in Oceanography*, 216, (2024), 105391

Lyons, Philip P, Turnbull, James F, Dawson, Karl A, and Crumlish, Margaret, 'Effects of Low-Level Dietary Microalgae Supplementation on the Distal Intestinal Microbiome of
Farmed Rainbow Trout *Oncorhynchus mykiss* (Walbaum)', *Aquaculture Research*, 48/5 (2017), 2438–52

Ma, Mingyang, Hu, Qiang, 'Microalgae as feed sources and feed additives for sustainable aquaculture: Prospects and challenges', *Reviews in Aquaculture*, 16/2 (2024), 818-835

Machado, Luís, Carvalho, Gonçalo, and Pereira, Ricardo N., 'Effects of Innovative Processing Methods on Microalgae Cell Wall: Prospects towards Digestibility of
Protein-Rich Biomass', *Biomass*, 2/2 (2022), 80–102

Malanovic, Nermina, and Lohner, Karl, 'Antimicrobial Peptides Targeting Gram-Positive
590 Bacteria', *Pharmaceuticals*, 9/3 (2016), 59

Matos, Ângelo Paggi, Feller, Rafael, Moecke, Elisa Helena Siegel, Oliveira, José Vladimir
592 de, Junior, Agenor Furigo, Derner, Roberto Bianchini, et al., 'Chemical Characterization of Six Microalgae with Potential Utility for Food Application' *Journal*
594 *of the American Oil Chemists' Society*, 93/7 (2016) 963-72

Md Nadzir, Syafiqah, Yusof, Norjan, Nordin, Norazela, Kamari, Azlan, and Yusoff, Mohd
596 Zulkhairi Mohd, 'A Review of Microalgal Cell Wall Composition and Degradation to Enhance the Recovery of Biomolecules for Biofuel Production', *Biofuels*, 14/9 (2023),
598 979–97

Medina-Félix, Diana, Garibay-Valdez, Estefanía, Vargas-Albores, Francisco, and Martínez-
600 Porchas, Marcel, 'Fish Disease and Intestinal Microbiota: A Close and Indivisible Relationship', *Reviews in Aquaculture*, 15/2 (2023), 820–39

602 Mladineo, Ivona, Bušelić, Ivana, Hrabar, Jerko, Radonić, Ivana, Vrbatović, Anamarija, Jozić, Slaven, et al., 'Autochthonous Bacterial Isolates Successfully Stimulate In Vitro
604 Peripheral Blood Leukocytes of the European Sea Bass (*Dicentrarchus labrax*)', *Frontiers in Microbiology*, 7 (2016), 1244

606 Morrison, Douglas J., and Preston, Tom, 'Formation of Short Chain Fatty Acids by the Gut Microbiota and Their Impact on Human Metabolism', *Gut Microbes*, 7/3 (2016), 189–
608 200

Naylor, Rosamond L., Hardy, Ronald W., Buschmann, Alejandro H., Bush, Simon R., Cao,
Ling, Klinger, Dane H., et al., 'A 20-Year Retrospective Review of Global Aquaculture', *Nature*, 591/7851 (2021), 551–63

Nikouli, Eleni, Meziti, Alexandra, Antonopoulou, Efthimia, Mente, Eleni, and Kormas, Konstantinos A., 'Gut Bacterial Communities in Geographically Distant Populations of
Farmed Sea Bream (*Sparus aurata*) and Sea Bass (*Dicentrarchus Labrax*)', *Microorganisms*, 6/3 (2018), 92

Nikouli, Eleni, Meziti, Alexandra, Antonopoulou, Efthimia, Mente, Eleni, and Kormas, Konstantinos Ar, 'Host-Associated Bacterial Succession during the Early Embryonic
Stages and First Feeding in Farmed Gilthead Sea Bream (*Sparus aurata*)', *Genes*, 10/7 (2019), 483

Nikouli, Eleni, Meziti, Alexandra, Smeti, Evangelia, Antonopoulou, Efthimia, Mente, Eleni, and Kormas, Konstantinos Ar., 'Gut Microbiota of Five Sympatrically Farmed Marine
Fish Species in the Aegean Sea', *Microbial Ecology*, 81/2 (2021), 460–70

Oliva-Teles, A., Enes, P., and Peres, H., '8 - Replacing Fishmeal and Fish Oil in Industrial
624 Aquafeeds for Carnivorous Fish', in *Feed and Feeding Practices in Aquaculture*, ed. by D. Allen Davis, Woodhead Publishing Series in Food Science, Technology and
626 Nutrition (Oxford, 2015), 203–233

Péron, Guillaume, François Mittaine, Jean, and Le Gallic, Bertrand, 'Where Do Fishmeal and
628 Fish Oil Products Come from? An Analysis of the Conversion Ratios in the Global Fishmeal Industry', Coping with global change in marine social-ecological systems,
630 *Marine Policy*, 34/4 (2010), 815–20

Phelps, Drake, Brinkman, Nichole E., Keely, Scott P., Anneken, Emily M., Catron, Tara R.,
632 Betancourt, Doris, et al., 'Microbial Colonization Is Required for Normal Neurobehavioral Development in Zebrafish', *Scientific Reports*, 7/1 (2017), 11244

634 Quast, Christian, Pruesse, Elmar, Yilmaz, Pelin, Gerken, Jan, Schweer, Timmy, Yarza, Pablo, et al., 'The SILVA Ribosomal RNA Gene Database Project: Improved Data Processing
636 and Web-Based Tools', *Nucleic Acids Research*, 41/Database issue (2013), D590

Reis, Bruno, Gonçalves, Ana Teresa, Santos, Paulo, Sardinha, Manuel, Conceição, Luís E.
638 C., Serradeiro, Renata, et al., 'Immune Status and Hepatic Antioxidant Capacity of Gilthead Seabream *Sparus aurata* Juveniles Fed Yeast and Microalga Derived β-
640 Glucans', *Marine Drugs*, 19/12 (2021), 653

Ren, Lu-Jing, Ji, Xiao-Jun, Huang, He, Qu, Liang, Feng, Yun, Tong, Qian-Qian, et al.,
642 'Development of a Stepwise Aeration Control Strategy for Efficient Docosahexaenoic Acid Production by *Schizochytrium* sp', *Applied Microbiology and Biotechnology*, 87/5
644 (2010), 1649–56

Ringø, Einar, Li, Xuemei, Doan, Hien van, and Ghosh, Koushik, 'Interesting Probiotic
646 Bacteria Other Than the More Widely Used Lactic Acid Bacteria and Bacilli in Finfish', *Frontiers in Marine Science*, 9 (2022), 848037

648 Roeselers, Guus, Mittge, Erika K., Stephens, W. Zac, Parichy, David M., Cavanaugh, Colleen M., Guillemin, Karen, et al., 'Evidence for a Core Gut Microbiota in the Zebrafish',
650 *The ISME Journal*, 5/10 (2011), 1595–1608

Rognes, Torbjørn, Flouri, Tomáš, Nichols, Ben, Quince, Christopher, Mahé, Frédéric,
652 'VSEARCH: a versatile open source tool for metagenomics', *PeerJ* 4, e2584.

Rosenau, Simon, Oertel, Elisa, Mott, Alexander Charles, and Tetens, Jens, 'The Effect of a
654 Total Fishmeal Replacement by *Arthrospira platensis* on the Microbiome of African Catfish (*Clarias gariepinus*)', *Life (Basel, Switzerland)*, 11/6 (2021), 558

656 Salam, Mohammad Abdus, Islam, Md Ariful, Paul, Sulav Indra, Rahman, Md Mahbubur, Rahman, Mohammad Lutfar, Islam, Fatama, et al., 'Gut Probiotic Bacteria of
658 *Barbonymus gonionotus* Improve Growth, Hematological Parameters and Reproductive Performances of the Host', *Scientific Reports*, 11/1 (2021), 10692

660 Shah, Mahfuzur Rahman, Lutzu, Giovanni Antonio, Alam, Asraful, Sarker, Pallab, Kabir Chowdhury, M. A., Parsaeimehr, Ali, et al., 'Microalgae in Aquafeeds for a Sustainable
662 Aquaculture Industry', *Journal of Applied Phycology*, 30/1 (2018), 197–213

Shi, Yanyan, Cao, Xingyu, Ye, Zhidong, Xu, Yiyuan, Wang, Yiming, Li, Zhipeng, et al.,
'Role of Dietary *Schizochytrium* sp. in Improving Disease Resistance of Zebrafish through Metabolic and Microbial Analysis', *Aquaculture*, 539 (2021), 736631

Siddik, Muhammad A. B., Sørensen, Mette, Islam, S. M. Majharul, Saha, Newton, Rahman, Md Arifur, and Francis, David S., 'Expanded Utilisation of Microalgae in Global
Aquafeeds', *Reviews in Aquaculture*, 16/1 (2024), 6–33

Souza, Felipe Pinheiro de, Lima, Ed Christian Suzuki de, Urrea-Rojas, Angela Maria, Suphoronski, Suelen Aparecida, Facimoto, César Toshio, Bezerra Júnior, Jailton da Silva, et al., 'Effects of Dietary Supplementation with a Microalga (*Schizochytrium* sp.)
on the Hemato-Immunological, and Intestinal Histological Parameters and Gut Microbiota of Nile Tilapia in Net Cages', *PloS One*, 15/1 (2020), e0226977

Teixeira, Carla, Peixoto, Diogo, Hinzmann, Mariana, Santos, Paulo, Ferreira, Inês, Pereira, Gabriella V., et al., 'Dietary Strategies to Modulate the Health Condition and Immune
Responses in Gilthead Seabream (*Sparus aurata*) Juveniles Following Intestinal Inflammation', *Animals: An Open Access Journal from MDPI*, 12/21 (2022), 3019

Wan, Xu-zhi, Ai, Chao, Chen, Yi-han, Gao, Xiao-xiang, Zhong, Ru-ting, Liu, Bin, et al., 'Physicochemical Characterization of a Polysaccharide from Green Microalga
Chlorella pyrenoidosa and Its Hypolipidemic Activity via Gut Microbiota Regulation in Rats', *Journal of Agricultural and Food Chemistry*, 68/5 (2020), 1186–97

Wirth, Joseph S., and Whitman, William B., 'Phylogenomic Analyses of a Clade within the Roseobacter Group Suggest Taxonomic Reassignments of Species of the Genera
Aestuariivita, Citreicella, Loktanella, Nautella, Pelagibaca, Ruegeria, Thalassobius, Thiobacimonas and *Tropicibacter*, and the Proposal of Six Novel Genera',
International Journal of Systematic and Evolutionary Microbiology, 68/7 (2018), 2393–2411

Yilmaz, Pelin, Parfrey, Laura Wegener, Yarza, Pablo, Gerken, Jan, Pruesse, Elmar, Quast, Christian, et al., 'The SILVA and "All-Species Living Tree Project (LTP)" Taxonomic
Frameworks', *Nucleic Acids Research*, 42/Database issue (2014), D643–48

       Zanella, Lorenzo, and Vianello, Fabio, 'Microalgae of the Genus Nannochloropsis: Chemical
692   Composition and Functional Implications for Human Nutrition', *Journal of Functional Foods*, 68 (2020), 103919

694   Zhang, Dao-Feng, He, Wei, Shao, Zongze, Ahmed, Iftikhar, Zhang, Yuqin, Li, Wen-Jun, et al., 'Phylotaxonomic Assessment Based on Four Core Gene Sets and Proposal of a
696   Genus Definition among the Families Paracoccaceae and Roseobacteraceae', *International Journal of Systematic and Evolutionary Microbiology*, 73/11 (2023),
698   006156

       Zhang, Zhimin, Xi, Longwei, Liu, Haokun, Jin, Junyan, Yang, Yunxia, Zhu, Xiaoming, et al.,
700   'High Replacement of Fishmeal by Chlorella Meal Affects Intestinal Microbiota and the Potential Metabolic Function in Largemouth Bass (*Micropterus salmoides*)',
702   *Frontiers in Microbiology*, 13 (2022), 1016662

       Zorriehzahra, Mohammad Jalil, Delshad, Somayeh Torabi, Adel, Milad, Tiwari, Ruchi,
704   Karthik, K., Dhama, Kuldeep, et al., 'Probiotics as Beneficial Microbes in Aquaculture: An Update on Their Multiple Modes of Action: A Review', *Veterinary Quarterly*, 36/4
706   (2016), 228–41

Figure 1. Non-metric multidimensional scaling (nMDS) plot for all the bacterial communities of all
710 samples of the four dietary groups based on Bray-Curtis distances. PI samples are separated from FO
and SP samples. FO: Fish Oil, MI: *Microchloropsis + Isochrysis*, SP: *Schizochytrium +*
712 *Phaeodactylum*, PI: *Phaeodactylum + Isochrysis*.

716 **Figure 2.** Taxonomic composition of the bacterial families in each dietary group. Relative abundances were calculated adding the relative abundance (calculated based on the average samples reads) of all
718 OTUs belonging to each family. Only the families with relative abundance > 1% were considered for the diagram. Families with relative abundance < 4% are depicted in grey. FO: Fish Oil, MI:
720 *Microchloropsis + Isochrysis*, SP: *Schizochytrium + Phaeodactylum*, PI: *Phaeodactylum + Isochrysis*.

Figure 3. Most dominant OTUs in each dietary group (cumulative relative dominance > 80% and closest relative (Nucleotide BLAST) at the genus level. Relative abundances are % of reads. FO: Fish Oil, MI: *Microchloropsis + Isochrysis*, SP: *Schizochytrium + Phaeodactylum*, PI: *Phaeodactylum + Isochrysis*.

732 **Figure 4.** Relative abundance of the FO reads versus their respective values of the PI, MI and SP. OTUs in green dots are in ≥1% relative abundance in PI or MI or SP but not in FO, OTUs in red dots
734 are in ≥1% relative abundance in FO and PI, MI and SP concomitantly, OTUs in pale blue dots are in ≥1% relative abundance in FO but not in PI, MI and SP and OTUs in dark blue dots are in ≤1%
736 relative abundance in any treatment. The limit set (orange point) is the log of 1% of the total reads. FO: Fish Oil, MI: *Microchloropsis + Isochrysis*, SP: *Schizochytrium + Phaeodactylum*, PI:
738 *Phaeodactylum + Isochrysis*.

Figure 5. Scatter plots with every pathway in x axis and the log of the ratio PI/FO, MI/FO and SP/FO
of the Taxon function abundance in y axis. Pathways > 1.5 are considered overexpressed in PI, MI
and SP respectively compared to FO and pathways < -1.5 are considered underexpressed. FO: Fish
Oil, MI: *Microchloropsis* + *Isochrysis*, SP: *Schizochytrium* + *Phaeodactylum*, PI: *Phaeodactylum* +
Isochrysis.