The effect of dietary fish oil replacement by microalgae on the gilthead sea bream midgut bacterial microbiota

Stefanos Katsoulis-Dimitriou[1], Eleni Nikouli[1], Elli Zafeiria Gkalogianni[1], Ioannis T. Karapanagiotidis[1], Konstantinos A. Kormas[1,2*]

[1] Department of Agriculture Ichthyology & Aquatic Environment, University of Thessaly, Volos, Magnesia, Greece

[2] Agricultural Development Institute, University Research and Innovation Centre "IASON", Argonafton & Filellinon, 38221 Volos, Greece

* Corresponding author: kkormas@uth.gr

ABSTRACT

The aim of this work was to evaluate the effects of dietary microalgae blends as fish oil

20 replacers on the midgut bacterial microbiota of gilthead sea bream (*Sparus aurata*). The

control diet (FO) contained only fish oil as source of lipids, EPA and DHA fatty acids, while

22 three experimental diets were used where fish oil was replaced at 67% by one of the

following microalgae biomass blends: *Michrochloropsis gaditana* and *Isochrysis* sp.

24 (*Tisochrysis lutea*) (MI), *Phaeodactylum tricornutum* and *Isochrysis* sp. (PI) and

Schizochytrium sp. and *P. tricornutum* (SP). The midgut bacterial community composition

26 and the dominant OTUs indicated that the sea bream midgut bacterial communities were

altered compared to the control diet. Additional evidence from the presumptive bacterial

28 functional pathways suggests that the microalgae-based diets resulted in one overexpressed

and one underexpressed pathway. The overexpressed pathway was related to the metabolism

30 of fucose, a major carbohydrate of these microalgae species. This suggests that a new gut

microbiota profile was selected due to the microalgae inclusion in the provided diet. All these

32 data combined with the absence of mortality in fish, shows that the gilthead sea bream gut

microbiome can smoothly adapt its function according to the metabolic capacity of the

34 microalgae combinations that were used. The MI feed seems to promote several beneficial

bacteria with potential probiotic abilities in the fish gut.

Keywords: *Sparus aurata*, fish oil, aquaculture, microbiome, aquafeed

INTRODUCTION

The main sources of proteins and lipids for carnivorous farmed fish, such as the gilthead sea bream (*Sparus aurata)*, are fish meals and fish oils, respectively, which are added in the aquafeeds in order to satisfy fish nutritional requirements and to provide high nutritional quality to the final product (Siddik et al. 2023). Fish meals and fish oils originate mainly from wild small pelagic fish species, making the aquaculture sector dependent on wild fish stocks and reducing, thus, their sustainability (Peron, Mittaine & Le Gallic 2010). For this, several studies have focused on alternative sources of proteins and lipids to reduce the incorporation of fish meals and fish oils in aquafeeds (Oliva-Telles, Enes & Peres 2015, Naylor et al. 2021). Microalgae are suitable and sustainable alternative sources with studies so far showing promising results (Shah et al. 2017, Ahmad, Hassan & Banat 2022).

In fish, the composition of the gut microbiome is mainly determined by host selection (Roeselers et al. 2011) and diet (Liu et al. 2016), while environmental and other stochastic factors seem to be more important for wild fish (Kim et al. 2021, Kormas et al. 2023). The gut microbiome and its interactions with the host influence several important factors for fish health such as nutrition and metabolism (Liu et al. 2021), in maintaining the development and maturation of the immune system (Koch et al. 2018), the nervous system functioning and fish development (Phelps et al. 2017). Furthermore, commensal bacteria have the potential to protect the fish against pathogens (Kelly & Salinas 2017) and when a disease occurs there is a loss of gut microbiota diversity (Felix et al. 2022). For these reasons it is very important to investigate the changes that may occur in the composition of the gut microbiota with changes in the fish's diet.

In this research the following microalgae and oomycete species were used in gilthead sea bream aquafeeds: *Phaeodactylum tricornutum, Tisochrysis lutea* (here referred to as *Isochrysis* sp. due to its reclassification (Bendif et al. 2013))*, Schizochytrium* sp. and *Michrochloropsis gaditana* (formerly *Nannochloropsis gaditana*). *M. gaditana* (Fawley, Jameson & Fawley 2015) is a marine microalga of the class Eustigmatophyceae with high content of eicosapentaenoic acid (EPA) (usually 36% of total fatty acids) (Zanella & Vianello 2020). *P. tricornutum* is a marine diatom of the class Bacillariophyceae, rich in EPA (29% of total fatty acids) (Matos et al. 2016). The oomycete genus *Schizochytrium* belongs to the Thraustochytriaceae family. Members of this genus are especially rich in docosahexaenoic acid (DHA) (41% of total fatty acids) (Karapanagiotidis et al. 2022). *Isochrysis* is a genus of

marine unicellular microalgae that also produces high levels of DHA (49% of total fatty acids) (Ren et al. 2010).

Existing research has evaluated the use of *P. tricornutum* and *M. gaditana* in alternative feeds for gilthead seabream. In particular, *P. tricornutum* when used in experimental feeds, either alone or in combination with *Tetraselmis chuii* and *Bacillus subtilis*, has been linked with reduced gut microbial diversity in gilthead sea bream (Cerezuela et. al 2012). However, the above combination along with dietary inulin and the utilization of its b-glycans, has demonstrated immunostimulant effects, influencing intestinal gene expression (Cerezuela et al. 2013; Reis et al. 2021; Teixeira et al. 2022). The use of *M. gaditana* showed that gut microbiota composition and intestinal integrity remained unaltered (Cerezo-Ortega et al. 2021) or an increase in intestinal microbiota richness without affecting intestinal morphology and function (Jorge et al. 2019). When used in combination with other species (*Chlorella* sp., *Arthrospira* sp., *Tisochrysis lutea*, *Scenedesmus almeriensis*) it was shown to favour the presence of probiotic bacteria in the gut and reduce the presence of pathogenic species (Garcia-Márquez et al. 2023; Garcia-Márquez et al. 2023). To the best of our knowledge there are no studies about the effects of *Isochrysis* sp. and *Schizochytrium* sp. in gilthead sea bream gut microbiota.

These microalgae species when combined in fish diets, appear suitable for replacing the levels of EPA and DHA typically provided by fish oils. However, there are no studies evaluating the potential differences on the midgut microbial composition and function of gilthead seabream when fed with a diet containing these microalgae species in combination. Thus, the aim of this study was to investigate the potential effect on the midgut microbial composition and function of gilthead seabream when fed a diet containing these microalgae species in combination. It is hypothesised that a microalgal-based diets will promote gut microbiota related to the metabolism of these microalgae.

MATERIALS AND METHODS

Sampling. For the needs of the present study, 40 *Sparus aurata* specimens weighing 28-37g were collected from a feeding trial (Gkalogianni et al. 2023) conducted at the experimental facilities of the Department of Agriculture, Ichthyology and Aquatic Environment in 2022. The fish were divided into 4 nutritional groups with each group fed with a different diet for a period of 80 days. Fish of the control group were fed a diet

containing fish oil (FO) as the sole source of EPA and DHA, and fishmeal as a source of

108 marine animal protein. The remaining groups were fed experimental diets, in which 67% of

the fish oil used in the control diet was substituted by a mixture of microalgae species,

110 serving as a source of EPA and DHA. Specifically, the PI feed contained a microalgae

mixture of *P. tricornutum* and *Isochrysis sp.*, the SP a mixture of *Schizochytrium sp.* and *P.*

112 *tricornutum*, and the MI a mixture of *M. gaditana* and *Isochrysis sp.* Since the microalgae

used in the experimental diets were in the form of dry biomass (and not extracted oils), their

114 inclusion in the feed contributed to the overall protein content. For this reason, the percentage

of inclusion of fishmeal was reduced accordingly. At the end of the dietary trial, 10

116 individuals were randomly collected from each dietary group, euthanized with a strong dose

of the anesthetic phenoxyethanol and placed directly on ice. For the gut microbiota analysis,

118 the whole intestine of its individual was removed in aseptic conditions, with scissors and

scalpel after sterilization with 70% ethanol before each dissection and the midgut part was

120 stored at -80 °C until the DNA extraction procedure.

DNA extraction and sequencing. The DNA of seven gut samples of each feeding

122 group (FO, PI, SP, MI) was extracted using the Qiagen DNeasy PowerSoil Pro Kit (Qiagen,

Hilden, Germany), according to the manufacturer's instructions. Bacterial communities were

124 analysed by 16S rRNA amplicon sequencing targeting the V3-V4 hypervariable region with

the primer pair S-D-Bact-0341-b-S-17 and S-D-Bact-0785-a-A-21 (Klindworth et al. 2013).

126 PCR amplification and sequencing was performed at MRDNA Ltd.1 (Shallowater, TX,

United States) facilities on a MiSeq using paired end reads (2×300 bp) following the

128 manufacturer's guidelines.

Data analysis. The 16S rRNA sequencing raw data were processed using the

130 MOTHUR MiSeq SOP procedure (Schloss et al. 2013). Raw sequence reads have been

deposited in the Short Read Archive () with PRJNA1068122 BioProject accession number.

132 The operational taxonomic units (OTUs) at 97% cutoff similarity level were classified with

the SILVA database release 138.1 (Quast et al. 2013, Yilmaz et al. 2014). Identification of

134 the closest relatives of the most abundant OTUs was performed using a Nucleotide Blast

(http://blast.ncbi.nlm.nih.gov). The statistical analysis and graphic illustrations were

136 performed using PAleontological STudies (PAST) software (Hammer, Harper & Ryan 2001).

Presumptive functional analysis of midgut microbiota. The PICRUSt2

138 (Phylogenetic Investigation of Communities by Reconstruction of Unobserved STates)

(Douglas et al. 2020) software was used to predict the microbiota functional profile (Langille

140 et al. 2013). The predicted functions were searched in the MetaCyc database (Caspi et al.

2013), to be assigned to each metabolic pathway. Also, the KEGG (Kyoto Encyclopedia of
Genes and Genomes) database (Kanehisa & Goto 2000, Kanehisa et al. 2023) was used to
identify whether the genera of the most important OTUs have all the enzymes for specific
metabolic pathways of interest.

RESULTS

   Sequencing data and OTUs. After the quality filtering of the amplicon sequencing
of the 16S rRNA gene V3–V4 region, a total of 319.103 reads remained. The number of
reads ranged from 1.631 to 44.124 reads per sample. A subsample of 2.859 reads per sample
was used for the normalization of the data. As a result, two samples (one from each of
following treatments: FO and PI) with only 1.631 and 1.867 reads, respectively, were
excluded from the rest of the analyses. The total number of OTUs to which the reads were
assigned was 519.

   Alpha diversity indices. Taxa_S, Shannon_H, Simpson_1-D and Chao-1 indices
were calculated to assess the alpha diversity of the gut microbiota of gilthead seabream in the
four groups (Table 1). Compared to the control (FO), Shannon_H and Simpson_1-D values
showed a slight increase in PI, while they decreased in MI and SP. Chao-1 index values
showed a decrease compared to FO in all three groups, while Taxa_S increased in PI and SP
and showed a slight decrease in MI. Despite the existing differences, the Bray-Curtis
PERMANOVA (permutational analysis of variance) performed (Table S1) showed no
statistically significant differences between the indices values of the three experimental
groups with FO. The only statistically significant difference was observed in Taxa_S and
Shannon_H between PI and MI.

   Beta diversity. Non-metric multidimensional scaling analysis (nMDS) was used to
evaluate the beta diversity of the samples of the four feeding groups, based on a Bray-Curtis
distance matrix (Figure 1). The PI group is clearly separated from the samples of the FO and
SP group. The samples of the FO, MI and SP groups overlapped and there was also an
overlap between the samples of the PI and MI groups. Group PI was the only treatment with
no overlap with FO. PERMANOVA pairwise test (Table S2) was used to further analyze the
differences between dietary groups. Significant differences were observed between the PI and
FO and between the PI and SP groups.

174   The unique OTUs in FO, PI, SP and MI groups (Figure S1a) were 40 (7.7% of all

OTUs), 61 (11.8%), 148 (28.5%) and 56 (10.8%) respectively. 11.2% of the OTUs were

176   found in all groups (FO, PI, SP, MI). Furthermore, there were 91 unique OTUs in FO

compared to PI (30%), 95 unique OTUs compared to MI (34.2%) and 76 unique OTUs

178   compared to SP (19.5%) (Figure S1b). Shared OTUs were in a higher proportion between FO

and MI (34.5%) with FO and PI being very close (33%) and, lastly, FO and SP (29%). SP had

180   the highest number of unique OTUs (199 OTUs - 51%) compared to FO, with MI and PI

having smaller contributions of unique OTUs (87 OTUs - 31.3% and 112 OTUs -37%,

182   respectively).

Midgut bacterial taxonomic composition. Within the three most abundant phyla in

184   all groups were Proteobacteria and Actinobacteriota with different relative abundance rates

(Figure S2). In FO and MI the third most abundant phylum was Firmicutes, while in PI and

186   SP the 3rd most abundant phylum was Verrucomicrobiota and Bacteroidota respectively. In

FO, MI and SP, despite differences in either the order of abundance of the phyla (FO-MI) or

188   the phyla present (FO-SP), their ratio values were quite close. In contrast, in PI there was a

large increase in the relative abundance of Proteobacteria, since Proteobacteria were x2.1

190   more abundant than Actinobacteriota and x8.1 than Verrucomicrobiota. Actinobacteriota also

had a large difference in relative abundance with the third most abundant phylum,

192   Verrucomicrobiota (x3.8). Regarding other phyla, only in MI the Fusobacteriota occurred

with a relative abundance of 4.3% and in PI and SP the Firmicutes with relative abundance of

194   5.9% and 13.9% respectively.

In all feeding groups the Mycobacteriaceae family was most abundant in the FO, MI

196   and SP and second most abundant in PI (Figure 2). In PI the most abundant family was

Rhodobacteraceae, which in MI and SP occured in very low abundances. These abundances

198   were due to the most dominant OTUs, which in FO, MI and SP was OTU001 and OTU004 in

PI (Table S3). Nucleotide BLAST search assigned OTU001 as *Mycolicibacterium aurum* and

200   OTU004 as *Yoonia litorea*. Compared to FO, in the other feeding groups the family

Rhizobiaceae was not present and the family Xanthomonadaceae was only present in MI

202   (<4% relative abundance). The family Vibrionaceae occurred in MI and PI but it was absent

in SP and FO. Treatment MI had the highest number of families contributing >4% relative

204   abundance (nine families), while SP had the lowest (five families).

Most important OTUs. In the dominant OTUs (cumulative relative dominance >

206   80% - Figure 3) 3 OTUs occured in all feeding groups (OTU001, OTU003, OTU012).

OTU003 and OTU012 occured in small proportions in PI (1.67% and 2.44%). From the

Nucleotide BLAST search OTU003 closest relative was the *uncultured Staphylococcus sp.* and OTU012 closest relative was *Cutibacterium acnes*. OTU004, which was the most dominant in PI, was among the dominants of FO and MI, but with low relative abundances (2.95% and 2.16% respectively). Within the dominant OTUs there were several unique ones that occurred in each feeding group.

The qualitative differences already observed in phyla and families were also observed at the level of dominant OTUs in each group. To explore this further, we compared key OTUs between the FO and the three experimental groups. This comparison unveiled which OTUs were added in high abundance, were common in both the FO and experimental feeds, or exclusive to the FO (Figure 4, Table S4). 11 common OTUs were highly abundant in FO compared to MI, PI and SP. Furthermore, among OTUs found in high abundance in both FO and MI, PI, SP 2 OTUs (OTU001, OTU003) were observed in all cases. In OTUs added in high abundance in MI, PI and SP compared to FO, most were unique to each experimental feed, with 2 OTUs (OTU006, OTU027) being common in FO-MI and FO-PI.

Presumptive functional pathways. Compared to FO, one metabolic pathway was considerably overexpressed (FUCCAT-PWY) and one considerably underexpressed (PWY-6470) in all three experimental feeding groups (Figure 5, Table S5). According to the MetaCyc database, the FUCCAT-PWY pathway corresponds to L-fucose degradation, and the PWY-6470 pathway corresponds to peptidoglycan biosynthesis. Several common pathways were overexpressed in MI and PI relatively to FO and the highest number of unique pathways were overexpressed in SP. From the KEGG database it was found that the genomes of most of the OTUs in each of the three experimental feeds (see Fig. 4) contained the genes encoding for the enzymes of the metabolic pathway L-Fucose degradation (Table S6).

DISCUSSION

In this study, it was hypothesized that the inclusion of microalgal-based diets in experimental reared gilthead sea bream would induce structural bacterial microbiota changes adapted to these novel feed ingredients. In similar studies with the use of microalgae in the diet of gilthead sea bream there were varying results. Cerezuela et al. (2012) reported that the dietary inclusion of 100g/kg *Tetraselmis chui* or *P. tricornutum* decreased the intestinal microbial richness with a lower Shannon index value than the control. In the current study, the Shannon index was decreased in fish fed the *Schizochytrium - P. tricornutum* (SP) diet

but was slightly increased in fish fed the *P. tricornutum – Isochrysis* diet. Nevertheless, these two diets had more OTUs (Taxa_S) than the control (FO). Jorge et al. (2019) using *M. gaditana* for fishmeal replacement reported a higher, but insignificant to the control, intestinal richness and diversity of the species, while Garcia-Marquez et al. (2023) using a microalgae blend of *Tisochrysis lutea* (member of the *Isochrysis* sp**p. group**), *M. gaditana* and *Scenedesmus almeriensis* at 5%, 15% and 25% dietary inclusion levels showed that the microalgae blend promoted an increase in bacterial species diversity and a distinct shift in microbiota fingerprinting as inclusion levels increased. In our results the diet containing *M. gaditana* (MI) showed a decrease in the values of all alpha diversity indices, compared to the control (FO), but not statistically significant. Cerezo-Ortega et al. (2021) also reported no statistically significant differences in microbial diversity indices (alpha and beta diversity) in *Sparus aurata* gut after the inclusion of 5% hydrolysed *M. gaditana*.

Studies with dietary inclusion of *Schizochytrium* sp. in Nile tilapia, zebrafish and rainbow trout (Souza et al. 2020, Shi et al. 2021, Lyons et al. 2016) also showed varied results. Specifically, in Nile tilapia a diet with 1.2% inclusion of *Schizochytrium sp.* led to higher values of the Chao richness index compared to the control group. The authors, reported also higher Inverse Simpson and Shannon values in the *Schizochytrium*-fed fish, but these were not significantly different to those in the control group. In zebrafish, the dietary inclusion at 0, 60 and 120g/kg of *Schizochytrium sp.* resulted in insignificant differences in Chao and Shannon indices among the three groups, but in the beta diversity nMDS showed that both the *Schizochytrium sp.* supplemented diets were separated from the control group (Shi et al. 2021). The diet containing Schizochytrium sp. in our study (SP) had lower values in Shannon-H, Simpson_1-D and Chao-1 indices than the control (FO), but these were not statistically significant and also there were no statistically significant differences in beta diversity compared to the control. Furthermore, a 5% inclusion of *Schizochytrium limacinum* for fish oil replacement in the diet of rainbow trout resulted in a greater level of microbial diversity (Lyons et al. 2016; a fact that was also observed in our results using the SP diet that had a higher OTUs (Taxa_S) index number than the control (FO).

Overall, the effect on diversity is certainly determined by the different species of microalgae and fish and the levels of inclusion of each microalga in the diet. The literature as discussed above in combination with the results in our study shows the various changes that occur as the gut microbiome adapts to the different components added by microalgae to the diet. However, differences in diversity alone cannot show the overall effects and investigation

of changes at the level of bacterial phyla, families, and specific OTUs that modulate the diversity is required.

Most important OTUs. The dominant phyla found (Proteobacteria, Actinobacteriota, Bacteroidota) are common in the gilthead seabream gut microbiome (Kormas et al. 2014, Nikouli et al. 2018, 2019, Cerezo-Ortega et al 2021) and also in other fish species gut microbiome (Llewellyn et al. 2016, Rosenau et al. 2021, Zhang et al. 2022). In PI, however, there was an increased occurrence of Verrucomicrobiota. The Mycobacteriaceae family is the one that is highly abundant in all groups, due to OTU001, which belongs to the genus *Mycolicibacterium*. From the available literature it seems that this genus is not common in the gilthead sea bream gut. It includes some species of pathogenic micro-organisms for humans (*Mycobacterium tuberculosis, Mycobacterium leprae*) but also several species that inhabit a diverse range of environments including water bodies, soil, and metalworking fluids (Gupta, Lo & Son 2018). *Mycobacterium* genus has also been found in the gut of freshwater species (Gallet et al. 2022). The closest relative, *M. aurum* has never been reported as a pathogen, but its possible functions in the gut remain unknown. The Rhodobacteraceae family member OTU004 is the dominant in the PI treatment. According to the Nucleotide BLASt search its closest relative is *Yoonia litorea* (Paracocacceae family). *Y. litorea* though has been recently classified in the Roseobacter clade within the Rhodobacteraceae family (Wirth & Whitman 2018). Later, it was proposed the family Roseobacteraceae for members of the Roseobacter clade (Liang et al. 2021). Misclassifications have been a reoccurring problem between the Paracocacceae and Roseobacteraceae families (Zhang et al. 2023), but *Yoonia* sp. isolates are plylogenetically placed in the Roseobacteraceae family (Feng & Xing 2023). Roseobacter-clade species have been isolated from seawater marine sediments and biofilms, and are often associated with phytoplankton, macroalgae and marine animals (Wirth & Whitman 2018). They have also been found in intestine of finfish and they seem to have probiotic effects (Ringø et al. 2022). OTU003 is highly abundant in FO and MI, PI and SP and its closest relative is *Staphylococcus* sp., while OTU012 is highly abundant in FO, MI and PI and its closest relative is *Cutibacterium acnes*. *Staphylococcus* sp. is a common bacterial genus in healthy fish gut microbiomes and some species may have probiotic effects (Zorriehzahra et al.2015, Borah et al. 2016, El-Saadony et al. 2021). *Cutibacterium acnes* has been found in other fish species and it has been suggested to have potential probiotic features (Nikouli et al. 2021). OTU006 and OTU027, that belong to the *Vibrio* genus are in high abundance in MI and PI compared to FO. *Vibrio* is a common bacterial genus found in fish gut with both pathogenic and non-pathogenic species. Some *Vibrio* species can act as

symbionts assisting in the breakdown of dietary components, with strains found to produce

310   amylase, cellulose and chitinase among others (Egerton et al. 2018). From the rest top

abundant OTUs, in MI OTU013, OTU031 and OTU041 are closely related to

312   *Pseudoalteromonas sp., Pseudomonas sp.* and *Bacillus sp.* respectively. Several species of

these genera are known to have probiotic effects in the fish gut (Mladineo et al. 2016,

314   Zorriehzahra et al. 2015, Salam et al. 2021). OTU030 was abundant in FO and MI and was

related to *Rhodopseudomonas* sp., a genus that is also reported to have probiotic effects in the

316   fish gut (Koga et al. 2022). Thus, from the point of structural changes of the bacterial

communities, the MI experimental feed seems to promote several beneficial bacterial species

318   in the gut of gilthead seabream.

L-fucose degradation was the only considerably overexpressed bacterial metabolic

320   pathway in the three experimental feeds in comparison to FO. Fucose is a deoxyhexose that is

found as a constituent of exopolysaccharides from microalgae (Ahmed et al. 2013, Liu et al.

322   2019, Wan et al. 2019). It is also commonly found in animal cell glycans, macroalgae,

cyanobacteria and microbial exopolysaccharides (Becker & Lowe 2003). Furthermore, the

324   search of the KEGG database confirms that most of the important bacterial genera found in

this study, have all five enzymes to carry out the L-fucose degradation. This suggests that the

326   microalgal-based diets induced a clear gut bacterial response, at the functional level, favoring

the proliferation of bacteria which benefit from specific microalgal compounds with potential

328   benefits to the host, too. The degradation of fucose is a likely beneficial-to-the-host service.

For example, in humans fucose degraders are known to induce the production of the

330   beneficial short-chain fatty acids (Morrison et al. 2016, Høgsgaard et al. 2023, Fusco et al.

2023). The metabolic fate of the fucose degradation and its possible effect on the sea bream's

332   physiology remains to be specifically investigated in the future.

The only pathway that was considerably underexpressed in the three experimental

334   feeds compared to FO is peptidoglycan biosynthesis. The cell wall of some microalgae

species contains peptidoglycan (Agboola et al. 2019, Machado et al. 2022), but the species

336   used in our research do not seem to contain (Domozych et al. 2012, Le Costaouëc et al. 2017,

Nadzir et al. 2023). Peptidoglycan forms around 90% of the dry weight of Gram-positive

338   bacteria but only 10% of Gram-negative strains (Malanovic & Lohner 2016). The relative

abundance of Gram-negative phyla and families like Vibrionaceae in PI and MI and

340   Bacteroidota in SP is higher than FO. Finally, SP was the only treatment with unique

overexpressed pathways along with the highest number of unique OTUs compared to FO.

342   however, it seems that the unique OTUs are not responsible for these specific pathways (data

not shown). This microalgal feed inclusion requires further and more focused investigation

344   regarding its impact on the sea bream's gut microbiome.

     In conclusion, this study showed that the inclusion of three microalgal-based diets

346   induced both structural and functional changes in the sea bream's gut microbiome. The

structural changes in each dietary treatment were considerable compared to the control diet.

348   However, the total midgut bacterial communities which were favoured by the microalgal

inclusion are inferred to be able to metabolise L-fucose, a major microalgal carbohydrate.

352   **AKNOWLEDGMENTS**

354   This work was financially supported by the English MSc Program "Host-Microbe Interactions of the University of Thessaly and co-financed by the European Union and Greek

356   co-funded project entitled "Use of insect protein and microalgal oil for fishmeal and fish oil replacement in the diets of gilthead seabream (*Sparus aurata*) and European seabass

358   (*Dicentrarchus labrax*) undernational funds through the Operational Programme "Competitiveness, Entrepreneurship and Innovation-EPAnEK 2014-2020" (project code:

360   MIS 5045804)).

Table 1. Alpha diversity indices for each treatment. OTUs: operational taxonomic units.

Dietary treatment	OTUs (Taxa_S)	Shannon H	Simpson 1-D	Chao-1
Fish oil	53 ± 9.4	2.02 ± 0.570	0.75 ± 0.170	152.6 ± 85.38
Phaeodactylum + Isochrysis	57 ± 14.2	2.07 ± 0.366	0.77 ± 0.081	115.7 ± 26.74
Schizochytrium + Phaeodactylum	64 ± 63.3	1.80 ± 0.984	0.65 ± 0.265	123.2 ± 70.58
Microchloropsis + Isochrysis	44 ± 8.0	1.41 ± 0.657	0.56 ± 0.254	104.3 ± 37.40

REFERENCES

Agboola, Jeleel O., Teuling, Emma, Wierenga, Peter A., Gruppen, Harry, and Schrama, Johan W., 'Cell Wall Disruption: An Effective Strategy to Improve the Nutritive Quality of Microalgae in African Catfish (Clarias Gariepinus)', Aquaculture Nutrition, 25/4 (2019), 783–97

Ahmad, Ashfaq, W. Hassan, Shadi, and Banat, Fawzi, 'An Overview of Microalgae Biomass as a Sustainable Aquaculture Feed Ingredient: Food Security and Circular Economy', Bioengineered, 13/4 (2022), 9521–47

Ahmed, Mehboob, Moerdijk-Poortvliet, Tanja C.W., Wijnholds, Anita, Stal, Lucas J., and Hasnain, Shahida, 'Isolation, Characterization and Localization of Extracellular Polymeric Substances from the Cyanobacterium Arthrospira Platensis Strain MMG-9', European Journal of Phycology, 49/2 (2014), 143–50

Becker, Daniel J., and Lowe, John B., 'Fucose: Biosynthesis and Biological Function in Mammals', Glycobiology, 13/7 (2003), 41R-53R

Bendif, E.M., Probert, I., Schroeder, D.C. and de Vargas, C. (2013) On the description of *Tisochrysis lutea* gen. nov. sp. nov. and *Isochrysis nuda* sp. nov. in the Isochrysidales, and the transfer of Dicrateria to the Prymnesiales (Haptophyta). *Journal of Applied Phycology* **25**, 1763-1776.

Borah, Debajit, Gogoi, Olee, Adhikari, Chanakya, and Kakoti, B. B., 'Isolation and Characterization of the New Indigenous Staphylococcus Sp. DBOCP06 as a Probiotic Bacterium from Traditionally Fermented Fish and Meat Products of Assam State', Egyptian Journal of Basic and Applied Sciences, 3/3 (2016), 232–40

Caspi, Ron, Billington, Richard, Ferrer, Luciana, Foerster, Hartmut, Fulcher, Carol A., Keseler, Ingrid M., et al., 'The MetaCyc Database of Metabolic Pathways and Enzymes and the BioCyc Collection of Pathway/Genome Databases', Nucleic Acids Research, 44/D1 (2016), D471-480

Cerezo-Ortega, I. M., Di Zeo-Sánchez, D. E., García-Márquez, J., Ruiz-Jarabo, I., Sáez-Casado, M. I., Balebona, M. C., et al., 'Microbiota Composition and Intestinal Integrity Remain Unaltered after the Inclusion of Hydrolysed Nannochloropsis Gaditana in Sparus Aurata Diet', Scientific Reports, 11/1 (2021), 18779

Cerezuela, Rebeca, Fumanal, Milena, Tapia-Paniagua, Silvana Teresa, Meseguer, José, Moriñigo, Miguel Angel, and Esteban, Ma Angeles, 'Histological Alterations and Microbial Ecology of the Intestine in Gilthead Seabream (Sparus Aurata L.) Fed Dietary Probiotics and Microalgae', Cell and Tissue Research, 350/3 (2012), 477–89

Cerezuela, Rebeca, Meseguer, José, and Esteban, M. Ángeles, 'Effects of Dietary Inulin, Bacillus Subtilis and Microalgae on Intestinal Gene Expression in Gilthead Seabream (Sparus Aurata L.)', Fish & Shellfish Immunology, 34/3 (2013), 843–48

Domozych, David S., Ciancia, Marina, Fangel, Jonatan U., Mikkelsen, Maria Dalgaard, Ulvskov, Peter, and Willats, William G. T., 'The Cell Walls of Green Algae: A Journey through Evolution and Diversity', Frontiers in Plant Science, 3 (2012), 82

Douglas, Gavin M., Maffei, Vincent J., Zaneveld, Jesse R., Yurgel, Svetlana N., Brown, James R., Taylor, Christopher M., et al., 'PICRUSt2 for Prediction of Metagenome Functions', Nature Biotechnology, 38/6 (2020), 685–88

Egerton, Sian, Culloty, Sarah, Whooley, Jason, Stanton, Catherine, and Ross, R. Paul, 'The Gut Microbiota of Marine Fish', Frontiers in Microbiology, 9 (2018) , 873

El-Saadony, Mohamed T., Alagawany, Mahmoud, Patra, Amlan K., Kar, Indrajit, Tiwari, Ruchi, Dawood, Mahmoud A. O., et al., 'The Functionality of Probiotics in Aquaculture: An Overview', Fish & Shellfish Immunology, 117 (2021), 36–52

Fawley, Marvin W., Jameson, Ian, and Fawley, Karen P., 'The Phylogeny of the Genus Nannochloropsis (Monodopsidaceae, Eustigmatophyceae), with Descriptions of N. Australis Sp. Nov. and Microchloropsis Gen. Nov.', Phycologia, 54/5 (2015), 545–52

Feng, Xiaoyuan, and Xing, Peng, 'Genomics of Yoonia Sp. Isolates (Family Roseobacteraceae) from Lake Zhangnai on the Tibetan Plateau', Microorganisms, 11/11 (2023), 2817

Fusco, William, Lorenzo, Manuel Bernabeu, Cintoni, Marco, Porcari, Serena, Rinninella, Emanuele, Kaitsas, Francesco, et al., 'Short-Chain Fatty-Acid-Producing Bacteria: Key Components of the Human Gut Microbiota', Nutrients, 15/9 (2023), 2211

Gallet, Alison, Yao, Eric Kouamé, Foucault, Pierre, Bernard, Cécile, Quiblier, Catherine, Humbert, Jean-François, et al., 'Fish Gut-Associated Bacterial Communities in a Tropical Lagoon (Aghien Lagoon, Ivory Coast)', Frontiers in Microbiology, 13 (2022), 963456

García-Márquez, Jorge, Rico, Rosa María, Acién, Francisco Gabriel, Mancera, Juan Miguel, Figueroa, Félix L., Vizcaíno, Antonio Jesús, et al., 'Dietary Effects of a Short-Term Administration of Microalgae Blend on Growth Performance, Tissue Fatty Acids, and Predominant Intestinal Microbiota in Sparus aurata', Microorganisms, 11/2 (2023), 463

Gkalogianni, Elli Zafeiria, Psofakis, Pier, Asimaki, Adamantia, Fountoulaki, Eleni, Henry, Morgane, and Karapanagiotidis, Ioannis T. (2023). Microchloropsis gaditana, Schizochytrium sp., Phaeodactylum tricornutum, AND Tisochrysis lutea as n-3 PUFA sources in the diet of juvenile gilthead seabream (Sparus aurata). Proceedings of Aquaculture Europe 2023, September 18 - 21, Vienna, Austria.

Gupta, Radhey S., Lo, Brian, and Son, Jeen, 'Phylogenomics and Comparative Genomic Studies Robustly Support Division of the Genus Mycobacterium into an Emended Genus Mycobacterium and Four Novel Genera', Frontiers in Microbiology, 9 (2018), 00067

Hammer, Oyvind, Harper, David, and Ryan, Paul, 'PAST: Paleontological Statistics Software Package for Education and Data Analysis', *Palaeontologia Electronica*, 4 (2001), 1–9

Høgsgaard, Karina, Vidal, Natalia P., Marietou, Angeliki, Fiehn, Oliver Gam, Li, Qing, Bechtner, Julia, et al., 'Fucose Modifies Short Chain Fatty Acid and H2S Formation through Alterations of Microbial Cross-Feeding Activities', *FEMS Microbiology Ecology*, 99/10 (2023), fiad107

Jorge, Sara S., Enes, Paula, Serra, Cláudia R., Castro, Carolina, Iglesias, Paula, Oliva Teles, Aires, et al., 'Short-Term Supplementation of Gilthead Seabream (Sparus Aurata) Diets with Nannochloropsis Gaditana Modulates Intestinal Microbiota without Affecting Intestinal Morphology and Function', *Aquaculture Nutrition*, 25/6 (2019), 1388–98

Kanehisa, M., and Goto, S., 'KEGG: Kyoto Encyclopedia of Genes and Genomes', *Nucleic Acids Research*, 28/1 (2000), 27–30

Kanehisa, Minoru, Furumichi, Miho, Sato, Yoko, Kawashima, Masayuki, and Ishiguro-Watanabe, Mari, 'KEGG for Taxonomy-Based Analysis of Pathways and Genomes', *Nucleic Acids Research*, 51/D1 (2023), D587–92

Karapanagiotidis, I. T., Metsoviti, M. N., Gkalogianni, E. Z., Psofakis, P., Asimaki, A., Katsoulas, N., et al., 'The Effects of Replacing Fishmeal by Chlorella Vulgaris and Fish Oil by Schizochytrium Sp. and Microchloropsis Gaditana Blend on Growth Performance, Feed Efficiency, Muscle Fatty Acid Composition and Liver Histology of Gilthead Seabream (Sparus Aurata)', *Aquaculture*, 561 (2022), 738709

Kelly, Cecelia, and Salinas, Irene, 'Under Pressure: Interactions between Commensal Microbiota and the Teleost Immune System', *Frontiers in Immunology*, 8 (2017), 559

Kim, Pil Soo, Shin, Na-Ri, Lee, Jae-Bong, Kim, Min-Soo, Whon, Tae Woong, Hyun, Dong-Wook, et al., 'Host Habitat Is the Major Determinant of the Gut Microbiome of Fish', *Microbiome*, 9/1 (2021), 166

Klindworth, Anna, Pruesse, Elmar, Schweer, Timmy, Peplies, Jörg, Quast, Christian, Horn, Matthias, et al., 'Evaluation of General 16S Ribosomal RNA Gene PCR Primers for Classical and Next-Generation Sequencing-Based Diversity Studies', *Nucleic Acids Research*, 41/1 (2013), e1

Koch, Bjørn E. V., Yang, Shuxin, Lamers, Gerda, Stougaard, Jens, and Spaink, Herman P., 'Intestinal Microbiome Adjusts the Innate Immune Setpoint during Colonization through Negative Regulation of MyD88', *Nature Communications*, 9/1 (2018), 4099

Koga, Aoi, Yamasaki, Takumi, Hayashi, Shuhei, Yamamoto, Shinjiro, and Miyasaka, Hitoshi, 'Isolation of Purple Nonsulfur Bacteria from the Digestive Tract of Ayu (Plecoglossus Altivelis)', *Bioscience, Biotechnology, and Biochemistry*, 86/3 (2022), 407–12

Kormas, Konstantinos A., Meziti, Alexandra, Mente, Eleni, and Frentzos, Athanasios, 'Dietary Differences Are Reflected on the Gut Prokaryotic Community Structure of Wild and Commercially Reared Sea Bream (Sparus Aurata)', *MicrobiologyOpen*, 3/5 (2014), 718–28

Kormas, Konstantinos, Nikouli, Eleni, Kousteni, Vasiliki, and Damalas, Dimitrios, 'Midgut Bacterial Microbiota of 12 Fish Species from a Marine Protected Area in the Aegean Sea (Greece)', *Microbial Ecology*, 86/2 (2023), 1405–15

Kozich, James J., Westcott, Sarah L., Baxter, Nielson T., Highlander, Sarah K., and Schloss, Patrick D., 'Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform', *Applied and Environmental Microbiology*, 79/17 (2013), 5112–20

Langille, Morgan G. I., Zaneveld, Jesse, Caporaso, J. Gregory, McDonald, Daniel, Knights, Dan, Reyes, Joshua A., et al., 'Predictive Functional Profiling of Microbial Communities Using 16S rRNA Marker Gene Sequences', *Nature Biotechnology*, 31/9 (2013), 814–21

Le Costaouëc, Tinaïg, Unamunzaga, Carlos, Mantecon, Lalia, and Helbert, William, 'New Structural Insights into the Cell-Wall Polysaccharide of the Diatom Phaeodactylum Tricornutum', *Algal Research*, 26 (2017), 172–79

Liang, Kevin Y. H., Orata, Fabini D., Boucher, Yann F., and Case, Rebecca J., 'Roseobacters in a Sea of Poly- and Paraphyly: Whole Genome-Based Taxonomy of the Family Rhodobacteraceae and the Proposal for the Split of the "Roseobacter Clade" Into a Novel Family, Roseobacteraceae Fam. Nov.', *Frontiers in Microbiology*, 12 (2021), 683109

Liu, Han, Guo, Xianwu, Gooneratne, Ravi, Lai, Ruifang, Zeng, Cong, Zhan, Fanbin, et al., 'The Gut Microbiome and Degradation Enzyme Activity of Wild Freshwater Fishes Influenced by Their Trophic Levels', *Scientific Reports*, 6/1 (2016), 24340

Liu, Qishun, Yao, Changhong, Sun, Yongxin, Chen, Wei, Tan, Haidong, Cao, Xupeng, et al., 'Production and Structural Characterization of a New Type of Polysaccharide from Nitrogen-Limited Arthrospira Platensis Cultivated in Outdoor Industrial-Scale Open Raceway Ponds', *Biotechnology for Biofuels*, 12/1 (2019), 131

Liu, Yaqiu, Li, Xinhui, Li, Jie, and Chen, Weitao, 'The Gut Microbiome Composition and Degradation Enzymes Activity of Black Amur Bream (Megalobrama Terminalis) in Response to Breeding Migratory Behavior', *Ecology and Evolution*, 11/10 (2021), 5150–63

Llewellyn, Martin S., McGinnity, Philip, Dionne, Melanie, Letourneau, Justine, Thonier, Florian, Carvalho, Gary R., et al., 'The Biogeography of the Atlantic Salmon (Salmo Salar) Gut Microbiome', *The ISME Journal*, 10/5 (2016), 1280–84

Lyons, Philip P, Turnbull, James F, Dawson, Karl A, and Crumlish, Margaret, 'Effects of Low-Level Dietary Microalgae Supplementation on the Distal Intestinal Microbiome of Farmed Rainbow Trout Oncorhynchus Mykiss (Walbaum)', *Aquaculture Research*, 48/5 (2017), 2438–52

Machado, Luís, Carvalho, Gonçalo, and Pereira, Ricardo N., 'Effects of Innovative Processing Methods on Microalgae Cell Wall: Prospects towards Digestibility of Protein-Rich Biomass', *Biomass*, 2/2 (2022), 80–102

Malanovic, Nermina, and Lohner, Karl, 'Antimicrobial Peptides Targeting Gram-Positive Bacteria', *Pharmaceuticals*, 9/3 (2016), 59

Matos, Ângelo Paggi, Feller, Rafael, Moecke, Elisa Helena Siegel, Oliveira, José Vladimir de, Junior, Agenor Furigo, Derner, Roberto Bianchini, et al., 'Chemical Characterization of Six Microalgae with Potential Utility for Food Application' *Journal of the American Oil Chemists' Society*, 93/7 (2016) 963-72

Md Nadzir, Syafiqah, Yusof, Norjan, Nordin, Norazela, Kamari, Azlan, and Yusoff, Mohd Zulkhairi Mohd, 'A Review of Microalgal Cell Wall Composition and Degradation to Enhance the Recovery of Biomolecules for Biofuel Production', *Biofuels*, 14/9 (2023), 979–97

Medina-Félix, Diana, Garibay-Valdez, Estefanía, Vargas-Albores, Francisco, and Martínez-Porchas, Marcel, 'Fish Disease and Intestinal Microbiota: A Close and Indivisible Relationship', *Reviews in Aquaculture*, 15/2 (2023), 820–39

Mladineo, Ivona, Bušelić, Ivana, Hrabar, Jerko, Radonić, Ivana, Vrbatović, Anamarija, Jozić, Slaven, et al., 'Autochthonous Bacterial Isolates Successfully Stimulate In Vitro Peripheral Blood Leukocytes of the European Sea Bass (Dicentrarchus Labrax)', *Frontiers in Microbiology*, 7 (2016), 1244

Morrison, Douglas J., and Preston, Tom, 'Formation of Short Chain Fatty Acids by the Gut Microbiota and Their Impact on Human Metabolism', *Gut Microbes*, 7/3 (2016), 189–200

Naylor, Rosamond L., Hardy, Ronald W., Buschmann, Alejandro H., Bush, Simon R., Cao, Ling, Klinger, Dane H., et al., 'A 20-Year Retrospective Review of Global Aquaculture', *Nature*, 591/7851 (2021), 551–63

Nikouli, Eleni, Meziti, Alexandra, Antonopoulou, Efthimia, Mente, Eleni, and Kormas, Konstantinos A., 'Gut Bacterial Communities in Geographically Distant Populations of Farmed Sea Bream (*Sparus aurata*) and Sea Bass (*Dicentrarchus Labrax*)', *Microorganisms*, 6/3 (2018), 92

Nikouli, Eleni, Meziti, Alexandra, Antonopoulou, Efthimia, Mente, Eleni, and Kormas, Konstantinos Ar, 'Host-Associated Bacterial Succession during the Early Embryonic Stages and First Feeding in Farmed Gilthead Sea Bream (Sparus Aurata)', *Genes*, 10/7 (2019), 483

Nikouli, Eleni, Meziti, Alexandra, Smeti, Evangelia, Antonopoulou, Efthimia, Mente, Eleni, and Kormas, Konstantinos Ar., 'Gut Microbiota of Five Sympatrically Farmed Marine Fish Species in the Aegean Sea', *Microbial Ecology*, 81/2 (2021), 460–70

Oliva-Teles, A., Enes, P., and Peres, H., '8 - Replacing Fishmeal and Fish Oil in Industrial Aquafeeds for Carnivorous Fish', in *Feed and Feeding Practices in Aquaculture*, ed. by D. Allen Davis, Woodhead Publishing Series in Food Science, Technology and Nutrition (Oxford, 2015), 203–33

Péron, Guillaume, François Mittaine, Jean, and Le Gallic, Bertrand, 'Where Do Fishmeal and Fish Oil Products Come from? An Analysis of the Conversion Ratios in the Global

562   Fishmeal Industry', *Marine Policy*, Coping with global change in marine social-ecological systems, 34/4 (2010), 815–20

564   Phelps, Drake, Brinkman, Nichole E., Keely, Scott P., Anneken, Emily M., Catron, Tara R., Betancourt, Doris, et al., 'Microbial Colonization Is Required for Normal

566   Neurobehavioral Development in Zebrafish', *Scientific Reports*, 7/1 (2017), 11244

      Quast, Christian, Pruesse, Elmar, Yilmaz, Pelin, Gerken, Jan, Schweer, Timmy, Yarza, Pablo,

et al., 'The SILVA Ribosomal RNA Gene Database Project: Improved Data Processing and Web-Based Tools', *Nucleic Acids Research*, 41/Database issue (2013), D590

Reis, Bruno, Gonçalves, Ana Teresa, Santos, Paulo, Sardinha, Manuel, Conceição, Luís E. C., Serradeiro, Renata, et al., 'Immune Status and Hepatic Antioxidant Capacity of

Gilthead Seabream Sparus Aurata Juveniles Fed Yeast and Microalga Derived β-Glucans', *Marine Drugs*, 19/12 (2021), 653

Ren, Lu-Jing, Ji, Xiao-Jun, Huang, He, Qu, Liang, Feng, Yun, Tong, Qian-Qian, et al., 'Development of a Stepwise Aeration Control Strategy for Efficient Docosahexaenoic

Acid Production by Schizochytrium Sp', *Applied Microbiology and Biotechnology*, 87/5 (2010), 1649–56

Ringø, Einar, Li, Xuemei, Doan, Hien van, and Ghosh, Koushik, 'Interesting Probiotic Bacteria Other Than the More Widely Used Lactic Acid Bacteria and Bacilli in

Finfish', *Frontiers in Marine Science*, 9 (2022), 848037

      Roeselers, Guus, Mittge, Erika K., Stephens, W. Zac, Parichy, David M., Cavanaugh, Colleen

582   M., Guillemin, Karen, et al., 'Evidence for a Core Gut Microbiota in the Zebrafish', *The ISME Journal*, 5/10 (2011), 1595–1608

584   Rosenau, Simon, Oertel, Elisa, Mott, Alexander Charles, and Tetens, Jens, 'The Effect of a Total Fishmeal Replacement by Arthrospira Platensis on the Microbiome of African

586   Catfish (Clarias Gariepinus)', *Life (Basel, Switzerland)*, 11/6 (2021), 558

      Salam, Mohammad Abdus, Islam, Md Ariful, Paul, Sulav Indra, Rahman, Md Mahbubur,

588   Rahman, Mohammad Lutfar, Islam, Fatama, et al., 'Gut Probiotic Bacteria of Barbonymus Gonionotus Improve Growth, Hematological Parameters and

590   Reproductive Performances of the Host', *Scientific Reports*, 11/1 (2021), 10692

      Shah, Mahfuzur Rahman, Lutzu, Giovanni Antonio, Alam, Asraful, Sarker, Pallab, Kabir

592   Chowdhury, M. A., Parsaeimehr, Ali, et al., 'Microalgae in Aquafeeds for a Sustainable Aquaculture Industry', *Journal of Applied Phycology*, 30/1 (2018), 197–213

594   Shi, Yanyan, Cao, Xingyu, Ye, Zhidong, Xu, Yiyuan, Wang, Yiming, Li, Zhipeng, et al., 'Role of Dietary *Schizochytrium* Sp. in Improving Disease Resistance of Zebrafish

596   through Metabolic and Microbial Analysis', *Aquaculture*, 539 (2021), 736631

      Siddik, Muhammad A. B., Sørensen, Mette, Islam, S. M. Majharul, Saha, Newton, Rahman,

598   Md Arifur, and Francis, David S., 'Expanded Utilisation of Microalgae in Global Aquafeeds', *Reviews in Aquaculture*, 16/1 (2024), 6–33

600   Souza, Felipe Pinheiro de, Lima, Ed Christian Suzuki de, Urrea-Rojas, Angela Maria, Suphoronski, Suelen Aparecida, Facimoto, César Toshio, Bezerra Júnior, Jailton da

602   Silva, et al., 'Effects of Dietary Supplementation with a Microalga (Schizochytrium Sp.) on the Hemato-Immunological, and Intestinal Histological Parameters and Gut

604   Microbiota of Nile Tilapia in Net Cages', *PloS One*, 15/1 (2020), e0226977

      Teixeira, Carla, Peixoto, Diogo, Hinzmann, Mariana, Santos, Paulo, Ferreira, Inês, Pereira,

606   Gabriella V., et al., 'Dietary Strategies to Modulate the Health Condition and Immune Responses in Gilthead Seabream (Sparus Aurata) Juveniles Following Intestinal

608   Inflammation', *Animals: An Open Access Journal from MDPI*, 12/21 (2022), 3019

      Wan, Xu-zhi, Ai, Chao, Chen, Yi-han, Gao, Xiao-xiang, Zhong, Ru-ting, Liu, Bin, et al.,

610   'Physicochemical Characterization of a Polysaccharide from Green Microalga

Chlorella Pyrenoidosa and Its Hypolipidemic Activity via Gut Microbiota Regulation in Rats', *Journal of Agricultural and Food Chemistry*, 68/5 (2020), 1186–97

Wirth, Joseph S., and Whitman, William B., 'Phylogenomic Analyses of a Clade within the Roseobacter Group Suggest Taxonomic Reassignments of Species of the Genera Aestuariivita, Citreicella, Loktanella, Nautella, Pelagibaca, Ruegeria, Thalassobius, Thiobacimonas and Tropicibacter, and the Proposal of Six Novel Genera', *International Journal of Systematic and Evolutionary Microbiology*, 68/7 (2018), 2393–2411

Yilmaz, Pelin, Parfrey, Laura Wegener, Yarza, Pablo, Gerken, Jan, Pruesse, Elmar, Quast, Christian, et al., 'The SILVA and "All-Species Living Tree Project (LTP)" Taxonomic Frameworks', *Nucleic Acids Research*, 42/Database issue (2014), D643–48

Zanella, Lorenzo, and Vianello, Fabio, 'Microalgae of the Genus Nannochloropsis: Chemical Composition and Functional Implications for Human Nutrition', *Journal of Functional Foods*, 68 (2020), 103919

Zhang, Dao-Feng, He, Wei, Shao, Zongze, Ahmed, Iftikhar, Zhang, Yuqin, Li, Wen-Jun, et al., 'Phylotaxonomic Assessment Based on Four Core Gene Sets and Proposal of a Genus Definition among the Families Paracoccaceae and Roseobacteraceae', *International Journal of Systematic and Evolutionary Microbiology*, 73/11 (2023), 006156

Zhang, Zhimin, Xi, Longwei, Liu, Haokun, Jin, Junyan, Yang, Yunxia, Zhu, Xiaoming, et al., 'High Replacement of Fishmeal by Chlorella Meal Affects Intestinal Microbiota and the Potential Metabolic Function in Largemouth Bass (Micropterus Salmoides)', *Frontiers in Microbiology*, 13 (2022), 1016662

Zorriehzahra, Mohammad Jalil, Delshad, Somayeh Torabi, Adel, Milad, Tiwari, Ruchi, Karthik, K., Dhama, Kuldeep, et al., 'Probiotics as Beneficial Microbes in Aquaculture: An Update on Their Multiple Modes of Action: A Review', *Veterinary Quarterly*, 36/4 (2016), 228–41

Figure 1. Non-metric multidimensional scaling (nMDS) plot for all the bacterial communities of all samples of the four dietary groups based on Bray-Curtis distances. The MI samples have the largest range, PI does not coincide with FO and SP.

Figure 2. Taxonomic composition of the bacterial families in each dietary group. Relative abundances were calculated adding the relative abundance (calculated based on the average samples reads) of all OTUs belonging to each family. Only the families with relative abundance > 1% were considered for the diagram. Families with relative abundance < 4% are depicted in grey.

Figure 3. Most dominant OTUs in each dietary group (cumulative relative dominance > 80%) and closest relative (Nucleotide BLAST).

Figure 4. Relative abundance of the FO reads versus their respective values of the PI, MI and SP. OTUs depicted in green are in >1% relative abundance in PI or MI or SP but not in FO, OTUs depicted in red are in high abundance in FO and PI, MI and SP concomitantly and OTUs depicted in pale blue are in high abundance in FO but not in PI, MI and SP. The limit set (orange point) is the log of 1% of the total reads.

Figure 5. Scatter plots with every pathway in x axis and the log of the ratio PI/FO, MI/FO and SP/FO of the Taxon function abundance in y axis. Pathways > 1.5 are considered overexpressed in PI, MI and SP respectively compared to FO and pathways < -1.5 are considered underexpressed.