SMAC mimetics overcome apoptotic resistance in ovarian cancer through MSLN-TNF alpha axis

Authors:

Coelho Ricardo[1], Seashore-Ludlow Brinton[2], Schütz Sarah[1], Lombardo Christopher Flavio[1], Moussaud-Lamodière Elisabeth[2], Casanova Ruben[3,4], Ficek-Pascual Joanna[4], Labrosse Brunhilde Kathrin[1,5], Michal Hensler[6], Lopez-Nunez Monica[1], Rimmer Natalie[1], Fedier Andre[1], Lima Renata[1], ʾMontavon Sartorius Céline[5,7], Kurzeder Christian[5,8], Singer Franziska[9,10], Bertolini Anne[9,10], [Tumor Profiler Consortium](), Jitka Fucikova[6,11], Rätsch Gunnar[4], Bodenmiller Bernd[3], Kallioniemi Olli[2], Östling Päivi[2], David Leonor[12,13,14], Heinzelmann-Schwarz Viola[1,5,7*], Jacob Francis[1*]

Affiliations:
1. Ovarian Cancer Research, Department of Biomedicine, University Hospital Basel and University of Basel, Basel, Switzerland
2. Department of Oncology and Pathology, Science for Life Laboratory, Karolinska Institute, Solna, Sweden
3. Department of Quantitative Biomedicine, University of Zürich, Switzerland
4. Department of Computer Science, Eidgenössische Technische Hochschule (ETH) Zürich, Zürich, Switzerland
5. Gynaecological Cancer Centre, University Hospital Basel and University of Basel, Basel
6. Sotio Biotech, Prague, Czech Republic
7. Hospital for Women, University Hospital Basel, Basel, Switzerland
8. Breast Center, University Hospital Basel, Basel, Switzerland
9. SIB Swiss Institute of Bioinformatics, Basel, Switzerland
10. ETH Zurich, NEXUS Personalized Health Technologies, Schlieren, Switzerland
11. Department of Immunology, Charles University, 2nd Faculty of Medicine and University Hospital Motol, Prague, Czech Republic
12. Institute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP), Porto, Portugal.
13. Faculty of Medicine, University of Porto, Porto, Portugal
14. Institute for Research and Innovation in Health (i3S) of the University of Porto

*equal contribution

Keywords: chemoresistance, live cell imaging, myeloid cells, patient- derived cultures, programmed cell death, targeted therapy, TNF alpha signaling

Abstract

Resistance to chemotherapy and PARPi inhibitors remains a critical challenge in the treatment of epithelial ovarian cancer, mainly due to disabled apoptotic responses in tumor cells. Given mesothelin's pivotal role in ovarian cancer and its restricted expression in healthy tissues, we conducted a drug-screening discovery analysis across a range of genetically modified cancer cells to unveil mesothelin's therapeutic impact. We observed enhanced cell death in cancer cells with low mesothelin expression when exposed to a second mitochondria-derived activator of caspases (SMAC) mimetics, and demonstrated a compelling synergy when combined with chemotherapy in *ex vivo* patient-derived cultures and zebrafish tumor xenografts. Mechanistically, the addition of the SMAC mimetics drug birinapant to either carboplatin or paclitaxel triggered the activation of the Caspase 8-dependent apoptotic program facilitated by TNF signaling. Multimodal analysis of neoadjuvant-treated patient samples further revealed an association between tumor-associated macrophages and the activation of TNF-related pathways. Our proposed bimodal treatment shows promise in enhancing the clinical management of patients by harnessing the potential of SMAC mimetics alongside conventional chemotherapy.

Introduction

Primary debulking surgery followed by platinum-based chemotherapy has been the cornerstone treatment for advanced high-grade serous ovarian cancer (HGSOC) *(1, 2)*. In recent years, maintenance therapies including poly (ADP-ribose) polymerase inhibitors (PARPi), have become a valuable addition in patients with somatic or germline *BRCA1/2* tumors or homologous recombination deficiency (HRD), *(3–6)*. Despite these therapeutic advancements, most patients with metastatic HGSOC inevitably experience disease recurrence and ultimately platinum-resistance disease *(7)*. Considering these challenges, the necessity for overcoming drug resistance remains a critical clinical priority for these patients.

A growing body of evidence indicates that mesothelin (MSLN), a glycoprotein physiologically expressed at the surface of mesothelial cells lining the peritoneal cavity and overexpressed in several malignancies *(8–10)* plays a central role in tumor progression *(11–14)*. Additionally, we identified MSLN as a key player in the multistep process of metastasis in EOC, where it promotes cell survival in suspension, invasion through the mesothelial cell layer, and peritoneal spread *(15)*. Given its restricted expression and its role in ovarian cancer progression, MSLN has emerged as a promising candidate for targeted therapies using either antibody-based conjugates *(16, 17)* or CAR-T cell therapies *(18, 19)*. Currently, several clinical trials (NCT02792114, NCT03816358, NCT04809766, NCT03814447, NCT03054298) are underway to assess the potential of these targeted therapies in combination with chemotherapy, an area that holds promise for preventing or overcoming drug resistance.

Despite the increasing promise of MSLN-targeted therapies in HGSOC, a notable gap exists in our understanding, particularly regarding the broader role of MSLN in overall drug responses, including its specific impact on chemotherapy responses. To fill this knowledge gap, we functionally address the question by conducting a comprehensive drug library screen on genetically manipulated cell lines with varying levels of MSLN expression. Furthermore, we identify drug synergies, specifically combining carboplatin or paclitaxel with SMAC mimetics (bimodal treatment) through a TNFa-caspase-8 dependent mechanism.

Results

Ovarian cancer cells with low MSLN expression are selectively targeted by SMAC mimetics

MSLN expression has been correlated with chemoresistance *(20,21,22)*. However, these studies primarily report on associations without functionally elucidating the role of MSLN in drug responses. To fill this gap, we utilized our previously established experimental model *(15)*, consisting of isogenic EOC cell lines with genetically manipulated MSLN expression, to a comprehensive drug sensitivity and resistant test (DSRT) comprising a 528 oncology drug library including kinase inhibitors, epigenetic modifiers, differentiating agents, and conventional chemotherapeutic compounds ([CHEMICAL COMPOUND LIBRARIES](), **Supplementary Table 1**). We evaluated cellular response to treatment by calculating the drug sensitivity score (DSS, *(23)*), which revealed a diverse spectrum across all cell lines tested (n=8). Despite marginal differences in DSS to conventional chemotherapeutics (**Figure 1A-B**), we identified SMAC mimetics as exhibiting a selective effect on MSLN$^{low/\Delta MSLN}$ EOC cell lines (**Figure 1C**). This increased sensitivity to SMAC mimetics was further validated for the SMAC mimetic drug birinapant in two EOC cell lines genetically engineered for MSLN expression, either by MSLN knockout (OVCAR3$^{\Delta MSLN}$) or MSLNoverexpression (OVCAR4 OE MSLN) (**Figure 1D**). To further prove the MSLN-specific response to SMAC mimetics, we conducted a co-culture time-lapse experiment exposing parental (MSLN^{high}) and *ΔMSLN* OVCAR3 cells to birinapant treatment. For cell death assessment, we employed POPO™-1 iodide, a nontoxic cell impermeant dye. We observed a significant increase in cell death for *ΔMSLN* compared to parental OVCAR3, supporting the hypothesis that SMAC mimetics preferentially target MSLN$^{\Delta MSLN}$ cells (**Figure 1E**).

The tumor microenvironment (TME) plays a crucial role in modulating drug responses *(24)*. Mesothelial cells represent a major cellular component of the peritoneal microenvironment, the location where EOC cells are predisposed to metastasize *(25–28)*. Recognizing the importance of mesothelial cells in cancer progression and drug responses, we assessed the efficacy of birinapant in a co-culture assay using aggregates of parental and *ΔMSLN* OVCAR3 alongside normal immortalized mesothelial cells (MeT5A). This co-culture assay unveiled a selective effect of birinapant on *ΔMSLN* OVCAR3 cells, with minor effects on the MeT5A cell viability (**Figure 1F**). Taken together, our data indicates that the SMAC mimetics selectively targets MSLN$^{low/\Delta MSLN}$ EOC cells.

SMAC mimetics restore chemotherapy sensitivity

Despite observing marginal differences in the cellular response to carboplatin and paclitaxel chemotherapeutics when comparing the DSS values among cell lines with differential MSLN

expression, we hypothesized that chemotherapy might induce changes in MSLN levels. This

hypothesis is based on the well-documented effect of chemotherapeutic agents in inducing

cellular alterations, including DNA damage *(29)*, and cell cycle arrest *(30)*. These alterations

ultimately lead to extensive modifications in gene expression and protein function *(31, 32)*.

Thus, we screened a panel of 18 EOC cell lines for MSLN expression and selected

representative ones with MSLN[high] expression for downstream assays (**Supplementary Figure**

1A). Following exposure to a gradient concentration of carboplatin and paclitaxel, the most

effective chemotherapy compounds in EOC, we observed a decrease in MSLN protein levels

across all cell lines tested (n=4). This reduction was accompanied by reduced cell viability, as

evidenced by the increased expression of cleaved PARP (**Figure 2A, Supplementary Figure**

1B). The treatment of carboplatin combined with paclitaxel had a less pronounced impact on

MSLN protein levels as if they were given as single compounds (**Supplementary Figure 1C**).

Given this finding, we proceeded with downstream experiments using only the combination of

SMAC mimetics with either carboplatin or paclitaxel. Of note, EOC cell lines used exhibited a

general resistance to carboplatin and paclitaxel (**Supplementary Figure 1D**).

Supported by the observed reduction in MSLN protein levels following exposure to either

carboplatin or paclitaxel and the specificity of SMAC mimetics in targeting MSLN[low] cancer cells,

we conducted a dose-response matrix test and assessed the synergy scores for either

carboplatin or paclitaxel in combination with SMAC mimetics (birinapant and LCL161) in a panel

of EOC cell lines (n=6). We found enhanced MSLN-dependent drug efficiency when combining

one chemotherapeutic compound (either carboplatin or paclitaxel) with an SMAC mimetic

(bimodal treatment) (**Figure 2B**). To systematically analyze drug interaction patterns, we next

employed the Zero Interaction Potency (ZIP) model *(33, 34)*. This analysis revealed additive

and synergistic effects for the bimodal treatments in six EOC cell lines tested in an MSLN-

dependent manner. Surprisingly, the clinically routine combination of carboplatin with paclitaxel

exhibited only a low interaction pattern compared to bimodal treatments. (**Figure 2C and D and**

Supplementary Figure 2).

To further validate the synergistic effects, we profiled five HGSOC samples exhibiting high

MSLN expression for *ex vivo* drug testing (**Supplementary Figure 3A and Supplementary**

Table 2). These samples exhibited ubiquitous *TP53* mutations and heterogeneous mutation

patterns in HR and other frequently altered HGSOC genes (**Figure 2E, Supplementary Table**

3). The dose-response matrix demonstrated a remarkable increase in the efficiency of SMAC

mimetics in combination with either carboplatin or paclitaxel. We found additive and synergistic

interactions of bimodal treatments for three out of five HGSOC patient-derived samples tested,

importantly, independent of the tumor cell content (**Figure 2F and supplementary Figures 3B**

and C). To assess the on-target tumor effect of the bimodal treatment, we next utilized a

scaffold-free 3D *ex vivo* culture system followed by immunofluorescence staining for E-cadherin

and cleaved caspase 3 (cCASP3). A significant increase in the percentage of apoptotic tumor
cells was observed for the bimodal treatments as compared with controls suggesting a cancer
cell-specific killing effect (**Figures 2G and H**). Cellular responses to our bimodal treatment were
also assessed using the zebrafish embryo tumor xenograft model. In this model, we
transplanted COV362 MSLN[high] cells (carboplatin and paclitaxel-resistant) stably expressing
dTomato fluorescent protein (COV362_Chili) into transgenic *Tg(kdrl:eGFP)* zebrafish embryos
(35, 36). Zebrafish embryos were then exposed to birinapant in combination with paclitaxel or to
the vehicle for 72 hours. The bimodal treatment significantly (p=0.021) reduced the percentage
of viable COV362_Chili (DAPI⁻) cells as compared with the control condition (DMSO) (**Figure**
2I). Additionally, to exclude any potential toxicity arising from off-target effects of the bimodal
treatment, zebrafish embryos were exposed to a gradient concentration of paclitaxel, birinapant,
and their combinations for 72 hours. Neither the mortality rate, malformations, missing organs,
organ deficiencies, nor altered mobility capacity were observed in zebrafish larvae exposed to
the different drugs and combinations, indicating low off-target toxicity of the tested compounds
(**Supplementary Figure 4**). In summary, these results suggest that SMAC mimetics
administered with either carboplatin or paclitaxel enhance tumor cell killing *in vitro*, *ex vivo*, and
in vivo while causing minimal off-target toxicity.
169

Birinapant promotes cancer cell apoptosis through TNFα signaling
SMAC mimetics have gained attention as a promising class of targeted therapies and are
currently under clinical investigation for both solid and hematological malignancies *(37)*.
Mechanistically, SMAC mimetics are suggested to bind to Baculovirus Inhibitor apoptotic
proteins Repeat domains (BIR) of Inhibitor of Apoptosis (IAP)s, activating NF-⬜B signaling
pathway and, induction of cell death in a TNF⬜-dependent manner *(38–40)*. However, the
detailed mechanism of action of SMAC mimetics, particularly regarding their synergistic effects
when combined with either carboplatin or paclitaxel, remains unclear. To assess the impact of
TNF⬜ signaling on drug synergies, we exposed MSLN[high] EOC cell lines (Kuramochi and
OVCAR3) to the bimodal treatments in the presence of a neutralizing TNF⬜ antibody or
supplementing the culture medium with human recombinant TNF⬜ protein. The results
demonstrated that birinapant, used alone or in combination with either carboplatin or paclitaxel,
significantly enhanced cancer cell death in a TNF⬜-dependent manner (**Figure 3A and B and**
Supplementary Figure 5A-C). This was supported by the significant increase in the *TNFA*
expression in EOC cells after exposure to carboplatin, paclitaxel, birinapant, and bimodal
treatment (**Supplementary Figure 5D**). Of note, the major target of SMAC mimetics, cellular
(c)IAP1 protein *(39)*, exhibited a cell line-dependent protein expression after applied treatments
(**Figure 3B and Supplementary Figure 5C and E**). These data suggest that birinapant may

target different IAP proteins in a cell line-dependent manner, yet consistently produces an
enhanced effect when combined with either carboplatin or paclitaxel.

SMAC mimetics have been reported to trigger cell death through distinct programmed cell death
(PCD) pathways, involving either caspase-dependent apoptosis or caspase-independent
necroptosis *(41–45)*. The induction of these different pathways of PCD depends on the specific
combination of SMAC mimetics and the cancer model under investigation *(41, 43, 46, 47)*. To
evaluate which PCD is induced by our bimodal treatments in EOC, we established a reporter for
Caspase 3 and 8 activity, leveraging a previously introduced FLIP-GFP system *(48)*. Both
reporter systems were validated by either immunofluorescence staining or pharmacological
inhibition (Z-IETD-FMK) for Caspase 3 and 8, respectively (**Supplementary Figure 6A and B**).
Flow cytometry data from Kuramochi and OVCAR3 cells, which stably expressed caspase
reporters, demonstrated a significant increase in Caspase 8 and 3 activities (EGFP+) when
TNF☐ supplementation was provided, either in combination with birinapant or as part of our
bimodal treatments. Importantly, the increase in caspase activity was specifically induced by
supplementation with TNF☐ and not by the IFNγ (**Figure 3C and D, Supplementary Figure**
6C).

To further validate apoptosis as the primary PCD pathway responsible for the observed
synergies, and to investigate the role of the NF-☐B pathway in the outcome of bimodal
treatment, we conducted a CRISPR-Cas9 cell competition assay *(49)* in OVCAR3 cells. In this
assay, we targeted *NIK*, *IKKb*, *CASP8*, and *RIPK3* genes, encoding key players in the non-
canonical and canonical NF-☐B pathway, apoptosis, and necroptosis, respectively. The
generated data support the finding that apoptosis is indeed the primary PCD pathway triggered
by the treatment with TNF☐, birinapant, and birinapant in combination with paclitaxel. This was
evidenced by the increased cell resistance to these treatments upon genetic deletion of
ΔCASP8 (**Figure 3E**). Enhanced sensitivity to the bimodal treatment- (birinapant/ paclitaxel)
was also observed in *ΔKKb and ΔIKKb* indicating the additional involvement of the NF-☐B
signaling pathway in the treatment response. The role of Caspase 8 in mediating cell death
upon the bimodal treatment (birinapant with paclitaxel and birinapant with carboplatin) was
further validated through clonogenic cell survival in *ΔCASP8* cells (**Figure 3F**) and by co-
treatment with a Caspase 8- specific inhibitor (Z-IETD-FMK) (**Supplementary Figure 6D**).
Paclitaxel monotherapy appeared to induce necroptotic cell death, as indicated by the increased
resistance to paclitaxel observed in Δ*RIPK3* OVCAR3 cells (**Figure 3E**). Taken together, these
results indicate that treatment of either paclitaxel or carboplatin simultaneously with birinapant
primarily induces cell death through Caspase 8- mediated apoptosis. Furthermore, the results
also suggest that activation of the NF-☐B pathway, whether through the non-canonical or
canonical pathways, may modulate cellular responses to the applied treatment.

TNF☐ pathway activation triggered by tumor-associated macrophages enhances cancer cell apoptosis

Despite initially responding to chemotherapy, 20% of HGSOC patients relapse within 6 months after completing primary chemotherapy treatment, while 60% of the patients relapse after 6 months *(50)*. Patients who relapse after a treatment-free interval of more than six months often receive additional carboplatin or paclitaxel-based chemotherapy *(51)*. However, multiple recurrences increase the likelihood of developing platinum resistance, consequently resulting in poor survival *(52, 53)*.

To evaluate the potential of our bimodal treatment in enhancing chemotherapy effectiveness for patients with multiple relapses, we analyzed viable HGSOC cells derived from longitudinal samples collected at two disease recurrences (**Figure 4A**). Ascites taken at the time of the first and second relapse were subjected to *ex vivo* drug testing for carboplatin, birinapant, carboplatin with paclitaxel, and bimodal treatments. Overall, bimodal treatments revealed the highest efficacy in inducing tumor cell-specific apoptosis (E-Cadherin$^+$, cCASP3$^+$) compared to the remaining treatments tested (**Figure 4B and C**). Interestingly, bimodal treatments were generally more effective in reducing tumor cell viability at the second relapse time point compared to the first relapse.

In the complex ecosystem of tumor tissues, it is well-described that the tumor and the surrounding cells within the TME may be active participants in both the production and response to TNF☐ cytokine *(54, 55)*. Considering the pivotal role of TNF☐ in the apoptosis-mediated drug synergies for our bimodal treatments, we performed an integrated analysis on samples derived from the Swiss Tumor Profiler *(56)*. This analysis aimed to shed light on the potential contribution of different cell types within the TME to TNF☐ signaling activation and treatment response. This investigation included bulk RNA sequencing, cytometry by time-of-flight (CyTOF), and imaging mass cytometry (IMC) from the same sampling sites, obtained from patients with HGSOC who underwent neoadjuvant treatment. Based on our prior findings, we hypothesized that these patient samples were enriched in TNF☐ production post-neoadjuvant chemotherapy (**Supplementary Figure 5D**). Results from this analysis revealed a very strong positive correlation (spearman rho= 0.93, p=*0.007*, FDR=0.0317) between the TNF☐ gene sets and the fraction of tumor-associated macrophages (TAMs) (**Figure 4D-H, and Supplementary Figure 7**). To explore deeper into the IMC findings on the correlation between TAMs and TNF☐ gene sets, we conducted a deconvolution analysis of matching bulk RNA sequencing focusing on the immune compartment using CIBERSORT for relative quantification of the immune compartment using LM22 signature matrix *(57)*. Our analysis revealed an enrichment of myeloid cells (monocytes and M2 macrophages) in samples exhibiting increased activation of TNF☐ gene sets (**Figure 5A**). Additionally, deconvolution analysis by applying the xCell score *(58)* and the Molecular Signatures Database (MSigDB) for Hallmarks signatures of TNF☐ signaling

(https://www.pnas.org/doi/10.1073/pnas.0506580102) on bulk RNA sequencing data from an
independent cohort of 97 HGSOC cases confirmed the enrichment of M2 macrophages and
increased TNF signaling activation in post-neoadjuvant-treated omentum metastasis tissue
compared with chemotherapy-naive samples (**Figure 5B and Supplementary Figure 8**).
These findings prompted us to examine the impact of macrophage on responses to bimodal
treatments in *ex vivo* cultures of longitudinal samples. Analysis of the percentage of
macrophages (CD68$^+$) across these samples revealed a higher CD68$^+$ cell fraction in the
second relapse, aligning with an enhanced cellular response to bimodal treatments compared to
the first relapse (**Figure 4C and Figure 5C**). Additionally, supplementing *ex vivo* cultures with
TNF could further enhance the efficiency of the bimodal treatments (**Figure 5D and E**). In
summary, these findings suggest a potential role for TAMs as a predictive biomarker for the
response to bimodal treatments by regulating the TNF pathway activation.

Discussion

HGSOC is recognized as an extremely heterogeneous disease, presenting unique treatment challenges. Emerging therapeutic approaches aim to enhance patient outcomes by precisely identifying patient subgroups that will ultimately benefit from the applied treatment. The identification of such groups is based on multiple features of each tumor, including TME composition, genetic, and molecular profiles. Our approach seeks to improve patient stratification while maximizing the effectiveness of novel therapeutic approaches, thereby enhancing the quality of life and prolonging progression and disease-free survival. In this study, we introduce a novel approach to enhance the effectiveness of chemotherapy in HGSOC through co-treatment with SMAC mimetics. Our findings demonstrate that SMAC mimetics, particularly birinapant, in combination with either carboplatin or paclitaxel, trigger a Caspase 8-dependent apoptotic program facilitated by increased TNF production and signaling activation. Through our analysis of neoadjuvant-treated HGSOC samples, we identified a positive correlation between TAMs and TNF-related gene sets. Moreover, our experiments with HGSOC *ex vivo* cultures revealed a significant association between high TAM abundance and enhanced responses to the bimodal treatments. This suggests a potential role for TAMs in modulating drug responses through the mediation of TNF signaling, highlighting their potential utility as predictive biomarkers for the proposed bimodal treatments.

Our study has uncovered a promising avenue in targeted therapy in combination with chemotherapy for HGSOC. Specifically, we have identified SMAC mimetics as a class of drugs that exhibit a preferential affinity for EOC cells exhibiting low MSLN expression. Additionally, our investigation has unveiled a compelling synergy between SMAC mimetics and standard chemotherapeutic agents with the potential to enhance the efficacy of treatment regimens in HGSOC patients. Notably, our results highlight the remarkable increases in TNF expression following exposure to first-line chemotherapy. This, combined with validation experiments, indicates that TNF acts as a driving force behind the observed synergies. Furthermore, we hypothesize that the reduction in MSLN protein levels in EOC cells, after exposure to carboplatin or paclitaxel chemotherapy, enhances cell death triggered by increased TNF production in the presence of SMAC mimetics. This hypothesis is supported by a previous report showing increased resistance to TNF-induced cell death in MSLN-expressing cancer cells *(11)*.

While the exact mechanism through which MSLN contributes to heightened resistance to TNF-induced cell death remains incomplete, previous studies have suggested that MSLN induces the activation of the NF-B pathway, resulting in the overexpression of multiple anti-apoptotic factors *(11, 59–61)*, thereby contributing to increased resistance to TNF-induced cell death *(11)*. These findings align with data presented in the current study, where *MSLN* knockout

cancer cells displayed enhanced sensitivity to TNF□, birinapant, and the combination of birinapant with paclitaxel. It is worth noting that, in contrast to TNF□, the supplementation with IFNγ did not result in increased activation of Caspase 3 and 8 after exposure to birinapant or bimodal treatments. However, previous studies reported that IFNγ supplementation increases caspase activity following exposure to SMAC mimetics (BV6) in breast and colon cell line cancer models (46). These differences between our results and the literature may suggest a cancer model-dependent effect of the combination of cytokines with SMAC mimetics, or dependent on the SMAC mimetic applied, considering their documented varying effects on target proteins (39). Additionally, as recently reported by Cremona *et al.*, the genetic background of cancer models can influence the response to SMAC mimetics. This influence was demonstrated recently by the enhanced response to SMAC mimetics in *BRCA1* mutant cells as compared with *BRCA* wild-type cells (62). Therefore, further investigation is necessary to elucidate the role of different inflammatory cytokines on the activation of different programs of PCD in combination with SMAC mimetics across various cancer models.

SMAC mimetics have been reported to activate both apoptosis and necroptosis, two distinct pathways of PCD (43). The pharmacological induction of necroptosis is considered a promising approach to overcome apoptosis evasion of tumor cells (41, 63, 64). However, necroptosis has a dual role in cancer, it can either promote or inhibit tumor growth, depending on the specific cancer type and context. As a fail-safe mechanism of PCD, necroptosis can prevent tumor growth in cases where apoptosis induction fails. Nonetheless, as a form of necrotic cell death, necroptosis can trigger inflammatory responses and reportedly promote cancer metastasis and immunosuppression (65, 66). In our present study, we demonstrated that the combination of the birinapant with either carboplatin or paclitaxel chemotherapy primarily induces cell death through Caspase-8-mediated apoptosis. These findings are inconsistent with a previous study that reported SMAC mimetics to enhance the antitumor effect of carboplatin combined with paclitaxel in ovarian cancer models through a Caspase 8- independent mechanism (67). In this work, the effect on cell death was reported in cells with Caspase 8 knockdown when a triple combination of birinapant, paclitaxel, and carboplatin was applied. However, when tested individually, birinapant exhibited synergy with paclitaxel but an antagonism effect with carboplatin in caspase 8-depleted cells (67). While we cannot dismiss that the inconsistencies between our results and the previous report might arise from different measurements and experimental setups, a possible explanation could relate to the use of a triple combination (birinapant, carboplatin, and paclitaxel). Of note, one of the two cell line models used, OVCAR8, has been reported to exhibit resistance to NF-□B inhibition (68), indicating reduced responsiveness to TNF□ signaling. Interestingly, Hernandez *et al.* also noted an increase in cell death for the combination of birinapant with either paclitaxel or carboplatin in Caspase 8 expressing models. While further investigations are necessary, these results imply marginal

advantages in utilizing a triple combination of birinapant, carboplatin, and paclitaxel compared

to the bimodal treatments.

The TME has been reported to influence tumor evolution by altering the milieu of growth factors

and cytokines following chemotherapy *(69–71)*. Changes in cytokine levels produced by stromal

cells after chemotherapy tend to favor cancer cells expressing the corresponding receptors *(72)*.

In the current study, we found a compelling positive correlation between the presence of TAMs,

specifically M2 like macrophages, and the activation of TNF□ gene sets in neoadjuvant-treated

HGSOC samples. Among the various cell types in the TME, macrophages are recognized as

the major contributor to TNF□ production and signaling activation *(73, 74)*, which supports the

findings of in the current study. Additional studies are required to fully understand the interaction

and contribution of various cell types, especially TAMs, in the TME and their influence on TNF□

signaling in tumor cells. This is particularly relevant in the context of the bimodal treatment

where the synergistic effect of SMAC mimetics in combination with either carboplatin or

paclitaxel depends on the TNFa signaling. By mechanistically unraveling the interplay between

MSLN expression, TNF□ signaling, TAMs, and the enhanced efficacy of bimodal treatments, we

provide valuable insights that may guide the application of these therapeutic approaches in

patients with recurrent HGSOC. While further research is needed to explore how different cell

types in the TME contribute to TNF□ signaling in tumor cells and the effects of combination

therapies, our study established the groundwork for an exciting avenue in the treatment of this

challenging disease. Specifically, our results suggest that TAMs enriched in the TME could

potentially act as biomarkers for the proposed bimodal treatment, assisting in better patient

stratification. Overall, our findings hold promise in improving patient outcomes, especially for

those dealing with recurrent HGSOC.

Material and methods

Patient-derived cells

All HGSOC patient-derived samples and associated data included in this study fully adhere to ethical standards. Ethical approval was granted by the Ethical Committee of Northwest and Central, Switzerland (EKNZ, BASEC IDs: 2017-01900 and 2023-00988). All patients' clinical characteristics are provided in **Supplementary Table 2**.

Human cell lines

Ovarian cancer cell lines (Kuramochi, COV362, COV318, OVCAR3, OVCAR4, OVCAR5, OVCAR8, BG1, SKOV3, SKOV3 IP, ES-2, OVSAHO, A2780, A2780 cisplatin-resistant, TYKnu, TYKnu cisplatin-resistant, IGROV1, and IGROV1 paclitaxel-resistant) obtained from different commercial sources (**Supplementary Table 4**) were cultured in RPMI 1640 (Sigma-Aldrich) containing 10% fetal bovine serum (FBS) (Sigma-Aldrich) and 1% of penicillin/streptomycin (Sigma-Aldrich). Immortalized human mesothelial cell line MeT5A (ATCC) was maintained in Medium 199 (Sigma-Aldrich) containing 10% FBS, 3.3 nM epidermal growth factor (EGF) (PeproTech), 400 nM hydrocortisone (Sigma-Aldrich), 870 nM Bovine insulin (Sigma-Aldrich), 20 nM HEPES (Sigma-Aldrich) and 1:1000 of trace elements B (Corning). All cell lines were maintained at 37°C and 5% CO_2 authenticated using short tandem repeat (STR) profiling (Microsynth AG) and regularly tested for the absence of mycoplasma *(75)*. To study the role of MSLN in drug sensitivity and resistance we used a previously established and characterized cell line models: parental (MSLN^high) and MSLN knockout (*ΔMSLN*) OVCAR8 and OVCAR3 cell lines, as well as parental (MSLN^low) and MSLN overexpressing (MSLN OE- MSLN^high) BG1 and OVCAR4 cells *(15)*. These EOC cell lines were initially selected based on their nature of MSLN expressing levels.

Drug sensitivity and resistance testing (DSRT)

We used the DSRT system developed at the Institute for Molecular Medicine Finland (FIMM) for drug library screening. The FIMM oncology library- FO (https://www.helsinki.fi/en/infrastructures/drug-discovery-chemical-biology-and-screening/infrastructures/high-throughput-biomedicine/chemical-compound-libraries), consists of 528 approved and investigational oncology drugs including target compounds such as kinase inhibitors, epigenetic modifiers, differentiating agents and conventional chemotherapeutics. The compounds were dispensed in 384-well plates (Corning) at five different concentrations covering 1 and 10000nM range, using acoustic liquid dispensing technology (Echo 550; Labcyte Inc.). Assay-ready plates were stored in pressurized StoragePods (Roylan Developments) under an inert atmosphere until used. Before DSRT, growth patterns, and rates were evaluated by viability assays to ensure logarithmic cell growth during the 72 hours of drug exposure. Using

a MultiDrop™ Combi dispenser (Thermo Fisher Scientific) 5µL of culture media (RPMI 1640

supplemented with 10% FBS and 1% Pen/strep) was dispensed into assay-ready plates

followed by a brief centrifugation. Next, ovarian cancer cells (OVCAR3, OVCAR8, BG1, and

OVCAR4) and mesothelial cells, MeT5A, were seeded at the density of 1.5 to 2×10^3 cells/well in

20µl of media using the Multidrop Combi dispenser (Thermo Fisher Scientific) into drug-

containing plates followed by brief centrifugation and incubated for 72 hours at 37°C in the

presence of 5% CO_2. Cell viability/cellular ATP levels were measured with CellTiter-Glo 2.0

(CTG, Promega) with detection on the Ensight plate reader (PerkinElmer). Raw data were

further processed to calculate the drug efficacies of individual drugs. Using in-house software,

data from the plate reader were normalized per plate to the percent of cell viability in control

wells (DMSO, and benzethonium chloride- BzCl). Concentration-response curves were fitted to

percent viability values using a four-parameter logistic model and processed further to a

sensitivity metric (drug sensitivity score; DSS) using a weighted area-under-the-curve

calculation *(76)*. For the validation experiments, hit compounds and controls (DMSO, 100 µM

BzCl) were pre-dispensed on 384-well plates (Corning) using the Echo 550 (Labcyte Inc). Each

compound was plated as quadruplicate in eight concentrations (3-fold dilution).

Dose-response matrix analysis and synergy scoring

For the synergy experiments, we used 384-well screening plates with five concentrations per

compound in duplicate (carboplatin in combination with paclitaxel) or triplicate (birinapant,

LCL161 in combination either with carboplatin or paclitaxel, all dissolved in the same solution).

Cell seeding, viability, and calculation of DSS values (for cell lines and patient-derived samples,

Supplementary Table 2) were performed as described above for DSRT. Response matrices

were then calculated using the ZIP reference model with the SynergyFinder web application

(version 3.0; synergyfinder.fimm.fi) *(34)*. Combinations with a score > 10 were considered to

exhibit a strong synergy and <-10 a strong antagonism. Scores between 5 and 10 were

considered moderately synergistic, between -5 to -10 as moderately antagonistic, and between

-5 and 5 were indicative of additive effects.

Evaluation of drug efficiency using live cell confocal microscopy

Parental and △*MSLN* OVCAR3 cells were manually seeded into 384-well CellCarrier culture

plates (PerkinElmer) at the density of 2×10^3 cells/well in 20µl of media (RPMI 1640

supplemented with 10% FBS and 1% Pen/strep) and incubated for 24 hours. For the co-culture

assay, equal amounts of each cell population (1000 cells/well) were seeded in 20µl of media.

After the incubation time, 5µl of POPO™-1 Iodide (Thermo Scientific) diluted in culture media

(final concentration of 0.2µM) was added to all the wells of the plate. Birinapant (4000nM to

1.8nM concentration) was added in 5μl of media to the wells in quadruplicate as well as a

negative (DMSO 0.1%) and positive control (BzCl 100μM) followed by time-lapse microscopy

using the OperaPhenix microscope (PerkinElmer) for 48 hours. Image acquisition and

quantification of POPO™-1 iodide staining intensity were done using the Harmony high-content

imaging and analysis software 4.8 (PerkinElmer). Mesothelial clearance was performed by

seeding MeT5A-GFP cells *(15)* at the density of 1.3x 10^4 cells/well into 384-well CellCarrier

plates (PerkinElmer), pre-coated with 10μg/ml collagen type I (Sigma-Aldrich), in 20μl of media.

In parallel, ovarian cancer cell aggregates were generated by seeding the cells into t25 micro

cavity flasks (Corning) followed by a 24-hour incubation. After incubating, aggregates (10μl ~7

aggregates) were transferred to the plate containing the MeT5A-EGFP cells and exposed to a

gradient concentration of birinapant (4000nM to 1.8nM). POPO™-1 Iodide death marker

(Thermo Fisher Scientific) was added to all the wells of the plate at the final concentration of

0.2μM followed by 24-hour time-lapse microscopy using the OperaPhenix microscope

(PerkinElmer). Image acquisition and quantification of POPO™-1 Iodide (Thermo Fisher

Scientific) staining intensity was done using the Harmony high-content imaging and analysis

software 4.8 (PerkinElmer).

Immunoblotting

EOC cell lines were lysed in 1x radioimmunoprecipitation assay buffer (RIPA, Cell Signaling

Technology, BioConcept) containing a proteinase inhibitor cocktail (Sigma-Aldrich). Lysates

were clarified by centrifugation at 18,000 g for 15 minutes at 4°C. Clarified lysates were

quantified using the Pierce TM BCA protein assay (Promega) and boiled in 1x sample buffer (50

mM Tris-HCl, 1% SDS, 100 mM DTT, and 10% glycerol) at 95°C for 5 min and resolved by

SDS-PAGE. Proteins were then transferred to a polyvinylidene difluoride (PVDF) membrane

(BioRad) and blocked with 5% (w/v) bovine serum albumin in TBS-T (20 mM Tris-Base, 150 mM

NaCl, pH 7.8, 0.1% Tween 20) for 1 hour at room temperature. The membrane was incubated

with one of the listed primary antibodies (**Supplementary Table 5**) diluted in 5 % (w/v) BSA in

TBS-T overnight at 4°C. After washing (3 times, 10 minutes) in TBS-T, membranes were

incubated with corresponding HRP-conjugated secondary antibodies (1:10000, Cell Signaling,

BioConcept) for 3 hours at room temperature. Finally, the membrane was incubated with Super

Signal West Dura Extended Duration Substrate (Thermo Fisher Scientific) for the detection of

the HRP signal. Western blot results were visualized by Gel Doc XR+TM (BioRad) and

analyzed by Image Lab™ software (BioRad). Original membrane images can be found in the

Supplementary Figure 9-13.

Ex vivo **patient-derived cultures**

Tissue or ascites samples were thawed (retrospective cohort), and digested into single cells

(tissues) using a mix of collagenase/dispase (Roche), accutase (Millipore), DNAse type IV (New

England Biolabs/ Bioconcept) in DMEM/F12 (Sigma-Aldrich) for 1 hour at 37°C under agitation.

After digestion, samples were filtered through a 70µm cell strainer and washed with PBS

(Sigma-Aldrich). Cells were then seeded into agarose chips, which were pre-made using

MicroTissues 3D petri dish micro-mold (Sigma-Aldrich), at a density of 2.5×10^5 cells/ chip in

200µl of Ovarian TumorMACS™ (Miltenyi Biotec) medium and incubated for 24 hours. Patient-

derived cells were then exposed to the different drugs and their combinations for 48 hours

followed by fixation with 4% formalin (Formafix AG) for 20 minutes at room temperature. After

fixation cells were washed with PBS, and to prevent cell loss in the subsequent steps, a thin

layer of liquid histogel (Epredia) was added to the top of the cells. Agarose chips were then

subjected to a standard paraffin embedding protocol and sectioned at 5µm thickness followed

by immunofluorescence staining.

Immunofluorescence

After deparaffinization, heat-induced (98°C) antigen retrieval was performed with a citrate buffer

(pH 6.0) (Thermo Fisher Scientific), and slides were incubated with Triton X-100 0.25% in PBS

for 5 minutes, washed in PBS for another 5 minutes followed by incubation with blocking

solution (5% FBS with 1% BSA in PBS 1% Triton-X 100) for 1 hour. After blocking, samples

were incubated with one of the listed primary antibodies (**Supplementary Table 5**) diluted in

1% BSA 0.1% Triton-X 100 in PBS and incubated overnight at 4°C. Primary antibodies were

detected using goat anti-mouse or anti-rabbit IgG (H+L) Alexa fluor 488 or 647 (Cell signaling

Technologies/ Bioconcept) secondary antibodies diluted 1:500 in 1% BSA 0.1% Triton-X 100 in

PBS for 3 hours in the dark at room temperature. Slides were mounted using ProLong® Gold

antifade reagent (Cell Signaling Technology, BioConcept) and a coverslip. Images were taken

using the confocal microscope Nikon CSU-W1, analyzed with Image J (2.3.0/1.53q) and Qupath

(0.3.0) software, and developed scripts for cell detection and annotations. Further analyses

were performed by R/Bioconductor.

Zebrafish embryo tumor xenograft

All zebrafish experiments and husbandry followed federal guidelines complied with the Swiss

Animal Protection Ordinance and were approved by the Cantonal Veterinary Commission of

Basel-Stadt. The xenotransplantation experiments were performed as previously described with

minor alterations *(35, 77)*. Zebrafish embryos were anesthetized in 0.02% tricaine (Sigma-

Aldrich) at 2- day post fertilization (dpf) and 250–350 COV362 cells (stable expressing dTomato

fluorescent protein) were microinjected into the vessel-free area of the yolk of a transgenic

Tg(*kdrl:eGFP*) line. Embryos harboring cells were incubated at 35°C in 1x E3 medium

containing either DMSO 0.1% or birinapant in combination with paclitaxel at the final

concentration of 5000nM and 1000nM, respectively. Three days after transplantation, zebrafish

larvae (8-9 per condition) were dissociated into single cells by applying enzymatic digestion
using 5µl of liberase (Sigma-Aldrich) (2.5mg/ml) in a final volume of 50µl in PBS and incubated
for 15 minutes at 32°C with agitation 800 rpm. This incubation step was repeated three times
with repeated vigorous pipetting in between resulting in a total digestion time of 60 minutes.
After complete digestion, Liberase was inactivated by adding 5µl of 5% BSA in PBS. An
additional 150 µl of PBS was added, followed by filtration through a 70µm filter. The filter was
rinsed once with PBS to collect all cells. Single cells were washed twice with PBS and then
stained with a DAPI (Biolegend) solution (0.2 µg/ml) and analyzed on a CytoFLEX S (Beckman
Coulter) for DAPI-negative and dTomato-positive cells. The normalization for the input was
performed by dividing the number of dTomato$^+$ cells by the number of eGFP$^+$ (zebrafish) cells.
Toxicity tests of the various compounds and combinations were performed in zebrafish embryos
with a modified protocol based on the previously described methods *(78, 79)*. Zebrafish
embryos (at 2 dpf) were incubated for 3 days at 35°C in 1x E3 medium containing a gradient
concentration of DMSO, paclitaxel, birinapant, and birinapant in combination with paclitaxel.
After the incubation period, the mortality rate and general behavior were macroscopically
evaluated (n=2 wells per condition, each with 5 larvae). At least two larvae per condition were
selected for microscopic assessment of the morphology using confocal microscopy (Nikon CSU-
W1). This evaluation included a detailed analysis of malformation, absence of organs, or signs
of organ dysfunction.
536
Lentivirus production and transduction
Lentivirus production was performed using the HEK29T cells that were cultured in RPMI 1640
with 10% FBS and 1% penicillin/streptomycin. One day before transfection 4x10^6 cells were
seeded into a 75cm^2 tissue culture flask. For each flask, 4µg of the plasmid encoding the
targeted information, pUltra (Addgene #24129), pUltra-Chili (Addgene #48687), LentiV-Cas9-
puro (Addgene #108100), LGR2.1 (Addgene #108098) encoding the gRNA targeting the gene
of interest, pLenti-CMV-Blast-FLIP-GFP-DEVD, and pLenti-CMV-Blast-FLIP-GFP-XEXD, and
2µg of packaging (pCMVR8.74, Addgene #22036) and envelope (pMD2.G Addgene #12259)
plasmids were transfected using 24µl of jetPEI reagent and 1 ml of 150mM NaCl solution
(Polyplus-transfection). Culture media was changed 24 hours after transfection. The
supernatant containing active lentivirus particles was collected 48 hours later, filtered with a
0.45µm filter (Sartorius), and stored at -80°C. Target cell lines were then transduced with the
desired lentiviral supernatant. If needed after 3 passages, cells were enriched for dTomato,
EGFP, or mCherry positive cells using the BD FACS Aria Cell sorter (BD Bioscience).
551
Generation of caspase 3 and caspase 8 reporters

Plasmid encoding the sequence of the cleavage site for Caspase 3 (DEVD) and FLIP-GFP
reporter (Addgene#   124428) was digested using the *NheI*_HF and *XmaI* (New England
Biolabs, Bioconcept) and cloned into the pLenti_CMV_GFP_Blast (Addgene #17445) backbone,
that has been previously digested with *XbaI* (New England Biolabs, Bioconcept) and *XmaI*,
followed by transformation into *Stbl3 E. coli* strain. The right insertion and sequence of
FLIP_GFP_DEVD   in   the   pLenti_CMV_GFP_Blast   backbone
(pLenti_CMV_GFP_Blast_FLIP_GFP_DEVD) was confirmed by Sanger DNA sequence
(Microsynth) using the CMV-F primer (**Supplementary table 6**). Caspase 8 reporter was
established by replacing the DEVD amino acid sequence with XEXD in
pLenti_CMV_GFP_Blast_FLIP_GFP_DEVD plasmid through the In-fusion Snap assembly
master   mix   (Takara   Bio)   using   the   following   oligos   forward: 5'-
GTCTGGAAACCGATGGTGGCGGTGGCAAGGTG-3'   and   reverse   5-
CATCGGTTTCCAGACCTGATGCATCGGTAATGCCAG-3', PCR reaction was conducted under
the following conditions: denaturation at 98°C for 10 seconds, followed annealing at 55°C for 5
seconds and 72°C for 50 seconds, repeated for a total of 35 cycles. The resulting PCR product
was then purified using the Macherey-Nagel NucleoSpin gel and PCR clean-up kit (Thermo
Fisher Scientific). Linearized vector (67.4ng) was then mixed with 5X In-Fusion Snap Assembly
Master Mix and nuclease-free water in a final volume of 10ul. This mixture was incubated at
50°C for 15 minutes followed by transformation into Stellar™ competent cells a *HST08 E.coli*
strain. The protein-encoding XEXD DNA sequence was confirmed by the Sanger DNA
sequence using the CMV_F primer (**Supplementary Table 6**).
574
Caspase 3 and Caspase 8 activity analysis
Caspase 3 and Caspase 8 activity was visualized by flow cytometry in BG1, COV362, COV318,
OVCAR4, OVCAR3, and OVCAR8 cells stably expressing the FLIP-GFP reporter for caspase 3
and 8 activity. Cells ($5x10^4$/well) were seeded into 48-well plates and incubated for 24 hours,
followed by exposure to carboplatin, paclitaxel, birinapant, TNF☐, IFNy, and co-treatments for
24 hours. Cells were then harvested and stained with DAPI (0.1µg/ml) solution. Caspase 3 and
8 activity (EGFP$^+$ cells) was analyzed using the CytoFLEX Flow Cytometer (Beckman Coulter)
and Flowjo v10 BD (Becton Dickinson) software. Caspase 3 and 8 activity was represented as
the percentage of EGFP$^+$ cells within the mCherry positive cells (cells with stable expression of
the caspase 3 or caspase 8 reporter). For each cell line, two to four independent experiments
were performed.
586
Molecular cloning for the single RNAs
Single guide RNAs (sgRNA) targeting protein-coding DNA sequences of target genes was
designed using Benchling (Biology Software, 2021, retrieved from https://benchling.com).

SgRNAs with high-quality scores were selected for editing of the target gene (**Supplementary**
Table 7). Single-strand oligonucleotides were purchased from Sigma-Aldrich annealed and
cloned into LRG2.1 (addgene, #108098) using the *BsmBI* endonuclease restriction site and
T4_DNA ligase (Promega) for subsequent expression of the sgRNA together with EGFP
fluorescent protein. Ligations were transformed into *Stbl3 E.coli* following ampicillin selection,
ZR Plasmid Miniprep Classic plasmid purification (ZYMO Research, Lucerna-Chem), and
Sanger DNA sequencing (Microsynth) to confirm insertion of respective sgRNA using the
human U6 primer (**Supplementary Table 6**).
598

CRISPR-Cas9 cell competition assay
CRISPR-Cas9 cell competition assay was performed as previously described *(49)*. Stable Cas9-
expressing ovarian cancer cells were transduced with lentivirus particles containing sgRNAs
targeting selected genes (**Supplementary Table 7**). SgRNAs targeting the same gene were
pooled and further diluted in a 1:5 or 1:8 ratio with a complete culture growth medium. The
medium was changed 24 hours after transduction and cells were incubated for 2 additional days
until measuring the EGFP-positive cells. The percentage of fluorescence-positive cells was
determined initially (baseline reference, passage 0, 3 days after transduction) and all following
passages using the CytoFLEX Flow Cytometer (Beckman Coulter). In each passage, cells were
washed with FACS-wash (FW-1% FBS in PBS) in 96-well V-shape plates and stained with DAPI
(0.1 µg/ml) (Biolegend) for 5 minutes. All investigated cell lines were gated individually to
exclude debris, doublets, and dead (DAPI$^+$) cells.
Delivery of compounds and combinations was initiated at passage 1 and cells were kept under
treatment until passage 5 (20-23 days after transduction). The following concentrations were
used: 50 ng/ml, 1-2uM, 1-2nM, and 2-5nM of TNF□, carboplatin, paclitaxel, and birinapant.
Dropout values represent the fold-change of the percentage of EGFP$^+$ cells at each passage,
relative to passage 0, and were used as a readout of effects on cell fitness or drug sensitivity
conferred by the CRISPR-*Cas9* mediated gene mutations. Data analysis was performed using
FlowJo v10 BD (Becton Dickinson) and R/Bioconductor scripts. Relative drug sensitivity was
calculated by dividing the dropout values for the control condition (targeted gene without
treatment) by the values for the target gene under treatment.
Of note, due to the high number of guides applied in this study, each assay was performed at
least in two independent experiments with the negative control *AAVS1* (non-essential gene) and
at least one positive control (*RPA3* or *PCNA*). The robustness of the competition assay is
described in *(49)*.
624

Colony formation assay

Drug sensitivity was determined by the colony formation assays. OVCAR3-Cas9 cells were
transduced with lentivirus supernatant encoding sgRNAs targeting *AAVS1* or *CASP8*. Three
passages after lentivirus transduction cells were enriched for EGFP$^+$ cells using the BD
FACSAria Cell sorter (BD Bioscience). Cells were then seeded into 12 well plates at a density of
600 cells/well and exposed to the different drug combinations at the final concentration of 1µM,
1nM, and 5nM for carboplatin, paclitaxel, and birinapant, respectively. Complete growth media
with or without drugs was replaced every 3 days. After two weeks, colonies were fixed at room
temperature for 1 hour and stained with 0.05% crystal violet (Sigma-Aldrich) in 4% formalin
(Formafix AG) the plates were then rinsed 3-4 times with water, dried, and images of the well
were taken using the Fusion FX7 Edge Imaging System (Witek AG).
636
RT-qPCR
To extract RNA, 3x10^5 cells were seeded in 12-well plates and incubated for 48 hours. OVCAR3
and Kuramochi cells were then exposed to the following compounds and combinations for 24
hours: carboplatin (10µM), paclitaxel (10nM), birinapant (500nM), and birinapant (500nM)
combined with either carboplatin (10µM) or paclitaxel (10µM). Total RNA extraction was
performed using the RNeasy Plus universal mini kit (Qiagen). RNA was eluted in 30µl of
RNase-free water and the concentration was measured using a NanoDrop (Thermo Fisher
Scientific). A total of 1µg of RNA in 20µl reaction volume was reverse transcribed using the
iScript Reverse Transcription Supermix for RT-PCR (Bio-Rad). RT-qPCR was performed for
TNFA, and reference genes *SDHA* and *HSPCB (80)* in a 10µl reaction containing 25ng cDNA
(initial total RNA), 400nM forward and reverse primer (**Supplementary Table 6**), nuclease-free
water, and 1x GoTaq Master mix with ROX as a reference dye (Promega) on a Thermo ABI
7500 fast real-time PCR machine (Applied Biosystem, Thermo Fisher Scientific). Quantitative
PCR was performed in triplicates and analyzed using the 2–∆∆Ct method.
651
TNF☐ supplementation and neutralization
OVCAR3 and Kuramochi cells were seeded at a density of 5x10^5 cells/ well in a 6-well plate and
incubated for 48 hours. Cells were then exposed for 24 hours to the following compounds and
combinations: carboplatin, paclitaxel, birinapant, carboplatin and birinapant, paclitaxel and
birinapant either supplemented with TNF☐ (50ng/ml, Peprotech) or with TNF☐ neutralizing
antibody (1µg/ml, Cell Signalling Technologies), following protein extraction and immunoblot
analysis. Rabbit IgG (Santa Cruz Technologies) was used as a non-specific binding control.
659
Whole exome sequencing

Exonic regions were captured using the Twist Biosciences Human Comprehensive Exome
Panel for pair-end 100bp reads, as described by the manufacturer, and sequenced on an
Illumina NovaSeq 6000. Post-sequencing analysis of Whole Exome Sequencing (WES)
genomic data included the following steps: Initially, adapter trimming was performed using
SeqPurge (v2022_04), retaining reads with a minimum length of 15 bp. Alignment to the
reference genome GRCh38 was executed with BWA MEM (v0.7.17), followed by post-
processing via Samtools (v1.11) and Picard (v2.27.1) for removal of PCR duplicates and
secondary alignments. Base recalibration was carried out using the GATK suite (v4.2.6.1).
Quality control metrics were assessed using the following tools: FastQC (v0.11.9), Picard,
GATK, Samtools, and QualiMap (v2.2.2d), with MultiQC (v1.9) for aggregating the QC results.
Variant calling was performed with a combination VarScan2 (v2.4.4), Mutect2 (GATK v4.2.6.1),
and Strelka2 (2.9.10), with variants reported by at least two of three variant callers retained for
downstream analyses. Variants from all samples were merged using BCFTools (v1.15.1) and
annotated using the Ensembl Variant Effect Predictor (VEP) (v108) with databases including
ClinVar, dbSNP, gnomAD, dbNSFP, and COSMIC.
676
Suspension and imaging mass cytometry
Cytometry by the time of flight and imaging mass cytometry experiments (CyTOF and IMC
respectively) were performed as described before *(81)*. For CyTOF, single cells were identified
using a random forest classifier trained on a subset of manually gated cells using the Cytobank
platform *(82)*. For IMC, cell types were identified using a series of cell clustering and random
forest classification steps.
683
684 **Sequencing, alignment, and counting**
685 Bulk RNA libraries were ribo-depleted and sequenced with the Illumina NovaSeq 6000 system.
686 Alignment to the reference genome (GRCh38) was performed with STAR *(83)*. The raw
687 expression counts were obtained using simple counting, an in-house pipeline accessible
688 through github:gromics (at commit #67e1f4e).
689
Sample quality control
Quality control of samples was performed based on their RNA Integrity Score (RIN) and the
following FASTQC metrics: "Per Sequence GC Content", "Overexpressed Sequences", "Per
Base Sequence Quality", "Per Sequence Quality Scores" and "Per Base N Content". If multiple
FASTQ files were generated per sample, e.g., due to sequencing separately forward and
reverse reads or sequencing in multiple lanes or flowcells, the FASTQC metrics were
aggregated per sample, with at least one FAIL/WARN resulting in FAIL/WARN for this metric for
the considered sample. To determine the overall sample QC, taking into account all listed

FASTQC metrics as well as RIN score, the following criteria were adopted: RIN below six and

three or more failed FASTQC modules resulted in classifying a sample as FAIL, RIN score

above 7, and less than two failed FASTQC modules classified samples as PASS, anything in-

between or with missing information categorized sample as WARN. Applying these criteria to

the cohort resulted in 3 x FAIL, 11 x WARN, and 38 x PASS, with both WARN and PASS

samples being considered for preprocessing and all downstream analyses.

Data preprocessing

Only protein-coding genes, according to Gencode release v32 (GRCh38.p13) were kept, except

for mitochondrial genes, and further filtered to a set of genes with more or equal to 10 counts in

the cohort (sum across all samples). For pathway analysis, three subsequent preprocessing

steps were taken: a variance stabilizing transformation (VST), batch-effect correction, and

regressing out effects of confounding variables. VST, which internally adjusts for differences in

library sizes between samples, was performed with DESeq2 *(84)*, without taking any design

information into account. Moreover, batch correction with remove BatchEffect from the limma

package *(85)* was carried out using a sequencing date as a batch (with a total of 6 batches).

Furthermore, the first principal component capturing differences in RIN score (Pearson's r: -

0.68), as well as tumor content (estimated using CyTOF data) were regressed using the remove

Batch Effect function.

Pathway activation scoring

The pathway definitions, in terms of gene lists, were obtained from MSigDB (link) and KEGG

(link) databases, accessed on 29.07.2023. Pathway activation scores were computed using the

Gene Set Variation Analysis (GSVA, *(86)* with the minimum gene set size per pathway set to

five, Gaussian kernel, and using the normalized enrichment score (mx.diff=TRUE). The pathway

activation scores were then z-scored across the whole cohort, before subsetting the data to

neo-adjuvant cases only.

Bulk RNA sequencing

RNA-seq data was generated using the Illumina TruSeq Stranded mRNA Library Prep with 100

paired-end 100bp reads, according to the manufacturer's protocol, and sequenced on an

Illumina NovaSeq 6000. The analytical pipeline was built around the nf-core RNAseq pipeline

(v3.8.1). In brief, FASTQ files underwent trimming and quality control using Trim Galore

(v0.6.7), and Cutadapt (v.3.4) followed by mapping to the GRCh38 primary assembly using

STAR (v 2.7.10a), and the Gencode v41 primary annotation. Subsequent quantification was

performed with Salmon (v1.5.2). Quality control (QC) metrics were evaluated using

featureCounts (subread v2.0.1) for biotype counts, DESeq2 (v1.28.0) for Principal Component

Analysis (PCA) and sample similarity, Picard MarkDuplicates (v1.28.0) for identifying and
marking duplicate reads, Samtools (v1.15.1) for various alignment and post-alignment metrics,
FastQC (v0.11.9) for basic quality metrics, and RseQC (v3.0.1) for assessing RNA-seq data
quality. The QC results were collated into a singular report using MultiQC (v1.11). RNA-seq
data were normalized using the variance stabilizing transformation of the DESeq2 R (v1.36.0)
package to correct for technical variances. On this data, the deconvolution analysis was
performed by using a Docker container pulled locally to perform matrix deconvolution using the
LM22 signature matrix, allowing for the quantification of relative abundances in immune cells,
based on the bulk expression matrix of about 600 genes provided by the LM22 file.
744

Statistical analysis
Statistical analysis and figures were obtained using the software R Studio version 3.6.1 and 4.2
both versions were used interchangeably (www.R-project.org) and the {tidyverse} R software
suite. All negative and positive controls were performed on multiple replicates at least 3 times in
each cell line. All experiments were performed at least in duplicate and statistical evaluation
was performed using Prism 9 software (https://www.graphpad.com/scientific-software/prism/) or
R/Bioconductor. Where applicable, evaluation was done using two-way ANOVA with correction
for multiple comparison tests using BH (Benjamini-Hochberg correction) or the Wilcoxon test. p-
values of <0.05 were considered statistically significant and presented as a value or as
**$*p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.
755

Acknowledgments
We are grateful to the Flow Cytometry Core Facility (Stella Stefanova and former members),
Microscopy Facility (Mike Abanto, Loïc Sauteur, and Ewelina Bartoszek), Histology Facility
(Diego Calabrese), Martina Konantz, Markis Affolter group (Biozentrum, University Hospital
Basel, Basel, Switzerland) for their assistance and support during the zebrafish experiments,
and to the TuPro consortium and Miltenyi Biotec for providing essential support and reagents.
762

Funding Information
The work was funded by grants Strategic Focal Area "Personalized Health and Related
Technologies" grants of the ETH Domain (PHRT #2017-510 to VHS and FJ) and Krebsliga
Schweiz (KFS-5389-08–2021 to FJ), Swiss National Science Foundation (#320030_207975/1 to
VHS), and the Department of Biomedicine, University Hospital Basel and the University of
Basel, EACR travel fellowship (5787) and EMBO short-term fellowship (8099) both to RCo. FCT
project: Portuguese funds through FCT–Fundac̄ão para a Cie̅ncia e a Tecnologia/ Ministério
da Cie̅ncia, Tecnologia e Inovac̄ão in the framework of the project PTDC/MEC-ONC/29503/
2017 (to L.D.),

Author contributions

Conceptualization: RCo, FJ, LD, and BS-L

Methodology: RCo, BS-L, NR, FCL, and FJ

Investigation: RCo, SS, FCL, RL, MNL, NR, JF, RC, KBL, FS, AB, JF, MH and ELML

Data curation: RCo, FCL, and FJ

Writing—original draft preparation: RCo and FJ

Writing—review, and editing: all authors read and approved the final manuscript

Visualization: RCo, FJ, FCL, RC, and JF

Supervision: LD, PO, OK, BS-L, VHS, and FJ

Project administration: LD, VHS, and FJ.

Funding acquisition: RCo, LD, VHS.

Conflict of interest

The authors declare no competing interests.

ORCiDs

Ricardo Coelho: 0000-0002-9340-3612

Francis Jacob: 0000-0002-0446-1942

Viola Heinzelmann-Schwarz: 0000-0002-4056-3225

Sarah Schütz: 0009-0009-8679-3059

Kathrin Labrosse: 0000-0002-7195-5360

Franziska Singer: 0000-0002-6017-1595

Anne Bertolini: 0000-0003-0691-0489

Brinton Seashore-Ludlow: 0000-0001-8658-5967

Michal Hensler   0000-0002-8767-0722

Bibliography

1. I. Tsibulak, A. G. Zeimet, C. Marth, Hopes and failures in front-line ovarian cancer therapy., *Crit. Rev. Oncol. Hematol.* **143**, 14–19 (2019).

2. R. D. Mahmood, R. D. Morgan, R. J. Edmondson, A. R. Clamp, G. C. Jayson, First-Line Management of Advanced High-Grade Serous Ovarian Cancer., *Curr. Oncol. Rep.* **22**, 64 (2020).

3. N. Colombo, J. A. Ledermann, ESMO Guidelines Committee. Electronic address: clinicalguidelines@esmo.org, Updated treatment recommendations for newly diagnosed epithelial ovarian carcinoma from the ESMO Clinical Practice Guidelines., *Ann. Oncol.* **32**, 1300–1303 (2021).

4. M. R. Mirza, R. L. Coleman, A. González-Martín, K. N. Moore, N. Colombo, I. Ray-Coquard,
S. Pignata, The forefront of ovarian cancer therapy: update on PARP inhibitors., *Ann. Oncol.* **31**,
1148–1159 (2020).

5. A. Musella, C. Marchetti, M. L. Gasparri, L. Salerno, A. Casorelli, L. Domenici, L. Imperiale, I.
Ruscito, T. Abdul Halim, I. Palaia, V. Di Donato, F. Pecorini, M. Monti, L. Muzii, P. B. Panici,
PARP inhibition: A promising therapeutic target in ovarian cancer., *Cell Mol Biol (Noisy-le-*
grand) **61**, 44–61 (2015).

6. S. N. Banerjee, C. J. Lord, First-line PARP inhibition in ovarian cancer - standard of care for
all?, *Nat. Rev. Clin. Oncol.* **17**, 136–137 (2020).

7. R. Summey, D. Uyar, Ovarian cancer resistance to PARPi and platinum-containing
chemotherapy., *Cancer Drug Resist.* **5**, 637–646 (2022).

8. P. Argani, C. Iacobuzio-Donahue, B. Ryu, C. Rosty, M. Goggins, R. E. Wilentz, S. R.
Murugesan, S. D. Leach, E. Jaffee, C. J. Yeo, J. L. Cameron, S. E. Kern, R. H. Hruban,
Mesothelin is overexpressed in the vast majority of ductal adenocarcinomas of the pancreas:
identification of a new pancreatic cancer marker by serial analysis of gene expression (SAGE).,
Clin. Cancer Res. **7**, 3862–3868 (2001).

9. K. Chang, I. Pastan, Molecular cloning of mesothelin, a differentiation antigen present on
mesothelium, mesotheliomas, and ovarian cancers., *Proc Natl Acad Sci USA* **93**, 136–140
(1996).

10. H. F. Frierson, C. A. Moskaluk, S. M. Powell, H. Zhang, L. A. Cerilli, M. H. Stoler, H. Cathro,
G. M. Hampton, Large-scale molecular and tissue microarray analysis of mesothelin expression
in common human carcinomas., *Hum. Pathol.* **34**, 605–609 (2003).

11. U. Bharadwaj, C. Marin-Muller, M. Li, C. Chen, Q. Yao, Mesothelin confers pancreatic
cancer cell resistance to TNF-α-induced apoptosis through Akt/PI3K/NF-κB activation and IL-
6/Mcl-1 overexpression., *Mol. Cancer* **10**, 106 (2011).

12. U. Bharadwaj, C. Marin-Muller, M. Li, C. Chen, Q. Yao, Mesothelin overexpression
promotes autocrine IL-6/sIL-6R trans-signaling to stimulate pancreatic cancer cell proliferation.,
Carcinogenesis **32**, 1013–1024 (2011).

13. M.-C. Chang, C.-A. Chen, P.-J. Chen, Y.-C. Chiang, Y.-L. Chen, T.-L. Mao, H.-W. Lin, W.-H.
Lin Chiang, W.-F. Cheng, Mesothelin enhances invasion of ovarian cancer by inducing MMP-7
through MAPK/ERK and JNK pathways., *Biochem. J.* **442**, 293–302 (2012).

14. W. F. Cheng, C. Y. Huang, M. C. Chang, Y. H. Hu, Y. C. Chiang, Y. L. Chen, C. Y. Hsieh, C.
A. Chen, High mesothelin correlates with chemoresistance and poor survival in epithelial
ovarian carcinoma., *Br. J. Cancer* **100**, 1144–1153 (2009).

15. R. Coelho, S. Ricardo, A. L. Amaral, Y.-L. Huang, M. Nunes, J. P. Neves, N. Mendes, M. N.
López, C. Bartosch, V. Ferreira, R. Portugal, J. M. Lopes, R. Almeida, V. Heinzelmann-
Schwarz, F. Jacob, L. David, Regulation of invasion and peritoneal dissemination of ovarian
cancer by mesothelin manipulation., *Oncogenesis* **9**, 61 (2020).

16. H. L. Kindler, S. Novello, A. Bearz, G. L. Ceresoli, J. G. J. V. Aerts, J. Spicer, P. Taylor, K.
Nackaerts, A. Greystoke, R. Jennens, L. Calabrò, J. A. Burgers, A. Santoro, S. Cedrés, P.
Serwatowski, S. Ponce, J. P. Van Meerbeeck, A. K. Nowak, G. Blumenschein, J. M. Siegel, D.
A. Fennell, Anetumab ravtansine versus vinorelbine in patients with relapsed, mesothelin-
positive malignant pleural mesothelioma (ARCS-M): a randomised, open-label phase 2 trial.,

Lancet Oncol. **23**, 540–552 (2022).

17. X.-Y. Zhao, B. Subramanyam, N. Sarapa, S. Golfier, H. Dinter, Novel antibody therapeutics
targeting mesothelin in solid tumors., *Clin. Cancer Drugs* **3**, 76–86 (2016).

18. X. Liu, M. Onda, N. Watson, R. Hassan, M. Ho, T. K. Bera, J. Wei, A. Chakraborty, R.
Beers, Q. Zhou, A. Shajahan, P. Azadi, J. Zhan, D. Xia, I. Pastan, Highly active CAR T cells that
bind to a juxtamembrane region of mesothelin and are not blocked by shed mesothelin., *Proc*
Natl Acad Sci USA **119**, e2202439119 (2022).

19. T. Tokatlian, G. E. Asuelime, J.-Y. Mock, B. DiAndreth, S. Sharma, D. Toledo Warshaviak,
M. E. Daris, K. Bolanos, B. L. Luna, M. S. Naradikian, K. Deshmukh, A. E. Hamburger, A.
Kamb, Mesothelin-specific CAR-T cell therapy that incorporates an HLA-gated safety
mechanism selectively kills tumor cells., *J. Immunother. Cancer* **10** (2022), doi:10.1136/jitc-
2021-003826.

20. K. Nagata, E. Shinto, T. Shiraishi, M. Yamadera, Y. Kajiwara, S. Mochizuki, K. Okamoto, T.
Einama, Y. Kishi, H. Ueno, Mesothelin Expression is Correlated with Chemoresistance in Stage
IV Colorectal Cancer., *Ann. Surg. Oncol.* **28**, 8579–8586 (2021).

21. S. Takamizawa, S. Yazaki, Y. Kojima, H. Yoshida, R. Kitadai, T. Nishikawa, T. Shimoi, K.
Sudo, H. S. Okuma, M. Tanioka, E. Noguchi, M. Uno, M. Ishikawa, T. Kato, Y. Fujiwara, K.
Yonemori, High mesothelin expression is correlated with non-squamous cell histology and poor
survival in cervical cancer: a retrospective study., *BMC Cancer* **22**, 1215 (2022).

22. Y. Li, W. Tian, H. Zhang, Z. Zhang, Q. Zhao, L. Chang, N. Lei, W. Zhang, MSLN correlates
with immune infiltration and chemoresistance as a prognostic biomarker in ovarian cancer.,
Front. Oncol. **12**, 830570 (2022).

23. B. Yadav, T. Pemovska, A. Szwajda, E. Kulesskiy, M. Kontro, R. Karjalainen, M. M.
Majumder, D. Malani, A. Murumägi, J. Knowles, K. Porkka, C. Heckman, O. Kallioniemi, K.
Wennerberg, T. Aittokallio, Quantitative scoring of differential drug sensitivity for individually
optimized anticancer therapies., *Sci. Rep.* **4**, 5193 (2014).

24. M.-Z. Jin, W.-L. Jin, The updated landscape of tumor microenvironment and drug
repurposing., *Signal Transduct. Target. Ther.* **5**, 166 (2020).

25. J. Qian, B. L. LeSavage, K. M. Hubka, C. Ma, S. Natarajan, J. T. Eggold, Y. Xiao, K. C. Fuh,
V. Krishnan, A. Enejder, S. C. Heilshorn, O. Dorigo, E. B. Rankin, Cancer-associated
mesothelial cells promote ovarian cancer chemoresistance through paracrine osteopontin
signaling., *J. Clin. Invest.* **131** (2021), doi:10.1172/JCI146186.

26. H. A. Kenny, C.-Y. Chiang, E. A. White, E. M. Schryver, M. Habis, I. L. Romero, A. Ladanyi,
C. V. Penicka, J. George, K. Matlin, A. Montag, K. Wroblewski, S. D. Yamada, A. P. Mazar, D.
Bowtell, E. Lengyel, Mesothelial cells promote early ovarian cancer metastasis through
fibronectin secretion., *J. Clin. Invest.* **124**, 4614–4628 (2014).

27. M. Estermann, R. Coelho, F. Jacob, Y.-L. Huang, C.-Y. Liang, A. B. Faia-Torres, D.
Septiadi, B. Drasler, B. B. Karakocak, I. M. Dijkhoff, A. Petri-Fink, V. Heinzelmann-Schwarz, B.
Rothen-Rutishauser, A 3D multi-cellular tissue model of the human omentum to study the
formation of ovarian cancer metastasis., *Biomaterials* **294**, 121996 (2023).

28. M. Estermann, Y.-L. Huang, D. Septiadi, D. Ritz, C.-Y. Liang, F. Jacob, B. Drasler, A. Petri-
Fink, V. Heinzelmann-Schwarz, B. Rothen-Rutishauser, Patient-derived and artificial ascites
have minor effects on MeT-5A mesothelial cells and do not facilitate ovarian cancer cell

adhesion., *PLoS ONE* **15**, e0241500 (2020).

29. P. Bouwman, J. Jonkers, The effects of deregulated DNA damage signalling on cancer
chemotherapy response and resistance., *Nat. Rev. Cancer* **12**, 587–598 (2012).

30. H. Yang, R. M. Villani, H. Wang, M. J. Simpson, M. S. Roberts, M. Tang, X. Liang, The role
of cellular reactive oxygen species in cancer chemotherapy., *J. Exp. Clin. Cancer Res.* **37**, 266
(2018).

31. J. H. Fine, P. Chen, A. Mesci, D. S. J. Allan, S. Gasser, D. H. Raulet, J. R. Carlyle,
Chemotherapy-induced genotoxic stress promotes sensitivity to natural killer cell cytotoxicity by
enabling missing-self recognition., *Cancer Res.* **70**, 7102–7113 (2010).

32. A. Koussounadis, S. P. Langdon, D. J. Harrison, V. A. Smith, Chemotherapy-induced
dynamic gene expression changes in vivo are prognostic in ovarian cancer., *Br. J. Cancer* **110**,
2975–2984 (2014).

33. B. Yadav, K. Wennerberg, T. Aittokallio, J. Tang, Searching for Drug Synergy in Complex
Dose-Response Landscapes Using an Interaction Potency Model., *Comput. Struct. Biotechnol.*
J. **13**, 504–513 (2015).

34. L. He, E. Kulesskiy, J. Saarela, L. Turunen, K. Wennerberg, T. Aittokallio, J. Tang, Methods
for High-throughput Drug Combination Screening and Synergy Scoring., *Methods Mol. Biol.*
1711, 351–398 (2018).

35. F. Jacob, S. Alam, M. Konantz, C.-Y. Liang, R. S. Kohler, A. V. Everest-Dass, Y.-L. Huang,
N. Rimmer, A. Fedier, A. Schötzau, M. N. Lopez, N. H. Packer, C. Lengerke, V. Heinzelmann-
Schwarz, Transition of Mesenchymal and Epithelial Cancer Cells Depends on α1-4
Galactosyltransferase-Mediated Glycosphingolipids., *Cancer Res.* **78**, 2952–2965 (2018).

36. C. Cumin, Y.-L. Huang, C. Rossdam, F. Ruoff, S. P. Céspedes, C.-Y. Liang, F. C.
Lombardo, R. Coelho, N. Rimmer, M. Konantz, M. N. López, S. Alam, A. Schmidt, D. Calabrese,
A. Fedier, T. Vlajnic, M. von Itzstein, M. Templin, F. F. R. Buettner, A. Everest-Dass, F. Jacob,
Glycosphingolipids are mediators of cancer plasticity through independent signaling pathways.,
Cell Rep. **40**, 111181 (2022).

37. S. Fulda, Promises and challenges of smac mimetics as cancer therapeutics., *Clin. Cancer*
Res. **21**, 5030–5036 (2015).

38. E. Varfolomeev, J. W. Blankenship, S. M. Wayson, A. V. Fedorova, N. Kayagaki, P. Garg, K.
Zobel, J. N. Dynek, L. O. Elliott, H. J. A. Wallweber, J. A. Flygare, W. J. Fairbrother, K.
Deshayes, V. M. Dixit, D. Vucic, IAP antagonists induce autoubiquitination of c-IAPs, NF-
kappaB activation, and TNFalpha-dependent apoptosis., *Cell* **131**, 669–681 (2007).

39. L. Bai, D. C. Smith, S. Wang, Small-molecule SMAC mimetics as new cancer therapeutics.,
Pharmacol. Ther. **144**, 82–95 (2014).

40. H. Wu, J. Tschopp, S.-C. Lin, Smac mimetics and TNFalpha: a dangerous liaison?, *Cell*
131, 655–658 (2007).

41. C. Safferthal, K. Rohde, S. Fulda, Therapeutic targeting of necroptosis by Smac mimetic
bypasses apoptosis resistance in acute myeloid leukemia cells., *Oncogene* **36**, 1487–1502
(2017).

42. G. Brumatti, C. Ma, N. Lalaoui, N.-Y. Nguyen, M. Navarro, M. C. Tanzer, J. Richmond, M.
Ghisi, J. M. Salmon, N. Silke, G. Pomilio, S. P. Glaser, E. de Valle, R. Gugasyan, M. A.

Gurthridge, S. M. Condon, R. W. Johnstone, R. Lock, G. Salvesen, A. Wei, J. Silke, The caspase-8 inhibitor emricasan combines with the SMAC mimetic birinapant to induce necroptosis and treat acute myeloid leukemia., *Sci. Transl. Med.* **8**, 339ra69 (2016).

43. S. McComb, J. Aguadé-Gorgorió, L. Harder, B. Marovca, G. Cario, C. Eckert, M. Schrappe, M. Stanulla, A. von Stackelberg, J.-P. Bourquin, B. C. Bornhauser, Activation of concurrent apoptosis and necroptosis by SMAC mimetics for the treatment of refractory and relapsed ALL., *Sci. Transl. Med.* **8**, 339ra70 (2016).

44. S. L. Petersen, M. Peyton, J. D. Minna, X. Wang, Overcoming cancer cell resistance to Smac mimetic induced apoptosis by modulating cIAP-2 expression., *Proc Natl Acad Sci USA* **107**, 11936–11941 (2010).

45. Y. Gong, Z. Fan, G. Luo, C. Yang, Q. Huang, K. Fan, H. Cheng, K. Jin, Q. Ni, X. Yu, C. Liu, The role of necroptosis in cancer biology and therapy., *Mol. Cancer* **18**, 100 (2019).

46. M. J. Cekay, S. Roesler, T. Frank, A.-K. Knuth, I. Eckhardt, S. Fulda, Smac mimetics and type II interferon synergistically induce necroptosis in various cancer cell lines., *Cancer Lett.* **410**, 228–237 (2017).

47. A. K. Srivastava, S. Jaganathan, L. Stephen, M. G. Hollingshead, A. Layhee, E. Damour, J. P. Govindharajulu, J. Donohue, D. Esposito, J. P. Mapes, R. J. Kinders, N. Takebe, J. E. Tomaszewski, S. Kummar, J. H. Doroshow, R. E. Parchment, Effect of a Smac Mimetic (TL32711, Birinapant) on the Apoptotic Program and Apoptosis Biomarkers Examined with Validated Multiplex Immunoassays Fit for Clinical Use., *Clin. Cancer Res.* **22**, 1000–1010 (2016).

48. Q. Zhang, A. Schepis, H. Huang, J. Yang, W. Ma, J. Torra, S.-Q. Zhang, L. Yang, H. Wu, S. Nonell, Z. Dong, T. B. Kornberg, S. R. Coughlin, X. Shu, Designing a green fluorogenic protease reporter by flipping a beta strand of GFP for imaging apoptosis in animals., *J. Am. Chem. Soc.* **141**, 4526–4530 (2019).

49. R. Coelho, A. Tozzi, M. Disler, F. Lombardo, A. Fedier, M. N. López, F. Freuler, F. Jacob, V. Heinzelmann-Schwarz, Overlapping gene dependencies for PARP inhibitors and carboplatin response identified by functional CRISPR-Cas9 screening in ovarian cancer., *Cell Death Dis.* **13**, 909 (2022).

50. M. Luyckx, J.-L. Squifflet, A. M. Bruger, J.-F. Baurain, in *Ovarian Cancer*, S. Lele, Ed. (Exon Publications, Brisbane (AU), 2022).

51. T. Baert, A. Ferrero, J. Sehouli, D. M. O'Donnell, A. González-Martín, F. Joly, J. van der Velden, P. Blecharz, D. S. P. Tan, D. Querleu, N. Colombo, A. du Bois, J. A. Ledermann, The systemic treatment of recurrent ovarian cancer revisited., *Ann. Oncol.* **32**, 710–725 (2021).

52. P. G. Rose, Ovarian cancer recurrence: is the definition of platinum sensitivity modified by PARPi, bevacizumab or other intervening treatments?□: a clinical perspective., *Cancer Drug Resist.* **5**, 415–423 (2022).

53. S. Pignata, S. C Cecere, A. Du Bois, P. Harter, F. Heitz, Treatment of recurrent ovarian cancer., *Ann. Oncol.* **28**, viii51–viii56 (2017).

54. E. Dondossola, A. S. Dobroff, S. Marchiò, M. Cardó-Vila, H. Hosoya, S. K. Libutti, A. Corti, R. L. Sidman, W. Arap, R. Pasqualini, Self-targeting of TNF-releasing cancer cells in preclinical models of primary and metastatic tumors., *Proc Natl Acad Sci USA* **113**, 2223–2228 (2016).

55. H. Zhao, L. Wu, G. Yan, Y. Chen, M. Zhou, Y. Wu, Y. Li, Inflammation and tumor

progression: signaling pathways and targeted intervention., *Signal Transduct. Target. Ther.* **6**, 263 (2021).

56. A. Irmisch, X. Bonilla, S. Chevrier, K.-V. Lehmann, F. Singer, N. C. Toussaint, C. Esposito, J. Mena, E. S. Milani, R. Casanova, D. J. Stekhoven, R. Wegmann, F. Jacob, B. Sobottka, S. Goetze, J. Kuipers, J. Sarabia Del Castillo, M. Prummer, M. A. Tuncel, U. Menzel, M. P. Levesque, The Tumor Profiler Study: integrated, multi-omic, functional tumor profiling for clinical decision support., *Cancer Cell* **39**, 288–293 (2021).

57. A. M. Newman, C. L. Liu, M. R. Green, A. J. Gentles, W. Feng, Y. Xu, C. D. Hoang, M. Diehn, A. A. Alizadeh, Robust enumeration of cell subsets from tissue expression profiles., *Nat. Methods* **12**, 453–457 (2015).

58. D. Aran, Z. Hu, A. J. Butte, xCell: digitally portraying the tissue cellular heterogeneity landscape., *Genome Biol.* **18**, 220 (2017).

59. C. Marin-Muller, D. Li, U. Bharadwaj, M. Li, C. Chen, S. E. Hodges, W. E. Fisher, Q. Mo, M.-C. Hung, Q. Yao, A tumorigenic factor interactome connected through tumor suppressor microRNA-198 in human pancreatic cancer., *Clin. Cancer Res.* **19**, 5901–5913 (2013).

60. M.-C. Chang, C.-A. Chen, C.-Y. Hsieh, C.-N. Lee, Y.-N. Su, Y.-H. Hu, W.-F. Cheng, Mesothelin inhibits paclitaxel-induced apoptosis through the PI3K pathway., *Biochem. J.* **424**, 449–458 (2009).

61. Z. Tang, M. Qian, M. Ho, The role of mesothelin in tumor progression and targeted therapy., *Anticancer Agents Med Chem* **13**, 276–280 (2013).

62. M. Cremona, C. J. Vandenberg, A. M. Farrelly, S. F. Madden, C. Morgan, R. Kalachand, J. N. McAlpine, S. Toomey, D. G. Huntsman, L. Grogan, O. Breathnach, P. Morris, M. S. Carey, C. L. Scott, B. T. Hennessy, BRCA mutations lead to XIAP overexpression and sensitise ovarian cancer to inhibitor of apoptosis (IAP) family inhibitors., *Br. J. Cancer* **127**, 488–499 (2022).

63. Z. Su, Z. Yang, L. Xie, J. P. DeWitt, Y. Chen, Cancer therapy in the necroptosis era., *Cell Death Differ.* **23**, 748–756 (2016).

64. S. Hannes, R. Karlowitz, S. J. L. van Wijk, The Smac mimetic BV6 cooperates with STING to induce necroptosis in apoptosis-resistant pancreatic carcinoma cells., *Cell Death Dis.* **12**, 816 (2021).

65. L. Seifert, G. Werba, S. Tiwari, N. N. Giao Ly, S. Alothman, D. Alqunaibit, A. Avanzi, R. Barilla, D. Daley, S. H. Greco, A. Torres-Hernandez, M. Pergamo, A. Ochi, C. P. Zambirinis, M. Pansari, M. Rendon, D. Tippens, M. Hundeyin, V. R. Mani, C. Hajdu, G. Miller, The necrosome promotes pancreatic oncogenesis via CXCL1 and Mincle-induced immune suppression., *Nature* **532**, 245–249 (2016).

66. B. Strilic, L. Yang, J. Albarrán-Juárez, L. Wachsmuth, K. Han, U. C. Müller, M. Pasparakis, S. Offermanns, Tumour-cell-induced endothelial cell necroptosis via death receptor 6 promotes metastasis., *Nature* **536**, 215–218 (2016).

67. L. F. Hernandez, A. B. Dull, S. Korrapati, C. M. Annunziata, Smac-mimetic enhances antitumor effect of standard chemotherapy in ovarian cancer models via Caspase 8-independent mechanism., *Cell Death Discov.* **7**, 134 (2021).

68. L. Hernandez, M. K. Kim, A. M. Noonan, E. Sagher, H. Kohlhammer, G. Wright, L. T. Lyle, P. S. Steeg, M. Anver, D. D. Bowtell, Australian Ovarian Cancer Study Group 57, C. M. Annunziata, A dual role for Caspase8 and NF-κB interactions in regulating apoptosis and

necroptosis of ovarian cancer, with correlation to patient survival., *Cell Death Discov.* **1**, 15053
(2015).

69. L. R. Sanchez, L. Borriello, D. Entenberg, J. S. Condeelis, M. H. Oktay, G. S. Karagiannis,
The emerging roles of macrophages in cancer metastasis and response to chemotherapy., *J.*
Leukoc. Biol. **106**, 259–274 (2019).

70. D. W. Edwardson, A. M. Parissenti, A. T. Kovala, Chemotherapy and Inflammatory Cytokine
Signalling in Cancer Cells and the Tumour Microenvironment., *Adv. Exp. Med. Biol.* **1152**, 173–
215 (2019).

71. R. Hughes, B.-Z. Qian, C. Rowan, M. Muthana, I. Keklikoglou, O. C. Olson, S. Tazzyman, S.
Danson, C. Addison, M. Clemons, A. M. Gonzalez-Angulo, J. A. Joyce, M. De Palma, J. W.
Pollard, C. E. Lewis, Perivascular M2 Macrophages Stimulate Tumor Relapse after
Chemotherapy., *Cancer Res.* **75**, 3479–3491 (2015).

72. E. Hirata, E. Sahai, Tumor microenvironment and differential responses to therapy., *Cold*
Spring Harb. Perspect. Med. **7** (2017), doi:10.1101/cshperspect.a026781.

73. F. Balkwill, Tumour necrosis factor and cancer., *Nat. Rev. Cancer* **9**, 361–371 (2009).

74. B.-Z. Qian, J. W. Pollard, Macrophage diversity enhances tumor progression and
metastasis., *Cell* **141**, 39–51 (2010).

75. C. C. Uphoff, H. G. Drexler, Detection of mycoplasma contaminations., *Methods Mol. Biol.*
290, 13–23 (2005).

76. S. Potdar, F. Ianevski, A. Ianevski, Z. Tanoli, K. Wennerberg, B. Seashore-Ludlow, O.
Kallioniemi, P. Östling, T. Aittokallio, J. Saarela, Breeze 2.0: an interactive web-tool for visual
analysis and comparison of drug response data., *Nucleic Acids Res.* **51**, W57–W61 (2023).

77. Y.-L. Huang, C.-Y. Liang, D. Ritz, R. Coelho, D. Septiadi, M. Estermann, C. Cumin, N.
Rimmer, A. Schötzau, M. Núñez López, A. Fedier, M. Konantz, T. Vlajnic, D. Calabrese, C.
Lengerke, L. David, B. Rothen-Rutishauser, F. Jacob, V. Heinzelmann-Schwarz, Collagen-rich
omentum is a premetastatic niche for integrin α2-mediated peritoneal metastasis., *eLife* **9**
(2020), doi:10.7554/eLife.59442.

78. T. R. Henry, J. M. Spitsbergen, M. W. Hornung, C. C. Abnet, R. E. Peterson, Early life stage
toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in zebrafish (Danio rerio)., *Toxicol. Appl.*
Pharmacol. **142**, 56–68 (1997).

79. G. E. Elonen, R. L. Spehar, G. W. Holcombe, R. D. Johnson, J. D. Fernandez, R. J.
Erickson, J. E. Tietge, P. M. Cook, Comparative toxicity of 2,3,7,8-tetrachlorodibenzo- *p* -dioxin
to seven freshwater fish species during early life-stage development, *Environ. Toxicol. Chem.*
17, 472–483 (1998).

80. F. Jacob, R. Guertler, S. Naim, S. Nixdorf, A. Fedier, N. F. Hacker, V. Heinzelmann-
Schwarz, Careful selection of reference genes is required for reliable performance of RT-qPCR
in human normal and cancer cell lines., *PLoS ONE* **8**, e59180 (2013).

81. R. Casanova, S. Xu, S. Sivapatham, A. Jacobs, S. Engler, Tumor Profiler Consortium, M. P.
Levesque, R. Dummer, B. Bodenmiller, S. Chevrier, Standardization of suspension and imaging
mass cytometry readouts for clinical decision making, *BioRxiv* (2023),
doi:10.1101/2023.03.19.531228.

82. N. Kotecha, P. O. Krutzik, J. M. Irish, Web-based analysis and publication of flow cytometry

experiments., *Curr. Protoc. Cytom.* **Chapter 10**, Unit10.17 (2010).

83. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson,
T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner., *Bioinformatics* **29**, 15–21 (2013).

84. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2., *Genome Biol.* **15**, 550 (2014).

85. M. E. Ritchie, B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, G. K. Smyth, limma powers
differential expression analyses for RNA-sequencing and microarray studies., *Nucleic Acids*
Res. **43**, e47 (2015).

86. S. Hänzelmann, R. Castelo, J. Guinney, GSVA: gene set variation analysis for microarray
and RNA-seq data., *BMC Bioinformatics* **14**, 7 (2013).

1076

Figure 1. SMAC mimetics selectively target MSLN^{low} EOC cell lines. *A.* Heatmap depicting heterogeneous drug responses using a library of 528 compounds tested across 8 EOC cell lines with (△*MSLN for knockout; OE for overexpression*) and without genetic manipulation of MSLN expression. The represented values are drug sensitivity scores (DSS) colors indicating either sensitivity (red) or resistance (blue). **B.** Heatmap with drug responses of chemotherapeutic compounds commonly applied in EOC treatment. Visualization and hierarchical clustering of rows (1-Person correlation) was performed using Morpheus software. **C.** Dot plot illustrating the enhanced effectiveness of SMAC mimetics in cell lines characterized by low MSLN expression. **D**. Dose-response curves for birinapant across 4 different EOC cell lines. The data presented is the result of two independent experiments conducted in quadruplicate. **E.** Representative fluorescence images and quantification of a co-culture system using parental (EGFP) and △*MSLN* (dTomato) OVCAR3 cells treated with an increasing concentration of birinapant. The corresponding line chart summarizes the quantification of POPO™-1 iodide staining intensity after 48 hours of birinapant treatment. Scale bar 50μm. **F.** Representative fluorescent images of the mesothelial clearance assay in parental (left) and △*MSLN* (right) OVCAR3 cells with and without birinapant treatment. Cell viability was calculated based on POPO™-1 Iodide staining intensity and normalized against controls. Data is presented as mean ± SD of two independent experiments done in quadruplicate (D, E) or sextuplicates (F) *p* values were determined using two-way ANOVA followed by Sidak's multiple comparison test. (*, *p* <0.05, ***, *p*<0.001, **** *p*<0.0001). Scale bar 50μm.

Figure 2. SMAC mimetics enhance chemotherapy sensitivity. A. Representative Western blots depicting reduction of MSLN following exposure to carboplatin and paclitaxel in Kuramochi and OVCAR3 cells. **B.** Heatmap with increasing DSS values for the combination of SMAC mimetics with either carboplatin or paclitaxel across six EOC cell lines, sorted by MSLN expression levels. **C** and **D.** Synergy scores for SMAC mimetics in combination with either carboplatin or paclitaxel. **E.** Oncoprint for HGSOC samples treated *ex vivo*. Matched genomic data obtained from whole-exome sequencing, only genes with alterations in at least one patient sample are visualized. **Supplementary Table 3**. **F.** Tumor content and matched synergy scores for SMAC mimetics in combination with either carboplatin or paclitaxel in HGSOC patient-derived cells. **G** and **H.** Representative immunofluorescent images and quantification of "on target" tumor effect of the different treatments using an *ex vivo* culture system in three HGSOC samples. Carboplatin, paclitaxel, and birinapant were used at the final concentration of 10µM, 10nM, and 500nM, respectively. At least 20 cell aggregates were imaged and quantified for each condition in each patient sample resulting in 122,805 cells analyzed. Scale bar 50 µm. *P* values were determined using the Wilcoxon comparison test *, $p < 0.05$, ***, $p < 0.001$, **** $p < 0.0001$. **I.** Representative immunofluorescent images and quantification of COV362-Chili zebrafish xenografts, demonstrating a reduction in live COV362-Chili cells after 72 hours of exposure to the combination of birinapant (5000nM) with paclitaxel (1000nM). Data presented as a result of three independent experiments, each done in duplicate, *p* values were determined using the Wilcoxon comparison test. Scale bar 100µm.

1122
Figure 3. Birinapant restores the apoptotic effect of chemotherapy in a TNF⊡-dependent
manner. A and **B.** Representative Western blots demonstrate the TNF⊡ dependency for the

enhanced efficacy of birinapant in combination with either carboplatin or paclitaxel in Kuramochi
and OVCAR3 cell lines after 24 hours. Carboplatin, paclitaxel, and birinapant were used at the
final concentration of 10μM, 10nM, and 500nM, respectively. **C** and **D**. Representative
histograms and quantification of a real-time apoptosis reporter assay using FLIP-GFP as a
readout supporting the specificity of TNF☐ in increasing cell apoptosis mediated through
caspase 8 and 3 when combined with birinapant, and birinapant in combination with either
carboplatin or paclitaxel. Carboplatin, paclitaxel, and birinapant were used at the final
concentration of 10μM, 10nM, and 500nM, respectively. The data presented is the result of at
least three independent experiments. *P* values were determined by Wilcoxon comparison, * *p*
<0.05, ***, *p*<0.001, **** *p*<0.0001. **E.** Illustration of TNF☐ signaling pathway and histograms for
CRISPR-Cas9 cell competition assay over five consecutive passages depicting increased
resistance and sensitivity across different treatments upon gene editing of *MSLN* and TNF☐
pathways-related genes *NIK*, *IKKb, CASP8,* and *RIPK3*. **F.** Colony formation assay with
quantified colonies to assess treatment-dependent survival upon deletion of *AAVS1* (negative
control) or *CASP8* in OVCAR3-Cas9$^+$ cells. Carboplatin, paclitaxel, and birinapant were used at
the final concentration of 1μM, 1nM, and 5nM, respectively. The data presented is the result of
two independent experiments done in triplicates. *P* values were determined by Wilcoxon
comparison test.

1143

***Figure 4.* Macrophages in the TME promote TNF☐ pathway activation in HGSOC samples.**
A. Serum levels of CA125 in a patient with HGSOC during 36 months of follow-ups**. B** and **C.**
Representative immunofluorescence images and box plots depicting increased cancer cell
apoptosis (% of E-cadherin, cCASP3 positive cells) after exposure to birinapant in combination
with either carboplatin or paclitaxel. At least 20 cell aggregates were imaged and quantified for
each condition in each sample, in a total of 47'413 cells were analyzed. Scale bar 50µm. *P*
values were determined by the Wilcoxon comparison test. *p <0.05, **p <0.01, ***p<0.001. **D.**
Pathway activation scores (z-scored) for neoadjuvant HGSOC-derived patient samples (n=8)
with correction for the tumor content determined by CyTOF. Data generated from bulk RNAseq**.**
E. UMAPs depicting the major cell type composition across seven different HGSOC samples
using IMC. **F** and **G**. Heatmap and correlation matrix showing a significant association of
macrophage abundance and TNF☐ -related pathways activation. A grey asterisk indicates
significant correlations (r=0.82, *p*=0.034). **H.** Representative immunohistochemistry stainings in
HGSOC samples with high and low CD68 cellular content. Scale bar 100µm.
1158

1159

1160

Figure 5. Macrophages and TNF□ enhance response to bimodal treatments *ex vivo*. A.
Heatmap displaying the enrichment of myeloid cells (monocytes and M2 macrophages) in
samples with higher (green) and low (red) TNF□ gene set activation. Data sourced from bulk
RNA sequencing, based on CIBERSORT relative quantification of the immune compartment
using LM22 signature matrix. **B.** Boxplots depicting enrichment of M2 macrophages
accompanied with increased TNF□ pathway activation in omental metastasis in post-NACT
samples. Data source from bulk RNA sequencing of 97 samples based on xCell score, Primary
chemo_naive (n=30), Primary_NACT (n=24), Omentum metastasis_chemo_naive (n=19), and

Omentum_metastasis_NACT (n=24). **C** and **D** Representative immunofluorescence images and
corresponding box plots illustrate increased CD68$^+$ cell percentage in the 2nd relapse as
compared with the 1st relapse in PID_001. Quantification involved at least 16 cell aggregates
that were imaged and analyzed for each condition in every sample, in a total of 3'354 cells

analyzed. Scale bar: 50 µm, P values determined by t-test. **C** and **F** Representative

immunofluorescence images and box plots showing increased cancer cell apoptosis (% of E-

cad, cCASP3 positive cells) following bimodal treatments with TNFα supplementation as

compared with carboplatin in combination with paclitaxel. Quantification included ≥ 20 cell

aggregates that were imaged and quantified for each condition in each sample, in total 114'048

cells were analyzed. Scale bar: 50 µm. P values were determined by the Wilcoxon comparison

test. *p <0.05, **p <0.01.