ER-located PIN5 transporter generates potent auxin sinks driven by the IAA decarboxylation

pathway

Milada Čovanová, Karel Müller, Nikoleta Klierová, Nayyer Abdollahi Sisi, Petr Skůpa, Ondřej

Smetana, Petre Ivanov Dobrev, Kamil Růžička*, Jan Petrášek*

Institute of Experimental Botany, Czech Academy of Sciences, Rozvojová 263, 165 00 Prague

6, Czech Republic

*For correspondence (e-mail kamil.ruzicka@ueb.cas.cz, petrasek@ueb.cas.cz)

ER-resided PIN5 triggers IAA decarboxylation and PCD

ABSTRACT

Auxin is an essential and well-investigated regulator of plant development. Still, many aspects

determining its (sub-)cellular distribution, as well as its metabolic turnover, are obscure. PIN5

is a transporter that resides on the endoplasmic reticulum and is presumed to influence

internal auxin homeostasis by direct sequestration and subsequent degradation. Due to its

distinct expression pattern and incomplete metabolomics analyses, the exact role of PIN5

protein and the identity of downstream auxin degradation products show significant gaps. To

this end, we utilized morphologically homogeneous tobacco BY-2 cell cultures. We show that

the expression of *Arabidopsis thaliana AtPIN5* in the BY-2 system phenocopies so-called auxin

starvation defects. Moreover, we reveal that the activity of *At*PIN5 leads, in extreme cases, to

the broad range of processes accompanying programmed cell death (PCD). Notably, based on

the recently updated knowledge on auxin metabolism, we also show that a significant part of

auxin metabolites downstream of the *At*PIN5 activity are part of the re-emerged auxin

decarboxylation pathway. Taking together, we report the direct induction of PCD by auxin

stimulus and propose the physiological framework of the auxin decarboxylation route.

INTRODUCTION

The phytohormone auxin, or indole-3-acetic acid (IAA), is vital for plant growth and

development. The pattern formation at the level of individual tissues and organs is achieved

by the formation of local auxin maxima or minima, orchestrated by the cell-to-cell transport

of IAA (Weijers *et al.*, 2021). Among the mediators of auxin flow emerge the transporters of

the PIN-formed (PIN) family. Showing a remarkable asymmetrical distribution on the plasma

membrane (PM), they coordinate auxin fluxes across the tissues, thereby controlling the

downstream developmental programs (Adamowski & Friml, 2015).

Besides the well-investigated auxin fluxes between cells, mediated by the PM-located PINs,

another group of PIN proteins was described as residing at the endoplasmic reticulum (ER).

The so-called ER-PINs regulate internal auxin levels by conveying auxin into the ER lumen for

subsequent metabolization (Mravec *et al.*, 2009; Ding *et al.*, 2012). The metabolic products

downstream of the ER-located PIN activity have been examined several times and with various

degrees of consistency (Mravec *et al.*, 2009; Ding *et al.*, 2012; Abdollahi Sisi & Růžička, 2020).

In addition, the pathways accompanying the IAA breakdown have recently undergone revision

(Müller *et al.*, 2021; Hayashi *et al.*, 2021; Dobrev *et al.*, 2023), highlighting the IAA

decarboxylation as another relevant auxin degradation route (Bandurski *et al.*, 1995; Dobrev

et al., 2023). Thus, many aspects of the IAA homeostasis controlled by ER-PINs remain unclear

(Abdollahi Sisi & Růžička, 2020; Ung *et al.*, 2023a).

The connection between auxin and programmed cell death (PCD) has been repeatedly

proposed (Kacprzyk *et al.*, 2022). It was shown that auxin pulses produced from the cells

undergoing PCD in the outermost layers of the root cap are required for priming lateral root

primordia in the primary root meristem (Xuan *et al.*, 2015, 2016). It was also revealed that

auxin can suppress certain instances of PCD (Kerchev *et al.*, 2013). It has even been

hypothesized that auxin is a product of tryptophan degradation occurring in defunct cells

(Sheldrake, 2021). Nevertheless, the authentic effect of auxin activity to trigger PCD has never

been presented.

Cell-suspension cultures have proved to be an excellent model system, which enables

studying the effects of growth substances directly and in a homogeneous manner (Petrášek &

Zazímalová, 2006; Petrasek *et al.*, 2006). Here, we explore the impact of conditional

expression of the well-characterized ER-located *Arabidopsis thaliana At*PIN5 (Mravec *et al.*,

2009) in the BY-2 tobacco suspension cultures. We comprehensively analyze the immediate

IAA metabolites downstream of PIN5 activity and reveal the remarkably elevated presence of

products of the IAA decarboxylation pathway (Dobrev *et al.*, 2023). We also describe the

corresponding phenotypic consequences, including the extreme instances of PCD.

RESULTS

Inducible expression of *At*PIN5 in BY-2 cells phenocopies the symptoms of auxin starvation

The growth and proliferation of tobacco BY-2 cells depends on the exogenous supply of

auxin to the media. Its lack is characterized by enhanced cell elongation and lowered cell

division, termed collectively as symptoms of auxin starvation (Winicur *et al.*, 1998). We

presented earlier that interfering with auxin transport leads to similar morphological defects.

The induction of the transgene encoding a PM-located *Arabidopsis thaliana At*PIN7 auxin

carrier enhances cell elongation and division arrest, consistent with that occurring on the

auxin-free medium (Fig. 1, A to C and Supplemental Fig. S1C; Mravec *et al.*, 2008; Müller *et al.*,

2019). We observed similar auxin-deficient phenotypes by perturbing the nuclear auxin

signaling by the inducible expression of the stabilized version of the auxin signaling repressor

*At*IAA17, *At*axr3-1 (Leyser *et al.*, 1996; Knox *et al.*, 2003; Figs. 1D-F). We were also able to

pharmacologically reduce cell proliferation by the dose-dependent application of IAA

antagonist PEO-IAA, which competitively binds the TIR1/AFBs auxin receptors (Nishimura *et*

al., 2009; Fig. S1B). This indicates that the auxin starvation phenotypes are dependent on the

TIR1/AFB signaling, in addition to the availability of auxin in the cells.

Next, to study the impact of internal auxin fluxes on the cell culture phenotypes, we

expressed in the BY-2 cells the characterized *Arabidopsis thaliana* gene *AtPIN5* (Mravec *et al.*,

2009; Abdollahi Sisi & Růžička, 2020). We employed two independent transcription inducible

systems, XVE (Zuo *et al.*, 2000; Petrasek *et al.*, 2006) and GVG (Aoyama & Chua, 1997; Mravec

et al., 2008), triggering the expression of the transgene at the beginning of the subculture

interval. Upon the induction of *At*PIN5, the cells became remarkably expanded (Fig. 1G and

1H; Supplemental Fig. S1D and S1E), and their cell division reduced (Fig. 1I and Supplemental

Fig. S1F), akin to the auxin starvation phenotypes (Fig. 1 A to F and Supplemental Fig. S1C). In

addition, the auxin-deficient phenotypes caused by the ER-located *At*PIN5 corresponded to

the lowered expression of the auxin reporter *DR5::GFP* (Supplemental Fig. S1G; see also

Barbez *et al.*, 2013). This indicates that the *At*PIN5 activity is capable of changing the internal

availability of auxin to control the growth and expansion of the BY-2 cells.

IAA enhances the symptoms of *At*PIN5-induced auxin starvation

The dependence of the BY-2 tissue cultures on the addition of the synthetic auxin (2,4-

dichlorophenoxyacetic acid or 2,4-D) to the media is ascribed to the low levels of endogenous

IAA in the cells, caused by its rapid metabolization (Simon *et al.*, 2013; Seifertová *et al.*, 2014;

Müller *et al.*, 2021). Accordingly, substituting 2,4-D with IAA also leads to auxin starvation

phenotypes (Simon *et al.*, 2013; Fig. 2A). To test the role of internal auxin fluxes under

conditions where only native auxin is present, we cultivated the cells expressing *AtPIN5* under

the supply of IAA, too. The control lines proliferated less intensively and resembled the auxin-

starved ones. Surprisingly, the induction of *At*PIN5 led to even stronger auxin starvation

phenotypes (Fig. 2, B and C). Thus, it seems that the PIN5-like transporters can enhance rapid

IAA turnover in BY-2 cells.

To follow the original idea and to compensate for the fast IAA metabolization more

efficiently, we were continuously adding IAA to the media every 12 h during the subcultivation

interval. We observed no apparent signs of auxin deficiency (Fig. 2, D and E). Strikingly, though,

the *AtPIN5* induction caused strong auxin starvation phenotypes (Fig. 2, D and E). This further

underlines that PIN5 is able to generate a potent intracellular IAA sink.

ER-PINs are capable of transporting both IAA and 2,4-D (Ung *et al.*, 2022). Hence, we

measured the cellular accumulation of radioactively labeled [^{3}H]-IAA and [^{3}H]-2,4-D following

the induction of *AtPIN5*. The lines showed a strong cellular uptake of [³H]-IAA (Fig. 2F).

Interestingly, we observed no change in the *At*PIN5-mediated [³H]-2,4-D accumulation (Fig.

2G). Altogether, our data suggest that *At*PIN5 efficiently channels IAA from the cytosol to be

delivered to likely highly active metabolic enzymes inside the ER lumen.

Induction of *AtPIN5* promotes programmed cell death

To validate whether the strong auxin withdrawal from the cytosol by *At*PIN5 corresponds

to a native downstream process, we analyzed the viability of the cells examined. The trypan

blue exclusion test (van Wees, 2008), revealed an increased ratio of cells showing a dye-

positive cytoplasm shrinkage as a sign of decreased cell viability following the *At*PIN5 induction

(Fig. 3, A and B and Supplemental Fig. S2, A and B). Moreover, the intracellular trypan blue

staining was even more apparent in the cells cultured on IAA (Fig. 3, A and B). Thus, the PIN5-

mediated funneling of auxin from the cytosol can lead to increased mortality of BY-2 cells.

Further, we analyzed the global transcriptional changes downstream of the *At*PIN5 activity

to hint at the possible relevant physiological processes related to auxin starvation and lower

cell viability. Indeed, using the PANTHER overrepresentation test (Mi *et al.*, 2019), we found a

remarkable overlap with the previously published transcriptional profiles from the BY-2 cells

cultured in the auxin-free medium (Müller et al., 2021; Fig. 3C and Supplemental Fig. S2, C and

D). This included pronounced changes in the expression of stress-related genes, including

peroxidases (Supplemental Table S1). We further examined the transcriptional response

exclusively linked with the *At*PIN5 expression (Supplemental Table S2), revealing a single

enriched gene ontology term (GO:0000041) supported by 6 individual transcripts coding for

transporters of transition metals (Peñarrubia *et al.*, 2015). This suggests that the auxin

starvation is associated with the enhanced expression of the stress-related peroxidases and

that the *AtPIN5* expression leads to the enhanced metal ion transport, presumably activating

the response preceding the cell death.

The increased expression of peroxidases following the *AtPIN5* induction suggests elevated

activity of produced reactive oxygen species (ROS). The role of auxin has been implicated in

the crosstalk with ROS homeostasis (Tognetti *et al.*, 2012; Chen *et al.*, 2014b). Hence, we

utilized a sensor carboxy-2',7'-dichloro-dihydro-fluorescein diacetate (H2DCFDA), which

indicates the intracellular presence of various ROS, including H_2O_2 (Akter *et al.*, 2021). Indeed,

upon induction of *AtPIN5*, the cells displayed significantly increased H2DCFDA fluorescence,

corroborating the enhanced ROS activity (Fig. 3, D and E and Supplemental Fig. S2E). PIN5 was previously shown to be a part of ER stress adaptive mechanisms (Chen *et al.*, 2014a). Although we did not notice that the repeated additions of IAA into the media would influence the ER localization of *At*PIN5-GFP or lead to remarkable changes in the overall ER structural pattern (Supplemental Fig. S1G and S2F), it was earlier reported that pathways involving ROS production and ER stress are often preceding PCD (Petrov *et al.*, 2015). This thereby further highlights that the lower viability of the *At*PIN5-GFP cells accompanied by ROS can perhaps be ascribed to the initiation of processes related to PCD.

Among the main features of PCD are vacuolar leakage and subsequent acidification of cytoplasm (Valandro *et al.*, 2020). To this end, we utilized 2',7'-Bis-(2-carboxyethyl)-5-(and-6)-carboxyfluorescein acetoxymethyl ester (BCECF/AM), a fluorescent dye used for analyzing lowered cytosolic pH as a mark of disrupted vacuolar integrity (Scheuring *et al.*, 2015). Under control conditions, the cells contained several smaller vacuoles. The induction of *At*PIN5 led to the enlargement of central vacuoles, their occasional collapse and acidification of the whole cytoplasm and nucleus (Fig. 3, F and G). Finally, we tested the nuclear DNA fragmentation, a hallmark of PCD, using terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) (Tripathi *et al.*, 2017). The expression of *AtPIN5* (Fig. 3, H and I, and Supplemnetal Fig. S3A) resulted in an increased number of TUNEL-stained nuclei, indicating that DNA fragmentation takes part in the signaling cascade downstream of the rapid depletion of auxin in the cytoplasm. However, the decreased cell viability does not appear a general result of deficient auxin response. We also examined the TUNEL signal in cells expressing *AtPIN7* and *Ataxr3-1* in the medium supplemented with IAA in the 12-hour interval. We observed no relevant staining in any of these lines (Supplemental Fig. S3B).

We also tested whether the possibly disintegrated cell content can trigger (and enhance) the marks of auxin-induced starvation and PCD. We cultured wild-type BY-2 cells in a medium supplemented with the homogenized cell debris. The cell division rate remained unchanged (Supplemental Fig. S4A), and no obvious signs of increased cell mortality were observed (Supplemental Fig. S4B). This further evidences that the cell death induced by PIN5 expression is a result of the internal cellular activity, rather than being caused by any compound originating from disintegrated cells.

182   Taken together, we conclude that the observed decreased cell viability is associated with

183 PCD, but not uncontrolled stress-related necrotic processes (Van Durme & Nowack, 2016),

184 and it is a specific hallmark of PIN5-mediated auxin starvation.

186 The activity of *At*PIN5 stimulates auxin degradation by the IAA decarboxylation pathway

To get insight into how the ER-mediated IAA conversion affects internal auxin homeostasis

in light of recent findings (Müller *et al.*, 2021; Hayashi *et al.*, 2021; Dobrev *et al.*, 2023; Fig.

4A), we have tracked the expanded list of IAA metabolites by liquid chromatography-mass

spectrometry (LC-MS) following the *AtPIN5* induction. At the time point of adding IAA into

media (t_0), its degradation products were barely detectable and their amounts showed

negligible differences between the mock and chemical inducer-treated cells (Supplemental

Fig. S5A). 2 hours later, we observed that IAA was converted predominantly to amino-acid

conjugates of IAA (Fig. 4B; see also Seifertová *et al.*, 2014; Müller *et al.*, 2021; Dobrev *et al.*,

2023). The induction of *AtPIN5* usually led to a further increase in IAA conversion in this

experimental setup (Fig. 4B and Supplemental Fig. S5B). These measurements illustrate rapid

catabolism of IAA in BY-2 cells and evidence that the majority of detected metabolic

compounds indeed originate from the supplied IAA.

Next, we quantified the IAA breakdown in the cells cultivated in a media supplied with IAA

every 12 hours from the beginning of the subcultivation interval. 2 hours after addition, IAA

was converted in the control lines into its conjugates as well (Fig. 4C). Surprisingly, the prior

induction of *AtPIN5* dramatically reduced the internal IAA levels to close to trace amounts.

Although we were also able to detect the activity of the common oxidative IAA degradation

pathway, we found remarkably increased conversion towards IAA decarboxylation in these

lines (Fig. 4C), as documented by the elevated levels of ascorbigen (Abg) and

dihydroascorbigen (DiHAbg) (Dobrev *et al.*, 2023; Fig. 4C). We therefore conclude that the

PIN5-mediated auxin sinks are at least partially accompanied by the increased activity of the

auxin decarboxylation pathway.

210 The exogenous application of I3C raises the symptoms of cell death in BY-2 cells

211   It has been proposed earlier that indole-3-carbinol (I3C), an unstable immediate IAA

212 decarboxylation product and precursor of Abg (Fig. 4A) (Hrnčiřík *et al.*, 1998; Chhajed *et al.*,

2020; Dobrev *et al.*, 2023), can function as an auxin antagonist and can stimulate vacuolar disintegration and processes related to autophagy in *Arabidopsis* in consequence of the defense reaction (Katz *et al.*, 2015; Katz & Chamovitz, 2017). We therefore incubated the cells with I3C to relate its role to the PIN5-mediated programmed cell death in the BY-2 cultures. Despite the default presence of 2,4-D in the media, I3C treatment affected the cell expansion and proliferation rates (Fig. 5, A and B). I3C also slightly lowered their overall viability (Fig. 5, C and D). However, it remarkably increased the frequency of PCD, as documented by TUNEL staining (Fig. 5, E and F). Taken together, we propose that ER-resided PINs are capable of generating powerful IAA sinks to activate IAA decarboxylation activity and, hereafter, pathways leading to PCD.

DISCUSSION

A possible mechanism underlying PIN5-mediated auxin sinks

The classical concept of polar auxin transport has premised the existence of efflux carriers asymmetrically localized at PM as the central prerequisite for the directional flow of IAA (Rubery & Sheldrake, 1974; Raven, 1975). Later, the PIN auxin transporters were discovered, matching henceforth the proposed attributes of the main IAA carriers (Gälweiler *et al.*, 1998; Petrasek *et al.*, 2006). ER-located PINs were found subsequently, and the relevant works highlighted their capability to influence internal auxin homeostasis (Mravec *et al.*, 2009; Ding *et al.*, 2012; Abdollahi Sisi & Růžička, 2020). Further, the detailed crystallographic studies evidenced that the transporting mechanism employed by both PM and ER-located PINs is identical (Su *et al.*, 2022; Ung *et al.*, 2022; Yang *et al.*, 2022).

Yet, many aspects of the PIN-mediated transport remain obscure. Compelling arguments supported by the structural models suggest that IAA is transported from the cytosol against concentration gradients at the expense of membrane potential (Su *et al.*, 2022; Ung *et al.*, 2022, 2023a,b; Yang *et al.*, 2022; Andersen *et al.*, 2023). Indeed, in the root apical meristem, PM-located PINs concentrate auxin in the uttermost root tip (Adamowski & Friml, 2015). They also establish and maintain auxin maxima, determining the position of flower primordia in the shoot apical meristem in an equivalent manner (Reinhardt *et al.*, 2003). In contrast, the canalization hypothesis proposes that auxin is funneled away from the site of its external application, i.e., source, towards the sink in the established vascular tissue determining the

connective vascular strands (Mitchison, 1980; Sachs, 1991; Ravichandran *et al.*, 2020).
Similarly, the root or hypocotyl tropic reactions depend on the transport outwards from the
tissue with higher auxin concentration (Friml *et al.*, 2002; Ottenschläger *et al.*, 2003; Rakusová
et al., 2011; Band *et al.*, 2012). The expression of PIN5 proteins leads to a remarkable auxin
deficiency in BY-2 cells. In the conditions when 2,4-D does not interfere with the internal IAA
fluxes (Fig. 5B), the excess of IAA leads to even enhanced PIN5-mediated auxin degradation
rates. Hence, in this system, the PIN5-like auxin transporters likely operate in the source-sink
mode, too, perhaps by the mechanism of facilitated diffusion, as earlier presumed for auxin
canalization (Mitchison, 1980; Rolland-Lagan & Prusinkiewicz, 2005).

Auxin metabolization as a prompt for PCD

Along with the auxin-dependent processes, PCD is an inherent part of plant development.
The connections between auxin and PCD pathways have been proposed, but mostly indirectly
(reviewed in Van Durme & Nowack, 2016; Sheldrake, 2021; Kacprzyk *et al.*, 2022). A chemical
screen performed by Kerchev *et al.* (2013) revealed that the H_2O_2-dependent cell death
caused by defective photorespiration can be reverted by both synthetic auxins and IAA. Auxin
was demonstrated as a central regulator of xylem patterning, where PCD gives rise to
conductive vessels (Escamez & Tuominen, 2014; Ruzicka *et al.*, 2015); direct
transdifferentiation of the tissue cultures into xylem vessels can also be induced by external
cues (Fukuda & Komamine, 1980; Kubo *et al.*, 2005; Ruzicka *et al.*, 2015). In addition, various
stages of lateral root development are determined by auxin activity in parallel to PCD
(Casimiro *et al.*, 2001; Benková *et al.*, 2003; Swarup *et al.*, 2008; Escamez *et al.*, 2020). Auxin
serves as a signal produced by the cells undergoing PCD in the root cap to prime the
prospective lateral roots in the primary root meristem (Xuan *et al.*, 2015, 2016). Ultimately, Di
Mambro et al. (2019) presented the model that the activity of *PIN5* in the lateral root cap
affects root meristem size, but the direct influence of PIN5 on PCD in the lateral root cap has
not been investigated.

We show that the ER-mediated auxin sinks can trigger PCD in BY-2 cultures. We have not
noticed any cells resembling forming xylem vessels following the *AtPIN5* induction. Instead,
we report phenotypes associated with common PCD and present lines of evidence that they
are linked with the auxin decarboxylation activity. I3C is among native metabolites of

glucosinolates, a diverse group of small compounds produced predominantly in *Brassicales*,

functioning as agent in the insect defense (Fahey *et al.*, 2001). It is also a product of auxin

decarboxylation pathway in a multitude of the (non-)*Brassicales* plant species (Bandurski *et*

al., 1995; Kerk *et al.*, 2000; Dobrev *et al.*, 2023). I3C can induce some symptoms of PCD in

Arabidopsis, in a competitive manner to IAA (Katz *et al.*, 2015; Katz & Chamovitz, 2017). Hence,

our data suggest that the two pathways determining PCD overlap and share an unexpectedly

common and likely evolutionarily conserved functional output.

IAA decarboxylation pathway downstream of PIN5-mediated auxin sequestration

PIN5 transports auxin into the ER in order to promote auxin degradation (Mravec *et al.*,

2009; Ung *et al.*, 2023a). The levels of IAA-aspartate and IAA-glutamate conjugates are

elevated following *At*PIN5 overexpression (Mravec *et al.*, 2009). Paradoxically, the GRETCHEN

HAGEN (GH3) enzymes catalyzing the IAA conjugation are generally seen in the cytoplasm

(Ludwig-Müller *et al.*, 2009; Ostrowski *et al.*, 2014; Di Mambro *et al.*, 2019; Abdollahi Sisi &

Růžička, 2020; Zou *et al.*, 2022). We have recently uncovered a new level of complexity of the

IAA metabolome. Among known IAA breakdown routes emerges the auxin decarboxylation

pathway, which includes Abg and DiHAbg. Even if occurring across angiosperms, the purpose

of auxin decarboxylation is unknown (Dobrev *et al.*, 2023). Strikingly, we found that the

relevant metabolic compounds are related to increased AtPIN5 activity. Hence, we conclude

that IAA decarboxylation is the main and potent auxin degradation pathway downstream of

PIN5.

EXPERIMENTAL PROCEDURES

Plant material, growth conditions and treatments

The tobacco cell lines BY-2 (*Nicotiana tabacum* L., cv. Bright Yellow-2; Nagata *et al.*, 1992)

were standardly cultivated in a liquid medium supplemented with 1 µM 2,4-D as described in

Petrasek *et al.* (2006) unless stated otherwise. The transformation of cell lines, RT-qPCR

validation of the transgene expression, and cell line maintenance were done as described

earlier (Petrasek *et al.*, 2006; Müller *et al.*, 2021). By rule, the cells in the exponential growth

stage were used, typically 4-d-old, unless specified otherwise.

The expression of the *AtPIN7, AtPIN5* and *Ataxr3-1* genes was induced by 1 µM
dexamethasone (30 mM stock solution, GVG lines) or 3 µM β-estradiol (30 mM stock solution,
XVE lines) at the beginning of the subculture interval. In addition, 2-(1H-indol-3-yl)-4-oxo-4-
phenyl-butyric acid (PEO-IAA; OlChemIm, Olomouc, Czech Republic, 10 mM stock solution)
and indole-3-carbinol (I3C; Sigma-Aldrich, Merck KGaA, Darmstadt, Germany, 160 mM stock
solution) were used. The respective compounds were applied to the cells at the beginning of
the subculture interval, and the corresponding amount of the solvent (DMSO) was added to
the controls.

Molecular transgenic work

The dexamethasone-inducible gene constructs, *35S>>GVG::AtPIN5* and
35S>>GVG::AtPIN7, including the corresponding cell cultures, have been described earlier
(Petrasek *et al.*, 2006; Mravec *et al.*, 2009). For creating *G10-90>>XVE::AtPIN5*, the
corresponding *AtPIN5* open reading frame was cloned with the Gateway system (Invitrogen,
Thermo Fisher Scientific, Waltham, MA, USA) into the pMDC7 vector (Curtis & Grossniklaus,
2003). The respective Gateway entry clones were generated either by the amplification of
wild-type DNA template or that originating from the *PIN5-GFP*[MGS] plants (Verna *et al.*, 2015)
(primer sequences AAAAAGCAGGCTATATGATAAATTGTGGAGAT and
AAGAAAGCTGGGTTTCAATGAATAAACTCC). The *XVE::AtPIN5-GFP/DR5::RFP* cell line was
obtained by transformation of the *XVE::AtPIN5-GFP* line with the *DR5::RFP* construct (Barbez
et al., 2013). The *G10-90>>XVE::Ataxr3-1* line was generated by the transformation of BY-2
cells with the construct kindly provided by Jiří Friml.

Microscopy and image analysis

Nomarski differential interference contrast (DIC) microscopy was performed with the
Nikon Eclipse E600 microscope (Nikon, Tokyo, Japan) equipped with a color digital camera
(DVC 1310C, Austin, Texas, USA). Cell length and diameter were measured manually using the
Fiji image analysis software (Rueden *et al.*, 2017). 200 cells in each data set were measured.
Cell density was determined by counting the cells with the Fuchs-Rosenthal hemocytometer
slide, typically in the fourth day of subcultivation or periodically during the entire period;
individual values represent the average of at least six aliquots of every sample. For the
fluorescence imaging, Zeiss LSM900 laser scanning microscope (Axio Observer 7, inverted)
with Airyscan 2 with Multiplex mode (Carl Zeiss, Jena, Germany) with appropriate filter sets

to visualize reporter genes GFP (excitation 488 nm and emission 509 nm wavelengths) and/or mRFP (excitation 590 nm and emission 599 nm wavelengths) was used. Detailed images were obtained with the 40x or 63x water/oil immersion objectives.

Assays on cell viability, ROS content and PCD

The trypan blue exclusion test was used to visualize the penetration of the non-viable cells by the dye (van Wees, 2008). Approximately 10 µl of the 0.4 % trypan blue solution (Sigma-Aldrich) was added to the 0.5 ml cell suspension and recorded subsequently. Cell viability was determined as the percentage of dye-negative cells relative to the total number of cells.

For the detection of reactive oxygen species (ROS), the cells were incubated with the 10 µM H2DCFDA (2',7'-dichlorfluorescein-diacetate; Thermo Fisher Scientific, Waltham, Massachusetts, USA) solution (Akter *et al.*, 2021) for 15 min and washed twice with the BY-2 medium. The relative ROS accumulation is presented as the mean pixel intensity of cells determined by the Zeiss Zen software. In total, about four hundred cells were counted from three independent experiments. The integrity of the vacuoles was determined after 10 min of staining with 10 µM BCEF-AM ((2',7'-bis-2-carboxyethyl)-5-(and-6) carboxyfluorescein; Sigma-Aldrich; Scheuring *et al.*, 2015). The number of cells with aberrant vacuolar integrity was assessed as a ratio of cells with the cytoplasmic signal to that of the cells showing a normal vacuolar signal. About five hundred cells were processed in three independent experiments. In both cases, the confocal images were acquired with the Plan-Apochromat 10x/0.45 DIC M27 objective.

TUNEL (terminal deoxynucleotidyl transferase dUTP nick end labeling) assay (Tripathi *et al.*, 2017) was performed in 2-day-old BY-2 cells according to the manufacturer's protocol (TMR red in situ cell death detection kit, Roche Diagnostics GmbH, Heidelberg, Germany) with modifications as described earlier (Smetana *et al.*, 2012). The labeled cells were examined using a color digital camera (DVC 1310C, Austin, Texas, USA) at the Nikon Eclipse E600 microscope. All cells were analyzed from at least 10 view fields, covering approximately 100 cells each. The quantification is expressed as a percentage of cells carrying the TUNEL signal divided by the total cell number in the visual field. Positive controls, where the samples were additionally treated with DNAse I to detect the positive TUNEL reaction, and negative controls, where the TdT enzyme was omitted from the procedure, were imaged alongside with every experiment.

367   To treat wild-type BY-2 cells with cell debris, one-quarter volume of the total 2-day-old cell

368   culture was collected. The cells were ground with a pestle in a mortar in liquid nitrogen. The

369   homogenized cell content was then washed out by the original medium, and the flowthrough

370   was then returned to the growing cell culture. The cells were subsequently cultivated for two

371   more days before examining their physiological properties.

Auxin uptake assays

Auxin uptake assays were performed in 3-day-old cells as described earlier (Delbarre *et al.*,

1996; Petrásek *et al.*, 2003). [^{3}H]-2,4-D (10 Ci.mmol^{-1}) or [^{3}H]-IAA (25 Ci.mmol^{-1}) (American

Radiolabeled Chemicals, Inc., St Louis, MO, USA) were added to the equilibrated cell

suspension at the beginning of the accumulation assay to the final concentration of 2 nM. The

presented values are means of each triplicate from a representative experiment out of three

biological repetitions.

IAA metabolic profiling

The harvesting of the 2-d-old BY-2 cells was done 2 hours after the addition of 1 μM IAA

into the media, unless stated otherwise. Auxin metabolites were extracted according to

previously described procedures (Dobrev & Kamínek, 2002; Dobrev *et al.*, 2023), and

determined by the LC-MS quantitative analysis further detailed in Dobrev *et al.* (2023). The

samples were processed in triplicates. The results are presented as a mean of at least three

independent biological repetitions.

Transcriptome analysis

The total RNA was isolated using RNeasy Plant Mini kit (Qiagen, Düsseldorf, Germany) from

50-100 mg of cells cultured in medium for 48 h with IAA added every 12 h, 2 h after the last

addition of IAA. Isolated RNA was treated with DNA-Free kit (Thermo Fisher Scientific). RNA

purity, concentration and integrity were evaluated on 0.8% agarose gels (v/w) and with the

Bioanalyzer RNA Nano 6000 Assay Kit (Agilent Technologies, Santa Clara, CA, USA). RNA

sequencing was performed using Illumina NovaSeq 6000 platform. The analysis resulted in at

least 15 million 150 bps read pairs. The raw reads were quality-filtered using Rcorrector and

Trim Galore scripts (Song & Florea, 2015). Transcript abundances (transcripts per million –

TPM) were determined using the Salmon software package (Patro *et al.*, 2017) with

parameters --posBias, --seqBias, --gcBias, --numBootstraps 30. The index was built from the

Nicotiana tabacum v1.0 cDNA dataset (Edwards *et al.*, 2017), by including the *AtPIN5*

(*AT5G16530*) gene sequence. Visualization, quality control of data analysis and determination

of differentially expressed genes were determined using the sleuth package in R (Love *et al.*,

2014; Pimentel *et al.*, 2017) (version 0.29.0). Transcripts with the q-value <= 0.05 and $\log_2$ fold

change >=1 (upregulated) or <= -1 (down-regulated) were considered to be significantly

differentially expressed. Other criteria for selection of gene list of interest are specified in

corresponding result section.

Gene ontology analysis (statistical overrepresentation test) was done using the Panther

Classification System (Mi *et al.*, 2019). Raw sequencing data are available through GEO Series

accession number GSE260945 at the Gene Expression Omnibus

(https://www.ncbi.nlm.nih.gov/geo/). The grouping of differentially expressed genes was

visualized with the DiVenn software (Sun *et al.*, 2019).

ACKNOWLEDGEMENTS

We thank Jiří Friml for kindly providing the *Ataxr3-1* construct, Enrico Scarpella for the *PIN5-*

GFP^{MGS} lines, Zuzana Vondráková and Nikola Drážná for technical assistance, and Kateřina

Malínská for help with image analysis. The work was supported by the Czech Science

Foundation (project 19-23773S) to KR, JP and PID. Microscopy was done at the Imaging Facility

of the Institute of Experimental Botany and was supported by the Ministry of Education, Youth

and Sports of the Czech Republic (LM2023050 Czech-Bioimaging).

AUTHOR CONTRIBUTIONS

JP and KR conceived the research. JP, KR and MČ designed the experiments. MČ performed

most experiments. KM did RT-qPCR and transcriptome analysis. NK and NAS prepared DNA

constructs. PS provided preliminary data essential for the initiation of the study. OS performed

part of the PCD assays. PID did metabolic profiling. KR, JP and MČ wrote the manuscript. KR

acquired funding. All authors have exercised their right to comment on the manuscript.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Abdollahi Sisi N, Růžička K. **2020**. ER-Localized PIN Carriers: Regulators of Intracellular Auxin Homeostasis. *Plants* **9**: 1527.

Adamowski M, Friml J. **2015**. PIN-Dependent Auxin Transport: Action, Regulation, and Evolution. *The Plant Cell* **27**: 20–32.

Akter S, Shahneawz Khan M, N. Smith E, Flashman E. **2021**. Measuring ROS and redox markers in plant cells. *RSC Chemical Biology* **2**: 1384–1401.

Andersen CG, Bavnhøj L, Pedersen BP. **2023**. May the proton motive force be with you: A plant transporter review. *Current Opinion in Structural Biology* **79**: 102535.

Aoyama T, Chua NH. **1997**. A glucocorticoid-mediated transcriptional induction system in transgenic plants. *The Plant Journal: For Cell and Molecular Biology* **11**: 605–612.

Band LR, Wells DM, Larrieu A, Sun J, Middleton AM, French AP, Brunoud G, Sato EM, Wilson MH, Péret B, *et al.* **2012**. Root gravitropism is regulated by a transient lateral auxin gradient controlled by a tipping-point mechanism. *Proceedings of the National Academy of Sciences* **109**: 4668–4673.

Bandurski RS, Cohen JD, Slovin JP, Reinecke DM. **1995**. Auxin biosynthesis and metabolism. In: Davies PJ, ed. Plant Hormones. Physiology, Biochemistry and Molecular Biology. Dordrecht: Springer, 39–65.

Barbez E, Laňková M, Pařezová M, Maizel A, Zažímalová E, Petrášek J, Friml J, Kleine-Vehn J. **2013**. Single-cell-based system to monitor carrier driven cellular auxin homeostasis. *BMC plant biology* **13**: 20.

Benková E, Michniewicz M, Sauer M, Teichmann T, Seifertová D, Jürgens G, Friml J. **2003**. Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* **115**: 591–602.

Casimiro I, Marchant A, Bhalerao RP, Beeckman T, Dhooge S, Swarup R, Graham N, Inzé D, Sandberg G, Casero PJ, *et al.* **2001**. Auxin transport promotes Arabidopsis lateral root initiation. *The Plant Cell* **13**: 843–852.

Chen Y, Aung K, Rolčík J, Walicki K, Friml J, Brandizzi F. **2014a**. Inter-regulation of the unfolded protein response and auxin signaling. *The Plant Journal: For Cell and Molecular Biology* **77**: 97–107.

Chen C, Letnik I, Hacham Y, Dobrev P, Ben-Daniel B-H, Vanková R, Amir R, Miller G. **2014b**. ASCORBATE PEROXIDASE6 Protects Arabidopsis Desiccating and Germinating Seeds from Stress and Mediates Cross Talk between Reactive Oxygen Species, Abscisic Acid, and Auxin. *Plant Physiology* **166**: 370–383.

Chhajed S, Mostafa I, He Y, Abou-Hashem M, El-Domiaty M, Chen S. **2020**. Glucosinolate Biosynthesis and the Glucosinolate–Myrosinase System in Plant Defense. *Agronomy* **10**: 1786.

Curtis MD, Grossniklaus U. **2003**. A gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiology* **133**: 462–469.

Delbarre A, Muller P, Imhoff V, Guern J. **1996**. Comparison of mechanisms controlling uptake and accumulation of 2,4-dichlorophenoxy acetic acid, naphthalene-1-acetic acid, and indole-3-acetic acid in suspension-cultured tobacco cells. *Planta* **198**: 532–541.

Di Mambro R, Svolacchia N, Dello Ioio R, Pierdonati E, Salvi E, Pedrazzini E, Vitale A, Perilli S,
Sozzani R, Benfey PN, *et al.* 2019. The Lateral Root Cap Acts as an Auxin Sink that Controls Meristem
Size. *Current Biology* **29**: 1199-1205.e4.

Ding Z, Wang B, Moreno I, Dupláková N, Simon S, Carraro N, Reemmer J, Pěnčík A, Chen X, Tejos R,
***et al.* 2012**. ER-localized auxin transporter PIN8 regulates auxin homeostasis and male gametophyte
development in *Arabidopsis*. *Nature Communications* **3**: 941.

Dobrev PI, Filepová R, Lacek J, Vondráková Z, Müller K, Maršík P, Drašarová L, Talacko P, Hošek P,
Petrášek J. **2023**. Study of auxin metabolism using stable isotope labeling and LCMS; evidence for in
planta auxin decarboxylation pathway. : 2023.06.02.543384.

Dobrev PI, Kamínek M. **2002**. Fast and efficient separation of cytokinins from auxin and abscisic acid
and their purification using mixed-mode solid-phase extraction. *Journal of Chromatography A* **950**:
21–29.

Edwards KD, Fernandez-Pozo N, Drake-Stowe K, Humphry M, Evans AD, Bombarely A, Allen F,
Hurst R, White B, Kernodle SP, *et al.* 2017. A reference genome for Nicotiana tabacum enables map-
based cloning of homeologous loci implicated in nitrogen utilization efficiency. *BMC Genomics* **18**:
448.

Escamez S, André D, Sztojka B, Bollhöner B, Hall H, Berthet B, Voß U, Lers A, Maizel A, Andersson
M, *et al.* 2020. Cell Death in Cells Overlying Lateral Root Primordia Facilitates Organ Growth in
Arabidopsis. *Current Biology* **30**: 455-464.e7.

Escamez S, Tuominen H. **2014**. Programmes of cell death and autolysis in tracheary elements: when
a suicidal cell arranges its own corpse removal. *Journal of Experimental Botany* **65**: 1313–1321.

Fahey JW, Zalcmann AT, Talalay P. **2001**. The chemical diversity and distribution of glucosinolates
and isothiocyanates among plants. *Phytochemistry* **56**: 5–51.

Friml J, Wiśniewska J, Benková E, Mendgen K, Palme K. 2002. Lateral relocation of auxin efflux
regulator PIN3 mediates tropism in Arabidopsis. *Nature* **415**: 806–809.

Fukuda H, Komamine A. **1980**. Establishment of an Experimental System for the Study of Tracheary
Element Differentiation from Single Cells Isolated from the Mesophyll of Zinnia elegans. *Plant*
Physiology **65**: 57–60.

Gälweiler L, Guan C, Müller A, Wisman E, Mendgen K, Yephremov A, Palme K. **1998**. Regulation of
polar auxin transport by AtPIN1 in Arabidopsis vascular tissue. *Science (New York, N.Y.)* **282**: 2226–
2230.

Hayashi K, Arai K, Aoi Y, Tanaka Y, Hira H, Guo R, Hu Y, Ge C, Zhao Y, Kasahara H, *et al.* 2021. The
main oxidative inactivation pathway of the plant hormone auxin. *Nature Communications* **12**: 6752.

Hrnčiřík K, Valušek J, Velíšek J. **1998**. A study on the formation and stability of ascorbigen in an
aqueous system. *Food Chemistry* **63**: 349–355.

Kacprzyk J, Burke R, Schwarze J, McCabe PF. **2022**. Plant programmed cell death meets auxin
signalling. *The FEBS Journal* **289**: 1731–1745.

Katz E, Chamovitz DA. **2017**. Wounding of Arabidopsis leaves induces indole-3-carbinol-dependent
autophagy in roots of Arabidopsis thaliana. *The Plant Journal: For Cell and Molecular Biology* **91**: 779–
787.

Katz E, Nisani S, Yadav BS, Woldemariam MG, Shai B, Obolski U, Ehrlich M, Shani E, Jander G,
Chamovitz DA. **2015**. The glucosinolate breakdown product indole-3-carbinol acts as an auxin
antagonist in roots of Arabidopsis thaliana. *The Plant Journal* **82**: 547–555.

Kerchev P, Mühlenbock P, Denecker J, Morreel K, Hoeberichts FA, Van Der Kelen K, Vandorpe M,
Nguyen L, Audenaert D, Van Breusegem F. **2013**. Activation of auxin signalling counteracts
photorespiratory H2O2-dependent cell death. *Plant, Cell & Environment* **38**: 253–265.

Kerk NM, Jiang K, Feldman LJ. **2000**. Auxin Metabolism in the Root Apical Meristem. *Plant Physiology*
122: 925–932.

Knox K, Grierson CS, Leyser O. **2003**. AXR3 and SHY2 interact to regulate root hair development.
Development (Cambridge, England) **130**: 5769–5777.

Kubo M, Udagawa M, Nishikubo N, Horiguchi G, Yamaguchi M, Ito J, Mimura T, Fukuda H, Demura
T. **2005**. Transcription switches for protoxylem and metaxylem vessel formation. *Genes &*
Development **19**: 1855–1860.

Leyser HMO, Pickett FB, Dharmasiri S, Estelle M. **1996**. Mutations in the AXR3 gene of Arabidopsis
result in altered auxin response including ectopic expression from the SAUR-AC1 promoter. *The Plant*
Journal **10**: 403–413.

Love MI, Huber W, Anders S. **2014**. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biology* **15**: 550.

Ludwig-Müller J, Jülke S, Bierfreund NM, Decker EL, Reski R. **2009**. Moss (Physcomitrella patens)
GH3 proteins act in auxin homeostasis. *New Phytologist* **181**: 323–338.

Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. **2019**. PANTHER version 14: More genomes, a
new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids Research* **47**:
D419–D426.

Mitchison GJ. **1980**. A Model for Vein Formation in Higher Plants. *Proceedings of the Royal Society of*
London. Series B, Biological Sciences **207**: 79–109.

Mravec J, Kubeš M, Bielach A, Gaykova V, Petrášek J, Skůpa P, Chand S, Benková E, Zažímalová E,
Friml J. **2008**. Interaction of PIN and PGP transport mechanisms in auxin distribution-dependent
development. *Development* **135**: 3345–3354.

Mravec J, Skůpa P, Bailly A, Hoyerová K, Krecek P, Bielach A, Petrásek J, Zhang J, Gaykova V,
Stierhof Y-D, *et al.* 2009. Subcellular homeostasis of phytohormone auxin is mediated by the ER-
localized PIN5 transporter. *Nature* **459**: 1136–1140.

Müller K, Dobrev PI, Pěnčík A, Hošek P, Vondráková Z, Filepová R, Malínská K, Brunoni F, Helusová
L, Moravec T, *et al.* 2021. DIOXYGENASE FOR AUXIN OXIDATION 1 catalyzes the oxidation of IAA
amino acid conjugates. *Plant Physiology* **187**: 103–115.

Müller K, Hošek P, Laňková M, Vosolsobě S, Malínská K, Čarná M, Fílová M, Dobrev PI, Helusová M,
Hoyerová K, *et al.* **2019**. Transcription of specific auxin efflux and influx carriers drives auxin
homeostasis in tobacco cells. *The Plant Journal: For Cell and Molecular Biology* **100**: 627–640.

Nagata T, Nemoto Y, Hasezawa S. **1992**. Tobacco BY-2 Cell Line as the "HeLa" Cell in the Cell Biology
of Higher Plants. In: Jeon KW, Friedlander M, eds. International Review of Cytology. Academic Press,
1–30.

Nishimura T, Nakano H, Hayashi K, Niwa C, Koshiba T. **2009**. Differential downward stream of auxin
synthesized at the tip has a key role in gravitropic curvature via TIR1/AFBs-mediated auxin signaling
pathways. *Plant & Cell Physiology* **50**: 1874–1885.

Ostrowski M, Świdziński M, Ciarkowska A, Jakubowska A. **2014**. IAA–amido synthetase activity and
GH3 expression during development of pea seedlings. *Acta Physiologiae Plantarum* **36**: 3029–3037.

Ottenschläger I, Wolff P, Wolverton C, Bhalerao RP, Sandberg G, Ishikawa H, Evans M, Palme K.
2003. Gravity-regulated differential auxin transport from columella to lateral root cap cells.
Proceedings of the National Academy of Sciences **100**: 2987–2991.

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. **2017**. Salmon provides fast and bias-aware
quantification of transcript expression. *Nature Methods* **14**: 417–419.

Peñarrubia L, Romero P, Carrió-Seguí A, Andrés-Bordería A, Moreno J, Sanz A. **2015**. Temporal
aspects of copper homeostasis and its crosstalk with hormones. *Frontiers in Plant Science* **6**: 255.

Petrásek J, Cerná A, Schwarzerová K, Elckner M, Morris DA, Zazímalová E. **2003**. Do phytotropins
inhibit auxin efflux by impairing vesicle traffic? *Plant Physiology* **131**: 254–263.

Petrasek J, Mravec J, Bouchard R, Blakeslee JJ, Abas M, Seifertová D, Wisniewska J, Tadele Z, Kubes
M, Covanová M, *et al.* **2006**. PIN proteins perform a rate-limiting function in cellular auxin efflux.
Science (New York, N.Y.) **312**: 914–918.

Petrášek J, Zazímalová E. **2006**. The BY-2 Cell Line as a Tool to Study Auxin Transport. In: Nagata T,
Matsuoka K, Inzé D, eds. Biotechnology in Agriculture and Forestry. Tobacco BY-2 Cells: From Cellular
Dynamics to Omics. Berlin, Heidelberg: Springer Science & Business Media, 107–118.

Petrov V, Hille J, Mueller-Roeber B, Gechev TS. **2015**. ROS-mediated abiotic stress-induced
programmed cell death in plants. *Frontiers in Plant Science* **6**.

Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. **2017**. Differential analysis of RNA-seq
incorporating quantification uncertainty. *Nature Methods* **14**: 687–690.

Rakusová H, Gallego-Bartolomé J, Vanstraelen M, Robert HS, Alabadí D, Blázquez MA, Benková E,
Friml J. **2011**. Polarization of PIN3-dependent auxin transport for hypocotyl gravitropic response in
Arabidopsis thaliana. *The Plant Journal* **67**: 817–826.

Raven JA. **1975**. Transport of indoleacetic acid in plant cells in relation to pH and electrical potential
gradients, and its significance for polar IAA transport. *New Phytologist* **74**: 163–172.

Ravichandran SJ, Linh NM, Scarpella E. **2020**. The canalization hypothesis – challenges and
alternatives. *New Phytologist* **227**: 1051–1059.

Reinhardt D, Pesce E-R, Stieger P, Mandel T, Baltensperger K, Bennett M, Traas J, Friml J,
Kuhlemeier C. **2003**. Regulation of phyllotaxis by polar auxin transport. *Nature* **426**: 255–260.

Rolland-Lagan A-G, Prusinkiewicz P. **2005**. Reviewing models of auxin canalization in the context of
leaf vein pattern formation in Arabidopsis. *The Plant Journal* **44**: 854–865.

Rubery PH, Sheldrake AR. **1974**. Carrier-mediated auxin transport. *Planta* **118**: 101–121.

Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, Eliceiri KW. **2017**. ImageJ2:
ImageJ for the next generation of scientific image data. *BMC Bioinformatics* **18**: 529.

Ruzicka K, Ursache R, Hejátko J, Helariutta Y. **2015**. Xylem development – from the cradle to the
grave. *New Phytologist* **207**: 519–535.

Sachs T. **1991**. Cell polarity and tissue patterning in plants. *Development* **113**: 83–93.

Scheuring D, Schöller M, Kleine-Vehn J, Löfke C. **2015**. Vacuolar Staining Methods in Plant Cells. In:
Estevez JM, ed. Methods in Molecular Biology. Plant Cell Expansion: Methods and Protocols. New
York, NY: Springer, 83–92.

Seifertová D, Skůpa P, Rychtář J, Laňková M, Pařezová M, Dobrev PI, Hoyerová K, Petrášek J,
Zažímalová E. **2014**. Characterization of transmembrane auxin transport in Arabidopsis suspension-
cultured cells. *Journal of Plant Physiology* **171**: 429–437.

Sheldrake AR. **2021**. The production of auxin by dying cells. *Journal of Experimental Botany* **72**:
2288–2300.

Simon S, Kubeš M, Baster P, Robert S, Dobrev PI, Friml J, Petrášek J, Zažímalová E. **2013**. Defining
the selectivity of processes along the auxin response chain: a study using auxin analogues. *New*
Phytologist **200**: 1034–1048.

Smetana O, Široký J, Houlné G, Opatrný Z, Chabouté M-E. **2012**. Non-apoptotic programmed cell
death with paraptotic-like features in bleomycin-treated plant cells is suppressed by inhibition of
ATM/ATR pathways or NtE2F overexpression. *Journal of Experimental Botany* **63**: 2631–2644.

Song L, Florea L. **2015**. Rcorrector: efficient and accurate error correction for Illumina RNA-seq reads.
GigaScience **4**: s13742-015-0089-y.

Su N, Zhu A, Tao X, Ding ZJ, Chang S, Ye F, Zhang Y, Zhao C, Chen Q, Wang J, *et al.* 2022. Structures
and mechanisms of the Arabidopsis auxin transporter PIN3. *Nature* **609**: 616–621.

Sun L, Dong S, Ge Y, Fonseca JP, Robinson ZT, Mysore KS, Mehta P. **2019**. DiVenn: An Interactive and
Integrated Web-Based Visualization Tool for Comparing Gene Lists. *Frontiers in Genetics* **10**: 421.

Swarup K, Benková E, Swarup R, Casimiro I, Péret B, Yang Y, Parry G, Nielsen E, De Smet I, Vanneste
S, *et al.* 2008. The auxin influx carrier LAX3 promotes lateral root emergence. *Nature Cell Biology* **10**:
946–954.

Tognetti VB, Mühlenbock P, Van Breusegem F. **2012**. Stress homeostasis – the redox and auxin
perspective. *Plant, Cell & Environment* **35**: 321–333.

Tripathi AK, Pareek A, Singla-Pareek SL. **2017**. TUNEL Assay to Assess Extent of DNA Fragmentation
and Programmed Cell Death in Root Cells under Various Stress Conditions. *Bio-Protocol* **7**: e2502.

Ung KL, Schulz L, Kleine-Vehn J, Pedersen BP, Hammes UZ. **2023a**. Auxin transport at the
endoplasmic reticulum: roles and structural similarity of PIN-FORMED and PIN-LIKES. *Journal of*
Experimental Botany: erad192.

Ung KL, Schulz L, Stokes DL, Hammes UZ, Pedersen BP. **2023b**. Substrate recognition and transport
mechanism of the PIN-FORMED auxin exporters. *Trends in Biochemical Sciences*.

Ung KL, Winkler M, Schulz L, Kolb M, Janacek DP, Dedic E, Stokes DL, Hammes UZ, Pedersen BP.
2022. Structures and mechanism of the plant PIN-FORMED auxin transporter. *Nature* **609**: 605–610.

Valandro F, Menguer PK, Cabreira-Cagliari C, Margis-Pinheiro M, Cagliari A. **2020**. Programmed cell
death (PCD) control in plants: New insights from the Arabidopsis thaliana deathosome. *Plant Science*
299: 110603.

Van Durme M, Nowack MK. **2016**. Mechanisms of developmentally controlled cell death in plants.
Current Opinion in Plant Biology **29**: 29–37.

Verna C, Sawchuk MG, Linh NM, Scarpella E. **2015**. Control of vein network topology by auxin
transport. *BMC Biology* **13**: 94.

van Wees S. **2008**. Phenotypic analysis of Arabidopsis mutants: trypan blue stain for fungi,
oomycetes, and dead plant cells. *CSH protocols* **2008**: pdb.prot4982.

Weijers D, Ljung K, Leyser O. **2021**. *Auxin Signaling: From Synthesis to Systems Biology, Second*
Edition. Cold Spring Harbor Laboratory Press.

Winicur ZM, Zhang GF, Staehelin LA. **1998**. Auxin deprivation induces synchronous Golgi
differentiation in suspension-cultured tobacco BY-2 cells. *Plant Physiology* **117**: 501–513.

Xuan W, Audenaert D, Parizot B, Möller BK, Njo MF, De Rybel B, De Rop G, Van Isterdael G,
Mähönen AP, Vanneste S, *et al.* 2015. Root Cap-Derived Auxin Pre-patterns the Longitudinal Axis of
the Arabidopsis Root. *Current Biology* **25**: 1381–1388.

Xuan W, Band LR, Kumpf RP, Damme DV, Parizot B, Rop GD, Opdenacker D, Möller BK, Skorzinski
N, Njo MF, *et al.* 2016. Cyclic programmed cell death stimulates hormone signaling and root
development in Arabidopsis. *Science* **351**: 384–387.

Yang Z, Xia J, Hong J, Zhang C, Wei H, Ying W, Sun C, Sun L, Mao Y, Gao Y, *et al.* 2022. Structural
insights into auxin recognition and efflux by Arabidopsis PIN1. *Nature* **609**: 611–615.

Zou W, Lin P, Zhao Z, Wang D, Qin L, Xu F, Su Y, Wu Q, Que Y. **2022**. Genome-Wide Identification of
Auxin-Responsive GH3 Gene Family in Saccharum and the Expression of ScGH3-1 in Stress Response.
International Journal of Molecular Sciences **23**: 12750.

Zuo J, Niu QW, Chua NH. **2000**. Technical advance: An estrogen receptor-based transactivator XVE
mediates highly inducible gene expression in transgenic plants. *The Plant Journal: For Cell and*
Molecular Biology **24**: 265–273.

 FIGURE LEGENDS

Figure 1. The conditional expression of *XVE::AtPIN5* in BY-2 cells leads to the auxin starvation

phenotypes. **A** to **C)** The induction of the *XVE::AtPIN7* transgene results in increased cell

elongation **A)**, plotted on **B)** (*P* < 0.001 by Student's *t*-test), and decreased growth, monitored

by the cell density in the media **C)**. **D** to **F)** The induction of *XVE::Ataxr3-1* phenocopies the

symptoms of auxin starvation documented by representative image on **D)**, quantification of

increased cell elongation (*P* < 0.001 by Student's *t*-test) **E)** and reduced proliferation **F)**. **G** to **I)**

The induction of *XVE::AtPIN5* in the cell line cultured in a standard medium with 2,4-D leads

to increased cell elongation **G)**, plotted on **H)** (*P* < 0.05 by Student's *t*-test), and reduced

proliferation rate **I)**. On the dot plots, the open symbols are averages ± S.E. *******P* < 0.01, ********P* <

0.001 by Student's *t*-test. Scale bars: 50 µm.

Figure 2. Replacing 2,4-D with its native analog IAA enhances the auxin starvation effects

brought about by the induction of *XVE::AtPIN5*. **A)** The comparison of the cell proportions of

the control *XVE::AtPIN5* cells cultivated in the medium supplemented with 2,4-D or IAA reveals

the increase of the elongation growth in the presence of IAA (*P* < 0.001 by Student's *t*-test). **B**

to **C)** The proliferation rate of cells containing the induced *XVE::AtPIN5* transgene is even lower

in the medium with IAA added at the beginning of subcultivation, the representative image

on **B)**, the corresponding quantification of cell density on **C)**. **D** and **E)** The phenotype of the

XVE::AtPIN5 cells cultured in a medium continuously supplemented with IAA every 12 hours

during the monitored cultivation period leads to the partial reversion of the starvation

phenotypes in the control lines, however, the induction of the *XVE::AtPIN5* transgene leads to

strong auxin starvation phenotypes, as documented at the representative image on **D)** and by

the quantification of the cell growth on **E)**. **F** and **G)** The induction of *XVE::AtPIN5* results in

higher uptake of [^{3}H]-IAA **F)**, but shows no effect on uptake of [^{3}H]-2,4-D **G)**. Scale bars: 50 µm.

On the plots, the open symbols are averages ± S.E. *******P* < 0.01, ********P* < 0.001 by Student's *t*-

test.

Figure 3. An increased incidence of symptoms of PCD following induction of *XVE::AtPIN5*. **A**

and **B)** The replacement of 2,4-D with IAA leads to more frequent cytoplasm shrinkage and

hereafter increased accumulation of the trypan blue dye inside cells (arrowheads), where

XVE::AtPIN5 was induced, as documented by representative images on **A)** and the inverse

frequency of the trypan-blue intracellular staining on **B)**. **C)** DiVenn diagram indicating that a

large part of differentially expressed genes following the *XVE::AtPIN5* induction is shared with

those from the cultures undergoing auxin starvation; red, significantly upregulated genes;

blue, downregulated genes; yellow, opposite expression changes. A single dot denotes a group

of 10 genes. **D** and **E)** An increased fluorescence intensity on confocal images of the H2DCFDA

staining of *XVE::AtPIN5* lines detects an elevated presence of ROS following the transgene

induction **D)**, as quantified in **E)**. **F** and **G)** Confocal images of cells stained by BCECF-AM to

mark acidic areas **F)**, as quantified in **G)**, document that the *XVE::AtPIN5* induction leads to

decreased acidity in vacuoles but increased acidity in the cytoplasm and nucleus area

(arrowheads). **H** and **I)** Representative images of TUNEL staining, marking the disintegration

of nuclear DNA (red channel: TUNEL, blue channel: nuclear DAPI staining) **H)**, shows an

enhanced labeling frequency in the lines where *XVE::AtPIN5* was induced **I)**. For **D)** and **F)**, the

color intensity is indicated by the rainbow color lookup table (inset on **D)**). Scale bars: 50 μm.

On the dot plots, the open symbols are averages ± S.E. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ by

Student's *t*-test.

Figure 4. Inducible *XVE::AtPIN5* expression in BY-2 cells promotes IAA deactivation by the

decarboxylation pathway. **A)** A scheme of current understanding of main IAA degradation

pathways (products of IAA oxidation pathway in blue, IAA decarboxylation in red). **B)** LC-MS

quantification of IAA and its metabolites in the cells expressing *XVE::AtPIN5* cultured on the

standard 2,4-D-supplemented media shows a moderate decrease of IAA and its most

degradation products. **C)** The induction of *XVE::AtPIN5* in the cells cultivated in media with IAA

added every 12 hours leads to a dramatic drop of IAA and its conjugates, while the compounds

belonging to the IAA decarboxylation pathway show elevated levels. On **B)** and **C)**, the open

symbols indicate the mean from three independent measurements; the bars are ± S.E. *$P <$

0.05, **$P < 0.01$, ***$P < 0.001$, by Student's *t*-test, n. s.: not significant at $\alpha_{0.05}$. IAA: indole-3-

acetic acid, IAA-Asp: IAA-aspartate, IAA-GE: IAA-glucose ester, OxIAA-Asp: Oxo-IAA-aspartate,

Abg: ascorbigen, DiHAbg: dihydroascorbigen, I3M-SG: indole-3-methylglutathione. The cells

were harvested 2 hours after the addition of 1 μM IAA.

Figure 5. The I3C treatment leads to growth arrest and reduced viability of BY-2 cells. **A)** The

addition of 50 μM I3C to the media leads to increased elongation of BY-2 cells ($P < 0.001$ by

Student's *t*-test) and **B)** decreased proliferation, monitored by the cell density in the media. **C**

to **F)** The I3C treatment also results in increased cell mortality monitored by trypan blue **C** and

D) and TUNEL staining (red channel: TUNEL, blue channel: nuclear DAPI staining) **E** and **F)**.

Scale bars: 50 µm. On the dot plots, the open symbols are averages ± S.E. ***$P < 0.001$ by

Student's *t*-test.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Additional aspects of auxin deficiency of tobacco BY-2 cells. **A)** The induction of the

GVG::AtPIN7 transgene leads to decreased cell density, reflecting their lower proliferation rate

during the whole subcultivation interval. **B)** The cell density of the wild-type BY-2 cells

decreases with the growing concentration of supplied PEO-IAA, an inhibitor of auxin-

dependent gene expression ($F_{(5,48)} = [11.112]$, $P < 0.001$ by one-way ANOVA). **C)** The enhanced

cell expansion is typical for cultivation on the auxin-free medium (shown on the *GVG::AtPIN7*

line without the chemical inducer). **D to F)** The conditional expression of *GVG::AtPIN5* in the

cell lines cultured in standard medium with 2,4-D leads to increased cell expansion **D** and **E)**

($P < 0.001$ by Student's t-test) and reduced proliferation rates **F)**. **G to H)** The induction of

XVE::AtPIN5-GFP in the cells co-expressing auxin *DR5::RFP* reporter results in a reduction of

the fluorescence in cells in the upper-intensity quantiles **G)**, and PIN5-GFP colocalizes with the

ER-targeted RFP **H)**. On **A)**, **B)** and **F)**, the open symbols are averages ± S.E. On **G)**, the vertical

error bars represent median ± 95% confidence intervals. Scale bars: 50 µm on **C)**, **D)** and **H)**.

Figure S2. Increased frequency of symptoms of cell death and its relationship to auxin

starvation following induction of *AtPIN5*. **A and B)** The representative images of incorporation

of the trypan blue inside the cells following induction of *GVG::AtPIN5* **A)**, quantified on **B)**. **C**

and **D)** Quality control of RNA-Seq data processing, a heat-map representing Jensen-Shannon

divergence between individual samples (sorted by internal sample IDs) **C)**, projections of

individual samples onto the principal components (PC) **D)**. **E)** An increased fluorescence on

confocal images of the *GVG::AtPIN5* cells stained by 10 µM H2DCFDA to detect ROS, following

the transgene induction. **F)** The pattern and intensity of the *At*PIN5-GFP foci are comparable

in both standard 2,4-D-containing medium and that continuously supplemented with IAA

every 12 hours. The signal intensity is indicated by the green fire blue rainbow lookup table

on **F)**. On **B)**, the dot plot of a data set of 2000 cells in total, the open symbols are averages ±

S.E. , ***$P < 0.001$ by Student's *t*-test. Scale bars: 25 µm on **A)**, 100 µm on **E)**, and 10 µm on **F)**.

Figure S3. Detection of the DNA fragmentation during auxin deficiency. **A)** TUNEL staining (left: TUNEL, middle: control nuclear DAPI staining, right: merged false-colored images) marks an enhanced disintegration of nuclear DNA in the cells where *GVG::AtPIN5* was induced. **B** to **C)** Negative TUNEL stainings in *GVG::AtPIN7* and *XVE::Ataxr3-1* cells evidence the absence of the PCD, both under the standard conditions **B)**, or when supplemented with IAA every 12 hours during the monitored cultivation period **C)**. Upper panels: control nuclear DAPI staining, lower panels: TUNEL. Ctrl (+): control cells additionally treated with DNAse I to detect the positive TUNEL reaction, Ctrl (-): staining where the TdT enzyme action was omitted from the control reaction. Scale bars: 50 µm.

Figure S4. WT BY-2 cells cultured in a medium containing cell debris exhibit neither decrease of proliferation activity **A)** nor signs of decreased viability monitored by trypan blue staining **B)**. Scale bar: 100 µm on **B)**.

Figure S5. Additional experiments examining the activity of the IAA decarboxylation pathway following the *AtPIN5* expression. **A)** Representative LC-MS analysis of IAA and its metabolites in cells expressing *XVE::AtPIN5* determined immediately after 1 µM IAA supply (t_0). While the exogenously applied IAA is up-taken immediately, the levels of its metabolites are still barely detectable. **B)** LC-MS analysis of IAA and its metabolites in cells expressing *GVG::AtPIN5* cultured on the standard 2,4-D-supplemented media, 2 hours after adding 1 µM IAA. Open symbols indicate the mean of three technical measurements; the bars are ± S.E. **$P < 0.01$, ***$P < 0.001$ by Student's *t*-test, n. s.: not significant at $\alpha_{0.05}$. IAA: indole-3-acetic acid, IAA-Asp: IAA-aspartate, IAA-GE: IAA-glucose ester, OxIAA-Asp: Oxo-IAA-aspartate, Abg: ascorbigen, DiHAbg: dihydroascorbigen.

Table S1. Results of PANTHER Overrepresentation test (GO - Molecular function) of transcripts significantly upregulated in both auxin-starved and the *XVE::AtPIN5*-expressing cells.

777   **Table S2.** Results of PANTHER Overrepresentation test (GO - Molecular function) of transcripts

778   significantly upregulated in BY-2 cells carrying induced *XVE::AtPIN5*, compared to untreated

779   controls and auxin-starved cells.

A

B

C

A

B

	Ctrl (+)	Ctrl (−)	*AtPIN7* Nonind.	*AtPIN7* Ind.	*Ataxr3-1* Nonind.	*Ataxr3-1* Ind.
DAPI						
TUNEL						

C

DAPI IAA/12						
TUNEL IAA/12						

A

B

A IAA metabolites (standard medium) at t_0

B IAA metabolites (standard medium)