1 **Promoted extension of an epidermal cell by semaphorin in *C. elegans* revealed by a mosaic**

2 **gene expression analysis by using the IR-LEGO single-cell gene induction system**

4 Motoshi Suzuki[1] and Shin Takagi

9 Division of Biological Science, Graduate School of Science, Nagoya University,

10 1: To whom correspondence should be addressed

12 Short title: Extension of a *C. elegans* epidermal cell by semaphorin

13 Keywords: Semaphorin signaling /epidermis/ *C. elegans/* IR-LEGO gene induction system

Abstract

Semaphorin signaling regulates the cell shape in context-dependent manners. In the nematode *C. elegans*, membrane-bound semaphorins SMP-1/2 (SMPs) regulate the morphology of epidermal cells via their receptor plexin, PLX-1. In the larval male tail of the SMPs/PLX-1 signaling mutants, the border between two epidermal cells, R1.p and R2.p, is displaced anteriorly, resulting in the anterior displacement of the anterior most ray, ray 1, in the adult male. To elucidate how intercellular signaling mediated by SMPs defines the position of the cell border, we performed mosaic gene expression analyses by using IR-LEGO (InfraRed Laser Evoked Gene Operator). We show that PLX-1 expressed in R1.p and SMP-1 expressed in R2.p during a defined period of ray development is required for proper positioning of ray 1. The result suggests that in a ray-lineage epidermal cell, SMPs signaling promotes extension, rather than retraction, of the PLX-1-expressing cell.

1 INTRODUCTION

Semaphorin is a major group of signaling molecules of the animal conserved both in

vertebrates and invertebrates (Jongbloets and Pasterkamp, 2014). Semaphorins and the

receptor plexins are involved in regulation of diverse developmental and physiological

processes including neural circuit formation, vascular morphogenesis, bone development,

tumorigenesis and regulation of immuno-response (Tran et al., 2007; Pasterkamp, 2012;

Iragavarapu-Charyulu et al., 2020; Kang and Kumanogoh, 2013; Tamagnone, 2012;

Kumanogoh and Kikutani, 2013). In axon guidance, semaphorin was first identified as a

repellent for the growth cone, but later studies showed that the response of the growth cone to

semaphorin is context-dependent and may turn to attraction depending on factors internal

and external to the cell, such as second messengers, the FAK/src signaling, CNG channel or

proteoglycans (Song et al., 1998; Falk et al., 2005; Togashi et al., 2008; Kantor et al., 2004).

A nematode *Caenorhabditis elegans (C. elegans)* has three semaphorins and two plexins. In

larvae, Class I transmembrane-type semaphorins (SMP-1 and SMP-2: SMPs) bind to PlexinA

(PLX-1) to regulate the shape of epidermal cells that generate adult organs such as the vulva

in hermaphrodites and rays, sensory structures, in the male tail (Fujii et al., 2002; Ginzburg

et al.,2002; Dalpé et al., 2004, 2005; Liu et al., 2005). Defective SMPs signaling affects the

arrangement of larval epidermal cells. In vulva precursor cells (VPCs) of the SMPs-PLX-1

signaling mutants, VPCs failed to terminate the longitudinal extension, resulting in the

overshoot and lateral overlaps among VPCs (Liu et al., 2005). In the larval male tail, the

defective SMPs signaling, mainly mediated by SMP-1, causes an anterior shift in the position

of the boundary between a pair of epidermal cells called R1.p and R2.p, which eventually

results in anterior displacement of the anterior-most ray, ray 1, in the adult (Fujii et al., 2002).

While the SMPs-PLX-1 system appears to mediate a stop signal for cell extension in VPCs (Liu et al., 2005; Suzuki et al., 2022), how the semaphorin-plexin signal is transmitted among epidermal cells of the larval male tail and regulates their behavior is poorly understood. The receptor PLX-1 is expressed in all epidermal ray precursor cells, R(n)s, and their descendants including R1.p and R2.p (Fujii et al., 2002). Thus, although it is plausible that SMPs signaling between R1.p and R2.p keeps the boundary between them at a proper position, it is not clear how and in which of these two adjacent cells the function of SMPs and PLX-1 molecule is required: the anterior shift of the boundary between R1.p and R2.p observed in the SMPs-PLX-1 signaling mutants can be explained either by the failure of retraction of the anterior part of the posterior cell, R2.p, or the failure of extension of the posterior part of the anterior cell, R1.p. While in these simple models SMPs, transmembrane-type semaphorins, and PLX-1 are assumed to interact in a cell-to-cell contact-mediated manner, this, however, has not been tested experimentally; a previous study suggested that SMPs expressed in the hook primordium, which is located on the ventral midline distant from R1.p and R2.p, determine the position of ray 1 (Dalpé et al., 2004).

In order to address this issue, it would be useful to induce expression of PLX-1 and/or SMPs, SMP-1 in particular, in a single R(n) or its single descendant cell of each mutant, and examine the ray 1 phenotype. Whereas tissue-specific promoters, such as the promoter of *mab-5*, *egl-5*, *lin-32* and *efn-4* (Salser and Kenyon, 1996; Ferreira et al., 1999; Portman and Emmons, 2000; Hahn and Emmons, 2003) could be used to force the expression of the transgenes in R(n)s and their descendant cells, the expression is not restricted to a single cell and the timing of gene expression cannot be regulated as we desire. To express PLX-1 and

SMP-1 in a targeted single cell at a desired developmental stage, here we used IR-LEGO

(InfraRed Laser Evoked Gene Operator) (Kamei et al., 2009; Suzuki et al., 2013). Our mosaic

gene expression analysis indicates that expression of PLX-1 in R1.p and expression of SMP-1

in R2.a or R2.p at a certain period of development is important for proper positioning of the

boundary between R1.p and R2.p, suggesting that the SMPs signal promotes the extension of

a PLX-1-expressing R1.p.

MATERIALS AND METHODS

C. elegans strains

Standard techniques for *C. elegans* culture and genetics were used as described by Brenner

(1974).

The following strains were used.

N2, [LGI] *smp-1(ev715)*, *smp-2(ev709)*; [LGIV] *plx-1(nc37)*, *him-8(e1489)* [LGV] *him-5(e1490)*.

The transgenes used in this study include

jcIs1[ajm-1::gfp; rol-6](su1006)] (Mohler et al., 1998) ,

ncIs13[ajm-1::gfp] (Liu et al., 2005) ,

ncIs17[hsp::gfp] (Kamei et al., 2009),

ncIs19[hsp::plx-1; hsp::gfp; rol-6(su1006)] (Suzuki et al., 2022),

ncIs23[hsp::smp-1; hsp::gfp; rol-6(su1006)] (Suzuki et al., 2022),

ncEx2030[lin-32p::plx-1; rol-6(su1006)] (Nukazuka et al, 2008).

ncIs24[hsp::smp-1; hsp::gfp; rol-6(su1006)] was generated similarly to *ncIs23* (Suzuki

et al., 2022).

ncIs211[hsp::gfp] was generated similarly to *ncIs17* (Kamei et al., 2009).

ncIs210[hsp::plx-1;hsp::smp-1; hsp::gfp; rol-6(su1006)] is a chromosomally integrated

transgene generated by the gamma-ray irradiation (Shioi et al., 2001) to *ncEx2016[hsp::plx-1;*

hsp::smp-1; hsp::gfp; rol-6(su1006)] worms (Suzuki et al., 2022).

To generate the rescue construct *lin-32p::smp-1,* a *smp-1 cDNA* was cloned in the *pPD49.26*

vector (provided by A. Fire) carrying the *lin-32* promoter *(lin-32p).* The plasmid was mixed

with *pRF4* in the ratio of 1:2 (in weight) and was used to generate *ncEx2038[lin-32p::smp-1;*

rol-6(su1006)] by microinjection (Mello and Fire, 1995).

For analysis of male tails, the strains carried the *him-5* or *him-8* mutation. The strains used

for the IR-irradiation experiments contained *ajm-1::gfp* for visualizing the ray-lineage cells.

Some stains also carried *rol-6(su1006),* which confers the Roller phenotype, as a transgene to

facilitate the observation of rays.

NW1358 *smp-1(ev715)* and NW1335 *smp-2(ev709)* were provided by Joe Culotti (Ginzburg et

al., 2002) and SU93 *jcIs1[ajm-1::gfp; rol-6(su1006)]* were provided by Jeff Simske (Mohler et

al., 1998).

IR-laser irradiation

Live males at the mid-L3 stage were placed on a 6% agar pad mounted with 5% levamisole for

immobilization, and covered with a coverslip with a thickness of approximately 0.2 mm

(Matsunami, Japan). The IR-LEGO system (Olympus IX-70 equipped with the IR-laser set)

was described in the previous papers (Kamei et al., 2008; Suzuki et al., 2022). Specimens were

observed with a 100 X mono-layer coated objective lens (UplanApo 100 X NA 1.3, Olympus,

Japan) immersed in oil (Immersion oil type-F, Olympus, Japan). After irradiating worms, the

coverslips were immediately removed and worms were recovered on a NGM plate. Three

hours after irradiation, we checked GFP expression in the targeted cell and then observed

rays one day after irradiation. In typical experiments, ray-lineage cells at stage 2 were

irradiated with IR-laser beam at 13 mW for 0.25 sec a total of 4 times. About 15% of

irradiated R(n).as and R(n).ps were induced to express GFP. We used the condition at 11 mW

for 1 sec to irradiate R(n)s at stage 1, and R1+2.p at stage 3. About 30% of irradiated cells

expressed GFP. Irradiated cells were sometimes damaged and failed to generate rays.

RESULTS

Background of the experiments: Development of rays and SMPs-PLX-1 signaling

Rays are sensory organs in the adult male tail, and play an important role during copulation

(Figure 1(a), 3(a)). For readers who are not familiar with this structure, the development of

epidermal cells in the male tail will be briefly summarized. Rays are generated from a row of

epidermal cells called seam cells on the lateral side of the larval body (Emmons, 2005). Three

bilateral pairs of seam cells, V5, V6 and T, in the posterior part of a male body produce the

nine pairs of ray precursor cells, R1, R2-R6, R7-R9, at the third larval (L3) stage, respectively.

(Figure 2(a)). A R(n) divides into two daughter cells, R(n).a and R(n).p (Figure 2(b), (i)), and

then R(n).a generates three cells, which form a ray precursor cluster (Figure 2(c), (d), (e)) that

eventually produces a single ray (n) (n=1 - 9). Meanwhile, R(n).ps, after shifting their position

dorso-laterally to R(n).as, become enlarged, and fuse to each other. The cell-fusion occurs first

between R1.p and R2.p (Figure 2(f), (g)), and then proceed posteriorly to eventually form a

syncytial tail seam (set) (Figure 2(h)) (Sulston et al., 1980; Emmons, 2005). In this paper, for

the convenience of description, we designate R(n)s and their descendants collectively as

ray-lineage cells, and define the developmental stages as follows: stage 1: the stage prior to

the cell division of R(n) (Figure 2(a)); stage 2: the stage after the cell division of R(n) and prior

to the cell-fusion of R1.p and R2.p (Figure 2(b)-(f)); stage 3: the stage after the cell-fusion of

R1.p and R2.p (Figure 2(g)). The cell generated by the fusion of R1.p and R2.p is referred to as

R1+2.p in this paper (Figure 2(g)).

In the majority of wild-type animals, the anterior most ray 1 is juxtaposed to the

neighboring ray 2 (Figure 1(a), 3(a)), which we define as "Level 1" for the position of ray 1.

Since the position of ray precursor cluster 1 in *plx-1* and *smp-1 smp-2* mutants shifts

anteriorly compared to that in wild-type animals at the L4 stage (Figure 2(g)) (Fujii et al.,

2002; Nukazuka et al., 2008), the mutant adult males show the highly penetrant defect in

which the position of ray 1 is displaced anteriorly. In many cases, ray 1 is either located

outside of the fan (Level 3) (Figure 1(c), the right side of the tail in Figure 3(b)) or inside of the

fan but is separated from ray 2 (Level 2) (Figure 1(b), the left side of the tail in Figure 3(b)).

PLX-1 is expressed in the all ray-lineage cells (Fujii et al., 2002) and the defective positioning

of ray 1 in *plx-1* mutants was rescued by expression of a PLX-1 cDNA under the control of the

lin-32 promoter (Figure 1(d)), which drives expression of transgenes in the all ray-lineage

cells (Portman and Emmons, 2000), as reported previously (Nakao et al., 2007; Nukazuka et

al., 2008). Similarly, the defective positioning of ray 1 in *smp-1 smp-2* mutants was rescued by

expression of a SMP-1 cDNA under the control of the *lin-32* promoter with the transgene

ncEx2038[lin-32p::smp-1; rol-6(su1006)] (Figure 1(d)).

While these rescue experiments indicate that SMPs-PLX-1 signaling among ray-lineage cells

determines the proper position of ray 1, the cells that send and/or receive the signal remain to

be specified. To examine further where and when the SMPs signal is required, we induced to

express SMP-1 and PLX-1 in a single cell of each mutant animal by using IR-LEGO. For

induction of PLX-1 and SMP-1, we used the transgenes, *ncIs19[hsp::plx-1; hsp::gfp]* and

ncIs24[hsp::smp-1; hsp::gfp], which contains *hsp::plx-1* and *hsp::smp-1*, respectively, and

hsp::gfp as an induction marker. In this experiment, we irradiated a ray-lineage cell at

different developmental stages (Figure 3(c)) and checked the GFP expression 3 hours after

irradiation (Figure 3(d)). Then, one day after irradiation we scored the position of ray 1

(Figure 3(e)).

Induction of PLX-1 in R1.p rescued the ray 1 defect of *plx-1* mutants

First we tried to express PLX-1 in R1 (a V5 descendant) of the *plx-1; ncIs19[hsp::plx-1;*

hsp::gfp] worm at stage 1. In worms that successfully expressed PLX-1 in R1, the proportion

of each level of the ray 1 position did not differ significantly from that in *plx-1* worms (n=31.

Level 1: 20%, Level 3: 42%) (Figure 3(f)), indicating that the ray 1 displacement defect was not

rescued. Expression of PLX-1 in the other R(n)s at stage 1 also did not rescue the ray 1 defect

of *plx-1* mutants (Level 1: 6% n=35, 4% n=26, 7% n=14 for induction in R2, R3 and R4

respectively) (Figure 3(f)).

Then worms were irradiated at stage 2, when R(n) had already completed cell division to

give rise to R(n).a and R(n).p. Induction of PLX-1 expression in R1.p of *plx-1* mutants rescued

the ray 1 defects significantly: the proportion of Level 1, which represents normal positioning

(Figure 3(e)), increased to more than 60% (n=38, p<0.01) while that of Level 3, which

represents severe anterior displacement, decreased to less than 10% (Figure 3(f)). In contrast,

induction of PLX-1 expression in the other cells at stage 2 did not lead to significant rescue of

the ray1 defects. Induction of GFP alone, which serves as a control of IR-irradiation, in R1.p of

plx-1 mutants did not affect the position of ray 1 (Figure S1).

When we irradiated R1.p or R2.p at stage 2, we occasionally found the GFP expression in

R1+2.p as a result of fusion of R1.p with R2.p by 3 hours after irradiation. Such cases will be

referred to induction at "stage 2/3" in this paper. Induction of PLX-1 expression in R1.p at

stage 2/3 also rescued the ray 1 defect strongly (Level 1: 75% n=20, p<0.01) (Figure 3(f)). On

the other hands, induction in R2.p at stage 2/3 did not rescue the ray 1 defect significantly

(n=20). When worms were irradiated at stage 3, induction of PLX-1 expression in R1+2.p or

other cells (data not show) did not rescue the ray 1 defect.

These results indicate that function of PLX-1 in R1.p at stage 2 or 2/3 is required for

proper positioning of ray 1 (Figure 3(g)).

Induction of SMP-1 in R1.p, R2.a or R2.p rescued the ray 1 defect of *smp-1 smp-2* mutants

Next, we tried to examine the functional site of the semaphorin ligand involved in positioning

of ray 1. While two membrane-bound semaphorins, SMP-1 and SMP-2, both bind to PLX-1,

and are involved in regulation of the positioning of ray 1 (Fujii et al., 2002), SMP-1 has a

dominant role; In *smp-2* single mutants the ray 1 displacement defect is negligible, and

expression of SMP-1 cDNA under the control of the *lin-32* ray-lineage specific promoter is

sufficient for rescuing the ray 1 defect of *smp-1 smp-2* double mutants nearly completely

(Figure 1(d)). Therefore, we focused on SMP-1 and induced it in a single ray-lineage cell in

smp-1 smp-2; ncIs24[hsp::smp-1; hsp::gfp] worms by using IR-LEGO (Figure 4(a)). The

induced expression of SMP-1 in ray-lineage cells at stage 1 or stage 3 did not rescue the

phenotype of ray 1 displacement. Expression of SMP-1 in a single R1.p, R2.a or R2.p at stage

2 rescued the ray 1 defect (Level 1: 50% n=34, 53% n=15, and 53% n=19, for induction in R1.p,

R2.a or R2.p, respectively. p<0.01). Induction of GFP alone in these cell of *smp-1 smp-2*

mutants did not affect the position of ray 1 significantly (Figure S2), confirming that

expression of SMP-1 in either R1.p, R2.a or R2.p is required for proper positioning of ray 1

(Figure 4(b)). In another control experiment, induced expression of PLX-1 in R1.p of *smp-1*

smp-2 mutants did not affect the ray 1 phenotype (Level 1: 4% n=69).

On the other hand, expression of SMP-1 in R1.a or daughter cells of R3 at stage 2 did not

rescue the ray 1 displacement defect (Level 1: 0% n=20 and 5% n=21 for induction in R1.a and

R3d, respectively). In the case of induction at stage 2/3, proportion of Level 1 appeared to be

reduced by expressing SMP-1 in either R1.p or R2.p, (Level 1: 36% n=14 for R1.p; 23% n=22

for R2.p), though the results are considered to be statistically non-significant.

As shown in the preceding section, expression of PLX-1 in R1.p appears critical for proper

positioning of ray 1. Both R2.a and R2.p, in which induction of SMP-1 rescued the ray 1 defect,

are positioned posterior to and in contact with R1.p during a certain period of the

development of the male tail (Figure 6(c), (d)). In contrast to this, induced expression of

SMP-1 in R1.a, which is in contact with but positioned anterior to R1.p, failed to restore

proper positioning of ray 1. These results suggest that activation of SMPs-PLX-1 signaling in

the posterior part of R1.p via the adjacent cells is critical for proper positioning of ray 1,

presumably by shifting the position of the boundary between R1.p and R2.p posteriorly.

Therefor we propose a model, in which SMP-1 on R2.a or R2.p promotes or permits the

extension of the contacting edge of the neighboring R1.p (Figure 6(c), (d), (f)).

Co-induction of SMP-1 and PLX-1 in R1.p rescued the ray 1 defect of *smp-1 smp-2; plx-1*

mutants partly

The model described above, however, does not explain another finding that induced SMP-1 in

R1.p also rescued the ray 1 defect. We think that this apparent contradiction could be

resolved by assuming the *cis*-interaction between SMP-1 and PLX-1. Our previous study

showed that SMP-1, which was overexpressed in a VPC by using IR-LEGO, interacts with

co-expressed PLX-1 in *cis* and triggers the signal downstream to PLX-1 (Suzuki et al., 2022).

Similarly, the present finding that SMP-1 expression in R1.p rescues the ray 1 defect in *smp-1*

mutants might be explained if induced SMP-1 *cis*-activates PLX-1 on the same R1.p cell. To

examine this possibility, we induced the expression of both SMP-1 and PLX-1 in a single

ray-lineage cell in *smp-1 smp-2; plx-1* triple mutants background by using *smp-1 smp-2; plx-1;*

ncIs210[hsp::plx-1; hsp::smp-1; hsp::gfp; rol-6(su1006)] worms. We focused our analysis on the

effect of induction in R1.a, R1.p, R2.a or R2.p at stage 2. We found that IR-irradiation to R1.p

rescued the ray 1 displacement defect in 22% of the *smp-1 smp-2; plx-1* mutant worms (n=22

p<0.05.). IR-irradiation to the other cells did not affect the ray 1 phenotype (Level 1: less than

5% n=54) (Figure 5(a)). These data are consistent with the idea that SMP-1 induced by

IR-LEGO *cis*-activates PLX-1 in R1.p (Figure 6(e), (g)).

DISCUSSION

By using IR-LEGO, we have successfully induced the expression of functional genes in a

single epidermal cell of the larval male tail of *C. elegans*. We showed that expression of PLX-1

in R1.p is sufficient for rescuing the ray 1 displacement defect in *plx-1* mutants. We also

showed that the SMPs signal emanating from R2.p or R2.a, the cells located adjacent to and

posterior to R1.p during a certain period of development, is sufficient for positioning ray 1

properly. Although a previous study suggested that the SMPs signal emanating from remote

cells affects ray-lineage cells (Dalpé et al., 2004), the present results suggest that SMPs signal

mediated in a cell-cell contact-dependent manner is sufficient for determining the position of

ray 1.

It was suggested that the position of a ray precursor cluster, which later generate a ray, is

defined by the position of the boundary between two adjacent R(n).p cells and Hyp 7; the ray

precursor cluster appears retained at the position where the surface of the three cells

intersect (Baird et al., 1991) (Figure 6(a)). Accordingly, our previous studies confirmed that

ray 1 precursor cluster is localized at the junction between R1.p, R2.p, and Hyp 7 in both WT

and *plx-1* worms (Fujii et al., 2002; Nukazuka et al., 2008). Thus, the anterior displacement of

ray 1 in *plx-1* mutants is attributable to the shift of the boundary between R1.p and R2.p

anteriorly to its normal position (Figure 6(b)). Our finding is consistent with a model in which

the SMPs signal emanating from R2.p and R2.a acts on, and promotes or permits the

extension of the posterior edge of R1.p to keep the boundary between R1.p and R2.p at a

proper posterior position (Figure 6(f)).

Induction of PLX-1 expression at stage 2 or 2/3, when the shape of R1.p and R2.p changes

dramatically, rescued the ray 1 defects strongly. Earlier induction of PLX-1 or SMP-1 at stage

1 failed to rescue the defects in the *plx-1* and *smp-1 smp-2* mutants, respectively. It might be

that the proteins induced at stage 1 are no longer present by stage 2.

Unexpectedly, we found that induced expression of SMP-1 in R1.p, the cell on which we

suggest the SMPs signal acts, efficiently rescues the ray 1 positioning defect in *smp-1 smp-2*

mutants. We also found that induction of both SMP-1 and PLX-1 in R1.p of *smp-1 smp-2;*

plx-1 mutants partly rescues the ray 1 defect. These findings suggest that SMP-1, a cell

membrane-bound protein, induced by IR-LEGO *cis*-activates PLX-1 which is co-expressed on

R1.p (Figure 6(g)). The *cis-* interaction between SMP-1 and PLX-1 was reported previously

(Mizumoto and Shen, 2012). Our previous study also showed that SMP-1 *cis*-activates PLX-1

in VPCs only when SMP-1 was overexpressed; when being expressed at a physiological level,

SMP-1 failed to *cis*-activate PLX-1 (Suzuki et al., 2022). Although it is likely that induction

with IR-LEGO leads to overexpression of SMP-1 in R1.p, whether SMP-1 expressed at a

physiological level on R1.p can *cis*-activates PLX-1 remains to be determined.

The results suggest that SMPs signaling promotes or permits, rather than inhibits, the

extension of a cell. This, however, is in contrast to our previous finding on VPCs that SMPs

signaling act as a cell contact-dependent stop signal to arrest the extension of neighboring

cells. It is not clear why the consequence of activation of SMPs signaling differs between these

two types of epidermal cells of *C. elegans*. It is also intriguing that SMPs-PLX-1 signaling

mutants do not exhibit apparent morphological defects in ray-lineage cells other than R1.p,

although our previous studies indicated that SMPs signaling is activated in other ray-lineage

cells in addition to R1.p (Nukazuka et al., 2008; Tanaka et al., 2020). What cellular

mechanisms confer R1.p the specific morphological response to SMPs signaling remains to be

investigated. In the vertebrates, it is well known that some semaphorins act not only as

repulsive but also as attractive axon guidance cues depending on the situation (Song et al.,

1998; Falk et al., 2005). Therefore, context-dependence of the output on the cell shape appears

to be a general feature of the semaphorin signaling systems.

ACKNOWLEDGMENTS

We thank professor Yoichi Oda and the members of the Oda Laboratory for discussion and

comments, Hiromichi Omiya of Sigma Koki Co. Ltd for technical supports, Yuji Kohara for

cDNA clones, Andrew Fire for pPD29.26 and pPD49.78, and Joe Culotti and Jeff Shimske for

strains. Some strains were provided by the Caenorhabditis Genetic Center, which is funded

by the National Institute for Health National Center for Research Resources (P40 OD010440).

This work was supported by grants from the Ministry of Education, Science, Sports Culture

and Technology (MEXT), Japan (S.T.), and a grant from The Naito Memorial Foundation

(S.T.). M.S. is a Doctoral Course (DC1) Research Fellow of the Japan Society for the

Promotion of Science.

Figure legends

Figure 1.   The male tail defect of *smp-1(ev715) smp-2(ev709)* and *plx-1(nc37)* mutants. (a) A

lateral view of the tail of a wild-type adult male. Ray 1 (arrow) is normally located adjacent to

the neighboring ray 2 (arrowhead)(Level 1). (b) Ray 1 is located inside the fan but is separated

from ray 2 (Level 2). (c) The worm showed the severe displacement of ray 1, which is located

outside of the fan (Level 3). (d) The penetrance of the ray 1 phenotype in each mutant. The

lin-32 promoter drives gene expression in R(n) cells and their descendants (ray-lineage cells).

All worms carry *him-5* mutation except for *smp-1; ncEx2038[lin-32p::smp-1]* and *smp-1 smp-2,*

ncEx2038[lin-32p::smp-1], which carry the *him-8* mutation, to increase the incidence of male.

Anterior is left, and dorsal is top. Scale Bars = 10 µm.

Figure 2.   Development of lateral epidermis of the larval male tail. The apical boundary of

epidermal cells was visualized with an apical junction marker, *AJM-1::GFP*. R(n) cells (a) and

their descendants (b-e, h) in the wild-type male. The schematic drawing of each picture is

shown on the right. R(1), R(2), and R(3) and their descendants are marked in green, magenta,

and yellow, respectively. (f, g) The descendants of R(n) cells of a wild-type (WT) (left) and a

plx-1 mutant (right) male between the stages shown (e) and (h). The schematic drawing of

each picture is shown below.

(a) Nine R(n) cells are located on the lateral side of the larval male tail. (b) Each R(n) cell

divides into two daughter cells, R(n).a and R(n).p. (c) The arrangement of R(n).a and R(n).p

changes so that R(n).p is positioned dorsolateral to R(n).a. (d) R(n).a divides into two daughter

cells, R(n).aa and R(n).ap. (e) R(n).aa and R(n).ap each divides again, producing R(n).aaa,

R(n).aap, and R(n).apa, R(n).app, respectively. (f left) R(n).aap executes programmed cell

death, while the remaining three cells, R(n).aaa, R(n).apa and R(n).app, form a ray precursor

cluster. Then the apical surface of a ray precursor cluster shrinks while that of R(n).p

enlarges. Arrow indicates the apical junction between R1.p and R2.p. (g) R1.p fuses with R2.p

to form R1+2.p (light blue). (h) After formation of R1+2.p, the remaining R(n).ps gradually

fuse together to form a syncytial tail seam (set)

(f,g right) In a *plx-1* mutant male, the boundary between R1.p and R2.p shifts anteriorly

(arrow), and the position of ray 1 precursor cluster is dislocated anteriorly.

(i) Cell lineages of R(n) and its descendant cells.   ×: programmed cell death. Anterior is left,

and dorsal is top. Scale Bars = 10 µm.

Figure 3.   Induced expression of PLX-1 in R1.p efficiently rescued the ray 1 defect in *plx-1*

mutants.

A DIC photomicrograph of the ventral view of adult rays of (a) a wild-type and (b) a *plx-1*

mutant animal. (a) Ray 1 (arrow) is juxtaposed to ray 2 (triangle). (b) In the *plx-1* mutant, ray

1 (arrows) on the right side (Level 3) and on the right side (Level 2) were both shifted

anteriorly, separated from ray 2 (triangles). (c) R1.p cell (white arrow) of a *plx-1;*

ncIs19[hsp::plx-1; hsp::gfp; rol-6(su1006)]; ncIs13[ajm-1::gfp] worm was IR-irradiated. (d)

Three hours after irradiation, the expression of GFP, an induction marker, was found only in

the cytoplasm of the targeted R1.p (red arrow). The area below the ray-lineage cell shows the

green signal due to autofluorescence. (e) Induced expression of PLX-1 in the left R1.p at stage

2 rescued the ray 1 defect on the irradiated side (white arrow). Arrows indicate ray 1, and

triangles ray 2 (a, b, e). Scale Bars = 10 µm. (f) The ray 1 phenotype in the adult tail of

IR-irradiated worms. *plx-1; ncIs19[hsp::plx-1; hsp::gfp; rol-6(su1006)]; ncIs13[ajm-1::gfp]*

worms were used. The number of worms that are successfully induced and scored are shown

in the parenthesis. "stage - / PLX-1 –" indicates the ray 1 phenotype of non-irradiated worms.

Stage "2/3" indicates the cases where worms that were irradiated at stage 2 but had

developed into stage within 3 three hours after irradiation. R3 d indicates the combined

scores of induction to R3 daughter cells, R3.a and R3.p. PLX-1 expression in R1.p at stages 2

and 2/3 rescued the ray 1 defect (** $p<0.01$, Ryan's method for multiple comparison for the

proportion of the Level 1 phenotype). (g) Summary of induced expression of PLX-1. Induction

in R1.p (blue) at stage 2 rescued the ray 1 defect in *plx-1* mutants.

Figure 4.　Induced expression of SMP-1 in either R1.p, R2.a or R2.p efficiently rescued the

ray 1 defect in *smp-1 smp-2* mutants.

  (a) The ray 1 phenotype in the adult tail of IR-irradiated *smp-1 smp-2; ncIs24[hsp::smp-1;*

hsp::gfp; rol-6(su1006)]; ncIs13[ajm-1::gfp] worms was scored. SMP-1 was induced in each

R(n) cell of *smp-1 smp-2* double mutants. The number of worms that are successfully induced

and scored are shown in the parenthesis (* $p<0.05$, Ryan's method for multiple comparison for

the proportion of the Level 1 phenotype). (b) Summary of induced expression of SMP-1. SMP-1

expression in R1.p, R2.a or R2.p (in yellow) at the early stage 2 rescued the ray 1 defect.

Figure 5.　Induce expression of SMP-1 and PLX-1 in *smp-1 smp-2; plx-1* mutants of single

R1.p partly rescued the ray 1 displacement.

(a) *smp-1 smp-2; plx-1; ncIs210 [hsp::plx-1; hsp::smp-1; hsp::gfp; rol-6(su1006)];*

ncIs13[ajm-1::gfp] worms at stage 2 were used in this experiment ($p<0.01$, Ryan's method for

multiple comparison for the Level 1 phenotype). The total number of IR-irradiated worms are

shown in the parenthesis. (b) Summary of induced expression of SMP-1 and PLX-1 in *smp-1*

smp-2; plx-1 mutants.

Figure 6.   A model of SMPs signaling among ray-lineage cells.

(a, b) Schema of arrangement of R1.p, R2p and ray 1 precursor cluster in a wild-type (a) and a

plx-1 or *smp-1 smp-2* mutant (b) animal. The position of the ray 1 precursor cluster shifts

anteriorly in the mutant. The blue line shows the apical boundary between R1.p and R2.p,

which determine the position of the ray 1 precursor cluster. (c-e) Summary of the rescue

experiments by using IR-LEGO. Expression of PLX-1 in R1.p and SMP-1 in R2.a/p (c, d) is

sufficient for rescuing the ray 1 defect in mutants, suggesting that the *trans*-interaction

between SMP-1 and PLX-1 (f) promotes extension of the posterior part of R1.p. Induced

expression of PLX-1 and SMP-1 in R1.p (e) also partly rescued the ray 1 defect in the mutants,

suggesting that the *cis*-interaction between SMP-1 and PLX-1 (g) can also promotes the

extension of R1.p.

References

Baird, S. E., Fitch, D. H., Kassem, I. A. & Emmons, S. W. (1991). Pattern formation in the

nematode epidermis: determination of the arrangement of peripheral sense organs in the C.

elegans male tail. Development. 113:515-526.

Brenner, S. (1974) The genetics of Caenorhabditis elegans. Genetics. 77:71-94

Dalpé, G., Brown, L. & Culotti, J. G. (2005) Vulva morphogenesis involves attraction of plexin

1-expressing primordial vulva cells to semaphorin 1a sequentially expressed at the vulva

midline. Development 132:1387-400

Dalpé, G., Zhang, L. W., Zheng, H. & Culotti, J. G. (2004) Conversion of cell movement

responses to Semaphorin-1 and Plexin-1 from attraction to repulsion by lowered levels of

specific RAC GTPases in C. elegans. Development. 131:2073-88.

Emmons, S.W. (2005) Male development. In *WormBook* (ed. The *C. elegans* Research

Community), *WormBook*, doi: 10.1895/wormbook.1.33.1, http://www.wormbook.org

Falk, J., Bechara, A., Fiore, R., Nawabi, H., Zhou, H., Hoyo-Becerra, C., Bozon, M., Rougon, G.,

Grumet, M,, Püschel, A. W., Sanes, J. R., & Castellani, V. (2005) Dual functional activity of

semaphorin 3B is required for positioning the anterior commissure. Neuron.48:63-75.

Ferreira, H. B., Zhang, Y., Zhao, C., & Emmons, S. W. (1999) Patterning of Caenorhabditis

elegans posterior structures by the Abdominal-B homolog, egl-5. Developmental Biology.

207:215-28.

Fujii, T., Nakao, F., Shibata, Y., Shioi, G., Kodama, E., Fujisawa, H & Takagi, S. (2002) Caenorhabditis elegans PlexinA, PLX-1, interacts with transmembrane semaphorins and regulates epidermal morphogenesis. Development. 129:2053-63.

Ginzburg, V. E., Roy, P. J. & Culotti, J. G. (2002) Semaphorin 1a and semaphorin 1b are required for correct epidermal cell positioning and adhesion during morphogenesis in C. elegans. Development. 129:2065-78.

Hahn, A. C., & Emmons, S. W. (2003) The roles of an ephrin and a semaphorin in patterning cell-cell contacts in C. elegans sensory organ development. Developmental Biology. 256:379-88.

Iragavarapu-Charyulu, V., Wojcikiewicz, E., & Urdaneta, A. (2020) Semaphorins in angiogenesis and autoimmune diseases: Therapeutic targets? Front. Immunol. 11:346. doi: 10.3389/fimmu.2020.00346.

Jongbloets, B. C. & Pasterkamp, R. J. (2014) Semaphorin signalling during development. Development. 141(17):3292-7. doi: 10.1242/dev.105544.

Kamei, Y., Suzuki, M., Watanabe, K., Fujimori, K., Kawasaki, T., Deguchi, T., Yoneda, Y., Todo, T., Takagi, S., Funatsu, T., & Yuba, S. (2009) Infrared laser-mediated gene induction in targeted single cells in vivo. Nature Methods 6:79-81

Kang, S. & Kumanogoh, A. (2013). Semaphorins in bone development, homeostasis, and

disease. Semin. Cell Dev. Biol. 24, 163-171.

Kantor, D. B., Chivatakarn, O., Peer, K. L., Oster, S. F., Inatani, M., Hansen, M.J., Flanagan,

J. G., Yamaguchi, Y., Sretavan, D. W , Giger, R. J., & Kolodkin, A. L. (2004) Semaphorin 5A is

a bifunctional axon guidance cue regulated by heparan and chondroitin sulfate proteoglycans.

Neuron.44:961-75.

Kumanogoh, A., & Kikutani, H. (2013). Immunological functions of the neuropilins and

plexins as receptors for semaphorins. Nat. Rev. Immunol. 13, 802-814.

Liu, Z., Fujii, T., Nukazuka, A., Kurokawa, R., Suzuki, M., Fujisawa, H., & Takagi, S. (2005) C.

elegans PlexinA PLX-1 mediates a cell contact-dependent stop signal in vulval precursor cells.

Developmental Biology 282:138-51.

Mello, C. & Fire, A. (1995) DNA transformation. Methods Cell Biol. 48, 451–482.

Mizumoto, K., & Shen, K. (2013). Interaxonal interaction defines tiled presynaptic

innervation in C. elegans. Neuron, 77, 655–666.

Mohler, W. A., Simske, J. S., Williams-Masson, E. M., Hardin, J. D., & White, J. G. (1998)

Dynamics and ultrastructure of developmental cell fusions in the Caenorhabditis elegans

hypodermis. Curr. Biol. 8, 1087–1090.

Nakao, F., Hudson, M. L., Suzuki, M., Peckler, Z., Kurokawa, R., Liu, Z., Gengyo-Ando, K.,

Nukazuka, A., Fujii, T., Suto, F., et al. (2007) The plexin PLX-2 and the ephrin EFN-4 have

distinct roles in MAB-20/Semaphorin 2A signaling in Caenorhabditis elegans morphogenesis.

Genetics 176:591–1607.

Nukazuka, A., Fujisawa, H., Inada, T., Oda, Y. & Takagi, S. (2008) Semaphorin controls

epidermal morphogenesis by stimulating mRNA translation via eIF2alpha in Caenorhabditis

elegans. Genes & Development 22:1025–1036

Portman, D. S., & Emmons, S. W. (2000) The basic helix-loop-helix transcription factors

LIN-32 and HLH-2 function together in multiple steps of a C. elegans neuronal sublineage.

Development. 127:5415-26.

Pasterkamp, R. (2012) Getting neural circuits into shape with semaphorins. Nat Rev

Neurosci 13, 605–618. https://doi.org/10.1038/nrn3302

Salser, S. J. & Kenyon, C. (1996) A C. elegans Hox gene switches on, off, on and off again to

regulate proliferation, differentiation and morphogenesis. Development.122:1651-61.

Shioi, G., Shoji, M., Nakamura, M., Ishihara, T., Katsura, I., Fujisawa, H., & Takagi, S. (2001).

Mutations affecting nerve attachment of Caenorhabditis elegans. Genetics, 157, 1611–1622.

Song, H., Ming, G., He, Z., Lehmann, M., McKerracher, L., Tessier-Lavigne, M., & Poo, M.

(1998) Conversion of neuronal growth cone responses from repulsion to attraction by cyclic

nucleotides. Science. 281:1515-8.

Sulston, J. E., Albertson, D. G., & Thomson, J. N. (1980) The Caenorhabditis elegans Male:

Postembryonic Development of Nongonadal Structures. Developmental Biology 78: 542-576,

doi: 10.1016/0012-1606(80)90352-8

Suzuki, M., Toyoda, N., Shimojou, M., & Takagi, S. (2013). Infrared laser-induced gene

expression in targeted single cells of Caenorhabditis elegans. Dev. Growth Differ.

55(4):454-61. doi: 10.1111/dgd.1206

Suzuki, M., Nukazuka, A., Kamei, Y., Yuba, S., Oda, Y., & Takagi, S. (2022). Mosaic gene

expression analysis of semaphorin–plexin interactions in Caenorhabditis elegans using the

IR-LEGO single-cell gene induction system. Development, Growth & Differentiation, 1–13.

https://doi.org/ 10.1111/dgd.12793

Tamagnone, L. (2012). Emerging role of semaphorins as major regulatory signals and

potential therapeutic targets in cancer. Cancer Cell 22, 145-152.

Tanaka, H., Kanatome, A., & Takagi S. (2020) Involvement of the synaptotagmin/stonin2

system in vesicular transport regulated by semaphorins in Caenorhabditis elegans epidermal

cells. Genes to Cells 25(6):391-401.

Togashi, K., von Schimmelmann, M. J., Nishiyama, M., Lim C. S., Yoshida, N., Yun B., Molday,

1 R. S., Goshima, Y, & Hong, K. (2008) Cyclic GMP-gated CNG channels function in

2 Sema3A-induced growth cone repulsion. Neuron. 58:694-707.

4 Tran, T. S., Kolodkin, A. L. & Bharadwaj, R. (2007) Semaphorin regulation of cellular

5 morphology. Annu. Rev. Cell Dev. Biol. 23, 263–292

Supplementary materials

Supplementary Figure S1

(G) The ray 1 phenotype in the adult tail of IR-irradiated *plx-1; ncIs19[hsp::plx-1; hsp::gfp];*

ncIs13[ajm-1::gfp] worms compared with that of *plx-1; jcIs1[ajm-1::gfp; rol-6(su1006)];*

ncIs211[hsp::gfp] worms that serve as controls. Induced expression of GFP alone in R1.p of

control worms did not affect the ray phenotype (** p<0.01, Fisher's exact test).

Supplementary Figure S2

The ray 1 phenotype in the adult tail of IR-irradiated *smp-1 smp-2; ncIs24[hsp::smp-1;*

hsp::gfp]; ncIs13[ajm-1::gfp] worms compared with that of *smp-1 smp-2; ncIs13[ajm-1::gfp];*

ncIs17[hsp::gfp] worms that serve as control. Induced expression of GFP alone in R1.p, R2.a

or R2.p of control worms did not affect the ray phenotype.   (** p<0.01, Fisher's exact test).

Supplementary Table 1

The ray 1 phenotype and the statistical analysis

Supplementary Table 2

List of transgenes used in this paper.

(d)

Ray 1 phenotype

- level 3 (severely defective)
- level 2 (mildly defective)
- level 1 (normal)

Wild-type (151)
plx-1 (110)
smp-1 (109)
smp-2 (108)
smp-1 smp-2 (106)
plx-1;Ex[lin-32p::plx-1] (100)
smp-1;Ex[lin-32p::smp-1] (104)
smp-1 smp-2;Ex[lin-32p::smp-1] (112)

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

The ray 1 phenoype

Mutants

genotype	number				ratio		
	Level 1	Level 2	Level 3	total	Level 1	Level 2	Level 3
WT	142	9	0	151	0.940397	0.059603	0
plx-1	5	49	56	110	0.045455	0.445455	0.509091
smp-1	55	41	13	109	0.504587	0.376147	0.119266
smp-2	104	3	1	108	0.962963	0.027778	0.009259
smp-1 smp-2	3	39	64	106	0.028302	0.367925	0.603774
plx-1; ncEX[lin-32p::plx-1]	63	33	4	100	0.63	0.33	0.04
smp-1; ncEX[lin-32p::smp-1]	94	10	0	104	0.903846	0.096154	0
smp-1 smp-2; ncEX[lin-32p::smp-1]	82	30	0	112	0.732143	0.267857	0

plx-1; Is[hsp::plx-1; hsp::gfp]

stage	target	Level 1	Level 2	Level 3	total	Level 1	Level 2	Level 3
–	–	30	120	100	250	0.12	0.48	0.4
1	R1	6	12	13	31	0.193548	0.387097	0.419355
2	R1.a	6	9	12	27	0.222222	0.333333	0.444444
2	R1.p	23	10	5	38	0.605263	0.263158	0.131579
2/3	R1.p	15	4	1	20	0.75	0.2	0.05
1	R2	2	14	19	35	0.057143	0.4	0.542857
2	R2.a	1	5	5	11	0.090909	0.454545	0.454545
2	R2.p	7	16	13	36	0.194444	0.444444	0.361111
2/3	R2.p	6	12	6	24	0.25	0.5	0.25
3	R1+2.p	3	7	6	16	0.1875	0.4375	0.375
1	R3	1	10	15	26	0.038462	0.384615	0.576923
2	R3	1	14	14	29	0.034483	0.482759	0.482759

plx-1; Is[hsp::gfp]

stage	target	Level 1	Level 2	Level 3	total	Level 1	Level 2	Level 3
–	–	6	15	20	41	0.146341	0.365854	0.487805
2	R1.p	2	4	11	17	0.117647	0.235294	0.647059
2/3	R1.p	3	6	8	17	0.176471	0.352941	0.470588

smp-1 smp-2; Is[hsp::smp-1; hsp::gfp]

stage	target	Level 1	Level 2	Level 3	total	Level 1	Level 2	Level 3
–	–	18	81	80	179	0.100559	0.452514	0.446927
1	R1	2	10	18	30	0.066667	0.333333	0.6
2	R1.a	0	14	6	20	0	0.7	0.3

stage	target	Level 1	Level 2	Level 3	total	Level 1	Level 2	Level 3
2	R1.p	17	13	4	34	0.5	0.382353	0.117647
2/3	R1.p	5	8	1	14	0.357143	0.571429	0.071429
1	R2	3	18	16	37	0.081081	0.486486	0.432432
2	R2.a	8	7	0	15	0.533333	0.466667	0
2	R2.p	10	5	4	19	0.526316	0.263158	0.210526
2/3	R2.p	5	14	3	22	0.227273	0.636364	0.136364
3	R1+2.p	3	15	23	41	0.073171	0.365854	0.560976
1	R3	1	8	8	17	0.058824	0.470588	0.470588
2	R3	1	11	9	21	0.047619	0.52381	0.428571

smp-1 smp-2; Is[hsp::gfp]

stage	target	Level 1	Level 2	Level 3	total	Level 1	Level 2	Level 3
–	–	15	55	47	117	0.128205	0.470085	0.401709
2	R1.p	0	1	9	10	0	0.1	0.9
2	R2.a	1	2	5	8	0.125	0.25	0.625
2	R2.p	1	2	5	8	0.125	0.25	0.625

smp-1 smp-2; plx-1; Is[hsp:smp-1; hsp::plx-1; hsp::gfp]

stage	target	Level 1	Level 2	Level 3	total	Level 1	Level 2	Level 3
	–	2	42	38	82	0.02439	0.512195	0.463415
2	R1.a	1	5	10	16	0.0625	0.3125	0.625
2	R1.p	5	11	6	22	0.227273	0.5	0.272727
2	R2.a	1	6	8	15	0.066667	0.4	0.533333
2	R2.p	1	8	14	23	0.043478	0.347826	0.608696

Statistics　the ray 1 defect (Level 1)

Mutants

	genotype		total	Level 1
#1	*WT*		151	142
#2	*plx-1*		110	5
#3	*smp-1*		109	55
#4	*smp-2*		108	104
#5	*smp-1 smp-2*		106	3
#6	*plx-1; Ex[lin-32p::plx-1]*		100	63
#7	*smp-1; Ex[ln-32p::smp-1]*		104	94
#8	*smp-1 smp-2; Ex[ln-32p::smp-1]*		112	82

Ryan's method for multiple comparison (alpha=0.01)
　　　p[5]=0.02830 vs. p[4]=0.96296 : diff.= 0.93466, RD=0.20842 : P=0.00000, alpha'=0.00179
　　　p[5]=0.02830 vs. p[1]=0.94040 : diff.= 0.91210, RD=0.19358 : P=0.00000, alpha'=0.00208
　　　p[2]=0.04545 vs. p[4]=0.96296 : diff.= 0.91751, RD=0.19345 : P=0.00000, alpha'=0.00208
　　　p[5]=0.02830 vs. p[7]=0.90385 : diff.= 0.87554, RD=0.20829 : P=0.00000, alpha'=0.00250
　　　p[2]=0.04545 vs. p[1]=0.94040 : diff.= 0.89494, RD=0.18159 : P=0.00000, alpha'=0.00250
　　　p[3]=0.50459 vs. p[4]=0.96296 : diff.= 0.45838, RD=0.16735 : P=0.00000, alpha'=0.00250
　　　<mark>p[5]=0.02830 vs. p[8]=0.73214 : diff.= 0.70384, RD=0.19508 : P=0.00000, alpha'=0.00313</mark>
　　　p[2]=0.04545 vs. p[7]=0.90385 : diff.= 0.85839, RD=0.20068 : P=0.00000, alpha'=0.00313
　　　p[3]=0.50459 vs. p[1]=0.94040 : diff.= 0.43581, RD=0.15931 : P=0.00000, alpha'=0.00313
　　　p[6]=0.63000 vs. p[4]=0.96296 : diff.= 0.33296, RD=0.14901 : P=0.00000, alpha'=0.00313
　　　p[5]=0.02830 vs. p[6]=0.63000 : diff.= 0.60170, RD=0.18242 : P=0.00000, alpha'=0.00417
　　　p[2]=0.04545 vs. p[8]=0.73214 : diff.= 0.68669, RD=0.19208 : P=0.00000, alpha'=0.00417
　　　p[3]=0.50459 vs. p[7]=0.90385 : diff.= 0.39926, RD=0.18136 : P=0.00000, alpha'=0.00417
　　　p[6]=0.63000 vs. p[1]=0.94040 : diff.= 0.31040, RD=0.14319 : P=0.00000, alpha'=0.00417
　　　p[8]=0.73214 vs. p[4]=0.96296 : diff.= 0.23082, RD=0.12166 : P=0.00000, alpha'=0.00417
　　　p[5]=0.02830 vs. p[3]=0.50459 : diff.= 0.47629, RD=0.14745 : P=0.00000, alpha'=0.00625
　　　<mark>p[2]=0.04545 vs. p[6]=0.63000 : diff.= 0.58455, RD=0.18389 : P=0.00000, alpha'=0.00625</mark>
　　　p[3]=0.50459 vs. p[8]=0.73214 : diff.= 0.22756, RD=0.17829 : P=0.00048, alpha'=0.00625
　　　p[6]=0.63000 vs. p[7]=0.90385 : diff.= 0.27385, RD=0.16440 : P=0.00001, alpha'=0.00625
　　　p[8]=0.73214 vs. p[1]=0.94040 : diff.= 0.20825, RD=0.11598 : P=0.00000, alpha'=0.00625
　　　p[2]=0.04545 vs. p[3]=0.50459 : diff.= 0.45913, RD=0.15055 : P=0.00000, alpha'=0.01250
　　　p[8]=0.73214 vs. p[7]=0.90385 : diff.= 0.17170, RD=0.13212 : P=0.00117, alpha'=0.01250

plx-1; Is[hsp::plx-1; hsp::gfp]

	stage	induced cell	total	Level 1
#1	–	–	250	30
#2	1	R1	31	6
#3	2	R1.a	27	6
#4	2	R1.p	38	23
#5	2/3	R1.p	20	15
#6	1	R2	35	2
#7	2	R2.a	11	1
#8	2	R2.p	36	7
#9	2/3	R2.p	24	6
#10	3	R1+2.p	16	3
#11	1	R3	26	1
#12	2	R3	29	1

Ryan's method for multiple comparison (alpha= 0.01)
　　　p[12]=0.03448 vs. p[5]=0.75000 : diff.= 0.71552, RD=0.42848 : P=0.00000, alpha'=0.00015
　　　p[12]=0.03448 vs. p[4]=0.60526 : diff.= 0.57078, RD=0.34410 : P=0.00000, alpha'=0.00017
　　　p[11]=0.03846 vs. p[5]=0.75000 : diff.= 0.71154, RD=0.44326 : P=0.00000, alpha'=0.00017
　　　p[11]=0.03846 vs. p[4]=0.60526 : diff.= 0.56680, RD=0.35912 : P=0.00000, alpha'=0.00019

p[6]=0.05714 vs. p[5]=0.75000 : diff.= 0.69286, RD=0.42140 : P=0.00000, alpha'=0.00019
p[6]=0.05714 vs. p[4]=0.60526 : diff.= 0.54812, RD=0.33344 : P=0.00000, alpha'=0.00021
p[7]=0.09091 vs. p[5]=0.75000 : diff.= 0.65909, RD=0.57109 : P=0.00002, alpha'=0.00021
p[7]=0.09091 vs. p[4]=0.60526 : diff.= 0.51435, RD=0.49296 : P=0.00013, alpha'=0.00024
==p[1]=0.12000 vs. p[5]=0.75000 : diff.= 0.63000, RD=0.35211 : P=0.00000, alpha'=0.00024==
==p[1]=0.12000 vs. p[4]=0.60526 : diff.= 0.48526, RD=0.24927 : P=0.00000, alpha'=0.00028==

plx-1; Is[hsp::plx-1; hsp::gfp]

		induced cell	Level 1	Level 2	Level 3
#1		−	30	120	100
#4	2	R1.p	23	10	5
#5	2/3	R1.p	15	4	1

plx-1; Is[hsp::gfp]

#13		−	6	15	20
#14	2	R1.p	2	4	11
#15	2/3	R1.p	3	6	8

Fisher's exact test

       #1 vs #13   P= 0.381527
       ==#4 vs #14   P= 0.000173==141
       ==#5 vs #15   P= 0.000766==052

smp-1 smp-2; Is[hsp::smp-1; hsp::gfp]

	stage	induced cell	total	Level 1
#1	−	−	179	18
#2	1	R1	30	2
#3	2	R1.a	20	0
#4	2	R1.p	34	17
#5	2/3	R1.p	14	5
#6	1	R2	37	3
#7	2	R2.a	15	8
#8	2	R2.p	19	10
#9	2/3	R2.p	22	5
#10	3	R1+2.p	41	3
#11	1	R3	17	1
#12	2	R3	21	1

Ryan's method for multiple comparison (alpha=0.05)
    p[3]=0.00000 vs. p[7]=0.53333 : diff.= 0.53333, RD=0.42446 : P=0.00002, alpha'=0.00076
    p[3]=0.00000 vs. p[8]=0.52632 : diff.= 0.52632, RD=0.38200 : P=0.00000, alpha'=0.00083
    p[12]=0.04762 vs. p[7]=0.53333 : diff.= 0.48571, RD=0.42449 : P=0.00013, alpha'=0.00083
    p[3]=0.00000 vs. p[4]=0.50000 : diff.= 0.50000, RD=0.31647 : P=0.00000, alpha'=0.00093
    p[12]=0.04762 vs. p[8]=0.52632 : diff.= 0.47870, RD=0.38152 : P=0.00003, alpha'=0.00093
    p[11]=0.05882 vs. p[7]=0.53333 : diff.= 0.47451, RD=0.44729 : P=0.00044, alpha'=0.00093
    p[3]=0.00000 vs. p[5]=0.35714 : diff.= 0.35714, RD=0.34239 : P=0.00063, alpha'=0.00104
    p[12]=0.04762 vs. p[4]=0.50000 : diff.= 0.45238, RD=0.31507 : P=0.00000, alpha'=0.00104
    p[11]=0.05882 vs. p[8]=0.52632 : diff.= 0.46749, RD=0.40419 : P=0.00015, alpha'=0.00104
    p[2]=0.06667 vs. p[7]=0.53333 : diff.= 0.46667, RD=0.39973 : P=0.00013, alpha'=0.00104
    p[11]=0.05882 vs. p[4]=0.50000 : diff.= 0.44118, RD=0.33839 : P=0.00002, alpha'=0.00119
    p[2]=0.06667 vs. p[8]=0.52632 : diff.= 0.45965, RD=0.35491 : P=0.00003, alpha'=0.00119
    p[10]=0.07317 vs. p[7]=0.53333 : diff.= 0.46016, RD=0.38456 : P=0.00011, alpha'=0.00119
    p[2]=0.06667 vs. p[4]=0.50000 : diff.= 0.43333, RD=0.28473 : P=0.00000, alpha'=0.00139
    p[10]=0.07317 vs. p[8]=0.52632 : diff.= 0.45315, RD=0.33811 : P=0.00002, alpha'=0.00139
    p[6]=0.08108 vs. p[7]=0.53333 : diff.= 0.45225, RD=0.39594 : P=0.00026, alpha'=0.00139
    p[10]=0.07317 vs. p[4]=0.50000 : diff.= 0.42683, RD=0.26459 : P=0.00000, alpha'=0.00167
    p[6]=0.08108 vs. p[8]=0.52632 : diff.= 0.44523, RD=0.34822 : P=0.00006, alpha'=0.00167

smp-1 smp-2; Is[hsp::smp-1; hsp::gfp]

	stage	induced cell	Level 1	Level 2	Level 3
#1	–	–	18	81	80
#4	2	R1.p	17	13	4
#7	2	R2.a	8	7	0
#8	2	R2.p	10	5	4

smp-1 smp-2; Is[hsp::gfp]

	stage	induced cell	Level 1	Level 2	Level 3
#13	–	–	15	55	47
#14	2	R1.p	0	1	9
#15	2	R2.a	1	2	5
#16	2	R2.p	1	4	6

Fisher's exact test

    #1 vs #13   P=0.636436
    #4 vs #14   P= 4.55293e-06
    #7 vs #15   P= 0.00375678
    #8 vs #16   P= 0.0465011

smp-1 smp-2; plx-1; Is[hsp::plx-1; hsp::smp-1; hsp::gfp]

	stage	induced cell	total	Level 1
#1		–	82	2
#2	2	R1.a	16	1
#3	2	R1.p	22	5
#4	2	R2.a	15	1
#5	2	R2.p	23	1

Ryan's method for multiple comparison (alpha= 0.01)

    p[1]=0.02439 vs. p[3]=0.22727 : diff.= 0.20288, RD=0.19237 : P=0.00052, alpha'=0.00100

Trransgenes

jcIs1	*ajm-1::gfp; rol-6](su1006)*	Mohler et al., 1998
ncIs13	*ajm-1::gfp*	Liu et al., 2005
ncIs17	*hsp::gfp*	Kamei et al., 2009
ncIs19	*hsp::plx-1; hsp::gfp rol-6(su1006)*	Suzuki et al., 2022
ncIs23	*hsp::smp-1; hsp::gfp rol-6(su1006)*	Suzuki et al., 2022
ncIs24	*hsp::smp-1; hsp::gfp; rol-6(su1006)*	This paper
ncIs210	*hsp::plx-1;hsp::smp-1; hsp::gfp; rol-6(su1006)*	This paper
ncIs211	*hsp::gfp*	This paper
ncEx2030	*lin-32p::plx-1; rol-6(su1006)*	Nukazuka et al, 2008
ncEx2038	*lin-32p::smp-1; rol-6(su1006)*	This paper