Figure 1

Figure 2

Figure 6

Blue: DAPI Red: CALR Pink: IL1R1 Cyan: IFNB1 Green: IFNG

An Evaluation of the Tumor Microenvironment through CALR, IL1R1, IFNB1, and IFNG to Assess Prognosis and Immunotherapy Response in Bladder Cancer Patients

Lilong Liu[1#], Zhenghao Liu[1#], Lei Fan[2#], Zhipeng Yao[1#], Junyi Hu[1], Yaxin Hou[1], Yang Li[1], Yuhong Ding[1], Yingchun Kuang[1], Ke Chen[1*] ,Yi Hao[3*], Zheng Liu[1*]

[1]Department of Urology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

[2]Department of Urology, Xiangyang Central Hospital, Affiliated Hospital of Hubei University of Arts and Science, Xiangyang, China.

[3]Department of Pathogen Biology, School of Basic Medicine, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

[#]These authors contributed equally.

[*]Correspondence: Ke Chen: shenke@hust.edu.cn, Yi Hao: haoyi@hust.edu.cn, Zheng Liu: lz2013tj@163.com

Abstract

Immunogenic cell death (ICD) is a type of cell death sparking adaptive immune responses, can reshape the tumor microenvironment (TME). Exploring key ICD-related genes in bladder cancer (BLCA) could enhance personalized treatment. TCGA BLCA patients were divided into two ICD subtypes: ICD-high and ICD-low. High ICD expression linked to increased immune cell infiltration and longer survival, but with potentially suppressed immune function. The high ICD group responded better to PD1-targeted therapy. A risk-scoring model with four ICD-related genes (CALR, IL1R1, IFNB1, IFNG) was validated across TCGA, GEO datasets, and tissue samples, showing higher risk-score correlated with weaker anti-tumor immune function, more tumor-promoting elements, lower immunotherapy response rates, and shorter patient survival.This study connects ICD-related genes to BLCA prognosis and immune infiltration, offering a vital tool for personalized treatment guidance.

Keywords：Immunogenic cell death; bladder cancer; risk signature; prognosis; immunotherapy

1 Introduction

Bladder cancer (BLCA) is the tenth most common cancer globally, divided into non-muscle-invasive (NMIBC) and muscle-invasive (MIBC) types[1]. NMIBC patients often have high recurrence after surgery, so postoperative treatment like bladder instillation with BCG or chemotherapy is recommended[1]. MIBC is treated with radical cystectomy, often combined with platinum-based neoadjuvant chemotherapy for better results[2]. New immunotherapy using PD-1/PD-L1 inhibitors is showing

promise for BLCA. In 2017, the FDA approved atezolizumab and pembrolizumab for advanced cases intolerant to platinum-containing chemotherapy[3]. Response rates of 11.68% and 24.05% were seen with atezolizumab and pembrolizumab[4,5]. Chemotherapy can affect the tumor immune environment[6], and ongoing trials are exploring combining immune checkpoint inhibitors (ICIs) with chemotherapy (ClinicalTrials.gov Identifier: NCT04383743, NCT04630730)[7]. Research in BLCA immunotherapy is in early stages, highlighting the need for innovative approaches combining chemotherapy and immunotherapy.

Immunogenic cell death (ICD) is a regulated cell death that, in an active immune system, triggers an adaptive immune response by exposing antigens from dying cells[8]. Certain chemotherapy drugs, like cisplatin[9] and gemcitabine[10], can induce ICD. While studied in preclinical BLCA models, the potential benefits of ICD-based therapies in BLCA lack conclusive evidence[11]. It is crucial to explore the impact of ICD in clinical settings for BLCA treatment, be it through chemotherapy or immunotherapy. In this study, we discovered a connection between ICD-related genes and the prognosis and immune infiltration in BLCA patients using TCGA-BLCA, GEO datasets, and tissue microarray staining. Our validated risk-scoring model effectively evaluates immune infiltration, prognosis, immunotherapy response, and drug sensitivity in BLCA, providing guidance for personalized treatment and future research.

2 Materials and Methods

2.1 Identification of Differentially Expressed ICD-related genes

ICDs were sourced from a previous article[12]. We found differentially expressed genes (DEGs) in TCGA-BLCA using the "Limma" R package. This involved comparing gene expression in normal and tumor samples. DEGs interacting with ICDs were identified. Enrichment scores of ICDs were assessed using ssGSEA analysis with the "GSVA" R package. Patients were then grouped as "high" or "low" based on these scores. Finally, DEGs were visualized through volcano and heatmaps using R packages "pheatmap" and "ggplot2".

2.2 Functional Enrichment Analysis

We compared ICD-high and ICD-low groups by analyzing Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) using the "clusterProfiler" R package. Gene Set Enrichment Analysis (GSEA) assessed differences in expressed gene sets between the groups using the MSigDB collection.

2.3 Analysis of Somatic Mutation

BLCA patient mutation data from TCGA were visualized using "Maftools" in waterfall plots. Boxplots summarized tumor mutational burden (TMB), and the 20 most frequently mutated genes were shown in waterfall plots.

2.4 Analysis of Immune Infiltration in the TME

Used the "ESTIMATE" R package to predict the tumor microenvironment (TME)

scores for BLCA samples and assess immunocyte proportions. XCELL, TIMER, QUANTISEQ, MCPcounter, EPIC, CIBERSORT and CIBERSORT-ABS were used to explore immune cell infiltration, visualizing differences with a violin diagram and analyzing with the Wilcoxon signed-rank test. Evaluated immunotherapeutic response using the IPS algorithm and compared immune checkpoint and HLA-related gene expression in different groups. Also, accessed IPS data from The Cancer Immunome Atlas (TCIA) to compare responses to ICIs across BLCA patient groups.

2.5 Construction and validation of risk-scoring model

To explore the prognostic value of ICD-related genes in BLCA. Using TCGA-BLCA data, we conducted Cox regression analysis and LASSO Cox regression to build a prognostic model. Risk-score=CALR*(0.5378) + IFN1B*(-0.6349) + IFNG*(-0.2028) + IL1R1*(0.0919). Risk-scores were calculated, and patients were divided into low-risk and high-risk groups. We validated the model using the GSE13507 data and BLCA tissue arrays (HBlaU079Su01, Shanghai Outdo Biotech Co., Ltd). Kaplan-Meier curves for overall survival (OS) in low-risk and high-risk groups were created with the R packages "Survminer" and "Survival". Nomogram was created using "rms" package to integrate risk-scores and other clinicopathological information to assess survival probability. ROC analysis was used to evaluate the nomogram's ability to predict survival. The R package "pRRophetic" was used to evaluate the drug sensibility. The Remmele and Stegner's semiquantitative immunoreactive score (IRS) scale was employed to assess the expression levels of each marker, as detailed in earlier study[13]. The R umap package was utilized to conduct the UMAP analyses of the individual cells. A two-sided P <0.05 was considered valuable. The reagents used are as follows: anti-IFNB1 (cat #: bs-0787R, BIOSS), anti-IL1R1 (cat #: bs-20697R, BIOSS), anti-IFNG (cat #: MAB48116, Bioswamp), and anti-CALR (cat #: A1066, ABclonal).

3 Results

3.1 Stratification and pathway enrichment analysis of TCGA-BLCA based on ICD-related genes

We conducted unsupervised clustering analysis on 34 ICD-related genes to identify two ICD-associated subtypes. Subsequently, we defined ICD-high and ICD-low groups based on the expression levels of these genes and compared their ICD gene expression profiles (**Figure 1A**). A clinical heatmap demonstrated differences between the two subtypes in terms of Grade (**Figure 1B**). Furthermore, survival analysis (**Figure 1C**) indicated that the ICD-high group exhibited significantly longer survival times compared to the ICD-low group (p=0.039). Subsequently, we found that patients in the ICD-high group exhibited a significantly higher TMB (p < 0.05) (**Figure S1A**). We also found that TP53, TTN, KMT2D, MUC16, and ARID1A are the most frequently mutated genes in both groups (**Figure S1B, C**).

Next, we examined the differential gene expression between the two subgroups,
identifying 4,321 DEGs, consisting of 2,177 downregulated and 2,144 upregulated
genes. The heatmap displayed the top 50 upregulated and 50 downregulated genes
(**Figure 1D, E**). To further identify pathways associated with immune activation
specific to ICD-high and ICD-low groups, GO, KEGG, and GSEA were conducted.
The results indicate a close association between elevated ICD expression and the
activation of the immune microenvironment (**Figure1F-L**).
3.2 The immune characteristics within the TME in ICD-high and ICD-low
group.
We employed the ESTIMATE algorithm for analysis. The ICD-high group exhibited
higher ESTIMATE scores, immune scores, and stromal scores compared to the
ICD-low group, while the ICD-high group demonstrated lower tumor cell purity
(**Figure 2A-D**). Further analysis of immune cell infiltration between the two subtypes
revealed significantly higher levels of T cells CD8+, T cells CD4+ memory activated,
and T cells follicular helper in the ICD-high group, while Tregs displayed the opposite
trend (**Figure 2E**). Furthermore, we compared the expression of immune
function-related pathways, immune checkpoint genes, and HLA-related genes
between the two subtypes. The results indicate that the ICD-high group exhibited
elevated expression in nearly all 13 immune pathways (**Figure 2F**). In the 8 immune
checkpoint genes, PDCD1LG2, CTLA4, PDCD1, CD274, HAVCR2, LAG3, TIGIT
had higher expression levels, while SIGLEC15 had lower expression (**Figure 2G**).
Among the 24 HLA-related genes, the ICD-high group had higher expression levels
for almost all genes (**Figure 2H**). These results suggest that ICD-high patients have a
significantly higher quantity of tumor-infiltrating immune cells, but they appear to be
in a state of functional exhaustion. Thus, we evaluated the effects of ICIs PD1 and
CTLA4 on the ICD-high and ICD-low groups. Treated with PD1 and CTLA4 or PD1
alone, ICD-high group had better outcomes. With CTLA4 alone, both subtypes had
similar results. Without PD1 or CTLA4 treatment, ICD-high group had significantly
worse outcomes than ICD-low group (**Figure 2I**). Demonstrating that targeting PD-1
is crucial to enhance immune cell functionality and improve the prognosis of
ICD-high patients.
3.3 Construction and validation of risk-scoring model
In Cox univariate analysis, we identified 4 ICD-related genes significantly associated
with OS of patients: CALR (p=0.003), IFNB1 (p=0.037), IFNG (p=0.022), IF1R1
(p=0.047) (**Figure 3A**). Subsequently, these 4 ICD-related genes were tested and
selected in the LASSO regression analysis for constructing the predictive model
(**Figure 3B, C**). Subsequently, patients are categorized into high-risk and low-risk
groups according to the median (the boundary value is 4.372) of the risk-scores.
Survival analysis indicated that, in both the TCGA-BLCA dataset and the GSE13507

dataset, the high-risk group exhibits significantly shorter survival times than the low-risk group (P<0.05) (**Figure 3D, E**). Next, we analyzed somatic mutations and generated waterfall plots for the top 20 mutated genes (**Figure S2A, B**) and calculated the TMB for both groups. Among the ICD-high-risk group and ICD-low-risk group, TP53, TTN, KMT2D, MUC16, and ARID1A were the most frequently mutated genes, but the TMB does not differ between the ICD high-risk group and the ICD low-risk group (**Figure S2C**). Survival analysis also indicated longer survival times in the high TMB group (P<0.001), with the best prognosis observed in the low-risk + high-TMB group and the worst prognosis in the high-risk + low-TMB group (**Figure S2D, E**).

We further conducted a comprehensive analysis using the TCGA-BLCA dataset (**Figure 3F-H**) as the training set and the GSE13507 dataset (**Figure 3I-K**) as the validation set. Based on the median of the risk-scores, we categorized the datasets from TCGA and GEO into high-risk and low-risk groups. We analyzed the correlation between risk-scores and survival status, showing that higher risk-scores increased the risk of patient mortality. Additionally, we examined the expression of the four genes in the risk signature, finding higher expression of CALR and IL1R1 in the high-risk group, and higher expression of IFNB1 and IFNG in the low-risk group (**Figure 3F-K**). Overall, these results suggest a less favorable prognosis for high-risk patients.

3.4 The association between risk-scores and clinical features

We analyzed the association between the four genes composing the risk-score and patient prognosis (**Figure 4A-D**). Patients with low expression of CALR, low expression of IF1R1, high expression of IFNB1, and high expression of IFNG had better prognosis, which is consistent with CALR and IF1R1 being lowly expressed and IFNB1 and IFNG being highly expressed in the low-risk group (**Figure 3H, K**). The clinical heatmap demonstrates differences between the two subtypes in terms of Stage and Grade (**Figure 4E**). In the high-risk group, there is a higher proportion of Stage III and Stage IV cases, while Stage I and Stage II cases are less common (**Figure 4F**). Additionally, the high-risk group has a higher proportion of high-grade cases and a lower proportion of low-grade cases (**Figure 4G**).

To aid clinicians in making more intuitive predictions of individual patient survival probabilities at 1, 3, and 5 years, we created a nomogram that incorporated gender, grade, age, and risk-score (**Figure 4H**). The AUC values for the 1-year, 3-year, and 5-year ROC curves are 0.630, 0.635, and 0.651, respectively (**Figure 4I**). The AUC values for risk-score, age, gender, and stage are 0.630, 0.669, 0.488, and 0.637, respectively (**Figure 4J**). Additionally, we found that the risk-score can serve as an independent prognostic factor (p < 0.001) (**Figure 4K, L**). This suggests that the risk-score is an effective indicator for predicting patient prognosis.

We checked risk-scores and survival in different subgroups. Higher risk-scores were seen in advanced stages (high-grade, stage III, stage IV, and M1 cases),

suggesting an association with advanced disease (**Figure S3A-G**). Analyzing the relationship with patient prognosis, significant survival differences were noted between high-risk and low-risk groups in most subgroups (**Figure S3H-S**). This implies that high-risk classification is linked to shorter survival, especially in advanced stages compared to early stages.To explore the link between risk-scores and the immune microenvironment, we used the ESTIMATE algorithm. The high-risk group showed significantly lower immune scores and ESTIMATE scores than the low-risk group, while there was no notable difference in stromal scores between the two groups (**Figure 4M**). In the context of 13 immune-related pathways, the high-risk group demonstrated lower pathway activity (**Figure S4A**). Further analysis of immune cell infiltration between the two subtypes revealed that the high-risk group had significantly lower levels of T cells CD8, T cells CD4 memory activated, M1, and NK cells resting (**Figure 4N**). Correlation analysis between risk-scores and immune cells indicated a negative correlation with anti-tumor immune cells. Conversely, it showed a positive correlation with immune inhibitory cells (**Figure S5A-P**, **Figure S4B**). In addition, a significant correlation was observed between the 4 individual genes (CALR, IFN1B, IFNG and IL1R1) comprising the risk-score and immune cells, immune checkpoints or immune regulatory pathways (**Figure S6A-P**, **Figure S7A-C**). These analyses suggest that higher risk-scores are linked to weaker anti-tumor immune function and more tumor-promoting elements, partly explaining the shorter survival in the high-risk group.

3.5 The association between the risk-scores and the sensitivity to chemotherapy and immunotherapy

One of the primary factors inducing ICD is chemotherapy drugs, and targeted therapy plays a crucial role in BLCA treatment. Therefore, we further explored the correlation between the risk-scores and the sensitivity to chemotherapy drugs and targeted therapies. The high-risk group exhibited higher sensitivity (lower IC50 values) to sorafenib, epothilone B, docetaxel, elesclomol, MG132, lapatinib, and others. Conversely, the ICD-high risk group showed increased resistance (higher IC50 values) to lenalidomide, sunitinib, gefitinib, methotrexate, and camptothecin (**Figure 5A-P**). Furthermore, we investigated the prognostic characteristics of both groups in response to immunotherapy. The outcomes indicate that without anti-CTLA4 and anti-PD1 treatment, there was no significant difference in survival time between the groups. However, with at least one form of immunotherapy, the low-risk group had a significantly higher response rate than the high-risk group (**Figure 5Q-T**). These findings highlight the clinical potential of the risk-score in predicting sensitivity to various therapies and guiding treatment decisions for better outcomes.

3.6 Validate the risk-scoring model through BLCA tissue arrays and analyze the gene expression with single-cell sequencing

To validate the reliability of the aforementioned results, we conducted an analysis of
the expression levels of CALR, IFN1B, IFNG and IL1R1 and their associations with
prognosis and immune function using BLCA tissue arrays. Immunofluorescence
staining revealed that CALR expression in tumor tissues was higher than in normal
tissues, while the other three genes, IL1R1, IFNB1, and IFNG, showed no significant
differences (**Figure 6A, B**, **Figure S8A**). This observation is consistent with the
results from the TCGA-BLCA dataset exported using the GEPIA 2 tool (**Figure
S8B-E**). Patients were divided into high and low expression groups based on the
median expression levels (IRS scores) of CALR, IFN1B, IFNG, and IL1R1. It was
found that patients in the CALR high-expression group and IL1R1 high-expression
group had shorter survival times, while those in the IFNB1 high-expression group and
IFNB high-expression group had longer survival times (**Figure 6C**), consistent with
the results of TCGA-BLCA dataset (**Figure 4A-D**). Patients' risk-scores were
similarly calculated using the risk-scoring model: Risk-score = CALR-IRS*(0.5378)
+ IFN1B-IRS *(-0.6349) + IFNG-IRS *(-0.2028) + IL1R1-IRS *(0.0919). The
high-risk group (50% vs. 50%) had significantly shorter survival times than the
low-risk group (**Figure 6D**), corroborating the previous results (**Figure 3D, E**). More
importantly, we found a significant positive correlation between the risk-score and
CD39, a molecule we previously confirmed to be associated with immune
suppression[14]. Additionally, the risk score showed a significant positive correlation
with immune exhaustion (CD8+LAG3+), while having a negative correlation with
CD8+ T cell infiltration (the IRS scores are based on our previous research[14]).
However, there was no correlation with tumor volume and T-stage (**Figure 6E-I**).
This suggests a strong link between higher risk scores and immune suppression.
Finally, using our previous single-cell sequencing data[14] and the public dataset
GSE135337, we analyzed the cellular expression patterns of CALR, IFN1B, IFNG,
and IL1R1. CALR is expressed in nearly all cells in the TME, IL1R1 is mainly
expressed in tumor-associated fibroblasts and endothelial cells, and the expression
levels of IFNG and IFNB1 are minimal, primarily by T lymphocytes. This suggests a
crucial role of cancer-associated fibroblasts in remodeling the immune
microenvironment of BLCA (**Figure 6J, K**, **Figure S8F, G**).

4 Disscussion

ICD, a regulated cell death type, can be triggered by internal or external antigens,
activating an adaptive immune response[8]. Some chemotherapies, such as cisplatin and
gemcitabine, can induce ICD, thereby activating the immune microenvironment to
achieve antitumor effects[9,10]. ICIs are a key focus in cancer research, blocking tumors
by reactivating immune cells through targeting checkpoints like PD1/PD-L1 and
CTLA4[15]. Combining chemotherapy with ICI treatment is expected to enhance
immune function and inhibit tumors, given ICD's role in activating the immune

microenvironment, and some studies targeting key genes in the ICD process or
combining chemotherapy with immunotherapy have also achieved encouraging
results[6,16,17]. However, the results of clinical trials indicate that the response rate of
BLCA to ICIs is significantly lower than expected. The tumor immunosuppressive
microenvironment is undoubtedly an important influencing factor. Considering the
complexity of the TME and the impact of ICD, we investigate and confirm the
clinical utility of ICD-related genes using BLCA public databases, BLCA tissue
microarrays, and single-cell sequencing data. This offers theoretical guidance for
applying ICD in clinical settings.

Through unsupervised clustering analysis, we categorized TCGA-BLCA patients
into two groups: ICD-high and ICD-low. Pathway enrichment analysis revealed that
the ICD-high group predominantly clustered in pathways related to the immune
system, consistent with previous literature reports[18,19]. In the analysis of immune
infiltration, we found that the ICD-high group showed higher immune cell infiltration,
coupled with lower tumor purity, indicating a lower invasive ability of the tumor and
an activated immune state[20]. We checked if ICD boosts ICI effectiveness by studying
the connection between ICD-related gene expression and immunotherapy response
rates. Turns out, the ICD-high group showed a much better response compared to the
ICD-low group. This suggests that ICI treatment could work better in people with
high ICD. Thus, we created a risk-scoring model using four ICD-related genes
(CALR, IL1R1, IFNB1, and IFNG) identified through Cox regression and LASSO
regression analyses. This model helps classify BLCA and predict immune infiltration,
prognosis, and response to immunotherapy. Of them, IFNB1 and IFNG, both
belonging to the interferon family, play crucial roles in anti-tumor immunity[21,22].
IL1R1 is a receptor for IL1α and IL1β, a member of the interleukin family, and plays
an immunosuppressive role in immune regulation[23]. In ICD, the CALR
calcium-binding protein moves to the cell membrane to aid antigen presentation, a
widely accepted process[24]. Yet, the general increase in CALR levels does not
necessarily have a positive regulatory effect. Studies indicate that elevated CALR
expression in breast cancer can enhance tumor cell metastasis and resistance to
chemotherapy[25]. Similarly, our study found that high CALR expression in BLCA has
an inhibitory effect. Survival analysis showed that high CALR and IL1R1 expression
was detrimental to patient survival, while high IFNB1 and IFNG expression was
beneficial.

To verify our risk-scoring model, we used univariate and multivariate Cox
regression analyses. The results indicate that the ICD scoring system can
independently predict the prognosis of BLCA patients. Subgroup analysis based on
clinical characteristics showed no difference between high-risk and low-risk groups in
the early stages of the disease. However, in later stages, the high-risk group had

shorter survival times. This may be due to changes in the TME at different disease
stages, with a decrease in immune-killing molecules and an increase in inhibitory
molecules as the tumor advances[26]. Additional analysis of the correlation between the
risk-scores and immune cell infiltration confirmed this: as the risk-score increased, the
infiltration of cells with anti-tumor effects decreased, while the infiltration of M2-type
macrophages and cancer-associated fibroblasts increased. These findings confirm the
efficacy of our risk-scoring model in assessing patients' immune infiltration,
prognosis, and tumor progression.

In treating BLCA, chemotherapy, targeted therapy, and immunotherapy are
effective but have limitations in sensitivity and specificity[27]. Our risk signature
predicts drug sensitivity. High-risk group responds better to chemotherapy, and
low-risk group shows higher response in immunotherapy (PD1 or CTLA4). This
matches the higher immune infiltration in the low-risk patient group, emphasizing that
immune infiltration level is crucial for successful immunotherapy. However, we find
here that the quantity of immune cells in the tumor does not directly decide the
patient's prognosis. This is because many infiltrating immune cells are functionally
exhausted, and only immunotherapy can activate the immune response within the
tumor, improving prognosis. This is confirmed in BLCA tissue microarray stains.
Specifically, the risk-score shows a negative correlation with CD8+ T cell infiltration
but a positive correlation with T cell exhaustion (CD8+LAG3+ or CD39). This shows
our risk scoring model can categorize patients based on immune infiltration,
specifically identifying a high-risk group (lower immune infiltration) and a low-risk
group (higher immune infiltration). Personalized treatment is crucial for patients in
this context. Additionally, in BLCA single-cell sequencing, we found IL1R1 is
abundantly expressed in cancer-associated fibroblasts, linked to tumor promotion and
immunosuppression[28]. This suggests a crucial role for these fibroblasts in shaping the
immune microenvironment of BLCA, guiding our next research.

5 Conclusions

ICD, a type of cell death triggered in an immunogenic environment, influences
immune response. Our study links ICD-related genes to BLCA prognosis and tumor
immune infiltration. We created a validated risk-scoring model with four key genes
(CALR, IL1R1, IFNB1, and IFNG) using TCGA-BLCA, GEO datasets, and tissue
microarray staining. This model effectively assesses immune infiltration, prognosis,
immunotherapy response, and drug sensitivity in BLCA, guiding personalized
treatment and future research.

Acknowledgement

None.

CRediT authorship contribution statement

Lilong Liu：Funding acquisition;Data curation; Formal analysis;Writing - original draft; and Writing - review & editing.Zhenghao Liu, Lei Fan and Zhipeng Yao：Data curation; Formal analysis; Writing - original draft; and Writing - review & editing. Junyi Hu, Yaxin Hou,Yang Li, Yingchun Kuang, and Yuhong Ding: Data curation and Writing - review & editing. Yi Hao played crucial roles in Funding acquisition,conceptualization and Writing - review & editing. Zheng Liu, and Ke Chen played crucial roles in conceptualization and Writing - review & editing. All authors approved the final manuscript for publication.

Ethics approval and consent to participate

Patients provided informed consent for this work. All experimental procedures were approved by the Institutional Review Board of Tongji Medical College, Huazhong University of Science and Technology (IRB: S116).

Consent for publication

Not applicable.

Declaration of interest statement

None.

Funding

This research was supported by the National Natural Science Foundation of China (No.82303623), the National Key Research and Development Project of China (No.2019YFA0905600), the China Postdoctoral Science Foundation (No.2023M731199), and the Postdoctoral Innovation Research Post of Hubei Provincial Department of Human Resources and Social Security (No.331048). The authors acknowledge that the views expressed in this work are personal and may not necessarily represent the perspectives of official institutions.

Data Availability Statement

We obtained TCGA data from https://portal.gdc.cancer.gov/ and GEO data from https://www.ncbi.nlm.nih.gov/geo/. Access our single-cell sequencing data[14] at https://ngdc.cncb.ac.cn/gsa-human/browse/HRA000212 or https://www.ncbi.nlm.nih.gov/bioproject/PRJNA662018/.

References

1. Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA: a cancer journal for clinicians.* 2021;71(3):209-249.
2. Ma Z, Li Z, Mao Y, et al. AhR diminishes the efficacy of chemotherapy via suppressing STING dependent type-I interferon in bladder cancer. *Nature communications.* 2023;14(1):5415.
3. Suzman DL, Agrawal S, Ning YM, et al. FDA Approval Summary: Atezolizumab or Pembrolizumab for the Treatment of Patients with Advanced Urothelial Carcinoma Ineligible for Cisplatin-Containing Chemotherapy. *The oncologist.* 2019;24(4):563-569.
4. Necchi A, Joseph RW, Loriot Y, et al. Atezolizumab in platinum-treated locally advanced or metastatic urothelial carcinoma: post-progression outcomes from the phase II IMvigor210

study. *Annals of oncology : official journal of the European Society for Medical Oncology.*
2017;28(12):3044-3050.

5.   Balar AV, Castellano D, O'Donnell PH, et al. First-line pembrolizumab in cisplatin-ineligible
patients with locally advanced and unresectable or metastatic urothelial cancer
(KEYNOTE-052): a multicentre, single-arm, phase 2 study. *The Lancet Oncology.*
2017;18(11):1483-1492.

6.   Galluzzi L, Buqué A, Kepp O, Zitvogel L, Kroemer G. Immunogenic cell death in cancer and
infectious disease. *Nature reviews Immunology.* 2017;17(2):97-111.

7.   Calleris G, Rouprêt M, Seisen T, et al. Design and rationale of a single-arm phase II study of
neoadjuvant Durvalumab and Gemcitabine associated with Cisplatin or Carboplatin for upper
urinary tract urothelial cancer: the iNDUCT trial (NCT04617756). *World journal of urology.*
2023.

8.   Galluzzi L, Vitale I, Warren S, et al. Consensus guidelines for the definition, detection and
interpretation of immunogenic cell death. *Journal for immunotherapy of cancer.* 2020;8(1).

9.   Chen JL, Yang SJ, Pan CK, et al. Cisplatin and Albumin-Based Gold-Cisplatin Nanoparticles
Enhance Ablative Radiation Therapy-Induced Antitumor Immunity in Local and Distant Tumor
Microenvironment. *International journal of radiation oncology, biology, physics.*
2023;116(5):1135-1149.

10.   McDonnell AM, Lesterhuis WJ, Khong A, et al. Tumor-infiltrating dendritic cells exhibit
defective cross-presentation of tumor antigens, but is reversed by chemotherapy. *European*
journal of immunology. 2015;45(1):49-59.

11.   Xu L, Su B, Mo L, et al. Norcantharidin Induces Immunogenic Cell Death of Bladder Cancer
Cells through Promoting Autophagy in Acidic Culture. *International journal of molecular*
sciences. 2022;23(7).

12.   Garg AD, De Ruysscher D, Agostinis P. Immunological metagene signatures derived from
immunogenic cancer cell death associate with improved survival of patients with lung, breast
or ovarian malignancies: A large-scale meta-analysis. *Oncoimmunology.* 2016;5(2):e1069938.

13.   Remmele W, Stegner HE. [Recommendation for uniform definition of an immunoreactive
score (IRS) for immunohistochemical estrogen receptor detection (ER-ICA) in breast cancer
tissue]. *Der Pathologe.* 1987;8(3):138-140.

14.   Liu L, Hou Y, Deng C, et al. Single cell sequencing reveals that CD39 inhibition mediates
changes to the tumor microenvironment. *Nature communications.* 2022;13(1):6740.

15.   Balar AV, Kamat AM, Kulkarni GS, et al. Pembrolizumab monotherapy for the treatment of
high-risk non-muscle-invasive bladder cancer unresponsive to BCG (KEYNOTE-057): an
open-label, single-arm, multicentre, phase 2 study. *The Lancet Oncology.* 2021;22(7):919-930.

16.   Jafari S, Lavasanifar A, Hejazi MS, Maleki-Dizaji N, Mesgari M, Molavi O. STAT3 inhibitory
stattic enhances immunogenic cell death induced by chemotherapy in cancer cells. *Daru :*
journal of Faculty of Pharmacy, Tehran University of Medical Sciences. 2020;28(1):159-169.

17.   Peng J, Xiao Y, Yang Q, et al. Intracellular aggregation of peptide-reprogrammed small
molecule nanoassemblies enhances cancer chemotherapy and combinatorial immunotherapy.
Acta pharmaceutica Sinica B. 2021;11(4):1069-1082.

18.   Jiang S, Dong Y, Wang J, et al. Identification of immunogenic cell death-related signature on
prognosis and immunotherapy in kidney renal clear cell carcinoma. *Frontiers in immunology.*
2023;14:1207061.

19.   Kuang Y, Jiang B, Zhu H, et al. Classification related to immunogenic cell death predicts
435   prognosis, immune microenvironment characteristics, and response to immunotherapy in
436   lower-grade gliomas. *Frontiers in immunology.* 2023;14:1102094.

437   20.   Dong B, Wu Y, Zhang J, et al. A novel immunogenic cell death-related subtype classification
438   and risk signature for predicting prognosis and immunotherapy efficacy in gastric cancer.
439   *Frontiers in immunology.* 2023;14:1162876.

440   21.   Gaidt MM, Morrow A, Fairgrieve MR, Karr JP, Yosef N, Vance RE. Self-guarding of MORC3
441   enables virulence factor-triggered immunity. *Nature.* 2021;600(7887):138-142.

442   22.   Boehm U, Klamp T, Groot M, Howard JC. Cellular responses to interferon-gamma. *Annual*
443   *review of immunology.* 1997;15:749-795.

444   23.   Liu NN, Yi CX, Wei LQ, et al. The intratumor mycobiome promotes lung cancer progression via
445   myeloid-derived suppressor cells. *Cancer cell.* 2023.

446   24.   Qian Y, Mao J, Leng X, et al. Co-delivery of proanthocyanidin and mitoxantrone induces
447   synergistic immunogenic cell death to potentiate cancer immunotherapy. *Biomaterials*
448   *science.* 2022;10(16):4549-4560.

449   25.   Liu X, Xie P, Hao N, et al. HIF-1-regulated expression of calreticulin promotes breast
450   tumorigenesis and progression through Wnt/β-catenin pathway activation. *Proceedings of*
451   *the National Academy of Sciences of the United States of America.* 2021;118(44).

452   26.   Li K, Shi H, Zhang B, et al. Myeloid-derived suppressor cells as immunosuppressive regulators
453   and therapeutic targets in cancer. *Signal transduction and targeted therapy.* 2021;6(1):362.

454   27.   Tran L, Xiao JF, Agarwal N, Duex JE, Theodorescu D. Advances in bladder cancer biology and
455   therapy. *Nature reviews Cancer.* 2021;21(2):104-121.

456   28.   Liao Z, Tan ZW, Zhu P, Tan NS. Cancer-associated fibroblasts in tumor microenvironment -
457   Accomplices in tumor malignancy. *Cellular immunology.* 2019;343:103729.

Figure legend

Figure 1. Stratification and pathway enrichment analysis of TCGA-BLCA based on

ICD-related genes. A-C. Gene expression heatmap (A), clinical heatmap (B), and

survival analysis (C) of ICD-high and ICD-low groups. D-E. Differential gene

heatmap (D) and volcano plot (E) of ICD-high and ICD-low groups. F-G. GO

enrichment analysis. H. KEGG enrichment analysis. I-L. GSEA enrichment analysis.

Figure 2. The immune characteristics within the tumor microenvironment vary

between ICD-high and ICD-low groups. ESTIMATE scores (A), immune scores (B),

stromal scores (C), and tumor cell purity (D) of ICD-high and ICD-low groups. E.

Immune cell infiltration in the two subgroups. F-H. Expression profiles of the two

subgroups in 13 immune-related pathways (F), 8 immune checkpoints (G), and HLA

gene family (H). I. Efficacy of immunotherapy in the two subgroups.

Figure 3. Construction and validation of risk-scoring model. A. Hazard ratio of four

ICD-related genes identified by Cox regression analysis. B-C. LASSO regression

analysis for constructing the prognostic model. D-E. Survival analysis of high-risk

and low-risk groups in TCGA (D) and GEO (E) databases. Analyzed the correlation

between risk-scores and patients, along with four key genes, in both TCGA (F-H) and
GEO databases (I-K).
Figure 4. The association between risk-scores and clinical features and immune
characteristics. A-D. The association between the four genes composing the
risk-scores and patient prognosis. E. Clinical heatmap of the two subgroups. F-G.
Differences in stage (F) and grade (G) between the two subgroups. H-J. Constructed
nomogram (H) and ROC curve (I-J). K-L. Cox regression analysis.M.Stromal scores,
immune scores, and ESTIMATE scores of the high-risk and low-risk groups. N.
Immune infiltration in the two subgroups.
Figure 5. The association between the risk-score and the sensitivity to chemotherapy
(A-P) and immunotherapy (Q-T).
Figure 6. Validate the risk-scoring model through tissue arrays and single-cell
sequencing. A-B. Analyzing the expression of CALR, IFN1B, IFNG, and IL1R1 using
immunofluorescence staining. C-D. Survival analysis of the four genes (C) and
risk-scores (D) in patients. E-I. The relationship between risk-scores and CD39, CD8+,
CD8+LAG3+, Tumor Size, and T Stage. J-K. Expression analysis of CALR, IFN1B,
IFNG, and IL1R1 in single-cell sequencing data.

A

B

	CALR	IL1R1	IFNB1	IFNG
Normal				
Tumor				

Blue: DAPI   Red: CALR   Pink: IL1R1   Cyan: IFNB1   Green: IFNG

C

CALR-Low / CALR-High — p<0.0001

IL1R1-Low / IL1R1-High — p<0.0001

IFNB1-Low / IFNB1-High — p<0.0001

IFNG-Low / IFNG-High — p<0.0001

D

Low risk / High risk — p<0.0001

E CD39 IRS — R=0.44, p<0.001

F CD8 — R=-0.38, p=0.002

G CD8+LAG3+ IRS — R=0.28, p=0.027

H Tumor Size (cm) — R=-0.23, p=0.080

I T Stage — R=0.23, p=0.083

J — UMAP: Epithelial, Endothelial, Mast cell, T cell, Myeloid, B cell, mCAF, iCAF
Legend: Myeloid, Epithelial, mCAF, iCAF, Endothelial, T cell, B cell, Mast cell

K — CALR, IL1R1, IFNG, IFNB1