A Deep Learning Pipeline for Mapping in situ Network-level Neurovascular
Coupling in Multi-photon Fluorescence Microscopy

Matthew Rozak[1,2], James Mester[1,2], Ahmadreza Attarpour[1,2], Adrienne Dorr[2], Maged
Goubran[1,2,3,*], Bojana Stefanovic[1,2,3,*]

1 - Department of Medical Biophysics, University of Toronto, Toronto, Ontario, Canada
2 - Physical Sciences, Sunnybrook Research Institute, Toronto, Ontario, Canada
3 - Hurvitz Brain Sciences, Sunnybrook Research Institute, Toronto, Ontario, Canada
* co-senior authors
9
10 Corresponding author emails: matthew.rozak@mail.utoronto.ca, bojana.stefanovic@utoronto.ca

1 Abstract

Functional hyperaemia is a well-established hallmark of healthy brain function, whereby local brain blood flow adjusts in response to a change in the activity of the surrounding neurons. Although functional hyperemia has been extensively studied at the level of both tissue and individual vessels, vascular network-level coordination remains largely unknown. To bridge this gap, we developed a deep learning-based computational pipeline that uses two-photon fluorescence microscopy images of cerebral microcirculation to enable automated reconstruction and quantification of the geometric changes across the microvascular network, comprising hundreds of interconnected blood vessels, pre and post-activation of the neighbouring neurons. The pipeline's utility was demonstrated in the Thy1-ChR2 optogenetic mouse model, where we observed network-wide vessel radius changes to depend on the photostimulation intensity, with both dilations and constrictions occurring across the cortical depth, at an average of 16.1 ± 14.3 μm (mean±stddev) away from the most proximal neuron for dilations; and at 21.9 ± 14.6 μm away for constrictions. We observed a significant heterogeneity of the vascular radius changes within vessels, with radius adjustment varying by an average of $24 \pm 28\%$ of the resting diameter, likely reflecting the heterogeneity of the distribution of contractile cells on the vessel walls. A graph theory-based network analysis revealed that the assortativity of adjacent blood vessel responses rose by $152 \pm 65\%$ at 4.3 mW/mm^2 of blue photostimulation *vs.* the control, with a 4% median increase in the efficiency of the capillary networks during this level of blue photostimulation in relation to the baseline. Interrogating individual vessels is thus not sufficient to predict how the blood flow is modulated in the network. Our computational pipeline, to be made openly available, enables tracking of the microvascular network geometry over time, relating caliber adjustments to vessel wall-associated cells' state, and mapping network-level flow distribution impairments in experimental models of disease.

2 Introduction

To support healthy brain functioning, the cerebrovascular network undergoes continual adjustments in vessel calibers [1]–[3]. Neurovascular coupling refers to the change in blood flow following changes in the level of neuronal activity: under physiological conditions, a generous buffer of nutrients is granted to activated parenchyma via the capillary network [1], [4]. This buffer is maintained through finely tuned regulation of flow through changes in the vessel caliber, mediated via contractile cells in the vessel walls. In the absence of such tuning, pockets of tissue could experience inadequate access to metabolites [5]. Alterations in smooth muscle cells, pericytes, and astrocytes may lead to compromises in vessels' dilatory capacity and thus deficits in neurovascular coupling [6]–[9]. In various brain pathologies, including Alzheimer's disease, stroke, and trauma, regional blood flow regulation gets impaired through vessel loss and/or dysfunction of the vessels' dilatory capacity, resulting in regions of ischemia/hypoxia [10], [11]. Previous studies have examined either individual vessels or the tissue level responses, with little attention having been paid to the vascular network, though network dysfunction frequently is associated with accelerated disease progression and long-term symptomatology [3], [12]–[20].

While there is copious data on the functioning of individual vessels, interrogation of the microvascular network remains a challenge, in terms of both data acquisition and analysis [6], [7], [21]–[23]. To date, studies on the brain vasculature have been done by sparsely imaging individual blood vessels at the cellular scale [6], [7], [21], [24]–[32], thereby severely undersampling the microvascular network; or by evaluating the averaged flow over many vessels at the mesoscopic scale, thus failing to discern the flow through individual vessels. A critical gap in the field is the characterization of flow across hundreds of individual vessels, while imaging the network structure that links them together to determine how the vascular response is coordinated across the network. This gap is particularly significant as studies investigating blood flow across several vessels at a time (imaged by varying the line acquisition pattern [6], [24]) have shown highly heterogeneous responses among capillaries. Neuronal function impairments arise wherever local metabolite supply becomes inadequate, notwithstanding the physiological level of flow across the network as a whole, making mapping of vessel changes across the network of particular importance.

To address the limitations of previous work, we developed a novel deep learning (DL)-based pipeline for mapping changes to the geometry of the brain vascular network following neuronal activation, from a time series of volumetric two-photon fluorescence microscopy (2PFM) data. Neuronal activation was elicited by photostimulation of pyramidal neurons expressing Channelrhodopsin-2 (ChR2) [33] in the Thy1-ChR2-YFP mouse model [34]. Our DL pipeline enabled automatic and accurate segmentation, registration and network analysis of large 2PFM datasets across time. We applied our pipeline in a dataset of 17 Thy1-ChR2-YFP mice to map photostimulation-induced changes across the microvascular network - at the level of individual vessels and at the level of vertices spaced every micrometre along the vessels - in

81 relation to the distance to the closest pyramidal neurons expressing the optogenetic actuator and
82 across the cortical depth. Our findings demonstrate the utility of our pipeline for studying in situ
83 microvascular morphology and function to address various neuroscientific hypotheses.

3 Methods

3.1 Cohort

Animals

All experimental procedures in this study followed the ARRIVE 2.0 guidelines [35]. They
were approved by the Animal Care Committee of the Sunnybrook Research Institute, which
adheres to the policies and guidelines of the Canadian Council on Animal Care and meets all the
requirements of the Provincial Statute of Ontario, Animals for Research Act, and the Canadian
Federal Health of Animals Act. We used 15 male and 13 female Thy1-ChR2-YFP mice (#007612,
line 18, Jackson Laboratory) at 6-12 months of age (283.9 ± 67.0 days), weighing 28.8 ± 7.1g on
the day of imaging. The mice were bred in-house and housed under a 12-hr light/dark cycle [34].
In this mouse line, Channelrhodopsin-2 is expressed preponderantly by pyramidal neurons with
soma in cortical layer 5 and dendrites projecting to the cortical surface, enabling depolarization of
their cellular membranes and action potential generation upon blue light illumination [30], [34].
An attrition schematic is provided in Supplementary Figure 1.

Surgical Preparation

On the day of imaging, mice were induced with 5% isoflurane, transferred to a rectal probe
feedback-regulated heating pad (CWE Inc, Ardmore PA) to maintain a temperature of 37 °C, and
maintained under 2% isoflurane. A subcutaneous injection of 1 mL of lactated Ringer's solution
was administered at the start of surgery. Throughout the surgical preparation and imaging, we
monitored breath rate, heart rate, arterial blood oxygen saturation, pulse distention, and breath
distention via a pulse oximeter, with a probe mounted on the thigh (MouseOx, STARR Life
Sciences) (Supplementary Table 1). For fine control over respiration, the mice were
tracheostomized with an endotracheal tube (20 Ga catheter) and ventilated with a gas mixture of
20-30% O_2 and medical air using a small animal ventilator (SAR 830/P, CWE Inc, Ardmore PA)
set to 115-130 breaths per minute at an inspiratory/expiratory ratio of 1:3-4. Their heads were
immobilized via ear bars during the placement of the cranial window and imaging. The tail vein
was cannulated with a 26 Ga catheter to enable fluorophore and anesthetic delivery. A cranial
window was implanted over the forelimb region of the primary somatosensory cortex (AP
0.25mm, ML 2.0 mm); the dura was excised, and 1% agarose was applied between the brain and
the glass coverslip. A well was built using dental cement to allow water immersion of the objective.
Texas Red dextran (70 kDa MW, Thermo Fisher Scientific Inc, Waltham MA) was diluted in PBS

and injected through the tail vein catheter at a concentration of 33 mg/kg for a total injection
volume between 0.1 and 0.2 mL. A line containing 0.01 g/ml alpha chloralose was connected to
the tail vein catheter. The animal was then positioned underneath the microscope objective, and
the structural 2PFM scan acquired, as detailed below. Seventeen mice (nine male and eight female)
made it through the surgical preparation (cf. Supplementary Figure 1). Following the structural
acquisition, the isoflurane was discontinued, and the continuous infusion of alpha chloralose
commenced at 40 mg/kg/hr.

3.2 Imaging

Two-photon Fluorescence Imaging and Optogenetic Stimulation

Mice were imaged on an FVMPE-RS microscope (Olympus, Japan) using a 25x/1.05NA
objective. An Insight tunable Ti:Saphire near-infrared laser (SpectraPhysics, USA) was used for
900 nm excitation of Texas Red-labelled vasculature and YFP-labeled ChR2-expressing pyramidal
neurons. Two visible light stimulation lasers were used to excite ChR2 at either 458 nm or 552 nm
(control). The optical setup is shown in Figure 1. The imaging field-of-view was selected to include
at least one penetrating artery and vein yet avoid major pial vessels and thus signal loss in the
underlying cortex. Structural scans were acquired under 2% isoflurane using a Galvano scanner,
with 2x averaging, a z-step of 0.99 μm, and a nominal lateral resolution of 0.99 μm. Following
structural scanning, mice were allowed to rest for 5 minutes to let blood flow equilibrate from
brain exposure to room light. The five-minute resting period was observed due to prior reports of
red blood cell velocities following optogenetic ChR2 stimulation in the same mouse strain
remaining altered for a minute following photostimulation and due to potential pericyte-mediated
responses via cytoskeleton reorganization occurring over a minute [6], [30].
Functional scans were acquired over baseline and post-stimulus periods, flanking a period
of blue light illumination. The haemodynamic response to optogenetic stimulation in the Thy1-
ChR2-EYFP mouse model was previously observed to last about 45 seconds [30], similar to data
collected in other optogenetic models of mural cell stimulation where arteries were observed to
dilate for approximately 60 and 20 seconds respectively, and capillaries were observed to dilate
on the scale of minutes [6], [23]. We thus constrained our volumetric acquisitions to 45 seconds.
Each volumetric scan lasted for 42.98 seconds and used the resonant scanner with 5x frame
averaging (resulting in a signal-to-noise ratio on the vasculature channel of 6.9±1.6), a nominal
lateral resolution of 0.99 μm, a z-step of 2.64 μm, and traversed 250.8 μm of cortical depth. This
z-step was chosen because larger z-steps led to discontinuities in the capillary bed segmentation.
We imaged the brain in sections from the cortical surface to a depth of 250 μm and from 250 μm
to 500 μm of cortical depth. The acquisitions were split into two slabs to maximize the volume
coverage without creating discontinuities in the capillary segmentation masks while maintaining a
scan duration of under 45 seconds. Stacks were acquired either from the cortical surface down or
from the bottom of the stacks toward the cortical surface on different paired acquisitions, so that
the post-stimulus delay of image acquisition at different cortical depths was not constant. A 239-

um diameter circular photostimulation region was positioned 250 μm beneath the cortical surface. The photostimulation light was raster scanned over this ROI for 5 seconds between imaging scans, with a pixel dwell time of 4 μs for each pixel of the stimulation ROI. Each pixel in the photostimulation ROI was thus excited every 501.55 milliseconds, i.e. at about 2Hz. We used two laser powers for 458-nm photostimulation: 1.1 mW/mm^2 and 4.3 mW/mm^2. The 1.1 mW/mm^2 is just above the threshold for ChR2 photoactivation and is expected to elicit a small degree of neuronal activity, whereas the 4.3 mW/mm^2 photoactivation elicits more neuronal activity while still being below ChR2 saturation level [33], [36]. Low powers were used to activate ChR2-expressing neurons to minimize tissue heating [37]. We also performed control experiments via 552-nm photostimulation (where ChR2 excitation falls to 4% of its peak) at 4.3 mW/mm^2 [38]. Mice rested for approximately 5-7 minutes between scan pairs for the vascular tone to return to its resting state. Photostimulation parameters were presented in randomized order for each mouse.

Figure 1: Photostimulation setup: The excitation and stimulation light pass through a FV30-NDM690 dichroic mirror with two notch filters, at 458 nm and 552 nm, to excite TexasRed, EYFP and ChR2 within the mouse. The emitted light passes through the objective, is reflected off the FV30-NDM690 dichroic mirror, and passes through a 650 nm barrier filter before reaching a 570 nm long pass filter (LPF) separating emitted light from EYFP and TexasRed, which respectively pass through 495-540 nm and 575-630 nm barrier filters to be collected via GaAsP detectors.

3.3 Segmentation and Graph Extraction

Ground truth generation

To generate ground truths, we used ilastik's pixel classification workflow, which relies on random forests, to annotate blood vessels and pyramidal excitatory neurons soma's in 42 volumes from 25 Thy1-ChR2-EYFP mice for training our DL model (24 images from 16 Thy1-ChR2-EYFP mice from other studies were included in the training, validation, and test cohorts to increase our dataset size). Images were semi-automatically segmented in groups of up to 4 each, with a size of 507x507x250 μm each, due to ilastik's inability to handle large amounts of annotations over more than a few volumes. The rater labelled targets (neurons vs. blood vessels), corrected mistakes, and classified the pixels where the output was uncertain (as shown using ilastik's uncertainty guidance feature). During the manual annotation, feature selection was repeatedly optimized using ilastik's suggest feature function (wrapper method), leading to different features being used for the random forest model for each set of images. Small (under 50 pixels) isolated vascular components were removed with connected component analysis in Python using scikit-image and connected-components-3d. Testing data was withheld until the final model was selected based on the models' performance evaluation on the validation data set.

Data preprocessing and augmentation

We used the MONAI python package for data preprocessing, augmentation, model training, and prediction [39]. All images and ground truth segmentation masks were up-sampled to an isotropic voxel size using bilinear and nearest neighbour interpolation. Raw 10-bit image intensities were normalized to range from 0 to 1.0. Each volume had eight 128x128x128 pixel patches randomly cropped out for training. Data augmentation, transformations and parameters are listed in Supplementary Table 2. Spatial transformations were selected to expand data variety via cropping, rotations, and mirroring hence exposing the network to images that would be acquired on different positioning of the animal under the microscope. Zooming and deformation transformations were included to expose the network to small changes in the size and morphology of the vasculature. Intensity and Gaussian transformations exposed the network to signal intensity and contrast variations. Dropping pixels was included as the resonant scanner acquisition occasionally yields images with some relatively low signal pixels.

Model Architecture

We trained a state-of-the-art 3D vision transformer (UNETR) model and the U-Net model for baseline comparison, as implemented in PyTorch using the MONAI library [39]. UNETR is a U-Net style architecture with a transformer-based multi-attention head encoder and a CNN decoder. [40], [41]. The encoder takes an input image (or patch from each batch) and breaks it down into a sequence of non-overlapping patches, each of size 16x16x16 pixels, which are weighted differently to account for variation in signal intensities and patterns within an image. The

sequence of patches is then passed through a multi-head self-attention and multilayer perceptron encoder to capture self-attention between different pixels of the patch to encode long-range relationships between patches. The encoder is then connected to a CNN decoder with skipped connections to the encoder to map features back onto the original image at multiple spatial scales [40]. The model was trained on 2-channel images (EYFP-expressing neurons and Texas Red-labelled vasculature) as inputs to segment neurons and vasculature. We chose the following parameters for the UNETR architecture: a 12 multi-attention head encoder, 16 convolutional features in the first layer of the encoder, a hidden layer size of 768, a multilayer perceptron size of 3072, and Monte Carlo dropout, where on each run of the model 10% of weights were zeroed [40]–[42].

The U-net model used is based on the original 2D U-net proposed by Ronneberger et al. in 2015 and extended into 3D via 3D convolutional operations and features residual blocks, parametric rectifying linear units, and instance normalization as described by Kerfoot et al. [41], [43]–[47]. The residual blocks were implemented for better gradient flow during the network training. At the same time, the parametric ReLU activation functions were used to improve the ability of the model to adjust its weights during training. Instance normalization was implemented to reduce contrast differences between images fed into the network to improve model robustness. The model was implemented with five layers featuring 16, 32, 64, 128, and 256 channels and dropout. The U-net model was trained using the same data augmentation employed during the UNETR model training.

Model Hyperparameter Optimization, Training, and Prediction

We used a grid search to determine optimal hyperparameters for the UNETR and UNet models within the following parameter space [43], [48]: loss functions (Dice Loss, Dice + Cross Entropy Loss, Dice + Focal Loss, or Tversky Loss), dropout rates (0.1, 0.2, or 0.3), learning rates (5e-3, 1e-3, 5e-4, 1e-4, or 1e-5), and the number of residual units for the Unet model (2 or 3) [49], [50]. We utilized the Adam optimizer [51], and the best model during hyperparameter optimization was selected based on the validation Dice similarity coefficient (DSC) and trained for a maximum of 2400 epochs. The Dice score was the principal evaluation metric, to maximize the overlap between ground truth and prediction masks. Precision and recall were used as secondary metrics to achieve balanced precision and recall where over-segmented results were produced. For early stopping, the epoch with the best performance of the DSC on the validation dataset was selected. The final model that had the best Dice score during hyper-parameter optimization was trained with a Dice + Cross Entropy Loss function, a dropout rate of 0.1, 2 residual units, a learning rate of 1e-5, and a batch size of 1 image with 8 crops per image. Training and optimization were performed on the Narval cluster of Calcul Quebec and the Digital Research Alliance of Canada, with each node using 498 GB of RAM and 4 Nvidia A100 GPUs, each with 40GB HBM2 VRAM.

Ilastik utilized a random forest model from the Vigra library with the default parameters and 100 trees [52], [53]. We initially utilized all default 3D Colour/Intensity/Texture/Edge features during ilastik feature selection and added 2D Colour/Intensity/Texture/Edge features with a sigma

of 20.0 pixels. These 2D features were added as a strong, largely planar artifact was observed
towards the surface of the images in the neuron channel. We used the wrapper method for feature
selection with a set size penalty of 0.10 to determine the optimal features from the starting set. The
model was then trained using live updates on the Narval cluster of Calcul Quebec and the Digital
Research Alliance of Canada, with each node using 249 GB of RAM and 2 x AMD Rome 7532
CPUs with 64 cores.
For the ilastik model, a subset of 3 training images from 3 mice was used as ilastik's
random forest was unable to train a model with more data even on compute nodes with large
memory and CPU resources. The model could never complete training within the time limits of
the resource allocations. This inability to complete model optimization was expected as ilastik's
documentation does not recommend training random forest models with full annotation datasets,
and increasing the number of annotations does not necessarily lead to better predictions for the
pixel classification workflow [52]. The ilastik random forest model was trained with whole images
rather than with patches.

Model Comparison

To compare models, we employed several metrics evaluating the similarity between the
ground truth and prediction: the Dice Similarity Coefficient (DSC), Precision, and Recall, as well
as surface-based metrics: 95% Hausdorff distance (HD95) and mean surface distance. We
specifically focused on mean surface distance and HD95 distance since the centerline extraction
used in the downstream analysis was highly sensitive to surface irregularities. The model with
the lowest standard deviation on the mean surface distance was to be selected absent statistically
significant differences in the average mean surface distance between different models.
Consistency in surface placement was key for extracting good centerlines and graphs from the
segmentation masks. The metrics utilized are defined as follows:
273

$$DSC = \frac{2|X \cap Y|}{|X| + |Y|} \tag{1}$$

$$Precision = \frac{True\ Positives}{True\ Positives\ +\ False\ Positives} \tag{2}$$

$$Recall = \frac{True\ Positives}{True\ Positives\ +\ False\ Negatives} \tag{3}$$

$$HD95 = Max_{95\%}\{sup_{x \in X} d(x, Y), sup_{y \in Y} d(X, y)\} \tag{4}$$

$$Mean\ Surface\ Distance \tag{5}$$

$$= \frac{1}{n_X + n_Y}\left(\sum_{p=1}^{n_X} d(p, n_Y) + \sum_{q=1}^{n_Y} d(q, n_X)\right)$$

The DSC measures the overlap between the prediction and ground truth with a value of 1 corresponding to complete overlap. Precision measures the rate of correctly returned predicted values, whereas recall accesses the rate of return of targeted values. Hausdorff 95% distance (HD95) provides the maximum of the 95th percentile of the minimum surface distance between the boundaries of the ground truth and the predicted segmentation mask. The mean surface distance measures the average minimum distance between the surface of the ground truth and the surface of the predicted segmentation mask. Together, the HD95 and mean surface distance assess model performance at the boundaries of the segmentation masks.

Graph extraction of cerebrovascular networks

Graph extraction was performed in Python except for the centerline calculation, which was performed in MatLab (R2021a). Upsampled image acquisitions (0.99 μm isotropic voxel size) were registered with ANTsPy using the rigid registration method with a total sigma of 2 pixels (for smoothing within the registration function) and mean squared error as the similarity metric as input parameters of the *registration* function (Figure 2A,B). We selected the first baseline scan from each region of interest that was scanned from the bottom to the top to serve as a reference to which all other images from the same region were aligned. The calculated transforms were used to transform images from the same ROI to the space of the reference image using linear interpolation [54]. We then generated segmentation masks for each aligned image using our trained UNETR model with sliding window inference, and we retained Monte Carlo dropout during prediction to create an ensemble of 20 UNETR models, so as to assess the model uncertainty (Figure 2D) [55].

To extract the centerlines, we assumed that no background was present within the vessels, as regions of background within vessels disrupt the centerline extraction algorithm. Background labelled pixels surrounded by blood vessel segmentations were thus filled in using connected component analysis. Disconnected vascular components under 50 pixels were assumed to be noise and removed. Each aligned binary segmentation mask was dilated three times with a footprint defined by scikit-image's disk function with a radius of 1 pixel. Next, we computed the union of all segmentation masks from the same ROI generated from registered images (Figure 2E). This common segmentation mask for all time points was eroded to a centerline via a medial axis transformation (*bwskel* function in Matlab (R2021a)). Critically, using the union of all time points minimized the sensitivity of centreline identification to RBC stalls at individual time points. "Hair" segments shorter than 20 micrometres and terminal on one end were removed to reduce false positive vascular branches [56]. The vascular centerlines were next used to construct vascular graphs (with *sknw*) for each ROI, rendering coordinates of vertices along the vessel's centreline as edges, and branch points as nodes [57].

Based on the vascular graphs, we computed vascular radius estimates at each vertex at each timepoint and then calculated the vertex's distance to the nearest YFP-expressing pyramidal neuron (as measured by the distance transformation at the vertex to the nearest labelled neuron, with the radius of the vessel subtracted off). We employed a two-stage approach to estimate the radius for each point on the centerline. First, using the binary segmentation mask, we calculated

the distance from every vessel pixel in the mask to the nearest background pixel. These values
were averaged for each vessel to estimate its radius. The radius estimate defined the size of the
Gaussian kernel that was convolved with the image to smooth the vessel: smaller vessels were thus
convolved with narrower kernels.

In the second stage, the registered raw image was deconvolved with the point spread
function of the microscope, as measured via FluoSpheres carboxylate-modified Microspheres
(Cat# F8803, Thermo Fisher Scientific Inc, Waltham MA), using Richardson-Lucy deconvolution.
To low pass filter the centreline in advance of computing the tangent vector at each vertex, the
coordinates of the vertices along the centerline were smoothed using a Gaussian with a sigma of
3. Next, the tangent vector to the centerline was specified by calculating the gradient of the
centerline path. Given this local tangent to the vessel's direction of travel at a given vertex, we
extracted image intensities in the orthogonal plane from the deconvolved raw registered image. A
2D Gaussian kernel with sigma equal to 80% of the estimated vessel-wise radius was used to low-
pass filter the extracted orthogonal plane image and find the local signal intensity maximum
searching, in 2D, from the center of the image to the radius of 10 pixels from the center. The
orthogonal plane image was sampled every 10 degrees (as finer radial sampling did not improve
the estimation) along radial lines emanating from the local signal intensity maximum closest to the
center of the image and 5x bicubic upsampled to extract thirty-six 1D signal intensity profiles at
that vertex, as shown in Figure 2H. These line profiles were then convolved with a 1D Gaussian
kernel with a sigma of 80% of the estimated radius of that vessel, and the gradient of each profile
was calculated as shown in Figure 2I, J. The vessel boundary was then placed at the local minimum
of the gradient of each profile. The mean and standard deviation of the boundary distances for the
thirty-six 1D line profiles were calculated, and boundary points greater than 2 standard deviations
away from the mean were excluded (Figure 2K). This radius estimation procedure was repeated
for all vertices of all vessels.

In the last step, we computed the distance from each image voxel to the nearest YFP-
expressing pyramidal neuron by computing the distance from every image pixel not belonging to
the neuron mask to the closest YFP-expressing neuron's soma boundary (based on the neuron's
binary segmentation mask). At the vessel vertices, these distances were adjusted by subtracting the
vessel's radius. We thereby captured the distance from the vessel surface to the closest YFP-
expressing neuron.

344
345

Figure 2. Computational analysis pipeline: **A.** The stacks of 2PFM slices were registered using ANTS rigid registration and aligned to the reference time point. **B.** Images were upsampled using bicubic interpolation to an isotropic resolution of 0.99x0.99x0.99 μm. **C.** An ensemble of UNETR deep learning models with dropout generated segmentation masks at each time point, producing probability maps. **D.** The mean and standard deviation of the probability of each pixel being vasculature were computed and used to create binary vascular segmentation masks. **E.** The union over the vascular segmentation masks for all time points was computed, and background pixel clusters within vessel masks were removed. **F.** The vascular segmentation mask was thinned down to centerlines and rendered as a graph, where edges were vessel segments connecting branch points (nodes). This skeleton was overlaid on the vasculature channel from which the neuron channel was subtracted. **G.** The plane orthogonal to the tangent to the vessel's travel direction was computed every micrometre along the centerline. **H,I.** 1D signal intensity profiles at each centerline vertex were computed in the orthogonal plane every 10°. **J.** The boundary for each profile was placed at the minimum of the signal gradient for that signal intensity profile. **K.** The raw intensity image with the detected boundary points, where outlier boundary points (in green) were defined as points over 2 standard deviations from the mean and excluded. **L.** Visualization of the changes in vertex-wise radii on a sample vascular network.

Boundary detection validation

To assess the uncertainty in the radius estimates at each vertex, we simulated changes to the vascular diameter by "resizing" the extracted orthogonal plane image and adding Gaussian noise to it. These steps were undertaken to validate the ability of the boundary detection method to estimate diameter changes and evaluate the robustness of the estimates to increased amounts of noise. For the diameter change estimation, images were resampled from the orthogonal plane by a random factor, uniformly distributed in the range of 0.5x to 2x. Then, the aforementioned boundary detection methods were used to estimate vertex-wise calibre changes. The end goal of the assessment was to see how close the boundary detection method-based radius change matched the prescribed diameter change. In the second task, we added Gaussian noise with a sigma randomly chosen from a uniform distribution ranging from 0 to 500 SU to the orthogonal plane image. The noise was added to the image after deconvolution with the PSF and the extraction of the orthogonal plane but before Gaussian smoothing. We then reported on the percent change in the radius as a result of resizing or adding noise in relation to the baseline radius.

3.4 Vascular Network Analysis

Leveraging the graph representation of the vasculature in our pipeline, we next used graph theory to better understand the networks' behaviour upon neuronal activation. We looked at morphometrics including vascular segment count density, vascular length density, and mean vessel length for comparison with other work. To demonstrate the benefits of extracting a graphical representation of the vasculature in situ, we looked at graph theory metrics including the assortativity of vascular radius changes and changes to the global efficiency of the capillary (below 5 μm in radius) network following optogenetic stimulation. The assortativity measures the tendency for one vessel to dilate or constrict by a similar amount as its neighbours. The assortativity (q) of radius changes in response to stimulation is defined as the Pearson correlation coefficient of these changes on connected vessels:

$$\text{Assortativity} \qquad q = \frac{\sum_{xy} \ xy(e_{xy} - a_x b_y)}{\sigma_a \sigma_b} \qquad (6)$$

where e_{xy} represents the fraction of vessels (edges) in the network that join together nodes with values x and y (i.e. radius changes with values x and y); a_x and b_y are the percentages of edges connecting nodes with values x and y, and $\boldsymbol{\sigma}_a$ and $\boldsymbol{\sigma}_b$ are the standard deviations of a_x and b_y [58]. The efficiency, in turn, captures how easily the graph can be traversed. For vascular graphs, efficiency can be conceptualized as the average of the inverse of the total resistive distance of the shortest paths of all combinations of vascular junctions, with the hydraulic resistivity serving as the distance between them. The resistivity (equation 7) is summed along the shortest paths, and the inverse of this sum is then averaged across all shortest paths to compute the efficiency (equation 8). Finally, pPhotostimulation-induced change in the efficiency, from its baseline level, is reported.

Resistivity
$$\rho = \frac{8\mu L}{\pi R^4}$$
(7)

Efficiency
(8)
$$E = \frac{1}{N(N-1)} \sum_{i \neq j \in V} \frac{1}{\rho_{ij}}$$

In computing resistivity, we assumed a fluidic viscosity (μ) of 4 cP which is within the
physiological range [59]; L was the length of the capillary; and R was the radius of the capillary.
Vessels greater than 10 μm in diameter were excluded from the efficiency calculation as we wanted
to examine blood flow through the capillary nexus between arteries and veins where blood flow
may be reversible [60]. The sum of the resistivities along the least resistive path specified ρ_{ij} in
the efficiency calculation: this sum quantifies how hard it is for blood to move through the said
microvascular bed. N refers to the number of nodes in the network. The efficiency was calculated
as the average of the inverse of the least resistive paths between all pairs of nodes. The efficiency
increases with increasing radius on the shortest paths through the network.

3.5 Statistical models

To statistically compare deep learning model performance, we used the Wilcoxon signed-
rank test as implemented in *scipy* (1.9.3) [61]. Statistics for vascular radius and network metrics
were performed in R (4.3.1) using restricted maximum likelihood mixed effects models as
implemented in lme4 (1.134) [62], with post hoc comparisons done using *emmeans* (1.8.8) [63].
We ran the following linear mixed effects model, separately on dilating and constricting vessels
that exhibited radius changes above two standard deviations of the vessel's baseline radius, to
examine the effects of photostimulation power on microvascular radius changes in responding
vessels:

ΔRadius ~ Stimulation + (1|Vessel)

Due to the difference in their vessel wall ultrastructure, larger microvessels (above 5 μm in
radius) were examined separately from capillaries (radius < 5 μm).

For graph metrics, we included nesting of the field of view within each subject as a random
effect so as to account for differences in vascular network architecture within an individual. The
linear mixed effects models for the graph metrics used were as follows:

Assortativity ~ Stimulation + (1|Subject/Field of View)

Δ Efficiency ~ Stimulation + (1|Subject/Field of View)

In Tables and text, all values have been quoted as mean ± standard deviation, unless otherwise
specified.

4   Results

Application of our computational pipeline resulted in robust segmentation of the
vasculature and neurons from 4D in situ 2PFM images and rendering of the microvasculature as a
graph. The vessel-wise and vertex-wise calibres were tracked across stimulation conditions and
related to the cortical depth and the distance to the closest YFP-expressing neuron, mapping the
network-level vascular responses to ChR2 activation and revealing the coordination of the
microvascular responses following neuronal activation.

4.1 Segmentation model results and comparisons

We compared an ensemble of UNETR models, an ensemble of U-Net models, and an
ilastik random forest model on a test dataset of 9 images (507x507x250 μm each) from 6 mice.
Examples of segmentation masks produced by each of the models are shown in Figure 4. When
evaluating model performance, we paid close attention to the smoothness of the surface of the
segmentation masks due to the sensitivity of the centerline extraction algorithms to irregularities
in the surface of the masks: smoother vascular segmentation masks resulted in fewer falsely
identified vessel branches. Ilastik tended to over-segment vessels, i.e. the model returned
numerous false positives, having a high recall (0.89±0.19) but low precision (0.37±0.33) (Figure
3, Supplemental Table 3). When comparing the UNETR and U-Net models, we focused on the
surface-based mean surface distance and HD95 distance metrics. Since we observed no significant
differences in these metrics between the two models, we selected the UNETR model as our final
model because it produced more consistent segmentations on visual inspection and showed
significantly better performance than ilastik on HD95 for both vessel and neuron segmentation (p
< 0.05).

Figure 3. Model performance metrics: The Dice, precision, recall, mean surface distance, and
HD95 distance for the vascular (**A**) and neuron (**B**) channels**.** Each model was evaluated on the
same test dataset composed of 9 images (250x507x507 μm each) from 6 mice. A Wilcoxon
signed-rank test was used to compare the model's performance on each performance metric for
images from the test dataset. * p < 0.05, ** p < 0.005, and *** p < 0.0005. p-values were not
adjusted.

Figure 4. Visual model comparison: A. Raw images of the vascular channel with the neuron channel subtracted to facilitate vessel visualization. The first and last stacks in each row span from the cortical surface to 250 μm below the surface, while the middle stack spans from 250 μm below the surface to 500 μm below the surface. All images were from the test dataset, which was unseen during model training. **B.** Ground truth segmentation masks for the vasculature were generated by a rater who utilized ilastik-assisted manual segmentation. **C.** Ilastik predictions generated via a random forest model. **D.** Binary segmentation masks generated by an ensemble of 3D UNet Models. **E.** Binary segmentation masks generated by an ensemble of 3D UNETR models.

4.2 Vessel extraction improvements via image registration

Rigid registration across all time points from the same field of view improved the ability to trace vascular paths from in situ 2PFM data. Firstly, registration decreased the mean squared error (MSE) between acquisitions from 1306±747 to 0.008±0.003 Signal Units. The number of images acquired per field of view ranged from a total of 2 to 10 depending on how many repeats were able to be acquired. Registration enabled the computation of the union of segmentation masks from all time points. This increased the number of vessel segments identified in each field of view from 241±174 based on a single time point to 412±281 vessel segments per field of view (507x507x250 μm, n=107 fields of view). Taking the union of segmentation masks of an image stack across all time points substantially decreased the incidence of gaps in capillaries, likely arising due to "transient RBC plugs." The pipeline's ability to reconstruct the cortical vascular network was thus significantly improved by registering data obtained at different time points.

4.3 Validation of pipeline sensitivity to geometric changes

To evaluate the ability of our computational pipeline to detect vessel caliber changes of various magnitudes, we simulated a range of vascular caliber changes and injected various levels of Gaussian noise. Across >100,000 simulations, the fit of the estimated radius following rescaling against the simulated radius had an R^2 value of 0.68. Figure 5B presents a heatmap of the estimated radius post-scaling vs. simulated radius, across different vertices of vessel centerlines, highlighting the ability of our pipeline to estimate vascular radii accurately. The addition of Gaussian noise revealed the robustness of the pipeline: radius estimates remained stable with increasing noise levels, until the addition of noise with a standard deviation of over 200 SU (with the intensity of the images ranging from 0 to 1023 SU).

492
Figure 5. Estimation of simulated radii changes: A. An image in the plane orthogonal to the local tangent to a capillary with the detected boundary (in blue) and with the estimated radius of 2.28 μm. On the right, this image was resized (upsampling, via bicubic interpolation, by 1.10 times) to simulate dilation. **B.** The plot shows correspondence between the estimated radius following scaling and the simulated level of scaling. **C.** An image in the plane orthogonal to the local tangent of a capillary with the detected boundary (in blue) and with the estimated radius of 3.65 μm. On the right, Gaussian noise with a sigma of 205.36 SU was added to the image. **D.** The estimated % change in the vessel's radius after the addition of varying levels of Gaussian noise, demonstrating the robustness of the radius estimated to noise.

4.4 Vascular morphology and heterogeneity within and among vessels

Segmentation coupled with graph extraction enabled a detailed characterization of the microvascular network properties. The morphological properties of extracted networks are listed in **Table 1**, with the probability densities of the vessel length, baseline vessel radius, mean vessel segment depth, and vessel branch point depth shown in Supplementary Figure 3. On the extracted graphs, the vascular radius and distance to labelled neurons were sampled every 1-1.73 μm, enabling detailed analysis of the relationship between the vessel radius change and the proximity to the YFP-expressing neurons. The radius was tracked across different time points, permitting the analysis of the stimulation-induced change in the vascular caliber. To highlight the ability of the

pipeline to detect vessels that significantly change their radius after stimulation, Figure 6A shows
the standard deviation of the average radii on each vessel segment during baseline frames for three
mice. There was a large difference in this standard deviation across various blood vessels,
showcasing the model's ability to reveal baseline variations within each subject. We examined the
average change in the vascular radius of each vessel segment after vs. before photostimulation
(Figure 6B), with even finer spatial patterns detected by analyzing the vertex-wise radius changes
(Figure 6C). The vascular diameter changes were related to the distance from the vessel's surface
to the closest pyramidal neuron at each vertex of the centerline (Figure 6D). The vertex-wise radii
estimation allowed the assessment of the variations in radii changes within individual blood vessels
(Figure 7). Notably, capillary radius varied along the vessel length across the baseline frames, by
$24 \pm 28\%$ of the mean resting radius. Consequently, point measurements in vessel calibers - that
are widely reported in the literature - do not permit accurate estimation of the microvessel volume
changes. Together, the within- and across-vessel radii estimations illustrate the pipeline's ability
to capture spatial variations in the vascular reactivity and relate it to other morphological features
(e.g., the distance to the closest activated neuron).

Table 1: S1FL Vascular Network Morphological Properties

Metric	Mean ± SD	N = 17 Mice (9M/8F)
Number of Individual Vessels per Volume	368 ± 239	32 FOVs
Vessel Density	5705 ± 3705 mm^{-3}	32 FOVs
Number of Vascular Junctions per Volume	207 ± 154	32 FOVs
Vascular Junction Density	3215 ± 2385 mm^{-3}	32 FOVs
Number of Terminal Vessels per Volume	128 ± 52	32 FOVs
Individual Vessel Length	70.7 ± 61.1 μm	12555 vessel segments
Cumulative Vessel Length Density	0.40 ± 0.22 m/mm^3	32 FOVs
Baseline Vessel Radius	2.19 ± 1.66 μm Range: 0.66-15.88μm	12555 vessel segments
Baseline Intra-Vessel Radius Standard Deviation	0.53 ± 0.47 μm	12555 vessel segments
Baseline Vascular Volume Density	0.010 ± 0.007 mm^3/mm^3	32 FOVs

Number of Pyramidal Neurons per Volume	313 ± 202 Neuronal Somas	32 FOVs
Pyramidal Neuron Density	4872 ± 3145 Neuronal Somas/mm^3	32 FOVs

Figure 6. Vascular graph examples: A. Baseline variability in vessel diameter estimated by the
standard deviation of each vessel's mean radius across baseline time frames. **B.** Mean change in
the vessel radius induced by optogenetic stimulation. **C.** Mean change in the vertexwise radius,
allowing the visualization of heterogeneity of radius changes within each vessel. **D.** Distance
from each vertex to the closest pyramidal neuron. Each row corresponds to the vascular graph of
a different mouse.

538
Figure 7. Vertex-wise radii along vessel lengths of a sample artery, capillary and venule at baseline vs. post-stimulation: A. MIP of an artery, vein, and capillary segments before (left) and after (right) optogenetic stimulation with 458nm light at 1.1 mW/mm². The artery and capillary dilated by 1.33±0.86 μm and 0.42±0.39 μm, respectively (for both p<1e-4, Mann-Whitney U test), whereas there was no significant change in the venular caliber upon photostimulation (p=0.22, Mann-Whitney U test). **B.** Estimates of the vertex-wise radius obtained along each of the three vessels' centrelines, before and after stimulation. **C.** Vertex-wise radii changes in response to optogenetic stimulation. **D.** The vertex-wise distance from the vascular surface to the closest YFP-expressing neuron.

4.5 Vascular reactivity to optogenetic stimulation

The ability of the pipeline to reveal novel spatial relationships between the vascular network reactivity and neuronal activation was demonstrated by examining the relationship between photostimulation-induced microvascular radii responses and 1) the closest YFP-labelled pyramidal neurons within 80μm, and 2) the cortical depth, at the vertex-wise level and across different photostimulations (Figure 8). Vessels were coarsely segregated into small (average radius < 5 μm) and large (average radius > 5 μm) vessels, as we expected them to respond differently due to their differential wall-associated cell composition. Only vessels that significantly responded following optogenetic stimulation were analyzed: a vessel was deemed a responder if its radius changed by more than twice the baseline standard deviation in the vessel's radius. The morphometric properties of the responders, under different stimulation conditions, are listed in Table 2. The average magnitude of the capillary radius change was 1.04 ± 1.11 μm. The variability in the radius change within the vessel was higher in the dilating vessels, 0.77 ± 0.61 μm, then in the constricting vessels, 0.69 ± 0.49 μm, for 458-nm, *4.3* $\frac{mW}{mm^2}$ photostimulation ($p<1e-4$) (with no statistically significant changes detected for 458-nm, *1.1* $\frac{mW}{mm^2}$ photostimulation, and 552-nm, *4.3* $\frac{mW}{mm^2}$ photostimulations). Excluding vessels that did not change their radius by over twice the baseline standard deviation removed almost all large vessels from this analysis (Notwithstanding, Supplemental Figure 3 depicts unfiltered large vessel constrictions and dilations.) We ran mixed effects models (at the vessel level) separately on constricting and dilating vessels to investigate the effect of optogenetic stimulation power on the vessel radius changes. Each of the models was run separately on small and large vessels.

Table 2: Details of responder vessels

Stimulation Condition	Total number of vessel estimates	Average Minimum distance to the closest neuron (μm)	Number of Dilators	Minimum distance from dilators to the closest neuron(μm)	Average vessel depth of dilators (μm)	Diameter change (μm)	Number of constrictors	Minimum distance from constrictors to the closest neuron(μm)	Average vessel depth of constrictors (μm)	Diameter change (μm)
All Vessels			**Dilators**				**Constrictors**			
Capillaires										
552 nm 4.3 $\frac{mW}{mm^2}$	5036	21.2 ± 16.2	144 (2.9%)	25.5 ± 19.0	186 ± 114	0.58 ± 0.92	49 (1.0%)	26.5 ± 19.5	247 ± 122	-0.37 ± 0.30
458 nm 1.1 $\frac{mW}{mm^2}$	10136	18.7 ± 14.5	317 (3.1%)	16.8 ± 13.5	196 ±138	0.90 ± 0.93	255 (2.5%)	22.7 ± 16.3	254 ± 126	-1.39 ± 1.51
458 nm 4.3 $\frac{mW}{mm^2}$	12537	20.6 ± 15.4	575 (4.6%)	16.1 ± 14.3	237 ± 146	0.90 ± 0.77	974 (7.7%)	21.9 ± 14.6	274 ± 103	-1.19 ± 1.13
Large Vessels										

552 nm 4.3 $\frac{mW}{mm^2}$	225	43.1 ± 19.5	1(0.4%)	75.4	82	13.98	0 (0%)	NA	NA	NA
458 nm 1.1 $\frac{mW}{mm^2}$	545	38.4 ±19.5	1 (0.2%)	26.1	402	1.97	1 (0.2%)	19.0	179	-3.65
458 nm 4.3 $\frac{mW}{mm^2}$	569	38.4 ± 20.1	2 (0.35%)	53.1 ± 6.3	84 ± 34	2.47 ± 2.93	6 (1.1%)	43.1 ± 16.3	290 ± 125	-6.07 ± 2.45

4.5.1 Vessels further away from activated neurons constrict while those closer to the activated neurons dilate

We examined the relationship between vascular radius changes and the distance to the closest labelled pyramidal neuron, as microvascular response is thought to result from neuronal activation-elicited generation of vasoactive molecules that diffuse to the neighbouring vessels. For the control condition (552nm, *4.3 $\frac{mW}{mm^2}$* photostimulation), 2.9% of small capillaries dilated while 1.0% of small capillaries constricted; in larger vessels, barely any responded (0.4% dilated). For this control condition, there was no significant difference in the distance from constrictors or dilators to the closest pyramidal neuron. For the 458 nm photostimulation, capillary constrictors were on average farther away than were dilators from the activated pyramidal neuron: dilations occurred 16.8 ± 13.5 μm away from activated neurons while constrictions occurred 22.7 ± 16.3 μm for *1.1 $\frac{mW}{mm^2}$* photostimulation (p=1.5e-3) whereas the *4.3 $\frac{mW}{mm^2}$* photostimulation had dilations occur 16.1 ± 14.3 μm away and 21.9 ± 14.6 μm for constrictors (p<1e-4). There was no significant shift between the distance from vessels to neurons for the 1.1 $\frac{mW}{mm^2}$ and 4.3 $\frac{mW}{mm^2}$ stimulations with 458 nm light.. Dilations in capillaries following 458 nm photostimulation were larger than those following the 552 nm control photostimulation: 0.90 ± 0.93 μm dilatations occurred with 1.1 $\frac{mW}{mm^2}$ and 0.90 ± 0.78 μm with 4.3 $\frac{mW}{mm^2}$ at 458 nm; vs. 0.58 ± 0.92 μm with 4.3 $\frac{mW}{mm^2}$ at 552 nm (p<1e-4). For constrictions, 458 nm photostimulations led to -1.39 ± 1.51 μm radius changes with 1.1 $\frac{mW}{mm^2}$ and -1.20 ± 1.13 μm radius changes with 4.3 $\frac{mW}{mm^2}$ (p=4.4e-3); whereas 552 nm photostimulation induced -0.37 ± 0.30 μm radius changes with 4.3 $\frac{mW}{mm^2}$ of power, which was smaller than the 458-nm induced responses (p=0.02).

4.5.2 Vascular radius changes at increasing cortical depths

Vascular responses were next segregated by the cortical depth of the vessel (i.e. the average vessel distance from the cortical surface) [64]. Dilators tended to be located closer to the cortical surface across all stimulation conditions. Constricting vessels were located at an average 58 ± 187 μm deeper than dilators for 458-nm stimulation at 1.1 $\frac{mW}{mm^2}$ (p=0.02), and 37 ± 179 μm

597   deeper for 458-nm photostimulation at 4.3 $\frac{mW}{mm^2}$ (p<1e-4) (with no change in the mean depth of

598   either constricting or dilating vessels with changes in the photostimulation power).

Figure 8. Optogenetic activation-induced changes in vessel-wise microvascular radii:
Capillary responses included both dilatations, shown in **A.** and constrictions, shown in **B.**, with
changes in the magnitude of the capillary response with increased photostimulation power. $* p <$
0.05, ** $p < 0.005$, and *** $p < 0.0005$. p-values were not adjusted. **C.** Changes to capillary
radii are displayed in relation to the closest pyramidal neurons. The proportion of vessels
constricting increased with the higher intensity of blue light stimulation, and constrictions tended
to occur further away from pyramidal neurons than did dilations. **D.** Mean cortical depth of
responding capillaries showed a tendency for dilators to be closer to the surface and for
constrictors to be deeper in the tissue.

4.5.3 Vascular Network Coordination Following Optogenetic Stimulation

We examined the coordination of changes in the microvascular network as a whole via

assortativity of radius changes and network efficiency changes. The vessel responses were

observed to be assortative, i.e., capillaries mirrored the responses in their neighbours. The

increases in stimulation power were accompanied by increases in the assortativity of capillary
responses: increasing stimulation level resulted in heightened coordination between adjacent
capillaries.

The efficiency increased only at the strongest blue photostimulation, i.e., only at this
level of stimulation did the resistivity along the average of all of the shortest paths between
junctions in the vascular network decrease, resulting in attenuated resistance to flow through the
network. The distribution of changes to the efficiency were highly skewed (with a coefficient of
skewness of -1.06 for green illumination, 2.92 for lower intensity blue photostimulation, and
4.87 for higher intensity blue photostimulation). The median increase in the efficiency induced
by the higher intensity blue photostimulation, of 4% (IQR: -8% to 38%), was significantly higher
than the median -6% (IQR=-9% to 4%) efficiency change following the control green
illumination

Figure 9. Microvascular Network Coordination Following Optogenetic Stimulation: A.
Graph representation of a vascular network of 425 vascular segments from a single image stack.
Vessel segments are depicted as nodes of the graph; vascular segments that are joined at
junctions are connected by edges. Nodes are coloured by the change in the mean vessel-wise
radius following photostimulation with 458 nm light at 4.3 mW/mm^2. **B.** Assortativity of
photostimulation-induced changes in mean capillary radius increased with increasing
photostimulation power. **C.** Photostimulation-induced changes in the efficiency of the capillary
network. The capillary network efficiency changed by a median -0.16 PΩ^{-1} (IQR: -0.39 to 0.10
PΩ^{-1}) in response to green light; -0.14 PΩ^{-1} (IQR: -0.55 to 0.27 PΩ^{-1}) in response to lower
intensity bluelight; and 0.22 PΩ^{-1} (IQR = -0.43;1.47 PΩ^{-1}) in response to higher intensity blue
light.. There was a significant increase (p = 0.03) in the capillary network efficiency post 458-nm
light at 4.3 mW/mm^2, when compared to that following the control green illumination. The
measurements came from 72 paired acquisitions of 32 image stacks acquired in 17 mice
(9M/8F). * p < 0.05, ** p < 0.005, and *** p < 0.0005. p-values were not adjusted.

5   Discussion

Recent studies have demonstrated temporal propagation and coordination in
cerebrovascular responses to neuronal activation, whereby arteries dilated after capillary
exposure to increased potassium ion concentration [65]; opposing geometric changes have been

reported by some studies in capillaries connected by intercapillary tunnelling nanotubes [24].
However, how the effects of these and other mechanisms influence in situ reactivity of the 3D
brain capillary network remains unknown. Here, we developed a pipeline for extracting graphs
of brain microvascular networks from in situ 2PFM and examine coordination within and across
capillaries. Capillary networks and their geometrical changes were imaged via 2PFM during
periods of baseline alternated with photostimulation of ChR2 in pyramidal neurons of transgenic
mice, and the microvascular mesh was evaluated every 1-1.73 μm. The vascular morphology was
then analyzed vertex- and vessel-wise across the entire network. All vessels exhibited significant
heterogeneity in caliber changes along their length. Neuronal activation induced both dilations
and constrictions of vessels and the incidence of constrictions increased with increasing cortical
depth. As the stimulation power increased, the tendency for vessels to change their radius by an
amount similar to their neighbours increased. Only the highest photostimulation intensity elicited
an increase in the network efficiency. Our findings reveal an intricate level of coordination
among brain microvessels and provide a computational analysis platform for interrogating a host
of hypotheses on cerebral microvascular reactivity.

Vascular Segmentation and Network Extraction
Intensity thresholding-based image processing pipelines have been used to examine
vascular networks and quantify vascular morphology, but they have not gained widespread use
due to difficulties in adapting them to highly heterogeneous levels of noise across samples [66]–
[69]. Deep learning-based methods for analyzing vascular morphology from 3D microscopy
images have become prevalent as they provide robust segmentation results over a wide range of
signal-to-noise ratios. Recent work has demonstrated steady improvements of segmentation
models' performance with respect to similarity-based metrics (i.e., Dice scores) [70]–[75];
though surface-based metrics may be better predictors of how amenable segmentation outputs
will be to subsequent morphological analysis of the microvascular network. Our final UNETR
model was selected based on a combination of performance metrics including mean surface
distance, Hausdorf 95% distance, and rater evaluation, to maximize the smoothness of the
surface of generated segmentation masks and reduce false positive branch points during
centerline extraction; thereby leading to higher fidelity rendering of microvascular networks.
Graph generation was greatly facilitated by computing the union of the vascular segmentation
masks across all time points as it enabled tracing of capillaries that had stalls at the individual
time points (since the accompanying loss of the fluorescent label otherwise resulted in graph
discontinuities).
Network morphological properties at baseline were in line with prior work. The vascular
length density measured from fixed tissue ranged from 0.44 to 1.10 m/mm^3 [66], [76]–[80]. Our
reported vascular density in the forelimb region of the primary somatosensory cortex was $0.40 \pm$
0.22 m/mm^3, with the low end of the range value expected due to fluorescence absorption by
hemoglobin in the large pial vessels leading to signal dropout (or shadowing) in the underlying
tissue. Our reported average capillary radius of 2.19 ± 1.66 μm was also in line with other

studies, where the mean capillary radius ranged from 1.75 to 2.2 µm as measured with confocal microscopy or 2PFM [7], [66].

In Situ Changes to Vessel Calibers Upon Neuronal Activation

Caliber changes at individual vertices along vessel centerlines exhibited significant heterogeneity. Such heterogeneity is expected due to non-uniformly distributed alpha smooth muscle actin-containing cells along vessel walls, as well as differential activations leading to heterogeneous metabolic demand within the tissue [1], [6], [81]–[84]. Many previous studies assumed vessel caliber changes to be uniform, compromising the accuracy of the estimates.

As expected, the control 552 nm stimulation led to minimal changes in vessel calibers. To probe for off-target effects, non-transgenic mice were also tested with the same optical setup and photostimulation, with no changes to vascular diameters observed at any of the photostimulation powers utilized (data not shown). In transgenic mice, we detected an average capillary dilation in significantly responding vessels of 70±83% with low-intensity 458 nm stimulation, and 67±61% with higher intensity 458 nm stimulation. Across photostimulation conditions, the capillary dilations ranged from 2% to 805%. These calibre changes were higher than those previously reported, which varied from 2% to 20% depending on the capillary branch order [6], [7], [23], [85], [86]. Far less data are available on constrictions. In the current work, constrictions averaged 47±20% for lower-intensity blue light stimulation and 47±17% for higher-intensity blue light stimulation, with a constriction range of 5% to 97% of the baseline radius. These are higher than the previously reported constrictions of 20% [6], [23], likely due to our identifying as responding vessels only those whose caliber changed by at least twice their baseline caliber variation.

458-nm photostimulation caused capillaries to dilate when pyramidal neurons were close, and constrict when they were further away. However, the stronger blue light stimulation led to an increased rate of constrictions, double that of the low-powered blue light stimulation. For larger vessels, both 458 nm stimulation powers led to a similar dilation level that diminished with increasing distance from activated pyramidal neurons. This tendency for vessels close to neurons to dilate and further away ones to constrict would be expected in flow redirection into regions of high level of neuronal activity. Stimulation power dependence in blood flow changes has previously been reported in optogenetic mouse models with diffuse stimulation via LED probes, and following transcranial alternating current stimulation [87], [88]. However, neither of the previously employed methods was able to discern the spatial relationship between the vascular caliber changes, or relate these changes to the distribution of the stimulated neurons.

Across all stimulations, constricting vessels tended to be located deeper than dilating vessels. As the blue light stimulation power increased, the mean depth of both constricting and dilating vessels increased, likely resulting from higher intensity light reaching pyramidal neurons deeper in the tissue [89], [90]. Our results underscore that the haemodynamic response following targeted neuronal activation is not uniformly distributed across the microvascular network: accurate neurovascular coupling assessment thus requires network-based analysis.

Vascular Network Reactivity

   To study the microvascular network response as a whole, we examined the assortativity between capillary radius changes and network efficiency changes following optogenetic stimulation. These two graph theory metrics were selected as they both leverage the knowledge of the vascular network structure. Assortativity sheds light on how the vascular network coordinates its responses, while efficiency provides insight into the extent to which those changes facilitate flow through the network. The assortativity revealed that as the stimulation power increased, the tendency of vessels to match their changes to those of their neighbours increased. Previously characterized assortative mechanisms include endothelial cell cation conduction via Kir2.1 channels to synchronize vascular responses [65], and spatial adjacency of pericytes on in vitro retinal preparation leading to assortative changes in neighbouring capillaries [81]. Disassortative (causing opposite changes) mechanisms of capillary coordination have also previously been observed in situ and may result from intercapillary nanotubules' signalling causing connected pericytes to undergo opposing changes [24]. While not ruling out the presence of disassortative control mechanisms, our results suggest that assortative mechanisms dominate capillary responses to neuronal activation in the somatosensory cortex.

   The network efficiency here can be thought of as paralleling mean transit time, i.e., the time it takes blood to traverse the capillary network from the arteries to the veins. In situ studies of mean transit time have revealed a high heterogeneity of plasma traversal of the capillary bed during stimulation, with stimulation reducing plasma transit times by 11% to 20% from its resting levels [86], [91], and simulations suggesting that capillary network geometry and locations of caliber changes exert a substantial influence on these responses [92]. The efficiency of the vascular network here increased significantly only with the strongest 458 nm stimulation. Small dilatations may thus not increase flow in the cortex. The differences in efficiency are likely due to the patterns of localized dilations and constrictions within the vascular network. Efficiency calculations are sensitive to bottlenecks when traversing meshes and certain locations constricting or dilating can have profound impacts on the shortest paths between nodes and the path's resistivity. The highest powered 458 nm stimulation increasing efficiency may have resulted from increased assortativity causing dilation in key locations within the microvascular network, leading to a significant reduction in shortest path resistivity.

Pipeline Limitations and Adaptability

   The segmentation model was trained only on vascular and neuronal labels, limiting its generalizability to segmenting alternative cells in the current state. However, it can easily be fine-tuned or retrained to label other brain cells (e.g., pericytes, astrocytes, or endothelial cells). In addition, the segmentation model performed poorly when significant bleeding occurred in the cranial window, compromising the vascular contrast. Our imaging protocol was challenged by the need to resolve individual vessel responses and capture the entire network within the span of the microvascular response to stimulation: we could thus not comment on the temporal evolution of the microvascular response. The temporal evolution of the vascular response in individual vessels, however, has been reported on already using line scanning acquisitions to measure red

blood cell velocity and flux [6], [9], [23], [30], [32], [86]. Moreover, the present analysis pipeline
can readily be applied to 2PFM data obtained with finer temporal (e.g., via a Piezo objective
positioner) or spatial resolution, and/or larger fields of view. Finally, microvascular networks in
different brain areas may show distinct spatiotemporal profiles of response to neuronal
activation. Future work is required to test the generalizability of present findings across different
brain regions.

6 Conclusion

We developed a novel deep learning-based computational pipeline for analysis of a time
series of 3D 2PFM images and investigation of spatial patterns in microvascular network
reactivity to neuronal activation. The microvascular network was represented as a graph,
allowing for the evaluation of network geometry changes over time. We tracked the size of blood
vessels throughout the network and related vessel radius changes to the distance from the
stimulated neurons and the cortical depth. Neuronal activation induced both dilatations and
constrictions of capillaries, and the magnitude of these responses increased with increased
photostimulation levels while showing significant heterogeneity within and between vessels. In
the analysis presented, vertex-wise measurements were aggregated for vessel-wise analysis
resulting in highly robust estimates of vessels' calibers and allowing ready comparisons to
literature. Notwithstanding, the pipeline also affords vertex-wise analysis and thus registration of
microvascular reactivity with other local morphological features, at an unprecedented spatial
scale. With increasing distance of the vessel from the most proximal activated neuron, dilatation
magnitude decreased and the incidence of constrictions increased. At the highest stimulation
level investigated, the incidence of vessel constrictions also increased with cortical depth. With
increasing activation levels, capillaries displayed diameter changes that were similar to their
immediate neighbours, while vascular network efficiency increased only under the strongest
stimulation. Our computational analysis pipeline permits probing microvascular network
reactivity and sheds light on the heterogeneity and coordination of vessel caliber changes across
the microvascular network.The pipeline will be made available to the research community to
propel future studies of neurovascular coupling and network reactivity.

Author contributions

Author contributions

M.W.R. contributed to experimental conceptualization and design, surgical preparation, imaging
and data acquisition, pipeline development, data analysis, and manuscript preparation. J.R.M.
contributed to the experimental design conceptualization and manuscript preparation. A.A.
contributed to pipeline development and manuscript preparation. A.D. contributed to surgical
preparation and manuscript preparation. M.G. and B.S. contributed to funding acquisition,
experimental conceptualization and design, data analysis, manuscript preparation, and supervised
all aspects of the study.

Acknowledgements
We are grateful to Calcul Québec and the Digital Research Alliance of Canada (alliancecan.ca)
for their allocation of compute resources that in part supported this research. This work was
supported by funding from Canadian Institute of Health Research grants PJT376309, PJT156179
and PJT178059. M.W.R was supported by the Queen Elizabeth II/Sunnybrook and Women's
College Health Sciences Centre Graduate Scholarships in Science and Technology. B.S. and
M.G. are supported by the Canada Research Chair program (CRC-2018-00042 and CRC-2021-
00374).
808

809 **Data and code availability statement**
810 Data are available upon request, and the code will be made available at a later date as part of a
811 later release of the MIRACL pipeline.
812

813 **Conflicts of interest**
814 The authors do not have any conflicts of interest regarding this manuscript, its preparation, or the
815 research that went into it.
816
817

818 **Supplementary Figure 1. Attrition:** Flow chart of animal numbers at each step of the
819 experiment.

820
821

Supplementary Figure 2. Vascular Network Characteristics: A. Probability density of the extracted mean radii of vascular segments. **B.** Probability density of the lengths of extracted vessel segments between branch points or terminal ends. **C.** Probability density of the mean vessel segment depths. **D.** Probability density of the depths of vessel branch points. Terminal nodes were excluded from this probability density. 12555 vessels, and 6421 vascular junctions from 17 mice (9M/8F) were used to estimate these PDFs.

Supplemental Figure 3. Large Vessels Radius Changes Following Optogenetic Stimulation:
All vascular responses in large vessels (radius > 5µm) separated into dilators **A.** and constrictors
B. showing an increase in the magnitude of response as stimulation power increases. **C.** Radius
changes to vascular radii in relation to the closest pyramidal neurons (within 80 µm). **D.** Mean
depth of responding large vessels.

Supplementary Table 1: Physiological Monitoring Data

	Mean Across Subjects During Imaging
End-tidal pCO2	15.93 ± 5.07 mmHg
Heart rate	318.30 ± 28.69 BPM
O2 Saturation	96.30 ± 3.50 %
Breath Rate	123.89 ± 12.47 BPM
Temperature	37.15 ± 0.30 °C
Weight	27.94 ± 6.03 g

843 **Supplementary Table 2 Data Augmentations:**

Category	Transformation	Parameters
Cropping	Random spatial cropping	Crops: 8 Crop size: 128x128x128
Rotations	Random rotation 90 degrees	Max rotations: 3
Mirroring	Random flipping	Axes: all
Rotations	Random affine transformation	Rotation restriction: 20°
Zoom	Random zooming	Min zoom: 0.3 Max zoom: 3
Deformations	Random 3D elastic deformation	Sigma: 1 and 3 SU Magnitude: 3-15 μm
	Random grid distortion	Cells: 8 Magnitude: -0.3 to 0.3 of cell width
Intensity Transformations	Random intensity shift	Offset: 0.4 SU
	Random contrast adjustment	Gamma: 0.5 to 5.5
	Random histogram shift	Control points: 4
Gaussian Transformation	Random Gaussian sharpening	Sigma1: 0.5 to 1 SU Sigma2: 0.5 SU Alpha: 10 to 30 SU
	Random Gaussian smoothing	Sigma: 0.25 to 1.25 SU
	Random Gaussian noise	Mean : 0 SU STD: 0.2 SU
Drop pixels	Random coarse dropout	Minimum holes: 50 Maximum holes: 1000 Spatial size: 2 pixels Fill: 0.0001 to 0.1 SU

Supplementary Table 3 Model performance comparisons:

	UNETR	UNet	Ilastik
Vessels			
Dice	0.763±0.096	0.790±0.088	0.449±0.375
Precision	0.811±0.089	0.813±0.076	0.377±0.347
Recall	0.750±0.172	0.797±0.165	0.893±0.198
Hausdorff 95%	13.567±12.131	12.116±12.144	38.921±35.706
Mean Surface Distance	1.900±1.545	1.737±1.622	1.675±1.355
Neurons			
Dice	0.712±0.062	0.766±0.044	0.487±0.196
Precision	0.628±0.111	0.698±0.091	0.435±0.239
Recall	0.841±0.053	0.860±0.061	0.814±252
Hausdorff 95%	23.654±19.299	20.340±22.67	50.000±26.660
Mean Surface Distance	1.248±0.569	2.409±3.901	4.450±7.315

References

[1] C. Iadecola, "The Neurovascular Unit Coming of Age: A Journey through Neurovascular
Coupling in Health and Disease," *Neuron*, vol. 96, no. 1, pp. 17–42, Sep. 2017, doi:
10.1016/j.neuron.2017.07.030.
[2] K. Kisler, A. R. Nelson, A. Montagne, and B. V. Zlokovic, "Cerebral blood flow regulation
and neurovascular dysfunction in Alzheimer disease," *Nat. Rev. Neurosci.*, vol. 18, no. 7,
Art. no. 7, Jul. 2017, doi: 10.1038/nrn.2017.48.
[3] K. Kenney *et al.*, "Cerebral Vascular Injury in Traumatic Brain Injury," *Exp. Neurol.*, vol. 275
Pt 3, pp. 353–366, Jan. 2016, doi: 10.1016/j.expneurol.2015.05.019.
[4] A. A. Phillips, F. H. Chan, M. M. Z. Zheng, A. V. Krassioukov, and P. N. Ainslie,
"Neurovascular coupling in humans: Physiology, methodological advances and clinical
implications," *J. Cereb. Blood Flow Metab.*, vol. 36, no. 4, pp. 647–664, Apr. 2016, doi:
10.1177/0271678X15617954.
[5] T. W. Secomb, R. Hsu, N. B. Beamer, and B. M. Coull, "Theoretical simulation of oxygen
transport to brain by networks of microvessels: effects of oxygen supply and demand on
tissue hypoxia," *Microcirc. N. Y. N 1994*, vol. 7, no. 4, pp. 237–247, Aug. 2000.
[6] D. A. Hartmann *et al.*, "Brain capillary pericytes exert a substantial but slow influence on
blood flow," *Nat. Neurosci.*, pp. 1–13, Feb. 2021, doi: inter.
[7] C. N. Hall *et al.*, "Capillary pericytes regulate cerebral blood flow in health and disease,"
Nature, vol. 508, no. 7494, pp. 55–60, Apr. 2014, doi: 10.1038/nature13165.
[8] J. R. Mester *et al.*, "Attenuation of tonic inhibition prevents chronic neurovascular
impairments in a Thy1-ChR2 mouse model of repeated, mild traumatic brain injury,"
Theranostics, vol. 11, no. 16, pp. 7685–7699, Jun. 2021, doi: 10.7150/thno.60190.
[9] C. Adams *et al.*, "Neurogliovascular dysfunction in a model of repeated traumatic brain
injury," *Theranostics*, vol. 8, no. 17, pp. 4824–4836, 2018, doi: 10.7150/thno.24747.
[10] W. J. Carroll, N. Maganti, and M. K. Gill, "Hypoperfusion of the deep capillary plexus
associated with acute on chronic cocaine use," *Am. J. Ophthalmol. Case Rep.*, vol. 18, p.
100684, Apr. 2020, doi: 10.1016/j.ajoc.2020.100684.
[11] Q. Yang *et al.*, "Cerebral small vessel disease alters neurovascular unit regulation of
microcirculation integrity involved in vascular cognitive impairment," *Neurobiol. Dis.*, vol.
170, p. 105750, Aug. 2022, doi: 10.1016/j.nbd.2022.105750.
[12] A. R. Mayer, M. V. Mannell, J. Ling, C. Gasparovic, and R. A. Yeo, "Functional connectivity
in mild traumatic brain injury," *Hum. Brain Mapp.*, vol. 32, no. 11, pp. 1825–1835, 2011,
doi: https://doi.org/10.1002/hbm.21151.
[13] E. Park, J. D. Bell, I. P. Siddiq, and A. J. Baker, "An analysis of regional microvascular loss
and recovery following two grades of fluid percussion trauma: a role for hypoxia-inducible
factors in traumatic brain injury," *J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood
Flow Metab.*, vol. 29, no. 3, pp. 575–584, Mar. 2009, doi: 10.1038/jcbfm.2008.151.
[14] V. Petkus, S. Krakauskaitė, A. Preikšaitis, S. Ročka, R. Chomskis, and A. Ragauskas,
"Association between the outcome of traumatic brain injury patients and cerebrovascular
autoregulation, cerebral perfusion pressure, age, and injury grades," *Medicina (Mex.)*, vol.
52, no. 1, pp. 46–53, Jan. 2016, doi: 10.1016/j.medici.2016.01.004.
[15] J. Ramos-Cejudo *et al.*, "Traumatic Brain Injury and Alzheimer's Disease: The
Cerebrovascular Link," *EBioMedicine*, vol. 28, pp. 21–30, Feb. 2018, doi:
10.1016/j.ebiom.2018.01.021.
[16] N. Franzmeier *et al.*, "Functional connectivity associated with tau levels in ageing,
Alzheimer's, and small vessel disease," *Brain*, vol. 142, no. 4, pp. 1093–1107, Apr. 2019,
doi: 10.1093/brain/awz026.
[17] J. S. Rabin *et al.*, "Vascular Risk and **β** -Amyloid Are Synergistically Associated with
Cortical Tau: Vascular Risk, Aβ, and Tau," *Ann. Neurol.*, vol. 85, no. 2, pp. 272–279, Feb.

2019, doi: 10.1002/ana.25399.

[18] N. A. Boehme *et al.*, "Axonopathy precedes cell death in ocular damage mediated by blast
exposure," *Sci. Rep.*, vol. 11, no. 1, p. 11774, Jun. 2021, doi: 10.1038/s41598-021-90412-
2.

[19] V. E. Koliatsos and V. Rao, "The Behavioral Neuroscience of Traumatic Brain Injury,"
Psychiatr. Clin. North Am., vol. 43, no. 2, pp. 305–330, Jun. 2020, doi:
10.1016/j.psc.2020.02.009.

[20] J. B. Ware *et al.*, "Relationship of Cerebral Blood Flow to Cognitive Function and Recovery
in Early Chronic Traumatic Brain Injury," *J. Neurotrauma*, vol. 37, no. 20, pp. 2180–2187,
Oct. 2020, doi: 10.1089/neu.2020.7031.

[21] R. A. Hill, L. Tong, P. Yuan, S. Murikinati, S. Gupta, and J. Grutzendler, "Regional Blood
Flow in the Normal and Ischemic Brain Is Controlled by Arteriolar Smooth Muscle Cell
Contractility and Not by Capillary Pericytes," *Neuron*, vol. 87, no. 1, pp. 95–110, Jul. 2015,
doi: 10.1016/j.neuron.2015.06.001.

[22] L. Lindvere, A. Dorr, and B. Stefanovic, "Two-photon fluorescence microscopy of cerebral
hemodynamics," *Cold Spring Harb. Protoc.*, vol. 2010, no. 9, p. pdb.prot5494, Sep. 2010,
doi: 10.1101/pdb.prot5494.

[23] P. J. O'Herron, D. A. Hartmann, K. Xie, P. Kara, and A. Y. Shih, "3D optogenetic control of
arteriole diameter in vivo," *eLife*, vol. 11, p. e72802, Sep. 2022, doi: 10.7554/eLife.72802.

[24] L. Alarcon-Martinez *et al.*, "Interpericyte tunnelling nanotubes regulate neurovascular
coupling," *Nature*, vol. 585, no. 7823, pp. 91–95, Sep. 2020, doi: 10.1038/s41586-020-
2589-x.

[25] S. Sakadžić *et al.*, "Cerebral blood oxygenation measurement based on oxygen-dependent
quenching of phosphorescence," *J. Vis. Exp. JoVE*, no. 51, p. 1694, May 2011, doi:
10.3791/1694.

[26] Y. Guo *et al.*, "Dynamic Volumetric Imaging of Mouse Cerebral Blood Vessels In Vivo with
an Ultralong Anti-Diffracting Beam," *Molecules*, vol. 28, no. 13, p. 4936, Jun. 2023, doi:
10.3390/molecules28134936.

[27] J. T. Giblin *et al.*, "Measuring capillary flow dynamics using interlaced two-photon
volumetric scanning," *J. Cereb. Blood Flow Metab.*, vol. 43, no. 4, pp. 595–609, Apr. 2023,
doi: 10.1177/0271678X221145091.

[28] K. P. McDowell, A.-A. Berthiaume, T. Tieu, D. A. Hartmann, and A. Y. Shih, "VasoMetrics:
unbiased spatiotemporal analysis of microvascular diameter in multi-photon imaging
applications," *Quant. Imaging Med. Surg.*, vol. 11, no. 3, pp. 969–982, Mar. 2021, doi:
10.21037/qims-20-920.

[29] G. Kim *et al.*, "Direct Blood Cell Flow Imaging in Microvascular Networks," *Small*, vol. n/a,
no. n/a, p. 2302244, doi: 10.1002/smll.202302244.

[30] J. R. Mester *et al.*, "In vivo neurovascular response to focused photoactivation of
Channelrhodopsin-2," *NeuroImage*, vol. 192, pp. 135–144, May 2019, doi:
10.1016/j.neuroimage.2019.01.036.

[31] T. N. Kim *et al.*, "Line-Scanning Particle Image Velocimetry: An Optical Approach for
Quantifying a Wide Range of Blood Flow Speeds in Live Animals," *PLOS ONE*, vol. 7, no.
6, p. e38590, Jun. 2012, doi: 10.1371/journal.pone.0038590.

[32] D. Kleinfeld, P. P. Mitra, F. Helmchen, and W. Denk, "Fluctuations and stimulus-induced
changes in blood flow observed in individual capillaries in layers 2 through 4 of rat
neocortex," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 95, no. 26, pp. 15741–15746, Dec. 1998,
doi: 10.1073/pnas.95.26.15741.

[33] E. S. Boyden, F. Zhang, E. Bamberg, G. Nagel, and K. Deisseroth, "Millisecond-timescale,
genetically targeted optical control of neural activity," *Nat. Neurosci.*, vol. 8, no. 9, pp.
1263–1268, Sep. 2005, doi: 10.1038/nn1525.

[34] B. R. Arenkiel *et al.*, "In vivo light-induced activation of neural circuitry in transgenic mice

expressing channelrhodopsin-2," *Neuron*, vol. 54, no. 2, pp. 205–218, Apr. 2007, doi:
10.1016/j.neuron.2007.03.005.
[35] N. P. du Sert *et al.*, "Reporting animal research: Explanation and elaboration for the
ARRIVE guidelines 2.0," *PLOS Biol.*, vol. 18, no. 7, p. e3000411, Jul. 2020, doi:
10.1371/journal.pbio.3000411.
[36] J. Y. Lin, M. Z. Lin, P. Steinbach, and R. Y. Tsien, "Characterization of Engineered
Channelrhodopsin Variants with Improved Properties and Kinetics," *Biophys. J.*, vol. 96,
no. 5, pp. 1803–1814, Mar. 2009, doi: 10.1016/j.bpj.2008.11.034.
[37] R. L. Rungta, B.-F. Osmanski, D. Boido, M. Tanter, and S. Charpak, "Light controls
cerebral blood flow in naive animals," *Nat. Commun.*, vol. 8, no. 1, Art. no. 1, Jan. 2017,
doi: 10.1038/ncomms14191.
[38] T. J. Lambert, "FPbase: a community-editable fluorescent protein database," *Nat.*
Methods, vol. 16, no. 4, Art. no. 4, Apr. 2019, doi: 10.1038/s41592-019-0352-8.
[39] M. Consortium, "MONAI: Medical Open Network for AI." Zenodo, Jul. 25, 2022. doi:
10.5281/zenodo.6903385.
[40] A. Hatamizadeh *et al.*, "UNETR: Transformers for 3D Medical Image Segmentation,"
ArXiv210310504 Cs Eess, Oct. 2021, Accessed: Mar. 01, 2022. [Online]. Available:
http://arxiv.org/abs/2103.10504
[41] E. Kerfoot, J. Clough, I. Oksuz, J. Lee, A. P. King, and J. A. Schnabel, "Left-Ventricle
Quantification Using Residual U-Net," in *Statistical Atlases and Computational Models of*
the Heart. Atrial Segmentation and LV Quantification Challenges, M. Pop, M. Sermesant,
J. Zhao, S. Li, K. McLeod, A. Young, K. Rhode, and T. Mansi, Eds., in Lecture Notes in
Computer Science. Cham: Springer International Publishing, 2019, pp. 371–380. doi:
10.1007/978-3-030-12029-0_40.
[42] N. Srivastava, G. Hinton, A. Krizhevsky, I. Sutskever, and R. Salakhutdinov, "Dropout: A
Simple Way to Prevent Neural Networks from Overfitting," *J. Mach. Learn. Res.*, vol. 15,
no. 56, pp. 1929–1958, 2014.
[43] P. Mojiri Forooshani *et al.*, "Deep Bayesian networks for uncertainty estimation and
adversarial resistance of white matter hyperintensity segmentation," *Hum. Brain Mapp.*,
vol. 43, no. 7, pp. 2089–2108, Jan. 2022, doi: 10.1002/hbm.25784.
[44] D. Ulyanov, A. Vedaldi, and V. Lempitsky, "Instance Normalization: The Missing Ingredient
for Fast Stylization." arXiv, Nov. 06, 2017. doi: 10.48550/arXiv.1607.08022.
[45] K. He, X. Zhang, S. Ren, and J. Sun, "Delving Deep into Rectifiers: Surpassing Human-
Level Performance on ImageNet Classification." arXiv, Feb. 06, 2015. doi:
10.48550/arXiv.1502.01852.
[46] K. He, X. Zhang, S. Ren, and J. Sun, "Identity Mappings in Deep Residual Networks."
arXiv, Jul. 25, 2016. doi: 10.48550/arXiv.1603.05027.
[47] Z. Zhang, Q. Liu, and Y. Wang, "Road Extraction by Deep Residual U-Net," *IEEE Geosci.*
Remote Sens. Lett., vol. 15, no. 5, pp. 749–753, May 2018, doi:
10.1109/LGRS.2018.2802944.
[48] M. Goubran *et al.*, "Hippocampal segmentation for brains with extensive atrophy using
three-dimensional convolutional neural networks," *Hum. Brain Mapp.*, vol. 41, no. 2, pp.
291–308, 01 2020, doi: 10.1002/hbm.24811.
[49] S. S. M. Salehi, D. Erdogmus, and A. Gholipour, "Tversky loss function for image
segmentation using 3D fully convolutional deep networks." arXiv, Jun. 18, 2017. doi:
10.48550/arXiv.1706.05721.
[50] T.-Y. Lin, P. Goyal, R. Girshick, K. He, and P. Dollár, "Focal Loss for Dense Object
Detection." arXiv, Feb. 07, 2018. doi: 10.48550/arXiv.1708.02002.
[51] D. P. Kingma and J. Ba, "Adam: A Method for Stochastic Optimization." arXiv, Jan. 29,
2017. doi: 10.48550/arXiv.1412.6980.
[52] S. Berg *et al.*, "ilastik: interactive machine learning for (bio)image analysis," *Nat. Methods*,

vol. 16, no. 12, Art. no. 12, Dec. 2019, doi: 10.1038/s41592-019-0582-9.

[53] "VIGRA Homepage." Accessed: Aug. 30, 2023. [Online]. Available:
https://ukoethe.github.io/vigra/

[54] B. B. Avants, N. J. Tustison, M. Stauffer, G. Song, B. Wu, and J. C. Gee, "The Insight
ToolKit image registration framework," *Front. Neuroinformatics*, vol. 8, p. 44, Apr. 2014,
doi: 10.3389/fninf.2014.00044.

[55] Y. Gal and Z. Ghahramani, "Dropout as a Bayesian Approximation: Representing Model
Uncertainty in Deep Learning," *ArXiv150602142 Cs Stat*, Oct. 2016, Accessed: Jun. 17,
2021. [Online]. Available: http://arxiv.org/abs/1506.02142

[56] T. C. Lee, R. L. Kashyap, and C. N. Chu, "Building Skeleton Models via 3-D Medial
Surface Axis Thinning Algorithms," *CVGIP Graph. Models Image Process.*, vol. 56, no. 6,
pp. 462–478, Nov. 1994, doi: 10.1006/cgip.1994.1042.

[57] J. Nunez-Iglesias, A. J. Blanch, O. Looker, M. W. Dixon, and L. Tilley, "A new Python
library to analyse skeleton images confirms malaria parasite remodelling of the red blood
cell membrane skeleton," *PeerJ*, vol. 6, p. e4312, Feb. 2018, doi: 10.7717/peerj.4312.

[58] M. E. J. Newman, "Mixing patterns in networks," *Phys. Rev. E*, vol. 67, no. 2, p. 026126,
Feb. 2003, doi: 10.1103/PhysRevE.67.026126.

[59] E. Nader *et al.*, "Blood Rheology: Key Parameters, Impact on Blood Flow, Role in Sickle
Cell Disease and Effects of Exercise," *Front. Physiol.*, vol. 10, p. 1329, Oct. 2019, doi:
10.3389/fphys.2019.01329.

[60] C. B. Schaffer *et al.*, "Two-Photon Imaging of Cortical Surface Microvessels Reveals a
Robust Redistribution in Blood Flow after Vascular Occlusion," *PLOS Biol.*, vol. 4, no. 2, p.
e22, Jan. 2006, doi: 10.1371/journal.pbio.0040022.

[61] P. Virtanen *et al.*, "SciPy 1.0: fundamental algorithms for scientific computing in Python,"
Nat. Methods, vol. 17, no. 3, Art. no. 3, Mar. 2020, doi: 10.1038/s41592-019-0686-2.

[62] "Fitting Linear Mixed-Effects Models Using lme4 | Journal of Statistical Software."
Accessed: Nov. 03, 2023. [Online]. Available: https://www.jstatsoft.org/article/view/v067i01

[63] R. V. Lenth *et al.*, "emmeans: Estimated Marginal Means, aka Least-Squares Means." Oct.
17, 2023. Accessed: Nov. 03, 2023. [Online]. Available: https://cran.r-
project.org/web/packages/emmeans/index.html

[64] J. DeFelipe, L. Alonso-Nanclares, and J. I. Arellano, "Microstructure of the neocortex:
comparative aspects," *J. Neurocytol.*, vol. 31, no. 3–5, pp. 299–316, 2002, doi:
10.1023/a:1024130211265.

[65] F. Dabertrand *et al.*, "PIP2 corrects cerebral blood flow deficits in small vessel disease by
rescuing capillary Kir2.1 activity," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 118, no. 17, Apr.
2021, doi: 10.1073/pnas.2025998118.

[66] P. S. Tsai *et al.*, "Correlations of Neuronal and Microvascular Densities in Murine Cortex
Revealed by Direct Counting and Colocalization of Nuclei and Vessels," *J. Neurosci.*, vol.
29, no. 46, pp. 14553–14570, Nov. 2009, doi: 10.1523/JNEUROSCI.3287-09.2009.

[67] M. Y. Rennie *et al.*, "Vessel tortuousity and reduced vascularization in the fetoplacental
arterial tree after maternal exposure to polycyclic aromatic hydrocarbons," *Am. J. Physiol.-
Heart Circ. Physiol.*, vol. 300, no. 2, pp. H675–H684, Feb. 2011, doi:
10.1152/ajpheart.00510.2010.

[68] J. Steinman, M. M. Koletar, B. Stefanovic, and J. G. Sled, "3D morphological analysis of
the mouse cerebral vasculature: Comparison of in vivo and ex vivo methods," *PLOS ONE*,
vol. 12, no. 10, p. e0186676, Oct. 2017, doi: 10.1371/journal.pone.0186676.

[69] L. Lindvere *et al.*, "Cerebral microvascular network geometry changes in response to
functional stimulation," *NeuroImage*, vol. 71, pp. 248–259, May 2013, doi:
10.1016/j.neuroimage.2013.01.011.

[70] R. Damseh, F. Cheriet, and F. Lesage, "Fully Convolutional DenseNets for Segmentation
of Microvessels in Two-photon Microscopy*," in *2018 40th Annual International Conference*

of the IEEE Engineering in Medicine and Biology Society (EMBC), Jul. 2018, pp. 661–665.
doi: 10.1109/EMBC.2018.8512285.

[71] V. Goodarzi Ardakani, A. M. Gambaruto, G. Silva, and R. Pereira, "A porosity model for
medical image segmentation of vessels," *Int. J. Numer. Methods Biomed. Eng.*, vol. 38, no.
4, p. e3580, 2022, doi: 10.1002/cnm.3580.

[72] M. Livne *et al.*, "A U-Net Deep Learning Framework for High Performance Vessel
Segmentation in Patients With Cerebrovascular Disease," *Front. Neurosci.*, vol. 13, p. 97,
2019, doi: 10.3389/fnins.2019.00097.

[73] M. R. K. Mookiah *et al.*, "A review of machine learning methods for retinal blood vessel
segmentation and artery/vein classification," *Med. Image Anal.*, vol. 68, p. 101905, Feb.
2021, doi: 10.1016/j.media.2020.101905.

[74] "Frontiers | DeepVesselNet: Vessel Segmentation, Centerline Prediction, and Bifurcation
Detection in 3-D Angiographic Volumes | Neuroscience." Accessed: Feb. 21, 2021.
[Online]. Available: https://www.frontiersin.org/articles/10.3389/fnins.2020.592352/full

[75] C. Poon, P. Teikari, M. F. Rachmadi, H. Skibbe, and K. Hynynen, "A dataset of rodent
cerebrovasculature from in vivo multiphoton fluorescence microscopy imaging," *Sci. Data*,
vol. 10, no. 1, Art. no. 1, Mar. 2023, doi: 10.1038/s41597-023-02048-8.

[76] M. I. Todorov *et al.*, "Machine learning analysis of whole mouse brain vasculature," *Nat.*
Methods, vol. 17, no. 4, Art. no. 4, Apr. 2020, doi: 10.1038/s41592-020-0792-1.

[77] J. A. Boero, J. Ascher, A. Arregui, C. Rovainen, and T. A. Woolsey, "Increased brain
capillaries in chronic hypoxia," *J. Appl. Physiol. Bethesda Md 1985*, vol. 86, no. 4, pp.
1211–1219, Apr. 1999, doi: 10.1152/jappl.1999.86.4.1211.

[78] E. Lugo-Hernandez *et al.*, "3D visualization and quantification of microvessels in the whole
ischemic mouse brain using solvent-based clearing and light sheet microscopy," *J. Cereb.*
Blood Flow Metab., vol. 37, no. 10, pp. 3355–3367, Oct. 2017, doi:
10.1177/0271678X17698970.

[79] L.-Y. Zhang *et al.*, "CLARITY for High-resolution Imaging and Quantification of Vasculature
in the Whole Mouse Brain," *Aging Dis.*, vol. 9, no. 2, pp. 262–272, Apr. 2018, doi:
10.14336/AD.2017.0613.

[80] A. Miettinen *et al.*, "Micrometer-resolution reconstruction and analysis of whole mouse
brain vasculature by synchrotron-based phase-contrast tomographic microscopy." bioRxiv,
p. 2021.03.16.435616, Mar. 16, 2021. doi: 10.1101/2021.03.16.435616.

[81] T. Kovacs-Oller, E. Ivanova, P. Bianchimano, and B. T. Sagdullaev, "The pericyte
connectome: spatial precision of neurovascular coupling is driven by selective connectivity
maps of pericytes and endothelial cells and is disrupted in diabetes," *Cell Discov.*, vol. 6, p.
39, 2020, doi: 10.1038/s41421-020-0180-0.

[82] H. Abdelazim *et al.*, "Pericyte Heterogeneity Identified by 3D Ultrastructural Analysis of the
Microvessel Wall," Cell Biology, preprint, Aug. 2022. doi: 10.1101/2022.08.08.503052.

[83] "Characterization of culture from smooth muscle cells isolated from rat middle cerebral
arteries - ScienceDirect." Accessed: Feb. 21, 2021. [Online]. Available:
https://www.sciencedirect.com/science/article/pii/S0040816620301002?via%3Dihub

[84] G. Wang, L. Jacquet, E. Karamariti, and Q. Xu, "Origin and differentiation of vascular
smooth muscle cells," *J. Physiol.*, vol. 593, no. 14, pp. 3013–3030, Jul. 2015, doi:
10.1113/JP270033.

[85] A. P. Del Franco, P.-P. Chiang, and E. A. Newman, "Dilation of cortical capillaries is not
related to astrocyte calcium signaling," *Glia*, vol. 70, no. 3, pp. 508–521, Mar. 2022, doi:
10.1002/glia.24119.

[86] B. Stefanovic *et al.*, "Functional reactivity of cerebral capillaries," *J. Cereb. Blood Flow*
Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab., vol. 28, no. 5, pp. 961–972, May 2008,
doi: 10.1038/sj.jcbfm.9600590.

[87] J. Lee *et al.*, "Opposed hemodynamic responses following increased excitation and

parvalbumin-based inhibition," *J. Cereb. Blood Flow Metab.*, vol. 41, no. 4, pp. 841–856,
Apr. 2021, doi: 10.1177/0271678X20930831.
[88] D. A. Turner, S. Degan, F. Galeffi, S. Schmidt, and A. V. Peterchev, "Rapid, Dose-
Dependent Enhancement of Cerebral Blood Flow by transcranial AC Stimulation in
Mouse," *Brain Stimul. Basic Transl. Clin. Res. Neuromodulation*, vol. 14, no. 1, pp. 80–87,
Jan. 2021, doi: 10.1016/j.brs.2020.11.012.
[89] E. L. Johnson *et al.*, "Characterization of light penetration through brain tissue, for
optogenetic stimulation." bioRxiv, p. 2021.04.08.438932, Apr. 08, 2021. doi:
10.1101/2021.04.08.438932.
[90] S. I. Al-Juboori, A. Dondzillo, E. A. Stubblefield, G. Felsen, T. C. Lei, and A. Klug, "Light
scattering properties vary across different regions of the adult mouse brain," *PloS One*, vol.
8, no. 7, p. e67626, 2013, doi: 10.1371/journal.pone.0067626.
[91] E. Gutiérrez-Jiménez *et al.*, "Effect of electrical forepaw stimulation on capillary transit-time
heterogeneity (CTH)," *J. Cereb. Blood Flow Metab.*, vol. 36, no. 12, pp. 2072–2086, Dec.
2016, doi: 10.1177/0271678X16631560.
[92] A. Lücker, T. W. Secomb, M. J. P. Barrett, B. Weber, and P. Jenny, "The Relation Between
Capillary Transit Times and Hemoglobin Saturation Heterogeneity. Part 2: Capillary
Networks," *Front. Physiol.*, vol. 9, p. 1296, Sep. 2018, doi: 10.3389/fphys.2018.01296.
1122
1123

A Baseline Radius STD

507 μm · 507 μm · 250 μm

B Vesselwise radius Δ(μm)

C Vertexwise radius Δ(μm)

D Vertexwise neuron distance(μm)