MuPET-Flow: Multiple Ploidy Estimation Tool from Flow cytometry data

Gómez-Muñoz C.*, Fischer G.

Laboratory of Computational and Quantitative Biology, Centre National de la Recherche Scientifique (CNRS), Institut de Biologie Paris-Seine (IBPS), Sorbonne Université, Paris F-75005, France

*Corresponding author

Email address: cintia.gomez_munoz@sorbonne-universite.fr

ABSTRACT

Summary:

Ploidy, representing the number of homologous chromosome sets, can be estimated from

flow cytometry data acquired on cells stained with a fluorescent DNA dye. This estimation

relies on a combination of tools that often require scripting, individual sample curation, and

additional analyses. To automate the ploidy estimation for multiple flow cytometry files,

we developed MuPET-Flow—a Shiny graphical user interface tool. MuPET-Flow allows

users to visualize cell fluorescence histograms, detect the peaks corresponding to the

different cell cycle phases, perform a linear regression using standards, make ploidy or

genome size predictions, and export results as figures and table files. The tool was

benchmarked with known ploidy datasets, yielding consistent ploidy results. MuPET-Flow

produces ploidy results comparable to those from other available tools; however, MuPET-

Flow stands out for its user-friendly interface, which allows the whole process to be carried

out in one place, considerably speeding up analysis, particularly in projects involving large

numbers of samples.

Availability and implementation: https://github.com/CintiaG/MuPET-Flow.

INTRODUCTION

Ploidy (n) is the number of homologous chromosome sets in a cell. Cells can be haploid if they have one set (1n), diploid if they have two (2n), or polyploid if they have three or more ($\geq$3n). Polyploidy exists in several branches of the tree of life (Otto and Whitton, 2000), notably in the *Saccharomyces* genus (Peter *et al.*, 2018; Gómez-Muñoz *et al.*, 2021), and in plants (Čertner *et al.*, 2022).

Flow cytometry (FC) allows measuring the fluorescence of single cells stained with a DNA intercalant, such as propidium iodide (PI). When this fluorescence is plotted against the cell count (histogram of fluorescence intensity), usually two peaks corresponding to the different cell cycle stages, G0/G1 and G2, are recognizable. The peaks' fluorescence intensity is proportional to the DNA content, which allows inferring both ploidy and genome size. To achieve this, the peaks' fluorescence of query samples needs to be identified and then compared to that of known standards. Such standards can be run separatedly (external standardization) or within the sample (internal standardization) (Todd *et al.*, 2018; Sliwinska *et al.*, 2022).

The available software tools for FC data analysis can be categorized into proprietary and open-source. Proprietary software comprise those included with the instruments, and independent software such as FlowJo (Becton Dickinson & Company, USA). The latter offers diverse analyses through a graphical user interface (GUI). However, licenses are usually tied to the equipment or require payment. For these reasons, several open-source software options have been developed in R and Python programming languages. These tools are freely available and have the potential to automate the production of publication-

ready images. However, they typically lack a GUI and involve scripting, which might be

daunting for some users.

For the specific case of peaks detection and ploidy estimation, two open-source tools

deserve special mention as they offer a GUI. On one hand, Cytoflow (Teague, 2022),

implemented in Python, is a versatile FC data analysis package. In this software, data can

be loaded and visualized through a GUI. After manual configuration and using the 1D

mixture model, the different peaks' fluorescence can be detected. However, the number of

peaks for the model needs to be specified, and the same number is applied to all samples.

Additionally, for ploidy estimation, the data needs to be exported and analyzed outside the

software (Table S1). On the other hand, flowPloidy, implemented in R (Smith *et al.*, 2018),

is a tool that provides powerful peaks and debris fitting, as well as gating, facilitated by the

inclusion of an R Shiny GUI (Chang *et al.*, 2023). Nevertheless, it still involves scripting

for file uploading, detecting channels, specifying the number of breaks for histograms, and

saving images. Furthermore, failed models need to be inspected and corrected individually.

Finally, flowPloidy is principally intended for calculating plant genome size with internal

standards. If external standars were used, the peaks' means need to be exported and the

ploidy calculated through additional operations (Table S1).

Given the importance of FC in ploidy estimation and the limitations of the existing

tools, we developed MuPET-Flow (Multiple Ploidy Estimation Tool from Flow cytometry

data). As described below, we benchmarked our tool using *Saccharomyces* datasets and

demonstrated its applicability in plant species for ploidy and genome size estimation.

MATERIALS AND METHODS

Flow cytometry data acquisition

S. cerevisiae strains included two natural polyploids CLQCA_17-111 (abbreviated as CRE; 3n) and CH10 (abbreviated as AVQ; 5n), used as test, and four laboratory strains, BY4742 (1n), BY4743 (2n), YPS128_3n (3n), and YPS128_4n (4n), serving as standards. Three biological replicates of yeast cells were cultured in YPD media (Sigma-Aldrich, USA), fixed in ethanol 70% and washed twice with Phosphate-buffered saline solution 1X. Subsequently, the cells were treated with RNAse A (EUROMEDEX, France) at 1 mg/mL at 37°C for 2h, and stained with PI (Thermo Fisher Scientific, USA) at 50 µg/mL. Fluorescence was measured in the MACSQuant VYB system (Miltenyi Biotec, Germany), in channel FL4-A (605-625 nm). Files were gated by FSC-A vs SSC-A and FSC-A vs FSC-H using CytoExploreR (Hammill, 2021). The FC files generated are available on MuPET-Flow's GitHub. *Saccharomyces pastorianus* files were obtained from Gómez-Muñoz *et al.* (2021), while *Solanum pseudocapsicum* files (Čertner *et al.*, 2022) were downloaded from the FlowRepository (ID FR-FCM-Z45W; Spidlen *et al.*, 2012).

Tool description and performance evaluation

MuPET-Flow is implemented in the R package Shiny (Chang *et al.*, 2023) and comprises three tabs (Fig. 1). On the first tab, 'Peaks', multiple local Flow Cytometry Standard (FCS) files can be uploaded through the GUI using internally flowCore (Hahne *et al.*, 2009). MuPET-Flow automatically scans the files for available channels and displays them in and dropdown menu, from which one must be selected. The tool then estimates the number of breaks (minimum 256) and automatically generates a histogram in the chosen channel for

all samples. A 'Local Polynomial Regression Fitting' is applied for histograms smoothing

(default 0.5). Peaks are detected using a local maximum algorithm by employing a sliding

window of a given width (default 50). Optionally, individual files can be selected through a

dropdown menu to display the sample's histogram and peak detection results, and the

smoothing level and window width adjusted by a numeric input menu. The minimun cell

count to call peak can be selected (default 5). The algorithm assumes that only the G0/G1

and G2 populations exist, and only two peaks can be used for the subsequent steps.

However, additional detected peaks can be explored and selected through a checkbox.

In the second tab, 'Regression', MuPET-Flow estimates ploidy or genome size using a

linear regression based on external standards. Therefore, the type of analysis, the number of

standards files, and their corresponding values should be specified. A minimum of two

distinct standards is necessary. The two peaks of the standards are considered for linear

regression, and the ploidy or genome size of both peaks of the test samples is predicted.

The linear regression results are displayed in a text box.

Finally, all generated histograms can be previewed in the third tab called 'Summary'. In

case not all of the samples were individually inspected and an error is found, it is possible

to return to the first tab to review any parameters and selected peaks, provided that

regression is redone as well. All used parameters, such as 'smoothing' and 'window', as

well as the mean ploidy or genome size (presented as decimal and rounded values), are

reported. The histograms can be saved as a figure (PNG or TIFF), and the table as a

comma-separated value file (CSV).

The peaks' fluorescence of the *S. cerevisiea* dataset was also determined using Cytoflow

(Teague, 2022) and flowPloidy (Smith *et al.*, 2018), following their instructions. The

ploidies were obtained by applying the same principle as MuPET-Flow (linear regression). These results were compared with a paired Wilcoxon signed rank exact test. To assess the tool's execution time, we generated a mock dataset by duplicating the files of the six *S. cerevisiae* analyzed strains for all replicates resulting in 36 FCS files (considered a manageable number for visualization) with a total size of 119 MB. Using the R 'system.time' function, the user, system, and elapsed mean times of three runs were obtained for the main application processes. Full computer specifications for developing and testing MuPET-Flow are in Table S2.

RESULTS

Benchmarking with newly generated *S. cerevisiae* FC data

To test our application, we determined the ploidy of two *S. cerevisiae* polyploids, CRE and AVQ, from newly obtained PI-stained cells FC data. The histograms of the two tested strains and four standards were visualized simultaneously (Fig. S1), and the smoothing and window were adjusted for a few samples (Table S3). The estimated peaks' intensities and the inferred ploidy by linear regression are reported in Table S3. The fluorescence intensity varied among the biological replicates; however, there was a good agreement in the mean ploidy for CRE and AVQ in the triplicates. Additionally, the rounded ploidy of the tested strains was identical to that reported by Peter *et al.* (2018).

To emphasize the advantages of MuPET-Flow's pipeline, we compared it with the pipelines of Cytoflow and flowPloidy, focusing on their ability to calculate ploidy over the *S. cerevisiea* dataset. MuPET-Flow automates various steps including the detection of channels and the number of breaks, as well as the calculation of peaks. Furthermore, it

allows the visualization of samples both individually and simultaneously, and to save the

histograms and peaks' intensities as a table, always with the aid of a GUI. Notably,

MuPET-Flow is the only tool that allows the in-app calculation of ploidy (Table S1). We

also examined whether the peaks' fluorescences and ploidies obtained with MuPET-Flow

differed from those obtained with Cytoflow and flowPloidy. We observed a significant

difference ($p < 0.05$) in the peaks' fluorescence between MuPET-Flow and flowPloidy, as

well as between Cytoflow and flowPloidy, but not between MuPET-Flow and Cytoflow.

However, no differences were observed in the inferred plody values across the three tools

(Table S4). Finally, we examined the execution time over the *S. cerevisiea* mock dataset

consisting of 36 files with MuPET-Flow. The total user time for MuPET-Flow's main

processes was less than 2 seconds (Table S2).

App testing with published data in other species

To further assess the app's performance, we challenged it with FC data originating from

different instruments, fluorophore technology, and species. We utilized two previously

published cytometry datasets, one from the yeast *S. pastorianus* (Gómez-Muñoz *et al.*,

2021) for genome size estimation and another from the plant *S. pseudocapsicum* (Čertner *et*

al., 2022) for ploidy determination. In the *S. pastorianus* dataset, we obtained a genome

size of the query strain *S. pastorianus* 790 of 58.64 Mb, close to the 60.1 Mb previously

estimated (Table S5; Gómez-Muñoz *et al.*, 2021). For *S. pseudocapsicum*, multiple peaks

were detected in the fruit skin file. The ploidy of the highest intensity peaks was

investigated, revealing correspondence with the 8n peak previously reported for this tissue

(Table S6; Čertner *et al.*, 2022).

DISCUSSION

MuPET-Flow is an easy-to-use tool that assembles all necessary actions for ploidy

estimation. It is primarily developed to estimate ploidy in yeast; however, it can also be

applied to measure genome size and in other species. MuPET-Flow shares several features

and advantages with existing tools (Table S1), but its novelty consists in combining

features that usually exist separtedly. For instance, MuPET-Flow allows the simultaneous

visualization of multiple samples, as in Cytoflow, and the capability to correct histograms,

somewhat similar to flowPloidy. These characteristics combined facilitates the

identification and curation of the few problematic samples, not feasible in Cytoflow, and

more efficient than flowPloidy which necessitates inspecting every one of them. An

additional feature of MuPET-Flow is the ability to detect multiple peaks due to its local

maxima algorithm, without prior knowledge of the number of peaks. The multiple peaks

present in certain samples can be explored and selected, as observed in the case of

endopolyploidy reported in *S. pseudocapsicum* (Čertner *et al.*, 2022).

Cytoflow, flowPloidy, and MuPET-Flow idividually present several advantages and

disadvantages (Table S1). Nevertheless, MuPET-Flow is an attractive option as it

automates and facilitates many steps in ploidy estimation. It is the only tool capable of

performing this process within the app in a straightforward manner, while reducing the

need for manual curation. Overall, MuPET-Flow minimizes the time required for ploidy

analysis, which is particularly beneficial for projects dealing with a large number of

samples routinely.

ACKNOWLEDGEMENTS AND FUNDING

This work was supported by the Agence Nationale de la Recherche [ANR-20-CE12-0020].

Flow cytometry data acquisition was performed at the IBPS Imaging Facility. The IBPS

Imaging facility is supported by Region-Île-de-France, Sorbonne Université and CNRS.

Conflict of Interest: none declared.

REFERENCES

Čertner, M., Lučanová, M., Sliwinska, E., Kolář, F., & Loureiro, J. (2022). Plant material selection, collection, preservation, and storage for nuclear DNA content estimation. Cytometry Part A, 101(9), 737–748. https://doi.org/10.1002/cyto.a.24482

Chang, W., Cheng, J., Allaire, J., Sievert, C., Schloerke, B., Xie, Y., Allen, J., McPherson, J., Dipert, A., & Borges, B. (2023). shiny: Web Application Framework for R. https://shiny.posit.co/, https://github.com/rstudio/shiny

Gómez-Muñoz, C., García-Ortega, L. F., Montalvo-Arredondo, J., Pérez-Ortega, E., Damas-Buenrostro, L. C., & Riego-Ruiz, L. (2021). Long-insert clone experimental evidence for assembly improvement and chimeric chromosomes detection in an allopentaploid beer yeast. G3 Genes|Genomes|Genetics, 11(7), jkab088. https://doi.org/10.1093/g3journal/jkab088

Hahne, F., LeMeur, N., Brinkman, R. R., Ellis, B., Haaland, P., Sarkar, D., Spidlen, J., Strain, E., & Gentleman, R. (2009). flowCore: A Bioconductor package for high throughput flow cytometry. BMC Bioinformatics, 10, 106. https://doi.org/10.1186/1471-2105-10-106

Hammill, D. (2021). CytoExploreR: Interactive Analysis of Cytometry Data. R package. https://github.com/DillonHammill/CytoExploreR

Otto, S. P., & Whitton, J. (2000). Polyploid incidence and evolution. Annual Review of Genetics, 34(1), 401–437. https://doi.org/10.1146/annurev.genet.34.1.401

Peter, J., Chiara, M. D., Friedrich, A., Yue, J.-X., Pflieger, D., Bergström, A., Sigwalt, A., Barre, B., Freel, K., Llored, A., Cruaud, C., Labadie, K., Aury, J.-M., Istace, B., Lebrigand, K., Barbry, P., Engelen, S., Lemainque, A., Wincker, P., Liti, G., Schacherer, J. (2018). Genome evolution across 1,011 *Saccharomyces cerevisiae* isolates. Nature, 556(7701), 339–344. https://doi.org/10.1038/s41586-018-0030-5

Sliwinska, E., Loureiro, J., Leitch, I. J., Šmarda, P., Bainard, J., Bureš, P., Chumová, Z., Horová, L., Koutecký, P., Lučanová, M., Trávníček, P., & Galbraith, D. W. (2022). Application-based guidelines for best practices in plant flow cytometry. Cytometry Part A, 101(9), 749–781. https://doi.org/10.1002/cyto.a.24499

Smith, T. W., Kron, P., & Martin, S. L. (2018). flowPloidy: An R package for genome size and ploidy assessment of flow cytometry data. Applications in Plant Sciences, 6(7), e01164. https://doi.org/10.1002/aps3.1164

Spidlen, J., Breuer, K., Rosenberg, C., Kotecha, N., & Brinkman, R. R. (2012). FlowRepository: A resource of annotated flow cytometry datasets associated with peer-reviewed publications. Cytometry Part A, 81A(9), 727–731. https://doi.org/10.1002/cyto.a.22106

Teague, B. (2022). Cytoflow: A Python Toolbox for Flow Cytometry (p. 2022.07.22.501078). bioRxiv. https://doi.org/10.1101/2022.07.22.501078

223   Todd, R. T., Braverman, A., & Selmecki, A. (2018). Flow Cytometry Analysis of Fungal

224   Ploidy. Current Protocols in Microbiology, 50(1), e58.

225   https://doi.org/10.1002/cpmc.58

FIGURES AND LEGENDS

Fig. 1. MuPET-Flow: A Shiny app for estimating ploidy in multiple samples. The MuPET-Flow workflow is divided into three tabs: 'Peaks', 'Regression', and 'Summary'. The main functionalities consist of User Interface (UI) inputs, processes executed in the Server, and UI outputs. Adjusting the inputs in the dashed box is optional. Created with BioRender.com.