1
Extracellular ATP is a signalling molecule in bacteria
3

Sophie Tronnet[1,2,3], Vikash Pandey[1,2,3], Miriam Lloret-Berrocal[1,2,3], Mario Pérez-del-
Pozo[1,2,3], Niklas Söderholm[1,4], Carlos Hernández-Ortego[1,2,3], Oliver Billker[1,2,3],
Anders Nordström[5,6] and Andrea Puhar[1,2,3,7]
[1]The Laboratory for Molecular Infection Medicine Sweden (MIMS), 901 87 Umeå,
Sweden
[2]Umeå Centre for Microbial Research (UCMR), 901 87 Umeå, Sweden
[3]Department of Molecular Biology, Umeå University, 901 87 Umeå, Sweden
[4]Department of Chemistry, Umeå University, 901 87 Umeå, Sweden
[5]Umeå Plant Science Centre, 901 87 Umeå, Sweden
[6]Swedish Metabolomics Centre, 907 36 Umeå, Sweden
[7]Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine,
Dentistry and Biomedical Sciences, Queen's University Belfast, BT9 7BL Belfast,
United Kingdom

Abstract
In animals and plants extracellular ATP (eATP) functions as signalling molecule and
regulates the immune response. During inflammation intestinal bacteria are exposed
to elevated eATP originating from the mucosa. Whether bacteria respond to eATP is
unclear. Here we show that non-pathogenic *Escherichia coli* responds to eATP at
physiologically relevant concentrations by modifying its transcriptional and metabolic
landscape. The use of a promoter library showed that the response to eATP is time-,
dose- and medium-dependent. Genes related to lipid, amino acid, or vitamin
metabolism were regulated. Metabolomics showed that eATP triggers enrichment of
molecules with bioactive properties on the host or bacteria. Combined genome-scale
modelling highlighted the global metabolic modifications. Moreover, eATP altered the
sensitivity to antibiotics and antimicrobial peptides. Finally, in pathogens eATP
controls the expression of virulence and fitness factors. Our results indicate that eATP
is a signalling molecule in prokaryotes which globally regulates physiology,
antimicrobial resistance, and virulence.

Introduction
Extracellular ATP (eATP) has emerged as a key signalling molecule in lower and
higher eukaryotes, including protozoa, animals, plants, and fungi (Li *et al*, 2021;
Medina-Castellanos *et al*, 2014; Medina-Castellanos *et al*, 2018; Pham *et al*, 2020;
Tanaka *et al*, 2014; Verkhratsky & Burnstock, 2014). In animals and plants eATP
regulates physiological responses (Tanaka *et al.*, 2014; Verkhratsky & Burnstock,
2014). Importantly, eATP controls damage, stress or inflammatory responses in
protozoa, animals, plants and fungi upon tissue injury or alteration (Bours *et al*, 2006;
Idzko Ferrari & Eltzschig, 2014; Medina-Castellanos *et al.*, 2014; Medina-Castellanos
et al., 2018; Pham *et al.*, 2020; Tanaka *et al.*, 2014; Verkhratsky & Burnstock, 2014).
In contrast, the existence of purinergic signalling in the earlier forms of life, Bacteria
and Archaea, remains largely unexplored (Verkhratsky & Burnstock, 2014).
Whether bacteria can respond to eATP is unclear. Only a handful of studies addressed
the effect of eATP on bacteria at the phenotypic level. In *Streptomyces* exogenous
ATP promoted differentiation, possibly via cytosolic ATP-consuming enzymes (Li *et*
al, 2016; Li *et al*, 2008; Meng *et al*, 2011). Other studies investigated the role of added

ATP in biofilms and/or growth in several organisms, partially with opposing results. Effects on biofilms were linked to release of biofilm-composing DNA upon eATP-induced lysis (Xi & Wu, 2010), inhibition of twitching motility by reducing surface-assembled Type IV pili via eATP hydrolysis (Nolan *et al*, 2015), chelation of biofilm-composing cations by eATP (Ding & Tan, 2016), or not further dissected (Baeza & Mercade, 2021). Effects on growth were not explained (Abbasian *et al*, 2019; Li *et al*, 2011; Malo *et al*, 2014; Xi & Wu, 2010), except for eATP-dependent chelation of the nutrient iron in one instance (Tatano *et al*, 2015). However, whether eATP acts as signalling molecule remains unsolved. Finally, ATP supplementation increased survival of non-growing bacteria (Mempin *et al*, 2013).

Until recently, a contribution of eATP to host-microbe interactions was not appreciated. However, eATP is present in the nano- to micromolar range in culture supernatants (Atarashi *et al*, 2008) or homogenates (Chiaranunt *et al*, 2023) of murine faeces and many isolated bacteria (Abbasian *et al.*, 2019; Baeza & Mercade, 2021; Chiaranunt *et al.*, 2023; Ding Quah & Tan, 2016; Hironaka *et al*, 2013; Ihssen *et al*, 2021; Ivanova *et al*, 2006; Iwase *et al*, 2010; Mempin *et al.*, 2013; Nolan *et al.*, 2015; Qiu *et al*, 2019), during osmotic shock (Berrier *et al*, 1992), treatment with antimicrobial peptides (Alvarez *et al*, 2017; Valat *et al*, 2003) or antibiotics (Abbasian *et al.*, 2019; Ihssen *et al.*, 2021; Nilsson, 1978; Proietti *et al*, 2019; Yu Lomakina & Ugarova, 2022). During homeostasis, eATP is found in the gut lumen at low concentration (Kusu *et al*, 2013; Malo *et al.*, 2014; Patel, 2014; Perruzza *et al*, 2017; Proietti *et al.*, 2019; Puhar *et al*, 2013; Scott *et al*, 2021; Tani *et al*, 2021). Importantly, intestinal eATP increases during bacterial, viral, or parasitic infection and protist colonisation. Indeed, enteropathogenic bacteria trigger connexin-dependent secretion of ATP from the intestinal epithelium, *via* activation of the mechanosensor PIEZO1, as an early alert response, which induces potent gut inflammation (Puhar *et al.*, 2013; Tadala *et al*, 2022). Rotavirus also initiates epithelial purinergic signalling (Chang-Graham *et al*, 2020). During Shiga toxin-producing enterohaemorrhagic *Escherichia coli* (Arvidsson *et al*, 2022) or *Citrobacter rodentium* infection (Chiaranunt *et al.*, 2023) and *Tritrichomonas musculis* colonisation (Chiaranunt *et al*, 2022) eATP is present in faeces. Helminths induce ATP release in the gut (Chen *et al*, 2022). Further, intestinal eATP rises during experimental inflammatory bowels diseases (IBD) (Di Virgilio Pinton & Falzoni, 2016; Kurashima *et al*, 2012; Scott *et al.*, 2021; Wan *et al*, 2016) and after irradiation in graft-versus-host-disease (Wilhelm *et al*, 2010) and small intestinal fibrosis (Takemura *et al*, 2018). Ultimately, any process causing cell damage results in ATP release (Verkhratsky & Burnstock, 2014). Together, this implies that in the gut lumen, during infection and inflammation, commensal and pathogenic microbes are exposed to heightened concentrations of eATP, which may serve as environmental signal.

Here we show that eATP regulates the transcriptional and metabolic landscapes of non-pathogenic *E. coli* at physiologically relevant concentrations. The response to eATP presents the characteristic of bacterial transcriptional and metabolic networks, and result in production of bioactive compounds and altered susceptibility to antimicrobials. We further show that several gut pathogens and pathobionts respond to eATP by regulating expression of virulence or fitness factors. Together, our results indicate that eATP is a signalling molecule that modifies the physiology of harmless and pathogenic bacteria, allowing them to adapt to a changing environment.

Results
eATP triggers a robust transcriptional response in non-pathogenic *E. coli*

To investigate the ability of bacteria to sense and respond to eATP, we tested whether eATP induces a transcriptional response using a plasmid-based fluorescent reporter library in non-pathogenic *E. coli* K-12 MG1655 (Zaslaver *et al*, 2006). This collection can unravel the dynamic activity of approximately 2000 promoters (75% of *E. coli* promoters), allowing accurate quantification of gene expression induction or repression seen as GFP fluorescence normalised to the number of bacteria.

We screened this library for a response to eATP in rich (LB) and minimal medium (M9) upon treatment with 1 mM ATP for 3 h (Fig EV1A). Addition of ATP led to up- or down-regulation of promoters in both media, with a higher fraction modulated in M9 than in LB (Fig 1A-D, examples in Fig EV1B-G, Table EV1). We identified 291 and 476 modulated promoters in LB and M9, respectively (Fig 1E), with 121 promoters modulated by eATP in both media, of which 46 genes were upregulated and 27 downregulated (Fig 1F, G). These screens were highly reproducible (Fig EV1H), and addition of ATP neither altered bacterial growth (Fig EV1J-L) nor caused changes in pH.

To investigate whether the transcriptional response to eATP in *E. coli* MG1655 is dose- and time-dependent, we selected three reporters that were significantly upregulated by eATP at 3 h in LB and/or M9: P*ybjX*, controlling the expression of a DUF535 domain-containing protein; P*rstA*, controlling the DNA-binding transcriptional regulator of the two-component system RstAB; and P*gadW*, controlling a DNA-binding transcriptional dual regulator. We applied increasing concentrations of eATP and measured the promoter activity in LB (Fig 1H-J) and M9 (Fig 1K-M) over time. This showed that the response to eATP was dose-dependent for all tested promoters in both media after 60 min of stimulation or longer, except for the promoter activity of *gadW*, which never increased with the addition of eATP in M9 (Fig 1L). We confirmed the activity of two upregulated promoters by constructing chromosomal fusions with superfolder GFP. Both endogenous promoters of *gadW* and *cbpA* (which controls the expression of a DnaK co-chaperone) were significantly upregulated (fold change≈2) 5 and 30 min after eATP addition, respectively (Fig EV1M-O).

To understand what the response to eATP in non-pathogenic *E. coli* MG1655 is, we analysed the function of eATP-modulated genes. Several of these genes encode unknown or poorly characterized proteins (Table EV1). KEGG analysis revealed 13 and 15 significantly enriched pathways in LB and M9, respectively, including the metabolism of fatty acids, vitamins, co-factors and proteinogenic amino acids (Fig 2A, b and Table EV1). Quorum sensing, riboflavin and nitrogen metabolism were enriched in both media. In contrast, the metabolism of different types of vitamins, amino acids and fatty acids was predicted in LB and M9. We further analysed the differentially expressed genes (DEGs) using the web-based tool Panther (Mi *et al*, 2013), which functionally classified them into four general gene ontology (GO) categories: biological processes, cellular components, molecular function, and protein class (Fig 2C, Table EV1). The DEGs modulated in LB and M9 upon stimulation with eATP were mainly found in the subcategories: binding, catalytic activity, cellular processes, metabolic process, cellular anatomical entity, and metabolite interconversion enzyme and to a lesser extent, transporter and gene-specific transcriptional regulator, confirming the prediction that eATP impacts bacterial gene regulation and metabolism.

In summary, these results show that the intestinal non-pathogenic bacterium *E. coli* responds to eATP at physiologically relevant concentrations by modifying its transcriptional landscape in a medium-, dose- and time-dependent manner.

eATP triggers a strong metabolic response in non-pathogenic *E. coli*

Our data indicated that the gene expression programmes induced by eATP have a
strong impact on bacterial physiology and metabolism. To validate this finding, we
assessed the changes in intracellular (pellet) and extracellular (supernatant)
metabolites of bacteria grown in LB or M9 upon exposure to eATP by untargeted
metabolomics using liquid chromatography–mass spectrometry (LC–MS/MS). We
only measured extracellular metabolites in M9 because LB cannot be employed in LC-
MS/MS.
In unstimulated *E. coli* MG1655, we detected 57 and 53 intracellular metabolites in LB
and M9, respectively, with 51 present in both media. Further, we identified 89
extracellular metabolites, with 61 only present in the supernatant but not the pellets,
while 23 metabolites were found in all conditions (Fig 3A and Table EV2). We then
compared the metabolome of bacteria stimulated with 1 mM ATP for 3 h with time-
matched unstimulated controls. Partial least square-discriminant analysis (PLS-DA)
distinguished untreated and eATP-treated *E. coli* (Fig 3B-D), indicating a clear effect
of eATP on the metabolome. Univariate analysis of these data revealed that eATP
triggers a potent shift in the metabolic landscape (Fig 3E-J, Table EV2), with 5–16%
of detected metabolites significantly modulated (≥ 1.5 FC, $P \leq 0.05$) in response to
eATP. The metabolic changes upon ATP addition in M9 were more substantial than
in LB with 9/53 and 4/57 significantly modulated metabolites in pellets, respectively
(Fig EV2A-D, Table EV2), confirming the medium-dependent effect of eATP we
observed in transcriptional assays (Fig 1). We found 11/89 significantly eATP-
modulated metabolites in the supernatant (Fig EV2E-F, Table EV2).
Overall, the most recurrently modulated metabolites were intermediates of fatty acid,
co-factor, small organic acid, pentose phosphate, vitamin, nucleotide and nucleoside,
proteinogenic and non-proteinogenic amino acid biosynthesis, and redox regulation
(Fig 3E-J, Table EV2). This is in keeping with the predicted regulation of metabolic
pathways based on the eATP-dependent transcriptional changes. To identify the most
characteristic metabolites in LB and M9, we calculated VIP scores, ranking the
metabolites according to their influence on the separation of the treatment groups (Fig
EV2B, D, F). To analyse these differences in depth, we constructed metabolic models
(see below). Among the metabolites not integrated in the metabolic model, ophthalmic
acid was significantly decreased in LB (Fig 3H, Table EV2). In M9, cAMP and lysine
were strongly decreased, while PC-caproyl (a membrane lipid) was upregulated by
eATP (Fig 3I, Table EV2). Further, the analysis showed that the upregulated biogenic
amine tyramine was the top ranked metabolite in the VIP analysis in the M9
supernatant (Fig 3H, Fig EV2E-F). Of note, allantoin, AMP, threonate and GTP also
accumulated in the supernatant.
To test whether the metabolic response to eATP is dose-dependent, alike the
transcriptional response, as proof of principle we assessed the changes in intracellular
metabolites of *E. coli* MG1655 grown in M9 upon stimulation with 100 µM ATP. This
showed that the metabolites in treated bacteria were easily discriminated from the
unstimulated control (Fig EV3A) and revealed shifts in the abundance of several
compounds from proteinogenic and non-proteinogenic amino acid, fatty acid, small
organic acid, pentose phosphate, nucleotide and nucleoside, cell envelope, co-factor
metabolism and redox regulation (Fig EV3B-F, Table EV2). Intracellular metabolites
in bacteria treated with 100 µM or 1 mM eATP were only partially different (Fig EV3G-
H). Two compounds (cAMP and L-lysine – Fig 3I and Fig EV3E, Table EV2) were
significantly negatively modulated by eATP at both concentrations, while
arginosuccinic acid and L-pipecolate decreased at 100 µM eATP (Fig EV3E).
Interestingly, tyramine accumulated in 100 µM eATP-treated bacteria, as observed for

1 mM eATP. These results show that the response to eATP is concentration-
dependent at quantitative level (seen as duration and strength of transcriptional
induction) and also leads to cells to different metabolic states.
Altogether, our results show that eATP triggers a strong metabolic response in a non-
pathogenic bacterium in a medium- and dose-dependent manner.
205
Comprehensive genome-scale modelling reveals medium-specific differences
in *E. coli* metabolic capabilities induced by eATP.
To gain a comprehensive understanding of the key properties of eATP-modulated
metabolic networks, we performed genome-scale metabolic modelling by integrating
transcriptional and intracellular metabolomic data from *E. coli* MG1655 exposed to
1mM ATP using REMI (Pandey Hadadi & Hatzimanikatis, 2019).
The analysis revealed that in LB 78% (141/181) and in M9 59% (138/233) of the omics-
based constraints were consistently integrable with the model (Table EV3), indicating
that nutrient-rich medium provides more metabolic flexibility. We performed differential
flux analysis and identified 316 and 430 reaction fluxes that were significantly (*P* value
<1e-3) modulated for LB and M9, respectively (Table EV3). ATP stimulation enriched
the following metabolic subsystems in both LB and M9: valine, leucine and isoleucine
metabolism; murein synthesis and recycling; and cell envelope biosynthesis.
Glycerophospholipid metabolism was modulated in LB, while in M9 the core
metabolism (pentose phosphate pathway and citric acid cycle), non-essential amino
acid metabolism (proline and arginine), purine and pyrimidine nucleotide biosynthesis
were modulated (Fig 4A-B, Table EV3). These analyses further highlight the medium-
specific differences in metabolic capabilities induced by eATP.
To understand the downstream effects of transcriptional, metabolite and reaction
perturbations, we calculated the abundance of 68 Biomass Building Blocks (BBBs),
which are important indicators of bacterial physiology, with focus on amino acids, lipids
and co-factors (Table EV3). eATP caused marked modulation (>1.5-fold) of 6 BBBs in
LB and 5 in M9 (Fig 4C). In both LB and M9, the production of L-leucine increased,
while L-lysine and KDO(2)-lipid IV(A), an essential component of lipopolysaccharide,
were overproduced only in LB. In M9 the production of L-valine and L-isoleucine
increased but decreased in LB. Oxidized flavin adenine dinucleotide (FAD) and
riboflavin production diminished in M9. These findings indicate that BBBs synthesis,
and thereby the chemical nature of bacterial structures such as the cell envelope, is
altered by eATP in a medium-dependent manner.
Finally, to obtain a metabolic snapshot of the global changes induced by eATP, we
identified subnetworks that contain both significantly modulated reaction fluxes and
metabolites or genes (Fig 4D and 5). This very stringent analysis highlights pathways
that are consistently altered at many levels. Most reactions were increased upon
addition of eATP in both media (seen in iterative quantitative simulations shown in
boxes), with a few exceptions: UMPK (see figure legend for abbreviations) and INSH
in LB (Fig 5) and GLUPRT in M9 (Fig 5). eATP modulated the nucleotide salvage
network in LB and M9, but the extent and pattern of deregulation differed. Purine-
nucleoside phosphorylase-dependent reactions (PUNP2, PUNP5 encoded by *b4384*,
and PUNP6 encoded by *b2407*) play a vital role in recycling purines. Expression of
b2407 was not altered by eATP in M9, while *b4384* was downregulated in LB (Table
EV1). In LB only inosine, UMP, uridine, and few reactions (PUNP5 and INSH) were
impacted by eATP. By contrast, in M9 the network showed modular deregulation, with
several metabolites (inosine, adenosine, adenine, UMP, uracil) and reactions
(PUNP2, PUNP6, DADA, and AMPN) being modulated by eATP. UMP was

upregulated in both media, but produced through UPPRT reaction in M9, whilst through NTPP8, UMPK or URIK1 in LB. Moreover, N-acetyl-glutamate, ADP and oxidized glutathione accumulated in M9 in presence of eATP. While there is no direct interaction between N-acetyl-glutamate and oxidized glutathione, they may influence each other indirectly, as the reaction fluxes of GTHS and ACGS were higher in M9 in presence of eATP. Further, D-ribose-5 phosphate increased, together with directly or indirectly linked reactions associated with nucleotide salvage pathways (AMPN), pentose-phosphate pathway (TKT1), co-factor and prosthetic group biosynthesis (NNAM) as well as purine/pyrimidine biosynthesis (GLUPRT). Of note, ATP addition led to a slight but significant decrease of nicotinamide (vitamin B3), a precursor of the essential redox co-factor nicotinamide adenine dinucleotide (NAD).

These results highlight the potent impact of eATP on whole metabolic pathways the metabolism of non-pathogenic *E. coli*, including on the production of crucial cellular components and of compounds with known bioactive properties on bacteria or the host. Moreover, these findings underscore the importance of considering the cellular context when analysing changes in metabolic networks induced by eATP.

eATP alters antibiotics and antimicrobial peptides susceptibility

During infection, bacteria may be exposed to eATP in conjunction with environmental insults such as antibiotic therapy and antimicrobial peptides from the host. Our results indicate that eATP triggers global physiological changes, some of which modify the cell envelope in non-pathogenic *E. coli*. Therefore, we tested the sensitivity of eATP-pretreated bacteria to selected antibiotics and antimicrobial peptides of diverse origin and various modes of action (Table EV5). Non-pathogenic *E. coli* showed enhanced susceptibility to aztreonam, mecillinam, and gentamicin and reduced susceptibility to ceftazidime and imipenem (Fig 6A). Bacteria treated with eATP showed a marked increased susceptibility to all tested antimicrobial peptides except CAP-18 (Fig 6B-H and Fig EV4A-G). Altogether, eATP alters the sensitivity to antibiotics or antimicrobial peptides.

The ability to respond to eATP is present among clinically relevant bacteria

We explored whether the capacity to respond to eATP by modifying the transcriptional landscape is present in bacteria other than non-pathogenic *E. coli* MG1655. We selected a panel of harmless, pathobiontic or pathogenic Enterobacteriaceae: non-pathogenic *E. coli* BW25113, adherent invasive *E. coli* (AIEC) LF82 (pathobiontic), probiotic *E. coli* Nissle 1917, enteropathogenic *E. coli* (EPEC), enteroinvasive *E. coli* (EIEC), pathogenic *Shigella flexneri* (enteroinvasive), S*almonella enterica* serovar Typhimurium (enteroinvasive) and *Klebsiella pneumoniae*.

We transformed these organisms with transcriptional fluorescent reporter plasmids carrying promoters from *E. coli* MG1655 that were activated by eATP (Fig 1, Fig EV1 and Table EV1). To estimate the chances of *E. coli* MG1655 promoters to function in these bacteria, we performed multiple sequence alignments and sequence identity analyses (Fig EV5A). This revealed a high percentage of identity among all *E. coli* strains and *S. flexneri*, whereas *S*. Typhimurium and *K. pneumoniae* showed lower identity compared to *E. coli* MG1655. All tested Enterobacteriaceae displayed a transcriptional response to eATP in LB and M9 that was within the same range as *E. coli* MG1655, with a few exceptions (Fig 6I, Fig EV5B); P*gadW-gfp* in *E. coli* Nissle 1917 was only activated in M9, while we detected no activation in *S. flexneri* or *S*. Typhimurium. None of the reporters responded to eATP in *K. pneumoniae* grown in

299 M9. We omitted experiments with *S. flexneri*, EIEC and *S.* Typhimurium in M9 owing
300 to their poor growth.
301 These results indicate that eATP is a signalling molecule for medically relevant
302 bacteria. While in Enterobacteriaceae the ability to respond to eATP is conserved, the
303 type of response is strain- and species-specific.
304
305 The expression of virulence factors is regulated by eATP
We showed that eATP elicits transcriptional responses in pathogenic bacteria,
exploiting target genes that are mostly present in both non-pathogenic *E. coli* and the
pathogens. To determine whether eATP could also regulate strain-specific virulence
genes, we cloned the promoter of selected virulence and fitness factors into the
transcriptional reporter plasmid, which we introduced into the bacteria of origin, and
monitored promoter activity upon eATP stimulation.
We selected the promoters of *S.* Typhimurium *mgtC*, *hilA*, and *ssrB*. In *Salmonella*,
pathogenicity islands 1 and 2 (SPI-1 and SPI-2) have crucial roles in early and later
stages of infection, respectively (Lou *et al*, 2019). HilA positively controls the
expression of SPI-1 genes (Ellermeier Ellermeier & Slauch, 2005), while the two-
component system SsrAB regulates the expression of several SPI-2 operons (Deiwick
et al, 1999). MgtC aids bacterial survival inside macrophages (Choi *et al*, 2019).
Addition of ATP significantly upregulated P*mgtC* and downregulated P*hilA* in a dose-
dependent manner (Fig 6J), whilst P*ssrB* was not regulated (Fig EV5C).
E. coli Nissle 1917 contains a *pks* genomic that mediates the synthesis of colibactin,
a genotoxin (Nougayrède *et al*, 2006). ClbR is a key transcriptional regulator of the
pks island (Wallenstein *et al*, 2020). Stimulation with high concentrations of eATP
resulted in a decrease of the *clbRA* promoter activity (Fig 6K).
We also tested the promoter of *E. coli* LF82 *qseBC*. This strain was isolated from a
mucosal lesion of a Crohn´s disease patient (Darfeuille-Michaud *et al*, 1998). It
adheres to and invades enterocytes, and replicates within macrophages. The
pathogenicity of LF82 and that of many *E. coli* depends on the quorum-sensing two-
component system QseBC (Rooks *et al*, 2017). Upon treatment with eATP, the
promoter activity of LF82 *qseBC* significantly decreased at all eATP concentrations
except 5 mM (Fig 6L).
Finally, we tested P*ler* and P*grlA* from EPEC E2348/69. EPEC encodes a
pathogenicity island, the locus of enterocyte effacement (LEE). LEE is activated by
the master regulators Ler and GrlA (Platenkamp & Mellies, 2018). While the activity of
P*grlA* strongly diminished in a dose-dependent manner (Fig 6M), P*ler* remained
unaltered (Fig EV5D).
Together, these findings indicate that eATP can regulate virulence and fitness in a
broad panel of clinically important intestinal bacteria.
338
339 Discussion
In lower and higher eukaryotes, eATP is a universal signalling molecule that acts as
danger signal (Bours *et al.*, 2006; Tanaka *et al.*, 2014; Verkhratsky & Burnstock, 2014).
Whether eATP functions as signalling molecule in prokaryotes has been a
longstanding question. Here we show that heightened eATP elicits extensive changes
in physiology and virulence by reprogramming transcriptional and metabolic
landscapes in a panel of Enterobacteriaceae. In non-pathogenic *E. coli*, this alters the
envelope composition and the sensitivity to antibiotics or antimicrobial peptides.
In eukaryotes, danger signals positively and negatively regulate defence mechanisms,
ultimately promoting survival. For example, in mammals eATP initiates immunity and

also promotes healing (Bours *et al.*, 2006). But how would the activity of a danger
signal manifest in bacteria? The eATP-mediated loss of viability upon exposure to
antimicrobial peptides and some antibiotics is difficult to reconcile with the expected
protective function of a bacterial danger signal. Rather, this effect points to an
additional host defence mechanism, where ATP released by stressed cells directly
weakens bacteria, besides inducing inflammation (Puhar *et al.*, 2013). Of note,
elevated eATP augments the production of gut antimicrobial peptides during
experimental shigellosis (Puhar *et al.*, 2013), setting the conditions to exploit this
enhanced sensitivity. Importantly, our data indicate that the potency of antimicrobials
may be altered in inflammatory environments, where eATP is abundant. This is of
therapeutic concern for the antibiotics showing reduced efficacy in the presence of
eATP.
By contrast, up- or downregulation of virulence and fitness factors downstream of
eATP seems in line with a function as danger signal to bacteria. The dose-dependency
of the response indicates that microbes may utilise eATP as a site- and infection stage-
characteristic environmental signal, which provides an additional regulatory layer to
finetune species-specific adaptation. Indeed, we speculate that different niches in the
gut have distinctive eATP concentrations during inflammation, akin to oxygen tension
gradients (Marteyn *et al*, 2010). In healthy tissue the extracellular space is virtually
devoid of ATP, while injured cells release up to millimolar ATP from their cytosolic 3-
10 mM ATP pool (Bours *et al.*, 2006). Hence, eATP is likely most abundant at its site
of release, the mucosa, and gradually diffuses into the gut lumen. Thus, harmless
microbes in the lumen might be exposed to lower eATP than pathobionts or pathogens
adhering to the epithelium. Invasive bacteria encounter secreted ATP during cellular
uptake (Tadala *et al.*, 2022) and peaking ATP after vacuolar escape (Malik-Kale
Winfree & Steele-Mortimer, 2012). Accordingly, we generally observed
downregulation of early-stage and upregulation of late-stage virulence genes, when
eATP is highest. However, a larger number of virulence factors must be tested.
Moreover, eATP is a ubiquitous inflammatory mediator, for example in the airways
(Idzko *et al*, 2007). Notably, *K. pneumoniae* is carried in the respiratory and
gastrointestinal tracts (Raffelsberger *et al*, 2021). Finally, heightened eATP in dense
microbial communities could regulate bacterial adaptation, for instance in survivors of
bacteriophage- or antibiotic-induced death (Abbasian *et al.*, 2019; Ihssen *et al.*, 2021;
Nilsson, 1978; Proietti *et al.*, 2019; Yu Lomakina & Ugarova, 2022). Altogether, this
warrants comprehensive investigation of the impact of eATP on diverse organisms in
various environments and its potential role as danger signal in prokaryotes.
In eukaryotes, eATP acts as a signal in the micro- to millimolar range (Bours *et al.*,
2006; Pham *et al.*, 2020; Tanaka *et al.*, 2014; Verkhratsky & Burnstock, 2014). We
provide evidence supporting a bacterial response at these eATP levels. As in
eukaryotes, we found that eATP modifies bacterial gene expression. In non-
pathogenic *E. coli*, eATP triggered only partially overlapping promoter activities in
different media. In fact, the output to an external cue is a global function of distinct
input components: the cue and the cellular state. The latter strongly depends on the
environment (Gianchandani *et al*, 2009), thus giving rise to media-specific responses.
Further, the existence of early and late gene expression programmes reflects the
hierarchical structure of transcriptional regulatory systems, which are comprised of
environmental stimuli at the top and multiple lower levels of transcriptional regulators
and target genes (Gianchandani *et al.*, 2009).
Likewise, in different media eATP induced only partially overlapping shifts in
compounds of diverse chemical nature and genome-scale metabolic networks. As for

transcriptional states, several regulatory mechanisms influence metabolic fluxes, including reactant concentration (depending on the environment and cellular state), enzyme abundance (depending on gene expression), and enzyme kinetic parameters (Covert & Palsson, 2002; Gerosa & Sauer, 2011), thus explaining medium-specific metabolite changes. Introduction of ATP as a substrate may contribute to the observed changes in intra- and extracellular purines. While eATP is neither transported into *E. coli* (Mempin *et al.*, 2013; Tjaden *et al*, 1998) nor serves to phosphorylate bacterial molecules (Mempin *et al.*, 2013), its components may be internalised after ATP hydrolysis. Indeed, carbon flux mapping in intracellular *S.* Typhimurium showed that ribose and nucleobases partly and fully originate from the host, respectively, rather than from *de novo* synthesis (Mitosch *et al*, 2023). Altogether, our data reveal that the eATP response presents the characteristics of bacterial gene regulatory networks and metabolic fluxes.

Among the metabolites modulated by eATP, we detected intracellular cAMP. This second messenger is produced downstream of environmental signals and regulates carbon metabolism, stress response pathways and virulence (McDonough & Rodriguez, 2011). Further, eATP stimulated the production of metabolic intermediates or end products with described bioactive properties on microorganisms or the host (Culp & Goodman, 2023; Krautkramer Fan & Bäckhed, 2021), for example antioxidants and the neurotransmitter tyramine (a complete list is provided in Table EV2). We found modulation of vitamins at the transcriptional level, but not by metabolomic analysis, possibly due to the extraction protocol (Krautkramer Fan & Bäckhed, 2021). Cross-feeding of vitamins is common among intestinal bacteria (Culp & Goodman, 2023). Likewise, quorum sensing was transcriptionally upregulated, but we didn't detect autoinducers. Altogether, this points to the activation of intracellular signalling pathways in bacteria downstream of sensing eATP, which culminate in the production of compounds potentially regulating intra- and inter-kingdom cell-to-cell communication.

The mechanism by which bacteria sense and respond to eATP remains to be elucidated. A molecular understanding of bacterial eATP signalling is key to predict which species encode the ability to utilise eATP as environmental cue and to assess, under well-controlled conditions, the role of eATP-mediated host-bacteria interactions in complex systems such as the gut. Our discovery of eATP-dependent transcriptional and metabolic regulation of bacteria, together with its molecular dissection, may open unfathomed opportunities for the prevention and treatment of infection and inflammation.

In conclusion, we show that eATP rewires physiology, antimicrobial sensitivity, and virulence in medically important intestinal bacteria. Thus, eATP emerges as a potent signalling molecule in prokaryotes.

Acknowledgements

A.P. acknowledges institutional support from MIMS (Vetenskapsrådet grant 2021-06602) and Umeå University Medical Faculty, Knut and Alice Wallenberg Foundation grant KAW 2015.0225, Umeå University Bioteknikmedel FS 2.1.6-1862.17 and Insamlingsstiftelse FS 2.1.6-452-20, Vetenskapsrådet project grant VR-MH 2022-00778, Carl Kempe Foundation grant SMK-1859, and institutional support from Queen's University Belfast. S.T. acknowledges Svenska Sällskapet för Medicinsk Forskning (SSMF) for postdoctoral grant P19-0098. We thank David A. Cisneros (Queen's University Belfast) for critical reading.

Conflict of Interests
The authors declare that they have no conflict of interest.

Contributions
Conceptualization: S.T. and A.P.
Data curation: S.T.
Formal analysis: S.T., V.P., M.L.B., M.P.P., C.H.O., A.N.
Funding acquisition: S.T. and A.P.
Investigation: S.T., M.L.B., M.P.P., C.H.O., A.N.
Methodology: V.P.
Project administration: A.P.
Resources: O.B.
Software: S.T., V.P. and N.S.
Supervision: S.T. and A.P.
Validation: S.T. and A.P.
Visualization: S.T.
Writing – Original Draft: S.T., V.P. and A.P.
Writing – Reviewing and editing S.T., V.P., M.L.B., M.P.P., N.S., C.H.O., O.B., A.N., A.P.

Material and methods

Reagents
Unless stated otherwise, all chemicals, antibiotics and primers were purchased from Sigma-Aldrich. Restriction and cloning enzymes were obtained from New England Biolabs (NEB). Unless stated otherwise, plasticware for bacterial culture was purchased from Techno Plastic Products (TPP) and Greiner. ATP sodium salt solution 100 mM was obtained from ThermoFisher (reference R0441). Antimicrobial peptides were purchased from Eurogentec. Antibiotic MIC strips were obtained from Liofilchem.

Bacterial strains and culture conditions
The bacterial strains used in this study are listed in Table EV4. Bacteria were preserved as glycerol stocks at -80°C. Bacterial cultures in rich medium were performed in Luria Broth (LB) 1X medium (yeast 5 g/l; tryptone 10 g/l; NaCl 5 g/l). Bacterial cultures in minimal medium were performed in M9 (M9 salts 1X; $CaCl_2$ 0,1 mM; $MgSO_4$ 2 mM; glucose 0,5 %). Kanamycin (25 µg/ml) was added to the medium when needed. If not specified, overnight bacterial cultures were grown at 37°C in the medium of choice, subcultured 1:100, and grown at 37°C in a shaking incubator at 150 RPM until mid-exponential phase (OD_{600nm} = 0.3-0.5).

Screen for genome-wide promoter activity in *E. coli* MG1655 using a transcriptional reporter library
The library of reporter strains, each carrying a low-copy plasmid (pUA66-*mut2GFP*) with a promoter of interest controlling fast-folding GFP (GFPmut2), was previously described (Zaslaver *et al.*, 2006) and purchased from Dharmacon. The library was replicated to accommodate positive and negative controls to create a working library, resulting in punctual changes in sample position with respect to the original library. The disposition of samples is provided in Table EV1. Reporter strains were inoculated in 96-well plates from frozen stocks and grown overnight in the medium of choice with kanamycin at 37°C in a shaking incubator. Overnight cultures were diluted 1:100 into

150 μl of fresh medium containing kanamycin in black 96-well plates with an optical
flat bottom (Greiner). The cultures were incubated at 37°C in a shaking incubator at
150 RPM. When the mid-exponential phase was reached, 50 μl of medium containing
4 mM ATP was added to each well to obtain a final concentration of 1 mM ATP in each
well. In each plate, a strain carrying a promoterless plasmid was included as a
background control. Optical density (OD$_{600nm}$) and fluorescence intensity (excitation:
485 nm; emission: 535 nm) were measured in an Infinite M200 or a Spark plate reader
(Tecan) just before (t=0 min) and after ATP addition (time=5 min) and then measured
at intervals of 60 min for 240 min, with incubation at 37°C in a shaking incubator at
150 RPM between measurements. The expression levels were calculated by
subtracting the background fluorescence (promoterless strain) and by dividing the
intensity of fluorescence by the corresponding OD$_{600nm}$. The reproducibility of the
screening was verified by plotting the normalised fluorescence of a random plate
measured on day 1 against the values on day 2. Promoters showing a final
fluorescence value lower than 100 in both ATP-treated and control samples were
considered as not expressed. The ratio of the fluorescent signal produced by eATP-
treated bacteria and untreated control bacteria was calculated. The screen was
performed twice in LB and in M9, hence statistical analysis was not performed.
517
Construction of chromosomal reporter fusions
Reporter fusions were constructed using Red recombination as previously described
(Gerlach *et al*, 2007). The kanamycin resistance cassette was amplified from pKD4
with ST1_aph_KpnI_F and ST2_aph_XhoI_R, cloned into a KpnI/XhoI digested
pBluescript plasmid and named pBluescript-kan (See Table EV4). The same
procedure was performed with the PCR-amplified msfGFP Addgene plasmid # 29750
and subcloned into EcoRI/HindIII-digested pBluescript-kan. This PCR fragment
contained a ribosome binding site upstream of *msfGFP*. The final plasmid construct
was verified by sequencing. PCR amplification of msfGFP-Kan was performed using
the newly cloned plasmid using ST5_cbpA_sfGFP_aph_F and
ST6_cbpA_sfGFP_aph_R as templates. The PCR product was digested with DpnI
and transformed into *E. coli* MG1566 carrying pKD56. Colonies were selected on
plates containing 50 μg/mL kanamycin and verified by sequencing. The kanamycin
resistance gene was removed by FLP-mediated recombination (Datsenko & Wanner,
2000; Gerlach *et al.*, 2007).
533
Measurement of promoter activity upon ATP addition at small scale
To determine promoter activities at different concentrations of added ATP, in *E. coli*
MG1655 carrying chromosomal fusion reporters and in all strains other than *E. coli*
MG1655 from the transcriptional reporter library (Figure 6), the experimental protocol
was adjusted to the smaller scale compared to the screening setup. The reporter
strains used in these experiments were directly inoculated in 3 ml of appropriate
medium with kanamycin for 16 h at 37°C and 150 RPM. The cultures were then diluted
to obtain a final OD$_{600nm}$=0,01 using the medium of choice at a final volume of 150 μl
per well in black 96-well plates with an optical flat bottom (Greiner) and, as previously
described, grown until mid-exponential phase. After incubation, 50 μl of medium
containing the appropriate ATP concentration was added to each well to obtain 100
μM, 500 μM, 1 mM, 2 mM or 5 mM in the final volume (200 μl). The optical density and
fluorescence were measured, and data were analysed as described above (screening
section). Experiments were performed in at least 3 biological replicates with 2 or 3
technical replicates.

Panther analysis and KEGG pathway analysis

To investigate the biological functions and gene ontology of differentially expressed genes (DEGs, FC>1,5) upon exposure to 1 mM ATP, DAVID Bioinformatics Resources version 6.8 (Huang da Sherman & Lempicki, 2009) and the PANTHER classification system version 17.0 (Thomas *et al*, 2003) were used to functionally annotate and classify genes. The reference list for the GO enrichment analysis contained 1,807 modulated genes of the transcriptional reporter library (Zaslaver *et al.*, 2006).

Sample preparation for untargeted metabolomics

E. coli MG1655 was grown overnight at 37°C in the medium of choice. The cultures were diluted to obtain a final OD_{600nm}=0,01 in 20 mL of appropriate medium and incubated until mid-exponential phase at 37°C under agitation. The subculture was divided into 5 ml samples for each condition. ATP was added (100 µM or 1 mM), and the tubes were incubated at 37°C under agitation for 3 h. The time-matched control samples were supplemented with an equal volume of distilled water. Samples of 2 ml were collected in 2 ml microtubes (Sarstedt, ref 72.690.001) 180 min after the addition of the compounds and centrifuged at 13000 x g for 2 min. The supernatant of M9-cultured bacteria was filtered (pore size: 0,22 µm, diameter: 13 mm, syringe-driven, Millex) and collected in a sample tube. The bacterial pellets were washed 3 times with ice-cold PBS keeping cells on ice between centrifugations. Both the supernatant and the pellet were stored at -20°C until analysis. At least 3 biological replicates were prepared for each condition.

Metabolite extraction for untargeted metabolomics

Bacterial pellets were solubilized using 200 µl 90:10 $MeOH:H_2O$ and were stored at -80°C or kept on dry ice. Acid-washed glass beads of 425-600 µm diameter (Sigma Aldrich) were added to samples kept on dry ice to constitute 50% v/v of the 90:10 $MeOH:H_2O$ cell suspension, and cells were disrupted by shaking at 30 Hz for 2 min (Mixer Mill MM400, Retsch) using prechilled (4°C) holding blocks. Samples were centrifuged at 4°C and 14000 x g for 10 min, followed by transfer of the supernatant into an LC–MS vial (Chromacol, 03-FISV, Thermo Fisher, MA, USA). Samples were evaporated in a centrifugal evaporator (Speed Vac, Savant) at room temperature until dry and stored at -80°C, and reconstituted with 20 µl 60:40 $MeOH:H_2O$ prior to LC-MS analysis.

LC–MS analysis of metabolites

Five microliters were injected into a 1290 Infinity II LC UHPLC system (Agilent, CA, USA) connected to either a 6550 Series or 6540 Series Q-TOF mass spectrometer (Agilent, CA, USA) with a Jet Stream electrospray ionization (ESI) source (Agilent, CA, USA). Data were collected in positive and negative ionization mode with ESI settings: gas temperature 150°C, drying gas flow 16 l/min, nebulizer pressure 35 psi, sheath gas temperature 350°C, sheath gas flow 11 l/min, Vcap 4000 V, nozzle voltage 300 V, Fragmentor voltage 380 V, Skimmer1 45 V and OctapoleRFPeak 750 V. Metabolites were separated using reversed-phase chromatography using an ACQUITY UPLC HSS T3 column with C18 chemistry, 1.8 µM particle size, and 100 Å pore size of 50 mm length and 2.1 mm inner diameter (Waters, MA, USA). For reversed-phase chromatography, elution solvents were (A) H_2O, 0.1% formic acid and (B) 75:25 acetonitrile:isopropanol, 0.1% formic acid. For separation, the following

599   linear gradient was used (flow rate 0.5 ml/min): min 0: 0.1% B, min 2: 10% B, min 7:
600   99% B, min 9: 99% B, min 9.3: 0.1% B, min 10.8: 0.1% B. Column re-equilibration
601   occurred from 9.8 to 10.7 min with an increase in flow rate to 0.8 ml/min.
602
Metabolite identification and relative quantification
Raw data was processed using Agilent Mass Hunter Profinder software and the
metabolites were identified with an inhouse retention time accurate mass library
constructed using synthetic standards on the same analytical setup. A retention time
window of 0.5 minutes and a mass accurace tolerance of 5 ppm was employed. A
statistics-based analysis of the metabolites was performed using MetaboAnalyst 5.0,
a web-based metabolomics processing tool (Chong Wishart & Xia, 2019). The peak
area for each metabolite was uploaded to the MetaboAnalyst program and normalised
using log transformed and autoscaling features (mean-centred and divided by the
standard deviation of each variable). If values were missing, an estimate was
performed using K-nearest neighbours' imputation. A one-way ANOVA using Tukey's
honestly significant difference test was used to determine significant features. Partial
least squares-discriminant analysis (PLS-DA) was used to examine differences
between the groups. Metabolites with a P value<0,05 and fold change (FC)>1,5 were
considered significantly modulated. The normalised peak area of each differential
metabolite was used to construct the bar plots. Heatmaps were constructed to provide
an overview of the metabolome changes, and PLS-DA was used to show the 25 most
important features. VIP (Variable Importance in Projection) values (≥1.0) and
Student's t test (P value <0.05) were combined to identify the differential metabolites.
The VIP score is a weighted sum of squares of the PLS loadings. The weights are
based on the amount of explained Y-variance in each dimension. Metabolites with
high VIP are more important in providing class separation, while those with small VIP
provide less contribution.
626
Susceptibility tests to antibiotics and antimicrobial peptides
E. coli MG1655 was grown overnight at 37°C in LB. The cultures were diluted to obtain
a final OD_{600nm}=0,01 in 10 mL of medium and incubated until mid-exponential phase
at 37°C under agitation. The subculture was divided into 3 ml samples for each
condition. ATP (1 mM) or mock were added, and the tubes were incubated at 37°C
under agitation for 3 h.
Resistance to various antibiotics was determined on agar using MIC Test Strips
(Liofilchem). Exponential cultures were adjusted to OD_{600nm}=0,1 in LB and
homogeneously spread onto an LB agar plate containing or not 1mM ATP. The strip
was applied once the plates had fully dried. Plates were incubated over night at 37°C
and the MIC at which the no-growth halo was observed was determined visually.
Susceptibility to antimicrobial peptides (Eurogentech) was tested by adjusting the
OD_{600nm}=0,04 in LB, 50 µl of culture were distributed into a 96-well plate (TPP),
followed by 150 µl containing serially diluted antimicrobial peptides together with
vehicle or 1 mM ATP. Plates were maintained at 37°C with 3 mm linear agitation in a
Spark multimode plate reader (Tecan) and the OD_{600nm} was acquired every 10 min.
Seven antimicrobial peptides were used: indolicidin, apidaecin IB, CAP-18, cecropin,
melittin, protegrin-1, C18G (see also Table EV4). The antimicrobial peptide stock
solutions were prepared beforehand in sterile water and kept at −80°C until usage,
except for protegrin-1 and C18G, which were prepared in 0,01% acetic acid before
freezing. Experiments were performed in biological triplicates with 2 or 3 technical
replicates.

Cloning of promoter regions into pUA66-*GFPmut2*

The high-fidelity polymerase Phusion was used for all PCRs. All plasmid constructs were verified by sequencing (Eurofins). Plasmids and oligos are listed in Table EV4. Promoter regions (positions with respect to start ATG) of *mgtC* (-480 to 113), *grlA* (-89 to 143), *hilA* (-700 to 457), *clbRA* (-537 to 156), *qseBC* (-480 to 109), *ler* (-407 to 108) and *ssrAB* (-431 to 112) were PCR amplified with primers containing either *XhoI* or *BamHI* restriction sites, using DNA extracted from the relevant organisms with the Wizard® genomic purification kit (Promega) according to manufacturer´s instructions. The purified PCR products were XhoI/BamHI digested. The plasmid pUA66-mut2GFP was XhoI/BamHI digested and purified in a 1% agarose gel run at 100 V for 40 min in TAE buffer, followed by isolation with the GeneJet gel extraction kit (ThermoScientific). The digested PCR products and backbone plasmid were ligated using T4 DNA ligase (Thermo Fisher Scientific). The mixture was transformed into CaCl$_2$-competent *E. coli* DH5α. Transformants were selected on LB agar plates containing 25 µg/ml kanamycin. A few colonies were picked and screened by colony PCR using primers (Table EV4) designed for the upstream and downstream regions of the two restriction sites. Verified recombinant plasmids were purified and transformed into the organisms of promoter origin by electroporation.

Generation of a phylogenetic tree and percentage identity calculation

A phylogenetic tree of the employed strains based on NCBI taxonomy was generated with PhyloT (https://phylot.biobyte.de/index.cgi). The visualization of the tree was achieved using iTOL (https://itol.embl.de/) (Letunic & Bork, 2021). Taxonomy numbers used for the tree are shown in Table EV4. To calculate the percentage identity of *gadW*, *ybjX*, *slyB* and *rstA* between the different strains, the nucleotide sequences were downloaded for every strain and aligned using MUSCLE v5 (https://www.ebi.ac.uk/Tools/msa/muscle/). The identity of each sequence compared to the selected reference sequence (*E. coli* MG1655) was calculated using The Sequence Manipulation Suite (https://www.bioinformatics.org/sms2/ident_sim.html) (Stothard, 2000).

Omics data integration into genome-scale models

To investigate the impact of eATP on *E. coli* MG1655 metabolism in depth, a publicly available genome-scale GEM called iML1515 (Monk *et al*, 2017) was utilised. Two different growth media, M9 and LB, were separately considered, and the model was adapted accordingly. We created a metabolic model for M9 using minimal growth medium features/nutrition input values, which we referred to as M9GEM. For LB, nutritional input values were assigned from a previous study (Marinos Kaleta & Waschina, 2020), and the adapted model was called LBGEM. The REMI method (Pandey Hadadi & Hatzimanikatis, 2019) was used to integrate genetic screen data and metabolomics data into M9GEM and LBGEM, allowing for a better understanding of the perturbation effect on metabolism while simultaneously integrating gene expression and metabolomics data. REMI assumes that the reaction fluxes associated with markedly regulated genes are deregulated, and it also integrates intracellular metabolites by constraining reaction fluxes corresponding to differentially regulated metabolites. The main goal of REMI is to ensure the consistency between gene expression, metabolite concentrations, and reaction fluxes. To determine the specific metabolic differences between two conditions (such as eATP and mock-treated control), REMI creates a separate model for each condition and computes a maximum

consistency (MCS) score. This score reflects the number of genes/metabolites whose
levels are consistent with fluxes, and the constraints can be incorporated into the
model. To generate at least 20 alternative sets for the MCS from the given set of
constraints, we employed mixed-integer linear programming (MILP). Due to the
computational complexity of the optimization process, we selected 20 solutions for
statistical analysis.
705
Differential reaction flux analysis
To quantify condition-specific changes in reaction fluxes, we obtained a score (R_i) for
each reaction by utilizing the means and variances of flux distributions in eATP-treated
and control conditions. To calculate scores for each reaction, we employed the REMI
analysis approach (Pandey Hadadi & Hatzimanikatis, 2019), which involves
considering alternative solutions of reaction fluxes for two conditions.
To identify markedly regulated reaction fluxes that were significantly altered, we
utilized a cut-off of $|R_i|>1$. Additionally, we calculated *P* values using a nonparametric
test, namely, the Wilcoxon signed rank test. We considered reactions with a *P* value
<1e-3 to be significantly changed.

716
$$R_i = \frac{Mean_{eatp}(v_i) - Mean_{control}(v_i)}{\sqrt{Var_{eatp}(v_i) - Var_{control}(v_i)}}$$

Pathway and metabolic subsystem enrichment analysis
We conducted pathway enrichment analysis based on modulated genes and
metabolites using KEGG pathways as the framework for analysis. To analyse
metabolic subsystem enrichment based on metabolic reactions, we utilized GEM
iML1515 (Monk *et al.*, 2017). To determine the significance of pathway or metabolic
subsystems based on genes, metabolites, or reactions, we employed the
hypergeometric probability density function (*P*), as defined by Equation (1):

725
$$P(k,N,K,n) = \frac{\binom{K}{k}\binom{N-K}{n-k}}{\binom{N}{n}} \qquad (1)$$

where k and K represent the number of markedly regulated genes, metabolites, or
reactions and the total number of genes/metabolites/reactions in a given pathway or
metabolic subsystem, respectively. Variables n and N indicate the total number of
markedly regulated genes, metabolites, or reactions in the given conditions and the
total number of genes/metabolites/reactions in the organism and metabolic context,
respectively.
To calculate the *p value* for each pathway or metabolic subsystem, we summed the
probabilities for greater than or equal to the enriched number of markedly regulated
genes, metabolites, or reactions (as shown in Equation 2):

735
$$pvalue = \sum_{k}^{K} P(x) \qquad (2)$$

736
737 **Biomass building block analysis**
738 Biomass building blocks (BBBs) are crucial components of genome-scale metabolic
739 models that include amino acids, nucleotides, lipids, and other compounds required

for cellular growth and maintenance. We used flux balance analysis (FBA), a commonly used constraint-based method, to explore the dynamics of these building components under various environmental conditions. FBA allowed us to examine BBB fluxes in various media, including those subjected to eATP perturbations, providing vital insights into how an organism's metabolism adapts and responds to these changes.

Identification of metabolic subnetworks

To gain a more thorough understanding of the metabolic remodelling that results from exposure to eATP, we focused our efforts on identifying subnetworks containing both differentially regulated metabolites (DRMs) and differentially regulated reaction fluxes (DRFs). To accomplish this, we first removed co-factors from the network, as they can be involved in multiple metabolic reactions and can artificially inflate connectivity. This allowed us to better identify important metabolic subnetworks and reveal the true structure of the network. Additionally, we examined the first and second neighbour reactions of each DRM. First neighbour reactions are those that are directly involved with a metabolite's first neighbours, while second neighbour reactions involve the metabolite's second neighbours, which are not directly connected to the metabolite but are connected to its first neighbours. Finally, we analysed the most perturbed or deregulated reactions to identify the metabolic subnetworks where the most deregulation occurs in terms of both metabolites and reactions.

Statistical analysis

Except for the metabolomic analysis, which was performed using Metaboanalyst (5.0), all statistical analyses were performed in GraphPad Prism (version 9.0.2). The type of analysis and test of significance is mentioned in the respective figure legends. The default P value system of GraphPad Prism was used in all cases and is indicated in figure legends.

Source code availability

The script for metabolic modelling can be found at https://gitlab.com/obilab1/externalatpmets

Softwares used

The metabolomic analysis as performed with MetaboAnalyst 5.0 (Chong Wishart & Xia, 2019). For the visualisation of enrichment pathways and BBB analysis, Python (3.9) and R (4.2) were used. For modelling simulation, IBM CPLEX 12.9 MILP solver and MATLAB r2019a (The Mathworks, Natick, MA, USA) were used. The figures 4D and 5 were created with Illustrator.

Abbasian B, Shair A, O'Gorman DB, Pena-Diaz AM, Brennan L, Engelbrecht K, Koenig DW, Reid G, Burton JP (2019) Potential Role of Extracellular ATP Released by Bacteria in Bladder Infection and Contractility. *mSphere* 4: e00439-00419

Alvarez CL, Corradi G, Lauri N, Marginedas-Freixa I, Leal Denis MF, Enrique N, Mate SM, Milesi V, Ostuni MA, Herlax V, Schwarzbaum PJ (2017) Dynamic regulation of extracellular ATP in Escherichia coli. *The Biochemical journal* 474: 1395-1416

Arvidsson I, Tontanahal A, Johansson K, Kristoffersson AC, Kellnerová S, Berger M, Dobrindt U, Karpman D (2022) Apyrase decreases phage induction and Shiga toxin release from E. coli O157:H7 and has a protective effect during infection. *Gut Microbes* 14: 2122667

Atarashi K, Nishimura J, Shima T, Umesaki Y, Yamamoto M, Onoue M, Yagita H, Ishii N, Evans R, Honda K, Takeda K (2008) ATP drives lamina propria T(H)17 cell differentiation. *Nature* 455: 808-812

Baeza N, Mercade E (2021) Relationship Between Membrane Vesicles, Extracellular ATP and Biofilm Formation in Antarctic Gram-Negative Bacteria. *Microb Ecol* 81: 645-656

Berrier C, Coulombe A, Szabo I, Zoratti M, Ghazi A (1992) Gadolinium ion inhibits loss of metabolites induced by osmotic shock and large stretch-activated channels in bacteria. *European journal of biochemistry* 206: 559-565

Bours MJL, Swennen ELR, Di Virgilio F, Cronstein BN, Dagnelie PC (2006) Adenosine 5′-triphosphate and adenosine as endogenous signaling molecules in immunity and inflammation. *Pharmacology & Therapeutics* 112: 358-404

Chang-Graham AL, Perry JL, Engevik MA, Engevik KA, Scribano FJ, Gebert JT, Danhof HA, Nelson JC, Kellen JS, Strtak AC *et al* (2020) Rotavirus induces intercellular calcium waves through ADP signaling. *Science* 370

Chen F, El-Naccache DW, Ponessa JJ, Lemenze A, Espinosa V, Wu W, Lothstein K, Jin L, Antao O, Weinstein JS *et al* (2022) Helminth resistance is mediated by differential activation of recruited monocyte-derived alveolar macrophages and arginine depletion. *Cell Rep* 38: 110215

Chiaranunt P, Burrows K, Ngai L, Cao EY, Liang H, Tai SL, Streutker CJ, Girardin SE, Mortha A (2022) NLRP1B and NLRP3 Control the Host Response following Colonization with the Commensal Protist Tritrichomonas musculis. *Journal of immunology (Baltimore, Md : 1950)* 208: 1782-1789

Chiaranunt P, Burrows K, Ngai L, Tai SL, Cao EY, Liang H, Hamidzada H, Wong A, Gschwend J, Flüchter P *et al* (2023) Microbial energy metabolism fuels an intestinal macrophage niche in solitary isolated lymphoid tissues through purinergic signaling. *Sci Immunol* 8: eabq4573

Choi S, Choi E, Cho YJ, Nam D, Lee J, Lee EJ (2019) The Salmonella virulence protein MgtC promotes phosphate uptake inside macrophages. *Nat Commun* 10: 3326

Chong J, Wishart DS, Xia J (2019) Using MetaboAnalyst 4.0 for Comprehensive and Integrative Metabolomics Data Analysis. *Curr Protoc Bioinformatics* 68: e86

Covert MW, Palsson B (2002) Transcriptional regulation in constraints-based metabolic models of Escherichia coli. *The Journal of biological chemistry* 277: 28058-28064

Culp EJ, Goodman AL (2023) Cross-feeding in the gut microbiome: Ecology and mechanisms. *Cell Host Microbe* 31: 485-499

Darfeuille-Michaud A, Neut C, Barnich N, Lederman E, Di Martino P, Desreumaux P, Gambiez L, Joly B, Cortot A, Colombel JF (1998) Presence of adherent Escherichia coli strains in ileal mucosa of patients with Crohn's disease. *Gastroenterology* 115: 1405-1413

Datsenko KA, Wanner BL (2000) One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci USA* 97

Deiwick J, Nikolaus T, Erdogan S, Hensel M (1999) Environmental regulation of Salmonella pathogenicity island 2 gene expression. *Mol Microbiol* 31: 1759-1773

Di Virgilio F, Pinton P, Falzoni S (2016) Assessing Extracellular ATP as Danger Signal
In Vivo: The pmeLuc System. *Methods in molecular biology (Clifton, NJ)* 1417: 115-
129

Ding Q, Quah SY, Tan KS (2016) Secreted adenosine triphosphate from
Aggregatibacter actinomycetemcomitans triggers chemokine response. *Mol oral*
Microbiol 31: 423-434

Ding Q, Tan KS (2016) The Danger Signal Extracellular ATP Is an Inducer of
Fusobacterium nucleatum Biofilm Dispersal. *Front Cell Infect Microbiol* 6: 155

Ellermeier CD, Ellermeier JR, Slauch JM (2005) HilD, HilC and RtsA constitute a feed
forward loop that controls expression of the SPI1 type three secretion system regulator
hilA in Salmonella enterica serovar Typhimurium. *Mol Microbiol* 57: 691-705

Gerlach RG, Hölzer SU, Jäckel D, Hensel M (2007) Rapid Engineering of Bacterial
Reporter Gene Fusions by Using Red Recombination. *Appl Environ Microbiol* 73:
4234-4242

Gerosa L, Sauer U (2011) Regulation and control of metabolic fluxes in microbes. *Curr*
Opin Biotechnol 22: 566-575

Gianchandani EP, Joyce AR, Palsson B, Papin JA (2009) Functional states of the
genome-scale Escherichia coli transcriptional regulatory system. *PLoS computational*
biology 5: e1000403

Hironaka I, Iwase T, Sugimoto S, Okuda K, Tajima A, Yanaga K, Mizunoe Y (2013)
Glucose triggers ATP secretion from bacteria in a growth-phase-dependent manner.
Appl Environ Microbiol 79: 2328-2335

Huang da W, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nature protocols* 4: 44-57

Idzko M, Ferrari D, Eltzschig HK (2014) Nucleotide signalling during inflammation.
Nature 509: 310-317

Idzko M, Hammad H, van Nimwegen M, Kool M, Willart MA, Muskens F, Hoogsteden
HC, Luttmann W, Ferrari D, Di Virgilio F *et al* (2007) Extracellular ATP triggers and
maintains asthmatic airway inflammation by activating dendritic cells. *Nat Med* 13:
913-919

Ihssen J, Jovanovic N, Sirec T, Spitz U (2021) Real-time monitoring of extracellular
ATP in bacterial cultures using thermostable luciferase. *PLOS ONE* 16: e0244200

Ivanova EP, Alexeeva YV, Pham DK, Wright JP, Nicolau DV (2006) ATP level
variations in heterotrophic bacteria during attachment on hydrophilic and hydrophobic
surfaces. *International microbiology : the official journal of the Spanish Society for*
Microbiology 9: 37-46

Iwase T, Shinji H, Tajima A, Sato F, Tamura T, Iwamoto T, Yoneda M, Mizunoe Y
(2010) Isolation and identification of ATP-secreting bacteria from mice and humans. *J*
Clin Microbiol 48: 1949-1951

Krautkramer KA, Fan J, Bäckhed F (2021) Gut microbial metabolites as multi-kingdom
intermediates. *Nat Rev Microbiol* 19: 77-94

Kurashima Y, Amiya T, Nochi T, Fujisawa K, Haraguchi T, Iba H, Tsutsui H, Sato S,
Nakajima S, Iijima H *et al* (2012) Extracellular ATP mediates mast cell-dependent
intestinal inflammation through P2X7 purinoceptors. *Nat Commun* 3: 1034

Kusu T, Kayama H, Kinoshita M, Jeon SG, Ueda Y, Goto Y, Okumura R, Saiga H,
Kurakawa T, Ikeda K *et al* (2013) Ecto-nucleoside triphosphate diphosphohydrolase 7
controls Th17 cell responses through regulation of luminal ATP in the small intestine.
Journal of immunology (Baltimore, Md : 1950) 190: 774-783

Letunic I, Bork P (2021) Interactive Tree Of Life (iTOL) v5: an online tool for
phylogenetic tree display and annotation. *Nucleic Acids Res* 49: W293-w296

Li M, Chen Y, Wu S, Tang Y, Deng Y, Yuan J, Dong J, Li H, Tang L (2016) TmcN is involved in ATP regulation of tautomycetin biosynthesis in Streptomyces griseochromogenes. *Biochem Biophys Res Commun* 478: 221-226

Li M, Kim TJ, Kwon HJ, Suh JW (2008) Effects of extracellular ATP on the physiology of Streptomyces coelicolor A3(2). *FEMS Microbiol Lett* 286: 24-31

Li M, Lee SK, Yang SH, Ko JH, Han JS, Kim TJ, Suh JW (2011) ATP modulates the growth of specific microbial strains. *Curr Microbiol* 62: 84-89

Li XP, Zhou LL, Guo YH, Wang JW (2021) The signaling role of extracellular ATP in co-culture of Shiraia sp. S9 and Pseudomonas fulva SB1 for enhancing hypocrellin A production. *Microbial cell factories* 20: 144

Lou L, Zhang P, Piao R, Wang Y (2019) Salmonella Pathogenicity Island 1 (SPI-1) and Its Complex Regulatory Network. *Front Cell Infect Microbiol* 9: 270

Malik-Kale P, Winfree S, Steele-Mortimer O (2012) The bimodal lifestyle of intracellular Salmonella in epithelial cells: replication in the cytosol obscures defects in vacuolar replication. *PLOS ONE* 7: e38732

Malo MS, Moaven O, Muhammad N, Biswas B, Alam SN, Economopoulos KP, Gul SS, Hamarneh SR, Malo NS, Teshager A *et al* (2014) Intestinal alkaline phosphatase promotes gut bacterial growth by reducing the concentration of luminal nucleotide triphosphates. *Am J Physiol Gastrointest Liver Physiol* 306: G826-838

Marinos G, Kaleta C, Waschina S (2020) Defining the nutritional input for genome-scale metabolic models: A roadmap. *PLOS ONE* 15: e0236890

Marteyn B, West NP, Browning DF, Cole JA, Shaw JG, Palm F, Mounier J, Prévost MC, Sansonetti P, Tang CM (2010) Modulation of Shigella virulence in response to available oxygen in vivo. *Nature* 465: 355-358

McDonough KA, Rodriguez A (2011) The myriad roles of cyclic AMP in microbial pathogens: from signal to sword. *Nat Rev Microbiol* 10: 27-38

Medina-Castellanos E, Esquivel-Naranjo EU, Heil M, Herrera-Estrella A (2014) Extracellular ATP activates MAPK and ROS signaling during injury response in the fungus Trichoderma atroviride. *Frontiers in plant science* 5: 659

Medina-Castellanos E, Villalobos-Escobedo JM, Riquelme M, Read ND, Abreu-Goodger C, Herrera-Estrella A (2018) Danger signals activate a putative innate immune system during regeneration in a filamentous fungus. *PLOS Genet* 14: e1007390

Mempin R, Tran H, Chen C, Gong H, Kim Ho K, Lu S (2013) Release of extracellular ATP by bacteria during growth. *BMC Microbiol* 13: 301

Meng L, Li M, Yang SH, Kim TJ, Suh JW (2011) Intracellular ATP levels affect secondary metabolite production in Streptomyces spp. *Bioscience, biotechnology, and biochemistry* 75: 1576-1581

Mi H, Muruganujan A, Casagrande JT, Thomas PD (2013) Large-scale gene function analysis with the PANTHER classification system. *Nature protocols* 8: 1551-1566

Mitosch K, Beyß M, Phapale P, Drotleff B, Nöh K, Alexandrov T, Patil KR, Typas A (2023) A pathogen-specific isotope tracing approach reveals metabolic activities and fluxes of intracellular Salmonella. *PLOS Biol* 21: e3002198

Monk JM, Lloyd CJ, Brunk E, Mih N, Sastry A, King Z, Takeuchi R, Nomura W, Zhang Z, Mori H *et al* (2017) iML1515, a knowledgebase that computes Escherichia coli traits. *Nat Biotechnol* 35: 904-908

Nilsson L (1978) New rapid bioassay of gentamicin based on luciferase assay of extracellular ATP in bacterial cultures. *Antimicrob Agents Chemother* 14: 812-816

Nolan LM, Cavaliere R, Turnbull L, Whitchurch CB (2015) Extracellular ATP inhibits
twitching motility-mediated biofilm expansion by Pseudomonas aeruginosa. *BMC*
Microbiol 15: 55

Nougayrède J-P, Homburg S, Taieb F, Boury M, Brzuszkiewicz E, Gottschalk G,
Buchrieser C, Hacker J, Dobrindt U, Oswald E (2006) Escherichia coli Induces DNA
Double-Strand Breaks in Eukaryotic Cells. *Science* 313: 848-851

Pandey V, Hadadi N, Hatzimanikatis V (2019) Enhanced flux prediction by integrating
relative expression and relative metabolite abundance into thermodynamically
consistent metabolic models. *PLoS computational biology* 15: e1007036

Patel BA (2014) Mucosal adenosine triphosphate mediates serotonin release from
ileal but not colonic guinea pig enterochromaffin cells. *Neurogastroenterology and*
motility : the official journal of the European Gastrointestinal Motility Society 26: 237-
246

Perruzza L, Gargari G, Proietti M, Fosso B, D'Erchia AM, Faliti CE, Rezzonico-Jost T,
Scribano D, Mauri L, Colombo D *et al* (2017) T Follicular Helper Cells Promote a
Beneficial Gut Ecosystem for Host Metabolic Homeostasis by Sensing Microbiota-
Derived Extracellular ATP. *Cell Rep* 18: 2566-2575

Pham AQ, Cho SH, Nguyen CT, Stacey G (2020) Arabidopsis Lectin Receptor Kinase
P2K2 Is a Second Plant Receptor for Extracellular ATP and Contributes to Innate
Immunity. *Plant Physiol* 183: 1364-1375

Platenkamp A, Mellies JL (2018) Environment Controls LEE Regulation in
Enteropathogenic Escherichia coli. *Front Microbiol* 9: 1694

Proietti M, Perruzza L, Scribano D, Pellegrini G, D'Antuono R, Strati F, Raffaelli M,
Gonzalez SF, Thelen M, Hardt WD *et al* (2019) ATP released by intestinal bacteria
limits the generation of protective IgA against enteropathogens. *Nat Commun* 10: 250

Puhar A, Tronchère H, Payrastre B, Tran Van Nhieu G, Sansonetti Philippe J (2013)
A Shigella Effector Dampens Inflammation by Regulating Epithelial Release of Danger
Signal ATP through Production of the Lipid Mediator PtdIns5P. *Immunity* 39: 1121-
1131

Qiu H, Pu F, Liu Z, Deng Q, Sun P, Ren J, Qu X (2019) Depriving Bacterial Adhesion-
Related Molecule to Inhibit Biofilm Formation Using CeO2 -Decorated Metal-Organic
Frameworks. *Small (Weinheim an der Bergstrasse, Germany)*: e1902522

Raffelsberger N, Hetland MAK, Svendsen K, Småbrekke L, Löhr IH, Andreassen LLE,
Brisse S, Holt KE, Sundsfjord A, Samuelsen Ø, Gravningen K (2021) Gastrointestinal
carriage of Klebsiella pneumoniae in a general adult population: a cross-sectional
study of risk factors and bacterial genomic diversity. *Gut Microbes* 13: 1939599

Rooks MG, Veiga P, Reeves AZ, Lavoie S, Yasuda K, Asano Y, Yoshihara K, Michaud
M, Wardwell-Scott L, Gallini CA *et al* (2017) QseC inhibition as an antivirulence
approach for colitis-associated bacteria. *Proc Natl Acad Sci U S A* 114: 142-147

Scott BM, Gutiérrez-Vázquez C, Sanmarco LM, da Silva Pereira JA, Li Z, Plasencia
A, Hewson P, Cox LM, O'Brien M, Chen SK *et al* (2021) Self-tunable engineered yeast
probiotics for the treatment of inflammatory bowel disease. *Nat Med* 27: 1212-1222

Stothard P (2000) The sequence manipulation suite: JavaScript programs for
analyzing and formatting protein and DNA sequences. *BioTechniques* 28: 1102, 1104

Tadala L, Langenbach D, Dannborg M, Cervantes-Rivera R, Sharma A, Vieth K,
Rieckmann LM, Wanders A, Cisneros DA, Puhar A (2022) Infection-induced
membrane ruffling initiates danger and immune signaling via the mechanosensor
PIEZO1. *Cell Rep* 40: 111173

Takemura N, Kurashima Y, Mori Y, Okada K, Ogino T, Osawa H, Matsuno H, Aayam L, Kaneto S, Park EJ *et al* (2018) Eosinophil depletion suppresses radiation-induced small intestinal fibrosis. *Sci Transl Med* 10

Tanaka K, Choi J, Cao Y, Stacey G (2014) Extracellular ATP acts as a damage-associated molecular pattern (DAMP) signal in plants. *Frontiers in plant science* 5: 446

Tani H, Li B, Kusu T, Okumura R, Nishimura J, Okuzaki D, Motooka D, Arakawa S, Mori A, Yoshihara T *et al* (2021) The ATP-hydrolyzing ectoenzyme E-NTPD8 attenuates colitis through modulation of P2X4 receptor-dependent metabolism in myeloid cells. *Proc Natl Acad Sci U S A* 118

Tatano Y, Kanehiro Y, Sano C, Shimizu T, Tomioka H (2015) ATP exhibits antimicrobial action by inhibiting bacterial utilization of ferric ions. *Sci Rep* 5: 8610

Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K, Muruganujan A, Narechania A (2003) PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res* 13: 2129-2141

Tjaden J, Schwöppe C, Möhlmann T, Quick PW, Neuhaus HE (1998) Expression of a plastidic ATP/ADP transporter gene in Escherichia coli leads to a functional adenine nucleotide transport system in the bacterial cytoplasmic membrane. *The Journal of biological chemistry* 273: 9630-9636

Valat C, Champiat D, N'Guyen TT, Loiseau G, Raimbault M, Montet D (2003) Use of ATP bioluminescence to determine the bacterial sensitivity threshold to a bacteriocin. *Luminescence* 18: 254-258

Verkhratsky A, Burnstock G (2014) Biology of purinergic signalling: its ancient evolutionary roots, its omnipresence and its multiple functional significance. *BioEssays : news and reviews in molecular, cellular and developmental biology* 36: 697-705

Wallenstein A, Rehm N, Brinkmann M, Selle M, Bossuet-Greif N, Sauer D, Bunk B, Spröer C, Wami HT, Homburg S *et al* (2020) ClbR Is the Key Transcriptional Activator of Colibactin Gene Expression in Escherichia coli. *mSphere* 5: e00591-00520

Wan P, Liu X, Xiong Y, Ren Y, Chen J, Lu N, Guo Y, Bai A (2016) Extracellular ATP mediates inflammatory responses in colitis via P2 x 7 receptor signaling. *Sci Rep* 6: 19108

Wilhelm K, Ganesan J, Müller T, Dürr C, Grimm M, Beilhack A, Krempl CD, Sorichter S, Gerlach UV, Jüttner E *et al* (2010) Graft-versus-host disease is enhanced by extracellular ATP activating P2X7R. *Nat Med* 16: 1434-1438

Xi C, Wu J (2010) dATP/ATP, a multifunctional nucleotide, stimulates bacterial cell lysis, extracellular DNA release and biofilm development. *PLOS ONE* 5: e13355

Yu Lomakina G, Ugarova NN (2022) Application of Bioluminescent Methods to Study the Effect of the Membrane-active Antibiotic Colistin on Bacterial Cells. *Photochem Photobiol* 98: 1077-1083

Zaslaver A, Bren A, Ronen M, Itzkovitz S, Kikoin I, Shavit S, Liebermeister W, Surette MG, Alon U (2006) A comprehensive library of fluorescent transcriptional reporters for Escherichia coli. *Nature methods* 3: 623-628

Figure Legends

Figure 1. eATP triggers a strong dose-, medium- and time- dependent transcriptional response in non-pathogenic *E. coli*.
A and **C**, Number of eATP-regulated promoters in percentage with respect to unstimulated controls from fluorescence reporter screens (n=2) performed in LB or minimal medium M9 in *E. coli* MG1655, 180 min after addition of 1 mM ATP.

B and **D**, Log fold change of promoter activity upon exposure to 1 mM eATP 180 min
after ATP addition in LB or M9 in *E. coli* MG1655, shown as mean of independent
experiments (n≥2).
E-**G**, Comparison of the number of (**E**) all differentially expressed promoters in *E. coli*
MG1655, (**F**) all up-regulated promoters, and (**G**) all down-regulated promoters 180
min after addition of 1 mM ATP in LB or M9.
H-**M**, Promoter activity of selected up-regulated promoters P*ybjX*, P*gadW* and P*rstA*
controlling the expression of the reporter GFP in the presence of increasing
concentrations of added ATP in *E. coli* MG1655. Strains were grown in LB (upper
panels) or M9 (lower panels). Shown is the ratio of GFP fluorescence to the number
of bacteria determined by absorbance at 600 nm over time (columns representing the
mean with s.e.m, left Y axes), as well as the bacterial growth (continuous lines
representing means with s.e.m, right Y axes). Statistical analysis: two-way ANOVA
with Dunnett's multiple comparison test; *P < 0.05, **P < 0.01, ***P < 0.001 and ****P
< 0.0001; ns, not significant.
1048
Figure 2. Pathway analysis of the eATP-triggered transcriptional response in
non-pathogenic *E. coli*.
A-**B**, KEGG pathway enrichment analysis on the up- and down-regulated genes in LB
or M9 180 min after 1 mM ATP addition in comparison to unstimulated bacteria.
C, Gene ontology analysis using the PANTHER classification system of up- or down-
regulated genes in LB and M9. Processes with a P value < 1×10^{-3} are shown.
1055
Figure 3. eATP triggers a strong metabolic response in non-pathogenic *E. coli*.
A, Comparison of the total number of common or unique detected metabolites from *E.*
coli MG1655 grown in LB or M9 after 180 min in the presence of 1 mM added ATP.
LB pellet: n=3 and 57 metabolites, M9 pellet: n=4 and 53 metabolites, M9 supernatant:
n=3 and 89 metabolites.
B-**D**, Partial Least Squares discriminant analysis (PLS-DA) scores plots of
components one and two, comparing metabolomic samples from *E. coli* MG1655
treated with 1 mM ATP for 180 min versus the unstimulated control in (**B**) the pellet in
LB (n=3), (**C**) the pellet in M9 (n=4), and (**D**) the supernatant in M9 (n=3).
E, Comparison of fold changes in abundance of top 25 metabolites based on PLS-DA
analysis from samples in (**B**).
F, Comparison of changes in abundance of top 25 metabolites based on PLS-DA
analysis from samples in (**C**).
G, Comparison of fold changes in abundance of top 25 metabolites based on PLS-DA
analysis from samples in (**D**).
H-**J**, Changes of relative concentrations of selected modulated intracellular
metabolites from samples in (**B** and **D**) and extracellular metabolites from samples in
(**D**). The boxplots show the normalised values (mean and s.d.). The Y-axis represents
the normalised peak area of metabolites. Black dots represent the normalised
concentrations of all samples of each metabolite. The notch indicates the 95%
confidence interval around the median. The mean concentration of each group is
shown by a yellow diamond. Statistical analysis: two-sided Welch's t test; raw P-values
are indicated in each panel.
1079
1080 **Figure 4. Model-based enrichment analysis of predicted reaction fluxes and**
1081 **analysis of biomass building blocks triggered by eATP in non-pathogenic *E.***
1082 ***coli*.**

A, Metabolic subsystems enrichment analysis for *E. coli* MG1655 stimulated with 1 mM eATP for 180 min in rich (LB) medium using differentially regulated reactions (DERs).

B, Metabolic subsystems enrichment analysis for *E. coli* MG1655 stimulated with 1 mM eATP for 180 min in minimal (M9) medium using differentially regulated reactions (DERs).

C, Differential fold changes of *in silico* production for eight biomass building block groups between *E. coli* MG1655 grown in rich (LB) and minimal (M9) medium in presence of 1 mM eATP for 180 min *versus* the unstimulated control. Each dot is an alternative solution predicted by the genome-scale metabolic model. A cut-off P-value <0,05 was used.

D, Identification of metabolic subnetworks based on differentially regulated reactions and differentially regulated intracellular metabolites triggered in *E. coli* MG1655 by addition of 1 mM ATP for 180 min in LB (n=3). Shown are the differentially regulated intracellular metabolites (labelled in red, fold change >1,5, see also Fig **3H**), modulated genes (labelled in orange, fold change >1,5, Table EV1) and differentially regulated reactions (labelled in capital purple and black circles, P-value < 1e-3). Other compounds are labelled in orange circles and written in black. Non-modulated reactions are labelled in capital grey and black circles. TRPtipp, L-tryptophan transport out (periplasm); INSH, Inosine hydrolase; UMPK, UMP kinase; GALUi, UTP-glucose-1-phosphate uridylyltransferase (irreversible).

Figure 5. Metabolic subnetwork analysis integrating transcriptional and metabolic changes triggered by eATP in non-pathogenic *E. coli* in M9.

Identification of metabolic subnetworks based on differentially regulated reactions and differentially regulated intracellular metabolites triggered in *E. coli* MG1655 by addition of 1 mM ATP for 180 min M9 (n=4). Shown are the differentially regulated intracellular metabolites (labelled in red, fold change >1,5, see also Fig **3I**), modulated genes (labelled in orange, fold change >1,5, Table EV1) and differentially regulated reactions (labelled in capital purple and black circles, P-value < 1e-3). Other compounds are labelled in orange circles and written in black. Non-modulated reactions are labelled in capital grey and black circles. DADA, Deoxyadenosine deaminase; PUNP6, Purine-nucleoside phosphorylase (Deoxyinosine); GLUPRT, Glutamine phosphoribosyldiphosphate amidotransferase; ACGS, N-acetylglutamate synthase; GTHS, Glutathione synthetase; NNAM, Nicotinamidase; CYTDK2, Cytidine kinase (GTP); TKT1, Transketolase; AMPN, AMP nucleosidase; PUNP2, Purine-nucleoside phosphorylase (Deoxyadenosine).

Figure 6. eATP alters antibiotics and antimicrobial peptides susceptibility, triggers a transcriptional response in several Enterobacteriacae and regulates the expression of virulence factors.

A, Relative antibiotic sensitivity of *E. coli* MG1655 grown on LB compared to LB + 1mM ATP plates in the presence of ampicillin (AMP), fosfomycin (FOS), aztreonam (ATM), ceftazidime (CAZ), imipenem (IMI), cefazolin (KZ), mecillinam (MEC), gentamicin (CN), plazomicin (PLZ), or ciprofloxacin (CIP). Shown is the mean with s.d. of 3 independent experiments. Statistical analysis: multiple t-test; *P < 0.05; ns, not significant.

B-H, Growth of *E. coli* MG1655 in LB upon treatment at time 0 with antimicrobial peptides or vehicle in the presence or absence of 1 mM ATP. Shown is the mean with s.d. of 3 independent experiments performed in duplicates or triplicates. Statistical

analysis: two-way ANOVA with Dunnett's multiple comparison test; *P < 0.05, **P <
0.01, ***P < 0.001 and ****P < 0.0001; ns, not significant.
I, Response ratio to eATP in percentage in selected bacteria transformed with reporter
plasmids encoding promoters that upregulate the expression of GFP in eATP-treated
commensal *E. coli* MG1655, which was used as reference (100%). Bacteria were
stimulated with 1 mM ATP for 180 min in LB or M9. Shown is the average ratio
calculated from at least 3 independent experiments performed in triplicates.
J-M, Promoter activity of selected virulence or fitness factors seen as GFP expression
normalised to number of bacteria upon addition of increasing concentrations of ATP
for 180 min in LB. Indicated promoter regions were cloned into GFP reporter plasmids
and transformed into indicated corresponding strains. Increasing concentrations of
eATP were added and the expression ratio was calculated comparing the
fluorescence/OD_{600nm} in LB without ATP with fluorescence/OD_{600nm} in LB + ATP.
Shown is the mean with s.e.m. of $\geq$ 3 independent experiments performed in
duplicates or triplicates. Statistical analysis: two-way ANOVA with Dunnett's multiple
comparison test; *P < 0.05, **P < 0.01, ***P < 0.001 and ****P < 0.0001; ns, not
significant.
1150
1151
1152   Expanded View Figure Legends
1153
Expanded view Fig 1. eATP triggers a strong transcriptional response in non-
pathogenic *E. coli*, related to Fig 1.
A, Overview of the protocol for the promoter library screening for a transcriptional
response to eATP. Created using BioRender.com.
B-G, Activity of indicated promoters controlling the expression of the reporter GFP
(plasmid construct) in the presence or absence of 1 mM ATP obtained in LB (upper
panels) or M9 (lower panels). Shown is the ratio of GFP fluorescence to number of
bacteria determined by absorbance at 600 nm over time (columns representing the
mean with s.e.m, left Y axes), as well as the bacterial growth (continuous lines
representing means with s.e.m, right Y axes).
H, Day-to-day reproducibility for a set of 96 random samples. Shown is the GFP
fluorescence divided by the number of bacteria determined by absorbance at 600 nm
in LB measured on day 1 and day 2.
I, Growth curves of *E. coli* MG1655 over 240 min with (blue line) or without (orange
line) 1mM ATP addition in LB obtained for a set of 2 random plates from the library.
The continuous lines represent the mean absorbance and the colour-matched
shaded areas the s.e.m.
J, Bacterial counts determined by plating before and 180 min after 1 mM ATP
addition in LB.
K, Same as in (**I**) but in M9.
L, Same as in (**J**) but in M9.
M, Promoter activity of MG1655 carrying the P*gadW*-mEGFP chromosomal fusion in
the presence and absence of 1 mM ATP obtained as in **B-G**. The mean fluorescence
value of the unstimulated control at 0 min was set to 1 to calculate relative changes of
the fluorescence. The activity of the *gadW* promoter obtained using the reporter
plasmid is shown in Fig **1I** for comparison.
N, Promoter activity (GFP/OD_{600nm}) of the reporter plasmid encoding P*cbpA-gfp* in the
presence of 1mM ATP in *E. coli* MG1655.

O, Promoter activity of *E. coli* MG1655 carrying the P*cbpA*-mEGFP chromosomal
fusion in the presence and absence of 1 mM ATP obtained as in **B-G**. The mean
fluorescence value of the unstimulated control at 0 min was set to 1 to calculate
relative changes of the fluorescence. The activity of the *cbpA* promoter obtained using
the reporter plasmid is shown in Fig **1I** for comparison.
Statistical analysis: two-way ANOVA with Dunnett's multiple comparison test; *P <
0.05, **P < 0.01, ***P < 0.001 and ****P < 0.0001; ns, not significant.
1189
Expanded view Fig 2. eATP triggers a strong metabolic response in non-
pathogenic *E. coli* related to Fig 3.
A, Volcano plot analysis of intracellular metabolites changes after addition of 1 mM
ATP in LB. Fold change (FC) ≥ 1.5 or ≤0.5 (p ≤ 0.05) is considered significant.
B, VIP (Variable Importance in Projection) score of PLS-DA analysis from samples in
(**A**).
C, Volcano plot analysis of intracellular metabolites changes after addition of 1 mM
ATP in M9. Fold change (FC) ≥ 1.5 or ≤0.5 (p ≤ 0.05) is considered significant.
D, VIP score of PLS-DA analysis from samples in (**C**).
E, Volcano plot analysis of extracellular metabolites changes after addition of 1 mM
ATP in M9. Fold change (FC) ≥ 1.5 or ≤0.5 (p ≤ 0.05) is considered significant.
F, VIP score of PLS-DA analysis from samples in (**E**).
1202
Expanded view Fig 3. Modulated intracellular metabolites upon exposure to
100 μM ATP in M9 in non-pathogenic *E. coli*, related to Fig 3.
A, PLS-DA scores plots of components one and two, comparing metabolomic samples
(pellet) from treatment with 100 μM ATP for 180 min versus the unstimulated controls
(n=3).
B, Comparison of fold changes in abundance of top 25 metabolites based on PLS-
DA analysis from samples in (**A**).
C, Volcano plot analysis of intracellular metabolites changes from samples in (**A**). Fold
change (FC) ≥ 1.5 or ≤0.5 (p ≤ 0.05) is considered significant.
D, VIP (Variable Importance in Projection) score of PLS-DA analysis from samples in
(**A**).
E, Boxplots of relative concentrations of selected modulated intracellular metabolites
from samples in (**A**). The bar plots show the normalised values (mean with s.d.). The
Y-axis represents normalised peak area of metabolites. Black dots represent the
normalised concentrations of all samples of each metabolite. The notch indicates the
95% confidence interval around the median. The mean concentration of each group
is shown by a yellow diamond. Statistical analysis: two-sided Welch's t test; raw P-
values are indicated in each panel.
F, KEGG pathway enrichment analysis of the up- and down-regulated intracellular
metabolites in M9 180 min after 100 μM ATP addition in comparison to unstimulated
bacteria.
G, Comparison of the number of all measured intracellular metabolites in *E. coli*
MG1655 180 min after addition of 1 mM or 100 μM ATP M9 medium.
H, Comparison of the number of all modulated intracellular metabolites in *E. coli*
MG1655 180 min after addition of 1 mM or 100 μM ATP M9 medium.
1228
1229 **Expanded view Fig 4. eATP enhances antimicrobial peptide susceptibility,**
1230 **related to Fig 6.**

A-G, Growth of *E. coli* MG1655 in LB upon treatment at time 0 with antimicrobial peptides or vehicle in the presence or absence of 1 mM ATP. Shown is the mean with s.d. of 3 independent experiments performed in duplicates or triplicates. Statistical analysis: two-way ANOVA with Dunnett's multiple comparison test; *P < 0.05; **P < 0.01, ***P < 0.001 and ****P < 0.0001; ns, not significant.

Expanded view Fig 5. eATP triggers a transcriptional response in non-pathogenic, pathobiontic or pathogenic bacteria and regulates the expression of virulence factors, related to Fig 6.

A, Sequence similarity in percentage of promoters P*gadW*, P*ybjX*, P*rstA* and P*slyB* promoters in the indicated bacteria compared to *E. coli* MG1655.

B, Promoter activity of the promoter P*rstA* controlling the expression of the reporter GFP in the presence and absence of 1 mM ATP in indicated strains grown in LB, as in Fig **6A**. Shown is the ratio of GFP fluorescence to number of bacteria determined by the absorbance at 600 nm over time (columns representing the mean with s.e.m of 3 independent experiments performed in triplicates, left Y axes), as well as the bacterial growth (continuous lines representing means with s.e.m, right Y axes). Statistical analysis: two-way ANOVA with Dunnett's multiple comparison test; *P < 0.05, **P < 0.01, ***P < 0.001 and ****P < 0.0001; ns, not significant.

C-D, Promoter activity of selected virulence or fitness factors seen as GFP expression normalised to number of bacteria upon addition of increasing concentrations of ATP for 180 min in LB. The expression ratio was calculated comparing the fluorescence/OD_{600nm} in LB without ATP with fluorescence/OD_{600nm} in LB + ATP. Shown is the mean with s.e.m. of $\geq$ 3 independent experiments performed in duplicates or triplicates. Statistical analysis: two-way ANOVA with Dunnett's multiple comparison test; ns, not significant.

Expanded view Tables legend

Expanded view Table 1: Transcriptional reporter library screening
Screening_LB: Summary of the data from the transcriptional library screening in LB. Shown is the average GFP fluorescence divided by OD_{600nm} of at least 2 biological replicates for each time point (0, 5, 30, 60, 120 and 180 min after ATP addition), the fluorescence ratio between the eATP-treated and the mock-treated samples and the log_2 of this ratio.
Screening_M9: Same as aforementioned but in M9.
Panther analysis: Results from the Panther analysis in LB and M9 of differentially expressed genes (DEGs, FC>1,5) upon exposure to 1 mM ATP for 180 min.
LB_genes_KEGG: Pathway enrichment analysis based on differentially expressed genes upon exposure to 1 mM ATP in LB for 180 min.
M9_genes_KEGG: Pathway enrichment analysis based on differentially expressed genes upon exposure to 1 mM ATP in M9 for 180 min.

Expanded view Table 2: Metabolomic data
Metabolites_LC_MS: Summary of all metabolites identified in *E. coli* MG1655 upon exposure to 100 µM or 1 mM ATP for 180 min and their associated FC and P-value compared to mock-treated bacteria.
Metabolites_role: Summary of modulated bioactive metabolites and their roles.

Expanded view Table 3: Metabolic and flux modelling

M9_diff_flux: List of reaction fluxes identified in *E. coli* MG1655 upon addition of 1 mM ATP for 180 min in M9.

M9_flux_enrichment: Metabolic subsystems enrichment analysis for *E. coli* MG1655 stimulated with 1 mM ATP for 180 min in M9 using differentially regulated reactions (DERs).

M9_BBB: List of the tested biomass building blocks (BBBs) and their associated production rate ratio in eATP-treated compared to mock-treated bacteria in M9 upon exposure to eATP for 180min.

LB_diff_flux List of reaction fluxes identified in *E. coli* MG1655 upon addition of 1 mM ATP for 180 min in M9.

LB_flux_enrichment: Metabolic subsystems enrichment analysis for *E. coli* MG1655 stimulated with 1 mM ATP for 180 min in LB using differentially regulated reactions (DERs).

LB_BBB: List of the tested biomass building blocks and their associated production rate ratio between the eATP treated bacteria and the mock in LB upon exposure to eATP for 180min.

Expanded view Table: Strains, plasmids, and primers
Strains: List of bacterial strains used in the study.
Plasmids: List of plasmids used in the study.
Primers: List of the primers used in the study.

Expanded view Table: List of antibiotics and antimicrobial peptides used in the study
Antibiotics: List of antibiotics used in the study and the MIC obtained on LB agar and LB agar supplemented with 1mM ATP.
AMPs: List of antimicrobial peptides used in the study.

References Expanded view Figures and Tables.

Allaoui, A., J. Mounier, M. C. Prévost, P. J. Sansonetti, and C. Parsot. 1992. 'icsB: a Shigella flexneri virulence gene necessary for the lysis of protrusions during intercellular spread', *Mol Microbiol*, 6: 1605-16.

Baba, T., T. Ara, M. Hasegawa, Y. Takai, Y. Okumura, M. Baba, K. A. Datsenko, M. Tomita, B. L. Wanner, and H. Mori. 2006. 'Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection', *Mol Syst Biol*, 2.

Bassil, A. K., S. Bourdu, K. A. Townson, A. Wheeldon, E. M. Jarvie, N. Zebda, A. Abuin, E. Grau, G. P. Livi, L. Punter, J. Latcham, A. M. Grimes, D. P. Hurp, K. M. Downham, G. J. Sanger, W. J. Winchester, A. D. Morrison, and G. B. Moore. 2009. 'UDP-glucose modulates gastric function through P2Y14 receptor-dependent and -independent mechanisms', *Am J Physiol Gastrointest Liver Physiol*, 296: G923-30.

Budnick, J. A., X. R. Bina, and J. E. Bina. 2021. 'Complete Genome Sequence of Klebsiella pneumoniae Strain ATCC 43816', *Microbiol Resour Announc*, 10.

Cliffe, E. E., and S. G. Waley. 1961. 'The mechanism of the glyoxalase I reaction, and the effect of ophthalmic acid as an inhibitor', *Biochem J*, 79: 475-82.

Colosimo, D. A., J. A. Kohn, P. M. Luo, F. J. Piscotta, S. M. Han, A. J. Pickard, A. Rao, J. R. Cross, L. J. Cohen, and S. F. Brady. 2019. 'Mapping Interactions of

Microbial Metabolites with Human G-Protein-Coupled Receptors', *Cell Host Microbe*, 26: 273-82.e7.

Darfeuille-Michaud, A., C. Neut, N. Barnich, E. Lederman, P. Di Martino, P. Desreumaux, L. Gambiez, B. Joly, A. Cortot, and J. F. Colombel. 1998. 'Presence of adherent Escherichia coli strains in ileal mucosa of patients with Crohn's disease', *Gastroenterology*, 115: 1405-13.

Darveau, R. P., J. Blake, C. L. Seachord, W. L. Cosand, M. D. Cunningham, L. Cassiano-Clough, and G. Maloney. 1992. 'Peptides related to the carboxyl terminus of human platelet factor IV with antibacterial activity', *The Journal of Clinical Investigation*, 90: 447-55.

Datsenko, K. A., and B. L. Wanner. 2000. 'One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products', *Proc Natl Acad Sci USA*, 97.

Di Stasi, L. C. 2021. 'Coumarin Derivatives in Inflammatory Bowel Disease', *Molecules*, 26.

Faas, M. M., T. Saez, and P. de Vos. 2017. 'Extracellular ATP and adenosine: The Yin and Yang in immune responses?', *Mol Aspects Med*.

Falla, T. J., D. N. Karunaratne, and R. E. Hancock. 1996. 'Mode of action of the antimicrobial peptide indolicidin', *J Biol Chem*, 271: 19298-303.

Garavaglia, M., E. Rossi, and P. Landini. 2012. 'The pyrimidine nucleotide biosynthetic pathway modulates production of biofilm determinants in Escherichia coli', *PLOS ONE*, 7: e31252.

Gobbo, M., M. Benincasa, G. Bertoloni, B. Biondi, R. Dosselli, E. Papini, E. Reddi, R. Rocchi, R. Tavano, and R. Gennaro. 2009. 'Substitution of the arginine/leucine residues in apidaecin Ib with peptoid residues: effect on antimicrobial activity, cellular uptake, and proteolytic degradation', *J Med Chem*, 52: 5197-206.

Hale, T. L., E. V. Oaks, and S. B. Formal. 1985. 'Identification and antigenic characterization of virulence-associated, plasmid-coded proteins of Shigella spp. and enteroinvasive Escherichia coli', *Infect Immun*, 50: 620-9.

Kim, E. K., K. A. Lee, D. Y. Hyeon, M. Kyung, K. Y. Jun, S. H. Seo, D. Hwang, Y. Kwon, and W. J. Lee. 2020. 'Bacterial Nucleoside Catabolism Controls Quorum Sensing and Commensal-to-Pathogen Transition in the Drosophila Gut', *Cell Host Microbe*.

Larrick, J. W., M. Hirata, Y. Shimomoura, M. Yoshida, H. Zheng, J. Zhong, and S. C. Wright. 1993. 'Antimicrobial activity of rabbit CAP18-derived peptides', *Antimicrobial Agents and Chemotherapy*, 37: 2534-9.

Lee, S. B., B. Li, S. Jin, and H. Daniell. 2011. 'Expression and characterization of antimicrobial peptides Retrocyclin-101 and Protegrin-1 in chloroplasts to control viral and bacterial infections', *Plant Biotechnol J*, 9: 100-15.

Liu, C., Y. Cheng, Y. Guo, and H. Qian. 2021. 'Magnesium-L-threonate alleviate colonic inflammation and memory impairment in chronic-plus-binge alcohol feeding mice', *Brain Res Bull*, 174: 184-93.

Luchini, A. C., P. Rodrigues-Orsi, S. H. Cestari, L. N. Seito, A. Witaicenis, C. H. Pellizzon, and L. C. Di Stasi. 2008. 'Intestinal anti-inflammatory activity of coumarin and 4-hydroxycoumarin in the trinitrobenzenesulphonic acid model of rat colitis', *Biol Pharm Bull*, 31: 1343-50.

O'Donnell, M. P., B. W. Fox, P. H. Chao, F. C. Schroeder, and P. Sengupta. 2020. 'A neurotransmitter produced by gut bacteria modulates host sensory behaviour', *Nature*, 583: 415-20.

Offord, R. E., J. Philippe, J. G. Davis, P. A. Halban, and M. Berger. 1979. 'Inhibition of
degradation of insulin by ophthalamic acid and by a bovine pancreatic
proteinase inhibitor', *Biochem J*, 182: 249-51.
Ohsu, T., Y. Amino, H. Nagasaki, T. Yamanaka, S. Takeshita, T. Hatanaka, Y.
Maruyama, N. Miyamura, and Y. Eto. 2010. 'Involvement of the calcium-
sensing receptor in human taste perception', *J Biol Chem*, 285: 1016-22.
Selsted, M. E., M. J. Novotny, W. L. Morris, Y. Q. Tang, W. Smith, and J. S. Cullor.
1992. 'Indolicidin, a novel bactericidal tridecapeptide amide from neutrophils', *J
Biol Chem*, 267: 4292-5.
Soga, T., R. Baran, M. Suematsu, Y. Ueno, S. Ikeda, T. Sakurakawa, Y. Kakazu, T.
Ishikawa, M. Robert, T. Nishioka, and M. Tomita. 2006. 'Differential
metabolomics reveals ophthalmic acid as an oxidative stress biomarker
indicating hepatic glutathione consumption', *J Biol Chem*, 281: 16768-76.
Sridharan, G. V., K. Choi, C. Klemashevich, C. Wu, D. Prabakaran, L. B. Pan, S.
Steinmeyer, C. Mueller, M. Yousofshahi, R. C. Alaniz, K. Lee, and A.
Jayaraman. 2014. 'Prediction and quantification of bioactive microbiota
metabolites in the mouse gut', *Nat Commun*, 5: 5492.
Steiner, M. R., J. S. Bomalaski, and M. A. Clark. 1993. 'Responses of purified
phospholipases A2 to phospholipase A2 activating protein (PLAP) and melittin',
Biochim Biophys Acta, 1166: 124-30.
Sun, Q., J. G. Weinger, F. Mao, and G. Liu. 2016. 'Regulation of structural and
functional synapse density by L-threonate through modulation of intraneuronal
magnesium concentration', *Neuropharmacology*, 108: 426-39.
Wang, J., K. Ma, M. Ruan, Y. Wang, Y. Li, Y. V. Fu, Y. Song, H. Sun, and J. Wang.
2018. 'A novel cecropin B-derived peptide with antibacterial and potential anti-
inflammatory properties', *PeerJ*, 6: e5369.
Zaslaver, A., A. Bren, M. Ronen, S. Itzkovitz, I. Kikoin, S. Shavit, W. Liebermeister, M.
G. Surette, and U. Alon. 2006. 'A comprehensive library of fluorescent
transcriptional reporters for Escherichia coli', *Nat Methods*, 3: 623-8.

1409
1410
1411
1412

Fig. 1 eATP triggers a strong transcriptomic response in *E. coli* in a medium-, dose- and time-dependent manner

Fig. 2 eATP modulates genes involved in critical pahtways, such as lipids, amino acid or vitamin metabolism

Fig. 3 Intracellular metabolites quantification upon exposure to eATP

Fig. 4. Reaction fluxes enrichment analysis and biomass building blocks analysis

Fig. 5 Metabolic map of eATP impact in LB

Fig. 6 Antimicrobial molecule susceptibility and response to eATP in other bacteria

Fig. EV1 Promoter library screening, figure related to Fig. 1

Fig. EV2 Intracellular metabolites quantification upon exposure to 1mM ATP in LB and M9

Fig.EV3 Intracellular metabolites modulation upon exposure to 100 µM ATP in M9

Fig.EV4 AMP susceptibility

Fig. EV5 - eATP triggers a transcriptional response in several bacteria and regulates the expression of virulence factors.

A Promoter region similarity

	gadW	*ybjX*	*rstA*	*slyB*
E. coli K-12 substr. MG1655	100.0	100.0	100.0	100.0
E. coli BW25113	100.0	82.2	100.0	100.0
E. coli O127 H6 E2348/69 (EPEC)	94.7	97.0	98.9	99.5
E. coli LF82	94.7	97.5	98.9	99.5
E. coli Nissle 1917	94.6	80.2	98.9	99.5
S. flexneri 5a M90T	99.3	81.7	91.3	100.0
S. Typhimurium SL1344		43.6	66.6	63.5
K. pneumoniae ATCC 43816			72.9	74.8