Development of a qPCR assay and a LAMP assay for *Verticillium longisporum*

detection and a triplex qPCR assay for simultaneous detection of *V. longisporum*,

***Leptosphaeria biglobosa* and *L. maculans* from canola samples**

Heting Fu[1], Yalong Yang[1], Junye Jiang[1], Greg C. Daniels[2], Blake Hill[2], Shiming Xue[1],

Kher Zahr[1], L. Stellar[2], Michael W. Harding[2], David Feindel[1], Carol Bvindi[3], Dilantha

Fernando[3], Lipu Wang[4], and Jie Feng[1]

[1]Alberta Plant Health Lab, Alberta Agriculture and Irrigation (AGI), Edmonton, AB,

T5Y 6H3, Canada.

[2]Crop Diversification Centre South, AGI, Brooks, AB, T1R 1E6, Canada.

[3]Department of Plant Science, University of Manitoba, Winnipeg, MB, R3T 2N2, Canada.

[4]Department of Plant Sciences, University of Saskatchewan, Saskatoon, SK, S7N 5A8,

Canada.

Corresponding author:

Jie Feng, email: jie.feng@gov.ab.ca

Abstract

Verticillium wilt, Verticillium stem striping, and Verticillium stripe, are common disease names that all denote infection caused by *Verticillium longisporum*, on canola, or other Brassica crops. In this study, a quantitative PCR (qPCR) assay and a loop-mediated isothermal amplification (LAMP) assay were developed for the detection of *V. longisporum* from canola stem samples. Both assays are specific to *V. longisporum* at the species level and ubiquitous at the strain level. The low limit for positive detection of the two assays is 1 pg fungal DNA in a 20-µL reaction or 1,400 fungal cells in 100-mg plant tissue. The qPCR assay was combined with the duplex qPCR assay for the two blackleg pathogens, *Leptosphaeria biglobosa* and *L. maculans* to constitute a triplex qPCR system for simultaneous detection of all three pathogens. The usefulness of this triplex qPCR system was verified on canola samples collected from various locations in Alberta, Canada. Using this triplex qPCR system, *V. longisporum* was detected from one sample, while the two blackleg pathogens were detected at higher frequencies. Since it is sometimes difficult to differentiate Verticillium stripe and blackleg on Alberta canola samples based on visual symptoms, the triplex qPCR system is an important tool for the detection of *V. longisporum*, especially when its presence is masked or obscured by symptoms of blackleg.

Key words:

Verticillium stem striping, Stripe, Wilt, Canola, Blackleg, Multiplex qPCR

Introduction

Verticillium longisporum is a serious pathogen on brassicaceous plants (Depotter et al.

2016). It has been reported in Europe (Karapapa et al. 1997), Russia (Pantou et al. 2005),

Japan (Ikeda et al. 2012), China (Yu et al. 2015), and Canada (CFIA 2016). On oilseed

rape (canola), the disease caused by *V. longisporum* has been called Verticillium wilt but

later renamed to Verticillium stem striping (Depotter et al. 2016). For consistency and

brevity, we will use the name "Verticillium stripe" throughout the remainder of the paper

understanding that any of the aforementioned common disease names may apply to

infections by this fungus.

In Canada, *V. longisporum* was first found in Manitoba in 2014, and a nationwide survey

conducted by the Canadian Food Inspection Agency and the Canola Council of Canada

indicated the presence of the pathogen in canola stubble from five other provinces:

British Columbia, Alberta, Saskatchewan, Ontario and Quebec (CFIA 2016). However,

thereafter *V. longisporum* on canola has not been confirmed in Alberta in annual canola

disease surveys. Inspectors observed symptoms reminiscent of Verticillium stripe but,

blackleg pathogens, *Leptosphaeria maculans,* and *L. biglobosa*, rather than *V.

longisporum*, were isolated from the suspect plant samples (Harding et al. 2022; 2023).

Different from many other fungal species, *V. longisporum* is amphidiploid that resulted

from hybridization between two of the following four haploid ancestors: genotype A1 of

an unknown species, genotype D1 of another unknown species, and genotypes D2 and

D3 of *V. dahliae*. The currently identified *V. longisporum* isolates can be classified into

three lineages based on the parental combination: A1/D1, A1/D2, and A1/D3. Among the

three lineages, A1/D1 is the most pathogenic lineage on canola, whereas lineage A1/D3

isolates are generally not pathogenic on this crop (Novakazi et al. 2015; Tran et al. 2013).

Lineage A1/D2 is known only from horseradish in Illinois (USA) (Eastburn and Chang

1994). In Canada, a recent study indicated that *V. longisporum* isolates from canola and

radish were in the A1/D1 group (Zou et al. 2020).

An effective disease management program is dependent on timely and proper

identification of the causal agent of the disease. However, due to the diploid nature of *V.*

longisporum and the fact that at least one of the parental species (i.e. *V. dahliae*) is a

commonly occurring and widely distributed plant pathogen, accurate detection of *V.*

longisporum has been difficult. Indeed, a DNA-based method for *V. longisporum*

identification from plant samples has never been published. A multiplex PCR protocol

developed by Inderbitzin et al. (2013) could be used to identify and differentiate *V.*

longisporum into the lineage level by using DNA from pure fungal cultures. However,

the use of this protocol for diagnosis of Verticillium stripe from plant samples has not

been reported so far.

Quantitative PCR (qPCR) is an efficient method for plant disease diagnosis. qPCR

especially probe-based qPCR is highly specific and sensitive. Another technique, loop-

mediated isothermal amplification (LAMP), has gained popularity because of its

capability for pathogen detection under a constant temperature and simple instrument

requirements (Notomi et al. 2000). In recent years, LAMP reaction master mix became

available commercially, which further facilitated the use of LAMP in diagnostics. Thus,

in this study, we developed and evaluated a probe-based qPCR assay and a LAMP assay

for specific detection of *V. longisporum* from infected canola plant samples. The qPCR

assay was combined with a previously developed duplex qPCR for the two blackleg

pathogens (Fu et al. 2023) to constitute a triplex qPCR system, which was then tested for

capacity to specifically detect the three pathogens from canola samples collected from

various fields in Alberta.

Materials and methods

Chemicals and standard techniques

All chemicals and instruments were purchased from Fisher Scientific Canada (Ottawa,

ON) unless otherwise specified. All primers, probes, and gBlock were synthesized by

Integrated DNA Technologies (Coralville, IA). All DNA extraction was conducted using

the DNeasy Plant Pro kits (Qiagen Canada, Toronto, ON) with a Qiacube (Qiagen

Canada). The extracted DNA was dissolved in 50 µL water. When needed, DNA

concentration was measured with a Nanodrop 1000. The whole-genome sequenced *V.*

longisporum strains were identified by searching "Verticillium longisporum" in the

National Center for Biotechnology Information (NCBI) genome database

(https://www.ncbi.nlm.nih.gov/datasets/genome/?taxon=100787). The average genome

size of all the whole-genome sequenced strains was calculated. The molecular weight of

the average genome was calculated using an online ssDNA Mass to Moles Convertor

(https://nebiocalculator.neb.com).

Multiplex PCR, qPCR, and LAMP reactions

Multiplex PCR was conducted in Promega Go Taq Master Mix or Thermo Scientific

Phire Plant Direct PCR Master Mix with a T100 thermocycler (Bio-Rad Canada,

Mississauga, ON). Each 20-µL reaction contained 0.25 µM of each primer and 1 ng DNA

from fungal cultures or 20 ng DNA from plants (infected or healthy). The PCR program

followed Zou et al. (2020). SYBR Green-based qPCR and probe-based qPCR were

conducted in SsoAdvanced universal SYBR green supermix (Bio-Rad Canada) and

PrimeTime gene expression master mix (Integrated DNA Technologies), respectively, in

a CFX96 touch real-time PCR detection system (Bio-Rad Canada). Each 20-µL qPCR

reaction contained 0.25 µM of each primer, 0.18 µM of each probe (probe-based qPCR

only), and 2 µL of each DNA template regardless of the concentrations. Each qPCR

reaction was conducted with three technical replicates. In each 96-well PCR plate, three

replicates of negative-control reactions were also included, in each of which 2 µL of

water was used as a template. The qPCR program consisted of an initial denaturation at

95°C for 3 min, followed by 40 cycles of denaturation at 95°C for 10 s and

annealing/extension at 60°C for 30 s. All LAMP reactions were conducted in the

WarmStart® Colorimetric LAMP Master Mix (NEB Canada, Whitby, ON). Each 20-µL

reaction contained 2 µL of template. Quantities of other components in each reaction

followed the NEB's instructions for the master mix. All reactions were conducted in

individual 200-µL PCR tubes. The reaction program consisted of only one step in which

the tubes were incubated at 65°C for 30 minutes. After the incubation, the reactions were

checked visually and the results were recorded by photography.

Selection of genes for primer designing

Candidate genes for qPCR and LAMP primer design were selected by the following

approach. All proteins of *V. longisporum* were accessed by searching the NCBI database

(https://www.ncbi.nlm.nih.gov) with the term 'Verticillium longisporum', which ended

up with 95,405 protein entries. From these proteins, those consisting of 50-60 amino

acids were retrieved, which ended up with 366 entries. After removing the 13 proteins

with predicted functions and 222 partial proteins, 131 protein entries remained. Blastp

was conducted with these 131 proteins as queries against NCBI protein database. After

removing the proteins that produced hit(s) from non-*V. longisporum* species, 66 proteins

remained. The genomic sequences (i.e. coding sequence and introns) were retrieved for

these 66 proteins (hereafter referred to as 66 genes). Two Blastn analyses were conducted

using the genomic sequences of these 66 genes as queries, one against the nucleotide

collection (nr/nt) database and the other against the whole-genome shotgun contigs (wgs)

database with the Organism set as Fungi (taxid:4751). After removing the genes that

produced hit(s) from non-*V. longisporum* species, four genes remained (GenBank

accession numbers CRK29573, CRK26624, CRK12486, and CRK45493). Among these

four genes, the first two are present in all the eight whole-genome sequenced *V.*

longisporum strains (strains PD589, VLB2, VL20, VL2, Vl32, VL1, Vl43, and Vl145c)

with 100% query cover and identical sequences. The other two were present in six whole-

genome sequenced strains with 100% query cover and identical sequences but were not

present in strains Vl32 and PD589. Both CRK29573 and CRK26624 had a coding

sequence of 180 nucleotides (nt), with the former having no intron and the latter having a

58-nt intron (thus the gene is 238 nt).

Primer, probe, and gBlock design for qPCR and LAMP

Based on the sequence of CRK29573, qPCR primer pairs were designed using Primer-

Blast (https://www.ncbi.nlm.nih.gov/tools/primer-blast). Each of the designed primer

pairs was tested in Primer 3 (https://primer3.ut.ee) for probe designing. Three

primers/probe sets were obtained. The sequences of the primers and the probe of each of

these three sets and the sequences of the primers/probe sets (P-Lm and P-Lb; Table 1)

used for the duplex qPCR for canola blackleg detection (Fu et al. 2023) were analyzed for

potential primer dimers using Multiple Primer Analyzer hosted by Thermo Fisher

(https://www.thermofisher.com). One set without any predicted primer dimer was

selected to be used in this study and named P-VL (Table 1). The probe in P-VL was

labeled with the fluorescent dye Cyanine5 (Cy5). A DNA sequence at 499 nt,

corresponding to nt 2669667 to 2670165 of the *V. longisporum* strain PD589 whole

genome sequence (GenBank accession number 200JAETXT010000011) and containing

the sequence of CRK29573, was retrieved from the NCBI database. A double-stranded

DNA fragment of this 499-nt sequence was synthesized as gBlock and named G-VL.

Based on the sequence of CRK26624, two sets of LAMP primers were designed using

Primer Explorer (https://primerexplorer.jp). After testing on *V. longisporum* genomic

DNA, only one set was used in further studies and named LAMP-VL (Table 1).

Specificity test of P-VL and LAMP-VL

The specificities of P-VL and LAMP-VL were tested on one known *V. longisporum*

isolate (APHL 189), and 27 isolates of other species including 18 fungi and one

oomycete (*Pythium intermedium*). APHL 189 was isolated from a canola sample

provided by Dr. Manika Pradhan (Manitoba Agriculture, Food and Rural Development)

in 2016. Other species are common pathogens of canola or have been isolated by Alberta

Plant Health Lab (APHL) from either diseased canola samples or samples of crops

rotated with canola in Alberta. All fungal and oomycete species were grown on 4% (w/v)

potato dextrose agar (PDA) plates. For each species, approximately 50 mg of fresh

mycelia was harvested and subjected to DNA extraction. DNA was also extracted from a

canola clubroot gall sample infected by *Plasmodiophora brassicae* pathotype 5. In

addition, four DNA samples were prepared as negative controls, including one from 7-

day-old healthy canola leaves and one from 7-day-old healthy canola roots. The third and

fourth negative controls were prepared from a pellet from 50 mL of YPG media (1%

yeast extract, 1% peptone, 2% glucose, w/v) and LB broth, respectively, after incubating

10 g of canola root tissue collected from a *Verticillium*-free field plot at 150 rpm 30°C for

48 hours (the pellet was prepared by centrifuging the culture after the root tissue being

removed at $4{,}000 \times g$ for 5 min). The concentration of all DNA samples was adjusted to

500 pg/µL. The qPCR primers/probe set P-VL and LAMP primer set LAMP-VL were

tested on all the above-mentioned DNA samples. In addition, to have a DNA quality and

quantity control, all DNA samples were tested with the universal primer pair ITS1/ITS2

(White et al. 1990; Table 1) by SYBR green-based qPCR. The qPCR analysis was

conducted twice.

Sensitivity test of qPCR and LAMP

The gBlock G-VL was dissolved in water and a 10 pM stock solution was prepared,

which roughly equals 6×10^6 double-stranded DNA copies/µL. A set of 10-fold serial

dilutions was prepared from 6×10^6 copies/µL to 6 copies/µL. The serial dilutions were

tested by probe-based singleplex qPCR using P-VL. The experiment was conducted twice

with alternative preparations of gBlock serial dilutions.

The sensitivity of P-VL was further tested on genomic DNA of a known *V. longisporum*

isolate APHL 189. Genomic DNA of APHL 189 was adjusted to 50 ng/µL and then a set

of 10-fold serial dilutions was prepared from 50 ng/µL to 0.5 fg/µL. The serial dilutions

were tested by probe-based singleplex qPCR using P-VL and by triplex probe-based

qPCR using P-VL, P-Lb, and P-Lm. P-Lb and P-Lm were developed by Fu et al. (2023)

for the detection of the two canola blackleg pathogens *L. biglobosa* and *L. maculans*,

respectively. The serial dilutions of the APHL 189 genomic DNA were also tested by

LAMP using the primer set LAMP-VL. The qPCR and LAMP experiments were

conducted twice with alternative preparations of genomic DNA serial dilutions.

Test of qPCR and LAMP on field samples

Twenty-four canola stem samples showing black lesions were collected from 15 counties

in central and southern Alberta (Fig. 1). In addition, two canola stem samples (MB 1 and

MB 2) were collected from Manitoba, in which MB 2 was confirmed to be infected by *V.*

longisporum (Dilantha Fernando, unpublished data). The samples were cut into 2-mm

cubes and air-dried overnight at 23°C. From each of the air-dried samples, a 100-mg

subsample was subjected to DNA extraction. The obtained DNA samples were tested by

the triplex qPCR using P-VL, P-Lb and P-Lm and LAMP using LAMP-VL. The

experiment was conducted twice on DNA samples from alternative 100-mg subsamples

for each plant sample.

Test of multiplex PCR on field samples and *V. longisporum* isolates

Selected field samples and the *V. longisporum* isolate APHL 189 were tested by the

multiplex PCR using the primer set P-Multi (Table 1), which was developed by

Inderbitzin et al. (2013) and included three primer pairs specific to the glyceraldehyde-3-

phosphate dehydrogenase gene (GAPDH), the elongation factor 1-alpha gene (EF1) and

the internal transcribed spacer region (ITS), respectively. According to Inderbitzin et al.

(2013), from the *V. longisporum* lineage A1/D1, the EF1 and ITS primers will produce a

310-base pair (bp) band and a 1020-bp band, respectively; from the lineage A1/D2, the

EF1 primers will produce the 310-bp band; from the lineage A1/D3, the GAPDH and

EF1 primers will produce a 490-bp band and the 310-bp band, respectively and from *V.*

dahliae, only the GAPDH primers will produce the 490-bp band. Thus, the primer set P-

Multi can be used to differentiate the three *V. longisporum* lineages and *V. dahliae* from

each other.

Results

Specificity of P-VL and LAMP-VL

In the qPCR analysis, the primer set ITS1/ITS2 produced signals from all the DNA

samples, as indicated by the quantification cycle (Cq) value from each reaction (Table 2),

which provided a reference for the quality and quantity of each DNA sample. P-VL

produced signals from the *V. longisporum* isolate APHL 189 (Table 2). In contrast, no

signal was produced from any of the triplicated qPCR reactions on DNA samples of all

the non-*V. longisporum* isolates and the four negative controls (Table 2). In the LAMP

analysis, LAMP-VL generated identical results as P-VL in the qPCR analysis: the signal

was produced from the *V. longisporum* isolate APHL 189 only (data not shown). Similar

results were obtained from the repeated experiment. These results confirmed that both P-

VL and LAMP-VL are specific to *V. longisporum*.

Sensitivity of P-VL in singleplex qPCR

The serial dilutions of the gBlock G-VL and *V. longisporum* genomic DNA were tested

in singleplex qPCR with P-VL. Based on the Cq values, standard curves were constructed

for P-VL against gBlock (Fig. 2a) and genomic DNA (Fig. 2b). The efficiencies of P-VL

against gBlock and genomic DNA were 1.01 (Fig. 2a) and 1.00 (Fig. 2b), respectively.

The low limit of P-VL for a positive detection in a 20-µL reaction was 24 ($10^{1.08} \times 2 = 24$)

single-stranded gBlock copies (Fig. 2a) or 1 pg genomic DNA (Fig. 2b). The average

genome size of the eight whole-genome sequenced *V. longisporum* stains is 69.7 Mb.

Based on the molecular weight calculation, one pg of the genomic DNA is composed of

28 genomes of *V. longisporum*. Thus the low detection limits of the qPCR assay using P-

VL were similar against gBlock and fungal genomic DNA.

Sensitivity of LAMP-VL

The serial dilutions of the *V. longisporum* genomic DNA were tested by LAMP using

LAMP-VL. The low limit of LAMP-VL for a positive detection in a 20-µL reaction was

1 pg genomic DNA (Fig. 3). This result indicated that the LAMP system has a similar

sensitivity as the qPCR system.

Sensitivity of P-VL in triplex qPCR

The serial dilutions of the *V. longisporum* genomic DNA were tested in triplex qPCR

using P-VL, P-Lb, and P-Lm. Based on the Cq values, standard curves were constructed

for each primer set (Fig. 4a-c). The efficiencies of P-VL, P-Lb, and P-Lm were 1.02, 1.00

and 0.97, respectively. The low limits of genomic DNA for a positive detection in a 20-

µL reaction were 1 pg for P-VL (Fig. 4a), 0.4 ($10^{-0.42} = 0.4$) pg for P-Lb (Fig. 4b) and 0.5

($10^{-0.27} = 0.5$) pg for P-Lm (Fig. 4c). This result indicated that the sensitivity of P-VL in

triplex qPCR was similar to that in singleplex qPCR.

Triplex qPCR and LAMP analyses on field samples

Twenty-six canola samples, 24 from Alberta and two from Manitoba were tested by the

triplex qPCR using the primer sets P-VL, P-Lb, and P-Lm (Table 3). *Verticillium*

longisporum was detected from one Alberta sample (23CAN-305B) and one Manitoba

sample (MB 2). Of the three pathogens, *V. longisporum*, *L. biglobosa,* and *L. maculans*,

at least one was detected from each of the 26 samples. From 13 of the 24 Alberta samples,

both *L. biglobosa* and *L. maculans* were detected. Among the 13 samples, the sample

positive for *V. longisporum* (23CAN-305B) was also positive for *L. biglobosa* and *L.*

maculans. In addition, all 26 canola samples were tested by LAMP using LAMP-VL. The

LAMP results were identical to the qPCR results for the yes/no detection of *V.*

longisporum (Table 3). These results indicated that both the triplex qPCR and LAMP

developed in this study can be used on field samples.

Confirmation of the triplex qPCR and LAMP results by multiplex PCR

With the Promega master mix, the multiplex PCR produced a single band close to 490 bp

from the two *V. dahliae* isolates, two bands close to 1020 bp and 310 bp, respectively,

from the *V. longisporum* isolate and three *V. longisporum*-infected canola samples (Fig.

5a). With the Thermo Fisher master mix, the multiplex PCR produced similar results as

with the Promega master mix except that two bands were produced from the two *V.*

dahliae isolates (Fig. 5b). Nevertheless, multiplex PCR confirmed the *V. longisporum*

identity of APHL 189 and the presence of *V. longisporum* in the three canola samples,

and also confirmed the specificity of our triplex qPCR and LAMP.

Discussion

Specificity of P-VL and LAMP-VL

The specificity of P-VL and LAMP-VL are reflected by the specificity of their target

genes, GenBank accession numbers CRK29573 and CRK26624, respectively. These two

genes are present in *V. longisporum* only. Blasting the NCBI nr/nt database with the two

genes did not produce any hit. When Blasting the two genes against the wgs database,

eight hits with 100% query cover and 100% similarity were produced for both genes.

These eight hits were from the eight whole-genome sequenced *V. longisporum* strains,

respectively, and these eight strains represent all the whole-genome sequenced strains

available in the database (Verified Jan 22, 2024). Thus, based on the sequence analysis,

the two genes are specific to *V. longisporum* at the species level and ubiquitous at the

strain level. This is in contrast with the other two *Verticillium*-specific genes (CRK12486

and CRK45493) identified in our initial candidate selection, which are present only in the

six whole-genome sequenced A1/D1 strains (VLB2, VL20, VL2, VL1, Vl43 and Vl145c),

but not in the two A1/D3 strains (PD589 and Vl32). It is likely that CRK29573 and

CRK26624 are derived from the A1 parent and CRK12486 and CRK45493 are derived

from the D1 parent. The specificity of P-VL and LAMP-VL was further demonstrated by

the test on non-*V. longisporum* species. No signal was produced from any reaction.

Sensitivity of P-VL and LAMP-VL

As a diploid, *V. longisporum* carries two copies of most genes, with each copy from one

parent (Tran et al. 2013). The exception is the nuclear ribosomal region (rDNA), for

which only one type is present in each lineage, as a result of DNA loss and/or concerted

evolution (Depotter et al. 2016). For some genes, the two copies may have similar

sequences that can be detected by the same set of primers. On the other hand, for many

other genes, the two copies may have significant differences. This would result in a

primer set for one copy unable to detect the other copy, but also Blast search of one copy

would fail to identify the other one. In addition, there may be other genes that are present

in only one parent and no homolog in the other parent. These genes are the real single-

copied genes. We do not know whether CRK29573 and CRK26624 are real single-copied

genes or genes with their homologs having different sequences. Nevertheless, based on

the Blast results, these two genes are the single targets of P-VL and LAMP-VL,

respectively, in the *V. longisporum* genome. Thus, the sensitivity of P-VL and LAMP-VL

was expected to be lower than primers that target multiple-copied genes or DNA regions.

In the sensitivity test, the low limits for P-VL and LAMP-VL in a positive reaction were

28 copies of genomic DNA. Given the fact that the efficiency of DNA extraction from

mycelia was not 100%, the low limit of the fungal cells for a positive reaction would be

larger than 28. According to Yang et al. (2021), the extraction efficiencies of pathogen

DNA from plant tissue was approximately 50%. Thus, we expected that the P-VL and

LAMP-VL could detect *V. longisporum* DNA from as less as 56 cells. With the

consideration that one DNA extraction generally results in 50 µL of DNA and 2 µL is

used in one qPCR or LAMP reaction, the low limit of *V. longisporum* cell numbers in a

sample from which the pathogen can be detected by our qPCR or LAMP assay is 1,400.

The usefulness of the triplex qPCR and the LAMP

Two typical Verticillium stripe symptoms are dark unilateral stripes on the stems and

microsclerotia in the stem cortex (Depotter et al. 2016). We have never observed these

two symptoms on a single stem sample collected in our surveys which sampled randomly

across Alberta and represented approximately 1% of canola acres in each County or

Municipal District (n = approximately 380 fields, annually). In 2021-2023, canola stem

samples showing symptoms suspicious for Verticillium stripe, such as discoloration on

the internal stem tissues in a 'starburst' pattern, rather than sectoring, or unilateral

striping on stems, were observed and tested for the presence of *V. longisporum*. However,

the results were always negative (Harding et al. 2021; 2022; 2023). The results presented

herein are the first confirmation of *V. longisporum* in our random Provincial survey.

However, even on the sample 23CAN-305B from which *V. longisporum* was detected,

the symptoms were not different from the symptoms of other samples from which only

one or both of the two blackleg pathogens were detected. This was largely because this

sample was infected by not only *V. longisporum* but also the two blackleg pathogens

(Table 3). On the other hand, this also reflected the difficulty of symptom-based

diagnosis of Verticillium stripe, at least on Alberta samples. Therefore, the triplex qPCR

will greatly aid in *V. longisporum* detection from canola samples in Alberta, even when

the two blackleg pathogens are not the detection targets. In our canola survey conducted

in 2023, we tested 306 canola stem samples (including the twenty 23CAN samples in

Table 2) with the triplex qPCR. All these samples had similar symptoms as the sample

23CAN-305B. *Verticillium longisporum* was detected from one sample only (23CAN-

305B). In contrast, *L. biglobosa* or *L. maculans* was detected from 235 samples. These

data again emphasized the difficulty in detecting *V. longisporum* visually when blackleg

symptoms are present and the importance of using the triplex qPCR to test canola

samples with ambiguous symptoms to discriminate between blackleg and Verticillium

stripe.

The LAMP assay developed in this study can be used to confirm the results from the

qPCR. Our data indicated that the LAMP had a similar sensitivity as the qPCR.

Compared to qPCR, LAMP was believed to be more specific because LAMP targets six

DNA regions by the six primers (Notomi et al. 2000). LAMP also has the advantage of

being simple to conduct and no requirement for special equipment. Thus, LAMP was

expected to be useful for on-site diagnosis. However, successful LAMP reactions are

indeed highly dependent on the quality of template DNA, which has been overlooked by

many researchers. DNA extracted by many simplified protocols (i.e. meant to be

conducted in the field) employing high concentrations of detergents or alkalinity, should

be avoided when producing template DNA for LAMP assays. Therefore, at the current

time and for diagnostic labs equipped with qPCR cyclers, qPCR assays should be

considered preferable to LAMP. This recommendation is further highlighted in the case

of *V. longisporum* detection because successful extraction of good quality DNA from

canola stems by simplified methods is more difficult than from other plant tissues.

In summary, in this study, we developed a triplex qPCR system that can be used to detect

V. longisporum, *L. biglobosa,* and *L. maculans* in a single reaction. We confirmed the

usefulness of this system on canola samples collected from different locations in Alberta.

In addition, a LAMP assay was also developed, which can be used to verify the results

from the triplex qPCR. We recommend the use of this triplex qPCR/LAMP combo

system to others interested in detecting *V. longisporum* from plant samples.

References

CFIA (Canadian Food Inspection Agency). 2016. Results of national survey for

Verticillium longisporum. Available at: https://inspection.canada.ca/plant-

health/invasive-species/plant-diseases/verticillium-stripe/notice-to-

industry/eng/1458134997626/1458135081349. Accessed: January 2024.

Depotter, J. R., Deketelaere, S., Inderbitzin, P., Von Tiedemann, A., Höfte, M., Subbarao,

K. V., Wood, T. A., and Thomma, B. P. 2016. *Verticillium longisporum*, the invisible

threat to oilseed rape and other brassicaceous plant hosts. Mol. Plant Pathol. 17:1004-

1016.

Fu, H., Yang, Y., Sarkes, A., Harding, M. W., Feindel, D., and Feng, J. 2023.

Development of a duplex qPCR system for detection and quantification of the two

canola blackleg pathogens *Leptosphaeria biglobosa* and *L. maculans*. Plant Dis.

107:2808-2815.

Harding, M. W., Daniels, G. C., Burke, D. A., Pugh, C. A., Hill, T. B., Zahr, K., Sarkes,

A., and Feng, J. 2021. Canola and mustard disease survey in Alberta, 2020. Can. Plant

Dis. Survey 101:109-113.

Harding, M. W., Daniels, G. C., Hill, T. B., Kennedy, M. A., Sarkes, A., Yang, Y., and

Feng, J. 2022. Canola disease survey in Alberta, 2021. Can. Plant Dis. Survey

102:110-113.

Harding, M. W., Daniels, G. C., Hill, T. B., Xue, S., Sarkes, A., Yang, Y., and Feng, J.

2023. Canola disease survey in Alberta, 2022. Can. Plant Dis. Survey 103:110-112.

Ikeda K., Banno S., Watanabe K., Fujinaga M., Ogiso H., Sakai H., Tanaka H., Miki S.,

Shibata S., Shiraishi T., and Fujimura M. 2012. Association of *Verticillium dahliae*

and *Verticillium longisporum* with Chinese cabbage yellows and their distribution in

the main production areas of Japan. J. Gen. Plant Pathol. 78:331-337.

Inderbitzin, P., Davis, R.M., Bostock, R.M., and Subbarao, K.V. 2013. Identification and

differentiation of *Verticillium* species and *V. longisporum* lineages by simplex and

multiplex PCR assays. PLoS One 8:e65990.

Karapapa, V. K., Bainbridge, B. W., and Heale, J. B. 1997. Morphological and molecular

characterization of *Verticillium longisporum* comb. nov., pathogenic to oilseed rape.

Mycol. Res. 101:1281-1294.

Notomi, T., Okayama, H., Masubuchi, H., Yonekawa, T., Watanabe, K., Amino, N., and

Hase, T. 2000. Loop-mediated isothermal amplification of DNA. Nuc. Acids Res.

28:63-70.

Novakazi, F., Inderbitzin, P., Sandoya, G., Hayes, R.J., Tiedemann, A. V., and Subbarao,

K. V. 2015. The three lineages of the diploid hybrid *Verticillium longisporum* differ in

virulence and pathogenicity. Phytopathology 105:662-673.

Pantou, M. P., Strunnikova, O. K., Shakhnazarova, V. Y., Vishnevskaya, N. A.,

Papalouka, V. G., and Typas, M. A. 2005. Molecular and immunochemical

phylogenyof *Verticillium* species. Mycol. Res. 109:889-902.

Tran, V. T., Braus-Stromeyer, S. A., Timpner, C., and Braus, G. H. 2013. Molecular

diagnosis to discriminate pathogen and a pathogen species of the hybrid *Verticillium*

longisporum on the oilseed crop *Brassica napus*. Appl. Microbiol. Biotechnol.

97:4467-4483.

White, T. J., Bruns, T., Lee, S., Taylor, J. 1990. Amplification and direct sequencing of

fungal ribosomal RNA genes for phylogenetics. In: Innes, M. A., Gelfand, D. H.,

Sninsky, J. J., and White, T. J. (eds) PCR Protocols: A Guide to Methods and

Applications. San Diego, Academic Press, pp. 315-322.

Yang, Y., Zhou, Q., Zahr, K., Harding, M. W., Feindel, D., and Feng, J. 2021. Impact of

DNA extraction efficiency on the sensitivity of PCR-based plant disease diagnosis and

pathogen quantification. Eur. J. Plant Pathol. 159:583-591.

Yu, S., Su, T., Chen, J., Wang, W., Zhang, F., Yu, Y., Zhang, D., Zhao, X., and Lu, G.

2015. Real-time PCR as a diagnostic tool for evaluating the resistance of Chinese

cabbage cultivars to Verticillium wilt. Eur. J. Plant Pathol. 143:549-557.

Zou, Z., Bisht, V., and Fernando, W. D. 2020. Identification and characterization of

Verticillium longisporum lineage A1/D1 from Brassica crops in Manitoba, Canada.

Internat. J. Mol. Sci. 21:3499.

Table 1. Primers and probes used in this study

Name[a]		Sequences (5'-3')[b]
P-VL	F	CGAGTCGTACTTGGATGGCT
	P	/5Cy5/CGTCCACTT/TAO/TACATAGCACGG/3IAbRQSp/
	R	CCAGGGAGAAGATGAGCTGT
P-Lb	F	GAAGAATGGCAAAATCACAGG
	P	/56-FAM/AGGAAGAAG/ZEN/CAGCCATAGGC/3IABkFQ/
	R	AGCTCTGCGCGACCTTTT
P-Lm	F	CCTCACACTCTCGACCCCTA
	P	/5HEX/CACAGCCAT/ZEN/ATCATCCTGCA/3IABkFQ/
	R	GCATGTTCTTGAACCGCTAC
LAMP-VL	F3	ATGAACTTACGCTGTCGA
	B3	AAACAAGCCCACAACGAC
	FIP	AGACGAACGTGACAGCATGCTCCACCTCAACAGAACAATG
	BIP	GTGTGACAGCCTCAACGAATACCTGATTACAGACTGTGTT
	LF	GCCTTATAAACCTTATCTCTGCGC
	LB	GTCCGCCTTGTGTAGAAAAGAA
ITS	ITS1	TCCGTAGGTGAACCTGCGG
	ITS2	GCTGCGTTCTTCATCGATGC
P-Multi	D1f	CCCCGGCCTTGGTCTGAT
	AlfD1r	TGCCGGCATCGACCTTGG
	A1f	AAGTGGAGCCCCGTATCTTGAAT
	A1r	CAACTGGCAACAGGGCTTGAAT
	Df	CCGGTCCATCAGTCTCTCTG
	Dr	CTGTTGCCGCTTCACTCG

461 [a]References of primers: P-VL and LAMP-VL, this study; P-Lb and P-Lm, Fu et al.
462 (2023); ITS, White et al. (1990); P-Multi, Inderbitzin et al. (2013).
463 [b]F = forward primer; R = reverse primer; P = probe.
464

465   **Table 2.** Specificity test of the qPCR primer set P-VL on canola pathogens

Species	Strain[a]	Cq values of qPCR[b]	
		ITS1/ITS2	P-VL
Alternaria infectoria	APHL 002	24.2±0.1	NS
Alternaria solani	APHL 004	21.0±0.2	NS
Botrytis cinerea	APHL 199	19.0±0.1	NS
Colletotrichum coccodes	APHL 029	28.2±0.3	NS
Fusarium oxysporum	APHL 069	24.3±0.1	NS
Leptosphaeria biglobosa	APHL 216	21.1±0.1	NS
Leptosphaeria biglobosa	LB19-7	17.4±0.1	NS
Leptosphaeria biglobosa	APHL 1170	17.1±0.1	NS
Leptosphaeria biglobosa	LB19SK-11-2	18.0±0.1	NS
Leptosphaeria maculans	LM19-11	18.2±0.2	NS
Leptosphaeria maculans	LM19SK-11-21	18.3±0.1	NS
Leptosphaeria sclerotioides	APHL 101	18.2±0.1	NS
Phoma aliena	APHL 128	20.3±0.1	NS
Phoma betae	APHL 130	20.1±0.1	NS
Phoma medicaginis	APHL 134	19.4±0.1	NS
Plasmodiophora brassicae	Pathotype 5	25.8±0.1	NS
Pythium intermedium	APHL 159	18.0±0.2	NS
Rhizoctonia solani	APHL 234	22.9±0.2	NS
Sclerotinia sclerotiorum	APHL 244	19.7±0.1	NS
Stagonospora nodorum	APHL 120	21.6±0.1	NS
Trichoderma sp.	APHL 242	23.6±0.1	NS
Verticillium albo-atrum	APHL 190	18.3±0.2	NS
Verticillium albo-atrum	APHL 191	20.2±0.2	NS
Verticillium albo-atrum	APHL 192	18.3±0.1	NS
Verticillium dahliae	APHL 232	17.4±0.1	NS
Verticillium dahliae	APHL 303	23.0±0.2	NS
Verticillium lecanii	APHL 032	22.7±0.3	NS
Verticillium longisporum	APHL 189	21.5±0.2	25.8±0.1
Verticillium tricorpus	APHL 1117	22.8±0.1	NS
Negative control	Canola leaf	21.2±0.1	NS
Negative control	Canola root in LB[c]	24.2±0.1	NS
Negative control	Canola root in YPG[c]	26.6±0.1	NS
Negative control	Canola root	20.4±0.1	NS

[a]All strains belong to Alberta Plant Health Lab plant pathogen culture collection, a
project funded by the Canadian Agricultural Partnership (no. 601322) to JF except:
Lb19SK-11-2 and LM19SK-11-21 were provided by Dr. Gary Peng from Agriculture and

Agri-Food Canada, Saskatoon Research and Development Centre, and LB19-7 and

LM19-7 were purchased from University of Manitoba.

[b]SYBR Green qPCR using the primer set ITS (ITS1/ITS2) and probe-based qPCR using

the primer set P-VL. Quantification cycle (Cq) data were shown as mean ± standard

deviation (n = 3). NS = no signal from all the three replicates.

[c]DNA from microorganisms grown in LB broth or YPG media (1% yeast extract, 1%

peptone, 2% glucose, w/v) after incubating 200 mg healthy canola root (150 rpm, 30°C)

for 48 hours.

Table 3. Triplex qPCR and LAMP analyses on canola samples

Samples	Location	Cq values of triplex qPCR[a]			LAMP[b]
		P-Lb	P-Lm	P-VL	
2025A	Alberta	25.1±0.1	24.7±0.1	NS	-
2025B	Alberta	31.5±0.2	29.0±0.3	NS	-
2025C	Alberta	32.3±0.2	NS	NS	-
2025D	Alberta	26.5±0.1	NS	NS	-
23CAN-023	Alberta	NS	23.8±0.1	NS	-
23CAN-042	Alberta	NS	23.5±0.1	NS	-
23CAN-079	Alberta	28.0±0.1	33.2±0.4	NS	-
23CAN-083	Alberta	37.2±0.1	24.1±0.1	NS	-
23CAN-099	Alberta	28.3±0.1	25.1±0.1	NS	-
23CAN-135	Alberta	37.0±0.7	NS	NS	-
23CAN-152	Alberta	31.6±0.2	NS	NS	-
23CAN-197	Alberta	24.8±0.1	28.4±0.1	NS	-
23CAN-199	Alberta	27.0±0.1	NS	NS	-
23CAN-207	Alberta	30.3±0.1	38.5±0.9[c]	NS	-
23CAN-207	Alberta	28.3±0.1	27.6±0.1	NS	-
23CAN-305B	Alberta	26.9±0.1	26.7±0.1	30.0±0.1	+
23CAN-305C	Alberta	29.7±0.1	23.0±0.1	NS	-
23CAN-310	Alberta	NS	25.5±0.1	NS	-
23CAN-321	Alberta	37.2±0.8[c]	23.0±0.1	NS	-
23CAN-325	Alberta	24.2±0.1	37.2[c]	NS	-
23CAN-333A	Alberta	NS	25.3±0.1	NS	-
23CAN-342	Alberta	36.4±0.6	NS	NS	-
23CAN-348	Alberta	24.2±0.1	NS	NS	-
23CAN-363	Alberta	31.4±0.1	22.2±0	NS	-
MB 1	Manitoba	NS	22.7±0.1	NS	-
MB 2	Manitoba	24.1±0.1	NS	23.2±0.1	+
Negative control[d]		NS	NS	NS	-
Water		NS	NS	NS	-

[a]Triplex probe-based qPCR using the primer sets P-Lb, P-Lm and P-VL. Quantification
cycle (Cq) data were shown as mean ± standard deviation (n = 3). NS = no signal from all
the three replicates.
[b]LAMP analysis using the primer set LAMP-VL. +, positive reaction; -, negative reaction.
[c]Only one (without a standard deviation value) or two (with a standard deviation value)
of the three technical replicates generated Cq values.
[d]DNA from root of a 14-day old healthy canola seedling.
486

Fig. 1. Geographic locations in Alberta counties of the 24 canola stem samples that were collected for *Verticillium longisporum* detection. In each county, the number of samples was indicated.

Fig. 2. Sensitivity of the qPCR primer set P-VL. a, on gBlock DNA. b, on *Verticillium longisporum* genomic DNA. The qPCR standard curves were generated from the mean of quantification cycle (Cq) values against log10 of gBlock double-stranded DNA copies (a) or picogram of genomic DNA (b) in one reaction. The R^2 score of the equation and the efficiency of the primers (E) are indicated over the curve. Efficiency was calculated as E = $-1+10^{(-1/slope)}$. Each data point is shown as mean of three technical replicates ± standard deviation.

Fig. 3. Sensitivity of the LAMP primer set LAMP-VL. The LAMP reactions were conducted against a set of serial dilutions of *Verticillium longisporum* genomic DNA, from 1 ng per reaction to 10 fg per reaction. Changing of the color of the reaction mixture from red to yellow indicated a positive reaction. Each tube was also indicated by a + for positive reaction or a – for negative reaction.

Fig. 4. Sensitivity of the triplex qPCR system for the detection of *Verticillium longisporum* by the primer set P-VL (a), *Leptosphaeria biglobosa* by the primer set P-Lb (b) and *L. maculans* by the primer set P-Lm (c) in one reaction. The qPCR standard curves were generated from the mean of quantification cycle (Cq) values against log10 of picograms of serial dilutions of genomic DNA from each of the three fungi. The R^2 score of the equation and the efficiency of the primers (E) are indicated over the curve. Efficiency was calculated as $E = -1 + 10^{(-1/slope)}$. Each data point is shown as mean of three technical replicates ± standard deviation.

Fig. 5. Analysis of selected fungal strains and canola samples by a multiplex PCR system developed by Inderbitzin et al. (2013). The reactions were conducted in Promega Go Taq Master Mix (a) or Thermo Scientific Phire Plant Direct PCR Master Mix (b). M, GeneRuler Express DNA Ladder. 1 and 2, *Verticillium dahliae* isolates APHL 232 and APHL 303, respectively. 3, *V. longisporum* isolate APHL 189. 4, The canola sample from which APHL 189 was isolated. 5, Canola sample 23CAN-305B. 6, Canola sample MB 2. 7, Canola sample 23CAN023. 8, Healthy canola.