# A microRNA that controls the emergence of embryonic movement

Jonathan A. C. Menzies, Andre M. Chagas and Claudio R. Alonso[*]

*Department of Neuroscience,*
*Sussex Neuroscience,*
*School of Life Sciences,*
*University of Sussex,*
*Brighton BN1 9QG,*
*UK*

*(\*)      To whom all correspondence should be addressed*

Tel:     +44 1273 876621 (office)
          +44 794 493 0572 (mobile)

Email: c.alonso@sussex.ac.uk

# SUMMARY

Movement is a key feature of animal systems, yet its embryonic origins are not fully understood. Here we investigate the genetic basis underlying the embryonic onset of movement in *Drosophila* focusing on the role played by small non-coding RNAs (microRNAs, miRNAs). To this end, we first develop a quantitative behavioural pipeline capable of tracking embryonic movement in large populations of fly embryos, and using this system, discover that the *Drosophila* miRNA *miR-2b-1* plays a role in the emergence of movement. Through the combination of spectral analysis of embryonic motor patterns, cell sorting and RNA *in situs*, genetic reconstitution tests, and neural optical imaging we define that *miR-2b-1* influences the emergence of embryonic movement by exerting actions in the developing nervous system. Furthermore, through the combination of bioinformatics coupled to genetic manipulation of miRNA expression and phenocopy tests we identify a previously uncharacterised (but evolutionarily conserved) chloride channel encoding gene – which we term *Janus* – as a genetic target that mechanistically links *miR-2b-1* to the onset of movement. Cell-specific genetic reconstitution of *miR-2b-1* expression in a null miRNA mutant background, followed by behavioural assays and target gene analyses, suggest that *miR-2b-1* affects the emergence of movement through effects in sensory elements of the embryonic circuitry, rather than in the motor domain. Our work thus reports the first miRNA system capable of regulating embryonic movement, suggesting that other miRNAs are likely to play a role in this key developmental process in *Drosophila* as well as in other species.

**Keywords:** *Drosophila, embryo, movement, microRNA, chloride channel, neuron*

**RESULTS AND DISCUSSION**

*A miRNA that impacts larval movement*

Movement is the main output of the nervous system allowing animals to walk, fly, crawl, swim and maintain their posture, so that they can find prey, mate partners, escape predators and relocate within habitats. Despite the key biological and adaptive roles of movement across animal systems, how developing embryos manage to organise the necessary molecular, cellular, and physiological processes to initiate patterned movement is still unknown. Although it is clear that the genetic system plays a role, how genes control the formation, maturation and function of the cellular networks underlying the emergence of motor control remains poorly understood.

Recent work in our laboratory has shown that miRNAs – which are short regulatory non-coding RNAs that repress the expression of target genes [1, 2] – have pervasive roles in the articulation of complex movement sequences such as those involved in body posture control in the young *Drosophila* larva [3-6]; these observations, as well as those from others in *Drosophila* and other systems [7-11] hinted at the possibility that these non-coding RNA molecules might also be involved in the control of more fundamental aspects of motor development and control. To explore this question, we first searched for miRNAs that might affect the simple locomotor patterns of the *Drosophila* first instar larva. The L1 larva which is a convenient model to investigate the genetics of movement given that: (i) the assembly of the machinery for movement must be fully completed by this developmental stage so as to satisfactorily propel the animal into the external world, and (ii) if analysed sufficiently early, for example during the first few minutes after hatching, the animal had no real chance of compensating or learning ways around a putative defect, increasing the probability of detection by means of a suitable motor test. To extract a signature of larval movement, we applied a whole animal imaging method based on frustrated total internal reflection (FTIR) [12, 13] which renders high resolution and high contrast movies to both normal and miRNA mutant first instar larvae. This led us to discover that a single miRNA, *miR-2b-1*, had a significant impact on larval movement (Figure 1B-1C). *miR-2b-1* belongs to the *miR-2* family [14] and *ΔmiR-2b-1* mutant larvae show a

substantial decrease in larval speed (Figure 1C) suggesting that absence of this specific genetic component compromises the ability of the larva to move normally. Given that these larval tests were conducted within a 30-min period post-hatching, we considered the possibility that the defects observed in the larvae stemmed from changes in earlier ontogenetic processes that occur in the embryo. More specifically, we decided to test the possibility that early embryonic movement patterns might be affected by the lack of normal expression of *miR-2b-1*.

Previous work had provided an excellent first characterisation of the onset of embryonic movement patterns in wild type embryos [15, 16]; these early studies were based on the manual annotation of representative videotaped muscle contractions [16] or GFP-labelled muscle Z-lines analysed under spinning disc confocal microscopy [15]. Despite their attributes, these approaches were highly labour intensive and lacked the necessary throughput required to simultaneously analyse large numbers of embryos enabling a sensitive genetic screen. In consequence, we developed a new automated approach capable of quantifying movement in large populations of embryos.

*A high throughput approach to quantify movement in Drosophila embryos*

To monitor the onset of embryonic movement, which, in normal embryos, occurs during the final third of embryogenesis (i.e. 14-16h after egg laying (AEL) [15, 16] [Figure 1D-1E] we developed a 3D printed chamber system capable of hosting multiple embryos submerged under a thin layer of halocarbon oil to ensure adequate oxygenation and hydration (Figure 1F-1G) compatible with digital imaging by a charge coupled device (CCD) camera (Figure 1H). The camera captures the reflected light after its physical contact with the embryo; in this setting, even a subtle movement performed by the embryo results in a change in the path of reflected light, leading to variations in signal intensity detected by individual pixels in the CCD sensor, allowing for an accurate measurement of embryonic movement (Figure 1I-1J). To extract quantitative movement information from individual embryos we applied an image segmentation protocol to define regions of interest (ROIs) corresponding to each embryo and collected pixel intensity values from all ROIs at 4 frames/sec (Figure 1K-1L). The data

allow us to plot variations in average grey pixel intensity over embryonic time, which provide a quantitative signature of the ontogeny of movement in the individual embryo (Figure 1M). From this, we were able to observe distinct phases of movement that are consistent with previous data: namely, the onset of a phase of disorganised movements ~16 hours after egg laying (hAEL) and its transition into a phase characterised by rhythmic bursts of activity and inactivity ~18 hAEL [15, 16]. These phases have been termed as 'myogenic' and 'neurogenic' based on their respective dependence on neural input (Figure 1N) [15, 17-19] , and have been observed in *Drosophila*, as well as in other systems, including vertebrates [20-24] strongly suggesting that this is a general feature of motor development. See Movie S1 for a high resolution recording of *Drosophila* embryo movement.

To establish whether the readings of motor activity at the neurogenic phase detected by our approach were indeed dependent on neural activity, we expressed the inwardly rectifying potassium channel (*Kir*) [25] in all embryonic neurons using the pan-neuronal driver *elav-Gal4* seeking to suppress action potentials across all embryonic neuronal types. The results of this experiment show that whilst movement patterns during the early chaotic phase remain unchanged by this treatment (Figure 2C, 2E), motor activity at the rhythmic phase is almost completely eliminated (Figure 2C, 2F) strongly indicating that the emergence of this latter phase depends on normal neural activity. This is in agreement with previous observations of the effects of embryonic neural activity inhibition by other methods [15, 18]. In addition, spectral analysis demonstrates that the movement frequencies that characterise the rhythmic phase (Figure 2B, 2G) do not emerge in *Kir* embryos (Figure 2D, 2G). Altogether these observations suggest that *miR-2b-1* might exert its roles on embryonic movement, at least in part, due to action within the developing embryonic nervous system.

### *The miRNA miR-2b-1 affects embryonic movement patterns*

Given that the machinery for larval movement is assembled during embryogenesis [26-29] we considered the possibility that *miR-2b-1* might have an impact on the emergence of movement in the fly embryo. To explore this, we applied the approach described above to normal and *ΔmiR-2b-1*

mutant embryos (Figure 2H, 2J). These experiments showed that *ΔmiR-2b-1* mutant embryos displayed a different pattern of embryonic movement when compared with their wild type counterparts. Although the distinct phases of embryonic movement are clearly recognisable in mutant embryos, the overall amount of movement appeared greatly reduced (Figure 2J).

Indeed, comparison of quantity of movement in wild type and *ΔmiR-2b-1* mutant embryos either during the earlier myogenic phase (Figure 2L) or during the neurogenic phase (Figure 2M) shows significantly lower levels of movement in mutants. Furthermore, spectral analyses of embryonic movement traces reveal that miRNA mutant embryos shift to a higher frequency of movement bouts during the rhythmic phase when compared to normal embryos (Figure 2I, 2K) and that average bout length is also shortened (Figure S1).

### *miR-2b-1 expression and roles in the embryonic nervous system*

The fact that removal of *miR-2b-1* impacts the neurogenic phase of embryonic movement suggests that this miRNA might be expressed in the nervous system and exert a functional role there. To further explore this model, we first conducted a spatial expression analysis in the developing embryo using fluorescence RNA *in situ* hybridisation (FISH). Taking advantage from the fact that *miR-2b-1* is a 'mirtron' [30] we prepared an *in situ* probe against *miR-2b-1*'s host gene *Btk*, which encodes the Bruton tyrosine kinase [31, 32] (Figure 3A). These FISH experiments show that the *miR-2b-1* precursor miRNA transcript (pre-miRNA) [2] is expressed in multiple embryonic tissues, including the CNS (Figure 3B). To further confirm the expression of *miR-2b-1* in the nervous system, we conducted fluorescence-activated cell sorting (FACS, Figure 3C) to isolate embryonic neuronal samples (labelled by means of *elav>GFP*) followed by RT-PCR for the mature miRNA transcript and observed *miR-2b-1* specific signal (Figure 3D). Therefore, the results of these two distinct and complementary methods provide strong evidence of neural expression of the miRNA.

To gain more insight on the neural roles of *miR-2b-1* in regard to embryonic movement, we conducted a genetic reconstitution experiment in which we analysed the consequences of restoring *miR-2b-1*

expression in the nervous system in an otherwise *ΔmiR-2b-1* null mutant (Figure 3E). Results in Figure 3F show that *elav*-driven expression of *miR-2b-1* in a *ΔmiR-2b-1* mutant background leads to a phenotypic rescue, producing embryos that display statistically indistinguishable movement patterns to those recorded in control embryos, indicating that neural expression of *miR-2b-1* is sufficient to restore a normal onset of embryonic movement. To further examine the biological roles of neural *miR-2b-1* expression, we assessed the impact of restoring miRNA expression in embryonic neurons on first instar larval locomotor patterns using the FIM approach described above (Figure 1B-C) and observed that when *ΔmiR-2b-1* mutant larvae are developmentally provided with pan-neuronal *miR-2b-1* expression, the characteristic miRNA larval mutant phenotype is rescued, with specimens moving at natural speed (Figure 3G, S2A). The experiments described above strongly indicate that expression of *miR-2b-1* in the nervous system is biologically relevant and sufficient to rescue the embryonic and larval movement defects observed in *ΔmiR-2b-1* mutants. In turn, this suggests that absence of *miR-2b-1* must impinge a morphological and/or a functional deficit in the developing nervous system of the embryo. To tease apart these potential biological effects, we examined the structure of the nervous system in normal and *ΔmiR-2b-1* mutant embryos by means of immunohistochemistry and confocal microscopy and observed no detectable differences (Figure S2B). In contrast, analysis of neural activity patterns in the embryo by means of GCaMP6 functional imaging using a movement distortion correction approach (i.e. tdTomato, Figure 3H-3J [17]), shows that miRNA mutant embryos have a reduced level of calcium dynamics when compared with their control counter parts (Figure 3K-3M); notably, this occurs during a previously identified 'critical period' when neural activity levels are of crucial importance to the development of stable neural circuits [33-35].

***A genetic link between miR-2b-1 and embryonic movement***

Our gene expression, genetic reconstitution, morphological and functional imaging data support a model in which *miR-2b-1* plays a physiological role in the developing embryonic nervous system.

This raises the question of how might this regulatory miRNA system interact with the physiological control of the neuron during embryogenesis. To explore the genetic elements that link *miR-2b-1* to its role in embryonic movement we searched for candidate *miR-2b-1* target genes using the ComiR bioinformatic platform (Figure 4B) [36, 37]. A common issue with bioinformatic predictions of miRNA targets is the generation of false positives; in this regard, ComiR integrates multiple miRNA target prediction algorithms – each one with its intrinsic strengths and weaknesses[38-41] – seeking to identify a set of consistent *bona fide* miRNA targets that satisfy the filters of multiple algorithms, thus reducing the generation of false positives[36, 37]. Applying ComiR to *miR-2b-1* produced a list of high probability targets organised in the form of an ascending ranking (Figure 4B). At the very top of the list was *CG3638*, an uncharacterised *Drosophila* gene predicted to encode a chloride channel protein; this highly ranked target was of interest to us because of its potential role in the physiological control of anionic conductances, and its broad evolutionary conservation across insects and mammals[42], including humans (Figure 4E-I).

A genuine genetic target for a given miRNA is predicted to: (i) be de-repressed (up-regulated) in the absence of the miRNA; and (ii) be down-regulated under miRNA ectopic expression. Analysis of *CG3638* expression shows that this target meets the predictions of a genuine *miR-2b-1* target in full: expression of *CG3638* is upregulated in *ΔmiR-2b-1* mutants (Figure 4B) and reduced under neural over-expression of *miR-2b-1* (Figure 4D). As mentioned above, phylogenetic analysis of *CG3638* reveals that it belongs to an evolutionarily conserved family of chloride channel genes[42, 43], with representatives across distantly related lineages of insects and vertebrates including mammals, strongly indicating a functional role (Figure 4E). Comparison of the properties of *CG3638* and its human orthologue reveal the characteristic seven trans-membrane domains with an external carboxyterminal topology (Figure 4F, 4H) further supporting orthology, and applying *AlphaFold* – an artificial intelligence computational method able to predict protein structures with atomic accuracy [44] – to the proteins encoded by the *Drosophila CG3638* gene and the human TTYH1 gene reveals the marked similarities between these two polypeptides (Figure 4G, 4I).

To explore the relationship between *CG3638* and the embryonic movement phenotype displayed by *ΔmiR-2b-1* mutants, we tested the effects of an artificial reduction of *CG3638* in the genetic background of the miRNA mutant (Figure 4K). In this scenario, should the levels of expression of *CG3638* be relevant to the triggering of the embryonic movement phenotype, we predict that a reduction in *CG3638* expression levels should compensate its cellular effects, and, accordingly, reduce or even rescue the embryonic phenotype. The results of this experiment show that this is indeed the case, with embryonic movement of the *ΔmiR-2b-1* mutant effectively rescued by a reduction in *CG3638* (Figure 4J). Based on this role of embryonic movement we termed *CG3638* as *Janus*, making reference to the Roman god of movement and the symbol of change and transitions.

*Mapping the focus of action of miR-2b-1 within the known networks underlying embryonic movement*

Having observed the effects of *miR-2b-1* on embryonic movement we wondered about the site of action of this miRNA in regard to circuit components previously linked to embryonic movement. In this respect, previous work has identified embryonic motor neuronal components, as well as interneurons and elements of the sensory system as playing key roles in the control of motor development. These include a motor component that includes all embryonic motor neurons which command the stereotypic array of muscles in the embryonic body wall [45, 46], as well as specific elements of the sensory system – in particular the chordotonal system – which detect early myogenic movements in the embryo and transmit the information to the pattern generators thus modulating motorpatterns (Figure 4L) [17, 19, 47, 48]. To determine which one of these known circuitry elements might be the principal focus of action of *miR-2b-1* in connection to embryonic movement control we artificially expressed *miR-2b-1* in each motor and sensory circuit elements – using drivers *OK371-Gal4* [49] and *iav-Gal4* [50], respectively – in an otherwise null mutant background for *miR-2b-1*, asking whether these genetic restorations were sufficient to improve or perhaps even rescue normal

movement patterns. To ensure that circuit-specific Gal-4 drivers were active and UAS-driven miRNA levels achieved the necessary cumulative values for biological activity we chose to measure effects on early larval movement patterns tested in 30-min old first instar larvae (L1s). The results of these experiments are shown in Figure 4M-N M-O. Here we observe that restoring expression of *miR-2b-1* in the motor neuronal domain defined by the OK371 driver is insufficient to affect the defective movement patterns observed in *miR-2b-1* null mutants (Figure 1B-C and Figure 4M). In contrast, re-establishing expression of the miRNA in all eight chordotonal organs leads to a full rescue of the motor phenotype (Figure 4N) suggesting that this aspect of the sensory system might be the one where *miR-2b-1* exerts relevant actions during the normal development of movement. In line with this, we observe that the mature *miR-2b1* transcript is indeed expressed in FACS-isolated embryonic chordotonal organs prepared from wild type embryos (Figure 4P) and, that the genetic target of *miR-2b-1*, *Janus*, is also expressed in these cells in normal embryos (Figure 4Q). In addition, expression of *Janus* in chordotonal organs prepared from *miR-2b1* null mutants is upregulated (Figure 4Q) lending further support to a model in which *Janus* is de-repressed in these specific sensory elements. Furthermore, artificial reduction of *Janus* implemented as a stratagem to decrease the effects of de-repression specifically within the chordotonal system is sufficient to rescue normal movement patterns (Figure 4R). Altogether, these findings strongly suggest that *miR-2b-1* impacts the emergence of embryonic movement, at least in part, via effects on the sensory circuit components that underlie motor development, rather than affecting the actual generation of motor patterns.

Our work identifies a miRNA system that plays a role in the emergence of embryonic movement in the fly embryo, and offers a new approach to analyse the roles of non-coding RNAs and protein coding genes at the critical period when patterned movement develops. Understanding the molecular elements controlling the onset of motor development in *Drosophila* will put us one step closer to understanding the molecular basis of embryonic movement in other species, including vertebrates, whose embryos seem to undergo remarkably similar transition phases to those reported here [20].

## ACKNOWLEDGEMENTS

We wish to thank Tom Baden for their contributions to the development of the embryonic movement pipeline and computational analysis of movement patterns, respectively. We also thank all members of the Alonso Lab for helpful discussions and feedback on this work. We are very grateful to Marta Zlatic for sharing fly lines, and to the Bloomington Stock Center and the Vienna *Drosophila* Resource Center for providing fly stocks. This research was funded by a Wellcome Trust Investigator Award (098410/Z/12/Z) made to C.R.A., and a UK Medical Research Council Project Grant (Ref: MR/S011609/1) given to C.R.A.

## COMPETING INTERESTS

The authors declare no competing interests.

**FIGURE LEGENDS**

**Figure 1**

*A novel approach for the quantification of embryonic movement*

(A) Schematic describing miRNA action upon target genes to control cell biology and behaviour.

(B) Larval movement tracks for $w^{1118}$ (left) and *ΔmiR-2b-1* (right).

(C) Quantification of larval speed.

(D) Timeline of *Drosophila* embryonic movement.

(E) *Drosophila* embryo performing characteristic movements.

(F-H) Experimental pipeline for recording embryo movements.

(I-J) Graphic description of how movements are detected by the experimental setup.

(K-M) Analysis pipeline for the assessment of embryonic movements.

(N) Schematic of an embryo movement trace.

**Figure 2**

*miR-2b-1 controls movements during a neurogenic phase of embryonic movement*

(A) Representative control (*UAS-Kir*) embryo movement trace.

(B) Control movement frequency spectrogram (FFT).

(C) Representative experimental (*elav-Gal4>UAS-Kir*) embryo movement trace.

(D) Experimental movement frequency spectrogram (FFT).

(E) Quantification of myogenic phase movement.

(F) Quantification of neurogenic phase movement.

(G) Distribution of movement across different periodicities, derived from the FFT frequency analysis

shown in panels B and D. Different bars within each period are individual embryos.

(H) Representative control ($w^{1118}$) embryo movement trace.

(I) Control movement frequency spectrogram (FFT).

(J) Representative experimental (*ΔmiR-2b-1*) embryo movement trace.

(K) Experimental movement frequency spectrogram (FFT).

(L) Quantification of myogenic phase movement.

(M) Quantification of neurogenic phase movement.

(N) Distribution of movement across different periodicities, derived from the FFT frequency analysis shown in panels I and K. Different bars within each period are individual embryos.

**Figure 3**

*miR-2b-1 acts within neurons to regulate embryonic and larval phenotypes*

(A) Gene region of *miR-2b-1* with fluorescence *in-situ* hybridisation (FISH) probe for host *Btk*.

(B) FISH experiment.

(C) Fluorescence activated cell sorting (FACS) workflow.

(D) PCR for reference gene *RP49* (left) and mature *miR-2b-1* transcripts (right) from embryonic neurons.

(E) Graphic describing genetic reconstitution of *miR-2b-1* in neurons.

(F) Quantification of neurogenic phase movement.

(G) Quantification of larval speed.

(H) Schematic of fluorescence imaging of calcium dynamics in embryos.

(I) Representative GCaMP6s trace from control ($w^{1118}$) embryos.

(J) Representative tdTomato trace from the same embryo as in panel I.

(K) Representative ΔF/F trace for control ($w^{1118}$) embryos.

(L) Representative ΔF/F trace for (*ΔmiR-2b-1*) mutant embryos.

(M) Quantification of sum ΔF/F signal.

**Figure 4**

*The genetic and cellular mechanisms that link miR-2b-1 to embryonic movement*

(A) Experimental workflow for FACS and RT-qPCR to assess ComiR miRNA target predictions.

(B) Expression (RT-qPCR) of top 10 potential *miR-2b-1* target genes.

(C) *CG3638* transcript with *miR-2b-1* target sites in the 3' untranslated region (UTR) indicated by orange bars.

(D) Expression (RT-qPCR) of *CG3638* in whole embryos over-expressing *miR-2b-1* in neurons.

(E) Evolutionary conservation of *CG3638*.

(F, H) Transmembrane domains of *Homo sapiens* TTYH1 protein (G) and *Drosophila CG3638* proteins (H).

(G, I) Atomic protein structures for *Homo sapiens* TTYH1 protein (G) and *Drosophila CG3638* protein (I).

(J) Quantification of neurogenic phase movement under genetic inhibition of *CG3638* in all neurons.

(K) Expression (RT-qPCR) of *CG3638* under RNAi.

(L) Schematic of a feedback loop for activity dependent motor-development.

(M) Quantification of larval speed for *miR-2b-1* reconstitution in motor neurons.

(N) Quantification of larval speed for *miR-2b-1* reconstitution in chordotonal organs.

(O) Workflow for FACS isolation of chordotonal organs.

(P) PCR for reference gene *RP49* (left) and mature *miR-2b-1* transcripts (right) from embryonic chordotonal organs.

(Q) Expression (RT-qPCR) of *Janus* (*CG3638)* in chordotonal organs in the *ΔmiR-2b-1* background compared to wild-type.

(R) Quantification of larval speed under genetic inhibition of *Janus* (*CG3638)* specifically in chordotonal organs.

(S) Model for the mechanism by which *miR-2b-1* acts to control embryonic and larval movement.

**REFERENCES**

1.  Alonso, C.R. (2012). A complex 'mRNA degradation code' controls gene expression during animal development. Trends Genet 28, 78-88.

2.  Bartel, D.P. (2018). Metazoan MicroRNAs. Cell 173, 20-51.

3.  Issa, A.R., Picao-Osorio, J., Rito, N., Chiappe, M.E., and Alonso, C.R. (2019). A Single MicroRNA-Hox Gene Module Controls Equivalent Movements in Biomechanically Distinct Forms of Drosophila. Curr Biol 29, 2665-2675 e2664.

4.  Klann, M., Issa, A.R., Pinho, S., and Alonso, C.R. (2021). MicroRNA-Dependent Control of Sensory Neuron Function Regulates Posture Behavior in Drosophila. J Neurosci 41, 8297-8308.

5.  Picao-Osorio, J., Johnston, J., Landgraf, M., Berni, J., and Alonso, C.R. (2015). MicroRNA-encoded behavior in Drosophila. Science 350, 815-820.

6.  Picao-Osorio, J., Lago-Baldaia, I., Patraquim, P., and Alonso, C.R. (2017). Pervasive Behavioral Effects of MicroRNA Regulation in Drosophila. Genetics 206, 1535-1548.

7.  Fricke, C., Green, D., Smith, D., Dalmay, T., and Chapman, T. (2014). MicroRNAs influence reproductive responses by females to male sex peptide in Drosophila melanogaster. Genetics 198, 1603-1619.

8.  Holm, A., Possovre, M.L., Bandarabadi, M., Moseholm, K.F., Justinussen, J.L., Bozic, I., Lemcke, R., Arribat, Y., Amati, F., Silahtaroglu, A., et al. (2022). The evolutionarily conserved miRNA-137 targets the neuropeptide hypocretin/orexin and modulates the wake to sleep ratio. Proc Natl Acad Sci U S A 119, e2112225119.

9.  Kadener, S., Menet, J.S., Sugino, K., Horwich, M.D., Weissbein, U., Nawathean, P., Vagin, V.V., Zamore, P.D., Nelson, S.B., and Rosbash, M. (2009). A role for microRNAs in the Drosophila circadian clock. Genes Dev 23, 2179-2191.

10. Lackinger, M., Sungur, A.O., Daswani, R., Soutschek, M., Bicker, S., Stemmler, L., Wust, T., Fiore, R., Dieterich, C., Schwarting, R.K., et al. (2019). A placental mammal-specific microRNA cluster acts as a natural brake for sociability in mice. EMBO Rep 20.

11. Weng, R., Chin, J.S., Yew, J.Y., Bushati, N., and Cohen, S.M. (2013). miR-124 controls male reproductive success in Drosophila. Elife 2, e00640.

12. Risse, B., Berh, D., Otto, N., Klambt, C., and Jiang, X. (2017). FIMTrack: An open source tracking and locomotion analysis software for small animals. PLoS Comput Biol 13, e1005530.

13. Risse, B., Thomas, S., Otto, N., Lopmeier, T., Valkov, D., Jiang, X., and Klambt, C. (2013). FIM, a novel FTIR-based imaging method for high throughput locomotion analysis. PLoS One 8, e53963.

14. Marco, A., Hooks, K., and Griffiths-Jones, S. (2012). Evolution and function of the extended miR-2 microRNA family. RNA Biol 9, 242-248.

15. Crisp, S., Evers, J.F., Fiala, A., and Bate, M. (2008). The development of motor coordination in Drosophila embryos. Development 135, 3707-3717.

16. Pereanu, W., Spindler, S., Im, E., Buu, N., and Hartenstein, V. (2007). The emergence of patterned movement during late embryogenesis of Drosophila. Dev Neurobiol 67, 1669-1685.

17. Carreira-Rosario, A., York, R.A., Choi, M., Doe, C.Q., and Clandinin, T.R. (2021). Mechanosensory input during circuit formation shapes Drosophila motor behavior through patterned spontaneous network activity. Curr Biol 31, 5341-5349 e5344.

18. Crisp, S.J., Evers, J.F., and Bate, M. (2011). Endogenous patterns of activity are required for the maturation of a motor network. J Neurosci 31, 10445-10450.

19. Zeng, X., Komanome, Y., Kawasaki, T., Inada, K., Jonaitis, J., Pulver, S.R., Kazama, H., and Nose, A. (2021). An electrically coupled pioneer circuit enables motor development via proprioceptive feedback in Drosophila embryos. Curr Biol 31, 5327-5340 e5325.

20. Hamburger, V. (1963). Some Aspects of the Embryology of Behavior. Quarterly Review of Biology 38, 342-365.

21. Hamburger, V., and Balaban, M. (1963). Observations and experiments on spontaneous rhythmical behavior in the chick embryo. Dev Biol 6, 533-545.

22. Hamburger, V., Balaban, M., Oppenheim, R., and Wenger, E. (1965). Periodic motility of normal and spinal chick embryos between 8 and 17 days of incubation. J Exp Zool 159, 1-13.

23. Ripley, K.L., and Provine, R.R. (1972). Neural correlates of embryonic motility in the chick. Brain Res 45, 127-134.

24. Saint-Amant, L., and Drapeau, P. (1998). Time course of the development of motor behaviors in the zebrafish embryo. J Neurobiol 37, 622-632.

25. Baines, R.A., Uhler, J.P., Thompson, A., Sweeney, S.T., and Bate, M. (2001). Altered electrical properties in Drosophila neurons developing without synaptic transmission. J Neurosci 21, 1523-1531.

26. Bate, M., and Martinez Arias, A. (1993). The Development of Drosophila melanogaster, (Plainview, N.Y.: Cold Spring Harbor Laboratory Press).

27. Clark, M.Q., Zarin, A.A., Carreira-Rosario, A., and Doe, C.Q. (2018). Neural circuits driving larval locomotion in Drosophila. Neural Dev 13, 6.

28. Landgraf, M., Jeffrey, V., Fujioka, M., Jaynes, J.B., and Bate, M. (2003). Embryonic origins of a motor system: motor dendrites form a myotopic map in Drosophila. PLoS Biol 1, E41.

29. Heckscher, E.S., Lockery, S.R., and Doe, C.Q. (2012). Characterization of Drosophila larval crawling at the level of organism, segment, and somatic body wall musculature. J Neurosci 32, 12460-12471.

30. Okamura, K., Hagen, J.W., Duan, H., Tyler, D.M., and Lai, E.C. (2007). The mirtron pathway generates microRNA-class regulatory RNAs in Drosophila. Cell 130, 89-100.

31. Theroux, S.J., and Wadsworth, S.C. (1992). Protein-tyrosine kinase activity of alternate protein products of the Drosophila Dsrc28C locus. FEBS Lett 311, 1-6.

32. Vincent, W.S., 3rd, Gregory, R.J., and Wadsworth, S.C. (1989). Embryonic expression of a Drosophila src gene: alternate forms of the protein are expressed in segmental stripes and in the nervous system. Genes Dev 3, 334-347.

33. Ackerman, S.D., Perez-Catalan, N.A., Freeman, M.R., and Doe, C.Q. (2021). Astrocytes close a motor circuit critical period. Nature 592, 414-420.

34. Baines, R.A., and Landgraf, M. (2021). Neural development: The role of spontaneous activity. Curr Biol 31, R1513-R1515.

35. Giachello, C.N., and Baines, R.A. (2015). Inappropriate Neural Activity during a Sensitive Period in Embryogenesis Results in Persistent Seizure-like Behavior. Curr Biol 25, 2964-2968.

36. Bertolazzi, G., Benos, P.V., Tumminello, M., and Coronnello, C. (2020). An improvement of ComiR algorithm for microRNA target prediction by exploiting coding region sequences of mRNAs. BMC Bioinformatics 21, 201.

37. Coronnello, C., and Benos, P.V. (2013). ComiR: Combinatorial microRNA target prediction tool. Nucleic Acids Res 41, W159-164.

38. Betel, D., Koppal, A., Agius, P., Sander, C., and Leslie, C. (2010). Comprehensive modeling of microRNA targets predicts functional non-conserved and non-canonical sites. Genome Biol 11, R90.

39. Kertesz, M., Iovino, N., Unnerstall, U., Gaul, U., and Segal, E. (2007). The role of site accessibility in microRNA target recognition. Nat Genet 39, 1278-1284.

40. McGeary, S.E., Lin, K.S., Shi, C.Y., Pham, T.M., Bisaria, N., Kelley, G.M., and Bartel, D.P. (2019). The biochemical basis of microRNA targeting efficacy. Science 366.

41. Miranda, K.C., Huynh, T., Tay, Y., Ang, Y.S., Tam, W.L., Thomson, A.M., Lim, B., and Rigoutsos, I. (2006). A pattern-based method for the identification of MicroRNA binding sites and their corresponding heteroduplexes. Cell 126, 1203-1217.

42. Suzuki, M. (2006). The Drosophila tweety family: molecular candidates for large-conductance Ca2+-activated Cl- channels. Exp Physiol 91, 141-147.

43. Suzuki, M., and Mizuno, A. (2004). A novel human Cl(-) channel family related to Drosophila flightless locus. J Biol Chem 279, 22461-22468.

44. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with AlphaFold. Nature 596, 583-589.

45. Landgraf, M., and Thor, S. (2006). Development of Drosophila motoneurons: specification and morphology. Semin Cell Dev Biol 17, 3-11.

46. Peron, S., Zordan, M.A., Magnabosco, A., Reggiani, C., and Megighian, A. (2009). From action potential to contraction: neural control and excitation-contraction coupling in larval muscles of Drosophila. Comp Biochem Physiol A Mol Integr Physiol 154, 173-183.

47. Caldwell, J.C., Miller, M.M., Wing, S., Soll, D.R., and Eberl, D.F. (2003). Dynamic analysis of larval locomotion in Drosophila chordotonal organ mutants. Proc Natl Acad Sci U S A 100, 16053-16058.

48. Hughes, C.L., and Thomas, J.B. (2007). A sensory feedback circuit coordinates muscle activity in Drosophila. Mol Cell Neurosci 35, 383-396.

49. Mahr, A., and Aberle, H. (2006). The expression pattern of the Drosophila vesicular glutamate transporter: a marker protein for motoneurons and glutamatergic centers in the brain. Gene Expr Patterns 6, 299-309.

50. Kwon, Y., Shen, W.L., Shim, H.S., and Montell, C. (2010). Fine thermotactic discrimination between the optimal and slightly cooler temperatures via a TRPV channel in chordotonal neurons. J Neurosci 30, 10465-10471.

**A** miRNA — miRNA regulatory networks — gene target expression level — Neural network computations — Movement

**B** $w^{1118}$ — $\Delta miR\text{-}2b\text{-}1$

**C** Larval speed ($\mu$m/second) — $w^{1118}$ — $\Delta miR\text{-}2b\text{-}1$ — ****

**D** Fertilisation — Embryogenesis — Tracheal gas-filling — Hatching — 0h — 5h — 10h — 15h — 21h

Period of movement

**E**

**F** Embryo dechorionation — Adhesion to imaging chamber

**G** Adhered embryos

**H** Leica DFC340FX — LED — LED

**I** CCD camera — $\theta_1$ — Reflected light

**J** CCD camera — $\theta_2$ — $\theta_1$

**K** Image segmentation and ROI assignment

**L** Tracheal gas-filling — Hatching — Pixel grey value extraction — Average pixel grey value — Hours after egg laying — Background (rolling average) subtraction, absolute values

**M** Embryo movement (grey value deviations from rolling average AU) — Control ($w^{1118}$) — Hours after egg laying

**N** NEUROGENIC PHASE — 51-100% post onset — MYOGENIC PHASE — 0-50% post onset — Movement onset — Movement bout — Muscle activity — Neuronal activity — Hours after egg laying — Hatching