Oviposition status and sugar availability influence measures of transmission

potential in malaria-infected mosquitoes.

Authors

Justine C. Shiau [1,2,3], Nathan Garcia-Diaz [4,5,8], Dennis E. Kyle [1,2,6,7], and Ashutosh K. Pathak [1,2,3,7*].

[1]Department of Infectious Diseases, University of Georgia, Athens, GA, USA

[2]Center for Tropical and Emerging Global Diseases (CTEGD), University of Georgia, Athens, GA, USA

[3]Center for the Ecology of Infectious Diseases, Athens, GA, USA

[4]The NSF-REU program, Odum School of Ecology, Athens, GA, USA

[5]Department of Biology, Willamette University, Salem, OR, USA

[6]Department of Cellular Biology, University of Georgia, Athens, GA, USA

[7]The SporoCore, CTEGD, University of Georgia, Athens, GA, USA

[8]Current address: Department of Epidemiology, Brown University School of Public Health, Providence,

RI, USA.

* Corresponding author: ash1@uga.edu.

Justine C Shiau: justine.shiau@uga.edu.

Nathan Garcia-Diaz: nathan_garcia-diaz@brown.edu

Dennis E Kyle: dennis.kyle@uga.edu

ABSTRACT

Background: Like other oviparous organisms, the gonotrophic cycle of mosquitoes is not complete until they have selected a suitable habitat to oviposit. In addition to the evolutionary constraints associated with selective oviposition behavior, the physiological demands relative to an organism's oviposition status also influences their nutrient requirement from the environment. Yet, studies that measure transmission potential (vectorial capacity or competence) of mosquito-borne parasites rarely consider if the rates of parasite replication and development could be influenced by these constraints resulting from whether mosquitoes have completed their gonotrophic cycle.

Methods: *Anopheles stephensi* mosquitoes were infected with *Plasmodium berghei* the rodent analog of human malaria and maintained on 1% or 10% dextrose and either provided oviposition sites ('oviposited' herein) to complete their gonotrophic cycle or forced to retain eggs ('non-oviposited'). Transmission potential in the four groups was measured up to 27 days post-infection as 1) the rates of vector survival, 2) rates of sporozoite migration to the salivary glands ('extrinsic incubation period' or EIP), and 3), sporozoite densities.

Results: In the two groups of oviposited mosquitoes, rates of sporozoite migration and densities in the salivary glands were clearly higher in mosquitoes fed 10% dextrose. In non-oviposited mosquitoes however, rates of sporozoite migration and densities were independent of sugar concentrations, although both measures were slightly lower than oviposited mosquitoes fed 10% dextrose. Rates of vector survival were higher in non-oviposited mosquitoes.

Conclusions: Taken together, these results suggest vectorial capacity for malaria parasites may be dependent on nutrient availability and oviposition/gonotrophic status and as such, argue for more careful consideration of this interaction when estimating transmission potential; costs to parasite fitness and vector survival were buffered against changes in nutritional availability from the environment in non-oviposited mosquitoes, but not oviposited mosquitoes. In general, however, these patterns suggest parasite fitness may be dependent on complex interactions between physiological (nutrition) and evolutionary (egg-retention) trade-offs in the vector, with potential implications for disease transmission and management. For instance, while reducing availability of oviposition sites and environmental sources of nutrition are key components of integrated vector management strategies, their abundance and distribution is under strong selection pressure from the patterns associated with climate change.

Keywords: Sporozoites, *Anopheles*, *Plasmodium*, Oviposition, Survival Rate, Malaria, Nutrients, Sugars, Mosquito Vectors, Follicular Atresia, Oviparity.

INTRODUCTION

Like other oviparous organisms, mosquitoes retain their eggs until they find a suitable habitat to oviposit and complete its gonotrophic cycle. Amongst oviparous organisms however, mosquitoes occupy an important niche: most species rely exclusively on hematophagy (blood feeding) of vertebrates for producing eggs, which enables them to vector several parasites with devastating consequences for the health and fitness of humans and wildlife. Following ingestion of an infected blood meal, egg and parasite development may be initiated at the same time in the mosquito. However, by the time a mosquito has completed egg development, parasite development may or may not be complete, although transmission to the next host is dependent on the mosquito locating an oviposition site to complete its 'gonotrophic cycle' [1]. While several studies suggest widespread changes in mosquito behavior and physiology associated with the gonotrophic cycle [2, 3, 4, 5, 6, 7, 8], few have determined if and how changes in oviposition status influences parasite development rates and transmission potential [9]; in general, measures of parasite fitness and transmission potential are performed in mosquitoes prevented from completing their gonotrophic cycle (e.g., see [10, 11, 12, 13] and references therein). A greater understanding of how parasite fitness is shaped by oviposition status of mosquitoes is important for at least two reasons. First, integrated vector management strategies either directly or indirectly alter the availability of oviposition sites, by managing water sources for instance, to identifying specific physical, chemical, and biological cues underlying selective oviposition behavior for targeting with chemical or biological means [1, 14, 15, 16, 17, 18, 19]. Second, the availability of oviposition sites will continue to change as a cumulative response to climatic factors such as global warming and changes in rainfall patterns to the increasing urbanization and deforestation [20, 21, 22, 23, 24]. By altering the number and distribution of oviposition habitats, both scenarios could select for variation in oviposition behavior and thus modify parasite/disease transmission in the process.

Like other oviparous organisms, adult, female mosquitoes also show selective oviposition behavior [1, 16]. Offspring survival and reproduction are benefited by ovipositing mothers choosing sites that fulfil several criteria, including for instance, habitat size, quality, risk of predation and presence of conspecifics. The ubiquity of selective oviposition behavior attests to its benefits to future

reproduction, with selection for this trait imposed at the cost of reduced fecundity for instance [25, 26,
27, 28, 29, 30]. While several studies have shown how variation in this behavior could modify the
oviparous organism's own fitness [26, 27, 28, 30, 31, 32, 33, 34, 35], few have assessed whether the
associated trade-offs could have downstream effects on the fitness of other organisms such as parasites,
whose life-history strategy is driven by its interaction with the oviparous host.

Organismal fitness is also dependent on how it allocates available resources from the
environment to meet the costs of various life-history traits [25, 26, 27, 29, 36]. For vector-transmitted
parasites, although its growth and reproduction are dependent on how they allocate resources from the
mosquito host, the host's own fitness is also dependent on how they allocate resources from
environmental sources of nutrition; in other words, nutritional availability from the environment can
shape life-history strategies of the vector and parasite [37, 38]. Amongst environmental sources of
nutrition, the effect of sugars on life-history traits of adult, female mosquitoes are particularly well
characterized, with their propensity for sugar feeding also exploited for controlling vector density (e.g.
toxic sugar baits) [39]; while the costs of parasite infection for the mosquito are more apparent at lower
concentrations of sugar, in general, sugar requirements are greater in gravid/non-oviposited mosquitoes
foraging for oviposition sites, carrying and/or maintaining eggs [25, 38, 40, 41, 42]. Differences in sugar
concentrations and/or availability to the mosquito have also been shown to affect some parasite life-
history traits, especially for *Anopheles* mosquitoes infected with the unicellular, eukaryotic
microparasites of the genus *Plasmodium* [38]; these parasites are the etiological agent of malaria, a
disease that continues to extract significant costs in morbidity and mortality in humans [43]. In malaria-
infected mosquitoes, *Plasmodium* oocyst densities in the midguts and mosquito survival were dependent
on dextrose ('D-glucose') concentrations [44]. Taken together, these observations suggest that
differences in nutritional requirements between oviposited and non-oviposited mosquitoes could
influence parasite fitness.

In principle, a single *Plasmodium* parasite ingested with a blood meal can invade the midgut wall
of an adult, female *Anopheles* and differentiate into an oocyst. Within the oocyst, in a process termed
'schizogony', several rounds of asexual replication culminate in the generation of sporozoites. Once
mature, the oocysts rupture and sporozoites are released into the mosquito hemocoel, from where they
migrate to the salivary glands, ready for injection during the next blood meal. In general, transmission to
the next host (i.e., future reproduction) is dependent on three traits: 1) the vector surviving past the time
required for sporozoites to invade the salivary glands, 2) rates of sporozoite migration to the salivary

gland ('extrinsic incubation period', EIP herein) [45, 46], and 3), the number of sporozoites in the salivary glands [47]. Although a single oocyst can produce thousands of sporozoites, all three fitness traits (vector survival, rates of migration, and the overall densities) are dependent on trade-offs with other traits in the parasite and vector's life history strategy [45, 48, 49, 50, 51]. The current study tests the hypothesis that parasite fitness would be dependent not only on whether the mosquito has oviposited or not (oviposition status), but also the amount of nutrition available. Female *An. stephensi* mosquitoes were exposed to an infectious blood meal containing *P. berghei* [49, 50]. Vector survival, rates of sporozoite migration and densities were measured in mosquitoes that were (or not) allowed to oviposit, and whether this was affected by differences in nutritional availability (1% or 10% dextrose) [40].

MATERIALS AND METHODS

All reagents and consumables were purchased from Thermo-Fisher Scientific, unless stated otherwise.

Study design.

For a visual overview of the study design, refer to Additional file 1: Fig. S1. Based on procedures described in detail previously [49, 50], 1500 adult, female *An. stephensi* were infected with *P. berghei* infected mice (0 days post-infection (dpi herein)). At day 1 post-infection, ~350 mosquitoes were transferred to each of 4 cages for the respective treatments. Two cages were provided a low nutrition diet (1% dextrose), while the remaining two were offered the high nutrition diet (10% dextrose). At 6 dpi, 1 cage from each nutrient treatment were provided oviposition sites until 9 dpi [50]. The overall study lasted 27 days and employed a fully crossed design wherein the same starting population of *P. berghei*-infected mosquitoes were separated across four treatment groups: 1) oviposited mosquitoes with low nutrient availability (1% dextrose), 2) oviposited mosquitoes with high nutrient availability (10% dextrose), 3) non-oviposited mosquitoes with low nutrient availability and finally, 4) non-oviposited mosquitoes with high nutrient availability (Additional file 1: Fig. S1A).

Vector mortality, rates of *P. berghei* sporozoite migration (referred to herein as EIP) and sporozoite densities in the salivary glands were quantified essentially as described previously [49, 50]. While vector mortality and EIP are well known higher order components of the vectorial capacity model [45, 46], sporozoite densities in the salivary glands are increasingly recognized as a critical fitness-conferring trait in the next vertebrate host [47]. Mosquito mortality was monitored daily in all 4 cages from day 1 post-infection [50]. To determine EIP and overall sporozoite prevalence [50], salivary glands from 15-20 mosquitoes from each group were checked for presence or absence of sporozoites at three-

day intervals starting 9 dpi (i.e., days 9, 12, 15, 18, 21, 24 and 27 post-infection). To enumerate

sporozoite densities [49], salivary glands from 10-15 mosquitoes from each group were sampled from

days 15 to 27 post-infection, also at three-day intervals (i.e., 15, 18, 21, 24 and 27 dpi); note that due to

differences in assay procedures (see below), EIP [50] and sporozoite densities [49] were obtained from

different mosquitoes. Finally, to determine whether mosquitoes had oviposited or not, ovaries of each

sampled mosquito was assessed for presence (non-oviposited) or absence (oviposited) of eggs

(Additional file 1: Fig. S1B).

Mosquito husbandry

An. stephensi mosquitoes (Walter Reed Army Institute of Research ca 2015, "Indian" strain) were

maintained in a level 2 Arthropod containment laboratory at the University of Georgia, as described

previously [52]. Colonies were maintained at $27°C \pm 0.5°C$, $80\% \pm 5\%$ relative humidity, and under a

12-hour day/night photoperiod. Female mosquitos were provided with human whole blood (Interstate

Blood-Bank, Memphis, TN) with glass membrane feeders (Chemglass Life Sciences, NJ) to support egg

production. Hatched larvae were dispensed at a density of 300 L1s/1000 ml of water and maintained on

pelleted Hikari cichlid gold diet (HikariUSA, Hayward, CA). Adult colonies were maintained on sugar

water composed of 5% dextrose (w/v) and 0.05% para-aminobenzoic acid (PABA) (w/v).

P. berghei **infections of mice**

All animal procedures described herein were performed as per AUP number A2020 01-013-Y3-A10,

approved by UGA IACUC. Four- to six-week-old male, Hsd:ICR(CD-1) mice (Envigo, Indianapolis,

IN) were infected with *P. berghei* GFP-LUC$_{CON}$, a transgenic strain expressing GFP and Luciferase

reporter genes in an ANKA background (referred to herein as *P. berghei*), as described previously [49].

Briefly, 7 mice were infected with a dose of $5x10^6$ parasites/ml in a 500µl inoculum. At 4 dpi, mice

were anesthetized using 1.8% tri-bromo ethanol (Avertin, Sigma-Aldrich, MO) just prior to mosquito

feeds.

P. berghei **infections of mosquitoes**

At three-to-seven-days post-emergence, host-seeking, female mosquitoes were sorted into a cage (32

cm^3, Bug Dorm, Taiwan) and transferred to a 20°C environmental chamber (Percival Scientific, Perry,

IA). As described previously [49], anesthetized *P. berghei*-infected mice were placed atop the cage.

Mosquitoes were allowed to feed on the mice for 15 minutes (day 0 post-infection). At 24 hours post-

infection (or day 1), mosquitoes were distributed into 4 separate cages and maintained for the duration

of the experiment as described above ('Study design'). Mosquitoes were maintained on either 1% or

10% dextrose (w/v) supplemented with 0.05% para-Aminobenzoic acid (w/v).

Data collection

While daily rates of mortality were recorded for all 4 treatments, starting 9 dpi and up to 27 dpi, the

salivary glands of mosquitoes from each group were checked for the presence or absence of sporozoites

as described previously [50]. Briefly, mosquitoes were aspirated directly into 70% ethanol (v/v) before

salivary glands were dissected from the mosquito and transferred to a 3μl drop of PBS on a clean glass

slide. Salivary glands were ruptured to release sporozoites by placing a cover glass directly on the drop

of PBS; presence or absence of sporozoites was assessed on an upright DIC microscope (Leica

DM2500) at 100x or 400x magnification. Salivary glands were sampled from 15 to 17 mosquitoes in

each group at days 9, 12, 15, 18, 21, 24 and 27 post-infection.

Sporozoite densities in the salivary glands were enumerated as described previously [49].

Briefly, at days 15, 12, 18, 21, 24 and 27 post-infection, ten mosquitoes from each group were

terminally sampled by aspirating directly into 70% ethanol. Salivary glands from each mosquito were

resuspended in 50μl of PBS supplemented with 0.1% bovine serum in a 0.65ml tube. Sporozoites were

released by homogenizing the salivary glands with 10 strokes of a microtube pestle. Sporozoites were

quantified in a 10μl aliquot of the homogenate as described previously [49]; while this approach can

detect ≥125 sporozoites, the relatively laborious and time-consuming nature of this assay meant

quantification of sporozoite densities was not initiated until 15 dpi when appreciable (and thus more

reliable) quantities of sporozoites would be available for enumeration.

Data analyses

All data analyses were performed with generalized linear models (GLM) in JMP Pro software (ver 16).

In general, prevalence data were modeled with a binomial distribution, and count data were modeled

with a type 2 negative binomial distribution. For sporozoite prevalence and sporozoite densities, time

was standardized and treated as a continuous fixed effect. Oviposition status and nutrient levels (1% vs

10% dextrose) were treated as categorical fixed effects. The effect of time was also modeled as a

quadratic effect ('humped') to account for saturation and/or decline in the dependent variable trends.

Gravid status (presence or absence of eggs) was analyzed with the same fixed effects described above

using a binomial distribution. Mosquito survival was analyzed using the Cox Proportional Hazards
model with the interaction between oviposition status and nutrient levels as fixed effects.

The analyses also considered the possibility that not all mosquitoes in the respective groups may
respond to the two oviposition treatments, and as such, could potentially confound analyses and
comparisons between groups. For instance, while some mosquitoes may still retain eggs despite being
offered oviposition sites, in the groups not offered an oviposition site, some mosquitoes may nonetheless
oviposit; assessing the contribution of these confounders is also relevant to the vector survival data
where it was not possible to quantify oviposition status post-hoc [50]. To address this, mean sporozoite
prevalence or mean sporozoites densities from all four groups at each time point were quantified from
all mosquitoes irrespective of whether they responded to the treatment, Pearson's product-moment
correlations were performed with the mean sporozoite prevalence and densities from the same groups,
but after excluding non-responder mosquitoes from the respective groups, i.e., by excluding non-
oviposited (or gravid) mosquitoes from the groups offered oviposition site, or oviposited mosquitoes
(not gravid) from the groups not offered oviposition sites.

RESULTS

Data and treatment summary

For the duration of the study, a total of 616 *An. stephensi* mosquitoes were sampled from all four groups,
with 416 sampled for EIP and 200 for sporozoite densities. To determine the effectiveness of the
oviposition treatments, gravid status from each mosquito (presence or absence of eggs) was assessed at
the time of dissection. In the groups that *were offered* oviposition sites ('oviposited' herein), 89.1% of
mosquitoes successfully laid eggs (or 271 of 304 mosquitoes dissected), with 90.5% (133 of 147
mosquitoes) and 87.9% (138 of 157 mosquitoes) oviposited mosquitoes recovered from low (1%
dextrose) and high (10% dextrose) nutrition groups, respectively, over the duration of the experiment
(Additional file 2: Fig. S2). While similar number of oviposited mosquitoes were recovered at each time
point (Chi-squared test (χ^2) = 0.02, df = 1, P = 0.88), recovery was also not affected by nutritional
availability over time (χ^2 = 0.22, df = 1, P = 0.64) or nutritional availability in general (χ^2 = 0.49, df = 1,
P = 0.48) (Additional file 3: Table S1).

In the groups that were *not offered* an oviposition site ('non-oviposited' herein) (Additional file
2: Fig. S2), 78.5% of the mosquitoes still retained eggs in their ovaries (or 245 of 312 mosquito ovaries

assessed), with the 84.8% of mosquitoes retaining eggs in the groups offered low nutrition (134 of 158
mosquitoes) higher than the 72.1% in the high nutrition group (111 of 154 mosquitoes) ($\chi^2 = 6.86$, $df =$
1, $P = 0.008$) (Additional file 3: Table S1). However, this difference was due to the apparent decline at
day 24 post-infection (right pane, Additional file 2: Fig. S2A): excluding this sampling point from the
analysis suggested that while nutrient availability did not affect recovery of non-oviposited mosquitoes
($\chi^2 = 0$, $df = 1$, $P = 0.9686$), similar numbers of non-oviposited mosquitoes were recovered at the chosen
sampling times ($\chi^2 = 0.22$, $df = 1$, $P = 0.6322$). As the reason for this discrepancy was unknown, all
statistical analyses were performed with this sampling point included in the dataset.

Despite the presence of these mosquitoes in the respective groups (i.e., oviposited individuals in
groups not offered an oviposition cup, or non-oviposited individuals in groups offered oviposition cups),
correlation analysis suggested that in general, these individuals should not confound downstream data
analyses (Additional file 4: Fig. S3): compared to the estimations of mean sporozoite prevalence and
densities from all four groups over time irrespective of oviposition status, excluding these individuals
did not significantly alter the estimation of prevalence (Pearson's correlation coefficient, $r_{(26)} = 0.99$,
$P<0.001$) (Additional file 4: Fig. S3A), or sporozoite densities (Pearson's $r_{(18)} = 0.96$, $P<0.001$)
(Additional file 4: Fig. S3B). Thus, all analyses below were performed with datasets that consider all
individuals irrespective of whether they responded to the oviposition sites as expected; furthermore, this
approach also ensures compatibility with the vector survival data where it was not possible to
discriminate between mosquitoes that may or may not have responded as expected to the oviposition
sites.

Time to infectiousness (EIP) and overall sporozoite prevalence in salivary glands is dependent on
nutritional availability in mosquitoes that have oviposited, but not those that retain eggs.

Of the 616 mosquitoes, salivary glands from a total of 416 mosquitoes from all four groups were
assessed for the presence or absence of sporozoites at 9, 12, 15, 18, 21, 24 and 27 dpi (Fig. 1).
Differences in nutritional availability to oviposited and non-oviposited mosquitoes affected the rates at
which the mosquitoes became infectious with sporozoites in the salivary glands (EIP) ($\chi^2 = 5.45$, $df = 1$,
$P = 0.02$) (row 8, Table 1) (Fig. 1A), as well as the overall proportion of sporozoite-positive salivary
glands ($\chi^2 = 5.45$, $df = 1$, $P = 0.02$) (row 5, Table 1) (Fig. 1B). This effect is particularly apparent in
oviposited individuals that were maintained at 1% dextrose (left pane, Fig. 1A, and Fig. 1B). Indeed,
pairwise comparisons of the marginal means predicted by the model suggested a consistently lower

prevalence in this group compared to the other three groups after 15 dpi, with differences in estimated
probabilities predicted to range from -0.55 to -0.23 (rows 1-21, Additional file 5: Table S2). While
nutritional availability did not appear to result in clear differences between non-oviposited mosquitoes,
for the groups maintained at 10% dextrose, mosquitoes were infectious earlier in the oviposited group,
with higher prevalence at 15 dpi (Estimated difference in probability (*Est.* herein) = 0.34, *standard error*
(SE) = 0.1, *P* = 0.004) and to a lesser yet significant extent at 18 dpi (*Est.* = 0.27, *SE* = 0.1, *P* = 0.048)
(rows 38 and 39, Additional file 5: Table S2).

The dynamics of sporozoite migration and overall prevalence were generally higher in oviposited
mosquitoes maintained at 10% dextrose compared to the other groups, albeit not as clearly as the
oviposited groups maintained on 1% dextrose (Additional file 5: Table S2). In other words, the low
prevalence in the 1% dextrose group was compensated by the high prevalence in the 10% dextrose
group such that the mean prevalence in oviposited groups (mean = 36.8%) were comparable to the non-
oviposited groups (mean = 43.4%), which could explain why oviposition status alone did not clearly
influence prevalence (χ^2 = 3.14, *df* = 1, *P* = 0.076) (Table 1). In contrast, the clear effect of nutrition on
prevalence (χ^2 =12.37, *df* = 1, *P* = 0.008) (Table 1) was likely driven by the strong reduction in
prevalence in the oviposited group maintained on 1% dextrose, which, despite the higher prevalence in
the non-oviposited mosquitoes also maintained on 1% dextrose, resulted in mean prevalence of 32.2%
that was lower than the mean of the groups maintained at 10% dextrose (47.4%). Finally, the analyses
also suggested that when averaged over all four groups, the initial increase in proportion of sporozoite-
positive salivary glands recovered (χ^2 = 33.94, *df* = 1, *P*<0.001) was eventually followed by saturation
and/or decline (χ^2 = 31.95, *df* = 1, *P*<0.001) (Table 1) after 18 dpi (Fig. 1A). Taken together, these
results suggest complex consequences of interactions between nutrient availability and oviposition status
of mosquitoes on sporozoite migration rates.

Sporozoite densities in the salivary glands are dependent on nutritional availability in mosquitoes
that have oviposited, but not those that retain eggs.

Of the 613 mosquitoes sampled in total, 200 salivary glands were assessed for sporozoite densities at 15,
18, 21, 24 and 27 dpi (Fig. 2); 142 individuals showed evidence of carrying ≥125 sporozoites (see
"Methods"). In general, the trends were similar to sporozoite prevalence, with differences in nutritional
availability to oviposited and non-oviposited mosquitoes affected the dynamics of sporozoites densities
in infected salivary glands (χ^2 = 5.36, *df* = 1, *P* = 0.021) (row 8, Table 1) (Fig. 2A), as well as overall

sporozoite densities (χ^2 = 12.94, *df* = 1, *P*<0.001) (row 5, Table 1) (Fig. 2B). Pairwise comparisons of
the marginal means suggested lower sporozoite densities in oviposited groups maintained at 1%
dextrose compared to the other three groups (rows 1-15, Additional file 6: Table S3). In general, while
sporozoite densities increased more rapidly in oviposited mosquitoes maintained on 10% dextrose,
overall densities in the salivary glands were also higher compared to the other groups (Additional file 6:
Table S3).

Survival is reduced in mosquitoes that have oviposited, but also due to low nutrition.

Survival was dependent on oviposition status (χ^2 = 15.37, *df* = 1, *P*<0.001) and to a lesser extent on
nutritional concentrations (χ^2 = 4.94, *df* = 1, *P* = 0.026) (Figure 3) (Additional file 7: Table S4), with
higher risk of mortality to mosquitoes that had already oviposited (risk ratio (*rr*) = 1.42, *SE* = 0.13,
P<0.001) and maintained at 1% dextrose (*rr* = 1.23, *SE* = 0.11, *P* = 0.025). However, mosquito survival
rates were not affected by the combination of nutrient concentrations and oviposition status (χ^2 = 2.36,
df = 1, *P* = 0.124) (Additional file 7: Table S4).

DISCUSSION

Nutrient availability and oviposition status of mosquitoes resulted in complex consequences for
Plasmodium fitness traits. Differences in nutrient availability from the environment influenced the rates
at which oviposited and non-oviposited became infectious (EIP), sporozoite densities in salivary glands,
and vector survival. In general, however, nutritional availability had less of an impact on the three traits
in mosquitoes that were prevented from ovipositing. Taken together, these results suggest that costs to
parasite fitness may be 'buffered' by tradeoffs experienced by mosquitoes carrying eggs, even in a
resource-poor environment. However, for assays measuring transmission potential, where mosquitoes
are generally not allowed to oviposit, both oviposition status and nutritional availability should receive
critical consideration. For instance, at the very least, depending on oviposition status and/or nutritional
availability, transmission potential may either be underestimated or overestimated.

The positive effect of environmental nutrition on parasite fitness (i.e., probability of transmission
to the next host) was clearly evident in the mosquito groups that were allowed to oviposit: while higher
sugar concentrations enhanced rates of sporozoite migration (Fig. 1A, left pane) and densities (Fig. 2A,
left pane) in the salivary glands, higher vector survival (Fig. 3A, left pane), especially after 15 days,
indicate a larger proportion of transmission-capable mosquitoes. In general, higher nutrient levels

enhance sporozoite migration and densities by increasing the rates of replication of sporozoites within
the oocysts lining the basal lamina of the mosquito's midgut [53]. While low glucose concentrations
adversely affected the survival of adult mosquitoes in general [40, 41], *An. gambiae* infected with *P.*
falciparum showed increased attraction to and uptake of sugars, presumably in response to the higher
energetic demands [38, 54, 55]. Despite the potential for compensatory feeding behavior, the
concentrations used in the current study (1% glucose) were likely too low to prevent the apparent
decline in mosquito survival in the oviposited mosquitoes after 15-18 dpi [40]. Although mounting
immune defenses is energetically costly in insects, it could explain reduced sporozoite rates and
densities in this group, with associated pathology resulting in reduced vector survival [28]. Clearly,
determining whether these costs to parasite fitness are due to a general lack of resources or how the host
allocates resources or energetic budgets is critical [25, 32].

Nutrition did not have such a strong influence on parasite fitness in the mosquito groups that
were not allowed to oviposit (right panes in Fig. 1A, Fig. 2A and Fig. 3A). This effect was especially
apparent when comparing the non-oviposited individuals maintained on a low nutrient diet with their
oviposited counterparts. Indeed, despite the low nutrition, overall rates of sporozoite migration (Fig. 1B)
and densities (Fig. 2B) were comparable to the high nutrient groups, with no clear costs to vector
survival (Fig. 3B). While this indicates alternative sources of nutrition were able to support sporozoite
replication and vector survival, reduced immune defenses due to reproduction-immunity tradeoffs in
non-oviposited mosquitoes may also result in the within-host conditions being more conducive to
parasite and vector survival [28]. In the absence and/or reduced availability of environmental sources of
nutrition, energetic demands in adult stages of holometabolous insects can be met by the nutrients stored
in the fat bodies, but also by resorption of nutrients from reproductive tissues (also referred to as
'follicular resorption', 'oosorption' or 'follicular atresia'), albeit at the expense of reduced fecundity [32,
56, 57, 58, 59]. Since the current study was undertaken with mosquitoes originating from a common
pool (Additional file 1: Fig. S1), the potential contribution of the fat bodies should be similar between
the four treatment groups; in other words, all else being equal, the difference in rates of parasite
replication and vector survival between the non-oviposited and oviposited groups maintained under low
nutrient conditions should, in principle, reflect the extent of nutrient resorption from reproductive
tissues.

In mosquitoes, nutrient resorption follows apoptotic death of ovarian follicles (referred to as
follicular resorption herein) [57, 58, 60]. Follicular resorption is a trait that is shared across several

orders of insects and allows the organism to adjust to physiological demands [56]. While the trait incurs
costs in reduced fecundity for the current generation, evolutionary theory suggests these costs are traded
off with enhanced life span, which in turn allows the insect more time to locate oviposition sites that can
benefit the fitness of future generations [25, 27, 30, 31, 32, 34, 35, 58]. In the current study, survival was
indeed higher in the non-oviposited group and particularly evident when compared to the oviposited
group maintained under low nutrient conditions (Fig. 3). Increased mortality in oviposited mosquitoes
maintained under low nutrient treatment, especially towards the later stages of the infection may account
for the weak but nonetheless significant, negative effect of reduced dextrose on survival. In general,
nutrition and oviposition status showed a more pronounced influence on the parasite traits of EIP and
densities than on vector survival (Fig. 1A, Fig. 2A and Fig. 3A), which could be due to the reduced costs
to vector survival at the lower temperature (20°C) [61] necessitated by this parasite-vector interaction
[49]. Taken together, while the results presented here are in line with evolutionary theory for insects in
general, they also indicate that the benefits of nutrient resorption to future fitness may not only outweigh
the costs of reduced fecundity to the mosquitoes, while also enhancing their capacity to vector disease.

Follicular resorption has been suggested to result in reduced fecundity of *An. stephensi* following
infections with another rodent malaria species, *P. yoelii nigeriensis* [60], and *An. maculipennis* infected
with microsporidia [62]. To the best of the authors' knowledge, whether follicular resorption directly
benefits parasite replication has not been assessed empirically but is nevertheless plausible, and likely
warrants confirmation. For instance, while greater availability of neutral lipids during oogenesis
enhanced rates of sporozoite replication and migration to the salivary glands ([63] and reviewed by
[64]), similar lipid profiles were recovered following follicular resorption in *Ae. aegypti* [58].
Furthermore, in previous studies with the major arbovirus vector *Ae. aegypti*, higher rates of follicular
resorption were observed in mosquitoes provided lower concentrations of sugar, likely to compensate
for the reduced nutritional availability (at the cost of reduced fecundity) [57, 58]. In principle, higher
rates of resorption in non-oviposited *An. stephensi* maintained in a low nutrition environment (1%
dextrose) could explain the earlier appearance of *P. berghei* sporozoites in the salivary glands (yellow
line, right pane, Fig. 1A), while continued availability of higher nutrient concentrations (10% dextrose +
follicular resorption) in the other non-oviposited group could explain the higher proportion of infected
mosquitoes over time, as a result of sustained sporozoite replication and/or migration (blue line, right
pane, Fig. 1A) (and possibly enhanced survival, right pane, Fig. 3A) [49, 50]. Yet, when considered in
the context of their oviposited counterparts, where a linear effect of dextrose on parasite fitness was

evident in oviposited mosquitoes (shortened EIP and higher rates of infection, vector survival and
sporozoite densities), it is unclear why the same 10-fold higher concentration of dextrose (and additional
nutrients, albeit low, available from follicle resorption) in non-oviposited mosquitoes did not enhance
rates of sporozoite replication and/or migration to the salivary gland, or support higher sporozoite
densities. Taken together, these results suggest parasite fitness may be dependent on complex, dynamic
interactions between physiological (nutrition) and evolutionary (egg retention) trade-offs in the
mosquito vector.

CONCLUSIONS

In summary, as with other insects, preventing mosquitoes from ovipositing (and completing their
gonotrophic cycle) enhanced survival, even under nutrient limiting conditions [25, 32], albeit at the cost
of promoting fitness of parasite species, whose analogs in humans cause malaria [43]. While increasing
nutrition (% dextrose) positively impacted parasite fitness in mosquitoes that had laid eggs, the results
from non-oviposited mosquitoes were consistent with the rates of follicular resorption compensating for
differences in nutritional availability, and in the process enhancing vectorial capacity due to higher
vector survival and earlier sporozoite appearance in the salivary glands (shorter EIP) as well as
sporozoite densities. By obtaining three critical measures of transmission potential [45, 46, 47], this
study argues for careful consideration of egg retention and nutritional availability on measures of
vectorial capacity and vector competence.

From the parasite's perspective, the overall patterns indicate an intrinsic ability to adjust its growth rates
to the nutritional status of the host. Future studies manipulating the length of the gonotrophic cycle and
duration of nutritional availability should reveal the extent of parasite's adaptive plasticity. Additionally,
from the vector's perspective, whether parasite infection and low nutrition reduce fecundity and/or
hatchability would be valuable to determine its effect on vector abundance, although neither trait
appears to be as critical as selective oviposition behavior, which prioritizes egg retention [25, 27, 30, 31,
32, 34, 35, 58]. In general, however, as these results suggest, the consequences of the interactions
between physiological (nutrition) and evolutionary (egg retention) trade-offs in the vector are likely to
be complex for parasite fitness. As such, this complexity further underscores the importance of
quantifying the contribution of selective oviposition behavior to disease transmission [41, 65], especially

since most integrated vector control strategies [1, 14, 15, 16, 17, 18, 19] and human-induced climate

change either directly or indirectly target this key mosquito behavior [20, 21, 22, 23, 24].

REFERENCES

1. Day JF. Mosquito Oviposition Behavior and Vector Control. Insects. 2016;7 4; doi: 10.3390/insects7040065. https://www.ncbi.nlm.nih.gov/pubmed/27869724.

2. Dieter KL, Huestis DL, Lehmann T. The effects of oviposition-site deprivation on *Anopheles gambiae* reproduction. Parasit Vectors. 2012;5:235; doi: 10.1186/1756-3305-5-235. https://www.ncbi.nlm.nih.gov/pubmed/23072301.

3. Govoetchan R, Sovi A, Aïkpon R, Salako A, Agbo FO, Asidi A, et al. The impact of oviposition-site deprivation in gravid females of *Anopheles gambiae* (Diptera: Culicidae) on fecundity, trophic behaviour and life expectancy. International Journal of Tropical Insect Science. 2013;33 4:207-15; doi: 10.1017/S1742758413000349. https://doi.org/10.1017/S1742758413000349.

4. Artis ML, Huestis DL, Lehmann T. The effects of oviposition-site deprivation on longevity and bloodfeeding rate in *Anopheles gambiae*. Parasit Vectors. 2014;7:163; doi: 10.1186/1756-3305-7-163. https://www.ncbi.nlm.nih.gov/pubmed/24708656.

5. Johnson BJ, Fonseca DM. The effects of forced-egg retention on the blood-feeding behavior and reproductive potential of *Culex pipiens* (Diptera: Culicidae). J Insect Physiol. 2014;66:53-8; doi: 10.1016/j.jinsphys.2014.05.014. https://www.ncbi.nlm.nih.gov/pubmed/24862157.

6. Toure DS, Ouattara AF, Kra KD, Kwadjo KE, Kone M, Doumbia M, et al. [Impact of egg laying delay on reproduction, gorging habit and mortality in gravid females *Anopheles gambiae* (Diptera Culicidae)]. Bull Soc Pathol Exot. 2017;110 5:318-25; doi: 10.1007/s13149-017-0582-x. https://www.ncbi.nlm.nih.gov/pubmed/29299880.

7. Chisulumi PS, Nampelah B, Yohana R, Philbert A, Kweka EJ. Diet and Oviposition Deprivation Effects on Survivorship, Gonotrophic Dissociation, and Mortality of *Anopheles gambiae* s.s. J Parasitol Res. 2022;2022:6313773; doi: 10.1155/2022/6313773. https://www.ncbi.nlm.nih.gov/pubmed/35761826.

8.   Talyuli OAC, Bottino-Rojas V, Polycarpo CR, Oliveira PL, Paiva-Silva GO. Non-immune Traits
Triggered by Blood Intake Impact Vectorial Competence. Front Physiol. 2021;12:638033; doi:
10.3389/fphys.2021.638033. https://www.ncbi.nlm.nih.gov/pubmed/33737885.

9.   Smartt CT, Richards SL, Anderson SL, Vitek CJ. Effects of forced egg retention on the temporal
progression of West Nile virus infection in *Culex pipiens quinquefasciatus* (Diptera: Culicidae).
Environ Entomol. 2010;39 1:190-4; doi: 10.1603/EN09172.
https://www.ncbi.nlm.nih.gov/pubmed/20146856.

10. Blagborough AM, Delves MJ, Ramakrishnan C, Lal K, Butcher G, Sinden RE. Assessing
transmission blockade in *Plasmodium* spp. Methods Mol Biol. 2013;923:577-600; doi:
10.1007/978-1-62703-026-7_40. https://www.ncbi.nlm.nih.gov/pubmed/22990806.

11. Wu VY, Chen B, Christofferson R, Ebel G, Fagre AC, Gallichotte EN, et al. A minimum data
standard for vector competence experiments. Scientific Data. 2022;9 1; doi: 10.1038/s41597-022-
01741-4.

12. Chen B, Sweeny AR, Wu VY, Christofferson RC, Ebel G, Fagre AC, et al. Exploring the
Mosquito-Arbovirus Network: A Survey of Vector Competence Experiments. Am J Trop Med Hyg.
2023;108 5:987-94; doi: 10.4269/ajtmh.22-0511. https://www.ncbi.nlm.nih.gov/pubmed/37037424.

13. Bousema T, Dinglasan RR, Morlais I, Gouagna LC, van Warmerdam T, Awono-Ambene PH, et al.
Mosquito feeding assays to determine the infectiousness of naturally infected *Plasmodium*
falciparum gametocyte carriers. PLoS One. 2012;7 8:e42821; doi: 10.1371/journal.pone.0042821.
https://www.ncbi.nlm.nih.gov/pubmed/22936993.

14. Mafra-Neto A, Dekker T. Novel odor-based strategies for integrated management of vectors of
disease. Curr Opin Insect Sci. 2019;34:105-11; doi: 10.1016/j.cois.2019.05.007.
https://www.ncbi.nlm.nih.gov/pubmed/31247410.

15. Ferguson HM, Dornhaus A, Beeche A, Borgemeister C, Gottlieb M, Mulla MS, et al. Ecology: a
prerequisite for malaria elimination and eradication. PLoS Med. 2010;7 8:e1000303; doi:
10.1371/journal.pmed.1000303. https://www.ncbi.nlm.nih.gov/pubmed/20689800.

16. Mwingira V, Mboera LEG, Dicke M, Takken W. Exploiting the chemical ecology of mosquito
oviposition behavior in mosquito surveillance and control: a review. J Vector Ecol. 2020;45 2:155-
79; doi: 10.1111/jvec.12387. https://www.ncbi.nlm.nih.gov/pubmed/33207066.

17. Reiter P. Oviposition, dispersal, and survival in *Aedes aegypti*: implications for the efficacy of
control strategies. Vector Borne Zoonotic Dis. 2007;7 2:261-73; doi: 10.1089/vbz.2006.0630.
https://www.ncbi.nlm.nih.gov/pubmed/17627447.

18. Thongsripong P, Hyman JM, Kapan DD, Bennett SN. Human-Mosquito Contact: A Missing Link
in Our Understanding of Mosquito-Borne Disease Transmission Dynamics. Ann Entomol Soc Am.
2021;114 4:397-414; doi: 10.1093/aesa/saab011. https://www.ncbi.nlm.nih.gov/pubmed/34249219.

19. Kumari A, Kaushik N. Oviposition Deterrents in Herbivorous Insects and their potential use in
Integrated Pest Management. Indian Journal of Experimental Biology. 2016;54 3:163-74. <Go to
ISI>://WOS:000372562700001.

20. LaDeau SL, Allan BF, Leisnham PT, Levy MZ. The ecological foundations of transmission
potential and vector-borne disease in urban landscapes. Funct Ecol. 2015;29:889-901; doi:
10.1111/1365-2435.12487. https://www.ncbi.nlm.nih.gov/pubmed/26549921.

21. Brown L, Medlock J, Murray V. Impact of drought on vector-borne diseases--how does one
manage the risk? Public Health. 2014;128 1:29-37; doi: 10.1016/j.puhe.2013.09.006.
https://www.ncbi.nlm.nih.gov/pubmed/24342133.

22. Franklinos LHV, Jones KE, Redding DW, Abubakar I. The effect of global change on mosquito-
borne disease. Lancet Infect Dis. 2019;19 9:e302-e12; doi: 10.1016/S1473-3099(19)30161-6.
https://www.ncbi.nlm.nih.gov/pubmed/31227327.

23. Brown JJ, Pascual M, Wimberly MC, Johnson LR, Murdock CC. Humidity - The overlooked
variable in the thermal biology of mosquito-borne disease. Ecol Lett. 2023;26 7:1029-49; doi:
10.1111/ele.14228. https://www.ncbi.nlm.nih.gov/pubmed/37349261.

24. Russell MC, Herzog CM, Gajewski Z, Ramsay C, El Moustaid F, Evans MV, et al. Both
consumptive and non-consumptive effects of predators impact mosquito populations and have
implications for disease transmission. Elife. 2022;11; doi: 10.7554/eLife.71503.
https://www.ncbi.nlm.nih.gov/pubmed/35044908.

25. Boggs CL. Understanding insect life histories and senescence through a resource allocation lens.
Functional Ecology. 2009;23 1:27-37; doi: 10.1111/j.1365-2435.2009.01527.x.

26. Nestel D, Papadopoulos NT, Pascacio-Villafan C, Righini N, Altuzar-Molina AR, Aluja M.
Resource allocation and compensation during development in holometabolous insects. J Insect

Physiol. 2016;95:78-88; doi: 10.1016/j.jinsphys.2016.09.010.

https://www.ncbi.nlm.nih.gov/pubmed/27650504.

27. Resetarits WJ. Oviposition site choice and life history evolution. American Zoologist. 1996;36

2:205-15. <Go to ISI>://WOS:A1996UJ51800010.

28. Schwenke RA, Lazzaro BP, Wolfner MF. Reproduction-Immunity Trade-Offs in Insects. Annu Rev

Entomol. 2016;61:239-56; doi: 10.1146/annurev-ento-010715-023924.

https://www.ncbi.nlm.nih.gov/pubmed/26667271.

29. Zera AJ, Harshman LG. The Physiology of Life History Trade-Offs in Animals. Annu Rev Ecol

Syst. 2001;32 1:95-126; doi: 10.1146/annurev.ecolsys.32.081501.114006.

30. Refsnider JM, Janzen FJ. Putting Eggs in One Basket: Ecological and Evolutionary Hypotheses for

Variation in Oviposition-Site Choice. Annual Review of Ecology, Evolution, and Systematics.

2010;41 1:39-57; doi: 10.1146/annurev-ecolsys-102209-144712.

31. Jaumann S, Snell-Rood EC. Trade-offs between fecundity and choosiness in ovipositing butterflies.

Animal Behaviour. 2017;123:433-40; doi: 10.1016/j.anbehav.2016.11.011.

32. Jervis MA, Boggs CL, Ferns PN. Egg maturation strategy and survival trade-offs in holometabolous

insects: a comparative approach. Biological Journal of the Linnean Society. 2007;90 2:293-302;

doi: DOI 10.1111/j.1095-8312.2007.00721.x. <Go to ISI>://WOS:000243909900008.

33. Rafferty AR, Reina RD. Arrested embryonic development: a review of strategies to delay hatching

in egg-laying reptiles. Proc Biol Sci. 2012;279 1737:2299-308; doi: 10.1098/rspb.2012.0100.

https://www.ncbi.nlm.nih.gov/pubmed/22438503.

34. Rosenheim JA, Heimpel GE, Mangel M. Egg maturation, egg resorption and the costliness of

transient egg limitation in insects. Proc Biol Sci. 2000;267 1452:1565-73; doi:

10.1098/rspb.2000.1179. https://www.ncbi.nlm.nih.gov/pubmed/11007333.

35. Rosenheim JA, Jepsen SJ, Matthews CE, Smith DS, Rosenheim MR. Time limitation, egg

limitation, the cost of oviposition, and lifetime reproduction by an insect in nature. Am Nat.

2008;172 4:486-96; doi: 10.1086/591677. https://www.ncbi.nlm.nih.gov/pubmed/18793093.

36. Garland T, Jr., Downs CJ, Ives AR. Trade-Offs (and Constraints) in Organismal Biology. Physiol
Biochem Zool. 2022;95 1:82-112; doi: 10.1086/717897.
https://www.ncbi.nlm.nih.gov/pubmed/34905443.

37. Peach DAH, Gries G. Mosquito phytophagy – sources exploited, ecological function, and
evolutionary transition to haematophagy. Entomologia Experimentalis et Applicata. 2020;168
2:120-36; doi: 10.1111/eea.12852.

38. Carvajal-Lago L, Ruiz-Lopez MJ, Figuerola J, Martinez-de la Puente J. Implications of diet on
mosquito life history traits and pathogen transmission. Environ Res. 2021;195:110893; doi:
10.1016/j.envres.2021.110893. https://www.ncbi.nlm.nih.gov/pubmed/33607093.

39. Kumar G, Gupta SK, Pasi S. Attractive toxic sugar baits: a magic bullet for control of malaria and
dengue in urban settings of India? J Vector Borne Dis. 2023;60 3:340-1; doi: 10.4103/0972-
9062.374242. https://www.ncbi.nlm.nih.gov/pubmed/37843248.

40. Foster WA. Mosquito sugar feeding and reproductive energetics. Annu Rev Entomol. 1995;40:443-
74; doi: 10.1146/annurev.en.40.010195.002303. https://www.ncbi.nlm.nih.gov/pubmed/7810991.

41. Stone CM, Foster WA. Plant-sugar feeding and vectorial capacity. Ecology of parasite-vector
interactions; 2013. p. 35-79.

42. Barreaux P, Barreaux AMG, Sternberg ED, Suh E, Waite JL, Whitehead SA, et al. Priorities for
Broadening the Malaria Vector Control Tool Kit. Trends Parasitol. 2017;33 10:763-74; doi:
10.1016/j.pt.2017.06.003. https://www.ncbi.nlm.nih.gov/pubmed/28668377.

43. WHO: World Malaria Report 2022. Geneva: World Health Organization; 2022.

44. Lambrechts L, Chavatte JM, Snounou G, Koella JC. Environmental influence on the genetic basis
of mosquito resistance to malaria parasites. Proc Biol Sci. 2006;273 1593:1501-6; doi:
10.1098/rspb.2006.3483. https://www.ncbi.nlm.nih.gov/pubmed/16777744.

45. Ohm JR, Baldini F, Barreaux P, Lefevre T, Lynch PA, Suh E, et al. Rethinking the extrinsic
incubation period of malaria parasites. Parasit Vectors. 2018;11 1:178; doi: 10.1186/s13071-018-
2761-4. https://www.ncbi.nlm.nih.gov/pubmed/29530073.

46. Brady OJ, Godfray HC, Tatem AJ, Gething PW, Cohen JM, McKenzie FE, et al. Vectorial capacity
and vector control: reconsidering sensitivity to parameters for malaria elimination. Trans R Soc

Trop Med Hyg. 2016;110 2:107-17; doi: 10.1093/trstmh/trv113.

https://www.ncbi.nlm.nih.gov/pubmed/26822603.

47. Churcher TS, Sinden RE, Edwards NJ, Poulton ID, Rampling TW, Brock PM, et al. Probability of

Transmission of Malaria from Mosquito to Human Is Regulated by Mosquito Parasite Density in

Naive and Vaccinated Hosts. PLoS Pathog. 2017;13 1:e1006108; doi:

10.1371/journal.ppat.1006108. https://www.ncbi.nlm.nih.gov/pubmed/28081253.

48. Lefevre T, Ohm J, Dabire KR, Cohuet A, Choisy M, Thomas MB, et al. Transmission traits of

malaria parasites within the mosquito: Genetic variation, phenotypic plasticity, and consequences

for control. Evol Appl. 2018;11 4:456-69; doi: 10.1111/eva.12571.

https://www.ncbi.nlm.nih.gov/pubmed/29636799.

49. Pathak AK, Shiau JC, Franke-Fayard B, Shollenberger LM, Harn DA, Kyle DE, et al. Streamlining

sporozoite isolation from mosquitoes by leveraging the dynamics of migration to the salivary

glands. Malar J. 2022;21 1:264; doi: 10.1186/s12936-022-04270-y.

https://www.ncbi.nlm.nih.gov/pubmed/36100902.

50. Pathak AK, Shiau JC, Freitas RCS, Kyle DE. Blood meals from 'dead-end' vertebrate hosts

enhance transmission potential of malaria-infected mosquitoes. One Health. 2023;17; doi:

10.1016/j.onehlt.2023.100582.

51. Childs LM, Prosper OF. The impact of within-vector parasite development on the extrinsic

incubation period. R Soc Open Sci. 2020;7 10:192173; doi: 10.1098/rsos.192173.

https://www.ncbi.nlm.nih.gov/pubmed/33204441.

52. Pathak AK, Shiau JC, Thomas MB, Murdock CC. Cryogenically preserved RBCs support

gametocytogenesis of *Plasmodium falciparum* in vitro and gametogenesis in mosquitoes. Malar J.

2018;17 1:457; doi: 10.1186/s12936-018-2612-y. https://www.ncbi.nlm.nih.gov/pubmed/30522507.

53. Habtewold T, Sharma AA, Wyer CAS, Masters EKG, Windbichler N, Christophides GK.

Plasmodium oocysts respond with dormancy to crowding and nutritional stress. Sci Rep. 2021;11

1:3090; doi: 10.1038/s41598-021-81574-0. https://www.ncbi.nlm.nih.gov/pubmed/33542254.

54. Nyasembe VO, Teal PEA, Sawa P, Tumlinson JH, Borgemeister C, Torto B. *Plasmodium*

falciparum infection increases *Anopheles gambiae* attraction to nectar sources and sugar uptake.

Curr Biol. 2014;24 2:217-21; doi: 10.1016/j.cub.2013.12.022.

https://www.ncbi.nlm.nih.gov/pubmed/24412210.

55. Zhao YO, Kurscheid S, Zhang Y, Liu L, Zhang L, Loeliger K, et al. Enhanced survival of

Plasmodium-infected mosquitoes during starvation. PLoS One. 2012;7 7:e40556; doi:

10.1371/journal.pone.0040556. https://www.ncbi.nlm.nih.gov/pubmed/22808193.

56. Bell WJ, Bohm MK. Oosorption in insects. Biol Rev Camb Philos Soc. 1975;50 4:373-96; doi:

10.1111/j.1469-185x.1975.tb01058.x. https://www.ncbi.nlm.nih.gov/pubmed/1106789.

57. Clifton ME, Noriega FG. Nutrient limitation results in juvenile hormone-mediated resorption of

previtellogenic ovarian follicles in mosquitoes. J Insect Physiol. 2011;57 9:1274-81; doi:

10.1016/j.jinsphys.2011.06.002. https://www.ncbi.nlm.nih.gov/pubmed/21708165.

58. Clifton ME, Noriega FG. The fate of follicles after a blood meal is dependent on previtellogenic

nutrition and juvenile hormone in Aedes aegypti. J Insect Physiol. 2012;58 7:1007-19; doi:

10.1016/j.jinsphys.2012.05.005. https://www.ncbi.nlm.nih.gov/pubmed/22626792.

59. Arrese EL, Soulages JL. Insect fat body: energy, metabolism, and regulation. Annu Rev Entomol.

2010;55:207-25; doi: 10.1146/annurev-ento-112408-085356.

https://www.ncbi.nlm.nih.gov/pubmed/19725772.

60. Hurd H. Evolutionary drivers of parasite-induced changes in insect life-history traits from theory to

underlying mechanisms. Adv Parasitol. 2009;68:85-110; doi: 10.1016/S0065-308X(08)00604-0.

https://www.ncbi.nlm.nih.gov/pubmed/19289191.

61. Miazgowicz KL, Shocket MS, Ryan SJ, Villena OC, Hall RJ, Owen J, et al. Age influences the

thermal suitability of *Plasmodium falciparum* transmission in the Asian malaria vector *Anopheles*

stephensi. Proc Biol Sci. 2020;287 1931:20201093; doi: 10.1098/rspb.2020.1093.

https://www.ncbi.nlm.nih.gov/pubmed/32693720.

62. Detinova TS, Bertram DS, World Health O: Age-grouping methods in diptera of medical

importance, with special reference to some vectors of malaria / T. S. Detinova ; [with] an Annex on

the ovary and ovarioles of mosquitos (with glossary) by D. S. Bertram. Geneva: World Health

Organization; 1962.

63. Costa G, Gildenhard M, Eldering M, Lindquist RL, Hauser AE, Sauerwein R, et al. Non-

competitive resource exploitation within mosquito shapes within-host malaria infectivity and

606  virulence. Nat Commun. 2018;9 1:3474; doi: 10.1038/s41467-018-05893-z.

607  https://www.ncbi.nlm.nih.gov/pubmed/30150763.

608 64. O'Neal AJ, Butler LR, Rolandelli A, Gilk SD, Pedra JH. Lipid hijacking: a unifying theme in

609  vector-borne diseases. Elife. 2020;9; doi: 10.7554/eLife.61675.

610  https://www.ncbi.nlm.nih.gov/pubmed/33118933.

611 65. Le Menach A, McKenzie FE, Flahault A, Smith DL. The unexpected importance of mosquito

612  oviposition behaviour for malaria: non-productive larval habitats can be sources for malaria

613  transmission. Malar J. 2005;4:23; doi: 10.1186/1475-2875-4-23.

614  https://www.ncbi.nlm.nih.gov/pubmed/15892886.

615 LEGENDS TO MAIN FIGURES AND TABLE.

616 **Fig. 1. A)** Time from infection till mosquitoes became infectious ('extrinsic incubation period' or EIP)
617 based on appearance of sporozoites in the salivary glands of oviposited (left pane) and non-oviposited
618 (right pane) mosquitoes, maintained on a diet of low (1% dextrose, yellow) or high nutrition (10%
619 dextrose). **B)** Mean sporozoite prevalence aggregated over the time points in A). Refer to Table 1 for
620 statistical modeling of the data set, and Additional file 5: Table S2 for post hoc pairwise comparisons of
621 the four groups at each day post-infection.

622 **Fig. 2**. **A)** Dynamics of sporozoite densities in the salivary glands of oviposited (left pane) and non-
623 oviposited (right pane) mosquitoes (infected mosquitoes only), maintained on a diet of low (1%
624 dextrose, yellow) or high nutrition (10% dextrose). **B)** Mean sporozoite densities aggregated over the
625 time points in A. Refer to Table 1 for statistical modeling of the data set, and Additional file 6: Table S3
626 for post hoc pairwise comparisons of the four groups at each day post-infection. The minimal limit of
627 detection with this technique is 125 sporozoites. ND = not done (for rationale, refer to Study design
628 under "Methods").

629 **Fig. 3**. **A)** Survival rates of oviposited (left pane) and non-oviposited (right pane) mosquitoes (infected
630 mosquitoes only), maintained on a diet of low (1% dextrose, yellow) or high nutrition (10% dextrose).
631 **B)** Mean survival rates aggregated over the time points in A).

632 **Table 1.** Statistical modeling of sporozoite prevalence ('y = sporozoite prevalence') (Figure 1) and
633 sporozoite densities ('y = sporozoite densities') (Figure 2). Binomial and negative binomial generalized
634 linear models respectively, were used to model sporozoite prevalence and densities.

LEGENDS TO SUPPLEMENTARY FIGURES AND TABLES.

Additional file 1: Fig. S1. A) Study design. **B)** Representative images of ovaries from mosquitoes that had oviposited successfully (left pane) or still retained eggs (right pane).

Additional file 2: Fig. S2. A) The proportion of mosquitoes that oviposited after being provided an oviposition site (left pane, referred to as 'oviposited' elsewhere in this manuscript) or not provided a site (right pane, referred to as 'non-oviposited' elsewhere), based on the gravid status (presence or absence of eggs as in Additional file 1: Fig. S1B) of mosquitoes at the time of dissection. **B)** Mean rates of gravidity recorded at the time points in A). Note that the data in this figure is representative of all 616 mosquitoes sampled in the study (Fig. 1 and Fig. 2) as the same approach was used to assess gravid status, prior to preparing the salivary glands to assess prevalence (Fig. 1) or densities (Fig. 2). For statistical analysis of the data, refer to Additional file 3: Table S1.

Additional file 3: Table S1. Statistical models of the proportion of mosquitoes that had oviposited after being provided an oviposition site ('oviposited') or not provided a site ('non-oviposited'), based on the gravid status ('y = gravid status (ovaries ± eggs)') at the time of dissections for data collection, as depicted in Additional file 1: Fig. S1.

Additional file 4: Figure S3. Excluding individuals that did not oviposit when offered an oviposition site, or oviposited despite not being offered an oviposition sites, does not influence the mean rates of sporozoite prevalence (**A**) or densities (**B**). Each data point represents the mean of mosquitoes sampled from all four groups at each time point (for prevalence, 4 groups * 7 dpi = 28 data points, for densities, 4 groups * 5 dpi = 20 data points); x-axes in both figures depicting the mean of all individuals (regardless of oviposition status), regressed against the mean of individual mosquitoes that were confirmed to have oviposited when provided a site ('oviposited'), or gravid (carrying eggs) due to not being provided an oviposition site ('non-oviposited').

Additional file 5: Table S2. Pairwise comparisons (post-hoc) of the estimated marginal means in sporozoite prevalence predicted by the statistical model in Table 1 ('y = sporozoite prevalence') (Fig. 1A). Comparisons were made between each of the four groups (oviposited and nutrient treatments) for the indicated day post-infection. Estimates ('Est. (= A-B)') indicate the difference of the predicted mean in group B subtracted from that of group A. Tukey's method was used to adjust for multiple comparisons.

Additional file 6: Table S3. Pairwise comparisons (post-hoc) of the estimated marginal means in sporozoite densities predicted by the statistical model in Table 1 ('y = sporozoite densities') (Fig. 2A). Comparisons were made between each of the four groups (oviposited and nutrient treatments) for the indicated day post-infection. Estimates ('Est. (= A-B)') indicate the difference of the predicted mean in group B, subtracted from that of group A. Tukey's method was used to adjust for multiple comparisons.

Additional file 7: Table S4. Statistical model of survival rates (Fig. 3). Survival was measured daily in all four groups until 27 dpi, and data modeled using Cox proportional hazards.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

CONSENT FOR PUBLICATION

Not applicable

AVAILABILITY OF DATA AND MATERIALS

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

COMPETING INTERESTS

The authors declare that they have no competing interests.

FUNDING

Georgia Research Alliance. JCS was funded by an NIH T32 award (AI1060546). NGD was funded by the National Science Foundation REU Site "The Population Biology of Infectious Diseases" at the University of Georgia, grant number 1659683.

AUTHORS' CONTRIBUTIONS

JCS: Design of the work, acquisition, analysis, interpretation, drafted the work.

NGD: acquisition, analysis.

DEK: acquisition.

AKP: Conception, design of the work, acquisition, analysis, interpretation, drafted the work and substantially revised.

All authors have approved the submitted version.

ACKNOWLEDGEMENTS

The authors would like to thank Courtnie Vickery and Rafael Freitas for help with data collection, Anne Elliot and Jared Cooper for their help with mosquito husbandry, and the staff at UGA animal resources for mouse husbandry.

Graphical abstract

Figure 1

Figure 2

A.

B.

Figure 3

A.

B.

Table 1

Row	Terms (x)	y = Sporozoite prevalence			y = Sporozoite densities		
		χ^2	df	P	χ^2	df	P
1	Oviposition status [a]	3.14	1	0.076	0.69	1	0.404
2	Nutrient levels [b]	12.37	1	<0.001*	6.95	1	0.008*
3	Dpi [c] (linear effect)	33.94	1	<0.001*	4.29	1	0.038*
4	Dpi (humped/'quadratic' effect)	31.95	1	<0.001*	1.78	1	0.183
5	Oviposition status * nutrient levels	10.67	1	0.001*	12.94	1	<0.001*
6	Oviposition status * dpi	2.86	1	0.091	1.22	1	0.269
7	Nutrient levels * dpi	1.87	1	0.171	2.50	1	0.114
8	Oviposition status * nutrient levels * dpi	5.45	1	0.02*	5.36	1	0.021*
	Number of salivary glands	416			142 (200 sampled)		

[a] Oviposited or Non-oviposited; [b] 1 or 10% dextrose; [c] salivary glands sampled at 3-day intervals from 9-27 dpi for sporozoite rates, and 15-27 dpi for densities.

Abbreviations: χ^2 = Chi-squared test (type 2); df = degrees of freedom; dpi = days post-infection; P = p-value.

* Statistical significance set at $P \leq 0.05$.