A senescence-specific lncRNA controls metabolic rewiring of senescent cells

Elena Grossi[1,2#], Francesco P. Marchese[1,2], Jovanna González[1,2], Enrique Goñi[1,2], Alicia Amadoz[1,2], Nicolás

Herranz[3], Marta Montes[1,2] *, Maite Huarte[1,2] *

[1] Center for Applied Medical Research, University of Navarra, Pamplona, Spain

[2] Institute of Health Research of Navarra (IdiSNA), Pamplona, Spain

[3] Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain; Vall d'Hebron Institute of Research (VHIR),

Barcelona, Spain

[#] Current address: Department of Oncological Sciences, Icahn School of Medicine at Mount Sinai, New

York, NY, USA; Department of Dermatology, Icahn School of Medicine at Mount Sinai, New York, NY,

USA.

* Co-corresponding authors: mmontesr@unav.es; maitehuarte@unav.es

ABSTRACT

Despite the classical view of senescence as passive growth arrest, it is an active process with profound

implications for cellular homeostasis. Senescent cells remain metabolically active to be able to cope with

the energetic demand of the senescence program, although the precise mechanisms underlying this

metabolic reprogramming are just beginning to emerge. Here we have identified *sin-lncRNA*, a previously

uncharacterized lncRNA, highly specific of senescent cells, and transcriptionally induced by C/EBPβ, the

master regulator of the senescence-associated secretory phenotype (SASP). While being strongly

activated in senescence, *sin-lncRNA* knockdown reinforces the senescence program by affecting oxidative

phosphorylation and mitochondrial function. *sin-lncRNA* interacts with the TCA enzyme dihydrolipoamide

S-succinyltransferase (DLST) to facilitate its proper. s*in-lncRNA* depletion increases DLST nuclear

translocation, favoring a metabolic shift from oxidative phosphorylation to a glycolytic phenotype.

Moreover, while *sin-lncRNA* expression remains low in highly proliferative cancer cells, it is strongly

induced upon cisplatin-induced senescence. Knockdown of *sin-lncRNA* in ovarian cancer cells results in

deficient oxygen consumption and increased extracellular acidification, sensitizing the cells to cisplatin

treatment. Altogether, these results indicate that *sin-lncRNA* is specifically induced in cellular senescence

to maintain metabolic homeostasis. Our findings reveal a new regulatory mechanism in which a lncRNA

contributes to the adaptive metabolic changes in senescent cells, unveiling the existence of an RNA-

dependent metabolic network specific to senescent cells.

INTRODUCTION

Senescence is a permanent state of growth arrest that can be induced by different types of stress, such as shortening of telomeres due to extensive replication, DNA damage, oxidative stress, chemotherapy treatment or oncogene overexpression [1]. Oncogene-induced senescence (OIS) was firstly described by Serrano *et al.* when they observed that expression of the HRASV12 oncogene in primary fibroblast induced a permanent cell-cycle arrest [2]. Later evidence has shown that OIS is a potent tumor suppressive mechanism that can repress malignant transformation by inducing a coordinated activation of tumor suppressor pathways, such as p53 and p16 [3, 4]. Senescent cells present characteristic morphological and biochemical features such as enlarged size, halted proliferation, activation of senescence-associated (SA) β-galactosidase activity and increased expression of cell cycle inhibitors [1]. Furthermore, in several types of senescence the cells secrete a combination of interleukins, metalloproteases and growth factors, collectively known as the senescence-associated secretory phenotype (SASP) [5-7]. While the SASP reinforces senescence, its chronic activation can have protumorigenic effects and contribute to several aging related pathologies [8]. In fact, the persistence of therapy-induced senescence cancer cells in the tumor has been linked to resistance to chemotherapy [9, 10]. There is therefore a critical need to develop combinatorial approaches to not only stop proliferation, but also induce the elimination of the senescent tumor cells.

Despite their decreased proliferative potential, senescent cells are metabolically active in order to cope with the energetic demands of the senescence program [11]. Indeed, metabolic reprogramming is considered a hallmark of senescence [12]. In general, senescent cells accumulate dysfunctional mitochondria resulting in an increase of reactive oxygen species (ROS) [13-15]. However, different types of senescence can lead to different types of metabolic alterations. For instance, OIS cells display a significant increase in tricarboxylic acid (TCA) cycle intermediates, oxidative phosphorylation (OXPHOS) [16] and mitochondrial fatty acid oxidation [17], while fibroblasts undergoing replicative senescence show increased glycolysis and reduced OXPHOS [18-20]. The fact that different triggers of senescence result in broad metabolic differences indicate that the interplay between senescent cells and metabolism is highly dynamic and context

dependent. Yet, the mechanisms underlying this regulatory process remain largely unexplored. In this

context, long non-coding RNAs (lncRNAs) represent a relatively uncharted territory for investigation.

LncRNAs are transcripts recently defined as longer than 500 nucleotides with exquisite cell-type specific

expression patterns that lack protein-coding potential [21]. They are tightly regulated during development or

in response to signaling pathways [22]. Although it is still unclear how many of the thousands of annotated

lncRNAs have a significant biological effect, several have been found to be essential for the regulation of

key cellular processes, such as proliferation and differentiation, as well as a broad range of diseases

including cancer [23, 24]. Notably, our previous work and other laboratories' have reported that specific

lncRNAs are differentially expressed during senescence induction [25, 26], regulating multiple senescence

aspects, from transcriptional response to SASP production [27-30]. Intriguingly, recent studies have involved a

number of lncRNAs in the regulation of cellular metabolism [31-34]; and ncRNAs are emerging as regulators of

metabolic enzymes [35]. However, the contribution of this plastic class of molecules to the metabolic

reprogramming of senescent cells remains unknown.

Here we report the identification of the senescence-specific lncRNA, *sin-lncRNA*, which is induced in

senescence to prevent uncontrolled metabolic alterations by regulating the function of a key metabolic

enzyme. Together, our results provide evidence of an RNA-dependent metabolic network specific of the

senescent cellular state.

RESULTS

Oncogene-induced senescence leads to a broad gene expression response, encompassing alterations

in lncRNAs

Although the importance of lncRNAs in a variety of cellular processes is now fully recognized, only a

handful of lncRNAs have been described as directly implicated in cellular senescence. To identify

noncoding regulators of this process, we analyzed RNA-seq data from a cellular system that recapitulates

senescence induction *in vitro* thanks to the controlled activation of an oncogene [36]. IMR90 non-

immortalized lung fibroblasts were engineered to express an oncogenic HRAS variant (HRAS[G12V]) fused to

the estrogen receptor (ER:RAS), which is activated following 4-hydroxy-tamoxifen (4OHT) administration

(Figure 1A) [5, 37]. Multiple senescence markers, such as arrest of proliferation, increased expression of pro-

inflammatory factors and tumor suppressors along with detectable β-galactosidase staining, were

observed, ensuring a proper senescence progression (Suppl. Figure 1A-E). Furthermore, we monitored the

chromatin remodeling linked to OIS by analyzing the presence of senescence-associated heterochromatin

foci (SAHF) as well as an increase in H3K9me3 and γH2AX foci (Suppl. Figure 1F, G). By mining RNAseq

data previously generated in this cell system [36,] we detected more than 4720 genes differentially expressed

upon induction of senescence (log2FC>1; padj<0.05), including not only many coding genes (3456) but also

lncRNAs (951) (Figure 1B and Suppl. Figure 1H). The majority of lncRNAs (62%) were downregulated,

which might reflect cell cycle inhibition during senescence, whereas the remaining 38% of lncRNAs were

upregulated. We focused our attention on lncRNAs highly induced upon OIS, as we speculated, they might

play a direct role in the senescence process, and we verified their upregulation at different time points of

OIS progression (Figure 1C and Suppl. Figure 1I).

sin-lncRNA is a senescence-specific lncRNA transcriptionally induced by C/EBPβ

Since several lncRNAs increased their expression in OIS, we sought to investigate their possible

contribution to this cellular process. Interestingly, we observed a previously uncharacterized lncRNA

(RIPOR3-as1) as one of the most significantly and strongly upregulated genes, with an induction

comparable to that of key senescence genes CKN2A, CXCL1 and IL6 (Figure 1B and Suppl. Table 1). We

consequently renamed it senescence-induced lncRNA or sin-lncRNA. sin-lncRNA is an intronic lncRNA

located in chromosome 20 and transcribed in the antisense orientation of the protein-coding gene RIPOR3

(Figure 1D, Suppl. Figure 2A, B). This noncoding transcript lacks coding potential (Suppl. Figure 2C) and

presents two poly-adenylated isoforms annotated by GENCODE (v41); a longer one (638 nts), composed

by three exons, and a shorter one (539 nts) of four exons. However, only the shorter isoform is expressed

and highly upregulated upon 4OHT treatment in IMR90 fibroblasts, according to both sequencing data and

RT-qPCR performed with different primer sets (Figure 1D and Suppl. Figure 2A). This observation was

confirmed by absolute quantification by RT-qPCR, which showed that sin-lncRNA is lowly expressed in

proliferating conditions with ~9 copies per cell, while, upon senescence induction, the expression of its

short isoform increases up to ~86 copies (Suppl. Figure 2D).

Interestingly, *sin-lncRNA* levels increased only at later stages of senescence (Figure 1C and Suppl. Figure

2A), prompting us to investigate whether *sin-lncRNA* upregulation was dependent on the full activation of

a senescence program or it was rather related to the induction of the HRAS oncogene in OIS. To this aim,

we analyzed *sin-lncRNA* expression levels in different types of senescence, such as replicative senescence -

achieved by prolonged culture of untransformed IMR90 fibroblasts - and senescence induced by ionizing

radiation (γ-irradiation). In both cases *sin-lncRNA* expression was strongly induced in the senescent

context, correlating with known senescence markers, such as $p16^{INK4a}$ and *IL6* (Figure 1E and Suppl. Figure

2E) and independently of HRAS or the approach used.

Due to the strong and highly specific *sin-lncRNA* induction in senescence, we speculated that senescence-

specific factors might regulate the expression of this lncRNA. We took advantage of publicly available

ChIP-seq data generated in IMR90 cells to examine *sin-lncRNA* genomic region, finding that the

transcription factor C/EBPβ was bound to several positions in the proximity of *sin-lncRNA* TSS and gene

body (Figure 1D). Interestingly, C/EBPβ exerts an important role in the later stages of senescence

induction, mainly by activating genes related to inflammation and SASP [38]. After confirming the previously

reported C/EBPβ upregulation in OIS [6] (Suppl. Figure 2F), we depleted C/EBPβ in senescent cells and

analyzed *sin-lncRNA* expression (Figure 1F, G and Suppl. Figure 2G). Notably, the induction of *sin-lncRNA*

in senescence was prevented by C/EBPβ depletion, similarly to what occurred to *IL1β*, *IL6* and *IL8,* other

canonical targets of this transcription factor (Figure 1G). To further investigate whether or not *sin-lncRNA*

is a C/EBPβ transcriptional target, we performed ChIP-qPCR of this transcription factor in proliferative and

senescent conditions (Figure 1H). C/EBPβ was found enriched at *sin-lncRNA* regulatory regions and its

binding increased in senescent cells (Figure 1H), supporting the evidences that the senescence driver

C/EBPβ controls *sin-lncRNA* expression at later stages of senescence induction and suggesting that the

lncRNA might exert a role in this response.

sin-lncRNA contributes to the senescent phenotype

To explore the function of *sin-lncRNA*, we depleted it in senescent IMR90 ER:RAS fibroblasts (Figure 2A

and Suppl. Figure 3A) and analyzed several senescence markers at 5 days of induction in order to detect

intermediate states before the cells enter a fully senescent phenotype [39]. Interestingly, *sin-lncRNA* KD

further decreased proliferation of senescent cells (Figure 2B, C and Suppl. Figure 3B), increased the

percentage of β-galactosidase positive cells (Figure 2D) and activated the p53 signaling pathway (Figure

2E), indicating that *sin-lncRNA* knockdown reinforces the senescent phenotype. Notably, the opposite

phenotype was observed when *sin-lncRNA* was overexpressed in senescent cells, alleviating the senescent-

mediated growth arrest (Figure 2H-I and Suppl. Figure 3H-I).

To corroborate the broader significance of these results, we took advantage of TIG3 hTERT lung

fibroblasts engineered to express an inducible and constitutively active portion of the human BRAF

oncogene to drive OIS (TIG3 ΔBRAF:ER) [40]. In this cell system, we again observed reduced proliferation and

increased β-galactosidase staining upon *sin-lncRNA* downregulation by RNAi (Suppl. Figure 3C-E),

indicating that *sin-lncRNA* depletion leads to the same phenotype in different models of cellular

senescence and regardless of the type of oncogene expressed. We also generated TIG3 ΔBRAF:ER cell

lines stably depleted of *sin-lncRNA* by CRISPR/Cas9 technology. Despite not being able to isolate viable

cell clones with *sin-lncRNA* biallelic deletion, we found that the removal of just one allele was sufficient to

decrease *sin-lncRNA* expression, leading to a consistent strengthening of the senescence phenotype

(Figure 2F-G and Suppl. Figure 3F-G). Altogether, these results suggest a role for *sin-lncRNA* as a negative

regulator of senescence induction.

***sin-lncRNA* depletion is linked to metabolic alterations and increased oxidative stress in senescent**

cells

To investigate which aspect of the complex senescence pathway is influenced by *sin-lncRNA*, we analyzed

the global changes in gene expression by RNA-seq of *sin-lncRNA*-depleted ER:RAS fibroblasts upon

senescence induction. RNA-seq analysis identified 295 upregulated and 137 downregulated genes upon

sin-lncRNA depletion (adj. P-value < 0.01; $\log_2$FC > ±1) (Figure 3A and Suppl. Table 2). After validating

some of the differentially expressed transcripts by RT-qPCR (Suppl. Figure 4A), we employed Gene Set

Enrichment Analysis (GSEA) to determine enriched pathways associated with the observed changes in

gene expression. *sin-lncRNA* knockdown led to the significant upregulation of genes associated to

epithelial-to-mesenchymal transition (EMT), as well as pathways generally related to response to stress,

such as interferon gamma (IFN-γ) response and p53 pathway. Conversely, growth promoting pathways,

such as MYC and E2F targets, were reduced upon *sin-lncRNA* knockdown, in line with the observed

changes in cell proliferation (Figure 3B and Suppl. Figure 4B). Interestingly, among the most significantly

reduced were several metabolic pathways, such as cholesterol homeostasis and fatty acid metabolism, as

well as those related to oxidative phosphorylation (Figure 3B-C).

The gene expression alterations linked to *sin-lncRNA* depletion pointed to changes in the metabolic state

of the cells. Metabolic rearrangements are considered a hallmark of senescence and in particular,

alterations in oxidative phosphorylation (OXPHOS) are known to be linked to senescence [16, 41]. In order to

analyze metabolic alterations upon *sin-lncRNA* depletion, we measured the oxygen consumption rate

(OCR) coupled to ATP production during OXPHOS by Seahorse assay [42]. Compared to control senescent

cells, *sin-lncRNA* knockdown led to the reduction of several key parameters of mitochondrial function,

such as Maximal Respiration rate, ATP production, Spare Respiratory Capacity and Coupling Efficiency

(Figure 3D, Suppl. Figure 4C). Accordingly, we also observed an increment in extracellular acidification rate

(ECAR), a reflection of increased glycolysis (Figure 3E).

Since alterations in OXPHOS are also linked to increased oxidative stress, we measured intracellular levels

of reactive oxygen and nitrogen species (ROS/RNS) and observed a significant enrichment of oxidant

species in *sin-lncRNA*-depleted senescent ER:RAS cells (Figure 3F), suggesting that *sin-lncRNA* is involved

in the control of oxidative stress in senescent cells. Moreover, since an increase in oxidant species and

metabolic dysfunctions can reinforce senescence [43, 44] we hypothesized that *sin-lncRNA* depletion might

exacerbate senescence-related mitochondrial alteration and promote oxidative stress, thereby leading to

the observed senescence reinforcement. Indeed, blocking ROS accumulation by N-acetyl-L-cysteine

(NAC) treatment[45, 46] restored cell growth and reduced SA-β-galactosidase activation in senescent cells

depleted for *sin-lncRNA* (Figure 3G and Suppl. Figure 4D). Together these data support the notion that *sin-*

lncRNA contributes to control the metabolic program and oxidative stress response of senescent cells.

sin-lncRNA interacts with DLST and controls its subcellular localization

LncRNAs have been previously associated with defects in oxidative stress and metabolic pathways, in

particular in cancer, where they alter gene programs through different mechanisms [22, 47]. To elucidate how

sin-lncRNA regulates metabolic response of senescent cells, we first analyzed *sin-lncRNA* subcellular

localization by RNA-FISH and cellular fractionation in cells depleted or not of the lncRNA. We found *sin-*

lncRNA to be mainly enriched in the cytoplasm of senescent fibroblasts (Figure 4A and Suppl. Figure 4E).

Therefore, *sin-lncRNA* likely acts in the cytoplasm to control proliferative and metabolic pathways of

senescent cells. We then reasoned that *sin-lncRNA* function would be mediated through interacting

proteins. We then set out to identify its protein interactors by performing RNA pulldown by incubating *in*

vitro transcribed *sin-lncRNA* or an unrelated control RNA with senescent IMR90 ER:RAS extracts. We

isolated a band of ~50-55 kDa unique for *sin-lncRNA*. Mass spectrometry (MS) analysis revealed several

proteins enriched (≥2 unique peptides) specifically associated with sin-lncRNA but not to the control RNA

MS (Figure 4B and Suppl. Table 3). Notably, the most abundant proteins retrieved by MS analysis

(Dihydrolipoamide S-succinyltransferase (DLST) and DEAD-Box Helicase 28 (DDX28)) have a recognized

role in mitochondrial function[48], in line with our findings linking sin-lncRNA to senescent cell metabolism.

DLST, identified as the main *sin-lncRNA* interactor, is the E2 component of the α-ketoglutarate (αKG)

dehydrogenase complex (KGDH), which is essential for the entry of glutamine into the tricarboxylic acid

(TCA) cycle [49] (Figure 4C). We independently confirmed the interaction between the endogenous DLST

and *sin-lncRNA* by RNA Immunoprecipitation in senescent fibroblasts (Figure 4D and Suppl. Fig 5B),

suggesting that they function coordinately. We therefore sought to investigate whether DLST depletion

could reproduce the phenotypic alterations observed upon *sin-lncRNA* knockdown in senescent cells, as

previous reports demonstrated increased ROS production upon DLST knockdown [50]. Indeed, DLST

knockdown in senescent fibroblasts led to an increase in the percentage of β-galactosidase positive cells

and decreased cell growth (Figure 4E, F and Suppl. Figure 4F), thus phenocopying *sin-lncRNA* knockdown.

Moreover, DLST knockdown resulted in similar expression changes of genes affected by *sin-lncRNA*

downregulation (Suppl. Figure 4G), suggesting that DLST and the lncRNA control the same cellular

pathways. Consistently with the increased ECAR observed in *sin-lncRNA* Knockdown cells (Figure 3E), we

observed an increase in lactate secretion in both *sin-lncRNA* and DLST-depleted senescent cells compared

to control cells, suggesting a shift to a glycolytic phenotype (Figure 4G).

Altogether, these results show that *sin-lncRNA* is upregulated during senescence to maintain a correct

balance in metabolic substrate consumption by regulating DLST engagement.

Our data suggest that *sin-lncRNA*, which physically interacts with DLST, is functionally connected to the

enzyme. However, DLST RNA and protein levels were not affected upon *sin-lncRNA* knockdown and

remained stable upon senescence induction (Suppl. Figure 4G-I), indicating that neither *sin-lncRNA* nor

cellular senescence modulate DLST expression. We then shifted our attention to the subcellular

localization of DSLT, which is encoded by a nuclear gene, and typically shuttles between different

subcellular compartments. While its main activity takes place in the mitochondria, it is also localized in the

cytoplasm and the nucleus of the cell [50-52]. We therefore investigated whether DLST localization might be

regulated by the lncRNA. Interestingly, cellular fractionation of senescent cells upon *sin-lncRNA* depletion

showed that DLST levels were significantly decreased in the cytoplasm and increased in the nuclear

fraction of senescent cells in absence of the lncRNA (Figure 4H). These results suggest that *sin-lncRNA*

might have an impact on the metabolic enzyme DLST in senescent cells by modulating its subcellular

localization.

***sin-lncRNA* regulates OXPHOS in ovarian cancer cells**

Our results indicate that *sin-lncRNA* is a regulator of the metabolic state of the senescent cells. To gain

insight into the potential significance of *sin-lncRNA* function in a physio-pathological context, we

interrogated gene expression data of tumor samples available through The Cancer Genome Atlas (TCGA)

portal. While *sin-lncRNA* presented low expression across cancer types (Suppl. Figure 5A), its expression

negatively correlated with that of genes involved in active cell division, such as cyclins, as *CCNA2*, and

members of the origin of replication complex (ORC), such as *ORC1* and *MCM6* [53] (Suppl. Figure 5B).

Accordingly, terms related to cell cycle and mitosis were the most enriched Gene Ontology pathways

associated with these genes (Suppl. Figure 5C). Since all these cell division-related genes presented a

negative correlation with the senescence-specific *sin-lncRNA*, we speculated that a high expression of this

noncoding transcript would predict cells with arrested cell cycle (*e.g.* senescent/quiescent cells), in line with

the upregulation of *sin-lncRNA* upon OIS.

While tumor cells are highly proliferative, it has been shown that chemotherapy can induce cellular

senescence, which can be linked to the resistance to treatment [10]. To investigate the potential

significance of *sin-lncRNA* in response to cancer therapy, we analyzed the expression of the

lncRNA in cisplatin-treated ovarian cancer cells, given their ability to undergo senescence upon

treatment with platinum-based agents [54]. 10 days after cisplatin treatment, COV362 cells displayed

a fully senescent phenotype, as confirmed by the expression of senescence markers, such as β-

galactosidase and the induction of cell cycle inhibitors and SASP components (Figure 5A-B).

Interestingly, we observed a strong increase of *sin-lncRNA* expression in the senescent cells,

whereas it was undetectable in untreated cells (Figure 5C). Cisplatin treatment also upregulated

C/EBPβ expression (Figure 5D), suggesting that this transcription factor might be responsible for the

transcriptional activation of *sin-lncRNA* also in the cancer cells. As previously shown for senescent

fibroblasts, knockdown of *sin-lncRNA* in cisplatin-treated COV362 cells reduced the expression of

genes involved in OXPHOS pathway (Figure 5E), which was reflected in alterations in the

metabolic functions, as a decreased OCR and increased ECAR (Figure 5F, G). Moreover,

knockdown of either *sin-lncRNA* or DLST further decreased cell growth compared to control

senescent cells (Figure 5H) indicating that depletion of the senescence-specific lncRNA sensitizes

ovarian cancer senescent cells to cisplatin treatment.

Altogether, these data provide evidence that a lncRNA uniquely expressed in senescent cells

contributes to the configuration of their metabolic state, also when senescence is induced in cancer

cells, thus representing a valuable marker and potential target in aging and cancer.

DISCUSSION

Although senescence has been traditionally considered a static terminal condition, it is now recognized as a dynamic stepwise process, which involves a plethora of factors that coordinately ensure its correct progress. We have identified a lncRNA that is specifically expressed at later stages of senescence induction to maintain the metabolic balance of these cells. Such precise induction is linked to a timely transcriptional activation mediated by the SASP (senescence-associated secretory phenotype) master regulator C/EBPβ; a transcription factor known to promote the later inflammatory phase of the senescent program [38]. Remarkably, depletion of *sin-lncRNA* is linked to the activation of some inflammatory signals as well as extracellular matrix remodeling pathways, gene networks that are induced during the second phase of senescence induction. *sin-lncRNA* depletion reinforces the senescence response, altering many of the canonical senescence markers. This is likely a consequence of the augmented ROS production, which feeds forward the senescent phenotype.

Mitochondria are the main source of ROS in the cell [55]. Interestingly, one of the most deregulated pathways upon *sin-lncRNA* depletion is oxidative phosphorylation (OXPHOS), which reflects mitochondrial dysfunction and results in high levels of ROS production [56]. High levels of ROS are critical for onset and maintenance of cellular senescence, although it remains difficult to decipher whether they are the cause or consequence of the senescence process [57]. Nevertheless, several factors are involved in feedback mechanisms to prevent an uncontrolled ROS increase [58]. We have found that *sin-lncRNA* interacts with DLST, an enzyme of the TCA cycle, a process that is tightly interconnected to OXPHOS and a main source of ROS [59]. Interestingly, it has been previously shown that the KGDH complex is sensitive to ROS and involved in ROS production [60], and inhibition of this enzyme might be critical in the metabolic deficiency induced by oxidative stress [61]. Accordingly, DLST depletion in senescent cells not only mimics *sin-lncRNA* knockdown by exacerbating growth arrest and β-galactosidase staining but also impairs cell metabolism. In fact, both *sin-lncRNA* and DLST downregulation result in increased extracellular acidification rate and

lactate production, suggesting a shift in the energy production pathway towards glycolysis. This may

reflect a mechanism for escaping the restrictions caused by the increased oxidative stress.

Where this interaction takes place remain unknow. However, although DLST function in the TCA occurs in

the mitochondria, we cannot exclude that DLST interacts with *sin-lncRNA* in the cytoplasm regulating the

availability of DLST for the TCA cycle. Of note, we have identified as a potential *sin-lncRNA* interactor

DDX28, which plays a role in OXPHOS and is also known to shuttle from the nucleus to the cytoplasm and

mitochondria [48]. Therefore, we cannot exclude that *sin-lncRNA* might regulate other aspects of

metabolism at the interface between cytoplasm and mitochondria or even other subcellular

compartments. Interestingly, TCA enzymes have been found to translocate into the nucleus where they

have roles in epigenetic regulation [62-65]. In particular, the KGDH complex which generates succinyl-CoA,

has been demonstrated to contribute to histone succinylation affecting expression of cell cycle genes [52]

and altering metabolic states [66]. Whether *sin-lncRNA* knockdown affects histone modification to regulate

transcription programs remains to be elucidated.

Platinum-based therapies, currently used as the first line of treatment for patients with ovarian

cancer, are known to induce senescence in cancer cells [67]. Although therapy-induced senescence

(TIS) has been considered a desirable outcome of cancer therapy, the activation of the

senescence program has been shown to induce the rewiring of cellular metabolism promoting the

pro-tumorigenic and pro-metatastic potential of ovarian cancer cells [68]. Here, we show that *sin-*

lncRNA is specifically induced in ovarian cancer cells that undergo senescence and contributes to

sustain the metabolic demands to maintain senescence in cancer cells upon drug treatment.

Emerging studies link metabolic reprogramming to cancer drug resistance [69]. Indeed, cisplatin-

resistant cells exhibit increased fatty acids (FA) uptake, accompanied by decreased glucose

uptake and lipogenesis, indicating reprogramming from glucose to FA dependent anabolic and

energy metabolism [70]. Importantly, other metabolic processes affected by *sin-lncRNA* depletion in

senescent fibroblasts seen at the transcriptional level include fatty-acid metabolism. Depletion of

sin-lncRNA might sensitize ovarian tumor cells to cisplatin by metabolic reprogramming at different

levels.

Metabolic enzymes required for the tricarboxylic acid cycle have been shown to affect senescence by

regulating p53[71]. Importantly, advances in the field of RNA interactome capture have demonstrated that

the interplay between metabolic enzymes and RNA is broader than previously considered[35,72]. These

studies have revealed the RNA binding capacity of many enzymes, although the precise mechanism or the

physiological relevance requires deeper characterization. Emerging evidence indicates that lncRNAs may

contribute to cellular processes in order to adapt and coordinate mitochondrial function to adjust their

activities to the environmental conditions[34,73,74]. We propose that distinct cellular states or stresses that

result in metabolic changes involve the function of ncRNAs that contribute to the re-configuration of the

cell metabolism. The exquisite specificity of *sin-lncRNA* in senescence highlights its potential role as a

therapeutic target. This discovery paves the way for better comprehension of therapy-induced

senescence and its associated implications.

MATERIALS AND METHODS

Cell culture, retroviral infection and treatments

IMR90 lung fibroblasts and HEK-293T cells were purchased from ATCC. TIG3 hTERT pMSCV-ER:ΔBRAF

lung fibroblasts[40] were kindly provided by Dr. Lund's laboratory (BRIC - Copenhagen, Denmark). All cell

lines were cultured in DMEM medium (GIBCO), supplemented with 10% fetal bovine serum (GIBCO) and 1x

penicillin/streptomycin (Lonza). Cells were maintained at 37°C in the presence of 5% CO_2 and tested for

mycoplasma contamination regularly, using the MycoAlert Mycoplasma Detection Kit (Lonza).

To generate OIS cell systems, viruses were first produced in HEK-293T cells transfected with 12µg of either

pLNC-ER:RAS or pLNCX (Empty) vectors, 6µg of gag-pol plasmid, 3µg of pVSVG vector and 1µg of pMAX-

GFP plasmid (to check transfection efficiency). Transfection reaction was carried out in opti-MEM medium

(GIBCO) using Lipofectamine 2000 (Invitrogen), following manufacturer's instructions. 48 hours after

transfection, filtered supernatant supplemented with 4µg/ml polybrene (Santa Cruz), was used to

transduce low passage IMR90. Cells were selected with Neomycin-G418 (Sigma) at a final concentration of

400µg/ml for at least one week.

For senescence induction studies, IMR90 ER:RAS or TIG3-hTERT ΔER:BRAF fibroblasts have been treated

with 200nM of 4-hydroxy-tamoxifen (4OHT) for three days unless specified otherwise in the text. When

longer time courses were needed, medium was replaced on day 3. Only normal fibroblasts cell lines with

less than 20-25 passages were used for experiments.

COV362 cells were obtained from SIGMA, kindly provided by Dr. Beatriz Tavira (CIMA) and cultured in

DMEM medium (GIBCO), supplemented with 10% fetal bovine serum (GIBCO) and 1x

penicillin/streptomycin (Lonza). For senescence induction the cells were treated with 10 μM cisplatin for 10

days.

N-Acetyl-L-cysteine (NAC, SIGMA) was added at 5 mM 1h prior 4-OHT treatment and for the duration of

the experiment.

Irradiation

To induce senescence by ionizing radiation, IMR90 cells were γ-irradiated at 5Gy and harvested at the

indicated times. Radiation was delivered at 180 cGy/min using a Siemens Oncor Impression Plus linear

accelerator equipped with 6MeV X rays.

RNAi and overexpression studies

For siRNA studies, cells were transfected once with a final concentration of 40nM siRNA using

Lipofectamine 2000 (Invitrogen) in opti-MEM medium (GIBCO), unless differently stated in the text.

Scrambled siRNA (siRNA CTRL in the text) was used as transfection control. All siRNAs employed

in this study were designed using BLOCK-iTTM RNAi Designer

(https://rnaidesigner.thermofisher.com/rnaiexpress/) and purchased from Sigma, except C/EBPβ

siRNA, a kind gift of Dr. Aragón (CIMA - Pamplona, Spain).

To induce senescence in transfected cells, transfection medium was replaced after 6 hours with complete

medium containing 200nM 4OHT. When a senescence induction longer than 3 days was needed,

transfected cells were trypsinized and re-plated to perform multiple analyses in parallel. All siRNAs used in

this study are listed in Table 4.

To achieve *sin-lncRNA* overexpression, its shorter isoform has been amplified from IMR90 cDNA with

specific primers and cloned in pcDNA3.1(+) vector (purchased from Addgene). For overexpression studies,

IMR90 ER:RAS cells has been transiently transfected with 500ng of *sin-lncRNA* or empty pcDNA3.1 plasmid

using Lipofectamine 2000 (Invitrogen) in opti-MEM medium (GIBCO). Primers employed for cloning are

described in Table 4.

CRISPR-Cas9 gene editing and CRISPR clonal cell lines generation

The CRISPR Design Tool from the Zhang Lab (http://crispr.mit.edu/) was used to find suitable target sites

for the Streptococcus pyogenes Cas9 (SpCas9) in *sin-lncRNA* genomic region. Specific guide RNAs

(sgRNA_up and sgRNA_dw, see Table 1) were cloned into the lentiviral plasmid lentiCRISPRv2 carrying

spCas9 gene and puromycin resistance, a kind gift from Dr. Agami's lab (NKI - Amsterdam, Netherlands)

and originally generated in Dr. Zhang's lab [282] (for details, see plasmid reference #5296113 on Addgene

website).

LentiCRISPR vectors carrying the two independent sgRNAs were separately transfected in HEK-293T cells

to generate viral particles, as described above. LentiCRISPR viruses were mixed (ratio 1:1) and used to

transduce TIG3-hTERT ΔER:BRAF fibroblasts, then selected with 1μg/ml of puromycin. Single puromycin-

resistant TIG3 cells were sorted in 96-well plates, using untreated TIG3 fibroblasts as feeder cells, which

were then eliminated by a second round of puromycin selection (0.5μg/ml). Genomic DNA was isolated

from puromycin-resistant clones using QuickExtract DNA extraction solution (Lucigen), PCR amplified and

tested on agarose gel for accurate deletion.

Cell proliferation assays

For crystal violet assay upon OIS, 1-2×10^4 untreated or transfected cells were plated in 6-well plates and

senescence induction was prolonged for 5 days in 4-OHT medium. Cells were fixed with 0.5%

glutaraldehyde for 15 min and stained with 0.1% Crystal Violet solution (Sigma) for 30 min. To quantify cell

density, cells were incubated with 500μl of 10% acetic acid (Sigma) and collected in ELISA plates.

Absorbance was measured by spectrophotometry at 570 nm in a SPECTROstar Nano equipment.

For BrdU incorporation assay, cells were labeled with 50 μM BrdU solution (BD Pharmingen) for 16-18

hours. In case of senescent cells, OIS was previously induced for 5 days before starting BrdU assay. Then,

cells were harvested by trypsinization and BrdU staining was performed using BrdU Flow Kits (BD

Pharmingen), according to manufacturer's instructions. Amount of BrdU incorporation was measured by

flow cytometry using a FACSCalibur Cell Analyzer (BD Bioscience).

Senescence Associated galactosidase (SA-β-gal) staining

To detect SA-β-gal activity, 1×10^3 of untreated or transfected cells were plated in 6-well plates and

senescence was induced for 8 days upon 4-OHT treatment. Cells were fixed with 0.5% glutaraldehyde

solution for 15 min at RT and washed with PBS containing 1mM MgCl2 (pH 6.0). Then, cells were incubated

overnight at 37°C in X-gal staining solution (1mg/ml X-gal [Thermo Fisher], 0.12 mM K3Fe(CN)6 [Merk],

0.12 mM K4Fe(CN)6 [Merk], PBS/MgCl2 [pH 6.0]). Cells were washed and percentage of blue cells was

counted manually using an inverted microscope.

ROS/RNS measurement assay

Senescent IMR90 ER:RAS fibroblasts transfected with *sin-lncRNA* targeting or control siRNAs were

collected by trypsinization and re-suspended at $0.5-1 \times 10^6$ cells/ml in Hypotonic Buffer (10mM Tris HCl [pH

7.5], 10mM NaCl, 1.5mM MgCl2). To facilitate membrane lysis, cells were incubated for 10 min at 4°C and

sonicated for 5 cycles (30''ON-30''OFF) in a Bioruptor sonication device. Cell extract was cleared by

centrifugation and assayed for ROS/RNS accumulation by fluorescence-based approach using OxiSelect in

vitro ROS/RNS Assay Kit (Cell Biolabs), following manufacturer's instructions. The amount of fluorescent

oxidized probe (2', 7' -dichlorodihydrofluorescein or DCF) was measured with a fluorescence plate reader

at 480nm excitation / 530nM emission. Data were interpolated to a DCF standard curve to obtain absolute

values.

Measurement of oxygen consumption and extracellular acidification

Cellular OCR was measured using a Seahorse XF Cell Mito Stress Test and a Seahorse Bioanalyzer XFe96

(Agilent Technologies) according to the manufacturer's standard protocol. Briefly, 200,000 IMR90 cells

were reverse transfected with the corresponding siRNAs in 6 well plates. After 24h cells were treated with

500 nM 4-OHT. After 3 days of treatment, cells were trypsinized and seeded on Seahorse XF96 Cell Culture

microplates to confluence (10000 cells/well) and were subjected to metabolic profiling the next day. For

Mito stress tests, cells were cultured for 1 h in a CO_2-free incubator at 37 °C. Oxygen consumption rates

(OCR) were monitored at basal conditions and, after sequential injections of 1 μM oligomycin to block the

mitochondrial ATP synthase, 1.5 μM FCCP to uncouple oxidative phosphorylation, 1 μM antimycin A and

1 μM rotenone were used to fully inhibit mitochondrial respiration and 50 mM 2-deoxygluocse (2-dG) to

block glycolysis. Two separate measurements of OCR and extracellular acidification rate (ECAR) from 4

independent experiments were taken after the addition of each inhibitor. Samples were normalized based

on the protein concentration. All reagents were purchased from Agilent Technologies.

Lactate measurements

IMR90 ER:RAS fibroblasts were transfected with *sin-lncRNA* targeting or control siRNAs and treated with

4-OHT for 6 days. The supernatant was harvested and centrifuged for 5 min at 500 g. Lactate levels were

measured using Cobas C311 Analyzer with LACT2 kits (Roche Diagnostics GmbH). Human Precinorm y

Precipath were used as controls.

Liquid Chromatographic -Mass Spectrometry (LC-MS) metabolomics

Isotope labelling

6-days senescent IMR90 ER:RAS fibroblasts transfected with siRNA control or siRNA against *sin-lncRNA* or

DLST were incubated with glutamine-free DMEM supplemented with 4 mM L-$^{13}C_5$-Glutamine (Cambridge

Isotopes Laboratories) . After 6 hours the cells were washed with PBS 1x and snap-freeze in liquid nitrogen.

Sample preparation and LC/MS methods

LCMS was performed at the Metabolomics Platform at CIC bioGUNE (Donostia, Spain). Cells from three

wells (1E6 cells/well) were pooled to obtain one LCMS sample. Therefore, cells were extracted with 200 μL

icecold extraction liquid per well. The volume from one well was transferred to the next and finally the

resulting 600 μL volume was passed over the three wells two times. The extraction liquid consisted of a

mixture of ice-cold methanol/water (50/50 %v/v). Subsequently 400 μL of the cell homogenate plus 400μL

of chloroform was transferred to a new aliquot and shaken at 1400 rpm for 60 minutes at 4 °C. Next the

aliquots were centrifuged for 30 minutes at 14000 rpm at 4 °C. 250μL of the aqueous phase was

transferred to a fresh aliquot. The chilled supernatants were evaporated with a speedvac in approximately

2h. The resulting pellets were resuspended in 150 μL water/MeOH (75/25 %v/v). Samples were measured

with a UPLC system (Acquity, Waters Inc., Manchester, UK) coupled to a Time-of-Flight mass

spectrometer (ToF MS, SYNAPT G2S, Waters Inc.). A 2.1 x 100 mm, 1.7 μm Phenyl-Hexyl column (Waters

Inc.), thermostated at 40 °C, was used to separate the analytes before entering the MS. Mobile phase

solvent A (aqueous phase) consisted of 99.5% water and 0.5% FA while solvent B (organic phase) consisted

of 99.5% MeCN and 0.5% FA. In order to obtain a good separation of the analytes the following gradient

was used: from 98% A to 0% A in 2 minutes in curved gradient (#8, as defined by Waters), constant at 0%

A for 1 minute, back to 98% A in 0.2 minutes. The flow rate was 0.250 mL/min and the injection volume

was 3 μL. After every 6 injections, a sample was injected. The MS was operated in negative electrospray

ionization in full scan mode. The cone voltage was 10 V, cone offset was 50V and capillary voltage was 400

V. Source temperature was set to 120 °C and capillary temperature to 500 °C. The flow of the cone,

desolvation and nebulizer gas (nitrogen) were set to 5 L/h, 6 L/h and 850 L/h, respectively. A 2 ng/mL

leucine-enkephalin solution in water/acetonitrile/formic acid (49.9/50/0.1 %v/v/v) was infused at 10 μL/min

and used for a lock mass which was measured each 36 seconds for 0.5 seconds. Spectral peaks were

automatically corrected for deviations in the lock mass. Extracted ion traces for relevant analytes were

obtained in a 10 mDa window in their expected m/z-channels. These traces were subsequently smoothed

and peak areas integrated with TargetLynx software. Signals of labeled analytes were corrected for

naturally occurring isotopes. Since we also had unlabeled experiments, for some metabolites we could

determine the naturally occurring isotope-distribution empirically. This was necessary for the first and

second isotopomers since correcting with theoretical values did not eliminate these isotopes from

unlabeled experiments. This is probably because signals are non-linear with respect to ion-counts, *i.e.*

higher ion-counts tend to have *relatively* lower signals than lower ion-counts due to saturation of the

detector. The calculated raw signals were adjusted by median fold-change (MFC) adjustment. This is a

robust adjustment factor for global variations in signal due to e.g. difference in tissue amounts, signal drift

or evaporation. The MFC is based on the total amount of detected mass spectrometric features (unique

retention time/mass pairs). The calculations and performance of the MFC adjustment factors are described

in the following publications [75, 76].

Immunofluorescence and RNA fluorescence in situ hybridization (RNA-FISH)

For fluorescence-based studies, IMR90 ER:RAS cells were collected by trypsinization after required

treatments (as specified in the text) and re-plated on top of cover glasses placed on clean 6-well plates.

The following day, cells were washed and fixed for 15 min at room temperature (RT) using 3% methanol-

free formaldehyde solution (Thermo Fisher).

For immunofluorescence studies, fixed cells were washed (PBS/0.5% NP-40) and incubated first with block

solution (10% FBS in wash buffer) for 20 min and then with primary antibody diluted in block solution for 1

hour at RT. Antibody solution was washed out and cells were incubated with fluorophore-conjugated

secondary antibody for 30 min. After extensive washes, cover glasses were mounted on microscope slides

using DAPI-containing mounting solution (Palex Medicals) and pictures were collected with an automated

optical microscope running Zen 2 core imaging software (Zeiss), according to standard procedures.

For RNA-ISH studies, fluorescein-labeled Locked Nucleic Acid (LNA) DNA probes were designed and

synthesized by Exiqon and were hybridized according to manufacturer's protocol with some modifications.

Briefly, fixed cells were first incubated with 70% ethanol for 1 hour and then with Acetylation Buffer (0.1M

Triethanol Amine, 0.5% (v/v) Acetic Anhydride) for 30 min at RT. To avoid non-specific probe binding, cells

were incubated with Hybridization buffer (10% dextran sulfate, 50% formamide, 2x saline-sodium citrate

[SSC] buffer) for 1 hour at 55°C. Specific LNA probes were denatured at 92°C for 4 min and then mixed

with Hybridization buffer at a final concentration of 20nM. Hybridization was carried out overnight at

55°C. The following day, probes residues were eliminated using 2x SSC wash buffer and fixed cells were

treated with 3% Hydrogen Peroxide for 30 min at RT. For fluorescein (FAM) detection, cells were first

incubated 1 hour with Blocking Buffer (10% heat-inactivated Goat Serum, 0.5% Blocking Reagent [Roche]

in PBS-0.5% Tween) and then 1 hour with 1.5U/ml αFAM-POD antibody (Roche) diluted in Blocking Buffer.

After extensive washes, fluorescent signal was developed by 10 min incubation with TSA-Cy3 solution

(Perkin Elmer). Residues were eliminated through stringent 4x SSC washing and slides were prepared for

microscope detection using DAPI mounting solution, as mentioned above.

Slides were imaged on a fluorescence inverted microscope using a 63x objective. At least 10 optical stacks

were acquired per image for DAPI (nuclear stain) and for the fluorescence channel corresponding to the

RNA FISH probe (Cy3). Image processing was performed using Zen software (version 2.3 - Zeiss) and for

each condition, signal was adjusted on a sample incubated with no probes and used as control for

background detection. ImageJ software was employed for stacks deconvolution.

Protein extraction and immunoblot analysis

Cells were lysed for 15 min in rotation at 4 □C using RIPA buffer (150mM NaCl, 25mM Tris HCl

[pH 7.5], 2mM EDTA, 0.1% sodium deoxycholate [Na-DOC], 0.1% SDS, 1% Triton X-100)

supplemented with 1x cOmplete Protease Inhibitor Cocktail [Roche]. For detection of

phosphoproteins, 1x PhosSTOP phosphatase inhibitor [Roche] was freshly added to RIPA

solution. Lysed cells were centrifuged at max speed for 10 min at 4 □C and the insoluble pellet

was discarded. Protein concentration was estimated by Pierce BCA Protein Assay Kit, using BSA

curve as reference and according to manufacturer's instructions. Proteins were separated on

denaturing SDS-PAGE gels and transferred to a nitrocellulose membrane [Biorad] following

standard procedures. Membranes were blocked using skim milk or BSA (VWR) and probed first for

primary and then for HRP-conjugated secondary antibody. Western Lightening ECL- Plus (Perkin

Elmer) was employed for chemiluminescence detection of proteins. The list of antibodies used in

this study can be found in Table 4.

Cellular Fractionation

IMR90 cells were reverse transfected with control or siRNA 1/3 *sin-lncRNA*. The day after transfection cells

were treated with 500 nM 4-OHT for 5 days. Cells were scrapped in cold PBS and centrifuge 5 min 0.2g.

Pellets were resuspended in NIB buffer (1.28 M Sucrose, 40 mM Tris pH 7.4, 20 mM MgCl2, 4% Triton X-

100), incubated on ice for 20 min and centrifuged 15 min at 2500rpm. The supernatant was kept as the

cytosolic fraction and further cleaned by an extra centrifugation step. Nuclear pellets were washed twice

with NWB (40 mM Tris pH 7.4, 20 mM $MgCl_2$) by centrifugation 15 min at 2500 rpm. RIPA (150 mM NaCL,

25 mM Tris pH 8, 0.1% SDS, 0.1 % Na-Deoxycholate, 2 mM EDTA, 1% Triton X-100) buffer was used to lyse

the nuclei. After 10 min centrifugation at max speed the supernatant was isolated as the nuclear fraction.

Immunoblot analysis was performed as described above.

Native RNA immunoprecipitation

20×10^6 cells were collected in cold PBS by scraping and lysed in RIP buffer (25 mM Tris-HCl pH 7.4,

150 mM KCl, 5mM EDTA, 0.5% Nonidet P-40, 20 U ml^{-1} RNAsin (Promega), 0,5 mM dithiothreitol (DTT)

and 1× protease inhibitor mixture Complete (Roche)). After centrifugation, the samples were precleared

with Dynabeads Protein A (Invitrogen) for 1 h. One per cent of the sample was used as the input control

and the remaining extracts were incubated with 10 µg DLST antibody (Bethyl) or IgG at 4 °C overnight.

RNA–antibody complexes were collected by incubation with Dynabeads Protein A. After extensive

washing, 1/5 of the sample was used for western blotting control and the remaining for RNA extraction and

qRT–PCR analysis, as described below.

In vitro **RNA pulldown**

RNA pulldown was performed according to [77]. Briefly, biotin-labeled *sin-lncRNA* and control RNA (cloned in

a pcDNA3.1(+) plasmid) were generated in vitro using T7 or T3 RNA polymerase, respectively (Applied

Biosystems). Biotinylated RNAs were treated with RNase-free DNase I (Promega), purified on G-50

MicroSpin columns (GE Healthcare) and heated to 65C for 10 min and cooled down slowly to 4C to allow

proper RNA folding. 10x106 IMR90 ER:RAS were collected after 6 days of senescence induction and total

protein extracts were prepared. Cell extracts were then incubated with 10µg of biotinylated RNA in Buffer

A (150mM KCl, 25mM Tris pH 7.4, 5mM EDTA, 0.5mM DTT, 0.5% NP40) supplemented with fresh protease

inhibitors and yeast tRNA (0.1µg/ml) and rotated overnight at 4C. Dynabeads MyOne Streptavidin T1

beads (Invitrogen) were used to isolate RNA-protein complexes. Beads were washed and boiled in 2X

Laemmli loading buffer, and retrieved proteins were loaded in a NuPAGE 4%–12% bis-Tris gel (Invitrogen).

Gel was stained with the SilverQuest Silver Staining Kit (Thermo Fisher) to identify differential bands

comparing *sin-lncRNA* and control RNA samples. Unique bands were cut and submitted for Mass

Spectrometry (MS) analysis to the Taplin MS Facility (Harvard University).

Chromatin Immunoprecipitation (ChIP)

C/EBPβ ChIP RT-qPCR was performed as previously described[78]. Briefly, 10×10^6 cells were crosslinked with

1% formaldehyde for 10 minutes at RT and nuclei were isolated with cell lysis buffer (5mM Tris pH8.0,

85mM KCl, 0.5% NP40, supplemented with fresh 0,5mM DTT and 1× protease inhibitor mixture Complete

(Roche)). Nuclear pellet was then resuspended in RIPA buffer (1X PBS, 1% NP40, 0.5% Na-deoxycholate,

0.1% SDS, supplemented with 1× protease inhibitor mixture Complete (Roche)) and chromatin was

sheared to a range of 200-500bp by using a Bioruptor instrument (Diagenode) with the following settings:

8 cycles LOW, 30"ON/30"OFF. Sonicated chromatin was incubated overnight with 6µg of C/EBPβ or IgG

antibody (listed in Table 1) followed by 2h incubation with Dynabeads Protein G beads (Invitrogen). After

five washes with LiCl buffer (100mM Tris pH 7.5, 500mM LiCl, 1% NP40, 1% Na-deoxycholate) and TE

buffer, DNA was reverse crosslinked by incubating beads in Elution buffer (1% SDS, 0.1M NaHCO3 and

Proteinase K (NEB)). C/EBPβ binding was analyzed by qPCR using specific primers (see Table 4).

RNA extraction and RT-qPCR analysis

Total RNA was isolated using TRI reagent (Sigma), treated with DNase I (Invitrogen) and reverse-

transcribed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystem) with random

hexamer primers following manufacturer's instructions. The obtained cDNA was analyzed by quantitative

PCR using iTaq Universal SYBR Green supermix (Bio-Rad) in a ViiATM 7 Real-Time PCR System machine

(Thermo-Fisher). All reactions were performed in triplicate or quadruplicate and HPRT1 (Hypoxanthine

Phosphoribosyl transferase 1) RNA levels were used for normalization, unless specified otherwise in figure

legend.

To calculate the number of *sin-lncRNA* molecules per cell, *sin-lncRNA* cloned in pcDNA3.1 vector was in

vitro transcribed, quantified and converted to number of copies based on its predicted molecular weight.

Dilutions of this transcript were retro-transcribed and used as standard curve for qPCR analysis along with

cDNA obtained from a precise number of IMR90 proliferating or senescent cells. All qPCR primer

sequences are provided in Table 4.

RNA sequencing and data analysis

Raw data of Poly-A+ RNA-seq from proliferating and senescent IMR90 fibroblasts were kindly provided by

Dr. Gil's laboratory (Imperial College London, UK) before their publication and are now available at the

Gene Expression Omnibus under the accession number GSE61130 [36]. For RNA-seq of IMR90 ER:RAS

fibroblasts treated with sin-lncRNA or control siRNA, cells were transfected and 4OHT-treated for 3 days,

as described above. Total RNA from samples prepared in biological triplicate was isolated with TRI

reagent, purified and treated with DNAse I using RNAeasy mini Kit (Qiagen). After quality evaluation by

Experion kit and quantification by Qubit, 1μg of RNA was used for library preparation and sequenced on

Illumina HiSeq 2000 (33×10^6 reads per sample, 50bp single-end sequencing modality). Sequenced reads

from three different replicates were aligned to hg19 genome using the BOWTIE2 algorithm (v.2.1.0) was

used to quantify the number of reads in annotated genes [79, 80] To identify enriched molecular pathways

associated with differences in gene expression, Gene Set Enrichment Analysis (GSEA, Broad Institute,

http://www.broadinstitute.org/gsea/index.jsp) was performed using the Hallmark gene sets in the Human

MSigDB database [81]. Enriched pathways were selected with Normalized Enrichment Score (NES) >±1 and

adj.Pvalue<1e^{-4}.

Sin-lncRNA expression analysis in tumor samples

TCGA RNA-seq expression data was retrieved using R bioconductor library TCGAbiolinks, and represented

using R.

Statistical analysis

Unless specified otherwise, experimental data were represented as mean ± standard deviation of at least

three biological replicates (unless otherwise stated in figure legends) and significance was determined by

two-tailed unpaired or Student's t-test or one sample using GraphPad software. Significant P values were

summarized as follows: not significant (ns); p-value<0.05 (*); p- value<0.01 (**); p-value<0.001 (***).

DATA AVAILABILITY

RNA-seq from proliferating and senescent IMR90 fibroblasts were retrieved from public data (GSE61130)

[36]. RNA-seq of *sin-lncRNA* KD and control senescent cells have been deposited in NCBI's Gene Expression

Omnibus and are accessible through Gene Expression Omnibus (GEO) [82] repository under the accession

number GSE241620 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE241620) with the

reviewers token mvylowqmjtotnwx. The mass spectrometry data have been deposited to the

ProteomeXchange Consortium via the PRIDE [83] partner repository with the dataset

identifier PXD044718 (Username: reviewer_pxd044718@ebi.ac.uk, Password: 8egDSySE).

ACKNOWLEDGEMENTS

We thank Eva Santamaría for her input and help with the seahorse experiments. We thank Beatriz Tavira

(CIMA, Universidad de Navarra) for providing the COD362 cells. We thank Lorea Válcarcel (UPV) and

Arkaitz Carracerdo (CIC bioGUNE) for their input on the metabolic experiments and Sebastiaan Martijn

Van Liempd (Metabolomic Platform, CIC bioGUNE) for his interpretation of the metabolomics results. We

thank Victor Segura and Eli Guruceaga, for the guilt-by-association analysis. We thank Jesus Gil (MRC,

London) for his input. We acknowledge the ENCODE project and The Cancer Genome Atlas database for

their valuable data sets. We thank the rest of the Huarte lab for contributing with helpful discussions. The

work was supported by Fondo Europeo de Desarrollo Regional/Ministerio de Ciencia, Innovación y

Universidades-Agencia Estatal de Investigación grant (BFU2017–82773-P), European Research

Council Consolidator grant (771425) and Worldwide Cancer Research grant (20-0204) and FP7-

PEOPLE-2013-ITN. M.M is funded by a MSCA fellowship (HORIZON-MSCA- 2021-PF-01: 101066499).

AUTHORS CONTRIBUTION

M.H. and E.G. conceived the study. E.G, M.M and M.H. designed the experiments, analyzed the data and

wrote the manuscript with the comments of co-authors. E.G., F.M., J.G. and M.M. performed the

experiments. E. G, E. Goñi and A.A. performed bioinformatic analyses. L.B. provided interpretation of the

metabolomics experiment.   N.H. contributed to the original set-up of the project.

COMPETING INTEREST

The authors declare no competing interest.

REFERENCES

1.   Kuilman, T., Michaloglou, C., Mooi, W.J. & Peeper, D.S. The essence of senescence. *Genes &*
development **24**, 2463-2479 (2010).
2.   Serrano, M., Lin, A.W., McCurrach, M.E., Beach, D. & Lowe, S.W. Oncogenic ras provokes premature
cell senescence associated with accumulation of p53 and p16INK4a. *Cell* **88**, 593-602 (1997).
3.   Michaloglou, C. *et al.* BRAFE600-associated senescence-like cell cycle arrest of human naevi. *Nature*
436, 720-724 (2005).
4.   Braig, M. *et al.* Oncogene-induced senescence as an initial barrier in lymphoma development.
Nature **436**, 660-665 (2005).
5.   Acosta, J.C. *et al.* Chemokine signaling via the CXCR2 receptor reinforces senescence. *Cell* **133**,
1006-1018 (2008).
6.   Kuilman, T. *et al.* Oncogene-induced senescence relayed by an interleukin-dependent inflammatory
network. *Cell* **133**, 1019-1031 (2008).
7.   Coppe, J.P. *et al.* Senescence-associated secretory phenotypes reveal cell-nonautonomous
functions of oncogenic RAS and the p53 tumor suppressor. *PLoS biology* **6**, 2853-2868 (2008).
8.   Coppe, J.P., Desprez, P.Y., Krtolica, A. & Campisi, J. The senescence-associated secretory
phenotype: the dark side of tumor suppression. *Annu Rev Pathol* **5**, 99-118 (2010).
9.   Nardella, C., Clohessy, J.G., Alimonti, A. & Pandolfi, P.P. Pro-senescence therapy for cancer
treatment. *Nat Rev Cancer* **11**, 503-511 (2011).
10.   Milanovic, M. *et al.* Senescence-associated reprogramming promotes cancer stemness. *Nature* **553**,
96-100 (2018).
11.   Dorr, J.R. *et al.* Synthetic lethal metabolic targeting of cellular senescence in cancer therapy. *Nature*
501, 421-425 (2013).
12.   Gonzalez-Gualda, E., Baker, A.G., Fruk, L. & Munoz-Espin, D. A guide to assessing cellular
senescence in vitro and in vivo. *FEBS J* **288**, 56-80 (2021).
13.   Kwon, S.M., Hong, S.M., Lee, Y.K., Min, S. & Yoon, G. Metabolic features and regulation in cell
senescence. *BMB Rep* **52**, 5-12 (2019).
14.   Lee, A.C. *et al.* Ras proteins induce senescence by altering the intracellular levels of reactive oxygen
species. *The Journal of biological chemistry* **274**, 7936-7940 (1999).
15.   Moiseeva, O., Bourdeau, V., Roux, A., Deschenes-Simard, X. & Ferbeyre, G. Mitochondrial
dysfunction contributes to oncogene-induced senescence. *Mol Cell Biol* **29**, 4495-4507 (2009).
16.   Kaplon, J. *et al.* A key role for mitochondrial gatekeeper pyruvate dehydrogenase in oncogene-
induced senescence. *Nature* **498**, 109-112 (2013).
17.   Quijano, C. *et al.* Oncogene-induced senescence results in marked metabolic and bioenergetic
alterations. *Cell Cycle* **11**, 1383-1392 (2012).

18.   Bittles, A.H. & Harper, N. Increased glycolysis in ageing cultured human diploid fibroblasts. *Biosci*
666   *Rep* **4**, 751-756 (1984).
667   19.   James, E.L. *et al.* Senescent human fibroblasts show increased glycolysis and redox homeostasis
668   with extracellular metabolomes that overlap with those of irreparable DNA damage, aging, and
669   disease. *J Proteome Res* **14**, 1854-1871 (2015).
670   20.   Kondoh, H. *et al.* Glycolytic enzymes can modulate cellular life span. *Cancer Res* **65**, 177-185 (2005).
671   21.   Mattick, J.S. *et al.* Long non-coding RNAs: definitions, functions, challenges and recommendations.
672   *Nat Rev Mol Cell Biol* **24**, 430-447 (2023).
673   22.   Statello, L., Guo, C.J., Chen, L.L. & Huarte, M. Gene regulation by long non-coding RNAs and its
674   biological functions. *Nat Rev Mol Cell Biol* **22**, 96-118 (2021).
675   23.   Qureshi, I.A. & Mehler, M.F. Emerging roles of non-coding RNAs in brain evolution, development,
676   plasticity and disease. *Nat Rev Neurosci* **13**, 528-541 (2012).
677   24.   Huarte, M. The emerging role of lncRNAs in cancer. *Nat Med* **21**, 1253-1261 (2015).
678   25.   Abdelmohsen, K. *et al.* Senescence-associated lncRNAs: senescence-associated long noncoding
679   RNAs. *Aging cell* **12**, 890-900 (2013).
680   26.   Montes, M. *et al.* The lncRNA MIR31HG regulates p16(INK4A) expression to modulate senescence.
681   *Nat Commun* **6**, 6967 (2015).
682   27.   Montes, M. & Lund, A.H. Emerging roles of lncRNAs in senescence. *FEBS J* (2016).
683   28.   Montes, M. *et al.* The long non-coding RNA MIR31HG regulates the senescence associated
684   secretory phenotype. *Nat Commun* **12**, 2459 (2021).
685   29.   Puvvula, P.K. *et al.* Long noncoding RNA PANDA and scaffold-attachment-factor SAFA control
686   senescence entry and exit. *Nat Commun* **5**, 5323 (2014).
687   30.   Grossi, E. *et al.* A lncRNA-SWI/SNF complex crosstalk controls transcriptional activation at specific
688   promoter regions. *Nat Commun* **11**, 936 (2020).
689   31.   Zhu, Y. *et al.* The long noncoding RNA glycoLINC assembles a lower glycolytic metabolon to
690   promote glycolysis. *Mol Cell* **82**, 542-554 e546 (2022).
691   32.   Wang, P., Xu, J., Wang, Y. & Cao, X. An interferon-independent lncRNA promotes viral replication by
692   modulating cellular metabolism. *Science* **358**, 1051-1055 (2017).
693   33.   Gandhi, M. *et al.* The lncRNA lincNMR regulates nucleotide metabolism via a YBX1 - RRM2 axis in
694   cancer. *Nat Commun* **11**, 3214 (2020).
695   34.   Ding, C. *et al.* A T(reg)-specific long noncoding RNA maintains immune-metabolic homeostasis in
696   aging liver. *Nat Aging* **3**, 813-828 (2023).
697   35.   Castello, A., Hentze, M.W. & Preiss, T. Metabolic Enzymes Enjoying New Partnerships as RNA-
698   Binding Proteins. *Trends Endocrinol Metab* **26**, 746-757 (2015).
699   36.   Herranz, N. *et al.* mTOR regulates MAPKAPK2 translation to control the senescence-associated
700   secretory phenotype. *Nat Cell Biol* **17**, 1205-1217 (2015).
701   37.   Vater, C.A., Bartle, L.M., Dionne, C.A., Littlewood, T.D. & Goldmacher, V.S. Induction of apoptosis by
702   tamoxifen-activation of a p53-estrogen receptor fusion protein expressed in E1A and T24 H-ras
703   transformed p53-/- mouse embryo fibroblasts. *Oncogene* **13**, 739-748 (1996).
704   38.   Hoare, M. *et al.* NOTCH1 mediates a switch between two distinct secretomes during senescence.
705   *Nat Cell Biol* **18**, 979-992 (2016).
706   39.   Innes, A.J. & Gil, J. IMR90 ER:RAS: A Cell Model of Oncogene-Induced Senescence. *Methods Mol*
707   *Biol* **1896**, 83-92 (2019).
708   40.   Woods, D. *et al.* Raf-induced proliferation or cell cycle arrest is determined by the level of Raf
709   activity with arrest mediated by p21Cip1. *Mol Cell Biol* **17**, 5598-5611 (1997).
710   41.   Hutter, E. *et al.* Senescence-associated changes in respiration and oxidative phosphorylation in
711   primary human fibroblasts. *Biochem J* **380**, 919-928 (2004).
712   42.   Zhang, J. & Zhang, Q. Using Seahorse Machine to Measure OCR and ECAR in Cancer Cells. *Methods*
713   *Mol Biol* **1928**, 353-363 (2019).
714   43.   Vizioli, M.G. *et al.* Mitochondria-to-nucleus retrograde signaling drives formation of cytoplasmic
715   chromatin and inflammation in senescence. *Genes & development* **34**, 428-445 (2020).

44. Velarde, M.C., Flynn, J.M., Day, N.U., Melov, S. & Campisi, J. Mitochondrial oxidative stress caused
by Sod2 deficiency promotes cellular senescence and aging phenotypes in the skin. *Aging (Albany*
NY) **4**, 3-12 (2012).
45. Macip, S. *et al.* Inhibition of p21-mediated ROS accumulation can rescue p21-induced senescence.
EMBO J **21**, 2180-2188 (2002).
46. Staal, F.J., Roederer, M., Herzenberg, L.A. & Herzenberg, L.A. Intracellular thiols regulate activation
of nuclear factor kappa B and transcription of human immunodeficiency virus. *Proc Natl Acad Sci U*
S A **87**, 9943-9947 (1990).
47. Leucci, E. *et al.* Melanoma addiction to the long non-coding RNA SAMMSON. *Nature* **531**, 518-522
(2016).
48. Tu, Y.T. & Barrientos, A. The Human Mitochondrial DEAD-Box Protein DDX28 Resides in RNA
Granules and Functions in Mitoribosome Assembly. *Cell Rep* **10**, 854-864 (2015).
49. Sheu, K.F. & Blass, J.P. The alpha-ketoglutarate dehydrogenase complex. *Ann N Y Acad Sci* **893**, 61-
78 (1999).
50. Shen, N. *et al.* DLST-dependence dictates metabolic heterogeneity in TCA-cycle usage among triple-
negative breast cancer. *Commun Biol* **4**, 1289 (2021).
51. Qattan, A.T., Radulovic, M., Crawford, M. & Godovac-Zimmermann, J. Spatial distribution of cellular
function: the partitioning of proteins between mitochondria and the nucleus in MCF7 breast cancer
cells. *J Proteome Res* **11**, 6080-6101 (2012).
52. Wang, Y. *et al.* KAT2A coupled with the alpha-KGDH complex acts as a histone H3
succinyltransferase. *Nature* **552**, 273-277 (2017).
53. Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased coverage,
supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* **47**, D607-
D613 (2019).
54. Alvarez-Abril, B., Garcia-Martinez, E. & Galluzzi, L. Platinum-based chemotherapy inflames the
ovarian carcinoma microenvironment through cellular senescence. *Oncoimmunology* **11**, 2052411
(2022).
55. Murphy, M.P. How mitochondria produce reactive oxygen species. *Biochem J* **417**, 1-13 (2009).
56. Hamanaka, R.B. & Chandel, N.S. Mitochondrial reactive oxygen species regulate cellular signaling
and dictate biological outcomes. *Trends Biochem Sci* **35**, 505-513 (2010).
57. Davalli, P., Mitic, T., Caporali, A., Lauriola, A. & D'Arca, D. ROS, Cell Senescence, and Novel
Molecular Mechanisms in Aging and Age-Related Diseases. *Oxid Med Cell Longev* **2016**, 3565127
(2016).
58. Wei, Z. *et al.* CUL4B impedes stress-induced cellular senescence by dampening a p53-reactive
oxygen species positive feedback loop. *Free Radic Biol Med* **79**, 1-13 (2015).
59. Martinez-Reyes, I. & Chandel, N.S. Mitochondrial TCA cycle metabolites control physiology and
disease. *Nat Commun* **11**, 102 (2020).
60. Hansen, G.E. & Gibson, G.E. The alpha-Ketoglutarate Dehydrogenase Complex as a Hub of Plasticity
in Neurodegeneration and Regeneration. *Int J Mol Sci* **23** (2022).
61. Tretter, L. & Adam-Vizi, V. Inhibition of Krebs cycle enzymes by hydrogen peroxide: A key role of
[alpha]-ketoglutarate dehydrogenase in limiting NADH production under oxidative stress. *J*
Neurosci **20**, 8972-8979 (2000).
62. Sutendra, G. *et al.* A nuclear pyruvate dehydrogenase complex is important for the generation of
acetyl-CoA and histone acetylation. *Cell* **158**, 84-97 (2014).
63. Kafkia, E. *et al.* Operation of a TCA cycle subnetwork in the mammalian nucleus. *Sci Adv* **8**,
eabq5206 (2022).
64. Li, W. *et al.* Nuclear localization of mitochondrial TCA cycle enzymes modulates pluripotency via
histone acetylation. *Nat Commun* **13**, 7414 (2022).
65. Chen, J. *et al.* Compartmentalized activities of the pyruvate dehydrogenase complex sustain
lipogenesis in prostate cancer. *Nat Genet* **50**, 219-228 (2018).
66. Mellid, S. *et al.* DLST mutations in pheochromocytoma and paraganglioma cause proteome
hyposuccinylation and metabolic remodeling. *Cancer Commun (Lond)* (2023).

67.   Ewald, J.A., Desotelle, J.A., Wilding, G. & Jarrard, D.F. Therapy-induced senescence in cancer. *J Natl*
Cancer Inst **102**, 1536-1546 (2010).

68.   Pardella, E. *et al.* Therapy-Induced Stromal Senescence Promoting Aggressiveness of Prostate and
Ovarian Cancer. *Cells* **11** (2022).

69.   Pranzini, E., Pardella, E., Paoli, P., Fendt, S.M. & Taddei, M.L. Metabolic Reprogramming in
Anticancer Drug Resistance: A Focus on Amino Acids. *Trends Cancer* **7**, 682-699 (2021).

70.   Tan, Y. *et al.* Metabolic reprogramming from glycolysis to fatty acid uptake and beta-oxidation in
platinum-resistant cancer cells. *Nat Commun* **13**, 4554 (2022).

71.   Jiang, P., Du, W., Mancuso, A., Wellen, K.E. & Yang, X. Reciprocal regulation of p53 and malic
enzymes modulates metabolism and senescence. *Nature* **493**, 689-693 (2013).

72.   Castello, A. *et al.* Comprehensive Identification of RNA-Binding Proteins by RNA Interactome
Capture. *Methods Mol Biol* **1358**, 131-139 (2016).

73.   Zhang, H. *et al.* LncRNA NEAT1 controls the lineage fates of BMSCs during skeletal aging by
impairing mitochondrial function and pluripotency maintenance. *Cell Death Differ* **29**, 351-365
(2022).

74.   Vendramin, R., Marine, J.C. & Leucci, E. Non-coding RNAs: the dark side of nuclear-mitochondrial
communication. *EMBO J* **36**, 1123-1133 (2017).

75.   Dieterle, F., Ross, A., Schlotterbeck, G. & Senn, H. Probabilistic quotient normalization as robust
method to account for dilution of complex biological mixtures. Application in 1H NMR
metabonomics. *Anal Chem* **78**, 4281-4290 (2006).

76.   Veselkov, K.A. *et al.* Optimized preprocessing of ultra-performance liquid chromatography/mass
spectrometry urinary metabolic profiles for improved information recovery. *Anal Chem* **83**, 5864-
5872 (2011).

77.   Athie, A. *et al.* Analysis of copy number alterations reveals the lncRNA ALAL-1 as a regulator of lung
cancer immune evasion. *J Cell Biol* **219** (2020).

78.   Marin-Bejar, O. *et al.* The human lncRNA LINC-PINT inhibits tumor cell invasion through a highly
conserved sequence element. *Genome Biol* **18**, 202 (2017).

79.   Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

80.   Langmead, B. & Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359
(2012).

81.   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell*
Syst **1**, 417-425 (2015).

82.   Edgar, R., Domrachev, M. & Lash, A.E. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res* **30**, 207-210 (2002).

83.   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based
proteomics evidences. *Nucleic Acids Res* **50**, D543-D552 (2022).

FIGURE LEGENDS

Figure 1. *sin-lncRNA* is a senescence-specific lncRNA transcriptionally induced by C/EBPβ. **A)** Schematic representation of normal fibroblasts undergoing OIS upon 4OHT-mediated activation of an oncogenic form of HRAS. **B)** Volcano plot depicting differential gene expression data (GSE61130) comparing IMR90 ER:RAS proliferating and senescent cells. Cut-off for significance $\log_2$FC>±1; adj.Pvalue < 0.05. A selection of canonical senescence markers (in black) and lncRNAs (in red) is shown. **C)** RT-qPCR analysis of a set of lncRNA candidates upregulated upon 4OHT time course (3 or 6 days) in IMR90 ER:RAS cells; *RIPOR3-as1* was amplified with primer set#2 (see Suppl. Figure 2A for details) Data are presented as mean values ± s.d. n=3/4. **D)** Genomic snapshot of *sin-lncRNA* locus with RNA-seq tracks of untreated and 4OHT-treated IMR90 ER:RAS fibroblasts, as well as C/EBPβ ChIP-seq ENCODE data generated in IMR90 cells. **E)** RT-qPCR analysis of "young" (<14 passages), "old" (>30 passages) IMR90 fibroblasts and 4OHT-induced "OIS" IMR90 ER:RAS cells, shows that *sin-lncRNA* upregulation is maintained in replicative senescence, compared to p16 positive control. (n=2) **F)** Western blot analysis of control and C/EBPβ-depleted senescent IMR90 ER:RAS cells. Proliferating fibroblasts were used as further control. **G)** RT-qPCR analysis of *sin-lncRNA*, as well as a set of C/EBPβ-target genes in control and C/EBPβ-depleted senescent IMR90 ER:RAS cells. Data are represented as mean values ± s.d. (n=2) **H)** ChIP-qPCR analysis of C/EBPβ in proliferating and senescent IMR90 ER:RAS cells upon 6 days of 4OHT treatment. IgG IP was used as IP control; C/EBPβ peaks at *sin-lncRNA* locus are labeled according to their distance from *sin-lncRNA* TSS; regions located in *DDX11*, *GAPDH* and *HPRT* promoters served as negative controls, *IL6* and *IL8* promoter regions as positive controls (n=2).

Figure 2. *sin-lncRNA* depletion reinforces the senescent phenotype. **A)** RT-qPCR analysis of control and *sin-lncRNA* depleted IMR90 ER:RAS cells after 3 days of senescence induction. Primer sets #1 was used to amplify *sin-lncRNA* (see Suppl. Table4). Proliferating control cells were used as further control. Values were normalized to senescent control cells. **B)** Crystal violet assay and **C)** BrdU analysis assessing cell growth of control and *sin-lncRNA* depleted IMR90 ER:RAS fibroblasts after 5 days of senescence induction.

D) β-galactosidase analysis of control and *sin-lncRNA* depleted IMR90 cells induced to senescence for 8 days. **E)** Western blot analysis of control and *sin-lncRNA* depleted IMR90 cells treated with 4OHT for 5 days. **F)** RT-qPCR analysis of TIG3 ΔBRAF cell clones infected with *sin-lncRNA* sgRNA or an empty plasmid. **G)** BrdU (left) and β-galactosidase (right) assays of control or *sin-lncRNA* sgRNA TIG3 ΔBRAFclones. **H)** RT-qPCR and **I)** BrdU analysis of IMR90 ER:RAS cells transiently transfected with empty or *sin-lncRNA* construct upon OIS.

Figure 3. *sin-lncRNA* KD alters oxidative stress response and metabolic pathways in senescent cells. **A)** Volcano plot depicting differentially expressed genes comparing *sin-lncRNA* KD vs control IMR90 ER:RAS senescent cells. Upregulated genes (adj.Pvalue < 0.01; log2FC>1) are labeled in red, downregulated genes (adj.Pvalue < 0.01; log2FC<-1) in blue. **B)** Hallmark Gene Set Enrichment Analysis (GSEA) reporting the most enriched pathways associated to the genes differentially expressed (DE) upon *sin-lncRNA* depletion (threshold: adj.Pvalue<1e-4). **C)** Heatmap highlighting expression changes across RNA-seq replicates of genes belonging to OXIDATIVE_PHOSPHORILATION GSEA pathway. **D)** Quantification of different parameters of the oxygen consumption rates (OCR) in control and *sin-lncRNA* depleted (siRNA#1/3) IMR90 ER:RAS, assessed through Seahorse technology. The graph represents the mean ± s.d. (n=4). Two tailed student's *t*-test, *p<0.05. A representative OCR curve for each cell line is included in (Supplementary Fig. 4). **E)** Basal extracellular acidification rate (ECAR) levels measured using the seahorse technology in control or *sin-lncRNA* depleted IMR90 ER:RAS senescent cells. Values were normalized to the total amount of protein per condition. **F)** Fluorometric analysis measuring the amount of the fluorescent oxidized probe 2',7'-dichlorofluorescein (DCF), as indicative of ROS/NOS intracellular levels in control and *sin-lncRNA* depleted IMR90 ER:RAS fibroblasts induced to senescence for 5 days. Proliferating control cells were used as further control. **G)** Absorbance levels of crystal violet solution measured at 560 nm from vehicle or N-acetyl-L-cisteine (NAC)-treated IMR90 ER:RAS fibroblasts transfected with siRNA control (CTRL) or against *sin-lncRNA* (sin#1/3) and induced to senescence for 5 days. The graph represents the mean ± s.d. (n=3). Two tailed student's *t*-test, *p<0.05.

Figure 4. *sin-lncRNA* interacts with DLST and affects metabolism of senescent cells by regulating DLST localization. **A)** RNA-FISH based on locked nucleic acid (LNA) technology on fixed IMR90 ER:RAS cells depleted or not of *sin-lncRNA* (siRNA_sin-lncRNA#1/3) and treated for 5 days with 4OHT. Two different LNAs were used. Incubation without LNA probe ("No_probe") was used as control. **B)** List of top candidate proteins identified by mass spectrometry after pulldown with in vitro biotinylated *sin-lncRNA*. **C)** Schematic representation of the TCA cycle that takes place in the mitochondria. DLST enzyme is highlighted in red. **D)** RNA immunoprecipitation analysis monitoring DLST binding to *sin-lncRNA, MALAT1* and *GAPDH* RNA in control and *sin-lncRNA* depleted (siRNA#1/3) senescent IMR90 ER:RAS, using anti-DLST antibody or IgG as a negative control. The graph represents the mean ± s.d. (n=2). **E)** *Left*, representative β-galactosidase images of untreated control cells and control, *sin-lncRNA* depleted or DLST-depleted IMR90 cells induced to senescence for 5 days. Scale bar: 50 μm. *Right*, quantification of the percentage of β-galactosidase positive cells in each condition normalized to control untreated cells set as 1. The graph represents the mean ± s.d. (n=4). Two tailed student's *t*-test, **p<0.01. **F)** Absorbance levels of crystal violet solution measured at 560 nm in the conditions described in E. The graph represents the mean ± s.d. (n=4). Two tailed student's *t*-test, *p<0.05. **G)** Lactate levels measured from the supernatant of IMR90 ER:RAS fibroblasts transfected with the indicated siRNA and induced with 4OHT for 3 or 6 days. The graph represents the mean ± s.d. (n=3). Two tailed student's *t*-test, *p<0.05; **p<0.01; ***p<0.001. **H)** *Left*, western blot analysis of DLST cellular fractionation in control and *sin-lncRNA* depleted (siRNA#1/3) senescent IMR90 ER:RAS. GAPDH and H3 were used as cytoplasmic and nuclear controls respectively. *Right*, percentage of nuclear and cytoplasmic distribution of DLST normalized to GAPDH (cyto) or H3 (nucl). The graph represents the mean ± s.d. (n=3). Two tailed student's *t*-test, **p<0.01.

Figure 5. *sin-lncRNA* regulates OXPHOS in ovarian cancer cells. **A)** Representative β-galactosidase images of untreated COV362 cells (upper panel) or treated for 10 days with μM cisplatin. Scale bar: 50 μm. Numbers represent the mean ± s.d. (n=3). **B)** RT-qPCR analysis of senescence markers genes in cisplatin treated COV362 for 10 days compared to untreated controls (set as 1). **C)** RT-qPCR analysis of *sin-lncRNA* in the cells described in B. The graph represents the mean ± s.d. (n=4). One sample *t*-test, **p<0.01. **D)**

Western blot analysis of C/EBPβ protein levels in the cells described in B. Numbers represent the mean ± s.d. (n=3). **E)** RT-qPCR analysis of OXPHOS GSEA pathway genes in control and *sin-lncRNA* depleted (siRNA#1/3) COV362 senescent cells treated with cisplatin for 10 days. **F)** Quantification of different parameters of the oxygen consumption rates (OCR) in proliferating cells compared to control or *sin-lncRNA* depleted (siRNA#1/3) senescent COV362 cells, assessed through Seahorse technology. The graph represents the mean ± s.d. (n=4). Two tailed student's *t*-test, *p<0.05; **p<0.01. **G)** Basal extracellular acidification rate (ECAR) levels measured using the seahorse technology in the same conditions as in F. The graph represents the mean ± s.d. (n=4). Two tailed student's *t*-test, *p<0.05. **H)** Absorbance levels of crystal violet solution measured at 560 nm in proliferating or senescent COV362 cells transfected with control, *sin-lncRNA* or DLST siRNA treated for 10 days with 10 μM cisplatin.

SUPPLEMENTARY FIGURE LEGENDS

Suppl. Figure 1. A) Change in proliferation rate comparing proliferating and senescent IMR90 ER:RAS fibroblasts, as measured by BrdU assay. **B)** Representative images of crystal violet staining in untreated or 4-OHT treated cells for 5 days. **C)** Representative images of the β-galactosidase staining in untreated cells or treated with 4-OHT for 8 days. **D)** qRT-PCR analysis of CXCL1 and IL1A upregulation in untreated or 4-OHT treated cells for 5 days. **E)** Western blot analysis of senescence markers p53, p21 and p16 in untreated or 4-OHT treated cells for 5 days. **F)** DAPI staining and **G)** Immunostaining images of gH2A.X, H3K96m3 and DAPI of untreated or 4OHT treated IMR90 ER:RAS for 5 days. **H)** Pie chart representation of the number of biotype distribution of transcripts deregulated during OIS (log_2-fold change >±1, adj.Pvalue <0.05). **I)** Heat map showing relative expression of senescence markers and lncRNAs deregulated in IMR90 ER:RAS untreated or treated with 4-OHT for 8 days.

Suppl. Figure 2. A) Genomic snapshot of *sin-lncRNA* (RIPOR3-as) locus PolyA sites annotated by Genecode v19 are indicated as well as the position of qPCR primers used to amplify the different isoforms. Bottom panel, qRT-PCR analysis of the different annotated isoforms of *sin-lncRNA* (RIPOR3-as) using different set of primers (set1-3) in a time course of 3 and 6 days of senescence induction in IMR90 ER:RAS cells. **B)** qRT-PCR analysis of protein-coding gene RIPOR3 in a time course of 3 and 6 days of senescence induction in IMR90 ER:RAS cells. **C)** CPAT software was interrogated to determine the coding probability of *sin-lncRNA*. GAPDH mRNA and MALAT1 lncRNA were used as coding and noncoding references, respectively. **D)** Number of copies per cell of *sin-lncRNA* in control untreated cells and in 4-OHT-treated cells measured by qRT–PCR using dilutions of a pcDNA3.1-*sin-lncRNA* plasmid as standard curve. **E)** RT-qPCR analysis showing *sin-lncRNA* upregulation upon treatment with γ-irradiation at 5Gy in IMR90 fibroblasts. *p16* and *IL6* levels served as positive controls. **F)** Western blot analysis of C/EBPβ of IMR90 ER:RAS cells treated or not with 4OHT for 3 or 6 days. **G)** qRT-PCR analysis of the levels of C/EBPβ in control and C/EBPβ-depleted senescent IMR90 ER:RAS cells

Suppl. Figure 3. A) RT-qPCR analysis of control and *sin-lncRNA* depleted IMR90 ER:RAS cells after 3 days of senescence induction. Proliferating control cells were used as further control. Primer sets #2 (Suppl. Figure 2A) were used to amplify *sin-lncRNA*. Values were normalized to senescent control cells. **B)** BrdU analysis measuring the proliferation of control and *sin-lncRNA* depleted IMR90 ER:RAS cells after 5 days of senescence induction. Proliferating control cells were used as further control. Values were normalized to senescent control cells. **C)** RT-qPCR analysis of control and *sin-lncRNA* depleted TIG3-hTERT:BRAF cells after 3 days of senescence induction. Proliferating control cells were used as further control. Primer sets #3 (Suppl. Figure 2A) were used to amplify *sin-lncRNA*. Values were normalized to senescent control cells. **D)** BrdU staining and **E)** relative growth measured by crystal violet at 570nm absorbance of control and *sin-lncRNA* depleted TIG3-hTERT:BRAF cells after 5 days of senescence induction. Proliferating control cells were used as further control. Values were normalized to senescent control cells. **F)** Schematic representation of the CIRSPR/Cas9 strategy applied to remove *sin-lncRNA* genetic loci. sgRNAs guiding Cas9 protein are indicated (sgRNA_up and sgRNA_dw), as well as inner and outer primer couples to screen engineered cell clones (fwd/rev_in and fwd/rev_out, respectively) **G)** Gel images showing PCR products obtained using outer (left) or inner primer sets (right) to amplify the genome of TIG3 BRAF single cell clones after infection with the lentiCRISPR carrying *sin-lncRNA* sgRNAs and selection (clone#n). TIG3 BRAF fibroblasts infected with empty lentiviral vector (Empty) or infected with vector carrying *sin-lncRNA* sgRNAs but without single cell separation and puro selection (CRISPR_unsorted) were used as control. **H)** Representation of *sin-lncRNA* isoform amplified from cDNA samples and cloned in pcDNA3.1 plasmid. **I)** BrdU analysis measuring the proliferation of IMR90 ER:RAS cells transiently transfected with empty or *sin-lncRNA* expression vector and 4OHT-treated for 5 days. Proliferating cells transfected with empty vector were used as further control.

Suppl. Figure 4. A) RT-qPCR analysis of the indicated transcripts in control and *sin-lncRNA* depleted IMR90 ER:RAS cells after 5 days of senescence induction. Proliferating control cells were used as further control. Values were normalized to senescent control cells. **B)** Gene Set Enrichment Analysis (GSEA) for genes belonging to epithelial to mesenchymal transition and E2F-targets pathways. **C)** Oxygen

Consumption Rate (OCR) measurements were obtained over time (min) using the seahorse technology from control or *sin-lncRNA* depleted IMR90 ER:RAS senescent cells. Values were normalized to the total amount of protein per condition. The graph shows a representative experiment. **D)** Percentage of β-galactosidase positive cells in control or *sin-lncRNA* depleted IMR90 ER:RAS senescent cells treated with 5mM vehicle (DMSO) or N-acetyl-L-cisteine (NAC). **E)** Percentage of nuclear and cytoplasmic RNA levels of *sin-lncRNA* measured by qRT–PCR after cell fractionation with two different set of primers (set#1 and set#2) in untreated or 4OHT-induced senescent cells. **F)** Western blot analysis of the immunoprecipitated DLST using anti-DLST antibody or IgG as a negative control. GAPDH was used as control. (n=2) **G)** Western blot analysis of control IMR90 ER:RAS cells or cells transfected with siRNAs against *sin-lncRNA* or DLST. Proliferating control cells were used as further control. **H)** RT-qPCR analysis in control, DLST or *sin-lncRNA* depleted IMR90 ER:RAS cells after 3 days of senescence induction. Values were normalized to senescent control cells. Data are presented as mean ± s.d (n=4). Two tailed student's *t*-test, *p<0.05; **p<0.01. **I)** Western blot analysis of DLST protein levels in untreated IMR90 ER:RAS cells or treated for 3 or 6 days. GAPDH was used as a loading control.

Suppl. Figure 5. A) *Sin-lncRNA* expression (as FPKM + pseudocount value) across tumor samples from the TCGA database. **B)** Protein-Protein Interaction network (*p-value*<$1.0e^{-16}$) by STRING analysis of genes inversely correlated with *sin-lncRNA* (92 genes; R< -0.5). **C)** Gene Ontology (GO) pathways associated with the inversely correlated genes.

A CTRL

11.3% ±9.8

Cisplatin (10d)

81.3% ±11.4

B CDKN2A CDKN1A IL8 CCNA2

C sin-lncRNA

D CTRL Cisplatin

C/EBPβ

GAPDH

1   1.84±0.43

E siRNA

F siRNA

G

H

A

B

PPI enrichment p-value < 1.0e⁻¹⁶

C

GO analysis - 92 inversely correlated genes (R< -0.5)

	ID	Name	p-value	FDR B&H
1	GO:0022402	cell cycle process	$6.52e^{-19}$	$1.63e^{-15}$
2	GO:0000278	mitotic cell cycle	$2.64e^{-16}$	$1.65e^{-13}$
3	GO:0051301	cell division	$6.14e^{-13}$	$3.07e^{-10}$