Ultrahigh Resolution Mass Spectrometry Imaging Maps Brain Lipidome Alterations Associated with Mild Traumatic Brain Injury

Dmitry Leontyev[1], Alexis N. Pulliam[2], Xin Ma[1], David A. Gaul[1,3], Michelle C. LaPlaca[2,3], Facundo M. Fernández[1,3]*

[1]School of Chemistry and Biochemistry, Georgia Institute of Technology, Atlanta, GA 30332 (USA)

[2]Coulter Department of Biomedical Engineering, Georgia Institute of Technology/Emory University, Atlanta, GA 30332 (USA)

[3]Parker H. Petit Institute for Bioengineering and Bioscience, Atlanta, GA 30332 (USA)

Dmitry Leontyev: dmitry.leontyev@gatech.edu

Alexis N. Pulliam: anpullia3@gatech.edu

Xin Ma: xin.ma@chemistry.gatech.edu

David A. Gaul: david.gaul@chemistry.gatech.edu

Michelle C. LaPlaca: michelle.laplaca@bme.gatech.edu

*Facundo M. Fernandez: facundo.fernandez@chemistry.gatech.edu 404-385-4432 311 Ferst Dr Atlanta, GA, USA 30332

Abbreviations: mTBI, mild traumatic brain injury; CCI, controlled cortical impact; ROI, region of interest; PLS-DA, partial least squares discriminant analysis; AUC, area under the curve; FC, fold change; PC, phosphatidylcholine; LPC, lysophosphatidylcholine; LPC-P, lysophosphatidylcholine-plasmalogens; PI-Cer, ceramide phosphoinositols; SM, sphingomyelin; PUFA, polyunsaturated fatty acid; DHA, docosahexaenoic acid.

Abstract

Background: Traumatic brain injury (TBI) is a global public health problem, with 50-60 million incidents per year, most of which are considered mild (mTBI). Despite its massive impact, the pathology of TBI is not fully understood, and there is a paucity of information on brain lipid dysregulation following mTBI. To gain more insight on mTBI pathology, a non-targeted spatial metabolomics workflow utilizing ultrahigh resolution mass spectrometry imaging was developed to measure brain region-specific lipid alterations in rats.

Results: Multivariate models were created for regions of interest including the hippocampus, cortex, corpus callosum, white matter and gray matter to identify lipids that discriminated between control and injured brains. The hippocampus model differentiated control and injured brains with an area under the curve of 0.994, using only four lipid markers. Lipid classes that were consistently selected for discrimination included polyunsaturated fatty acid -containing phosphatidylcholines (PC), lysophosphatidylcholines (LPC), LPC-plasmalogens (LPC-P), PC potassium adducts and ceramide phosphoinositols (PI-Cer). Many of the polyunsaturated fatty acid-containing PC, LPC-P, and PI-Cer selected have never been previously reported as altered in TBI.

Significance: The lipid alterations observed indicate that neuroinflammation, oxidative stress and disrupted sodium-potassium pumps are important pathologies that can explain cognitive deficits associated with mTBI. Therapeutics which target and attenuate these pathologies may be beneficial to limit persistent damage following a mild brain injury.

Keywords: Mild traumatic brain injury, mass spectrometry imaging, multivariate analysis, Fourier transform ion cyclotron resonance, lipidomics, modified controlled cortical impact

1. Introduction

TBI is a complex disease caused by excessive external force to the head that alters normal brain function. After the primary physical tissue damage, a secondary injury response cascade occurs, with damage following from neuroinflammation, oxidative stress, ion imbalance, disruption of the blood brain barrier and mitochondrial dysfunction[1]. This secondary damage can progress through a patient's life, causing long-term physical, emotional and cognitive disabilities[2] that include increased risk of Alzheimer's disease[3], Parkinson's disease[4], stroke[5], epilepsy[6] and chronic traumatic encephalopathy[7, 8].

Despite its widespread incidence, TBI pathology is not completely understood, partly due to the heterogeneity and complexity of the disease. TBI causes vary greatly and are dependent on premorbid health status, resulting in a wide clinical manifestation and unpredictable recovery trajectory. Furthermore, TBI varies in severity from mild to severe. Research has largely focused on severe TBI, yet over 90% of hospital cases are categorized as mild (mTBI)[9]. Repetitive mTBI (rmTBI) is a particular concern to sports like football and boxing where athletes sustain countless mild to moderate impacts to the head over the span of years. At the cellular and tissue level, heterogeneity is evident in the complexity of secondary injury cascades. These secondary

responses are reflected in lipid alterations since the brain is one of the most lipid rich organs[10] and lipid metabolism is known to be disrupted after injury. Brain lipid alterations associated with TBI and other neurodegenerative diseases have been shown to induce neuroinflammation and high abundance of polyunsaturated fatty acid (PUFA)-containing lipids make the brain more vulnerable to oxidative stress.

One of the key techniques used to map lipid composition in the brain is matrix-assisted laser desorption/ionization (MALDI) mass spectrometry imaging (MSI). Roux *et al.* tracked the lipid changes in rat brains over time using a silver nanoparticle MALDI matrix to discover significant time-dependent changes in ceramides (CER), sphingomyelins (SM) and diacylglycerols (DG) following TBI[11]. Other TBI MSI studies have reported decreases in cardiolipins (CL) and phosphatidylinositols (PI)[12], time-dependent changes in docosahexaenoic acid (DHA)-containing phospholipids[13], increases in lysophosphatidic acid (LPA)[14] and increases in acylcarnitines that colocalize to microglia[15]. While these studies have provided very valuable insights, the majority have focused on modeling severe injuries involving a craniotomy followed by a direct impact to the brain. Given the high incidence rate of mTBI and rmTBI, and the poor understanding of cellular mechanisms that render the brain vulnerable to long-term conditions, this study focused on rmTBI. rmTBI was induced by a modified controlled cortical impact (CCI) procedure where rats received three closed head impacts with a silicon padded pneumatic piston. We previously tracked serum lipid changes in rats over time following single mTBI and rmTBI using a lipidomics approach, building machine learning models able to classify injury status[16]. A study comparing brain lipid changes in rmTBI and Alzheimer's disease mice models found that lysophosphatidylcholine (LPC) increased in the hippocampus in both diseases[17].

Here, we applied ultrahigh mass resolution Fourier Transform Ion Cyclotron Resonance (FTICR) MSI to study rmTBI. The key advantages of FTICR MSI reside in its unparalleled mass resolving power and mass accuracy, leading to high confidence lipid annotation and highly specific select-ion images. With FTICR MSI, lipid elemental formula can be typically established with sub-ppm mass errors and mass resolving power of at least 300,000 (FWHM). Results presented here show highly-discriminant brain region-specific lipid alterations in poly unsaturated fatty acid (PUFA)-containing phosphatidylcholines (PC), LPC, LPC-plasmalogens (LPC-P), PC potassium adducts and ceramide phosphoinositols (PI-Cer) associated with rmTBI.

2. Experimental section

2.1. Chemicals and Materials

Isopentane (≥99%; Sigma-Aldrich, St. Louis, MO) was used to prepare dry ice cold baths for embedding. The embedding mixture was prepared using gelatin from bovine skin (Sigma-Aldrich), carboxymethyl cellulose sodium salt medium viscosity (Sigma-Aldrich) and ultrapure water with 18.2 MΩ cm resistivity (Barnstead Nanopure, Thermo Fisher Scientific, Waltham, MA). The MALDI matrix solution was made from 1,5-diaminonaphthalene (DAN) (97%; Sigma-Aldrich), LC-MS grade acetonitrile (Fisher Scientific International, Pittsburg, PA, USA) and ultrapure water. LC-MS grade methanol, LC-MS grade water, LC-MS grade acetonitrile, LC-MS grade isoproponal, formic acid, ammonium acetate and ammonium formate (Fisher Scientific

International, Pittsburg, PA, USA) were used to prepare mobile phases for LC-MS/MS experiments.

2.2. Animals and Modified Controlled Cortical Impact Procedures

All procedures performed on Sprague-Dawley rats were conducted in accordance with the guidelines set forth in the Guide for the Care and Use of Laboratory Animals (U.S. Department of Health and Human Services, Washington, DC, USA, Pub no. 85-23, 1985) and approved by the Georgia Institute of Technology Institutional Animal Care and Use Committee (protocol #A100188). The handling of rats and modified CCI procedure to induce mTBI was previously described in detail by our group[16]. Briefly, male Sprague-Dawley rats ($n=12$) (Charles River, Wilmington, MA, USA) weighing between 250-300 grams were kept under 12 h reverse light-dark cycles with food and water *ad libitum* and randomly assigned to the sham group ($n=6$) or the three-impact group ($n=6$). Prior to injury, rats were anesthetized and maintained with 2-3% isoflurane. Rats were then placed on 1-inch-thick ethylene-vinyl acetate foam (McMaster-Carr, Elmhurst, IL, USA). mTBI was induced by subjecting rats to three closed head impacts (2 min interval, 5 m s^{-1} velocity, 5 mm, 2 mm, and 2 mm head displacement) to the dorsal head surface using a CCI pneumatic injury device (Pittsburgh Precision Instruments, Pittsburgh, PA, USA). This device was equipped with a 1-cm diameter silicone stopper (Renovators Supply Manufacturing, Erving, MA, USA) attached to the piston tip. The sham group underwent identical procedures as the injured group, except for the impacts. Following the last impact, righting time was recorded, and rats were returned to their cages for recovery. Rats were anesthetized 72 hours post injury with 2-3% isoflurane and perfused with cold pH 7.4 phosphate buffer. Brains were then extracted, the cerebellum was excised, and the brains separated into hemispheres for sagittal sectioning. The hemispheres were snap frozen in a dry ice cold isopentane bath, embedded in a 5% gelatin/1% carboxymethyl cellulose mixture and stored at -80°C until sectioning.

2.3. Sectioning and Matrix Application

The Paxinos and Watson rat brain atlas[18] was used to guide sectioning with anatomical landmarks including the hippocampus and corpus callosum. Sections chosen for imaging were approximately 1500-2000 µm lateral to bregma. Sagittal 12-µm sections collected serially from right brain hemispheres using a cryostat (Thermo Shandon NX70 Cryostar, Waltham, MA) were mounted onto indium-tin-oxide (ITO) slides (Delta Technologies, Loveland, CO) and stored at -80°C until MALDI MSI. One brain was lost to sectioning issues that rendered the tissue not usable for imaging, leaving $n=5$ for the sham group and $n=6$ for the injury group. Prior to imaging, slides were placed in a desiccator for 15 minutes and sprayed with 8 passes of a 5 mg mL^{-1} DAN solution in 90% acetonitrile/10% water using an HTX TM-Sprayer (HTX Technologies, Chapel Hill, NC) at 30°C, 0.1 mL min^{-1} flow rate, 1200 mm min^{-1} velocity, 2.5 mm tracking speed, 10 psi, 2 L min^{-1} gas flow rate, 0 second drying time and 40 mm nozzle height.

2.4. Mass Spectrometry Imaging Parameters

MALDI imaging data were collected on a solariX 12T FTICR mass spectrometer (Bruker Daltonics, Bremen, Germany) in positive ion mode in the m/z 147-1500 range using 2M transients (~300,000 mass resolution at m/z 314). A 50 µm raster spacing in the x and y directions was used. The laser was set to 100 shots, small focus, 12 % power and 1000 Hz. Real time calibration with a lock mass of m/z 314.152598 from the DAN dimer and m/z 760.585082 from PC(34:1) was used to achieve optimal mass accuracy. Additional MS parameters are provided in Table S1 and Table S2.

2.5. Image Data Processing and Region of Interest Selection

Two sections from each of the five sham and six injured brains were examined by FTICR MSI. Serial sections were placed on the same ITO slide at the time of the experiment. Among the sections examined, two replicate images were eliminated from the dataset due to abnormally low ion abundances. A total of twenty images from eleven rats were used for multivariate image analysis with eight of those sections being from sham animals and twelve from injured animals. MS images were uploaded to and analyzed in SCiLS Lab Version 2022b Pro (Bruker Daltonics, Bremen, Germany). Regarding pre-processing options, no baseline correction or other notable options were used. For segmentation purposes, a feature list was created with the sliding window tool using the average mass spectrum from all brain sections and the lowest possible intensity threshold. This yielded a large peak list with a ± 3 ppm window for each ion. This feature list was used to computationally segment the brain images into molecularly similar areas and specific ROI were chosen for analysis. The parameters used for segmentation were the preliminary feature list, root mean square normalization, strong denoising, bisecting k-means and the Manhattan distance metric. Segmentation was performed on individual brain sections or regions. In a few cases, some ROI were not correctly picked out by the automated segmentation approach alone and were thus manually outlined following specific lipid distributions that helped delineate the ROI borders.

2.6. Region of Interest Feature List Creation and Refinement

For each ROI, feature lists were first created in SCiLS Lab with a ± 5 ppm feature tolerance. Receiver operating characteristic analysis was then conducted on these ROI-specific feature lists using the mean spectra. All ions with an area under the curve above 0.7 (*i.e.*, those with abundances larger in control brains), or those with AUC below 0.3 (more abundant in injured animals) were chosen. The remaining m/z values were filtered out from the input feature list. For ROI involving the gray matter, the white matter and the corpus callosum, AUC cutoff values were set to a stricter cutoff of 0.8 and 0.2, as the corresponding input feature lists contained a large abundance of ions. Extracted ion images for m/z values in the resulting feature lists were inspected to remove any species originating from the MALDI matrix, the embedding mixture, or ions with poor signal-to-noise ratios. The spectral profile for each feature was inspected to ensure that the interval chosen by SCiLS software was correctly aligned with the apex of each peak, and the feature list interval tolerance then lowered to ± 3 ppm.

2.7. Model Building

Multivariate models were created in Metaboanalyst[19] for each of the selected ROI using the respective feature lists. Averaged and root mean square-normalized ion abundances were extracted from each ROI and uploaded to Metaboanalyst. Pareto scaling was applied to all ion abundances. Partial Least Squares-Discriminant Analysis classification with t-statistic feature ranking led to the best performing classification models. The number of latent variables and features used in each model were optimized for each ROI to maximize AUC values. The resulting ROC curves were created by Monte-Carlo cross validation. With this approach, two thirds of the samples were used to determine feature importance, the top-ranking features were used to create models and the other one third of samples used to validate the model. This procedure was then repeated several times to determine overall model performance. Models were further optimized by trimming down the input feature list to only those species that led to the best AUC values. Any redundant isotopic features or adduct species in the feature lists were removed.

2.8. Feature Annotation and Liquid Chromatography-Tandem Mass Spectrometry

Feature annotation based on exact mass matches was conducted in LIPID MAPS[20] (https://www.lipidmaps.org) using a mass tolerance of ± 0.005 Da. Data used for annotation was collected from a sham brain section using a modified method. This modified method adjusted parameters including auto-calibration that improved the rigor and accuracy of the reported *m/z* values. The MS method parameters used in annotation experiments are listed in Table S2.

Reverse phase ultra-high-performance LC-MS/MS data was collected to confirm the identity of the lipids identified by LIPID MAPS. This reverse phase LC-MS/MS method was previously described in detail by our group[21]. Chromatography was performed with a Thermo Accucore C30, 150 × 2.1 mm, 2.6 µm particle size column. Mobile phase A was 10 mM ammonium acetate 0.1% formic acid with water/acetonitrile (40:60 v/v) and mobile phase B was 10 mM ammonium acetate 0.1% formic acid with isopropanol/acetonitrile (90:10 v/v). High-energy collision dissociation (HCD) and collision induced dissociation (CID) data was collected in an Orbitrap ID-X Tribrid mass spectrometer (ThermoFisher Scientific). Negative mode data was used to determine lipids fatty acid chain length and positive mode data was used to confirm the presence of the phosphocholine headgroup.

3. Results & Discussion

3.1. Region of interest selection and Multivariate classification

Figure 1 outlines the experimental workflow followed in this study. Representative MS images for a sham brain section and an injured section are given in Figure 2 (Figures S1 and S2 show the corresponding images from the remaining sections). The similarity in the anatomical features delineated by FTICR MSI and the robust reproducibility of data collected is exemplified in Figure S3.

The closed head injury used in this study resulted in no macroscopic signs of physical trauma. Neurologically, a statistically significant difference (p<0.05) in the righting time[22]

between sham and injured rats was observed (Figure S4), indicating that animals sustained detectable acute deficits via the inflicted rmTBI. The lack of visible physical trauma resulted in no overt injury region easily detectable by MALDI, as in previous open head TBI studies[11]. Therefore, regions of interest (ROI) were chosen to follow the outline of gross anatomical structures in the dorsal part of the sagittal brain section (Figure 3). The brains were segmented into white matter and gray matter (Figure 3B) and these regions were analyzed to examine lipidome changes at the level of the whole brain. Within the white matter and gray matter there are several regions known to be affected by TBI that were analyzed including the corpus callosum, the hippocampus and the cortex. The corpus callosum was selected using the segmentation tool, while the hippocampus was manually outlined with ion images that help show the regions' borders (Figure 3C). Segmentation of the ROI containing the cortex and the corpus callosum (white outline in Figure 3C) resulted in 8-9 separate segments that followed the general expected structure of the various cortical brain layers[23]. The outermost 1-2 segments from all brains were analyzed as this is the region where the most pronounced effects of the injury are expected. If a brain's outermost segment contained too few pixels, the layer under it was also selected as part of the outer cortex such that each ROI contained approximately 2500-3500 pixels. Outlines of the hippocampus, corpus callosum and outer cortex ROI for all imaged sections are given in Figure S5. The segmentation tool was able to select anatomical regions of interest consistently and reproducibly across different rat brains, despite the slight differences in the sizes and shapes of the anatomical features.

Using the average ion abundances within each ROI, Partial Least Squares-Discriminant Analysis (PLS-DA) models were created to compare sham and injured brains (Figure 4 and S6). Receiver operating characteristic (ROC) curves were generated by Metaboanalyst using Monte-Carlo cross-validation. During cross-validation, the top ion features are used to build PLS-DA models that are tested on an independent test set selected prior to model building. PLS-DA model performances were optimized by removing redundant and/or low-ranking ion features and by choosing models with features that resulted in the best area under the curve (AUC). Annotations for these features are given in Table 1. The relationship between important lipid classes is summarized in Figure S7.

3.2. PLS-DA models for regions of interest

The most pronounced alterations following TBI were in the hippocampus. The hippocampus model had an AUC of 0.994 using only 4 lipid features and 4 latent variables (Figure 5A). These were, in order of decreasing importance, PC(40:6), LPC(16:0), SM(42:2) $[M+K]^+$ and PC(36:1). The respective fold changes (FC) were 0.91, 1.27, 0.45 and 0.93 (Table 1). The hippocampus is especially susceptible to TBI[24], in agreement with these findings. Evidence shows synaptic reorganization and decreases in the number of interneurons in the hippocampus after TBI, altering the delicate balance between excitation and inhibition[24, 25]. This balance is important for proper brain function, and its disruption is thought to be related to long term deficits in TBI[24]. Furthermore, one of the most significant chronic deficits associated with TBI is in hippocampal-associated memory and learning impairment, and there is evidence that even subtle cellular changes in the hippocampus are related to cognitive and memory dysfunction[26, 27].

The outer cortex model had an AUC of 0.977 using 8 features and 5 latent variables (Figure 5B). There was an expectation for lipid distributions to be highly altered at the outer cortex since the rats are impacted on the dorsal head surface. The lipids in the model (in order of decreasing importance) were LPC(16:0), LPC(18:1), PC(38:4), PC(40:4), LPC(O-16:1), PC(32:1), LPC(O-18:2), and PC(38:6) (Figure 5B). The respective FC in the outer cortex were 1.20, 1.43, 0.95, 0.84, 1.35, 0.90, 0.76, and 1.06 (Table 1). All LPC species besides LPC(O-18:2) increased in the outer cortex after injury, whereas all PUFA-containing PC besides PC(38:6) decreased in the outer cortex after injury. There are several unique findings about this model including that all ions belonged to the PC or LPC classes, and that there are 3 PUFA-containing PC. The large AUC in the outer cortex PLS-DA model underscores the fact that the cortex is especially susceptible to TBI[24]. This is supported by evidence showing that, as with the hippocampus, there is synaptic reorganization and decreases in the number of interneurons in the neocortex after TBI[24, 25]. TBI can also induce neuronal hyperexcitation in the upper cortical layers[24].

The white matter model had an AUC of 0.96 using 8 features and 2 latent variables (Figure 5C). The features in the model, in order of decreasing importance, were PI-Cer(d40:0), LPC(16:0), PC(36:2) $[M+K]^+$, PC(36:1) $[M+K]^+$, LPC(O-16:1), SM(d40:1), LPC(O-18:2) and PC(36:4). Their FC are given in Table 1. The $[M+K]^+$ adduct ions were of particular interest. These PC potassium adducts were only selected for the white matter and corpus callosum models, suggesting that these regions showed disruption of membrane ion pumps, as discussed in the section devoted to PC lipids below. TBI is known to cause damage to the white matter, such as atrophy of white matter axons, a process known as traumatic axonal injury[28]. As axonal tracts are responsible for brain connectivity, white matter damage can also disrupt signaling between cortical areas, interfering with recovery from TBI[29]. White matter volume loss has been shown to be a predictor of recovery and a determinant for executive functions like verbal fluency after penetrating TBI[29].

The gray matter model had an AUC of 0.92 using 4 features and 3 latent variables (Figure 5D). The features in this model, in order of decreasing importance, were LPC(16:0), PI-Cer(d40:0), LPC(O-16:1), and LPC(18:1). Their FC are given in Table 1. All these ions increased in the injured gray matter as compared to controls. Three out of four ions in the gray matter model were LPC, highlighting their importance in mTBI as discussed in the LPC section below. Gray matter in the central nervous system has a crucial role in enabling humans' normal daily functions such as controlling movement, memory and emotions[30]. Various diffusor tensor imaging studies have shown increased fractional anisotropy in rat brain gray matter after TBI, indicating neurodegeneration and myelinated fiber loss[31,32,33].

The corpus callosum model had an AUC of 0.864 using 5 features and 1 latent variable (Figure S6). The performance of this model was the lowest of all the PLS-DA models. The main role of the corpus callosum is to connect and enable communication between the hemispheres. Damage to the corpus callosum has been shown to be associated with loss of social cognitive ability, which is complex and requires inter-hemispheric connection[34]. Damage to the corpus callosum in mTBI patients is also associated with prolonged neurological symptoms such as dizziness and gait disturbance[35]. Ionic gradient changes arise in the medial corpus callosum after TBI and may spread to white matter structure in lateral regions and induce lipid changes there[36]. One reason that could explain the lower performance of the corpus callosum model is that the

closed head injury examined in this study has less of an impact in this brain structure as in other investigated brain regions.

3.3. Phosphatidylcholines (PC) and Polyunsaturated fatty acids (PUFA)

Examination of all PLS-DA models created for various ROI showed that there were 9 PC used: PC(32:1), PC(36:4), PC(36:1), PC(38:6), PC(38:4), PC(36:2) $[M+K]^+$, PC(36:1) $[M+K]^+$, PC(40:6) and PC(40:4). LC-MS/MS was used to determine the fatty acid chain length of these lipids (Table 1) and several of these PC were found to contain PUFA. TBI MSI studies have reported alterations in DHA (FA(22:6)), arachidonic acid (FA(20:4)) and various PUFA-containing lipids including PI, PS, PE, DG, and lysophospholipids [11-13]. Reports of PUFA-containing PC changes are less common. PLS-DA models in our study selected PC(16:0/20:4), PC(16:0/22:6), PC(18:0/20:4), PC(18:0/22:6), and PC(16:0/24:4). Notably, PC(38:6) and PC(40:6) contained DHA, while PC(36:4) and PC(38:4) contained arachidonic acid. These chain length findings are supported by the ion images of these lipids (Figure 6), which show that PC(38:6) and PC(40:6) have similar distributions and so do PC(36:4) and PC(38:4). PC(40:6) was the 1st ranking ion in the hippocampus model based on the mean importance measure and PC(38:4) was ranked 3rd in the outer cortex model. Brain injury and neurodegenerative processes rely on PUFA, which are produced by PLA2 cleavage of phospholipids, to regulate inflammatory response[12]. DHA is thought to reduce neuroinflammation, reduce oxidative stress and activate cell survival pathways[13]. Additionally, DHA supplementation following TBI has been shown to counteract learning disability and restore mechanisms that maintain brain homeostasis[13].

FC values in the white matter and gray matter regions for the PC(36:2) $[M+K]^+$ adduct were 0.83 and 0.88, respectively. The FC values for the PC(36:1) $[M+K]^+$ adduct in these regions were 0.88 and 0.89. Both results suggested that PC potassium adducts systematically decreased after injury. The TBI literature has recognized a disruption and decrease in sodium-potassium pump (Na/K-ATPase) activity in the brain, leading to decreases in $[PC+K]^+$ adducts and increases in $[PC+Na]^+$[37, 38]. Hypokalemia, low levels of potassium in the blood, is common in TBI patients when compared to other trauma[39]. Loss of Na/K-ATPase activity impacts ion flux, creating an ion imbalance in the brain that disrupts homeostasis[39]. This imbalance is accompanied by an influx of extracellular fluid that causes edema[28]. This series of events may contribute to cognitive impairment after TBI[38]. The two PC potassium adducts were only selected by the corpus callosum and white matter models, suggesting that the Na/K-ATPase activity disruption affects the white matter to a larger extent than the gray matter.

3.3.1. Lysophosphatidylcholines (LPC)

LPC were top ranking discriminatory ions in all models. The LPC selected by PLS-DA models were LPC(O-16:1), LPC(16:0), LPC(O-18:2) and LPC(18:1). Of these, LPC(16:0) was the only ionic species present in all 5 PLS-DA models and was a top ranking ion in most models (Figure 5). In the white matter and gray matter ROI, the FC values for LPC(16:0) were 1.28 and 1.23, respectively. In the same regions, FC values for LPC(O-16:1), LPC(O-18:2), LPC(18:1) were 2.61 and 1.69, 2.64 and 2.50, and 1.22 and 1.45, respectively (Table 1). These indicated that LPC increased in injured brains. Some of these FC values were the highest amongst all ions in all models

(Table 1), highlighting their importance in mTBI processes. While LPC may also form *via* MALDI in-source fragmentation of PC, we concluded that this formation pathway was not predominant as imaging experiments were conducted at very low laser power (12%).

Cytosolic PLA2 (cPLA2) cleaves PC located on membranes at the sn2 position, releasing the sn2 chain that commonly contains PUFA, leaving LPC in the membrane, which alters membrane fluidity and permeability[40, 41]. cPLA2 activity increases after TBI which increasingly produces LPC that have been shown to induce the release of proinflammatory cytokines[42], induce apoptosis of neurons, induce immune cell migration[42], induce oligodendrocyte demyelination[43], mediate pericyte loss[43] and cause cognitive deficit[44]. Additionally, LPC have been associated with brain injury[17, 41, 45-47], Alzheimer's disease, multiple sclerosis, ischemic stroke[46], atherosclerosis, diabetes and liver disease[47]. Palafox-Sanchez *et al.* investigated how TBI induces systemic changes in the liver and alters lipid metabolism in the brain, identifying LPC as the most altered lipids in the hippocampus[47]. High levels of LPC(16:0) were associated with poorer memory performance, due to the proinflammatory action of LPC[47]. TBI is also associated with an increase of PUFA such as DHA that mediate neuroinflammation and oxidative stress. These PUFA can be produced from cPLA2 cleaving PUFA-containing PC. Interestingly, cPLA2 produces LPC that are proinflammatory and releases PUFA that are anti-inflammatory. However, the overall effects of increased cPLA2 activity are detrimental as demonstrated by multiple studies where inhibiting PLA2 showed beneficial outcomes[40, 48]. After TBI, cPLA2 is involved in lysosomal membrane permeabilization, which impairs autophagy and leads to neuronal cell death[40]. Inhibiting cPLA2 reduced these effects and improved neurological outcomes.

3.3.2. Ether-linked Lysophosphatidylcholines (LPC-O/LPC-P)

Two LPC species containing ether linkages were selected in several of the PLS-DA models. LPC(O-16:1) was selected for the corpus callosum, white matter, gray matter, and outer cortex models. LPC(O-18:2) was selected for the white matter and outer cortex models. The FC values in the white matter and gray matter for LPC(O-16:1) were 2.61 and 1.69. For LPC(O-18:2), the FC were 2.64 and 2.50 in these regions. Both lipid species significantly increased after injury. The double bond position of these LPC were not experimentally determined, however the LIPID MAPS database indicated that LPC(O-18:2) is a plasmalogen and can be referred to as LPC(P-18:1). Plasmalogens are unique lipids with vinyl-ether bonds, which are double bonds adjacent to their ether linkage.

MSI studies have reported decreases in ether-linked lipids such as PE(P-40:6) and PE(P-38:6) after TBI[11, 13], however ether-linked LPC and PC are less reported in TBI literature. This is likely because PE-P are concentrated in the brain, whereas PC-P are more abundant in the heart and kidney[49]. A study investigating the effect of induced ketosis in Alzheimer's patients reported improvements in cognitive function that were accompanied with increases in plasma LPC(P-18:0) and LPC(P-18:1)[50]. This last species is equivalent to LPC(O-18:2) detected here. PE-P deficiency in the blood and brain has been associated with Alzheimer's pathology and various other neurological disorders[51]. Orally supplementing with ether phospholipids has been shown to restore cognitive dysfunction in behavioral studies[51]. Reducing endogenous plasmalogens in mice hippocampi has been shown to induce neuroinflammation and has been associated with

significantly reduced learning and memory performance[49]. The beneficial cognitive observations associated with plasmalogens are likely partly due to their association with PUFA since plasmalogens commonly contain PUFA at the sn-2 position[51]. PC-P could be increasingly cleaved to release PUFA to help regulate inflammation and oxidative stress after TBI.

3.3.3. Sphingomyelins (SM) and Ceramides (CER)

The SM(d42:2) $[M+K]^+$ adduct ion was used in the hippocampus model and SM(d40:1) in the white matter model. SMs are formed by reaction of a PC and a CER, which also produces DG as a byproduct. The SM/CER signaling pathway is associated with apoptosis[52] and has been shown to be activated after ischemic brain injury and TBI[11, 53]. A MSI study found significant time dependent and chain length dependent changes in SM, CER and DG in rat brain after TBI including SM(d42:2)[11]. SM(d42:2) appears to be particuarly important to TBI based on previous literature reports. Interestingly, SM(d42:2)$[M+K]^+$ had a significant decrease in abundance with a 0.45 FC in the hippocampus, which can partly be attributed to the decreases of Na/K-ATPase activity after injury, as discussed above.

Alteration in specific ceramide phosphoinositol (PI-Cer) species was also observed. PI-Cer(d40:0) was used by the gray matter, white matter, and corpus callosum models. PI-Cer(d40:1) was used in the corpus callosum model. FC values in the white matter and gray matter for PI-Cer(d40:1) were 1.33 and 1.80, respectively. FC for PI-Cer(d40:0) were 1.21 and 1.73, respectively. These data indicate that PI-Cer increased after injury. PI-Cer(40:0) was the highest ranking ion, in terms of importance, in the corpus callosum and white matter models. PI-Cer are the sphingolipid analogue of PI and are formed from the reaction between PI and Cer. PI-Cer are present in plants, fungi and some bacteria, but they are uncommonly reported in mammals[54]. Although their identity could not be confirmed with LC-MS/MS here, a MSI study on mouse brains detected the $[M+H]^+$ of PI-Cer(d40:0), confirming the identity with MS/MS[55]. There is little information about the biological role of PI-Cer in the brain of mammals in general, however they are likely involved in the SM/CER signaling pathway. In plants, PI-Cer are thought to protect tissue from necrotic lesions[54]. If PI-Cer had an equivalent role in the brain, their substantial increase after injury could be rationalized as affecting a neuroprotective role.

This study was limited to a small sample size and to positive ionization mode, which allowed for examination of low abundance species that have biological significance. Another limitation was the multivariate models created focused on the optimum set of lipids that maximized discrimination between injured and non-injured brain tissue for specific regions of interest, which may exclude other lipids that could still be differential but may have lower fold changes. These models were primarily used as tools to determine which lipids could discriminate between control and injured region of interest and were built using cross-validation to help address overfitting.

4. Conclusions

In summary, we investigated the molecular level alterations in brain lipids following mTBI, using rats as a model and an exquisite level of mass resolving power. mTBI, which has no robust

accurate diagnostics, could be predicted by multivariate models harnessing the markers detected by ultrahigh resolution FTICR imaging, if these were also present in easily obtainable biofluids. Using FTICR MSI enabled ions with mDa mass differences to be distinguished, including low abundance species that are often not resolved from nearby higher abundance peaks and go undetected. Results showed significant alterations in numerous lipid classes in brain tissue, some never before reported in the literature. Amongst these were LPC, which are proinflammatory lipids that increase following TBI through the cleavage of PC. LPC(16:0), which is associated with lowered memory performance, was an ion of interest in all examined tissue regions that included the hippocampus, the outer cortex, the white matter, the gray matter, and the corpus callosum. For the first time ether linked LPC, such as LPC(O-16:1) and LPC(O-18:2), were shown to be increased in the white matter after mTBI. Alterations were also observed for DHA-containing PC such as PC(40:6) and PC(38:6). Enzymatic cleavage of these lipids releases DHA, which has been shown to protect the brain and aid in recovery after injury by reducing neuroinflammation and oxidative stress. Another notable finding was decreased abundances of PC and SM potassium adducts, likely due to disrupted Na/K-ATPase activity. These species reflect changes in ion flux that disrupt biological processes in the brain and lead to cognitive impairment. The brain lipid alterations observed in this study indicate that neuroinflammation, oxidative stress and disrupted Na/K-ATPase activity are important pathologies that can explain cognitive deficits associated with mTBI. Therapeutics which target and attenuate these pathologies may be beneficial to limit persistent damage following a brain injury.

Authorship contribution statement

Dmitry Leontyev: Investigation. Writing – original draft. Data curation. Formal analysis. **Alexis N. Pulliam:** Investigation**. Xin Ma:** Writing – review & editing. **David A. Gaul:** Supervision. **Michelle C. LaPlaca:** Project administration and conceptualization. **Facundo M. Fernandez:** Project administration and conceptualization.

Declaration of competing interest

The authors declare no competing financial interests or personal relationships that could have influenced the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

This project has been funded by the National Institute of Neurological Disorders and Stroke (NINDS) (R01NS101909). The authors also acknowledge support from the National Science Foundation (MRI CHE-1726528 grant) for the acquisition of a Fourier transform ion cyclotron resonance (FTICR) mass spectrometer for the Georgia Institute of Technology core facilities. This

material is also based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. (DGE-2039655: A.N.P.), and National Institute of Health under Grant No. (T32 GM145735: A.N.P.). The authors acknowledge support from Aqua Asberry from the Georgia Institute of Technology Histology Core for guidance in tissue sectioning.

Appendix A. Supplementary data

Supplementary data to this article can be found online at

Figure 1. Study workflow. (**A**) Male Sprague-Dawley rats were randomly assigned to a sham group (*n*=5) or the three-impact group (*n*=6). (**B**) Rats were sacrificed 72 h post injury, the brains were extracted, separated into hemispheres, flash frozen, embedded in a gelatin mixture and sagittally-sectioned at 12 μm thickness. (**C**) ITO slides were sprayed with 8 passes of a 5 mg mL^{-1} DAN solution. (**D**) MALDI MSI data were collected on a Bruker solariX 12T FTICR mass spectrometer in positive ion mode. (**E**) Imaging data were loaded into Bruker SCiLS Lab software and the segmentation tool was used to select ROI. (**F**) ROC analysis was used to refine the lipid feature list for each selected ROI. (**G**) PLS-DA classification models for specific ROI were created in Metaboanalyst. (**H**) For annotation purposes, tandem MS experiments were performed on the discriminant species of importance in multivariate models. Created with BioRender.

Figure 2. Representative images for rat brain tissue sections. Extracted ion images resulting from sham rat tissues (A,B) and injured rats (C,D) are shown. (A,C) show the distribution of the [M+H]+ ion for LPC(16:0) (*m/z* 496.34158 ± 3 ppm) and (B,D) display the distribution of the [M+K]+ ion for PC(36:1) (*m/z* 826.57584 ± 3 ppm). These sample images demonstrate an overall relative increase of LPC(16:0) and a decrease of PC(36:1) [M+K]+ in the injured brain. Root mean square normalization, hot spot removal and weak denoising were used. The average mass spectrum showcases the most prominent species observed in all brain images. Species below *m/z* 700 are primarily matrix and embedding mixture ions. Species from *m/z* 700-900 contain most of the lipids of interest.

Figure 3. Region of interest selection. **(A)** H&E stained section showing anatomical features relative to mass spectrometry images. The cortex, corpus callosum (CC) and hippocampus (HC) are outlined. The anterior side of the brain is labeled. **(B)** Segmentation of the white matter (WM) in yellow and gray matter (GM) in blue. The red segment surrounding the brain section corresponds to the area of embedding matrix adjacent to the tissue section itself. **(C)** Segmentation of the cortical area above the hippocampus resulted in segments resembling the corpus callosum and the layered outer cortex (OC). To compare the outer cortex layers across different brains, segments containing 2500-3500 pixels were used. The parameters used for segmentation were root mean square normalization, strong denoising, bisecting k-means and the Manhattan distance metric. The ion image for PC(36:1) at *m/z* 788.61738 (± 3 ppm) is shown in the background of panel (C). This image was used as a visual aid to outline the hippocampus.

Figure 4. PLS-DA model performance for each ROI using t-statistic ranking. ROC curves display 95% confidence intervals (CI). **(A)** hippocampus: 0.994 AUC using 4 features and 4 latent variables. **(B)** outer cortex: 0.977 AUC using 8 features and 5 latent variables. **(C)** white matter: 0.96 AUC using 8 features and 2 latent variables. **(D)** gray matter: 0.92 AUC using 4 features and 3 latent variables.

Figure 5. Mean importance measure for ions in each PLS-DA model. **(A)** PLS-DA model for the hippocampus ROI, **(B)** model for the outer cortex ROI, **(C)** model for the white matter ROI **(D)** model for the gray matter ROI. The importance measure for each lipid variable was calculated using Metaboanalyst. The higher the importance measure of each lipid species in each model, the higher their contribution to the overall model score. The labels on the right side of each plot denote the studied groups: Group A is the sham group and group B is the injured animal group. A blue square under group A indicates that the abundance of the corresponding ion is lower in the specific ROI for sham animals vs. injured animals. A red square under group A indicates that the corresponding ion abundance was higher in the specific ROI for sham animals vs. injured animals. See Figure S6 for the corpus callosum model.

Figure 6. Ion images for lipids in model. H&E-stained section to show anatomy relative to mass spectrometry images. Some ions such as LPC(O-18:2) shown here appear to be low intensity, however their intensity is higher in other brain sections. LPC(O-18:2) distribution resembles that of LPC(18:1) but is lower in intensity.

Table 1. Annotation of features used in PLS-DA models and their fold changes. Fold change (FC) was calculated as the ratio of injured/control (>1 means increase in injured). *m/z* and mass error values were calculated using a separate brain section. LC-MS/MS data was collected to confirm the headgroup of lipids and to determine chain length. ND: Not detected. HC: hippocampus, OC: outer cortex, WM: white matter, GM: gray matter, CC: corpus callosum.

m/z	mass error (ppm)	ID	LC-MS/MS	FC HC	FC OC	FC WM	FC GM	FC CC	Model
480.34319	0.20	LPC(O-16:1)	Phosphocholine	1.70	1.35	2.61	1.69	2.03	WM, GM, OC, CC
496.33905	0.23	LPC(16:0)	Phosphocholine	1.27	1.20	1.28	1.23	1.29	HC, WM, GM, OC, CC
506.35965	0.37	LPC(O-18:2)	Phosphocholine	1.74	0.76	2.64	2.50	1.70	WM, OC
522.35426	0.08	LPC(18:1)	Phosphocholine	1.49	1.43	1.22	1.45	1.20	GM, OC
732.55507	0.13	PC(32:1)	PC(16:0/16:1)	1.05	0.90	1.03	0.96	1.06	OC
782.56847	-0.18	PC(36:4)	PC(16:0/20:4)	1.01	0.99	1.03	1.00	1.03	WM
787.66851	-0.07	SM(d40:1)	Phosphocholine	0.45	1.48	1.47	1.18	2.10	WM
788.61681	0.39	PC(36:1)	PC(18:0/18:1)	0.93	0.96	0.97	1.02	0.96	HC
806.56772	-0.06	PC(38:6)	PC(16:0/22:6)	0.97	1.06	0.99	1.05	1.00	OC
810.59903	0.08	PC(38:4)	PC(18:0/20:4)	0.97	0.95	0.99	0.97	0.97	OC
824.55638	-0.14	PC(36:2) [M+K]$^+$	PC(18:1/18:1)	0.84	0.95	0.83	0.88	0.84	WM
826.57211	0.31	PC(36:1) [M+K]$^+$	PC(18:0/18:1)	0.83	0.92	0.88	0.89	0.92	WM, CC
834.60021	0.07	PC(40:6)	PC(18:0/22:6)	0.91	0.96	0.96	1.00	0.95	HC
838.62987	-0.19	PC(40:4)	PC(16:0/24:4)	0.94	0.84	1.01	0.95	1.06	OC
851.64284	0.02	SM(d42:2)[M+K]$^+$	Phosphocholine	0.45	1.17	0.99	1.03	1.07	HC
864.63126	-0.09	PI-Cer(d40:1)	-	0.70	ND	1.33	1.80	1.34	CC
866.64937	-0.18	PI-Cer(d40:0)	-	0.88	0.62	1.21	1.73	1.31	GM, WM, CC

References

1.   Khatri, N.; Thakur, M.; Pareek, V.; Kumar, S.; Sharma, S.; Datusalia, A. K., Oxidative Stress: Major Threat in Traumatic Brain Injury. *CNS Neurol. Disord. Drug Targets* **2018,** *17* (9), 689-695.

2.   Maas, A. I. R.; Menon, D. K.; Adelson, P. D.; Andelic, N.; Bell, M. J.; Belli, A.; Bragge, P.; Brazinova, A.; Büki, A.; Chesnut, R. M.; Citerio, G.; Coburn, M.; Cooper, D. J.; Crowder, A. T.; Czeiter, E.; Czosnyka, M.; Diaz-Arrastia, R.; Dreier, J. P.; Duhaime, A.-C.; Ercole, A.; van Essen, T. A.; Feigin, V. L.; Gao, G.; Giacino, J.; Gonzalez-Lara, L. E.; Gruen, R. L.; Gupta, D.; Hartings, J. A.; Hill, S.; Jiang, J.-y.; Ketharanathan, N.; Kompanje, E. J. O.; Lanyon, L.; Laureys, S.; Lecky, F.; Levin, H.; Lingsma, H. F.; Maegele, M.; Majdan, M.; Manley, G.; Marsteller, J.; Mascia, L.; McFadyen, C.; Mondello, S.; Newcombe, V.; Palotie, A.; Parizel, P. M.; Peul, W.; Piercy, J.; Polinder, S.; Puybasset, L.; Rasmussen, T. E.; Rossaint, R.; Smielewski, P.; Söderberg, J.; Stanworth, S. J.; Stein, M. B.; von Steinbüchel, N.; Stewart, W.; Steyerberg, E. W.; Stocchetti, N.; Synnot, A.; Te Ao, B.; Tenovuo, O.; Theadom, A.; Tibboel, D.; Videtta, W.; Wang, K. K. W.; Williams, W. H.; Wilson, L.; Yaffe, K.; Adams, H.; Agnoletti, V.; Allanson, J.; Amrein, K.; Andaluz, N.; Anke, A.; Antoni, A.; van As, A. B.; Audibert, G.; Azaševac, A.; Azouvi, P.; Azzolini, M. L.; Baciu, C.; Badenes, R.; Barlow, K. M.; Bartels, R.; Bauerfeind, U.; Beauchamp, M.; Beer, D.; Beer, R.; Belda, F. J.; Bellander, B.-M.; Bellier, R.; Benali, H.; Benard, T.; Beqiri, V.; Beretta, L.; Bernard, F.; Bertolini, G.; Bilotta, F.; Blaabjerg, M.; den Boogert, H.; Boutis, K.; Bouzat, P.; Brooks, B.; Brorsson, C.; Bullinger, M.; Burns, E.; Calappi, E.; Cameron, P.; Carise, E.; Castaño-León, A. M.; Causin, F.; Chevallard, G.; Chieregato, A.; Christie, B.; Cnossen, M.; Coles, J.; Collett, J.; Della Corte, F.; Craig, W.; Csato, G.; Csomos, A.; Curry, N.; Dahyot-Fizelier, C.; Dawes, H.; DeMatteo, C.; Depreitere, B.; Dewey, D.; van Dijck, J.; Đilvesi, Đ.; Dippel, D.; Dizdarevic, K.; Donoghue, E.; Duek, O.; Dulière, G.-L.; Dzeko, A.; Eapen, G.; Emery, C. A.; English, S.; Esser, P.; Ezer, E.; Fabricius, M.; Feng, J.; Fergusson, D.; Figaji, A.; Fleming, J.; Foks, K.; Francony, G.; Freedman, S.; Freo, U.; Frisvold, S. K.; Gagnon, I.; Galanaud, D.; Gantner, D.; Giraud, B.; Glocker, B.; Golubovic, J.; Gómez López, P. A.; Gordon, W. A.; Gradisek, P.; Gravel, J.; Griesdale, D.; Grossi, F.; Haagsma, J. A.; Håberg, A. K.; Haitsma, I.; Van Hecke, W.; Helbok, R.; Helseth, E.; van Heugten, C.; Hoedemaekers, C.; Höfer, S.; Horton, L.; Hui, J.; Huijben, J. A.; Hutchinson, P. J.; Jacobs, B.; van der Jagt, M.; Jankowski, S.; Janssens, K.; Jelaca, B.; Jones, K. M.; Kamnitsas, K.; Kaps, R.; Karan, M.; Katila, A.; Kaukonen, K.-M.; De Keyser, V.; Kivisaari, R.; Kolias, A. G.; Kolumbán, B.; Kolundžija, K.; Kondziella, D.; Koskinen, L.-O.; Kovács, N.; Kramer, A.; Kutsogiannis, D.; Kyprianou, T.; Lagares, A.; Lamontagne, F.; Latini, R.; Lauzier, F.; Lazar, I.; Ledig, C.; Lefering, R.; Legrand, V.; Levi, L.; Lightfoot, R.; Lozano, A.; MacDonald, S.; Major, S.; Manara, A.; Manhes, P.; Maréchal, H.; Martino, C.; Masala, A.; Masson, S.; Mattern, J.; McFadyen, B.; McMahon, C.; Meade, M.; Melegh, B.; Menovsky, T.; Moore, L.; Morgado Correia, M.; Morganti-Kossmann, M. C.; Muehlan, H.; Mukherjee, P.; Murray, L.; van der Naalt, J.; Negru, A.; Nelson, D.; Nieboer, D.; Noirhomme, Q.; Nyirádi, J.; Oddo, M.; Okonkwo, D. O.; Oldenbeuving, A. W.; Ortolano, F.; Osmond, M.; Payen, J.-F.; Perlbarg, V.; Persona, P.; Pichon, N.; Piippo-Karjalainen, A.; Pili-Floury, S.; Pirinen, M.; Ple, H.; Poca, M. A.; Posti, J.; Van Praag, D.; Ptito, A.; Radoi, A.; Ragauskas, A.; Raj, R.; Real, R. G. L.; Reed, N.; Rhodes, J.; Robertson, C.; Rocka, S.; Røe, C.; Røise, O.; Roks, G.; Rosand, J.; Rosenfeld, J. V.; Rosenlund, C.; Rosenthal, G.; Rossi, S.; Rueckert, D.; de Ruiter, G. C. W.; Sacchi, M.; Sahakian, B. J.; Sahuquillo, J.; Sakowitz, O.; Salvato, G.; Sánchez-Porras, R.; Sándor, J.; Sangha, G.; Schäfer, N.; Schmidt, S.; Schneider, K. J.; Schnyer, D.; Schöhl, H.; Schoonman, G. G.; Schou, R. F.; Sir, Ö.; Skandsen, T.; Smeets, D.; Sorinola, A.; Stamatakis, E.; Stevanovic, A.; Stevens, R. D.; Sundström, N.; Taccone, F. S.; Takala, R.; Tanskanen, P.; Taylor, M. S.; Telgmann, R.; Temkin, N.; Teodorani, G.; Thomas, M.; Tolias, C. M.; Trapani, T.; Turgeon, A.; Vajkoczy, P.; Valadka, A. B.; Valeinis, E.; Vallance, S.; Vámos, Z.; Vargiolu, A.; Vega, E.; Verheyden, J.; Vik, A.; Vilcinis, R.; Vleggeert-Lankamp, C.; Vogt, L.; Volovici, V.; Voormolen, D. C.; Vulekovic, P.; Vande Vyvere, T.; Van Waesberghe, J.; Wessels, L.; Wildschut, E.; Williams, G.; Winkler, M. K. L.; Wolf, S.; Wood, G.; Xirouchaki, N.; Younsi, A.; Zaaroor, M.; Zelinkova, V.; Zemek, R.; Zumbo, F., Traumatic brain injury:

integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol.* **2017,** *16* (12), 987-1048.

3.   Li, W.; Risacher, S. L.; McAllister, T. W.; Saykin, A. J., Traumatic brain injury and age at onset of cognitive impairment in older adults. *J Neurol* **2016,** *263* (7), 1280-5.

4.   Gardner, R. C.; Burke, J. F.; Nettiksimmons, J.; Goldman, S.; Tanner, C. M.; Yaffe, K., Traumatic brain injury in later life increases risk for Parkinson disease. *Ann Neurol* **2015,** *77* (6), 987-95.

5.   Liao, C. C.; Chou, Y. C.; Yeh, C. C.; Hu, C. J.; Chiu, W. T.; Chen, T. L., Stroke risk and outcomes in patients with traumatic brain injury: 2 nationwide studies. *Mayo Clin Proc* **2014,** *89* (2), 163-72.

6.   Walsh, S.; Donnan, J.; Fortin, Y.; Sikora, L.; Morrissey, A.; Collins, K.; MacDonald, D., A systematic review of the risks factors associated with the onset and natural progression of epilepsy. *Neurotoxicology* **2017,** *61*, 64-77.

7.   Lucke-Wold, B. P.; Turner, R. C.; Logsdon, A. F.; Bailes, J. E.; Huber, J. D.; Rosen, C. L., Linking traumatic brain injury to chronic traumatic encephalopathy: identification of potential mechanisms leading to neurofibrillary tangle development. *J Neurotrauma* **2014,** *31* (13), 1129-38.

8.   Hay, J.; Johnson, V. E.; Smith, D. H.; Stewart, W., Chronic Traumatic Encephalopathy: The Neuropathological Legacy of Traumatic Brain Injury. *Annu. Rev. Pathol.* **2016,** *11*, 21-45.

9.   Maas, A. I. R.; Menon, D. K.; Manley, G. T.; Abrams, M.; Akerlund, C.; Andelic, N.; Aries, M.; Bashford, T.; Bell, M. J.; Bodien, Y. G.; Brett, B. L.; Buki, A.; Chesnut, R. M.; Citerio, G.; Clark, D.; Clasby, B.; Cooper, D. J.; Czeiter, E.; Czosnyka, M.; Dams-O'Connor, K.; De Keyser, V.; Diaz-Arrastia, R.; Ercole, A.; van Essen, T. A.; Falvey, E.; Ferguson, A. R.; Figaji, A.; Fitzgerald, M.; Foreman, B.; Gantner, D.; Gao, G.; Giacino, J.; Gravesteijn, B.; Guiza, F.; Gupta, D.; Gurnell, M.; Haagsma, J. A.; Hammond, F. M.; Hawryluk, G.; Hutchinson, P.; van der Jagt, M.; Jain, S.; Jain, S.; Jiang, J. Y.; Kent, H.; Kolias, A.; Kompanje, E. J. O.; Lecky, F.; Lingsma, H. F.; Maegele, M.; Majdan, M.; Markowitz, A.; McCrea, M.; Meyfroidt, G.; Mikolic, A.; Mondello, S.; Mukherjee, P.; Nelson, D.; Nelson, L. D.; Newcombe, V.; Okonkwo, D.; Oresic, M.; Peul, W.; Pisica, D.; Polinder, S.; Ponsford, J.; Puybasset, L.; Raj, R.; Robba, C.; Roe, C.; Rosand, J.; Schueler, P.; Sharp, D. J.; Smielewski, P.; Stein, M. B.; von Steinbuchel, N.; Stewart, W.; Steyerberg, E. W.; Stocchetti, N.; Temkin, N.; Tenovuo, O.; Theadom, A.; Thomas, I.; Espin, A. T.; Turgeon, A. F.; Unterberg, A.; Van Praag, D.; van Veen, E.; Verheyden, J.; Vyvere, T. V.; Wang, K. K. W.; Wiegers, E. J. A.; Williams, W. H.; Wilson, L.; Wisniewski, S. R.; Younsi, A.; Yue, J. K.; Yuh, E. L.; Zeiler, F. A.; Zeldovich, M.; Zemek, R.; In, T. P.; Investigators, Traumatic brain injury: progress and challenges in prevention, clinical care, and research. *Lancet Neurol* **2022,** *21* (11), 1004-1060.

10.   Piomelli, D.; Astarita, G.; Rapaka, R., A neuroscientist's guide to lipidomics. *Nat Rev Neurosci* **2007,** *8* (10), 743-54.

11.   Roux, A.; Muller, L.; Jackson, S. N.; Post, J.; Baldwin, K.; Hoffer, B.; Balaban, C. D.; Barbacci, D.; Schultz, J. A.; Gouty, S.; Cox, B. M.; Woods, A. S., Mass spectrometry imaging of rat brain lipid profile changes over time following traumatic brain injury. *J. of Neurosci. Methods* **2016,** *272*, 19-32.

12.   Sparvero, L. J.; Amoscato, A. A.; Fink, A. B.; Anthonymuthu, T.; New, L. A.; Kochanek, P. M.; Watkins, S.; Kagan, V. E.; Bayir, H., Imaging mass spectrometry reveals loss of polyunsaturated cardiolipins in the cortical contusion, hippocampus, and thalamus after traumatic brain injury. *J Neurochem* **2016,** *139* (4), 659-675.

13.   Guo, S.; Zhou, D.; Zhang, M.; Li, T.; Liu, Y.; Xu, Y.; Chen, T.; Li, Z., Monitoring changes of docosahexaenoic acid-containing lipids during the recovery process of traumatic brain injury in rat using mass spectrometry imaging. *Sci Rep* **2017,** *7* (1), 5054.

14.   McDonald, W. S.; Jones, E. E.; Wojciak, J. M.; Drake, R. R.; Sabbadini, R. A.; Harris, N. G., Matrix-Assisted Laser Desorption Ionization Mapping of Lysophosphatidic Acid Changes after Traumatic Brain Injury and the Relationship to Cellular Pathology. *Am J Pathol* **2018,** *188* (8), 1779-1793.

15.   Mallah, K.; Quanico, J.; Raffo-Romero, A.; Cardon, T.; Aboulouard, S.; Devos, D.; Kobeissy, F.; Zibara, K.; Salzet, M.; Fournier, I., Matrix-Assisted Laser Desorption/Ionization-Mass Spectrometry Imaging of Lipids in Experimental Model of Traumatic Brain Injury Detecting Acylcarnitines as Injury Related Markers. *Anal Chem* **2019,** *91* (18), 11879-11887.

16.   Gier, E. C.; Pulliam, A. N.; Gaul, D. A.; Moore, S. G.; LaPlaca, M. C.; Fernandez, F. M., Lipidome Alterations following Mild Traumatic Brain Injury in the Rat. *Metabolites* **2022,** *12* (2).

17.   Ojo, J. O.; Algamal, M.; Leary, P.; Abdullah, L.; Mouzon, B.; Evans, J. E.; Mullan, M.; Crawford, F., Converging and Differential Brain Phospholipid Dysregulation in the Pathogenesis of Repetitive Mild Traumatic Brain Injury and Alzheimer's Disease. *Front Neurosci* **2019,** *13*, 103.

18.   Paxinos, G. W., C., *The Rat Brain in Sterotaxic Coordinates*. 6th ed.; Elsevier: 2006.

19.   Pang, Z.; Chong, J.; Zhou, G.; de Lima Morais, D. A.; Chang, L.; Barrette, M.; Gauthier, C.; Jacques, P. E.; Li, S.; Xia, J., MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional insights. *Nucleic Acids Res* **2021,** *49* (W1), W388-W396.

20.   Valerie B. O'Donnell, E. A. D., Michael J. O. Wakelam, Shankar Subramaniam4, LIPID MAPS: Serving the next generation of lipid researchers with tools, resources, data, and training. *Sci Signal* **2019,** *12* (563).

21.   Sah, S.; Ma, X.; Botros, A.; Gaul, D. A.; Yun, S. R.; Park, E. Y.; Kim, O.; Moore, S. G.; Kim, J.; Fernandez, F. M., Space- and Time-Resolved Metabolomics of a High-Grade Serous Ovarian Cancer Mouse Model. *Cancers (Basel)* **2022,** *14* (9).

22.   Grin'kina, N. M.; Li, Y.; Haber, M.; Sangobowale, M.; Nikulina, E.; Le'Pre, C.; El Sehamy, A. M.; Dugue, R.; Ho, J. S.; Bergold, P. J., Righting Reflex Predicts Long-Term Histological and Behavioral Outcomes in a Closed Head Model of Traumatic Brain Injury. *PLoS One* **2016,** *11* (9), e0161053.

23.   Palomero-Gallagher, N.; Zilles, K., Cortical layers: Cyto-, myelo-, receptor- and synaptic architecture in human cortical areas. *NeuroImage* **2019,** *197*, 716-741.

24.   Carron, S. F.; Alwis, D. S.; Rajan, R., Traumatic Brain Injury and Neuronal Functionality Changes in Sensory Cortex. *Front Syst Neurosci* **2016,** *10*, 47.

25.   Frankowski, J. C.; Kim, Y. J.; Hunt, R. F., Selective vulnerability of hippocampal interneurons to graded traumatic brain injury. *Neurobiol Dis* **2019,** *129*, 208-216.

26.   Ibrahim, S.; Hu, W.; Wang, X.; Gao, X.; He, C.; Chen, J., Traumatic Brain Injury Causes Aberrant Migration of Adult-Born Neurons in the Hippocampus. *Sci Rep* **2016,** *6*, 21793.

27.   Girgis, F.; Pace, J.; Sweet, J.; Miller, J. P., Hippocampal Neurophysiologic Changes after Mild Traumatic Brain Injury and Potential Neuromodulation Treatment Approaches. *Front Syst Neurosci* **2016,** *10*, 8.

28.   Bramlett, H. M.; Dietrich, W. D., Quantitative structural changes in white and gray matter 1 year following traumatic brain injury in rats. *Acta Neuropathol* **2002,** *103* (6), 607-14.

29.   Irene Cristofori, W. Z., Aileen Chau, Jeffrey Solomon, Frank Krueger, Jordan Grafman, White and gray matter contributions to executive function recovery after traumatic brain injury. *Neurology* **2015,** *84* (14).

30.   Anthony A. Mercadante, P. T., *Neuroanatomy, Gray Matter*. StatPearls Publishing: 2021.

31.   Laitinen, T.; Sierra, A.; Bolkvadze, T.; Pitkanen, A.; Grohn, O., Diffusion tensor imaging detects chronic microstructural changes in white and gray matter after traumatic brain injury in rat. *Front Neurosci* **2015,** *9*, 128.

32.   Bouix, S.; Pasternak, O.; Rathi, Y.; Pelavin, P. E.; Zafonte, R.; Shenton, M. E., Increased gray matter diffusion anisotropy in patients with persistent post-concussive symptoms following mild traumatic brain injury. *PLoS One* **2013,** *8* (6), e66205.

33.   Ling, J. M.; Klimaj, S.; Toulouse, T.; Mayer, A. R., A prospective study of gray matter abnormalities in mild traumatic brain injury. *Neurology* **2013,** *81* (24), 2121-7.

34.   McDonald, S.; Rushby, J. A.; Dalton, K. I.; Allen, S. K.; Parks, N., The role of abnormalities in the corpus callosum in social cognition deficits after Traumatic Brain Injury. *Soc Neurosci* **2018,** *13* (4), 471-479.

35.   Kim, D. S.; Choi, H. J.; Yang, J. S.; Cho, Y. J.; Kang, S. H., Radiologic Determination of Corpus Callosum Injury in Patients with Mild Traumatic Brain Injury and Associated Clinical Characteristics. *J Korean Neurosurg Soc* **2015,** *58* (2), 131-6.

36.   Hartnell, D.; Gillespie-Jones, K.; Ciornei, C.; Hollings, A.; Thomas, A.; Harrild, E.; Reinhardt, J.; Paterson, D. J.; Alwis, D.; Rajan, R.; Hackett, M. J., Characterization of Ionic and Lipid Gradients within Corpus Callosum White Matter after Diffuse Traumatic Brain Injury in the Rat. *ACS Chem Neurosci* **2020,** *11* (3), 248-257.

37.   Hankin, J. A.; Farias, S. E.; Barkley, R. M.; Heidenreich, K.; Frey, L. C.; Hamazaki, K.; Kim, H. Y.; Murphy, R. C., MALDI mass spectrometric imaging of lipids in rat brain injury models. *J Am Soc Mass Spectrom* **2011,** *22* (6), 1014-21.

38.   Lima, F. D.; Souza, M. A.; Furian, A. F.; Rambo, L. M.; Ribeiro, L. R.; Martignoni, F. V.; Hoffmann, M. S.; Fighera, M. R.; Royes, L. F.; Oliveira, M. S.; de Mello, C. F., Na+,K+-ATPase activity impairment after experimental traumatic brain injury: relationship to spatial learning deficits and oxidative stress. *Behav Brain Res* **2008,** *193* (2), 306-10.

39.   Astrid Sigel, H. S., Roland K.O. Sigel, The Alkali Metal Ions: Their Role for Life. In *The Alkali Metal Ions: Their Role for Life*, Springer: 2016; Vol. 16, pp 594-596.

40.   Sarkar, C.; Jones, J. W.; Hegdekar, N.; Thayer, J. A.; Kumar, A.; Faden, A. I.; Kane, M. A.; Lipinski, M. M., PLA2G4A/cPLA2-mediated lysosomal membrane damage leads to inhibition of autophagy and neurodegeneration after brain trauma. *Autophagy* **2020,** *16* (3), 466-485.

41.   Zheng, L.; Xie, C.; Zheng, J.; Dong, Q.; Si, T.; Zhang, J.; Hou, S. T., An imbalanced ratio between PC(16:0/16:0) and LPC(16:0) revealed by lipidomics supports the role of the Lands cycle in ischemic brain injury. *J Biol Chem* **2021,** *296*, 100151.

42.   Liu, P.; Zhu, W.; Chen, C.; Yan, B.; Zhu, L.; Chen, X.; Peng, C., The mechanisms of lysophosphatidylcholine in the development of diseases. *Life Sci* **2020,** *247*, 117443.

43.   Law, S. H.; Chan, M. L.; Marathe, G. K.; Parveen, F.; Chen, C. H.; Ke, L. Y., An Updated Review of Lysophosphatidylcholine Metabolism in Human Diseases. *Int J Mol Sci* **2019,** *20* (5).

44.   Wang, G.; Ding, L.; Gao, C.; Zhang, N.; Gan, D.; Sun, Y.; Xu, L.; Luo, Q.; Jiang, Z., Neuroprotective effect of l-serine against white matter demyelination by harnessing and modulating inflammation in mice. *Neuropharmacology* **2019,** *146*, 39-49.

45.   Tzekov, R.; Dawson, C.; Orlando, M.; Mouzon, B.; Reed, J.; Evans, J.; Crynen, G.; Mullan, M.; Crawford, F., Sub-Chronic Neuropathological and Biochemical Changes in Mouse Visual System after Repetitive Mild Traumatic Brain Injury. *PLoS One* **2016,** *11* (4), e0153608.

46.   Mavroudakis, L.; Stevens, S. L.; Duncan, K. D.; Stenzel-Poore, M. P.; Laskin, J.; Lanekoff, I., CpG preconditioning reduces accumulation of lysophosphatidylcholine in ischemic brain tissue after middle cerebral artery occlusion. *Anal Bioanal Chem* **2021,** *413* (10), 2735-2745.

47.   Palafox-Sanchez, V.; Ying, Z.; Royes, L. F. F.; Gomez-Pinilla, F., The interaction between brain and liver regulates lipid metabolism in the TBI pathology. *Biochim Biophys Acta Mol Basis Dis* **2021,** *1867* (4), 166078.

48.   Anthonymuthu, T. S.; Kenny, E. M.; Bayir, H., Therapies targeting lipid peroxidation in traumatic brain injury. *Brain Res* **2016,** *1640* (Pt A), 57-76.

49.   Hossain, M. S.; Mawatari, S.; Fujino, T., Plasmalogens, the Vinyl Ether-Linked Glycerophospholipids, Enhance Learning and Memory by Regulating Brain-Derived Neurotrophic Factor. *Front Cell Dev Biol* **2022,** *10*, 828282.

50.   Xu, Q.; Zhang, Y.; Zhang, X.; Liu, L.; Zhou, B.; Mo, R.; Li, Y.; Li, H.; Li, F.; Tao, Y.; Liu, Y.; Xue, C., Medium-chain triglycerides improved cognition and lipid metabolomics in mild to moderate

Alzheimer's disease patients with APOE4(-/-): A double-blind, randomized, placebo-controlled crossover trial. *Clin Nutr* **2020,** *39* (7), 2092-2105.

51.   Udagawa, J.; Hino, K., Plasmalogen in the brain: Effects on cognitive functions and behaviors attributable to its properties. *Brain Res Bull* **2022,** *188*, 197-202.

52.   Aureli, M.; Murdica, V.; Loberto, N.; Samarani, M.; Prinetti, A.; Bassi, R.; Sonnino, S., Exploring the link between ceramide and ionizing radiation. *Glycoconj J* **2014,** *31* (6-7), 449-59.

53.   Sergei A Novgorodov, T. I. G., Ceramide and mitochondria in ischemic brain injury. *Int J Biochem Mol Biol* **2011**.

54.   Alla Yu Frantova, A. E. S., Anatoly S. Bushnev, Elena N. Zvonkova, Vitaly I. Shvets, The synthesis of ceramide phosphoinositol. *Tetrahedron Letters* **1992,** *33* (24), 3539-3542.

55.   Heiles, S.; Kompauer, M.; Muller, M. A.; Spengler, B., Atmospheric-Pressure MALDI Mass Spectrometry Imaging at 213 nm Laser Wavelength. *J Am Soc Mass Spectrom* **2020,** *31* (2), 326-335.