Title

Bending the course of evolution: how mutualistic interactions affect macroevolutionary

dynamics of diversification in mimetic butterflies.

Authors

Chazot N. [1*], Pires Braga M. [1], Aubier T.G. [2], Llaurens V. [3,4], Willmott K. R. [5], Elias M [3,6].

[1] Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden.

[2] Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), Université de

Toulouse, CNRS, IRD, Toulouse INP, Université Toulouse 3 – Paul Sabatier (UT3),

Toulouse, France.

[3] Institut de Systématique, Evolution, Biodiversité, MNHN-CNRS-Sorbonne Université-

EPHE-Université des Antilles, Muséum national d'Histoire naturelle, 45 Rue Buffon, 75005,

Paris, France.

[4] Centre Interdisciplinaire de Recherche en Biologie, CNRS, INSERM, Collège de France, 11,

place Marcelin Berthelot, 75005 Paris, France.

[5] McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History,

University of Florida, Gainesville, FL, United States of America; and Instituto Nacional de

Biodiversidad, Quito, Ecuador (research associate).

[6] Smithsonian Tropical Research Institute, Panamá, Panamá.

Email addresses

Chazot N: nicolas.chazot@slu.se *

Pires Braga M: mpiresbr@gmail.com

Aubier TG: thomas.aubier@normalesup.org

Llaurens V: violaine.llaurens@mnhn.fr

Willmott K: kwillmott@flmnh.ufl.edu

Elias M: marianne.elias@mnhn.fr

* Corresponding author

Short running title

Macroevolutionary signature of mimicry

Key-words

mutualism, macroevolution, Müllerian mimicry, time-dependent diversification, butterflies,

color patterns.

Corresponding author

Dr Nicolas Chazot

Department of Ecology

Swedish University of Agricultural Sciences

Ulls väg, 16

75651 Uppsala

Sweden

Abstract

Evidence that species interactions can affect macroevolutionary dynamics of trait and species

diversification is scarce. Mutualistic Müllerian mimicry is a compelling example of example

of ecological interactions that has been shown to drive evolutionary convergence,

Here, we test how mutualistic Müllerian mimicry shapes macroevolutionary patterns of

diversification in the Ithomiini butterflies. We show that the age of color patterns is the

primary determinant of species richness within mimicry rings but not phylogenetic diversity.

We find pervasive phylogenetic signal in mimicry rings and in color patterns associated

within polymorphic species. Only a small set of mimicry rings show high phylogenetic

diversity. We identify patterns of saturation in the accumulation of new mimicry rings and in

the number of evolutionary convergences towards the most species-rich mimicry rings. Those

saturation patterns are likely caused by niche filling along various ecological dimensions,

within and among the mimetic communities living in sympatry. The time-dependent effects

detected in our study illustrate how neutral processes and ecological interactions interact and

shape species and phenotypic diversification.

Introduction

Identifying how microevolutionary processes and ecological interactions occurring at the community level can affect macroevolutionary dynamics has been a major focus of evolutionary biology (Reznick & Ricklef, 2009, Rolland et al. 2023). Despite a long interest, evidence for an effect of ecological interactions on macroevolutionary patterns of species diversification is scarce. Limitations lie primarily in the difficulty of testing such effects, in particular because species interactions are usually difficult to characterize, and because predicting the effects of processes happening at small spatial and temporal scale on macroevolutionary dynamics is challenging (Rolland et al. 2023). Different ecological interactions (e.g. competition, parasitism, mutualism) may indeed underlie drastically different macroevolutionary trends and involve one or several guilds of species, therefore raising questions about the relative effects of phylogenetic constraints *vs.* selection stemming from these ecological interactions.

Here, we focus on Müllerian mimicry, a widespread and well-documented mutualistic interaction that offers a remarkable opportunity to study the feedbacks between ecological interactions and macroevolutionary processes (Kunte et al. 2021). The evolutionary convergence of warning signals shared by co-occurring defended species, *i.e.* Müllerian mimicry, is one of the most compelling examples of ecological interactions driving trait evolution in multiple distantly related species. Müllerian mimicry among co-occurring species arises through positive frequency-dependent selection exerted by predators that more quickly learn to avoid locally abundant warning signals (Müller 1879, Sherratt 2006). Because the cost of predator learning is diluted among individuals of all the species that share the same warning signal, Müllerian mimicry is a mutualistic interaction; individuals from a given species benefit from co-occurring with individuals belonging to co-mimetic species, thereby

92   promoting wing color pattern convergences in distantly related species. Species that share a

93   common warning color pattern within communities are referred to as mimicry rings.

94   Müllerian mimicry has evolved independently multiple times in animals, including,

95   among others, frogs (e. g., Symula et al. 2001), snakes (e. g., Sanders et al. 2006), catfish

96   (Alexandrou et al. 2011), butterflies and moths (e. g., Beccaloni 1997), bees and wasps (e. g.,

97   Wilson et al. 2012, Ezray et al. 2019) and beetles (e. g., Muñoz-Ramírez et al. 2016). The

98   phenomenon is particularly well-studied in butterflies, in particular in the species-rich

99   neotropical tribe Ithomiini (396 species).

100   Numerous theoretical, experimental, and field studies have shown that predation is a

strong selective force driving the convergent evolution of warning signals such as color

patterns (e. g., Sherratt 2006, Elias et al. 2008, Gompert et al. 2011, Willmott et al. 2017,

Birskis-Barros et al. 2021). The strength of frequency-dependent selection resulting from

predation has been convincingly shown empirically in *Heliconius* (e. g., Mallet and Barton

1989, Kapan 2001, Chouteau et al. 2016) and ithomiine (Willmott et al. 2017) butterflies.

Importantly, this selection driven by ecological interactions has led to striking convergence

among sympatric species belonging to phylogenetically distant families, such as erebid moths

and nymphalid butterflies (Beccaloni 1997). Because mimicry involves strong resemblance

across phylogenetically distant species, it provides an excellent context to test the relative

importance of ecological interactions and phylogenetic constraints on macroevolutionary

dynamics.

112   Several lines of evidence show that mimicry drives the evolution of multidimensional

ecological niches in the species involved, including convergence in multiple traits (e.g. host-

plant preference, flight height and flight behaviour, wing morphology, altitudinal and climatic

niche) (Willmott and Mallet 2003, Elias et al. 2008, Hill 2010, Gompert et al. 2011, Chazot et

al. 2014, Aubier and Elias 2020, Hill 2021, Doré et al. 2023, Page et al. 2024). Mimicry is

thus an ecological interaction that plays a key role in the community assembly across multiple

sympatric species with various levels of phylogenetic divergence.

In addition to structuring communities, switches in mimetic patterns may act as a

driver of speciation in mimetic lineages, since differences in wing patterns among mimetic

species drive (Jiggins et al. 2001, Chamberlain et al. 2009, Merrill et al. 2011), and non-

mimetic hybrids between taxa belonging to distinct mimicry rings suffer stronger predation

(Merrill et al, 2012). Mimetic color patterns are thus often considered as so-called magic traits

(*i. e.*, traits involved in mate choice that are also under divergent ecological selection,

Gavrilets 2004, Servedio et al. 2011), and as one factor contributing to high diversification

rates in some clades of mimetic butterflies (Jiggins et al. 2006, Kozak et al. 2015). Mimicry

is also predicted to fuel species diversification through the formation of a spatial mosaic of

mimicry rings (Aubier et al. 2017), because such a mosaic should allow the persistence of

different species occupying similar ecological niches in different patches of the mosaic.

Here, we investigate whether mimetic interactions affect the macroevolutionary

dynamics of diversification of mimetic species and their warning signals, as well as the

species composition of mimicry rings. We specifically focus on: (1) the diversification

dynamics of colour patterns through time and (2) the consequences of such phenotypic

diversification on the variation in species richness and phylogenetic diversity among mimicry

rings. In order to address these two questions in the absence of a clearly established

framework of hypotheses we propose nine predictions to be tested (Figure 1). We used the

largest tribe of mimetic butterflies, namely the Ithominii, by combining phylogenetic

information with a dataset of color patterns for 339 Ithomiini species (Chazot et al. 2019,

Doré et al. 2021, 2022).

MATERIAL AND METHODS

Data

We used the time-calibrated phylogeny of Ithomiini from Chazot et al. (2019), which contains

339 species of Ithomiini (85% of species). We used Doré et al (2021)'s color pattern

classification and species data, resulting in 44 mimicry rings (Figure 2, Table S1). Mimicry

ring species richness in Ithomiini ranges from a single to over 70 species, and, due to local or

geographic wing pattern polymorphism, at least half of Ithomiini species belong to two or

more mimicry rings.

Ancestral state estimations

In our dataset, 200 out of the 339 species studied show color pattern polymorphism. We

therefore estimated ancestral color patterns across the Ithomiini tree using the method

described in Braga et al. (2020). This method, initially designed for estimating ancestral host-

parasite associations on a phylogenetic tree, allows current and ancestral species to belong to

multiple mimicry rings, due to wing color pattern polymorphism. We used the 2-state model

and the inference strategy of Braga et al. (2020) as implemented in RevBayes (Höhna et al.,

2016) to infer the history of color pattern evolution, based on the color patterns known for

each extant species. Model parameters and ancestral states were inferred using Markov Chain

Monte Carlo (MCMC). We ran two chains for 8×10^4 iterations, discarding the first 10% as

burn-in and sampling every 50 iterations, then confirmed that the chains converged to the

same posterior distribution by calculating the Gelman diagnostic (Gelman and Rubin, 1992)

in the R package *coda* (Plummer et al., 2006). Results from the MCMC chain with the highest

effective sample size are presented. All RevBayes and R scripts are available at

github.com/maribraga/Ithomiini_color.

Because polymorphic states can be inferred as ancestral states, the complexity of the

evolutionary history estimated greatly increased. We thus took advantage of the Bayesian

framework by performing all follow-up analyses on a posterior distribution of evolutionary

histories.

Dynamics of color pattern evolution through time

Prediction 1: The rate of emergence of new patterns decreases through time, because the

selective advantage of displaying an already prevalent color pattern should favor convergence

rather than innovation (Figure 1).

Prediction 2: The rate of color pattern shifts increases through time as the overall number of

color patterns rises, because an increasing number of color patterns provides new

opportunities for convergence (Figure 1).

To test these predictions, we extracted the timing of color pattern evolutionary events

(*i.e.*, gains and loss) for each independent history sampled from the posterior distribution of

the ancestral state estimations. We extracted the earliest timing of emergence of each color

pattern to calculate the cumulative sum of new color patterns through time within 1 My time

bins. Using a sliding-window analysis, we also counted the total number of color pattern

shifts (irrespective of color pattern identity) occurring within 1 My time bins to estimate the

number of shifts through time. Since both statistics are expected to increase proportionally

with branch length, we obtained the rate of new pattern emergence (Prediction 1) and the rate

of color pattern shift (Prediction 2) for each 1 My time-bin by dividing the number of new

patterns and the number of shifts, respectively, by the sum of branch lengths in that bin.

Prediction 3: There is a significant phylogenetic signal in color pattern associations within

species, caused by phylogenetically conserved developmental pathways of color patterns

and/or parallel co-evolution of color patterns across species, since polymorphic species can

simultaneously shift to different mimicry rings.

We evaluated whether polymorphic species exhibit random color pattern associations

or instead, whether they represent repeated association of the same color patterns. We used

statistical tools designed for ecological network analyses to test for modularity in the network

of color pattern associations, where modules represent clusters of color patterns frequently

associated in polymorphic species. We used the Beckett (2016) algorithm to identify the

network configuration maximizing the proportion of interactions within modules. We

repeated the module identification 100 times and calculated the frequency of pairwise

associations of color patterns within modules. We used the package *qgraph* to build the

network of pairwise associations.

We then tested whether the species composition of each module was phylogenetically

random or instead phylogenetically structured. We measured phylogenetic diversity in the

modules by comparing the mean phylogenetic distance (MPD) between species belonging to

the same module to the distribution of mean phylogenetic distances after permutations of

color patterns between modules. High MPD score indicates a module comprising more

distantly related species than expected by random community assemblage, *i.e.* resulting from

color pattern convergence between distantly related lineages. Conversely, a low MPD

indicates a module shared by species more closely related than a random combination of

species. MPD values falling into the 95% interval of the distribution obtained from the

permutations indicate no departure from random evolution of color pattern, *i.e.* driven neither

by convergence events nor by phylogenetic proximity. Since modules are not identical in

different iterations, we cannot use the entire posterior distribution for this test. We performed

the MPD test on one iteration of the module search algorithm.

Prediction 4: Phylogenetic signal of color patterns decreases with the age of the color pattern (Figure 1). Positive frequency-dependent selection on color patterns is expected to drive convergence among distantly related lineages, thereby decreasing phylogenetic signal. Greater time since the emergence of a color pattern allows for color pattern convergence involving more distantly related species (Kunte et al. 2021).

To assess the phylogenetic signal of color patterns, we computed the D statistics (Fritz & Purvis 2010) for each mimicry ring (while merging all other mimicry rings) with more than 3 species, with permutations to identify significantly conserved and overdispersed mimicry rings. A D value of 0 indicates Brownian motion evolution across the tree, and a value of 1 random evolution uncorrelated with phylogeny. D<0 indicates phylogenetic clustering (i. e., conservatism) while D>1 indicates phylogenetic overdispersion.

We obtained the age of color patterns by extracting the timing of color pattern gain and loss events for each independent history sampled from the posterior distribution of the ancestral state estimation analysis. The oldest timing of emergence of a color pattern, in absolute value, was considered as the age of the color pattern. To test our prediction, we fitted a linear model with D as response variable, and color pattern age as explanatory variable.

Species richness and phylogenetic diversity within mimicry rings

In Ithomiini butterflies there is a huge disparity in mimicry ring species richness (i. e., the number of species in a mimicry ring), which ranges from a single species to over 70 (Figure 2). We estimated mimicry ring phylogenetic diversity using an MPD test similar to the tests performed within modules. We compared the mean phylogenetic distance between species belonging to the same mimicry ring to the distribution of mean phylogenetic distances after permutations. Five color patterns were unique to a single species and therefore we did not perform the MPD test for the mimicry rings harboring these color patterns. MPD also strongly

varied across mimicry rings, from phylogenetically overdispersed to highly clustered mimicry

rings. A linear regression confirmed that MPD does not predict species richness (F= 2.20,

Df=37 , p=0.146 , R2=0.03, Supplementary Figure S2A). To explain such disparity in both

species richness and phylogenetic diversity, we estimated the contribution of four processes

by which variation in species richness or phylogenetic diversity between mimicry rings may

arise (Figure 1).

Prediction 5: Species richness and phylogenetic diversity of a mimicry ring increase with the

age of the associated color pattern (Figure 1). As a null hypothesis, species richness should

increase with time through the accumulation of either speciation without color pattern change

or speciation driven by convergence of color pattern. Greater time since the emergence of a

color pattern allows for color pattern convergence involving more distantly related species,

hence increasing phylogenetic diversity (Kunte et al. 2021).

Prediction 6: *Species richness and phylogenetic diversity increase with the number of*

*evolutionary convergence (*Figure 1). Differences in evolutionary gains among color patterns

may be observed, either (1) because they confer different advantages such as a higher

protection against predation (e. g., because they are locally abundant, or more memorable) or

benefits unrelated to aposematism (e.g., thermoregulation, crypsis), or (2) because fewer

mutations are required for this pattern to emerge from an ancestral one, for example when the

capacity to produce certain pigments and pattern elements is already present as an ancestral

state.

We summed the total number of species gain events for each mimicry ring across the

posterior distribution of ancestral state scenarios, similar to the calculation of the rate of shifts

through time. However, even under stochastic evolutionary processes, the number of gains

increases with time. Thus, we divided the gains by the age of the color pattern to obtain a rate.

Prediction 7: *Species richness decreases with the number of evolutionary losses of color*

patterns, while phylogenetic diversity should not be affected if losses are randomly distributed

across lineages (Figure 1). Here, evolutionary losses do not imply lineage extinction but

rather evolutionary shifts in color patterns leading to the loss of an ancestral color pattern. We

calculated a rate of color pattern loss in the same way we calculated the rate of gain.

Prediction 8: *Species richness and phylogenetic diversity of a mimicry ring increase if*

lineages belonging to that mimicry ring exhibit a higher rate of color pattern evolution

(Figure 1). Such lineages may lead to a high number of convergence events towards the same

color pattern, thereby increasing species richness and phylogenetic diversity via convergence

and subsequent diversification. While Prediction 6 focuses on tree-wide association between

rates of gains and losses, Prediction 8 tries to account for clade-specific rates of evolution that

may lead to higher species richness.

281   We estimated a tip rate of color pattern evolution – species-specific rate of evolution –

by first calculating the number of shifts occurring along each branch in the tree across the

posterior distribution of ancestral state estimations. The length of each branch of the

phylogenetic tree was then rescaled to this number of events. We calculated for every tip in

the tree the total rescaled branch length connecting the root to the tips. Because the number of

possible evolutionary changes is expected to increase with the number of nodes connecting

the root and the tips, we divided the sum of branch length of each tip by the number of

connecting nodes to obtain the tip rate of color pattern evolution. Finally, we calculated the

median tip rate of evolution for each color pattern.

Prediction 9: Species richness of a mimicry ring increases, but phylogenetic diversity

decreases, with the rate of diversification of the lineages belonging to it (Figure 1). Although

the reasons why a given color pattern could directly affect the probability of a speciation

event (without color pattern shift) are unclear, it may be indirectly associated with higher

diversification through correlation with other phenotypical, ecological or geographical traits

e.g., habitats that favor speciation or limit extinction, such as high-altitude Andean forests

(Chazot et al. 2016, 2018, 2019). Higher speciation rate within a mimicry ring will lead to a

larger number of species harboring the same color pattern, thereby increasing the species

richness of the mimicry ring, while decreasing its mean phylogenetic diversity because most

species joining that mimicry ring will be closely related.

To test Prediction 9, we calculated the Diversification Rate (DR) statistics (Jetz et al.,

2012) for each tip in the tree. DR is a tip rate of diversification calculated as the inverse of the

equal-split measure (Redding & Mooers 2006). We calculated the median DR for each

mimicry ring from all the species belonging to the same mimicry ring. We used MPD of

mimicry rings as before.

Statistical analyses – We tested for relationships between species richness or phylogenetic

diversity and each explanatory variable using linear regressions. However, the age of color

patterns correlated with many of our explanatory variables. In particular, both species richness

and the rate of shifts increased with age. Therefore, we also calculated the residuals of the

linear model between species richness and color pattern age, then repeated the statistical

analyses using those residuals as the dependent variable. We found no relationship between

MPD and color pattern age and thus did not perform the test for residual MPD. All statistical

tests were performed using the median predictor scores calculated over 100 posterior

evolutionary histories.

RESULTS

Color pattern evolution through time

The number of color pattern shifts increased over time, but the rate of shifts remained nearly

stable (Figure 3A and 3C), providing no support for our prediction that an increasing number

of colour patterns fuels colour pattern shifts (Prediction 2). The rate of new pattern emergence

decreases through time, moderately until about 15 My ago, and more abruptly after that, with

hardly any new pattern appearing during the last 5 My (Figure 3B and 3D), consistent with

saturation of colour patterns (in line with Prediction 1).

Patterns of species richness and modularity

There was high heterogeneity in extant species richness among mimicry rings, with a mean of

16.3 species per mimicry ring, ranging from 74 species sharing the same warning signal

(color pattern AGNOSIA*)*, to a single species (ACRISIONE, DERCYLLIDAS,

HUMBOLDT, PRAESTANS, VESTILLA) (Figure 2). We found modularity in the network

of interactions, indicating that certain combinations of color patterns within polymorphic

species occur more often than expected at random. A single iteration of the Beckett (2016)

algorithm identified nine modules (Figure 4A). The most species-rich mimicry rings were not

concentrated in a single module. For example, the six most species-rich mimicry rings were

distributed across four different modules. We found significantly low Mean Phylogenetic

Distance (MPD) scores for four of these modules (module 2, 3, 4, 9, Figure 4A). A single

module had significantly high MPD (module 6). This module contained most of the patterns

with black, orange and yellow opaque colors (often referred to as 'tiger' color patterns). The

generally low MPD score suggests that the association of different colour patterns within

species partially stems from phylogenetic proximity between species, therefore providing

partial support for Prediction 3. Nevertheless, not all modules were supported based on 100

iterations of the Beckett algorithm (Figure 5). Module 2, 4 and 6 were well supported and

module 3 and module 1 only partly supported (Figure 5). This analysis of the strength of color

patterns associations also clearly highlights a few color patterns generally not involved in

polymorphism, such as LIBETHRIS and DUILLIA. By contrast, DOTO appears to be

involved in many different iterations of polymorphism.

Phylogenetic signal within mimicry rings

Using the D statistics, we found significant phylogenetic signal for 14 color patterns

and overdispersion for 12 mimicry rings (Figure 6A). The D statistics increased as color

patterns emerged more recently, indicating higher phylogenetic signal in older mimicry rings,

i.e. opposite to Prediction 4 (Figure 6A).

Predictors of mimicry ring species richness and phylogenetic diversity

Using the MPD test, we found significantly low mean phylogenetic diversity within 20

mimicry rings (Figure 4B). By contrast, only four mimicry rings showed significantly high

mean phylogenetic diversity (EXCELSA, HERMIAS, MAELUS, MAMERCUS, 'tiger' color

patterns with black, orange and yellow opaque colors).

We found that the age of color patterns (Prediction 5), the rate of color pattern losses

(Prediction 7) and the tip rate of evolution (Prediction 8) were the main predictors of mimicry

ring species richness (Figure 1, 6, Table 1), but the observed patterns for Predictions 5 and 7

are opposite to those expected (Figure 6D&E). In addition, the smallest mimicry rings tended

to harbor the lowest gain rates, but the relationship between the rate of gains and species

richness was weak ($p = 0.05$, Figure 6C, Table1, Supplementary Figure S2). Diversification

rate (Prediction 9) seemed higher in species-rich mimicry rings but the relationship was not

significant (Figure 6F, Table 1). Overall, species-rich mimicry rings were older, experienced

higher rates of color pattern loss and assemblages of lineages with high rates of evolution

(Figure 1). When performed on richness-age residuals, the rate of gains, losses and tip rate of

evolution significantly predicted species richness (Supplementary Figure S2).

When testing the same predictors for phylogenetic diversity we found that neither

color pattern age (Prediction 5) nor the rate of losses (Prediction 7) predicted MPD (Figure1,

6, Table 1, Supplementary Figure S2). However, we found that mimicry rings with higher

phylogenetic diversity had higher rates of gain (Prediction 6) and higher tip rate of color

pattern evolution (Prediction 8, Figure 1, 6, Table 1). We initially found no relationship

between MPD and DR. However, we identified one outlier in our analysis (mimicry ring

PAVONII shared by only two species). When we performed the regression test again without

PAVONII, we found that high MPD mimicry rings were associated with low DR (Prediction

9, Figure 6, Table 1).

DISCUSSION

Strong phylogenetic signal and convergences in the formation of mimicry rings

Despite selection by predation favoring convergence among distantly-related species, we

found significant phylogenetic signal in 14 mimicry rings. Modularity in the network of

mimetic interactions also indicates that there is a non-random co-occurrence of color patterns

in polymorphic species. Out of the nine detected modules, half were phylogenetically

clustered (Figure 4), although the interpretation of this result is limited by the poor support we

found for some of these modules. Furthermore, polymorphic states at nodes were generally

estimated from our analyses. This suggests that phylogenetic inheritance is not limited to

single color patterns but also frequently affects the ability to produce sets of different patterns,

maybe generated by similar developmental pathways. It is also possible that ecological

similarities between co-mimetic species lead to repeated co-variation of color patterns when

these co-mimetic species co-occur over large geographical scale.

We predicted that phylogenetic signal should decrease with color pattern age

(Prediction 4, Kunte et al. 2021) but we found the opposite result (Figure 1, 6). Even though

the number of convergence events also increases with time (in line with Prediction 4), it is

probably outweighed by speciation without color pattern changes and constrained by the time-

dependent effects as discussed below.

Mean phylogenetic diversity was significantly higher than expected under a random

distribution of the color patterns within four mimicry rings (EXCELSA, HERMIAS,

MAELUS, MAMERCUS). All of them but EXCELSA are part of the same module, the only

module also phylogenetically overdispersed (Figure 4). While wing transparency is very

common in Ithomiini, with most of the mimicry rings having wings with transparent patches,

the four highly convergent color patterns mentioned above are all non-transparent and are

frequently found outside of Ithomiini, such as in heliconiine butterflies (Nymphalidae:

Heliconiini), danaines (Nymphalidae: Danainae: Danaini) or in pericopine tiger-moths

(Erebidae: Pericopina). The elevated number of distantly-related species displaying these

color patterns, both within and outside Ithomiini butterflies, may stem from several

developmental or selective factors: 1- the retention of genetic abilities that could be ancestral

to all Lepidoptera (phylogenetic signal hypothesis), 2- the fact that these patterns can be

produced by a variety of genetic toolkits, thus making them more likely to appear

independently (evolutionary convergence hypothesis), 3 – opaque patterns being more

conspicuous and therefore more easily learnt and remembered by predators (natural selection

hypothesis).

Color pattern evolution through time: evidence for ecological saturation?

We initially predicted that the frequency of color pattern shifts would increase through time as

a result of the increasing number of possibilities for convergence provided by the increasing

number of color patterns with time (Prediction 2, Figure 1). Yet, across all color patterns, we

found hardly any variation in the dynamics of shifts through time (when accounting for the

increasing number of lineages through time, Figure 3). Therefore, the increasing complexity

of the network of interactions over time did not result in an increase in the overall rate of

evolutionary shifts. The fact that the overall rate of shifts remained stable while the number of

patterns increased implies that the rate of gains and losses per pattern decreased through time.

In parallel, we found a strong dampening in the rate of appearance of new color

patterns, with very few new patterns appearing over the last 10 My, suggesting the existence

of a large-scale upper limit to the number of color patterns. Such dampening could stem from

a mimicry saturation effect. While novel color patterns tend to be eliminated by positive

frequency-dependent selection, they may still become established due to drift followed by

selection (shifting balance, Mallet and Joron 1999). Evolution of such novel color patterns is

especially favored when selection from predation is relaxed (Chouteau and Angers 2012), as

may be the case in habitats devoid of an established mimetic community (Sherratt 2006).

Therefore, mimicry diversity can increase as long as there are empty ecological niches (e. g.,

new geographical regions, new elevations, new microhabitats), with communities of naive

predators, to colonize (Aubier et al. 2017). When all niches have been colonized by mimetic

butterflies, even if there is still room to accommodate more species because resources are not

yet limiting, we can expect a strong filtering on color patterns, which are under selection to

match the local mimetic community (Kunte et al. 2021, Aubier et al. 2017, Sherratt 2006,

Mallet & Barton 1989). In Ithomiini, all geographical regions and likely most altitudinal

niches were already colonized five million years ago, but the number of ithomiine species at

that time was only about 25% of the number of extant species (Chazot et al 2019). Such

conditions likely provided opportunities for further species accumulation in those different

regions and habitats, either through speciation or colonization (Chazot et al. 2016, 2018).

Nevertheless, mimetic communities may have been well established already, thereby favoring

the establishment of species with an already existing color pattern locally. As species

continued to be formed, novel color patterns were likely selected against.

We also found evidence of saturation in the balance between the rates of color pattern

gains and losses (Supplementary Figure S2). The rate of losses increased with the age of

mimicry rings and in species-rich mimicry rings (opposite to our Prediction 6). However,

while the rate of losses increased linearly with species richness and equaled the rate of gains

for recent/species-poor mimicry rings, the rate of gains in the oldest/species-rich mimicry

rings were disproportionately low (Supplementary Figure S2). Overall, for mimicry ring

species richness to increase over time despite a high rate of evolutionary losses, the rate of

speciation without loss of ancestral color pattern must exceed the rate of color pattern loss.

This result further supports the retention of color patterns over evolutionary time. It is also

consistent with the significant decrease of tip rate of color pattern evolution in species-rich

mimicry rings (in a direction opposite to our Prediction 7). The rate of color pattern evolution

declined in lineages belonging to species-rich mimicry rings, and the decline in the rate of

gains highlighted above is probably responsible for that.

The dampening of the rate of evolutionary gains is also a plausible explanation for the

absence of a relationship between MPD and color pattern age. We predicted that MPD should

increase with color pattern age via the accumulation of convergence events. We indeed found

that the rate of evolutionary gains predicts MPD. But the time-dependency of this rate of

evolution probably constrains the accumulation of phylogenetic diversity as mimicry rings

accumulate species.

467   Altogether, our results point at a saturation effect within mimicry rings, which could

be due to resource limitation; co-mimetic species tend to share habitats (Chazot et al. 2014,

Doré et al. 2023), microhabitats (Elias et al. 2008, Willmott et al. 2017) and even larval

hostplants (Wilmott and Mallet 2003). After a mimicry ring becomes too species-rich the cost

of competition among co-mimics may outweigh the anti-predator benefits of joining a very

large mimicry ring.

The age of mimicry rings predicts species richness but not phylogenetic diversity

In our results, time emerged as a major factor determining mimicry ring species richness,

which altogether generates a saturation effect limiting new evolutionary convergence. We

found that mean phylogenetic diversity of mimicry rings is determined by different processes

than species richness. In agreement with our predictions, mean phylogenetic diversity was

high in mimicry rings where the color pattern is frequently acquired and whose lineages have

high rates of color pattern evolution. But phylogenetic diversity of a mimicry ring results from

the balance between diversification without color pattern shift and the rate of evolutionary

convergence and we found that MPD decreased with high diversification rates. This result

further supports the relative importance of color pattern retention during diversification. Even

though evolutionary shifts increased over time and MPD increased with the number of

evolutionary shifts, color pattern age did not predict MPD. This is likely another side effect of

the dampening of convergence events in species rich mimicry rings that indirectly bound

phylogenetic diversity in species rich mimicry rings.

Conclusion

Mimicry is a remarkable example of an interaction evolving through natural selection. How

such selection translates into the dynamics of phenotypic evolution and species accumulation

at macroevolutionary scale has previously been poorly studied (but see Aubier et al. 2017).

Our results on Ithomiini support mechanisms of diversification that are independent from

mimicry. Time alone is a strong predictor of the number of evolutionary events and species

richness in mimicry rings. We also found evidence of phylogenetic signal during

diversification, which directly influences the relative importance of speciation without color

pattern change over convergence in determining mimicry ring composition. Therefore, the

network of interactions and the composition of mimicry rings appear to be driven in a large

part by processes not associated with mimicry.

Nevertheless, we find evidence that ecological saturation possibly linked to mimetic

interactions did affect the dynamics of color pattern evolution and species diversification.

This conclusion comes from several lines of evidence: the strong decrease in the rate of new

color pattern appearance, the decrease in color pattern gains relative to losses in species-rich

mimicry rings, and the increasing rate of losses in species-rich mimicry rings. Ecological

limitations to the overall diversity of color patterns and to the number of species interacting

within a mimicry ring seem likely to change the dynamics and directionality of color pattern

evolution, and thereby the species composition of mimicry rings.

Our results also have implications for the global biogeography of species interactions.

Large networks of mimetic interactions seem to be primarily found at lower latitudes

(although there has not been any formal validation of this observation). Our study shows that

time is key to building a network of mimetic interactions, both in terms of color pattern

diversification and species composition of mimicry rings. Thus, our results support the idea

that complex networks of interactions are built over long geological times, such that specific

conditions such as the age of biomes or their climatic stability (Fischer 1960) may be key to

understanding latitudinal gradients in biotic interactions.

References

Alexandrou, M. A., Oliveira, C., Maillard, M., McGill, R. A., Newton, J., Creer, S., and

Taylor, M. I. (2011). Competition and phylogeny determine community structure in Müllerian

co-mimics. *Nature*, *469*(7328), 84-88.

Aubier, T.G. and Elias, M. (2020), Positive and negative interactions jointly determine the

structure of Müllerian mimetic communities. Oikos, 129: 983-997.

Aubier, T. G., Elias, M., Llaurens, V., & Chazot, N. (2017). Mutualistic mimicry enhances

species diversification through spatial segregation and extension of the ecological niche

space. *Evolution*, *71*(4), 826-844.

Beccaloni, G. W. (1997). Ecology, natural history and behaviour of ithomiine butterflies and

their mimics in Ecuador (Lepidoptera: Nymphalidae: Ithomiinae). *Tropical Lepidoptera*

Research, 103-124.

Beccaloni, G. W. (1997). Vertical stratification of ithomiine butterfly (Nymphalidae:

Ithomiinae) mimicry complexes: the relationship between adult flight height and larval host-

plant height. *Biological journal of the Linnean Society*, *62*(3), 313-341.

Beckett, S. J. (2016) Improved community detection in weighted bipartite networks. *Royal*

Society Open Science, 3, 140536.

Birskis-Barros, I., Freitas, A. V., and Guimarães Jr, P. R. (2021). Habitat generalist species

constrain the diversity of mimicry rings in heterogeneous habitats. *Scientific Reports*, *11*(1),

5072.

Braga, M. P., M. J. Landis, S. Nylin, N. Janz, and Ronquist, F. (2020). Bayesian Inference of

Ancestral Host-parasite Interactions under a Phylogenetic Model of Host Repertoire

Evolution. *Systematic Biology,* 69:1149–1162.

Chamberlain, N. L., Hill, R. I., Kapan, D. D., Gilbert, L. E., and Kronforst, M. R. (2009).

Polymorphic butterfly reveals the missing link in ecological speciation. *Science*, *326*(5954),

847-850.

Chazot, N., Willmott, K.R., Lamas, G., Freitas, A.V., Piron-Prunier, F., Arias, C.F., Mallet, J.,

De-Silva, D.L. and Elias, M. (2019). Renewed diversification following Miocene landscape

turnover in a Neotropical butterfly radiation. *Global Ecology and Biogeography*, *28*(8),

pp.1118-1132.

Chazot, N., Willmott, K. R., Santacruz Endara, P. G., Toporov, A., Hill, R. I., Jiggins, C. D.,

and Elias, M. (2014). Mutualistic mimicry and filtering by altitude shape the structure of

Andean butterfly communities. *The American Naturalist*, *183*(1), 26-39.

Chazot, N., Willmott, K.R., Condamine, F.L., De‑Silva, D.L., Freitas, A.V., Lamas, G.,

Morlon, H., Giraldo, C.E., Jiggins, C.D., Joron, M. and Mallet, J., (2016). Into the Andes:

multiple independent colonizations drive montane diversity in the Neotropical clearwing

butterflies Godyridina. *Molecular ecology*, *25*(22), pp.5765-5784.

Chazot, N., De‑Silva, D.L., Willmott, K.R., Freitas, A.V., Lamas, G., Mallet, J., Giraldo,

C.E., Uribe, S. and Elias, M., (2018). Contrasting patterns of Andean diversification among

three diverse clades of Neotropical clearwing butterflies. *Ecology and evolution*, *8*(8),

pp.3965-3982.

Chouteau, M., Arias, M., and Joron, M. (2016). Warning signals are under positive frequency-

dependent selection in nature. *Proceedings of the national Academy of Sciences*, *113*(8),

2164-2169.

Chouteau, M., & Angers, B. (2012). Wright's shifting balance theory and the diversification

of aposematic signals. *PloS one*, *7*(3), e34028.

Doré, M., Willmott, K., Leroy, B., Chazot, N., Mallet, J., Freitas, A. V. L., Hall, J. P. W.,

Lamas, G., Dasmahapatra, K. K., Fontaine, C. and Elias, M. (2021) Ithomiini mimicry

classification. *Zenodo*, doi: 10.5281/zenodo.5777463

Doré, M., Willmott, K., Leroy, B., Chazot, N., Mallet, J., Freitas, A.V., Hall, J.P., Lamas, G.,

Dasmahapatra, K.K., Fontaine, C. and Elias, M. (2022). Anthropogenic pressures coincide

with Neotropical biodiversity hotspots in a flagship butterfly group. *Diversity and*

Distributions, *28*(12), pp.2912-2930.

Doré, M., Willmott, K., Lavergne, S., Chazot, N., Freitas, AVL., Fontaine, C, and, Elias, M.

(2023) Mutualistic interactions shape global spatial congruence and climatic niche evolution

in Neotropical mimetic butterflies. *Ecology Letters*, 26, 843-857

Elias, M., Gompert, Z., Jiggins, C., and Willmott, K. (2008). Mutualistic interactions drive

ecological niche convergence in a diverse butterfly community. *PLoS biology*, *6*(12), e300.

Ezray, B. D., Wham, D. C., Hill, C. E., and Hines, H. M. (2019). Unsupervised machine

learning reveals mimicry complexes in bumblebees occur along a perceptual

continuum. *Proceedings of the Royal Society B*, *286*(1910), 20191501.

Fischer, A. G. (1960). Latitudinal variations in organic diversity. *Evolution,* 14:64–81.

Fritz, S. A. and Purvis, A. (2010). Selectivity in mammalian extinction risk and threat types: a

new measure of phylogenetic signal strength in binary traits. *Conservation Biology*,

24(4):1042-1051.

Gelman, A. and Rubin, D.B. (1992) Inference from iterative simulation using multiple

sequences. *Statistical Science*, 7, 457– 472

Gavrilets, S. (2004) Fitness landscapes and the origin of species. Princeton University Press;

Princeton, New Jersey, USA.

Gompert, Z., Willmott, K., and Elias, M. (2011). Heterogeneity in predator micro-habitat use

and the maintenance of Müllerian mimetic diversity. *Journal of theoretical biology*, *281*(1),

39-46.

Hill, R. I. (2010). Habitat segregation among mimetic ithomiine butterflies

(Nymphalidae). *Evolutionary ecology*, *24*(2), 273-285.

Hill, R. I. (2021). Convergent flight morphology among Müllerian mimic

mutualists. *Evolution*, *75*(10), 2460-2479.

Höhna, S., Landis, M. J., Heath, T. A., Boussau, B., Lartillot, N., Moore, B. R., Huelsenbeck,

J. P., and Ronquist, F. (2016). RevBayes: Bayesian Phylogenetic Inference Using Graphical

Models and an Interactive Model-Specification Language. *Systematic Biology,* 65:726–736.

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., and Mooers, A. O. (2012). The global

diversity of birds in space and time. *Nature*, *491*(7424), 444-448.

Jiggins, C D., Mallarino, R., Willmott, K. R., Bermingham, E. (2006) The phylogenetic

pattern of speciation and wing pattern change in neotropical *Ithomia* butterflies (Lepidoptera:

Nymphalidae). *Evolution*. 60(7), 1454-66

Jiggins, C. D., Naisbit, R. E., Coe, R. L., and Mallet, J. (2001). Reproductive isolation caused

by colour pattern mimicry. *Nature*, *411*(6835), 302-305.

Kapan, D. Three-butterfly system provides a field test of müllerian mimicry . (2001).

Nature, 409, 338–340.

Kozak, K. M., Wahlberg N., Neild A. F. E., Dasmahapatra K. K., Mallet J and Jiggins C. D.

(2015). Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic

Heliconius Butterflies, *Systematic Biology*, 64(3), 505–524

Kunte, K., Kizhakke, A. G., and Nawge, V. (2021). Evolution of mimicry rings as a window

into community dynamics. *Annual Review of Ecology, Evolution, and Systematics*, *52*, 315-

341.

Mallet, J. L. and Barton, N. H. (1989) Strong natural selection in a warning-color hybrid

zone. *Evolution,* 43, 421–431.

Mallet, J. and Joron, M. (1999). Evolution of diversity in warning color and mimicry:

polymorphisms, shifting balance, and speciation. *Annual Review of Ecology and Systematics,*

30: 201-233

Mallet, J., and Willmott, K. (2003). Taxonomy: renaissance or Tower of Babel?. *Trends in*

Ecology & Evolution, *18*(2), 57-59.

Merrill, R. M., Van Schooten, B., Scott, J. A., and Jiggins, C. D. (2011). Pervasive genetic

associations between traits causing reproductive isolation in Heliconius

butterflies. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1705), 511-518.

Merrill, R. M., Wallbank, R. W., Bull, V., Salazar, P. C., Mallet, J., Stevens, M., and Jiggins,

C. D. (2012). Disruptive ecological selection on a mating cue. *Proceedings of the Royal*

Society B: Biological Sciences, *279*(1749), 4907-4913.

Muñoz-Ramírez, C. P., Bitton, P. P., Doucet, S. M., and Knowles, L. L. (2016). Mimics here

and there, but not everywhere: Müllerian mimicry in Ceroglossus ground beetles?. *Biology*

Letters, *12*(9), 20160429.

Page E., Queste, L. M., Rosser, N., Salazar, P. A., Nadeau, N. J., Mallet, J., Srygley, R. B.,

McMillan, W. O., Dasmahapatra, K. K. (2024). Pervasive mimicry in flight behavior among

aposematic butterflies. Proc*eedings of the National Academy of Sciences*,

121(11):e2300886121.

Plummer, M., Best, N., Cowles, K., and Vines, K. (2006). CODA: convergence diagnosis and

output analysis for MCMC. *R News*, 6, 7–11.

Redding, D. W. & Mooers, A. O. (2006) Incorporating evolutionary measures into

conservation prioritization. *Conservation Biology,* 20, 1670–1678

Reznick, D. N., and Ricklefs, R. E. (2009). Darwin's bridge between microevolution and

macroevolution. *Nature*, *457*(7231), 837-842.

Rolland, J., Henao-Diaz, L. F., Doebeli, M., Germain, R., Harmon, L. J., Knowles, L. L., ... &

Schluter, D. (2023). Conceptual and empirical bridges between micro-and macroevolution.

Nature Ecology & Evolution, 7(8), 1181-1193.

Sanders, K. L., Malhotra, A., and Thorpe, R. S. (2006). Evidence for a Mullerian mimetic

radiation in Asian pitvipers. *Proceedings of the Royal Society B-Biological Sciences,*

273(1590): 1135-1141.

Servedio, M. R., Van Doorn, G. S., Kopp, M., Frame, A. M., and Nosil, P. (2011). Magic

traits in speciation:'magic'but not rare?. *Trends in ecology & evolution*, *26*(8), 389-397.

Sherratt, T. N. (2006) Spatial mosaic formation through frequency-dependent selection in

Müllerian mimicry complexes. *Journal of Theoretical Biology,* 240:165–174

Symula, R., Schulte, R., and Summers, K. (2001). Molecular phylogenetic evidence for a

mimetic radiation in Peruvian poison frogs supports a Mullerian mimicry hypothesis.

Proceedings of the Royal Society B-Biological Sciences, 268(1484): 2415-2421.

Willmott, K. R., Robinson Willmott, J. C., Elias, M., and Jiggins, C. D. (2017). Maintaining

mimicry diversity: optimal warning colour patterns differ among microhabitats in Amazonian

clearwing butterflies. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1855),

20170744.

Wilson J. S., Williams K. A., Forister M. L., von Dohlen C. D., and Pitts J. P. (2012).

Repeated evolution in overlapping mimicry rings among North American velvet ants. *Nature*

Communications, 3.

Table 1. Results of the regression analyses performed between either mimicry species

richness or MPD and five predictors of diversity: age, rate of gain, rate of loss, tip rate of

color pattern evolution, diversification rate (DR).

	F	df	R2	p
SR ~ Age	118.8	42	0.73	<0.001
MPD ~ Age	2.61	37	0.04	0,114
SR ~ Gain	3.99	42	0.06	0,05
MPD ~ Gain	7.77	37	0.15	0,008
SR ~ Loss	303.5	42	0.88	<0.001
MPD ~ Loss	0.49	37	0.01	0,488
SR ~ TipRate	4.88	42	0.08	0,033
MPD ~ TipRate	13.46	37	0.25	<0.001
SR ~ DR	1.35	42	0.01	0,252
MPD ~ DR	0.89	37	<0.01	0,35
MPD ~ DR*	14.94	36	0.27	<0.001

712   * Regression analysis performed after removing the mimicry ring PAVONII.

716
717 Figure 1. Graphical summary of our nine predictions tested in this study and the results we

718 obtained for Ithomiini butterflies. Predictions 5-9 are subdivided into predictions for species

719 richness and MPD. Predictions are shown with dashed lines and results in plain lines.

Figure 2. A. Example of a history of color pattern evolution drawn from the posterior

distribution of the Bayesian inference. Only the 11 most species-rich mimicry rings are

colored. Pie charts do not represent posterior probabilities, rather, they show all mimicry rings

that were inferred at a given internal node at this given sample from the posterior distribution.

B. Ranked species richness of mimicry rings. C. Phylogenetic tree of Ithomiini with branch

lengths rescaled according to the average number of shifts estimated across 100 ancestral

histories drawn from the posterior distribution.

731 Figure 3. Number and rate of color pattern shifts through time. Cumulative number of new

732 color patterns through time and rate of new color pattern appearance through time. Grey

733 polygons show the 95% confidence interval from 100 posterior ancestral state estimations,

734 line showing the median.

737
738   Figure 4. Phylogenetic diversity within modules (A.) and within mimicry rings (B.). Circles

739   indicate the observed mean phylogenetic distance (MPD) score. Bars indicate the 95%

740   confidence interval of MPD scores after 1000 permutations. Butterfly images show the color

741   pattern associated with each mimicry ring or module.

743
744　Figure 5. Graphical representation of the frequencies of pairwise associations of color patterns

745　within polymorphic species calculated from 100 iterations of the Beckett (2016) algorithm.

746　Only pairwise associations with a frequency higher than 0.25 are shown. Color pattern names

747　are provided with each picture. Labels M1-M9 correspond to the module assignments as

748　provided in Figure 4.

Figure 6. Predictors of mimicry ring species richness and phylogenetic diversity. (A.)

Relationship between phylogenetic signal for each color pattern (D score) and the time of first

appearance. (B. - F.) Species and MPD as functions of color pattern age (B.), rate of gain (C.),

rate of loss (D.), tip rate of color pattern evolution (E.), diversification rate (F.). Black dots

correspond to species richness, gray dots correspond to MPD. The linear models fitted for

each predictor individually are shown when each variable significantly predicted either

mimicry species richness or MPD. Results of the regression analyses performed are provided

in Table 1.