Title: Induction of NASH in the *Nwd1*[−/−] mouse liver via SERCA2-dependent endoplasmic reticulum stress

Author: Seiya Yamada[1]*, Kazuhiko Nakadate[2], Tomoya Mizukoshi[1], Kiyoharu Kawakami[2], Ryosuke Kobayashi[3], Takuro Horii[3], Izuho Hatada[3,4], and Shin-ichi Sakakibara[1]*

Affiliations:

[1]Laboratory for Molecular Neurobiology, Faculty of Human Sciences, Waseda University, 2-579-15 Mikajima, Tokorozawa, Saitama 359-1192, Japan.

[2]Department of Basic Biology, Educational and Research Center for Pharmacy, Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose-shi, Tokyo, 204-8588, Japan.

[3]Laboratory of Genome Science, Biosignal Genome Resource Center, Institute for Molecular and Cellular Regulation, Gunma University, 3-39-15 Showa-Machi, Maebashi, Gunma, 371-8512, Japan

[4] Viral Vector Core, Gunma University Initiative for Advanced Research (GIAR), Gunma, 371-8511, Japan.

*To whom correspondence should be addressed: Shin-ichi Sakakibara or Seiya Yamada, Laboratory for Molecular Neurobiology, Faculty of Human Sciences, Waseda University, Tokorozawa, Saitama 359-1192, Japan; sakakiba@waseda.jp or seiya.s.yamada@gmail.com; tel. +81-4-2947-7120

Abstract

The endoplasmic reticulum (ER) stores Ca^{2+} and plays crucial roles in protein folding, lipid transfer, and it's perturbations trigger an ER stress. In the liver, chronic ER stress is involved in the pathogenesis of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH). Previous studies revealed that dysfunction of sarco/endoplasmic reticulum calcium ATPase (SERCA2), a key regulator of Ca^{2+} transport from the cytosol to the ER, is associated with the induction of ER stress and lipid droplet formation. We previously identified NACHT and WD repeat domain-containing protein 1 (Nwd1), which is localized in the ER and mitochondria. However, the physiological significance of Nwd1 outside the central nervous system remains unclear. In this study, we revealed that *Nwd1* knockout mice exhibited pathological manifestations comparable to NASH. Nwd1 interacts with SERCA2 near ER membranes. *Nwd1$^{-/-}$* livers exhibited reduced SERCA2 ATPase activity and a smaller Ca^{2+} pool in the ER, leading to an exacerbated state of ER stress. These findings highlight the importance of SERCA2 activity mediated by Nwd1 in the pathogenesis of NASH.

Highlights

・*Nwd1$^{-/-}$* mice exhibited NASH-like liver steatosis.
・Elevated ER stress, fibrosis, and pyroptosis were observed in *Nwd1$^{-/-}$* livers.
・Nwd1 interacts with SERCA2, an ER membrane $Ca2^+$ pump.
・*Nwd1$^{-/-}$* livers exhibited reduced SERCA2 activity and smaller Ca^{2+} pools in the ER.

Keywords

NASH, Nwd1, SERCA2, liver, lipid droplet, ER, ER stress, vacuole

Introduction

Nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH) are

liver diseases that progress without symptoms, and they are linked to global public

health problems. Characterized by the accumulation of lipid droplets in the liver, NASH

55   evolves from steatosis to inflammation and cellular damage, including cell death, often

culminating in fibrosis, cirrhosis, and eventually hepatocellular carcinoma [1]. The

etiology of hepatic lipid accumulation involves various factors, including the excessive

uptake or synthesis of fatty acids, defective transport of very low-density lipoprotein

(VLDL), and impaired beta-oxidation of fatty acids [2,3]. Furthermore, spatially

60   controlled hepatic zonation is crucial for physiological liver function, encompassing the

metabolism of various endogenous products and xenobiotics [4-6]. A comprehensive

understanding of the intricate interplay of cellular processes in NASH pathogenesis is

imperative for developing targeted therapeutic interventions. Although several clinical

trials of pharmacotherapies for NASH treatment are ongoing [1], no effective treatment

65   has been developed.

NAFLD and NASH are considered polygenic disorders entailing a diverse array

of genes, including unidentified genes. However, the individual contributions of these

genes to the pathogenesis of these diseases are unclear. One crucial aspect of NASH

pathology is the dysregulation of endoplasmic reticulum (ER) homeostasis [7,8]. The

70   ER stores Ca^{2+} and plays crucial roles in protein folding, lipid transfer, and organelle

dynamics regulation. The accumulation of unfolded or misfolded proteins in the ER

activates a series of homeostatic responses collectively termed ER stress, and chronic

ER stress contributes to the pathogenesis of NAFLD and NASH, including liver

steatosis and lipid droplet formation [9,10]. The ER-localized transmembrane protein

75 BSCL2/SEIPIN, acting at ER–lipid droplet contact sites, facilitates the incorporation of

protein and lipid cargo into growing lipid droplets and helps connect newly formed lipid

droplets to the ER and stabilize ER–lipid droplet contacts [11]. Recent studies

suggested that the abnormal regulation of Ca^{2+} transport proteins is the basis of

NAFLD/NASH [12,13]. The expression of type II inositol 1,4,5-trisphosphate receptor,

80 the principal Ca^{2+}-release channel in hepatocytes, was dramatically decreased in

biopsied livers from patients with steatosis and NASH [14]. Consistently, dysfunction

of sarco/endoplasmic reticulum calcium ATPase (SERCA2), a key regulator of Ca^{2+}

transport from the cytosol to the ER, induces ER stress and lipid droplet formation [15].

Conversely, SERCA2 overexpression in obese mice reduced lipid synthesis gene

85 expression and triglyceride content, emphasizing its significance in dysregulated lipid

homeostasis and suggesting that SERCA2 could be targeted to treat NASH [16].

However, the molecular mechanism by which SERCA2 acquires its proper function in

ER remains unclear.

Our prior research identified the NACHT and WD repeat domain-containing

90 protein 1 (Nwd1) gene, a novel member of the signal transduction ATPases with

numerous domains (STAND) protein superfamily [17,18]. Nwd1 is localized in the ER

and mitochondria in neural stem/progenitor cells (NSPCs), and it is essential for normal

brain development via interactions with the purine-synthesizing enzyme PAICS to

regulate purinosome assembly in NSPCs [17,18]. Nwd1 is a multi-domain protein; i.e.,

95 it has a NACHT domain in the central region that is predicted to have nucleoside

triphosphatase activity, an effector domain at the N-terminus that binds to PAICS, and a

cluster of WD40 repeats at the C-terminus that is involved in interactions with other

proteins or ligands. Based on its domain structure, Nwd1 is classified into the STAND

superfamily, which is known to be involved in the formation of large protein complexes

100   required for various signaling cascades such as apoptosis (apoptosome) and

inflammation (inflammasome) and is conserved across species, including zebrafish,

mice, rats, monkeys, and humans [17,19,20]. We previously demonstrated that Nwd1 is

predominantly expressed in the liver in addition to the brain [17,19,20]; however, the

physiological role of Nwd1 in nonneural tissues remains unknown.

105   In this study, we revealed that the livers of $Nwd1^{-/-}$ mice exhibit pathological

manifestations resembling NASH, implying that $Nwd1$ plays a role in modulating liver

homeostasis by regulating the Ca^{2+} pump function of SERCA2. These findings could

provide new clues to understanding the pathogenesis of ER stress-induced NASH.

110   **Materials and methods**

Animals

C57BL/6J mice purchased from Japan SLC, Inc. (Shizuoka, Japan) were housed under

temperature- and humidity-controlled conditions on a 12-h/12-h light/dark cycle with *ad*

libitum access to food and water. The day of birth was designated as P0.

Generation of Nwd1$^{-/-}$ mice

All protocols were approved by the committees on the ethics of animal experiments at

Gunma University and Waseda University. *Nwd1^{flox}* mice were generated via

electroporation, as previously reported with a partial modification [21]. Corresponding

120   to the target DNA sequences (Nwd1L3, AATGGGACAGTCACTTGGGA; Nwd1R1,

ACAGAACCTGGATTTTGTGG), donor single-stranded oligodeoxynucleotides

(ssODNs) with 5′- and 3′-homology arms flanking loxP and a restriction site were designed (Nwd1L3loxpAS, cacacacacaTGCATGTGCTCACACATCCTAACCTGCCATGTGGGCTTTTGCAAG

125 CCATCATAACTTCGTATAATGTATGCTATACGAAGTTATGGATCCCCAAGTG ACTGTCCCATTTGTTAAACTTCACAGTCTGTGTCCCTCCGAACATGTTTCCCA ; Nwd1R1loxP, GCCACATGCATGCTGCTTCCTTCCTTTTTTGTAGGGATTTGGACAGAACCTG GATTTTGTATAACTTCGTATAATGTATGCTATACGAAGTTATAAGCTTGGAG

130 GTTAGTTATGTGTTGTGTGATTGAGCCTGGGCCCCAGACCtgcatgtagcctttgt; Fig. 1A). Pre-annealed crRNA (Alt-R CRISPR–Cas9 crRNA, Integrated DNA Technologies [IDT], Coralville, IA, USA])/tracrRNA (Alt-R CRISPR–Cas9 tracrRNA, IDT; 3 μM), recombinant Cas9 protein (100 ng/μl; GeneArt Platinum Cas9 Nuclease, Thermo Fisher Scientific, Waltham, MA, USA), and ssODNs (400 ng/μl; Ultramer, IDT) in Opti-MEM

135 I (Thermo Fisher Scientific) were used for electroporation. First, a left loxP site was introduced into intron 4 of *Nwd1* by electroporation using B6D2F1-derived zygotes, and then the edited embryos were transferred to the oviduct of pseudopregnant ICR female to obtain *Nwd1* left loxPed males. Next, *Nwd1^flox^* mice were generated by introducing a right loxP site into intron 5 of *Nwd1* using left loxPed male-derived zygotes. Floxed

140 alleles were confirmed by sequencing PCR products using the following primer set: 5′-CAGGTGTGATATGTAAATACTTCTTTG-3′ (F2) and 5′-CAGGCCTAACAGAGCCAGAC-3′ (R2). We further converted the floxed allele to a null allele (*Nwd1^−^*) by crossing *Nwd1^flox^* heterozygous mice with CAG-Cre driver mice (B6.Cg-Tg[CAG-Cre]CZ-MO2Osb, RBRC01828; RIKEN BRC, Tokyo, Japan)

145 harboring the CAG-promoter-driven Cre recombinase transgene [22]. Exon 5, which

encodes the NACHT domain, was eliminated by Cre/loxP-mediated excision. The genotyping of floxed and null alleles was performed by genomic PCR using the following primer sets: 5′-TCACACATCCTAACCTGCCA-3′ (F1) and 5′-AGTAGGCCAAGCTCGATCTC-3′ (R1) for the floxed allele and F1 and R2 primers

150   for the null allele.

Cell culture and transfection

HEK293T and HeLa cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan) containing 10%

155   fetal bovine serum (Biowest, Nuaillé, France), 1% penicillin/streptomycin (FUJIFILM Wako Pure Chemical), and 1% L-glutamine (FUJIFILM Wako Pure Chemical). Cells were seeded into poly-D-lysine–coated dishes (Merck Millipore, Burlington, MA, USA) and transfected with plasmid DNA and PEI MAX (Polysciences, Warrington, PA, USA) complexes (DNA to PEI MAX ratio, 1:3, w/w) formed in Opti-MEM (Thermo

160   Fisher Scientific) via incubation for 15 min at 25°C. DNA complexes were added to cell cultures with Opti-MEM for 3 h, followed by cultivation with serum-containing DMEM.

Electron microscopy

165   Anesthetized adult mice were perfused transcardially with saline solution, followed by a fixative containing 2% paraformaldehyde (PFA) and 2% glutaraldehyde (GA) in 0.1 M phosphate buffer (PB), pH 7.4. Liver tissue was dissected, cut into small pieces, washed several times with PB, and postfixed with 1% OsO_4 (TAAB Laboratories, Aldermaston, UK) for 2 h. After dehydration with ethanol and replacement with propylene oxide, the

170   sections were embedded in Epon-812 resin (TAAB Laboratories) and thermally

polymerized. After trimming, the samples were cut into 70-nm-thick sections, electron-

stained with uranium and lead, and observed and photographed using a transmission

electron microscope (HT7800, Hitachi, Tokyo, Japan). For immunoelectron

microscopy, HeLa cells expressing Flag-Nwd1 grown on glass coverslips were fixed

175   with 4% PFA and 0.05% GA in 0.1 M PB (pH 7.4). The cells were washed with PB and

treated with 30% sucrose in 0.1 M PB, and rapid freeze fractionation was performed

using liquid nitrogen. Cells were blocked with normal goat serum for 2 h and incubated

with anti-DDDDK-tag antibody for 2 h. After incubation with 1.4-nm gold-conjugated

secondary antibody (Nanoprobes, Yaphank, NY, USA) for 1 h, the signals were gold-

180   sensitized using a gold enhancing kit (Nanoprobes). After washing with H_2O, cells were

refixed with 1% OsO_4 for 1 h, treated with ethanol and propylene oxide, and embedded

in Epon-812 resin. Sections (70 nm thick) were electron-stained with uranium and lead

and photographed using a transmission electron microscope (HT7800).

185   ***Plasmids***

pCAG-EGFP-Nwd1 was cloned as described previously (Yamada et al., 2020). pN3-

3×Flag-Control was purchased from Addgene (Watertown, MA, USA, plasmid

#107717, deposited by Dr. Suske) [23]. To construct 3×Flag-Nwd1 (hereafter called

Flag-Nwd1), the full-length *Nwd1* cDNA was isolated using RT-PCR of total RNA

190   from E12 embryonic and adult mice brains and then subcloned in-frame into the pN3-

3×Flag-Control plasmid. *Nwd1* cDNA corresponding to the N-terminal or C-terminal

region of the protein (accession number BC082552; 4–1026 bp for the N-terminus and

2578–4563 bp for the C-terminus) was subcloned into C-Flag-pcDNA3 (Addgene,

plasmid # 20011, deposited by Dr. Smale) [24] and pN3-3×Flag-Control to construct

195 Flag-Nwd1-N and 3×Flag-Nwd1-C (hereafter called Flag-Nwd1-C). To construct Halo-

Nwd1, full-length Nwd1 cDNA was subcloned into the pFN21A-HaloTag-CMV Flexi-

vector (Promega, Madison, WI, USA). To construct 3×Flag-SERCA2 (hereafter called

Flag-SERCA2), the coding region of SERCA2b (Addgene, plasmid #75188, deposited

by Dr. Lytton) [25] was subcloned in-frame into a pN3-3×Flag-Control vector. The

200 deletion mutants 3×Flag-SERCA2-N (amino acids 1–787, hereafter called Flag-Flag-

SERCA2-N) and 3×Flag-SERCA2-C (amino acids 788–1044, hereafter called Flag-

SERCA2-C) were constructed from 3×Flag-SERCA2 using a Q5 Site-Directed

Mutagenesis Kit (New England BioLabs, Ipswich, MA, USA). mCherry-Sec61B

(Addgene, plasmid # 121160, deposited by Dr. Mayr) [26] was used for ER imaging.

Primary antibodies

The following primary antibodies were used: anti-cleaved caspase-3 (Asp175) (rabbit

polyclonal, #9661, RRID: AB_2341188, Cell Signaling Technology, Danvers, MA,

USA; 1:400 for immunohistochemistry [IHC]), anti-cleaved caspase-1 (Ala317) p10

210 (rabbit polyclonal, #AF4022, RRID: AB_2845464, Affinity Biosciences, Cincinnati,

OH, USA; 1:200 for IHC), anti-BSCL2/SEIPIN (rabbit polyclonal, #ab106793, RRID:

AB_10974250, Abcam, Cambridge, UK; 1:200 for IHC, 1:1000 for western blotting

[WB]), anti-KDEL (mouse monoclonal, #M181-3, RRID: AB_10693914, MBL,

Woburn, MA, USA; 1:400 for IHC, 1:400 for immunocytochemistry [ICC], 1:5000 for

215 WB), anti-LAMP1 (1D4B) (rat monoclonal, #sc-19992, RRID: AB_2134495, Santa

Cruz Biotechnology, Dallas, TX, USA; 1:250 for IHC, 1:1000 for WB), anti-LC3A/B

(D3U4C) XP (rabbit monoclonal, #12741, RRID: AB_2617131, Cell Signaling

Technology; 1:200 for IHC, 1:2000 for WB), anti-Nwd1 (rabbit polyclonal generated via immunization with the recombinant mouse Nwd1 protein [18]; 1:400 for ICC), anti-SERCA2 ATPase Clone IID8 (mouse monoclonal, #NB300-529, RRID: AB_531361, Novus Biologicals, Centennial, CO, USA; 1:200 for ICC, 1:2500 for WB), anti-DDDDK-tag (mouse monoclonal, #M185-3, RRID: AB_11126775, MBL; 1:10000 for WB, 1:10000 for ICC), anti-ADP/ATP translocase 1/2 (ANT1/2, rabbit polyclonal, #17796-1-AP, RRID: AB_2190358, Proteintech, Rosemont, IL, USA; 1:2000 for WB), anti-voltage-dependent anion-selective channel protein 1 (VDAC1)/porin (mouse monoclonal, #ab14734, RRID: AB_443084, Abcam; 1:2000 for WB), anti-α-tubulin (rabbit polyclonal, #11224-1-AP, RRID: AB_2210206, Proteintech; 1:5000 for WB), anti-GRP78/BIP (rabbit polyclonal, #11587-1-AP, RRID: AB_2119855, Proteintech; 1:5000 for WB), anti-GRP94 (rabbit polyclonal, #14700-1-AP, RRID: AB_2233347, Proteintech; 1:5000 for WB), anti-activating transcription factor 6 (ATF6, rabbit polyclonal, #24169-1-AP, RRID: AB_2876891, Proteintech; 1:4000 for WB, 1:200 for IHC), anti-CD11B/integrin alpha M (rabbit polyclonal, #21851-1-AP, RRID: AB_2878927, Proteintech; 1:200 for IHC 1:2000 for WB), anti-RAB5A (rabbit polyclonal, #11947-1-AP, RRID: AB_2269388, Proteintech; 1:2000 for WB, 1:250 for IHC), and anti-RAB7A (rabbit polyclonal, #55469-1-AP, RRID: AB_11182831, Proteintech; 1:2000 for WB, 1:250 for IHC).

Hematological and biochemical analysis of blood and plasma samples

Peripheral whole blood was collected and briefly mixed in Insepak II (Sekisui, Tokyo, Japan), followed by incubation at room temperature for 15–30 min. Subsequently, the mixture was centrifuged at 3500 rpm for 15 min, and the supernatant was collected as

the plasma. Peripheral blood was collected using MiniCollect K2E K2EDTA tubes (Greiner Bio-One, Kremsmünster, Austria). Hematological and biochemical analyses were conducted by the BioSafety Research Center Inc. (Shizuoka, Japan).

Oil red O staining, Sirius red staining, periodic acid-Schiff (PAS) staining, and Congo red staining

General staining of liver tissue sections was performed using an Oil Red O Stain Kit (for fat staining) (ScyTek Laboratories, Logan, UT, USA), Picro-Sirius Red Stain Kit

250 (for collagen staining) (ScyTek Laboratories), Periodic Acid-Schiff (PAS) Diastase Stain Kit (for polysaccharides staining) (ScyTek Laboratories), and Amyloid Stain Kit (Congo Red) (for amyloid staining) (ScyTek Laboratories) according to the manufacturer's protocol. Images were acquired using a DP72 CCD camera (Olympus, Tokyo, Japan) mounted on an ECLIPSE E800 (Nikon, Tokyo, Japan), BX53

255 (Olympus), or BZ-X800 microscope (Keyence, Osaka, Japan). Quantification of tissue images was performed using the hybrid cell count algorithm (Keyence).

Immunostaining and TUNEL staining

Fluorescent immunostaining was performed as previously described [17]. Anesthetized

260 adult mice were perfused transcardially with saline solution, followed by 4% PFA in 0.1 M PB (pH 7.4). Livers were dissected and postfixed in the same fixative overnight at 4°C. Fixed tissues were cryoprotect in 30% sucrose in 0.1 M PB (pH 7.4) overnight at 4°C and embedded in optimal cutting temperature compound (Sakura Finetek, Torrance, CA, USA). Frozen sections (14 μm thick) were collected on MAS-coated

265 glass slides (Matsunami Glass, Bellingham, WA, USA). Sections were blocked for 2 h

with 5% normal goat or donkey serum in 0.1% Triton X-100 in PBS (PBST), followed by incubation with primary antibodies in blocking buffer at 4°C overnight. After four washes with PBST, sections were incubated for 2 h with Alexa Fluor 488-, Alexa Fluor 555-, or Alexa Fluor 647-conjugated secondary antibodies (Thermo Fisher Scientific).

270   After counterstaining with 0.7 nM Hoechst 33342 (Thermo Fisher Scientific) for the nucleus, 9 µM Bodipy (Thermo Fisher Scientific) for lipid droplets, or 75 µM Filipin III (Santa Cruz Biotechnology) for cytosol, sections were mounted and imaged using a confocal microscope (FV3000, Olympus). Filipin dye was used to delineate the entire cytoplasm, excluding vacuolar structures. To immunostain cultured cells, fixed cells

275   were blocked for 1 h with 5% normal goat or donkey serum in PBST, followed by incubation with primary antibodies in blocking buffer for 1 h. After three washes with PBST, cells were incubated for 1 h with Alexa Fluor 488-, Alexa Fluor 555-, or Alexa Fluor 647-conjugated secondary antibodies. After counterstaining with 0.7 nM Hoechst 33342, the cells were mounted. To detect apoptosis, TUNEL staining was performed

280   using the In Situ Cell Death Detection Kit, TMR-red (Roche Diagnostics, Basel, Switzerland) according to the manufacturer's instructions.

*** ER fraction and WB ***

The ER faction was obtained as previously described with some modifications [27,28].

285   Dissected fresh livers were homogenized in fraction buffer containing 10 mM HEPES (pH 7.4), 220 mM mannitol, 90 mM sucrose, and protease inhibitor cocktails (Complete mini, Merck Millipore) using Potter–Elvehjem homogenizers and then further homogenized via passage through a 27-gauge needle. The crude postnuclear supernatant was further clarified via centrifugation twice at $7000 \times g$ for 10 min each and then at

290　10,000 × *g* for 10 min at 4°C. The pellet and supernatant were collected as the

mitochondrial and cytosolic fractions, respectively. The cytosolic fraction was further

separated into an ER-enriched pellet and cytosolic supernatant by ultracentrifugation

(Optima LE-80K, Beckman Coulter, Brea, CA, USA) at 100,000 × *g* for 1 h at 4°C. The

ER pellet was resuspended in the fraction buffer, and the total protein concentration was

295　measured using the Bradford method (Bio-Rad, Hercules, CA, USA). For WB, the

samples were treated with 2× sample buffer (125 mM Tris-HCl, pH 6.8, 4% SDS, 10%

sucrose, and 0.01% bromophenol blue) for 1 h at 37°C, resolved by 8%–12% SDS-

PAGE, and electroblotted onto an Immobilon-P membrane (Merck Millipore) using a

semidry transfer apparatus. After blocking with 1%–5% skim milk in TBST (150 mM

300　NaCl, 10 mM Tris-HCl, pH 7.4, 0.1% Tween 20), membranes were incubated with

primary antibody for 1 h, followed by incubation with horseradish peroxidase (HRP)-

conjugated secondary antibody (Cytiva [Marlborough, MA, USA] or Jackson

ImmunoResearch [West Grove, PA, USA]). The signal was detected using Immobilon

Western chemiluminescent HRP substrate (Merck Millipore). Blot images were

305　captured using the Fusion Solo S system (Vilber Lourmat, Marne La Vallée, France)

and quantified using the Quantification Software Module (Vilber Lourmat).

Proteomic analysis and protein binding assay

HEK293T cells expressing Halo-Nwd1 were washed with ice-cold PBS and lysed in

310　lysis buffer containing 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1 mM EDTA, 1%

NP-40, and protease inhibitor cocktail for 15 min at 4°C. The lysate was diluted 4-fold

with Halo tag protein purification buffer containing 1× PBS, 1 mM DTT, and 0.005%

NP-40 (Nacalai Tesque, Kyoto, Japan). The lysate was centrifuged at 15,000 rpm for 10

min at 4°C, and the supernatant was incubated with HaloLink Resin (Promega)

315   overnight at 4°C. After brief centrifugation, the pelleted resin was washed four times

with Halo tag protein purification buffer. To determine the proteins interacting with

Halo-Nwd1, the pelleted resin was treated with 2× sample buffer for 1 h at 37°C and

resolved by SDS-PAGE. The gel was stained using a Silver Stain MS Kit (FUJIFILM

Wako Pure Chemical) according to the manufacturer's instructions. Gel slices

320   containing the protein bands were excised for proteomic analysis on a nanoLC-MS/MS

system (Japan Proteomics Inc., Tokyo, Japan). Mass data acquisitions and peptide

identification were performed using Mascot software. To verify the purification of

recombinant halo-Nwd1 protein, the resin crosslinked with Halo-Nwd1 was incubated

with HaloTEV protease (Promega) overnight at 4°C. Eluted Nwd1 protein was resolved

325   by SDS-PAGE, followed by Coomassie brilliant blue (CBB) staining. For the protein

binding assay, HEK293T cells expressing Halo-Nwd1 along with Flag-SERCA2, Flag-

SERCA2-N, or Flag-SERCA2-C were lysed and purified using HaloLink Resin as

previously described. Halo-Nwd1-binding proteins were treated with 2× sample buffer

for 1 h at 37°C and resolved by SDS-PAGE, followed by WB using anti-Flag antibody.

Quantitative PCR

Quantitative PCR was performed as previously described [29]. Total RNA prepared

from adult mouse livers was reverse-transcribed using a PrimeScript II first-strand

cDNA Synthesis Kit (Takara, Shiga, Japan) with a random hexamer primer. According

335   to the manufacturer's protocol, PCR was performed using TB Green Ex Taq II Mix

(Takara) and the Thermal Cycler Dice Real-Time System (Takara).

ATPase activity measurement

The ATPase activity of SERCA2 was measured using a previously described protocol

340 with some modifications [15]. The ATPase activity of the ER fraction was measured

using a QuantiChrom ATPase/GTPase assay kit (BioAssay Systems, Hayward, CA,

USA) following the manufacturer's instructions and quantified as colorimetric

absorbance using a Nivo S luminometer (PerkinElmer, Shelton, MA, USA). The

SERCA2 protein content in the ER fraction was determined by WB, and ATPase

345 activity was normalized to that of total SERCA2 protein.

Ca^{2+} measurement

The Ca^{2+} concentrations in ER and mitochondrial fractions were measured using the

fluorescent indicator Fluo3. Each fraction was treated in buffer containing 100 mM

350 KCl, 10 mM MOPS (pH 7.2), 0.01% Pluronic F-127, 2.6 µM Fluo3 (Dojindo,

Kumamoto, Japan) in OptiPlate-96 Black plates (PerkinElmer). The fluorescent

intensity (excitation: 495 nm, emission: 530 nm) was measured using a Nivo S

luminometer. The total protein content of each fraction was determined using the

Bradford method.

Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis

We used "DAVID Bioinformatics Resources" [30,31] to perform GO enrichment

analysis (including biological process, cellular component, and molecular function) and

360 KEGG pathway enrichment analysis [32].

Statistical analyses

Statistical analyses were performed using Microsoft Excel version 16.77.1 (Microsoft, Redmond, WA, USA) and Prism version 9.5.0 (GraphPad, La Jolla, CA, USA). All numerical data are expressed as the mean ± SD. One-way analysis of variance (ANOVA) followed by Welch's *t*-test with Holm–Bonferroni correction was used in multiple-group comparisons. Welch's *t*-test was used to analyze between-group differences for the following variables: Hematoxylin and eosin (HE)$^+$ normal hepatocytes, oil red O$^+$ area, Sirius red$^+$ fibrotic area, PAS$^+$ area, the number of dead cells in the liver, the result of blood tests, the expression of ER stress-related proteins after tunicamycin (TM) treatment as determined by WB, relative ATPase activity, and relative Ca^{2+} concentrations. The paired *t*-test was used to compare liver and body weights and protein expression between littermates. The chi-squared test was used to compare the number of genotypes in the offspring of *Nwd1*$^{+/−}$ mating pairs. Significance was indicated by P < 0.05. We calculated the area of swollen hepatocytes, oil red O$^+$ area, Sirius red$^+$ fibrotic area, and PAS$^+$ area using hybrid cell count software (Keyence). The numbers of lipid droplets and vacuolar structures in the liver were manually measured by polygon selection using cellSens imaging software (Olympus). Lipid droplets that partially filled the vacuole and detached from the vacuolar membrane were classified as type C vacuoles. The area of the vacuoles was measured as the area of the vacuolar structures minus that occupied by the lipid droplets.

Results

Generation of *Nwd1*$^{−/−}$ mice

The mouse Nwd1 gene was mapped to chromosome 8, and it consisted of 19 exons
(Fig. 1A). To reveal the *in vivo* function of *Nwd1*, we generated *Nwd1*$^{-/-}$ mice. *Nwd1*^{flox}
mice were generated by CRISPR-Cas9 genome editing using electroporation in zygotes.
In total, 320 embryos were transferred to the recipient uteri. Of the 71 offspring born,
five floxed mice were obtained. Conditional knockout of the *Nwd1*^{flox} allele generated
by genome editing was confirmed by PCR using genomic DNA (Fig. 1A, 1B). To
obtain the mice harboring the *Nwd1*$^{-}$ null allele, heterozygous *Nwd1*$^{flox/+}$ mice were
crossed with the CAG-Cre deleter mouse line, which ubiquitously expresses Cre
recombinase, thereby allowing deletion of the loxP-flanked genomic region (Fig. 1A).
The *Nwd1*$^{-}$ allele lacked the genomic region encompassing exon 5, a pivotal exon
encoding the NACHT domain (Fig. 1A). Subsequent interbreeding of the heterozygous
mutant mice yielded homozygous mutant (*Nwd1*$^{-/-}$) pups. The accuracy of targeted
disruption of *Nwd1* was confirmed through genomic sequencing and PCR (Fig. 1C).
Quantitative reverse transcription (qRT)-PCR analysis confirmed the absence of *Nwd1*
mRNA expression in *Nwd1*$^{-/-}$ mice (Fig. 1D). Genotyping of 211 adult offspring
produced by breeding heterozygous *Nwd1*$^{+/-}$ mice revealed that the number of *Nwd1*$^{-/-}$
mice was 7% lower than expected from the Mendelian ratio (Fig. 1E). Histological
examination demonstrated that most *Nwd1*$^{-/-}$ adult mice exhibited normal development
and no evident abnormalities in any organs excluding the liver and lungs. The livers of
Nwd1$^{-/-}$ mice exhibited an approximately 1.4-fold increase in liver size versus that in
wild-type mice (wild-type, 1.5 ± 0.4; *Nwd1*$^{-/-}$, 2.1 ± 0.4) despite no significant
differences in body weight (wild-type, 29.0 ± 5.9; *Nwd1*$^{-/-}$, 30.2 ± 5.2; Fig. 1F–H).
Gross observation revealed abnormal speckled fatty-like patterns (Fig. 1F, arrowheads)

and white fibrotic-like structures (Fig. 1F, arrows) on the surface of $Nwd1^{-/-}$ hepatic lobes.

The $Nwd1^{-/-}$ liver exhibits NAFLD/NASH-like pathology

HE staining of $Nwd1^{-/-}$ livers revealed a striking increase in the number of enlarged and lucent hepatocytes with a lack of eosin staining compared to the findings in wild-type littermates (wild-type, 12.0% ± 3.1%; $Nwd1^{-/-}$, 30.6% ± 3.5%; Fig. 2A, B).

415   These swollen hepatocytes did not appear to have undergone programed cell death because their nuclei were not contracted or fragmented (Fig. 2A). We next performed oil red O staining on the liver tissues to detect the neutral lipid content in hepatocytes, finding that the size of the positively stained area was substantially increased in $Nwd1^{-/-}$ livers (wild-type, 0.5% ± 0.4%; $Nwd1^{-/-}$, 13.0% ± 2.6%). Many lipid droplets,

420 which are storage organelles for neutral lipids, accumulated in a zonate pattern around the central vein in the $Nwd1^{-/-}$ hepatic lobule (Fig. 2C, D). We further assessed the organization of collagen fibers in tissues using Sirius red staining. As presented in Fig. 2E and F, $Nwd1^{-/-}$ mice exhibited severe collagen fiber deposition in their livers (wild-type, 2.0% ± 1.1%; $Nwd1^{-/-}$, 9.2% ± 6.5%), indicating that the loss of $Nwd1$ resulted in

425 abnormal fibrosis that resembled NASH. Congo red staining (Fig. S2A) illustrated that $Nwd1^{-/-}$ mouse livers did not accumulate amyloid fibrils, an extracellular deposition of amyloid proteins frequently observed in hepatic amyloidosis. PAS staining, which detects polysaccharides such as glycogen and mucosubstances such as glycoproteins and mucins, was significantly reduced in swollen $Nwd1^{-/-}$ hepatocytes (wild-type,

430 98.2% ± 0.15%; $Nwd1^{-/-}$, 61.5% ± 18.6%; Figs. 2A and S1B, C). Because pretreatment

with α-amylase abolished PAS staining in liver sections, insoluble glycogen was not

accumulated in the $Nwd1^{-/-}$ mouse liver (Fig. S1B).

Based on the swelling of hepatocytes and detection of fibrosis, we explored the

possibility of accelerated apoptosis in the $Nwd1^{-/-}$ liver using double staining with anti-

435   cleaved caspase-3 antibody and TUNEL staining. As presented in Fig. 2G, although

many TUNEL$^+$ cells were observed in $Nwd1^{-/-}$ livers (wild-type, 8.7% ± 8.6%;

$Nwd1^{-/-}$, 49.2% ± 19.2%), few caspase-3$^+$ apoptotic cells were found in these livers

(wild-type, 0.1% ± 0.2%; $Nwd1^{-/-}$, 0.5% ± 0.5%). TUNEL staining is known to detect

different types of cell death [33], including pyroptosis, a form of programed necrotic

440   cell death activated upon inflammation [34,35]. A much larger percentage of cells was

positive for cleaved caspase-1, a pyroptosis marker, in $Nwd1^{-/-}$ mouse livers (wild-

type, 11.7% ± 9.6%; $Nwd1^{-/-}$, 41.7% ± 12.3%; Fig. 2H, I), indicating accelerated

proptosis. The observation that $Nwd1^{-/-}$ hepatocytes lacked nuclear fragmentation, a

morphological sign of apoptosis, was also consistent with the observation of

445   inflammatory pyroptosis in $Nwd1^{-/-}$ livers. To evaluate the inflammatory response and

lipid metabolism, hematological and biochemical analyses were performed using serum

and peripheral whole blood. The total number of leukocytes was approximately 3-fold

higher in $Nwd1^{-/-}$ mice than in wild-type mice (relative ratio = 2.85 ± 1.00; Fig. 2J).

Although $Nwd1^{-/-}$ mice had increased counts of each type of leukocytes, including

450   monocytes (relative ratio = 3.41 ± 4.29; Fig. 2J), lymphocytes, neutrophils, eosinophils,

and basophils, the proportion of each cell type was unchanged (Fig. S2A, B). Lipid

analysis revealed that the total content of triglycerides (relative ratio = 0.98 ± 0.26) and

cholesterol (relative ratio = 0.92 ± 0.16) in the peripheral serum remained unchanged

(Fig. 2K) despite the lipid accumulation in $Nwd1^{-/-}$ livers, suggesting impairment of the

455 lipid droplet efflux in $Nwd1^{-/-}$ mice. The pathological phenotype of $Nwd1^{-/-}$ mice,

including hepatomegaly, excessive lipid accumulation in hepatocytes, fibrosis, and

inflammatory pyroptosis, closely resembled that of NASH.

In addition to the liver, we performed histological analysis of various organs.

$Nwd1^{-/-}$ mouse lungs featured flattened alveoli (Fig. S3). We observed no

460 abnormalities in the structure or tissue architecture of the heart, intestine, kidneys,

skeletal muscle, spleen, and testes (Fig. S3).

Accumulation of lipid droplets and ER-derived vacuoles in $Nwd1^{-/-}$ hepatocytes

Ultrastructure analysis using electron microscopy revealed that $Nwd1^{-/-}$ hepatocytes

465 were occupied by numerous vesicular structures. These cytosolic vesicles varied in size,

and they were divided into two types based on the inner structure: gray vesicles filled

with electron-dense material and vacuoles that appeared vacant or contained small

amounts of lipid (Fig. 3A, B). Ultrastructurally, both types of vesicles were surrounded

by a single membrane (Fig. 3B). Based on these morphological characteristics, we

470 concluded that the gray vesicles filled with electron-dense materials corresponded to the

lipid droplets stained with oil red O. Notably, many tiny lipid droplets were evident in

the wild-type liver sinusoid (Fig. 3A, arrows). However, we rarely observed lipid

droplets in the $Nwd1^{-/-}$ sinusoid (Fig. 3B). To distinguish the vesicles, Bodipy staining

of lipid droplets was performed in conjunction with cytoplasmic staining. Consistent

475 with the result of electron microscopy, $Nwd1^{-/-}$ hepatocytes contained many vacant

vacuoles in addition to lipid droplets filled with Bodipy$^+$ triglycerides (Fig. 3C, D). We

classified the vesicular structures based on Bodipy staining into three types (Fig. 3E):

type A, a typical lipid droplet surrounded by a membrane; type B, a vacant vacuole

lacking Bodipy$^+$ lipids; and type C, a vacuole containing a small amount of Bodipy$^+$

480   lipids, representing a transitional morphology between types A and B. The $Nwd1^{-/-}$

liver exhibited increases levels of all subtypes (Fig. 3E). Quantification of the areas

occupied by lipid droplets and vacuoles in a single hepatocyte revealed a significant

increase in both lipid droplets (wild-type, $1.98 \pm 2.08\ \mu m^2$; $Nwd1^{-/-}$, $7.98 \pm 6.96\ \mu m^2$)

and vacuolar structures (wild-type, $3.95 \pm 2.46\ \mu m^2$; $Nwd1^{-/-}$, $12.18 \pm 9.13\ \mu m^2$) in

485     $Nwd1^{-/-}$ mice (Fig. 3F, H). The numbers of giant lipid droplets larger than $2.5\ \mu m^2$ (Fig.

3G) and vacuoles larger than $5\ \mu m^2$ (Fig. 3I) were significantly increased in $Nwd1^{-/-}$

mice. Based on the increased size and number of lipid droplets, the absence of lipid

droplets in the sinusoids (Fig. 3B), and the unchanged levels of total triglycerides and

cholesterols in peripheral blood (Fig. 2K), we assumed that $Nwd1^{-/-}$ mice have

490   defective lipid droplet efflux from the sinusoids, in addition to increased lipid synthesis

in hepatocytes.

       From where do these empty vacuoles originate? Previous studies revealed that

lipid droplet biogenesis is initiated in the ER, and ER-derived membrane structures

enclose the lipid during the maturation of lipid droplets [11,15,36-38]. Cells lacking

495   SEIPIN/BSCL2, a protein localized to ER–lipid droplet contact sites, had fewer ER–

lipid droplet contact sites, and the transport of lipid droplets occurred independently of

the ER [11]. Immunostaining for SEIPIN demonstrated that the $Nwd1^{-/-}$ liver exhibited

normal SEIPIN$^+$ zonation surrounding the central vein, and the area of SEIPIN$^+$

zonation remained unchanged (Fig. 4A, B). Hepatocytes play distinct metabolic roles

500   depending on their specific location along the lobular postcentral axis. This regional

segregation of hepatocytes is called zonation. Consistently, the accumulation pattern of

Bodipy$^+$ lipid droplets within $Nwd1^{-/-}$ hepatic lobules appeared to be distributed as

hepatic zonation surrounding a central vein (Fig. 4B). Immunostaining for the ER

marker KDEL illustrated that many vacuoles in $Nwd1^{-/-}$ hepatocytes were surrounded

505   by SEIPIN$^+$ KDEL$^+$ membranes, confirming that they were derived from the ER (Fig.

4C). Interestingly, RAB5, an early endosome marker, was frequently expressed in these

ER-derived vacuoles (Fig. 4D). By contrast, RAB7, an early-to-middle endosome

marker, exhibited little localization to the vacuoles (Fig. 4E). Moreover, the vacuolar

structures did not express LAMP1 or LC3A/B, indicating that they are unrelated to the

510   autophagy–lysosome–proteasome pathway (Figs. 4F, G and S4A, B).

Elevated ER stress in the $Nwd1^{-/-}$ liver

We examined the expression of proteins associated with ER and other intracellular

organelles, including mitochondria, endosomes, and lysosomes. Notably, the ER

515   membrane proteins SERCA2 and SEIPIN were upregulated in the $Nwd1^{-/-}$ liver

compared to their wild-type expression (Fig. 5A, B). We also confirmed that the

$Nwd1^{-/-}$ liver exhibited elevated expression of a 100-kDa KDEL protein (Fig. 5A, B).

Conversely, the expression of ANT2 and VDAC, which are proteins of the inner and

outer mitochondrial membranes, respectively, remained unchanged in the $Nwd1^{-/-}$ liver

520   (Fig. 5A, B). RAB5 and LAMP1 expression was slightly increased in the $Nwd1^{-/-}$ liver,

whereas RAB7 and LC3A/B expression remained unchanged (Fig. 5A, B). We

concluded that many ER-related proteins were upregulated in the $Nwd1^{-/-}$ mouse liver.

Because the induction of ER stress is known to be accompanied by increased

expression of ER-localized proteins such as KDEL proteins, it was speculated that ER

525   stress was enhanced in the livers of $Nwd1^{-/-}$ mice. A previous study illustrated that

tunicamycin (TM) treatment induced ER stress *in vivo* and the development of vacuolar

structures in the mouse liver [39]. To assess the susceptibility of $Nwd1^{-/-}$ mice to ER

stress, we intraperitoneally administered TM into wild-type and $Nwd1^{-/-}$ adult mice.

Although wild-type mice survived up to 48 h after TM injection, 40% of $Nwd1^{-/-}$ mice

530   died within 48 h (Fig. 5C), postulating that $Nwd1^{-/-}$ mice were more vulnerable to ER

stress or that the basal level of ER stress was already elevated in these mice. To directly

estimate whether ER stress is enhanced in the livers of $Nwd1^{-/-}$ mice, the ER fraction

was prepared from wild-type and $Nwd1^{-/-}$ livers following treatment with or without

TM, and the expression of ER stress-related proteins was quantified by WB (Fig. S4).

535   ATF6, GRP78 (BiP), GRP94, and KDEL are known to be upregulated upon the

induction of ER stress, and they have been used as markers of ER stress [40,41]. As

presented in Figure 5D, TM administration to wild-type mice increased the expression

of KDEL proteins harboring ER localization signal sequences, including GRP78 and

GRP94, confirming that TM treatment induces ER stress in hepatocytes. At this time,

540   the levels of these proteins were elevated in the ER fraction of $Nwd1^{-/-}$ livers regardless

of TM treatment (KDEL: wild-type, 1.00 ± 0.30; $Nwd1^{-/-}$, 2.90 ± 0.91; wild-type with

TM treatment, 2.29 ± 0.06; $Nwd1^{-/-}$ with TM treatment, 233 ± 0.48; GRP78: wild-type,

1.00 ± 0.11; $Nwd1^{-/-}$, 1.57 ± 0.19; wild-type with TM treatment, 3.88 ± 1.23; $Nwd1^{-/-}$

with TM treatment, 6.11 ± 1.12; and GRP94: wild-type, 1.00 ± 0.46; $Nwd1^{-/-}$, $1.33 \pm$

545   0.67; wild-type with TM treatment, 2.83 ± 0.64; $Nwd1^{-/-}$ with TM treatment, $4.25 \pm$

0.76; Fig. 5D, E). Additionally, a significant increase in the expression of ATF6, an ER

stress-responsive gene that encodes an ER-membrane-bound transcription factor

activated by ER stress-induced proteolysis [39,41], was observed in the $Nwd1^{-/-}$ liver

regardless of TM treatment (wild-type, 1.00 ± 0.61; $Nwd1^{-/-}$, 2.35 ± 0.87; wild-type

550   with TM treatment, 2.81 ± 0.98; $Nwd1^{-/-}$ with TM treatment, 3.02 ± 1.15; Fig. 5D, E).

Electron microscopy revealed that TM-mediated ER stress drove the accumulation of large lipid droplets and vacuoles in the wild-type liver (Fig. 5F). This hepatic pathology closely resembled that in the $Nwd1^{-/-}$ liver without TM administration (Figs. 3B and 5G). Bodipy staining confirmed that TM-mediated ER stress accelerated the

555   accumulation of lipid droplets in wild-type and $Nwd1^{-/-}$ livers (Fig. 5H, I). Immunostaining using ATF6 antibody illustrated that the nuclear translocation of ATF6 was induced in $Nwd1^{-/-}$ livers (Fig. 5K). ATF6 translocation was never detected in wild-type mice (Fig. 5J), indicating that ER stress is constitutively induced in the $Nwd1^{-/-}$ liver.

Nwd1 interacts with ER-associated proteins, including SERCA2

To understand the molecular mechanism underlying ER stress induced by Nwd1 depletion, Nwd1-binding proteins were screened using the pulldown assay and proteomic analysis. Nwd1-interacting proteins were copurified with halo-tagged Nwd1

565   protein from HEK293T cell lysate (Fig. 6A). Proteomic analysis using nanoLC-MS/MS identified 107 proteins as candidate Nwd1-binding proteins, including SERCA2, the mitochondrial outer membrane proteins VDAC1 and sideroflexin-1, the endosome membrane protein RAB7A, and chaperone proteins including calnexin and heat shock 70 kDa protein 1A (Fig. 6B and Tables S1–S3). GO analysis for biological processes

570   suggested a predominant role of Nwd1 in protein transport (Fig. 6C). According to the predicted distribution of the cellular components and the prediction of molecular function, many Nwd1-binding proteins were expected to localize to the ER membrane and various organelle membranes and function in protein binding (Fig. 6C). *In silico* pathway analysis using the KEGG database predicted that many Nwd1-binding proteins

575   were involved in protein processing in the ER (Fig. 6C). Collectively, these analyses

strongly predicted the involvement of Nwd1 in the membrane transport or processing of

ER proteins.

We examined whether Nwd1 is distributed in the ER. Double immunostaining

demonstrated the co-localization of Nwd1 and KDEL in cultured HEK293T cells (Fig.

580   6D). To analyze the association between Nwd1 and ER at higher resolution,

immunoelectron microscopy was performed using HeLa cells expressing Flag-Nwd1.

As presented in Figure 6E, many Flag-Nwd1 signals were detected on and near the

surface of the ER membrane but rarely in the lumen of the ER, confirming the close

association of Nwd1 with the ER membrane. To identify the protein region responsible

585   for ER localization, deletion mutants of Nwd1 were introduced into cultured cells

together with mCherry-Sec61B to visualize the ER architecture [26,29]. The

overlapping distribution of Flag-tagged full-length Nwd1 and mCherry-Sec61B verified

the predominant localization of Nwd1 to the ER. Flag-Nwd1-N, which lacks the central

NACHT domain and C-terminal WD40 repeat domain, localized to the cytoplasm or

590   nucleus (Fig. 6F). Conversely, Flag-Nwd1-C, which only contains the WD40 repeat

domain, was explicitly localized to the ER (Fig. 6F), indicating that the C-terminal

WD40 repeat domain of Nwd1 is crucial for its ER localization and thus for interactions

with ER proteins. Generally, the WD peptide motif folds into a β-propeller structure,

and proteins containing WD40 repeats are believed to serve as platforms for assembling

595   protein complexes or as mediators of transient interactions between other proteins [42].

Among the ER proteins identified as potential Nwd1-binding proteins, we focused

on SERCA2b, which had the highest spectral count in the pulldown assay (Fig. 6B).

SERCA2 is an ATPase pump embedded in the ER membrane that transports Ca^{2+} from

the cytosol to the ER lumen at the expense of ATP hydrolysis [15]. Immunostaining

600   revealed that the intracellular distribution of Nwd1 protein was identical to that of

endogenous and exogenous SERCA2 protein (Fig. 6H). The protein topology of

SERCA2b includes a cytoplasmic region comprising an actuator (A) domain (amino

acids 1–43 and 124–235), a nucleotide-binding (N) domain (amino acids 360–600), a

phosphorylation (P) domain (amino acids 330–359 and 601–739), and 11

605   transmembrane domains [43,44] (Fig. 6G). As previously described [45], we generated

two deletion constructs of SERCA2b named SERCA2-N and SERCA2-C. SERCA2-N

contains mainly the A, N, and P domains (amino acids 1–787), and SERCA2-C

corresponds to the region containing the ER lumen and the short cytoplasmic region

(amino acids 788–1044; Fig. 6G). SERCA2-N or SERCA2-C was introduced to

610   HEK293T cells with Flag-Nwd1, and their intracellular distribution was monitored. As

presented in Figure 6I, both SERCA2 deletion mutants localized to the ER similarly as

full-length SERCA2. *In vitro* protein binding assays using Halo-Nwd1 and Flag-

SERCA2 mutants demonstrated that Nwd1 binds to both the N-terminal region of

SERCA2 outside the ER and the C-terminal region (Fig. 6J). These results indicate that

615   Nwd1 recognizes a broad cytosolic region spanning the SERCA2 protein to regulate the

trafficking, conformational folding, or embedding of SERCA2 into the ER membrane.

**Decreased ATPase activity of SERCA2 and smaller Ca^{2+} pools in the *Nwd1$^{-/-}$* liver
ER**

620   How does the Nwd1–SERCA2 interaction contribute to ER stress or homeostasis? ER

fractions were prepared from wild-type and *Nwd1$^{-/-}$* livers, and SERCA2 ATPase

activity, which regulates Ca^{2+} transport from the cytosol to the ER lumen, was

measured. The results indicated that SERCA2 ATPase activity was significantly

reduced by approximately 70% in the $Nwd1^{-/-}$ liver (wild-type, 1.00 ± 0.64; $Nwd1^{-/-}$,

625   0.29 ± 0.09; Fig. 7A). Consistent with the decrease in ATPase activity, Ca^{2+} stores in

the $Nwd1^{-/-}$ ER were depleted by approximately 40% (wild-type, 1.00 ± 0.09; $Nwd1^{-/-}$,

0.63 ± 0.03; Fig. 7B). Contrarily, the amount of Ca^{2+} inside mitochondria did not differ

between wild-type and $Nwd1^{-/-}$ livers (wild-type, 1.00 ± 0.30; $Nwd1^{-/-}$, 0.92 ± 0.16;

Fig. 7C). Considering the finding that SERCA2 protein expression was drastically

630   increased by Nwd1 deficiency (Fig. 5), it was likely that the proportion of SERCA2

lacking pump activity is increased in $Nwd1^{-/-}$ livers. These results implied that Nwd1 is

required for the folding or topology of SERCA2 on the ER membrane, enabling

SERCA2 to acquire proper activity.

635   **Discussion**

Previous studies illustrated that ER stress contributes to hepatic steatosis by inducing

the accumulation of lipid droplets [8,9]. However, the mechanism by which ER stress in

the liver influences the development of hepatic steatosis is unclear. We demonstrated

that Nwd1 deficiency led to decreased Ca^{2+} storage in the ER *via* the reduction of

640   SERCA2 ATPase activity, leading to ER stress accompanied by diverse hepatic

pathologies, including the accumulation of lipid droplets and ER-derived vacuoles in

hepatocytes, hepatomegaly, hepatic fibrosis, and caspase-1–dependent inflammatory

pyroptosis along with an elevated number of leukocytes in peripheral blood (Fig. 7D).

Many of the pathologies in $Nwd1$-deficient mice are comparable to those in NASH in

645   humans. Quality control of newly synthesized proteins largely depends on the Ca^{2+}

concentration in the ER lumen. For instance, the calcium-binding protein GRP78 plays

a pivotal role in detecting the accumulation of misfolded proteins in the ER in cooperation with other ER transmembrane proteins such as ATF6, IRE1, and PERK [46,47]. Protein disulfide isomerases, a major group of ER-resident proteins, directly

650   interact with Ca^{2+} to serve as molecular chaperones for protein synthesis and maturation. Therefore, dysregulation of Ca^{2+} homeostasis in the ER leads to the collapse of protein quality control, resulting in the induction of ER stress [46]. It is well established that ER stress associated with unfolded proteins causes the excessive accumulation of lipid droplets surrounded by a monolayer membrane of ER-derived

655   phospholipids in the liver [7,9,41,48-50].

Despite the abundant accumulation of lipid droplets, no significant increases in serum triglyceride and total cholesterol levels were observed in the $Nwd1^{-/-}$ liver. Considering that lipid droplets in the sinusoid were rarely observed in the $Nwd1^{-/-}$ liver using electron microscopy, this NASH-like phenotype might be related to impaired

660   hepatic fat clearance in addition to augmented *de novo* fat synthesis in the $Nwd1^{-/-}$ liver. Consistently, a previous report in mice treated with TM disclosed the induction of vacuolar structures in the liver and concomitant decreases in total cholesterol and triglyceride levels in plasma, presumably attributable to the suppression of VLDL emission from the liver [39]. Furthermore, recent studies highlighted the involvement of

665   pyroptosis, a caspase-1–mediated programed cell death process, in the pathology of hepatic steatosis and NASH [51]. Pyroptosis is initiated by the activation of caspase-1 through inflammasome formation, leading to pore formation in the plasma membrane and the release of pro-inflammatory factors into the pericellular space. Specifically, the release of caspase-1–dependent inflammatory complexes in hepatocytes promotes liver

670   fibrosis and the excessive accumulation of extracellular matrix proteins, including

collagen [34]. Inhibiting pyroptosis has been reported to effectively reduce fat

deposition in the liver and reduce liver inflammation in NASH [52]. Along the same

lines, increased pyroptosis might have driven fibrosis in the $Nwd1^{-/-}$ liver. Deciphering

the molecular functions of Nwd1 could shed light on the novel aspects of the etiology of

675   diseases associated with increased ER stress and impaired Ca^{2+} homeostasis, such as

NAFLD and NASH.

Nwd1 might play an essential role as a co-chaperone in modulating the activity or

conformational regulation of SERCA2 on the ER membrane by binding to a specific

domain of SERCA2. Immunoelectron microscopy revealed that Nwd1 localizes on the

680   membrane or in close proximity to the cytoplasmic side of the ER, but it was rarely

found within the ER. *In vitro* binding assays illustrated that Nwd1 recognizes and binds

to a broad region of SERCA2, including cytosolic ATPase domains and the C-terminal

short region, which is involved in the Ca^{2+} affinity of SERCA2b [53]. A previous study

found that the topological fluctuation of SERCA2b is important for the transition of

685   Ca^{2+} affinity [44]. Nwd1 might affect the Ca^{2+} affinity of SERCA2 by regulating its

topology through binding to the C-terminal region, in addition to modulating ATPase

activity through binding to the A, N, and P domains of SERCA2. The elevated

expression of SERCA2 protein observed in $Nwd1^{-/-}$ livers might be attributable to a

compensatory mechanism in response to the increased proportion of SERCA2 lacking

690   proper pump activity.

Our previous study illustrated that Nwd1 regulates the formation of purinosomes

during cerebral cortex development by interacting with PAICS, an enzyme required for

de novo purine biosynthesis [18]. The purinosome is a macromolecular metabolic

complex consisting of six enzymes that catalyze *de novo* purine biosynthesis and

695　chaperone proteins necessary for the association of component proteins [54]. Nwd1 is

also reported to interact with several chaperones in prostate cancer cell lines [55]. These

facts led us to speculate that Nwd1 might function as a co-chaperone for protein folding,

allowing target proteins, including SERCA2, to acquire appropriate activity. Elucidating

the molecular machinery by which *Nwd1* is involved in hepatocellular pathology could

700　provide clues to understanding the mechanisms of NASH pathogenesis and developing

new therapeutic strategies for NASH associated with ER stress.

Abbreviations

ER, endoplasmic reticulum; NAFLD, nonalcoholic fatty liver disease; NASH,

705　nonalcoholic steatohepatitis; SERCA2, sarco/endoplasmic reticulum calcium ATPase;

Nwd1, NACHT and WD repeat domain-containing protein 1; VLDL, very low-density

lipoprotein; STAND, signal transduction ATPases with numerous domains; NSPCs,

neural stem/progenitor cells; ssODNs, single-stranded oligodeoxynucleotides; DMEM,

Dulbecco's Modified Eagle Medium; PFA, paraformaldehyde; GA, glutaraldehyde; PB,

710　phosphate buffer; PAS, periodic acid-Schiff; HRP, horseradish peroxidase; GO, Gene

Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; TM, tunicamycin; HE,

hematoxylin and eosin; ANT2, ADP/ATP translocase 2; VDAC, voltage-dependent

anion-selective channel protein 1; ATF6, activating transcription factor 6.

715　**Author contributions**

Study concept and design: S.Y. and S.S. Acquisition of data: S.Y., K.N., T.M., K.K.,

T.H., R.K., I.H., and S.S. Analysis and interpretation of data: S.Y., K.N., and S.S.

Drafting of the manuscript: S.Y., T.H., and S.S. Obtained funding: S.Y., I.H., and S.S

Conflict of interest

The authors declare no competing interests.

Acknowledgment

This work was funded by the Japan Society for the Promotion of Science grants-in-aid (KAKENHI) grant numbers 23K05996 (to S. S.) and 21K20701 (to S. Y.), the Research Support Project for Life Science and Drug Discovery (Basis for Supporting Innovative Drug Discovery and Life Science Research) from AMED under Grant Number JP21am0101120 (to I. H), Gout and Uric Acid Research Foundation 2020 (to S. S) and 2022 (to S. Y.), and Waseda University Grants for Special Research Projects 2021C-611 (to S. Y.), 2022C-611 (to S. Y.), 2022C-214 (to S. S.), and 2023C-207 (to S. S.). We would like to thank Ms. M. Mori and Ms. M Hidaka for their technical assistance. We also thank Enago (www.enago.jp) for English editing.

Figure legends

Fig. 1. Generation of *Nwd1* knockout mice

(A) Schematic diagram of the *Nwd1* wild-type allele (top) and the conditional allele of *Nwd1*^{flox} (middle), in which LoxP sequences were inserted into the introns on both sides (crRNA targeting sites: Nwd1L3 and Nwd1R1) of exon 5 *via* two rounds of genomic editing using ssODNs (Nwd1L3loxpAS and Nwd1R1loxP). Nwd1^{flox} mice were crossed with CAG-Cre transgenic mice harboring CAG promoter-driven Cre recombinase to generate the *Nwd1*$^{-}$ null allele (bottom). Upon elimination of the floxed *Nwd1* critical exon (exon 5), this allele generates a truncated RNA lacking the NACHT domain of the

open-reading frame of *Nwd1*. (B) PCR genotyping of wild-type or *Nwd1*flox/+ tail

genomic DNA. The specific primers (primer pair F1–R1 depicted in A) were used to

detect wild-type (189-bp amplicon) and *Nwd1*flox (229-bp amplicon) alleles. PCR

products were separated by 5% SDS-PAGE. (C) PCR genotyping of wild-type,

Nwd1+/−, and *Nwd1*−/− mice. Specific primers (primer pairs F1–R1 and F2–R2 depicted

in A) were used to detect wild-type (189-bp amplicon) and *Nwd1*− (431-bp amplicon)

alleles. PCR products were separated by 5% SDS-PAGE. (D) Loss of *Nwd1* mRNA

expression in *Nwd1*−/− livers. The mRNA isolated from wild-type and *Nwd1*−/− adult

livers were subjected to qRT–PCR, and *Nwd1* expression was normalized to that of *β-actin* mRNA. (E) The table presents the genotypes and birth rates observed in 116 adult

mice born from crosses between *Nwd1*+/− males and females. Numbers in parentheses

represent the expected number of offspring and birth rate according to the Mendelian

ratio. (F) Gross anatomical observation of the liver. *Nwd1*−/− livers exhibited

hepatomegaly (upper), fiber-like structures on the surface (middle, arrows), and

numerous abnormal hepatic lobules that turned whitish (lower, arrowheads). (G, H)

Liver (G) and body weight (H) of adult *Nwd1*−/− mice and their wild-type littermates

were compared by a paired *t*-test. ns, not significant; **, P < 0.01. Scale bars, 5 (F,

upper) or 2 mm (F, lower).

Fig. 2. NASH-like phenotype in the adult *Nwd1*−/− mouse liver

(A–I) Histological analyses of the adult mouse liver. HE (A, B), oil red O (C, D), and

Sirius red (E, F) staining of *Nwd1*−/− or wild-type livers. Nuclei were counterstained

with hematoxylin (C, upper). Welch's *t*-test was used to compare the number of swollen

hepatocytes (%) (B), oil red O+ area (%) (D), and Sirius red+ fibrotic area (%) (F). *, P

< 0.05; **, P < 0.01. Arrows denote swollen hepatocytes in (A) and Sirius red$^+$ fibrotic pathology in (E). (G) Double staining with anti-cleaved caspase-3 antibody (*green*) and TUNEL (*red*). (H) Immunostaining using anti-cleaved caspase-1 (*red*) antibody. Nuclei were stained with Hoechst dye (*blue*). (I) Comparison of the number of cleaved caspase-3$^+$ cells (%), TUNEL$^+$ cells (%), and cleaved caspase-1$^+$ cells (%) (Welch's *t*-test). ns, not significant; **, P < 0.01; ***, P < 0.001. (J) Analysis of leukocyte and monocyte counts in peripheral blood (relative number). (K) Relative comparison of triglyceride and total cholesterol levels in serum. Scale bars, 10 (A [upper]), 40 (A [lower]) and (E [lower]), 50 (C and E [upper]), or 100 μm (G and H).

Fig. 3. Accumulation of lipid droplets and vacuolar structures in *Nwd1*$^{-/-}$ hepatocytes

(A, B) Electron microscopy of wild-type (A) and *Nwd1*$^{-/-}$ (B) adult mouse liver. Arrows in (A) indicate the secretion of lipid droplets from hepatocytes into the wild-type liver sinusoid. (B) *Nwd1*$^{-/-}$ hepatocytes contained numerous large vacuolar structures (black arrowheads) partially or entirely filled with electron-dense lipid droplets (red arrows). (C, D) Lipid droplet staining with Bodipy dye (*green*) in wild-type (C) and *Nwd1*$^{-/-}$ livers (D). The cytosol (*magenta*) was counterstained. (E) Vesicular structures observed in *Nwd1*$^{-/-}$ hepatocytes were classified into three types: type A, a typical lipid droplet surrounded by a membrane; type B, a vacant vacuole lacking Bodipy$^+$ lipids; and type C, a lipid droplet partially or fully detached from the surrounding vacuolar membrane. The bars denote the distributions of types A, B, and C vesicles in wild-type and *Nwd1*$^{-/-}$ hepatocytes. (E–I) Quantification of the area occupied by lipid droplets (F), the number and size of lipid droplets (G), the area occupied by vacuolar space (H), and the number

of vacuoles (I) in wild-type and $Nwd1^{-/-}$ hepatocytes. Welch's *t*-test, ***, P < 0.001.

Scale bars, 2 (A and B), 50 (C [upper] and D [upper]), or 5 μm (C [lower] and D

[lower]).

795 **Fig. 4. Lipid droplets and vacuolar structures of $Nwd1^{-/-}$ hepatocytes were derived**

from the ER.

(A, B) Lipid droplets (Bodipy, *green*) and SEIPIN (*red*) staining of wild-type (A) and

$Nwd1^{-/-}$ livers (B). Note the SEIPIN⁺ zonation surrounding the central vein in the

$Nwd1^{-/-}$ hepatic lobule. (C) Magnified view of the SEIPIN⁺ (*red*) area in $Nwd1^{-/-}$

800 lobules simultaneously stained with Bodipy (*green*), anti-KDEL (*magenta*), and

cytoplasm (*cyan*). Lower panels present a high-magnification view of the $Nwd1^{-/-}$

hepatic cord. Many vacuolar structures were encapsulated by ER-derived membranes

that were positive for SEIPIN and KDEL (arrows). (D, E) Magnified view of the

hepatic cord immunostained with anti-RAB5 (D) or anti-RAB7 (E) together with anti-

805 KDEL (*blue*) antibody and cytosol staining (*cyan*). (F, G) $Nwd1^{-/-}$ hepatic cord

immunostained with anti-LAMP1 (F) or anti-LC3A/B (*green*) and anti-KDEL

(*magenta*) antibodies (G). The cytoplasm or nuclei were counterstained (*cyan*) (G).

Scale bars, 100 (A and B), 50 (C [upper]), or 5 μm (C [lower], D, E, F, and G).

810 **Fig. 5. Increased ER stress in $Nwd1^{-/-}$ livers**

(A) WB of wild-type and $Nwd1^{-/-}$ adult livers using the indicated antibodies. The blot

was probed with anti-α-tubulin antibody to confirm equal loading. (B) Quantitative

comparison of protein expression between an $Nwd1^{-/-}$ mouse and its wild-type

littermate (paired *t*-test). ns, not significant; *, P < 0.05; **, P < 0.01. (C) Survival

815　curves of *Nwd1*$^{-/-}$ and wild-type mice after TM treatment. (D) WB of ER stress-related

proteins in ER fractions prepared from livers with or without TM treatment. (E)

Quantitative comparison of the expression of each protein in (D). One-way ANOVA

followed by Welch's *t*-test with Holm–Bonferroni correction. ns, not significant; *, P <

0.05; **, P < 0.01. (F, G) Electron microscopy of *Nwd1*$^{-/-}$ and wild-type hepatocytes

820　treated with TM. A significantly increased number of small vacuoles (black

arrowheads), in addition to large lipid droplets (red arrows), was observed in both wild-

type and *Nwd1*$^{-/-}$ livers after TM treatment. (H, I) Lipid (Bodipy, *green*) and ER (anti-

KDEL, *magenta*) staining of wild-type and *Nwd1*$^{-/-}$ livers with or without TM

treatment. (J, K) ATF6 expression (*red*) in wild-type and *Nwd1*$^{-/-}$ livers. Lower panels

825　present magnified views. Arrows indicate nuclear translocation of ATF6, which

occurred only in *Nwd1*$^{-/-}$ livers. Nuclei were counterstained with Hoechst dye (*blue*).

Scale bars, 2 (F and G), 50 (H, I, J [upper], and K [upper]), or 5 μm in (J [lower] and K

[lower]).

830　**Fig. 6. Nwd1 binds to SERCA2**

(A) Purification of Halo-Nwd1 recombinant protein expressed in HEK293T cells. CBB

staining of the SDS-PAGE gel of the cell lysate and purified protein. (B) Identification

of Halo-Nwd1-binding proteins by nanoLC-MS/MS. The list of representative Nwd1-

binding proteins presents the gene symbols, the spectral counts in proteomic analysis,

835　and the UniProt protein database entry numbers. (C) GO enrichment and KEGG

pathway analyses of Nwd1-binding proteins. (D) HEK293T cells expressing Flag-

Nwd1. ICC with anti-Nwd1 (*green*) and anti-KDEL (*red*) antibodies revealing the co-

localization of Nwd1 with the ER. (E) Immunoelectron microscopy revealing the

localization of Flag-Nwd1 in the vicinity of the ER. HeLa cells were transfected with Flag-Nwd1 and immunostained with anti-Flag antibody (gold particle). Arrows indicate the Nwd1 signals on or near the ER membranes. MT, mitochondria. (F) Analysis of the Nwd1 region required for localization to the ER. HEK293T cells were transfected with mCherry-Sec61B and Flag-Nwd1, Flag-Nwd1-N, or Flag-Nwd1-C. Cells were stained with anti-Flag (*green*) antibody. (G) Structure and topology of mouse SERCA2b protein (1044 amino acids, accession number NP_733765.1). Blue boxes denote the cytosolic regions comprising the A (amino acids 1–43 and 124–235), N (amino acids 360–600), and P domains (amino acids 330–359 and 601–739), which are involved in ATPase activity. Black boxes denote 11 transmembrane domains, and white boxes denote the ER luminal regions. The red and yellow lines represent SERCA2b–N (amino acids 1–787) and SERCA2b–C (amino acids 788–1044), the deletion mutants of SERCA2b produced in this study. (H) Co-localization of Nwd1 and SERCA2 in the ER. HEK293T cells were transfected with EGFP-Nwd1 and stained with anti-SERCA2 (*red*) antibody. (I) HEK293T transfected with EGFP-Nwd1 and Flag-SERCA2, Flag-SERCA2-N, or Flag-SERCA2-C were stained with anti-Flag (*green*) antibody. (J) *In vitro* binding assay of Nwd1 with SERCA2. Cell lysate of HEK293T cells expressing Halo-Nwd1 and Flag-SERCA2, Flag-SERCA2-N, or Flag-SERCA2-C were pulled down using HaloLink Resin, and the binding with Halo-Nwd1 was analyzed by WB using anti-Flag antibody. The blot probed with anti-α-tubulin antibody confirmed equal input of each lysate. Scale bars, 5 μm (D, F, H, and I) or 100 nm (E).

Fig. 7. SERCA2 ATPase activity and ER Ca²⁺ storage are suppressed in the *Nwd1⁻ᐟ⁻* liver

(A) SERCA2 ATPase activity in the ER fractions of wild-type and $Nwd1^{-/-}$ livers. (B, C) Ca^{2+} storage in the ER (B) or mitochondrial (C) fractions of wild-type and $Nwd1^{-/-}$ livers. Welch's t-test, ns, not significant; **, $P < 0.01$; ***, $P < 0.001$. (D) Summary of NASH-like hepatic pathogenesis caused by $Nwd1$ deficiency and a model for dysfunction of the SERCA2 Ca^{2+} pump in the $Nwd1^{-/-}$ liver ER. Decreased activity of SERCA2 in the $Nwd1^{-/-}$ liver induces ER stress by reducing Ca^{2+} stores in the ER, leading to diverse hepatic pathologies, including the accumulation of ER-derived vacuolar structures with lipid droplets, fibrosis, and pyroptosis. The ER membranes of $Nwd1^{-/-}$ livers featured an increased population of SERCA2 with no or reduced pump activity (indicated by crosses and thin arrows), lowering Ca^{2+} stores inside the ER.

SUPPLEMENTAL INFORMATION

Fig. S1. Congo red and PAS staining in the $Nwd1^{-/-}$ liver

Congo red staining of wild-type and $Nwd1^{-/-}$ livers. Nuclei were counterstained with hematoxylin. (B) PAS staining with (lower) or without (upper) α-amylase in wild-type and $Nwd1^{-/-}$ livers. (C) Quantitative comparison of the PAS$^+$ area in wild-type and $Nwd1^{-/-}$ livers. Welch's t-test, **, $P < 0.01$. Scale bars, 50 (A) or 100 μm (B).

Fig. S2. The number of peripheral blood leukocytes was increased in $Nwd1^{-/-}$ mice

(A) The relative counts of lymphocytes, neutrophils, eosinophils, and basophils in whole peripheral blood from wild-type or $Nwd1^{-/-}$ adult mice. (B) The ratios (%) of

lymphocytes, monocytes, neutrophils, eosinophils, and basophils between wild-type and *Nwd1*$^{-/-}$ mice.

Fig. S3. HE staining of other organs in *Nwd1*$^{-/-}$ mice

890   (A–G) HE staining of lung, heart, intestine, kidney, skeletal muscle, spleen, and testis tissue from wild-type and *Nwd1*$^{-/-}$ adult mice. Scale bars, 50 (A, C, and F) or 20 μm (B, D, E, and G).

Fig. S4. Autophagy lysosomal pathway dysfunction was not observed in *the*

895   ***Nwd1*$^{-/-}$ liver**

(A, B) *Nwd1*$^{-/-}$ livers were immunostained with anti-LAMP1 (*red*) (A) or anti-LC3A/B (*green*) and anti-KDEL antibodies (*magenta*) (B). Scale bars, 50 μm.

Fig. S5. Validation of ER fractions in *the Nwd1*$^{-/-}$ liver

900   (A) WB of the total lysate and ER fractions of the liver using anti-SERCA2, anti-SEIPIN, anti-KDEL, and anti-α-tubulin antibodies.

Tables S1–S3. Halo-Nwd1-binding proteins identified by proteomic analysis

Each protein entry is presented with the respective protein name, gene symbol, spectral
905   count, and UniProt accession number.

References

[1] S.L. Friedman, B.A. Neuschwander-Tetri, M. Rinella, A.J. Sanyal, Mechanisms of NAFLD development and therapeutic strategies, Nature medicine 24 (2018) 908-922.

910   [2] N.L. Gluchowski, M. Becuwe, T.C. Walther, R.V. Farese Jr, Lipid droplets and liver disease: from basic biology to clinical implications, Nature reviews gastroenterology & hepatology 14 (2017) 343-355.

[3] E. Scorletti, R.M. Carr, A new perspective on NAFLD: Focusing on lipid droplets, Journal of hepatology 76 (2022) 934-945.

915   [4] S. Ben-Moshe, Y. Shapira, A.E. Moor, R. Manco, T. Veg, K. Bahar Halpern, S. Itzkovitz, Spatial sorting enables comprehensive characterization of liver zonation, Nature metabolism 1 (2019) 899-911.

[5] L. Planas-Paz, V. Orsini, L. Boulter, D. Calabrese, M. Pikiolek, F. Nigsch, Y. Xie, G. Roma, A. Donovan, P. Marti, The RSPO–LGR4/5–ZNRF3/RNF43 module controls

920   liver zonation and size, Nature cell biology 18 (2016) 467-479.

[6] W. Birchmeier, Orchestrating Wnt signalling for metabolic liver zonation, Nature cell biology 18 (2016) 463-465.

[7] A. Gonzalez-Rodriguez, R. Mayoral, N. Agra, M. Valdecantos, V. Pardo, M. Miquilena-Colina, J. Vargas-Castrillón, O. Lo Iacono, M. Corazzari, G. Fimia, Impaired

925   autophagic flux is associated with increased endoplasmic reticulum stress during the development of NAFLD, Cell death & disease 5 (2014) e1179-e1179.

[8] C.M. Flessa, I. Kyrou, N. Nasiri-Ansari, G. Kaltsas, A.G. Papavassiliou, E. Kassi, H.S. Randeva, Endoplasmic reticulum stress and autophagy in the pathogenesis of non-alcoholic fatty liver disease (NAFLD): current evidence and perspectives, Current

930   obesity reports 10 (2021) 134-161.

[9] A. Ajoolabady, N. Kaplowitz, C. Lebeaupin, G. Kroemer, R.J. Kaufman, H. Malhi, J. Ren, Endoplasmic reticulum stress in liver diseases, Hepatology 77 (2023) 619-639.

[10] H. Malhi, R.J. Kaufman, Endoplasmic reticulum stress in liver disease, Journal of hepatology 54 (2011) 795-809.

935   [11] V.T. Salo, I. Belevich, S. Li, L. Karhinen, H. Vihinen, C. Vigouroux, J. Magré, C. Thiele, M. Hölttä-Vuori, E. Jokitalo, Seipin regulates ER–lipid droplet contacts and cargo delivery, The EMBO journal 35 (2016) 2699-2716.

[12] E.S. Ali, G.Y. Rychkov, G.J. Barritt, Deranged hepatocyte intracellular Ca2+ homeostasis and the progression of non-alcoholic fatty liver disease to hepatocellular

940   carcinoma, Cell calcium 82 (2019) 102057.

[13] C.-C. Chen, L.-W. Hsu, K.-D. Chen, K.-W. Chiu, C.-L. Chen, K.-T. Huang, Emerging roles of calcium signaling in the development of non-alcoholic fatty liver disease, International journal of molecular sciences 23 (2021) 256.

[14] T. Khamphaya, N. Chukijrungroat, V. Saengsirisuwan, K.A. Mitchell-Richards,

945   M.E. Robert, A. Mennone, M. Ananthanarayanan, M.H. Nathanson, J.

Weerachayaphorn, Nonalcoholic fatty liver disease impairs expression of the type II inositol 1, 4, 5-trisphosphate receptor, Hepatology 67 (2018) 560-574.

[15] Y.G. Zhao, Y. Chen, G. Miao, H. Zhao, W. Qu, D. Li, Z. Wang, N. Liu, L. Li, S. Chen, The ER-localized transmembrane protein EPG-3/VMP1 regulates SERCA activity to control ER-isolation membrane contacts for autophagosome formation, Molecular cell 67 (2017) 974-989. e976.

[16] S.W. Park, Y. Zhou, J. Lee, J. Lee, U. Ozcan, Sarco (endo) plasmic reticulum Ca2+-ATPase 2b is a major regulator of endoplasmic reticulum stress and glucose homeostasis in obesity, Proceedings of the national academy of sciences 107 (2010) 19320-19325.

[17] S. Yamada, S. Sakakibara, Expression profile of the STAND protein Nwd1 in the developing and mature mouse central nervous system, Journal of comparative neurology 526 (2018) 2099-2114.

[18] S. Yamada, A. Sato, S. Sakakibara, Nwd1 regulates neuronal differentiation and migration through purinosome formation in the developing cerebral cortex, Iscience 23 (2020).

[19] D.D. Leipe, E.V. Koonin, L. Aravind, STAND, a class of P-loop NTPases including animal and plant regulators of programmed cell death: multiple, complex domain architectures, unusual phyletic patterns, and evolution by horizontal gene transfer, Journal of molecular biology 343 (2004) 1-28.

[20] S. Yamada, R. Furukawa, S. Sakakibara, Identification and expression profile of novel STAND gene Nwd2 in the mouse central nervous system, Gene expression patterns 46 (2022) 119284.

[21] T. Horii, R. Kobayashi, I. Hatada, Generation of Floxed Mice by Sequential Electroporation, Genome Editing in Animals: Methods and Protocols, Springer2023, pp. 135-147.

[22] H. Matsumura, H. Hasuwa, N. Inoue, M. Ikawa, M. Okabe, Lineage-specific cell disruption in living mice by Cre-mediated expression of diphtheria toxin A chain, Biochemical and biophysical research communications 321 (2004) 275-279.

[23] B. Stielow, F. Finkernagel, T. Stiewe, A. Nist, G. Suske, MGA, L3MBTL2 and E2F6 determine genomic binding of the non-canonical Polycomb repressive complex PRC1. 6, PLoS genetics 14 (2018) e1007193.

[24] S. Sanjabi, K.J. Williams, S. Saccani, L. Zhou, A. Hoffmann, G. Ghosh, S. Gerondakis, G. Natoli, S.T. Smale, A c-Rel subdomain responsible for enhanced DNA-binding affinity and selective gene activation, Genes & development 19 (2005) 2138-2151.

[25] J. Lytton, D. MacLennan, Molecular cloning of cDNAs from human kidney coding for two alternatively spliced products of the cardiac Ca2+-ATPase gene, Journal of Biological chemistry 263 (1988) 15024-15031.

985 [26] W. Ma, C. Mayr, A membraneless organelle associated with the endoplasmic reticulum enables 3′ UTR-mediated protein-protein interactions, Cell 175 (2018) 1492-1506. e1419.

[27] Y. Adachi, T. Kato, T. Yamada, D. Murata, K. Arai, R.V. Stahelin, D.C. Chan, M. Iijima, H. Sesaki, Drp1 tubulates the ER in a GTPase-independent manner, Molecular

990 cell 80 (2020) 621-632. e626.

[28] M.R. Wieckowski, C. Giorgi, M. Lebiedzinska, J. Duszynski, P. Pinton, Isolation of mitochondria-associated membranes and mitochondria from animal tissues and cells, Nature protocols 4 (2009) 1582-1590.

[29] S. Yamada, A. Sato, N. Ishihara, H. Akiyama, S. Sakakibara, Drp1

995 SUMO/deSUMOylation by Senp5 isoforms influences ER tubulation and mitochondrial dynamics to regulate brain development, Iscience 24 (2021).

[30] D.W. Huang, B.T. Sherman, Q. Tan, J. Kir, D. Liu, D. Bryant, Y. Guo, R. Stephens, M.W. Baseler, H.C. Lane, DAVID Bioinformatics Resources: expanded annotation database and novel algorithms to better extract biology from large gene lists,

1000 Nucleic acids research 35 (2007) W169-W175.

[31] D.W. Huang, B.T. Sherman, R.A. Lempicki, Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources, Nature protocols 4 (2009) 44-57.

[32] M. Kanehisa, S. Goto, KEGG: kyoto encyclopedia of genes and genomes, Nucleic

1005 acids research 28 (2000) 27-30.

[33] B.G. Kraupp, B. Ruttkay-Nedecky, H. Koudelka, K. Bukowska, W. Bursch, R. Schulte-Hermann, In situ detection of fragmented DNA (TUNEL assay) fails to discriminate among apoptosis, necrosis, and autolytic cell death: a cautionary note, Hepatology 21 (1995) 1465-1468.

1010 [34] S. Gaul, A. Leszczynska, F. Alegre, B. Kaufmann, C.D. Johnson, L.A. Adams, A. Wree, G. Damm, D. Seehofer, C.J. Calvente, Hepatocyte pyroptosis and release of inflammasome particles induce stellate cell activation and liver fibrosis, Journal of hepatology 74 (2021) 156-167.

[35] T. Bergsbaken, S.L. Fink, B.T. Cookson, Pyroptosis: host cell death and

1015 inflammation, Nature reviews microbiology 7 (2009) 99-109.

[36] D. Li, Y.G. Zhao, D. Li, H. Zhao, J. Huang, G. Miao, D. Feng, P. Liu, D. Li, H. Zhang, The ER-localized protein DFCP1 modulates ER-lipid droplet contact formation, Cell reports 27 (2019) 343-358. e345.

[37] Y. Combot, V.T. Salo, G. Chadeuf, M. Hölttä, K. Ven, I. Pulli, S. Ducheix, C. Pecqueur, O. Renoult, B. Lak, Seipin localizes at endoplasmic-reticulum-mitochondria contact sites to control mitochondrial calcium import and metabolism in adipocytes, Cell reports 38 (2022).

[38] D.G. Mashek, Hepatic lipid droplets: A balancing act between energy storage and metabolic dysfunction in NAFLD, Molecular metabolism 50 (2021) 101115.

[39] K. Yamamoto, K. Takahara, S. Oyadomari, T. Okada, T. Sato, A. Harada, K. Mori, Induction of liver steatosis and lipid droplet formation in ATF6α-knockout mice burdened with pharmacological endoplasmic reticulum stress, Molecular biology of the cell 21 (2010) 2975-2986.

[40] A.S. Lee, The ER chaperone and signaling regulator GRP78/BiP as a monitor of endoplasmic reticulum stress, Methods 35 (2005) 373-381.

[41] G. Zhu, A.S. Lee, Role of the unfolded protein response, GRP78 and GRP94 in organ homeostasis, Journal of cellular physiology 230 (2015) 1413-1420.

[42] B.P. Jain, S. Pandey, WD40 repeat proteins: signalling scaffold with diverse functions, The protein journal 37 (2018) 391-406.

[43] B. Chen, J.E. Mahaney, M.U. Mayer, D.J. Bigelow, T.C. Squier, Concerted but noncooperative activation of nucleotide and actuator domains of the Ca-ATPase upon calcium binding, Biochemistry 47 (2008) 12448-12456.

[44] Y. Zhang, M. Inoue, A. Tsutsumi, S. Watanabe, T. Nishizawa, K. Nagata, M. Kikkawa, K. Inaba, Cryo-EM structures of SERCA2b reveal the mechanism of regulation by the luminal extension tail, Science advances 6 (2020) eabb0147.

[45] H. Kim, T. Kim, B.-C. Jeong, I.-T. Cho, D. Han, N. Takegahara, T. Negishi-Koga, H. Takayanagi, J.H. Lee, J.-Y. Sul, Tmem64 modulates calcium signaling during RANKL-mediated osteoclast differentiation, Cell metabolism 17 (2013) 249-260.

[46] J. Krebs, L.B. Agellon, M. Michalak, Ca2+ homeostasis and endoplasmic reticulum (ER) stress: An integrated view of calcium signaling, Biochemical and biophysical research communications 460 (2015) 114-121.

[47] J.D. Malhotra, R.J. Kaufman, The endoplasmic reticulum and the unfolded protein response, Seminars in cell & developmental biology, 2007, pp. 716-731.

[48] C. Lebeaupin, D. Vallée, Y. Hazari, C. Hetz, E. Chevet, B. Bailly-Maitre, Endoplasmic reticulum stress signalling and the pathogenesis of non-alcoholic fatty liver disease, Journal of hepatology 69 (2018) 927-947.

[49] S. Martin, R.G. Parton, Lipid droplets: a unified view of a dynamic organelle, Nature reviews Molecular cell biology 7 (2006) 373-378.

[50] K. Tauchi-Sato, S. Ozeki, T. Houjou, R. Taguchi, T. Fujimoto, The surface of lipid droplets is a phospholipid monolayer with a unique fatty acid composition, Journal of Biological chemistry 277 (2002) 44507-44512.

[51] J.I. Beier, J.M. Banales, Pyroptosis: An inflammatory link between NAFLD and NASH with potential therapeutic implications, Journal of hepatology 68 (2018) 643-645.

[52] A.R. Mridha, A. Wree, A.A. Robertson, M.M. Yeh, C.D. Johnson, D.M. Van Rooyen, F. Haczeyni, N.C.-H. Teoh, C. Savard, G.N. Ioannou, NLRP3 inflammasome blockade reduces liver inflammation and fibrosis in experimental NASH in mice, Journal of hepatology 66 (2017) 1037-1046.

[53] J. Lytton, M. Westlin, S.E. Burk, G.E. Shull, D.H. MacLennan, Functional comparisons between isoforms of the sarcoplasmic or endoplasmic reticulum family of calcium pumps, Journal of Biological Chemistry 267 (1992) 14483-14489.

[54] J.B. French, S.A. Jones, H. Deng, A.M. Pedley, D. Kim, C.Y. Chan, H. Hu, R.J. Pugh, H. Zhao, Y. Zhang, Spatial colocalization and functional link of purinosomes with mitochondria, Science 351 (2016) 733-737.

[55] R.G. Correa, M. Krajewska, C.F. Ware, M. Gerlic, J.C. Reed, The NLR-related protein NWD1 is associated with prostate cancer and modulates androgen receptor signaling, Oncotarget 5 (2014) 1666.

Fig. 1

A

CDS:302-4867bp NM_176940.5

Nwd1 wild *allele*

1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19

F1 R1

4569-7722

crRNA — 4 ✕ 5 ✕ 6

Nwd1L3   Nwd1R1

ssODN

Nwd1L3loxpAS   Nwd1R1loxp

Nwd1^{flox}

loxp   loxp

4 | 5 | 6

×CAG-Cre

Nwd1[−]

loxp

4 | 6

F1 R2

B

wild   *Nwd1*^{flox/+}

bp

F1–R1

C

wild   *Nwd1*^{+/−}   *Nwd1*^{−/−}

bp

F1–R1

F1–R2

600
500
400

D

1.5

Nwd1 mRNA expression

1.0

0.5

wild   *Nwd1*^{-/-}

E

Parental cross: *Nwd1*^{+/−}× *Nwd1*^{+/−}	*Nwd1*^{+/+}	*Nwd1*^{+/−}	*Nwd1*^{−/−}	total	*p*
Number of observed (expected number)	57 (52.75)	116 (105.5)	38 (52.75)	211	0.064
Birth rate (%) (expected rate)	27.01(25)	54.98 (50)	18.01 (25)	100	

F

wild   *Nwd1*^{−/−}

G

Liver weight (g)

wild   *Nwd1*^{-/-}

**

H

Body weight (g)

wild   *Nwd1*^{-/-}

ns

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

A Halo-Nwd1

total lysate / purification

kDa

B

Protein	Gene Symbol	Spectral Count	Accession
Sarcoplasmic/endoplasmic reticulum calcium ATPase 2	ATP2A2 ATP2B SERCA2	411	P16615
Tubulin beta chain	TUBB	400	P07437
Voltage-dependent anion selective channel protein 1	VDAC1	278	P21796
Calnexin	CANX	261	P27824
Tubulin alpha-1B chain	TUBA1B	258	P68363
Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 48 kDa subunit	DDOST	229	P39656
Sideroflexin-1	SFXN1	206	Q9H9B4
Dolichyl-diphosphooligosaccharide-protein glycosyltransferase subunit 1	RPN1	199	P04843
NADH-cytochrome b5 reductase 3	CYB5R3	197	P00387
Sodium/potassium-transporting ATPase subunit alpha-1	ATP1A1	189	P05023
DNA-dependent protein kinase catalytic subunit	PRKDC	182	P78527
Ras-related protein Rab-7a	RAB7A	181	P51149
Prohibitin-2	PHB2	173	Q99623
Translocation protein SEC63 homolog	SEC63	167	Q9UGP8
Very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase 3	HACD3	167	Q9P035
Heat shock 70 kDa protein 1A	HSPA1A	164	P0DMV8
Membrane-associated progesterone receptor component 1	PGRMC1	160	O00264

C GO funcitonal analysis

-log10(P-value)

Biological process (red):
ER to Golgi vesicle-mediated transport; Golgi organization; protein transport; vesicle-mediated transport; protein N-linked glycosylation via asparagine; intracellular protein transport; cellular response to steroid hormone stimulus

Cellular component (green):
ER membrane; membrane; ER; integral component of membrane; melanosome; extracellular exosome; ER-Golgi intermediate compartment membrane; ER to Golgi transport vesicle; ER-Golgi intermediate compartment; mitochondrion

Molecular function (blue):
protein binding; GTPase activity; GTP binding; GDP binding; mannose binding; enzyme binding

KEGG pathway (yellow):
Protein processing in ER; Prion disease; N-Glycan biosynthesis; Phagosome; Alzheimer disease

D Nwd1 / KDEL / Merged — Flag-Nwd1

E Flag — Flag-Nwd1 — ER, MT, ER

F mCherry-Sec61B — Flag / mCherry / Merged — Flag-Nwd1, Flag-Nwd1-N, Flag-Nwd1-N, Flag-Nwd1-C

G SERCA2b protein domain structure (1044aa)

SERCA2b-N(1–787aa) / SERCA2b-C(788–1044aa)

P-domain / N-domain / A-domain

1–48 / 111–253 / 314–756 / 808–827 / 917–929 / 994–1012
70–89 / 274–295 / 777–786 / 851–896 / 949–963 / 1031–1044

Cytosolic region / Transmembrane domain / Luminal region

H EGFP / SERCA2 / Merged — EGFP-Nwd1

I EGFP-Nwd1 — EGFP / Flag / Merged — Flag-SERCA2, Flag-SERCA2-N, Flag-SERCA2-C

J

	Input					HaloTag Regin				
Halo-Nwd1	+	+	+	+	+	+	+	+	+	-
Flag-control	+	-	-	-	-	+	-	-	-	-
Flag-SERCA2	-	+	-	-	+	-	+	-	-	+
Flag-SERCA2-N	-	-	+	-	-	-	-	+	-	-
Flag-SERCA2-C	-	-	-	+	+	-	-	-	+	+

kDa

WB:Flag

WB:α-tubulin — 45

Fig. 7

A

Relative SERCA2 ATPase activity

**

wild $Nwd1^{-/-}$

B

Relative ER Ca²⁺ storage

***

wild $Nwd1^{-/-}$

C

Relative mitochondria Ca²⁺ storage

ns

wild $Nwd1^{-/-}$

D

wild liver

NASH-like pathogenesis in $Nwd1^{-/-}$ liver

Cytosol

Ca^{2+} Ca^{2+} SERCA2 Ca^{2+}

ER

Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+}

Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+} Ca^{2+}

· **Hepatomegaly** ⬆
· **Lipid droplet** ⬆
· **ER vesicle** ⬆
· **Fibrosis** ⬆
· **Pyroptosis** ⬆
· **ER stress** ⬆
· **SERCA2 ATPase activity** ⬇
· **Ca²⁺ storage** ⬇