Prolactin-mediates a lactation-induced suppression of arcuate kisspeptin neuronal

activity necessary for lactational infertility in mice

Eleni C.R. Hackwell[1,2], Sharon R. Ladyman[1,2,4], Jenny Clarkson[1,3], H. James McQuillan[1,2],

Ulrich Boehm[5], Allan E. Herbison[6], Rosemary S.E. Brown[1,3], David R. Grattan[1,2,4]

[1]Centre for Neuroendocrinology, [2]Department of Anatomy and [3]Department of Physiology,

School of Biomedical Sciences, University of Otago, Dunedin, NZ; [4]Maurice Wilkins Centre

for Molecular Biodiscovery, Auckland, NZ; [5]Saarland University School of Medicine, Centre

for Molecular Signalling (PZMS), Experimental Pharmacology, Homburg, Germany;

[6]Department of Physiology, Development and Neuroscience, University of Cambridge,

Cambridge, United Kingdom.

Corresponding author: David R. Grattan

Email: dave.grattan@otago.ac.nz

Author Contributions:

ECRH: Designed research, completed research, analysed data, wrote the first draft of the
manuscript, edited the manuscript.
SRL: Designed research, completed research, analysed data, edited the manuscript.
JC: Completed research, analysed data, edited the manuscript.
HJM: Completed research, edited the manuscript.
UB: Provided key resources, edited the manuscript.
AEH: Designed research, analysed data, edited the manuscript.
RSEB: Designed research, analysed data, edited the manuscript.
DRG: Designed research, provided key resources, analysed data, edited the manuscript.
25
26 **Competing Interest Statement:** The authors have no competing interests to declare.

Abstract

The specific role that prolactin plays in lactational infertility, as distinct from other suckling or metabolic cues, remains unresolved. Here, deletion of the prolactin receptor (Prlr) from forebrain neurons or arcuate kisspeptin neurons resulted in failure to maintain normal lactation-induced suppression of estrous cycles. Kisspeptin immunoreactivity and pulsatile LH secretion were increased in these mice, even in the presence of ongoing suckling stimulation and lactation. GCaMP fibre photometry of arcuate kisspeptin neurons revealed that the normal episodic activity of these neurons is rapidly suppressed in pregnancy and this was maintained throughout early lactation. Deletion of Prlr from arcuate kisspeptin neurons resulted in early reactivation of episodic activity of kisspeptin neurons prior to a premature return of reproductive cycles in early lactation. These observations show dynamic variation in arcuate kisspeptin neuronal activity associated with the hormonal changes of pregnancy and lactation, and provide direct evidence that prolactin action on arcuate kisspeptin neurons is necessary for suppressing fertility during lactation in mice.

Introduction

In mammals, lactation is accompanied by a period of infertility. This adaptive change establishes appropriate birth spacing to enable maternal metabolic resources to be directed towards caring for the new-born offspring, rather than supporting another pregnancy (1). Lactational infertility is characterized by a lactation-induced suppression of pulsatile luteinizing hormone (LH) secretion, and the temporary loss of the reproductive cycle (in rodents this is exhibited as an extended period of diestrus or anestrus) (2-5). Lactation is also characterised by chronically elevated levels of the anterior pituitary hormone prolactin, which is essential for milk production and promotes adaptive changes in maternal physiology and behaviour (1,2,4,5). Despite hyperprolactinaemia being a well-recognized cause of infertility, the specific role that prolactin plays in lactational infertility, as distinct from other suckling- or metabolic-related cues, is currently unclear (4,6).

Recent *in vivo* studies have confirmed that kisspeptin neurons in the arcuate nucleus of the hypothalamus are responsible for the periodic release of gonadotrophin-releasing hormone (GnRH) and subsequent pulsatile LH secretion that drives reproductive function (7-11). Studies using GCaMP fibre photometry in conscious mice have demonstrated that the arcuate kisspeptin neuronal population exhibits episodes of increased intracellular calcium levels coincident with, and immediately preceding, each pulse of LH secretion in intact and gonadectomised male and female mice (7-9). Miniscope investigations showed that individual kisspeptin neurons within the arcuate population act in a coordinated, synchronised, and episodic manner (10,11). Loss of pulsatile LH secretion during lactation and consequent lactational infertility may be caused by the loss of kisspeptin-mediated stimulation of GnRH secretion (12-18). Kisspeptin expression is markedly suppressed in lactation (12,16) and even when exogenously stimulated, kisspeptin neurons are unable to activate GnRH neurons during lactation, likely due to a lack of kisspeptin synthesis (13).

It is well established that hyperprolactinemia causes infertility, and thus, the elevated prolactin present in lactation seems a likely candidate to be involved in suppressing fertility during lactation. Prolactin administration acutely suppresses LH secretion (19), and chronic exposure to elevated prolactin reduces *Kiss1* mRNA expression in the arcuate nucleus (17,20,21). In lactating mice, suppressing endogenous prolactin secretion shortens the period of infertility (22), suggesting that prolactin is important for maintaining the suppression of pulsatile LH secretion during lactation. Such a role for prolactin is controversial, however

(4,6,23-27), with studies in a number of species suggesting that the neural stimulation of

suckling may be more important than prolactin in maintaining lactational infertility (28,29).

However, it has previously been difficult to disentangle the specific role of prolactin, as

suckling, prolactin, and milk production are so tightly linked that manipulating one ultimately

impacts the others, making it difficult to determine the contribution of any one element. Here,

using a conditional deletion strategy, we have blocked prolactin action in the brain leaving

suckling, lactation, and maternal behaviour intact. Using GCaMP fibre photometry

techniques, we have also documented arcuate kisspeptin neuron activity across pregnancy

and lactation transitions in the same mice and established that prolactin directly acts on these

neurons to suppress fertility in lactation.

Methods

Animals

All experiments were performed using adult female mice on a C57BL/6J background (8-20 weeks of age unless otherwise stated). Mice were housed under controlled temperature (22 °C ± 2°C) and lighting (12-hour light/12-hour dark schedule, with lights on at 0600 hours) with *ad libitum* access to food and water (Teklad Global 18% Protein Rodent Diet 2918; Envigo, Huntingdon, United Kingdom). Daily body weight was recorded and daily vaginal cytology was used to monitor the estrous cycle stage. All experiments were carried out with approval from the University of Otago Animal Welfare and Ethics Committee.

To establish pregnancies, mice were mated with male wild-type C57BL/6J mice (presence of sperm plug = day 1 pregnancy). Male mice were then removed once a sperm plug was seen, and no male mice were present at parturition. The first day a litter was seen was counted as day 1 of lactation and maternal mice were left undisturbed till day 3 of lactation, when vaginal monitoring would resume and litter size was normalised to 6 pups per animal, unless otherwise stated. The intention of the normalisation was to reduce variability in the degree of suckling stimulus each dam would receive.

Towards the end of each recording period, bilateral ovariectomy (OVX) was performed under isoflurane anaesthesia with pre- and post-operative Carprofen (5mg/kg, s.c.) administered for pain relief. This enabled us to examine post-OVX changes in activity of kisspeptin neurons and expression of kisspeptin in the arcuate nucleus.

Pulsatile hormone measurement

To monitor pulsatile secretion of LH, serial tail tip blood sampling and measurement of LH by ELISA was undertaken as reported previously (19,30,31). As novel exposure and restraint stress has been shown to suppress pulsatile LH secretion (32), all mice were habituated to the tail tip blood sampling procedure by picking up the mouse in a gentle restraint device (soft cardboard tube) or hand and lightly massaging the tail for approximately 5 minutes per day for at least 3 weeks prior to experimentation (33). Sequential whole blood samples (4μl) were collected in 6 minute intervals for 3 hours between 0900 and 1200 hours, unless otherwise stated. Samples were immediately diluted in 48ul 0.01M PBS/0.05% Tween 20, and frozen on dry ice before being stored at -20°C for subsequent LH measurement.

RNAscope

Mice were transcardially perfused with a micro-perfusion pump with 2% paraformaldehyde
(PFA). Brain sections (14μm-thick) were prepared, thaw mounted onto superfrost-plus
microscope slides and then stored at -80°C. RNAscope in-situ hybridization was performed
using the RNAscope 2.5 High definition Duplex Detection kit – chromogenic (Advanced Cell
Diagnostics, Hayward, CA) largely in accordance with manufacturer's instruction. The
channel 1 Prlr probe was custom designed to pick up only the long form of the prolactin
receptor. It was designed to transcript NM_011169.5 with a target sequence spanning
nucleotides 1107-2147 (Ref: 588621; Advanced Cell Diagnostics, Hayward, CA). The
channel 2 Kiss1 probe was custom designed to transcript NM_178260.3 with a target
sequence spanning nucleotides 5 to 485 (Ref: 500141-C2; Advanced Cell Diagnostics,
Hayward, CA). Sections were thawed at 55°C, postfixed for 3 minutes in 2% PFA, washed in
0.01M PBS for 5 minutes, and endogenous peroxidases were blocked with a hydrogen
peroxidase solution for 10 minutes. Tissue was washed in distilled water (3x 2 minutes), then
briefly immersed in 100% ethanol, air dried for 5 minutes, and a hydrophobic barrier was
applied. Tissue was permeabilized with RNAscope protease plus for 30 minutes at 40°C.
Sections were washed (2 x 2 minutes) and were hybridized with the Prlr and Kiss1 probes
(1:300 dilution, Prlr:Kiss1) or negative control probe (Cat#320751; Advanced Cell
Diagnostics, Hayward, CA) at 40°C for 2 hours. Amplification (Amp 1-6) was performed in
accordance with the manufacturer's instructions. Sections were then hybridized with a Fast-
RED (1:60, Fast-RED B:Fast-RED A) for 10 minutes at room temperature, before
undergoing further amplification steps (Amp 7-10) in accordance to manufacturer's
instructions. The final positive hybridization was detected by incubation with the secondary
detection reagents (1:50, Fast-GREEN B:Fast-GREEN) for 10 minutes at room temperature.
Sections were washed, counterstained with haematoxylin (25% Gills), dried at 60°C for 20
minutes, and cover-slipped with VectaMount (Vector laboratories, H-5000) before imaging
using an Olympus BX51 light microscope and Olympus UPlanSApo 10/20x lenses.

Quantification of the proportion of kisspeptin neurons co-expressing *Prlr* mRNA was
undertaken in FIJI software (National Institute of Health, Bethesda, Maryland, USA)
following image acquisition. Positive hybridization for *Kiss1* and *Prlr* was determined based
on scoring guidelines included in the manufacturer's protocol (Advanced Cell Diagnostics).

146   The total number of *Kiss1*-expressing cells and the total number of these that showed *Prlr*

147   mRNA expression were counted.

148   Effect of neuron-specific deletion of the prolactin receptor gene on the maintenance of

149   lactational infertility

To investigate whether prolactin action in the brain is required for lactational infertility,

neuron-specific *Prlr* knockout mice ($Prlr^{lox/lox}/Camk2a^{Cre}$) and their respective Cre-negative

controls ($Prlr^{lox/lox}$) were generated (as previously described in (34)). We have previously

shown that while $Prlr^{lox/lox}/Camk2a^{Cre}$ mice do not have a complete Prlr deletion in the

forebrain. There are areas of extensive deletion (as measured by reduced prolactin-induced

pSTAT5), such as the arcuate nucleus and ventromedial nucleus of the hypothalamus, and

there are also areas where Prlr is reduced by about 50% such as the medial pre-optic area

(34,35). In our experience with these animals, this is sufficient to retain normal maternal

behaviour in most animals (our unpublished data), and this was the case for the animals used

in the present study. RNAscope in-situ hybridization was done to confirm the degree of

knockout in intact, diestrous, nulligravid, non-lactating (henceforth termed NL) mice and 14

day post-OVX mice ($Prlr^{lox/lox}$ intact n = 6, $Prlr^{lox/lox}$ OVX n = 5, $Prlr^{lox/lox}/Camk2a^{Cre}$ intact n

= 5, $Prlr^{lox/lox}/Camk2a^{Cre}$ OVX n = 5; all aged 8-16 weeks). Both intact and OVX mice were

included as kisspeptin cell bodies are only visible in the rostral periventricular region of the

third ventricle (RP3V) of intact mice and in the arcuate nucleus of OVX mice, due to

estradiol regulation (36). $Prlr^{lox/lox}/Camk2a^{Cre}$ mice showed a significant decrease in the

percentage of *Kiss1*-expressing cells co-expressing *Prlr* compared to controls in both the

RP3V (p = <0.0001) and arcuate nucleus (p = 0.0009) (unpaired two-tailed t tests,

Supplementary Figure 1A-D).

$Prlr^{lox/lox}/Camk2a^{Cre}$ mice are hyperprolactinaemic due to impaired negative feedback of

prolactin on hypothalamic dopamine neurons (34) and therefore show disrupted estrous

cycles (showing recurrent pseudopregnancy-like cycles with long periods of diestrus of

approximately 14 days between estrous stages). However, these mice are able to become

pregnant and have normal pregnancies. All $Prlr^{lox/lox}/Camk2a^{Cre}$ mice were given a 250μl

subcutaneous injection of bromocriptine (5mg/kg, 5% ethanol/saline; Tocris Bioscience

Cat#0427) prior to being mated. This treatment was designed to reinstate an estrous cycle in

$Prlr^{lox/lox}/Camk2a^{Cre}$ mice. Bromocriptine is an agonist for the type 2 dopamine receptor and

inhibits prolactin secretion from the pituitary gland (37), thereby terminating the

pseudopregnancy-like state and bringing the mice into proestrus the following day. Following

treatment, all mice were then housed with a stud male.

For $Prlr^{lox/lox}/Camk2a^{Cre}$ mice, estrous cycles were monitored from day 3 of lactation until the

first day of diestrus following a day of estrus (proestrus and estrus had to be observed prior to

transcardial perfusion on the first day of diestrus). Brains were collected following

transcardial perfusion for assessment of kisspeptin immunoreactivity. For every lactating

$Prlr^{lox/lox}/Camk2a^{Cre}$ mouse (n = 8), the brain of a $Prlr^{lox/lox}$ control mouse (n = 8) of the

equivalent day (±1) of lactation was also collected. A group of NL mice of both genotypes (n

= 5-6) was also perfused for immunohistochemistry on diestrus.

To evaluate pulsatile LH secretion in early lactation (prior to the return of estrous cycles) and

to determine whether progesterone played any role in regulating pulsatile LH secretion in

lactation, additional groups of lactating $Prlr^{lox/lox}/Camk2a^{Cre}$ and $Prlr^{lox/lox}$ control mice were

generated. These mice were treated with either the progesterone receptor antagonist,

mifepristone (4mg/kg in sesame oil, s.c.; AK Scientific Inc Cat#J10622), or vehicle (n = 7-8

per group) on the morning of day 4 of lactation and day 5 of lactation. This dose was selected

as it was found to be sufficient to cause termination of pregnancy in wild-type C57BL/6J

mice (p = 0.0072, chi-squared test, Supplementary Figure 2A; pilot study, n = 6 both groups).

Neither vehicle nor mifepristone treatment had an effect on litter weight gain (interaction of

time x genotype & treatment p = 0.5322, two-way repeated measures ANOVA,

Supplementary Figure 2B). Mice underwent blood sampling for LH for 3 hours on day 5 of

lactation, 30 minutes after treatment.

Measurement of LH concentrations

An established sandwich ELISA method was used to determine LH concentration in diluted

whole blood samples collected from mice (30,31). Briefly, a 96-well high plate was

incubated with bovine monoclonal antibody (LHβ518b7, 1:1000 in 1xPBS; Dr. L. Sibley, UC

Davis, CA, USA) for 16 h at 4°C. Following incubation of standards, controls and

experimental samples for 2 hours, plates were incubated in rabbit polyclonal LH antibody

(AFP240580Rb; 1:10,000; National Hormone and Pituitary Program, NIH) for 90 minutes,

followed by incubation with polyclonal goat anti-rabbit IgG/HRP antibody (1:1000; DAKO

Cytomation) for 90 minutes. Finally, plates were incubated in OPD (o-phenylenediamine

capsules; Sigma-Aldrich Cat#P7288) for 30 minutes. A standard curve for the detection of

LH concentration was generated using serial dilutions of mouse LH-reference preparation

peptide (National Hormone and Pituitary Program, NIH). LH levels were read using a

standard absorbance plate reader (SpectraMax ABS Plus; Molecular Devices) at 490nm and

630nm wavelengths. The assay had a sensitivity of 0.04ng/ml to 4ng/ml, with an intra-assay

coefficient of variation of 4.40% and an inter-assay coefficient of variation of 8.29%.

PULSAR Otago was used to define LH pulses (38). Parameters used; Smoothing 0.7, Peak

split 2.5, Level of detection 0.04, Amplitude distance 3, Assay variability 0, 2.5, 3.3,

$G(1)=3.5$, $G(2)=2.6$, $G(3)=1.9$, $G(4)=1.5$, $G(6)=1.2$. Mean LH levels were calculated by

averaging all LH levels collected during the experiment. Individual LH profiles from mice

are shown in Supplementary Figure 3.

Assessment of kisspeptin expression
220
Perfusion and fixation of tissue

Mice were anaesthetised with sodium pentobarbital (15mg/mL, i.p.) and transcardially

perfused with 4% PFA. Brains were removed, postfixed in the same solution, and

cryoprotected overnight in 30% sucrose before being frozen at -80°C. Two sets of 30μm

thick coronal brain sections were cut using a sliding microtome, from Bregma 1.10mm to -

2.80mm. Brain sections were kept in cryoprotectant solution (pH = 7.6) at -20°C until

immunohistochemistry was performed.

Immunohistochemistry

Immunohistochemistry for kisspeptin in the RP3V and arcuate nucleus was performed as

previously described (39). Briefly, sections were incubated in polyclonal rabbit anti-

kisspeptin primary antibody (AC 566, 1:10,000; gift from A. Caraty, Institut National de la

Recherche Agronomique, Paris, France) for 48 hours at 4°C. Sections were then incubated

with biotinylated goat anti-rabbit IgG (1:200, Vector biolabs, Peterborough, GK) for 90 min

at room temperature, followed by incubation in an avidin-biotin complex (Elite vectastain

ABC kit, Vector laboratories). The bound antibody-peroxidase complex was visualised using

a nickel-enhanced diaminobenzidine (DAB) reaction, to form a black cytoplasmic precipitate.

Brain sections were imaged as described in RNAscope section above. Quantification of

kisspeptin neurons in the RP3V, was undertaken by manually counting all labelled neurons

present in all three subdivisions: the anteroventral periventricular nucleus (AVPV); rostral

preoptic periventricular nucleus (rPVpo); and caudal preoptic periventricular nucleus

(cPVpo). Two sections per subdivision per mouse were averaged for each animal. As

kisspeptin cell bodies in the arcuate nucleus were not easily observed, as previously reported

(40), kisspeptin fibre immunoreactivity was imaged using a Gryphax NAOS colour camera

(Jenoptik) and evaluated using FIJI software and the voxel counter function (National

Institutes of Health). Kisspeptin fibre density was measured in the arcuate nucleus across the

three subdivisions; rostral arcuate (rARC), middle arcuate (mARC), and caudal arcuate

(cARC). Two sections of each subdivision per animal were counted and then averaged across

each animal to get total number and reported as total amount of voxels per ROI (voxel

fraction).

Characterization of arcuate kisspeptin neuronal activity using GCaMP fibre

photometry

Stereotaxic surgery and AAV injections

Adult $Kiss1^{Cre}$ or $Prlr^{lox/lox}/Kiss1^{Cre}$ mice (2-3 months old) were anaesthetised with 2%

Isoflurane, given local Lidocaine (4mg/kg, s.c.) and Carprofen (5mg/kg, s.c.) and placed in a

stereotaxic apparatus. A custom-made unilateral Hamilton syringe apparatus holding one

Hamilton syringe was used to perform unilateral injections into the arcuate nucleus. The

needles were lowered into place (-0.14mm A/P, +0.04mm M/L, -0.56mm DV) over 2

minutes and left in situ for 3 minutes before injection was made. 1μl AAV9-CAG-FLEX-

GCaMP6s-WPRE-SV40 (1.3×10^{-13} GC/ml, University of Pennsylvania Vector Core,

Philadelphia, PA, USA) was injected into the arcuate nucleus at a rate of ~100nl/min with the

needles left in situ for 3 minutes prior to being withdrawn over a period of 6 minutes. This

was followed by implantation of a unilateral indwelling optical fibre (400 μm diameter, 6.5

mm long, 0.48 numerical aperture (NA), Doric Lenses, Canada, product code:

MFC_400/430-0.48_6.5mm_SM3*_FLT) at the same coordinates. Carprofen (5mg/kg body

weight, s.c.) was administered for post-operative pain relief. After surgery, mice received

daily handling and habituation to the fibre photometry recording conditions over 4-6 weeks

before experimentation began.

GCaMP fibre photometry

Photometry was performed as reported previously (9). Fluorescence signals were acquired

using a custom-built fibre photometry system made primarily from Doric components. Violet

(405nm) and blue (490nm) fibre-coupled LEDs were sinusoidally modulated at 531 and 211

Hz, respectively, and focused into a 400μm, 0.48 numerical aperture fibre optic patch cord

connected to the mouse. Emitted fluorescence was collected by the same fibre and focused

onto a femtowatt photoreceiver (2151, Newport). The two GCaMP6s emission signals were

collected at 10 Hz in a scheduled 5s on/15s off mode by demodulating the 405nm (non-

calcium dependent) and 490nm (calcium dependent) signals. The power output at the tip of

the fibre was set at 50μW. Fluorescent signals were acquired using a custom software

acquisition system (Tussock Innovation, Dunedin, New Zealand) and analysed using custom

templates created by Dr Joon Kim (University of Otago, Dunedin, New Zealand) based on

mathematics and calculations similar to those previously described (41,42). Briefly, the

fluorescent signal obtained after stimulation with 405nm light was used to correct for

movement artefacts as follows: first, the 405nm signal was filtered using a savitzky-golay

filter and fitted to the 490nm signal using least linear square regression. The fitted 405nm

signal was then subtracted and divided from the 490nm signal to obtain the movement and

bleaching corrected signal. The output of these templates is 490-adjusted405/adjusted405,

which was multiplied to get the final ΔF/F as a percentage increase. All photometry data is

reported as ΔF/F(%).

All recordings were obtained from freely behaving mice between 0800 hours and 1200 hours

(apart from 24 hours post weaning recording (0900 hours to 1700 hours), and day 18/19

pregnancy recording (1800 hours to 0800 hours the following day)). Synchronized events

(SEs) were defined as when ΔF/F exceeds 3 standard deviations (SD) above the trace mean.

Manual SE shape analysis was performed in addition to standard deviation method for certain

datasets where necessary. SEs were counted manually to determine frequency of SEs per 60

minutes. The between-animal variability in total signal means that changes in SE amplitude

can only be reported as relative changes within an animal. Relative SE amplitude was

calculated by using normalised ΔF/F data and then subtracting the peak of an SE from the

nearest nadir to the rise of the SE and averaging that for the number of SEs in a recording. To

obtain normalised ΔF/F, three pre-pregnancy datasets from each mouse were used to find the

average maximum ΔF/F for that mouse. All datasets were then divided by that normalisation

value to get normalised ΔF/F for each trace for each individual mouse. In these long-term

longitudinal studies, some data points are missing due to issues with recordings (e.g. dirty

fibre optic cannula or system battery issues), meaning that occasionally fewer mice are used

for analysis at various time points. During experimentation, no mice lost cannulas, but a

gradual reduction in amplitude of signal was observed over time.

Following experimentation, brains were collected in a similar way to that described above.

Postmortem analysis of cannula placement and GCaMP6 transfection in each animal was not

done, because with experience, the quality and characteristics of the SE recordings (including

corresponding pulsatile LH secretion) in each animal was indicative of successful cannula

placement and transfection of the GCaMP. In a preliminary trial the green fluorescent protein

(GFP) attached to the GCAMP was restricted to the caudal arcuate nucleus (n = 3 mice,

preliminary trial), with little evidence of spread into the rostral or middle arcuate. GFP was

seen exclusively ipsilaterally, with no GFP observed on the contralateral side. Cannula

placement in these trial animals was in the caudal arcuate.

Monitoring the activity of arcuate kisspeptin neurons across different reproductive stages in the same mice

Adult *Kiss1*^Cre mice were 8-10 weeks of age at the beginning of experiments, and up to 12

months old by the time of final recording (n = 8 during pregnancy; n = 6 during lactation, 2

mice were euthanised due to dystocia). Monitoring of vaginal cytology and weight was

continuous from 1 week pre-surgery till day 19 of pregnancy and resumed on day 3 of

lactation until the end of experiment. All handling stopped on day 19 of pregnancy to avoid

potential compromise of parturition and onset of maternal behaviour.

To investigate the activity of the arcuate kisspeptin population across different reproductive

states in the same animal, the following recording protocol was followed for all *Kiss1*^Cre

mice, unless otherwise stated; diestrus NL, day 4 of pregnancy, day 14 of pregnancy, day

18/19 of pregnancy (overnight), day 7 of lactation, day 14 of lactation, day 18 of lactation, 24

hours after weaning, first diestrus after estrous cycles begin following weaning, and 10 days

post-OVX. In addition, blood sample collection for paired LH measurement was done in

diestrus NL state, on day 14 of pregnancy (in 4 mice, maximum of 6 samples were collected

around small "peaks" in baseline), day 7 of lactation, and day 14 of lactation. During

sampling, blood samples were taken at the usual 6 minute intervals previously described,

however if a SE was observed and the next scheduled sampling was more than 3 minutes

away, an extra blood sample was taken 2 minutes after the SE. Following this additional

sampling, the next scheduled blood sample would be taken at its original time and then every

6 minutes after, till another SE was seen. Blood sampling was not carried out at additional

time points as blood sampling was undertaken at least a week apart, and stress from repeated

sampling was attempted to be kept at a minimum e.g., not blood sampling around the time of

birth. Fibre photometry recordings were usually between 2-4 hours in length. The only longer

recordings were undertaken on day 18/19 of pregnancy (14 hours) and 24 hours after

weaning (8 hours). These particular recording sessions were extended to determine whether

there were any longer-term changes occurring in the activity of the arcuate kisspeptin

population in the lead up to parturition or following weaning of pups (states closely followed

by postpartum estrus and resumption of normal estrous cycles, respectively).

Effect of arcuate kisspeptin neuron-specific deletion of the prolactin receptor gene on the

maintenance of lactational infertility and the activity of kisspeptin neurons during

lactation

Kisspeptin-specific prolactin-receptor knockout mice ($Prlr^{lox/lox}/Kiss1^{Cre}$) and their respective

Cre-negative controls ($Prlr^{lox/lox}$) were generated as previously described (19).

$Prlr^{lox/lox}/Kiss1^{Cre}$ (n = 27) and $Prlr^{lox/lox}$ control (n = 30) dams underwent estrous cycle

monitoring from day 3 of lactation onwards to determine whether mice showed an early

resumption of estrous cycles. RNAscope in-situ hybridization (as described above) was done

to confirm knockdown in a subgroup of these mice, with $Prlr^{lox/lox}/Kiss1^{Cre}$ mice showing a

significant decrease in the percentage of $Kiss1$-expressing cells co-expressing $Prlr$ compared

to controls in the arcuate nucleus (p = <0.0001, unpaired two-tailed t test, Supplementary

Figure 1E-G; $Prlr^{lox/lox}$ n = 6, $Prlr^{lox/lox}/Kiss1^{Cre}$ n = 8).

To determine whether the deletion of the prolactin receptor from arcuate kisspeptin neurons

led to early reactivation of these neurons during lactation, adult $Prlr^{lox/lox}/Kiss1^{Cre}$ mice (n =

5) and an additional $Kiss1^{Cre}$ control mouse (n = 1) (8 weeks old at the start of the

experiment, and up to 14 months at final recording timepoint) were set up for fibre

photometry, as described above. Monitoring of vaginal cytology and weight was continuous

from 1 week pre-surgery till day 18 of pregnancy and resumed on day 3 of lactation.

Recordings were undertaken in a similar timeline to that described above, however no

pregnancy recordings were done, and in early lactation recordings were performed every 2

days from day 3 to day 9 of lactation, before following the same protocol as described. No

blood samples were taken over the lactation period in these mice. As described previously,

recordings were kept between 2-4 hours, apart from the 24 hours after weaning recording (8

hours).

Statistical analysis

Data are presented as mean ± SEM and all statistical analysis was performed with PRISM

software 10 (GraphPad Software, San Diego, CA, USA) with a p value of < 0.05 considered

as statistically significant. Individual symbols in graphs represent individual mice.

Differences in kisspeptin cell number or fibre density was assessed using two-way ANOVAs

with Tukey's multiple comparisons tests or t tests, with both analyses using combined

averages of each animal (averaged number of cells or fibre density across the three

subdivisions of each nucleus to get total number reported). Resumption of estrous cycles was

analysed using Log-rank (Mantel-Cox) test chi-squared test. LH pulse frequency data and

mean LH data was analysed using two-way ANOVAs with Tukey's multiple comparisons

tests. SE frequency and amplitude throughout reproductive cycles was analysed using mixed

effect analysis (fixed type III) with Tukey's multiple comparisons tests where appropriate

and day 18/19 of pregnancy data was analysed using t tests. Correlation between SE

occurrence and LH pulses was assessed using chi-squared test. All fibre photometry data

used for quantitative analysis and comparison were from the first pregnancy and lactation. A

full list of probability values, inferential statistics, and degrees of freedom for all data can be

found in Supplementary Table 1.

Results

Prolactin action on forebrain neurons is necessary to maintain lactational infertility

Lactation has previously been associated with a marked decrease in *Kiss1* mRNA levels in

both RP3V and arcuate nucleus populations (13). To determine whether prolactin was

involved in the maintenance of lactational anestrus, the *Prlr* gene was knocked out of

Camk2a expressing neurons (most forebrain neurons) of female mice. Control *Prlr*^{lox/lox} mice

showed a marked reduction in kisspeptin cell body immunoreactivity in the RP3V of

lactating compared to NL mice (p = 0.0100, *Post hoc* Tukey's multiple comparisons test,

Figure 1A, C). In contrast, in *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice the lactation-induced suppression of

kisspeptin cell bodies in the RP3V was absent (p = 0.6409, *Post hoc* Tukey's multiple

comparisons test; interaction between reproductive state and genotype p = 0.0034, two-way

ANOVA, Figure 1A, C). Fibre density in the arcuate nucleus was significantly increased in

lactating *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice compared to lactating *Prlr*^{lox/lox} controls (p = 0.0020,

unpaired two tailed t test, measured as percentage voxels within the region of interest, Figure

1B, D).

Estrous cycles during lactation were significantly altered by deletion of Prlr in the forebrain,

with all *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice showing a return to estrus between day 6 and day 10 of

lactation (Figure 1E), while, as normal, estrus did not occur until day 20 (after weaning) in

control *Prlr*^{lox/lox} mice (p = <0.0001, log-rank (Mantel-Cox) test, Figure 1E). No differences

in litter weight gain from day 3 to day 8 of lactation were observed in either group (p =

0.3282, mixed analysis test, Supplementary Figure 4), indicating that the suckling stimulus

that mice received and lactating itself was maintained in the absence of *Prlr* expression in

Camk2a expressing neurons. Collectively, these data show that prolactin action in the brain is

absolutely required for the lactation-induced suppression of kisspeptin expression and to

maintain lactational infertility in mice.

In a separate cohort of *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice, pulsatile LH secretion was monitored in

early lactation, prior to the return of estrous cycles. To rule out a potential role for

progesterone in suppressing fertility during lactation (9,43,44), the progesterone receptor

antagonist mifepristone (RU486), was administered to these mice in early lactation. Vehicle-

treated *Prlr*^{lox/lox} control mice showed the expected near complete absence of pulsatile LH

secretion during lactation (Figure 2A, B). In contrast, nearly all *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice

showed a lack of the normal lactation-induced suppression of pulsatile LH secretion

demonstrated by a significant increase in frequency of LH pulses compared to controls (effect

of genotype p = 0.0024, two-way ANOVA with Tukey's multiple comparisons test, Figure

2A, B). There was no effect of mifepristone on the pattern of LH secretion, suggesting that

progesterone action is not required for the suppression of LH secretion (LH pulse frequency

(interaction genotype x treatment p = 0.2807; treatment p = 0.8588; Figure 2B); mean LH

levels (interaction genotype x treatment p = 0.8697; treatment p = 0.8586; Figure 2C); two-

way ANOVA with Tukey's multiple comparisons test). These data indicate that prolactin is

the primary signal responsible for the suppression of pulsatile LH secretion during lactation

in mice.

Episodic activity of arcuate kisspeptin neurons is suppressed during pregnancy and

most of lactation

To directly assess the role of prolactin in regulating kisspeptin neuron activity during

lactation, GCaMP fibre photometry was undertaken to monitor real-time activity of arcuate

kisspeptin neurons in freely behaving mice. We first undertook a longitudinal assessment of

changes in kisspeptin neuronal activity by tracking individual $Kiss1^{Cre}$ mice throughout

pregnancy, lactation, and following weaning (Figure 3).

Initially, GCaMP fibre photometry recordings were collected in the NL diestrous state, both

with and without serial blood sampling to measure LH concentrations. As can be seen in

Figure 4A, arcuate kisspeptin photometry recordings were characterized by discrete SEs of

elevated intracellular calcium (indicative of synchronous activity of the kisspeptin

population), with each SE consistently correlating to a single pulse of LH secretion in the

minutes following (p = <0.0001, chi-squared test). These were observed to be at a similar rate

to that described previously in diestrous mice using GCaMP photometry in a different $Kiss1$

mouse line (9), with periodic SEs occurring about once per hour (1.250±0.250/hr; Figure 3 &

4B).

The activity of the arcuate kisspeptin population in $Kiss1^{Cre}$ mice dynamically changed

depending on the reproductive state of the mouse (p = 0.0012, mixed-effect analysis, Figure 3

& 4B). On day 4 of pregnancy, SE frequency had markedly decreased (0.297±0.136/hr;

Figure 3 & 4B), indicating an early reduction in activity of arcuate kisspeptin neurons during

pregnancy. By day 14 of pregnancy, no SEs were seen (0±0/hr; Figure 3 & 4B) and this was

confirmed by a lack of pulsatile LH secretion (Supplementary Figure 5). In late pregnancy

(day 18), neuronal activity was monitored for 14 hours, during which time low amplitude,

SEs were unexpectedly observed at similar frequencies to NL levels (2.043±0.940/hr; Figure

3 & 4B-C). This unusual pattern of activity is illustrated in more detail in Figure 5 where,

alongside the resurgence of low amplitude SEs, a marked increase in baseline activity was

observed relative to recordings at other reproductive stages. This activity was reminiscent of

the miniature SEs observed to be caused by activation of subgroups of arcuate kisspeptin

cells using brain slice calcium imaging (11), but we are unable to resolve such SEs using the

present methods. Since our aim was to continue longitudinal assessment of the arcuate

kisspeptin population into lactation and weaning, mice were not disrupted by blood sampling

immediately prior to parturition as we were concerned that this additional stressor might

interfere with establishment of maternal behaviour. Hence, we are unable to report whether

these low amplitude SEs and elevated baseline activity were associated with LH secretion.

Evaluation of activity of arcuate kisspeptin neurons during lactation showed complete

suppression of activity on day 7 of lactation with a corresponding absence of pulsatile LH

secretion (0±0/hr; Figure 3 & 4B). This lactation-induced suppression of activity was

partially relieved by day 14 of lactation (0.583±0.083/hr; Figure 3 & 4B), with low frequency

SEs again corresponding to low frequency pulsatile LH secretion. Further increases in SE

frequency were seen on day 18 of lactation (0.850±0.100/hr; Figure 3 & 4B), including an

increase in baseline activity, similar to that seen in late pregnancy, and by 24 hours after

weaning (day 22 postpartum) the frequency of SEs had returned to close to non-pregnant

levels (1.375±0.114/hr; Figure 3 & 4B). SE frequency remained unchanged on the day of first

diestrus following a return to estrous cycles after weaning (1.533±0.226/hr; Figure 3 & 4B).

Ten days post-OVX, clusters of high amplitude arcuate kisspeptin population activity was

observed (4.778±0.222/hr; Figure 3 & 4B), consistent with previous reports following OVX

in nulliparous mice (45). Collectively, these observations show extensive, dynamic variation

in the activity of the arcuate kisspeptin neuronal population associated with pregnancy and

lactation.

Mice with an arcuate kisspeptin neuron-specific deletion have premature reactivation of

estrous cycles and neuronal activity in lactation

To determine whether the prolactin-induced suppression of estrous cycles and pulsatile LH

secretion was specifically mediated by kisspeptin neurons, mice were generated with an

arcuate-specific deletion of the Prlr from kisspeptin neurons (19). Similar to the data from

$Prlr^{\text{lox/lox}}/Camk2a^{\text{Cre}}$ mice, there was early resumption of estrous cycles in $Prlr^{\text{lox/lox}}/Kiss1^{\text{Cre}}$

mice during lactation (63% showing estrus by day 10 of lactation and 83% by day 19

lactation) compared to $Prlr^{\text{lox/lox}}$ controls (4% by day 19 lactation) (p = <0.0001, log-rank

(Mantel-Cox) test, Figure 6A). No difference in litter weight gain during lactation was

observed in either group indicating that suckling and/or lactation itself was not impaired (day

3-20 lactation weight gain, p = 0.6404, two-way ANOVA, Supplementary Figure 4H). *In*

vivo GCaMP fibre photometry in $Prlr^{\text{lox/lox}}/Kiss1^{\text{Cre}}$ mice showed early reactivation of the

arcuate kisspeptin population by day 5 of lactation (Figure 6B). This was accompanied by a

clear return to estrus within this early lactation window in four out of five mice. These data

demonstrate, in mice, that prolactin action specifically on arcuate kisspeptin neurons is

responsible for maintaining suppression of those neurons, and thereby fertility, during

lactation.

Discussion

We demonstrate here that prolactin action in arcuate kisspeptin neurons is necessary for the
maintained suppression of fertility during lactation in mice. Neuron-specific Prlr deletion
($Prlr^{lox/lox}/Camk2a^{Cre}$) resulted in premature return to estrus in early lactation, even in the
presence of ongoing suckling stimulus and the full metabolic consequences of milk
production. Accompanying the resumption of estrus was an absence of the normal lactation-
induced reduction in kisspeptin immunoreactivity (12,16,20). Pulsatile LH secretion was also
observed on day 5 of lactation prior to the premature estrus when it would normally have
been completely absent (46-48). To evaluate the specific role of kisspeptin neurons in
mediating the prolactin-induced suppression of fertility, we have comprehensively mapped
the activity of arcuate kisspeptin neurons throughout a full reproductive cycle: pregnancy,
lactation, and after weaning in individual animals. The data show an immediate suppression
of activity of arcuate kisspeptin neuronal activity during pregnancy, and this is maintained
throughout most of lactation, apart from a brief window of reactivation immediately prior to
parturition. Deleting Prlr specifically from arcuate kisspeptin neurons ($Prlr^{lox/lox}/Kiss1^{Cre}$)
prevented the suppression of activity in early lactation, resulting in premature induction of
episodic activation of kisspeptin neurons, and early onset of estrus. Combined, these data
provide direct evidence that prolactin action on kisspeptin neurons is necessary for lactation-
induced infertility in mice.

It is now well established that arcuate kisspeptin neurons form the GnRH "pulse generator",
and hence drive pulsatile release of GnRH from the hypothalamus and consequent pulsatile
secretion of LH from the pituitary that is required for fertility (49-51). This is the first study
to monitor activity of the arcuate kisspeptin neurons across different reproductive states in the
same animal, and the data largely match previously described patterns of LH secretion
(47,52). The frequency and dynamics of the SEs changed dramatically, initially due to the
pregnancy-induced changes in ovarian hormones. The abrupt decrease in arcuate kisspeptin
activity in early pregnancy is likely caused by rising levels of progesterone, known to
profoundly suppress activity of arcuate kisspeptin neurons and LH secretion (9,43,44).
Progesterone is elevated throughout pregnancy, gradually increasing until luteolysis and
progesterone withdrawal occurs in the lead up to parturition (53-56). Interestingly, we
observed a transient reactivation of the arcuate kisspeptin neurons in the night between days
18 and 19 of pregnancy. This was characterised by frequent, low amplitude SEs, and

increased baseline activity that may represent the intermittent synchroniszed activity of small

subsets of arcuate kisspeptin neurons that have not yet transitioned to full synchroniszation of

the whole population (11). It seems likely that this pattern of activity is associated with

progesterone withdrawal in late pregnancy and may be important in stimulating follicular

growth leading up to a postpartum ovulation (57,58).

In early lactation, episodic activity of arcuate kisspeptin neurons was absent, with sporadic

low-amplitude activity returning around day 14 of lactation. There was another period of

increased baseline activity in late lactation, similar to that seen in late pregnancy, potentially

representing a signature of reactivation of synchroniszed activity of the arcuate kisspeptin

neurons. This increase in arcuate kisspeptin population activity during late lactation mirrors

the increase in LH levels that has been reported as lactation progresses (59). Overall patterns

of SE activity rapidly returned to normal diestrus levels soon after weaning (59). However, in

the absence of Prlr in arcuate kisspeptin neurons, SE activity re-appeared as early as day 5 of

lactation, even in the presence of ongoing suckling. These data clearly show that prolactin

action in arcuate kisspeptin neurons is necessary to sustain lactational infertility in mice. The

observed disruption of lactational infertility in $Prlr^{lox/lox}/Kiss1^{Cre}$ mice is particularly

remarkable given that Prlr deletion is restricted to arcuate kisspeptin neurons in this model

(19), and prolactin action on RP3V kisspeptin neurons (21,60) and on gonadotrophs in the

pituitary gland (61-63) are unaffected.

The indispensable role for prolactin in mediating lactation-induced infertility in the mouse is

surprising, given the consensus of much work in other species concluding that other factors

may be more important (see (4,6,25,26,64)). This may reflect a level of redundancy amongst

contributing factors across all species, including ovarian hormones, metabolic cues and

neural inputs of suckling. Notably, the conditional deletion approach described here

distinguishes prolactin action from the neurogenic effects of suckling without altering the

process of lactation itself. Moreover, this approach avoids the potential confounding effects

of using dopamine agonists to suppress prolactin (28,29), given that dopamine can directly

inhibit GnRH neuronal activity (65).

While the effects of widespread neuronal deletion ($Prlr^{lox/lox}/Camk2a^{Cre}$) on fertility were

largely recapitulated by the arcuate kisspeptin-specific model ($Prlr^{lox/lox}/Kiss1^{Cre}$), it was

apparent that the global deletion was more effective at inducing the return to estrus during

lactation (in 100% of $Prlr^{lox/lox}/Camk2a^{Cre}$ mice by day 10), compared to the arcuate

kisspeptin-specific model (63% by day 10, and 83% by day 19). This may be due to the

absence of lactation-induced suppression of *Kiss1* expression in RP3V kisspeptin neurons of

Prlr^lox/lox^/*Camk2a*^Cre^ mice. Similarly, we cannot rule out the possibility that other populations

of prolactin-sensitive neurons, such as GABA or dopamine neurons (60), may contribute to

suppressing estrous cycles during lactation.

Nevertheless, our data collectively provide strong evidence that prolactin action on arcuate

kisspeptin neurons is the primary factor mediating lactation-induced infertility in mice. Given

that hyperprolactinemia induces infertility in many other mammalian species, including

humans (21,22,66-75), it is possible that a conserved mechanism could be contributing to

lactational infertility in these species.

Acknowledgments

We would like to acknowledge the research assistance of Zin Khant-Aung, genotyping by

Pene Knowles, and Dr Joon Kim for assistance with analysis of fibre photometry data.

Financial support: This work was supported by Health Research Council of New Zealand

(grant number: 21-560) and the Lions Club of Dunedin South - administered by Perpetual

Guardian (Otago Medical Research Foundation).

References

1. Short RV. Lactation--the central control of reproduction. *Ciba Foundation*
symposium. 1976(45):73-86.
2. McNeilly AS, Glasier AF, Howie PW, Houston MJ, Cook A, Boyle H. Fertility after
childbirth: pregnancy associated with breast feeding. *Clin Endocrinol (Oxf)*.
1983;19(2):167-173.
3. Fox SR, Smith MS. The suppression of pulsatile luteinizing hormone secretion during
lactation in the rat. *Endocrinology*. 1984;115(6):2045-2051.
4. McNeilly AS. Lactational control of reproduction. *Reproduction, fertility, and*
development. 2001;13(7-8):583-590.
5. Tsukamura H, Maeda K. Non-metabolic and metabolic factors causing lactational
anestrus: rat models uncovering the neuroendocrine mechanism underlying the
suckling-induced changes in the mother. *Progress in brain research*. 2001;133:187-
205.
6. McNeilly AS. Lactation and fertility. *Journal of mammary gland biology and*
neoplasia. 1997;2(3):291-298.
7. Clarkson J, Han SY, Piet R, McLennan T, Kane GM, Ng J, Porteous RW, Kim JS,
Colledge WH, Iremonger KJ, Herbison AE. Definition of the hypothalamic GnRH
pulse generator in mice. *Proc Natl Acad Sci U S A*. 2017;114(47):E10216-E10223.
8. Han SY, Kane G, Cheong I, Herbison AE. Characterization of GnRH Pulse Generator
Activity in Male Mice Using GCaMP Fiber Photometry. *Endocrinology*.
2019;160(3):557-567.
9. McQuillan HJ, Han SY, Cheong I, Herbison AE. GnRH Pulse Generator Activity
Across the Estrous Cycle of Female Mice. *Endocrinology*. 2019;160(6):1480-1491.
10. Moore AM, Coolen LM, Lehman MN. In vivo imaging of the GnRH pulse generator
reveals a temporal order of neuronal activation and synchronization during each
pulse. *Proceedings of the National Academy of Sciences*. 2022;119(6):e2117767119.
11. Han SY, Morris PG, Kim JC, Guru S, Pardo-Navarro M, Yeo SH, McQuillan HJ,
Herbison AE. Mechanism of kisspeptin neuron synchronization for pulsatile hormone
secretion in male mice. *Cell Rep*. 2023;42(1):111914.
12. Yamada S, Uenoyama Y, Kinoshita M, Iwata K, Takase K, Matsui H, Adachi S,
Inoue K, Maeda KI, Tsukamura H. Inhibition of metastin (kisspeptin-54)-GPR54
signaling in the arcuate nucleus-median eminence region during lactation in rats.
Endocrinology. 2007;148(5):2226-2232.
13. Liu X, Brown RS, Herbison AE, Grattan DR. Lactational anovulation in mice results
from a selective loss of kisspeptin input to GnRH neurons. *Endocrinology*.
2014;155(1):193-203.
14. Roa J, Vigo E, Castellano JM, Navarro VM, Fernández-Fernández R, Casanueva FF,
Dieguez C, Aguilar E, Pinilla L, Tena-Sempere M. Hypothalamic expression of KiSS-
1 system and gonadotropin-releasing effects of kisspeptin in different reproductive
states of the female Rat. *Endocrinology*. 2006;147(6):2864-2878.
15. Ladyman SR, Woodside B. Food restriction during lactation suppresses Kiss1 mRNA
expression and kisspeptin-stimulated LH release in rats. *Reproduction*.
2014;147(5):743-751.
16. True C, Kirigiti M, Ciofi P, Grove KL, Smith MS. Characterisation of arcuate nucleus
kisspeptin/neurokinin B neuronal projections and regulation during lactation in the
rat. *J Neuroendocrinol*. 2011;23(1):52-64.

17.   Brown RS, Herbison AE, Grattan DR. Prolactin regulation of kisspeptin neurones in
the mouse brain and its role in the lactation-induced suppression of kisspeptin
expression. *J Neuroendocrinol*. 2014;26(12):898-908.
18.   Xu J, Kirigiti MA, Grove KL, Smith MS. Regulation of food intake and
gonadotropin-releasing hormone/luteinizing hormone during lactation: role of insulin
and leptin. *Endocrinology*. 2009;150(9):4231-4240.
19.   Brown RSE, Khant Aung Z, Phillipps HR, Barad Z, Lein HJ, Boehm U, Szawka RE,
Grattan DR. Acute Suppression of LH Secretion by Prolactin in Female Mice Is
Mediated by Kisspeptin Neurons in the Arcuate Nucleus. *Endocrinology*.
2019;160(5):1323-1332.
20.   Araujo-Lopes R, Crampton JR, Aquino NS, Miranda RM, Kokay IC, Reis AM,
Franci CR, Grattan DR, Szawka RE. Prolactin regulates kisspeptin neurons in the
arcuate nucleus to suppress LH secretion in female rats. *Endocrinology*.
2014;155(3):1010-1020.
21.   Sonigo C, Bouilly J, Carre N, Tolle V, Caraty A, Tello J, Simony-Conesa FJ, Millar
R, Young J, Binart N. Hyperprolactinemia-induced ovarian acyclicity is reversed by
kisspeptin administration. *J Clin Invest*. 2012;122(10):3791-3795.
22.   Hackwell ECR, Ladyman SR, Brown RS, Grattan DR. Mechanisms of lactation-
induced infertility in female mice. *Endocrinology*. 2023;164(5):bqad049.
23.   McNeilly AS. Suckling and the control of gonadotropin secretion. *The physiology of
reproduction*. 1994.
24.   Sugimoto A, Tsuchida H, Nagae M, Inoue N, Uenoyama Y, Tsukamura H. Central
somatostatin-somatostatin receptor 2 signaling mediates lactational suppression of
luteinizing hormone release via the inhibition of glutamatergic interneurons during
late lactation in rats. *Journal of Reproduction and Development*. 2022.
25.   Tay CC, Glasier AF, McNeilly AS. Twenty-four hour patterns of prolactin secretion
during lactation and the relationship to suckling and the resumption of fertility in
breast-feeding women. *Human reproduction (Oxford, England)*. 1996;11(5):950-955.
26.   Diaz S, Seron-Ferre M, Cardenas H, Schiappacasse V, Brandeis A, Croxatto HB.
Circadian variation of basal plasma prolactin, prolactin response to suckling, and
length of amenorrhea in nursing women. *The Journal of clinical endocrinology and
metabolism*. 1989;68(5):946-955.
27.   Ordog T, Chen MD, O'Byrne KT, Goldsmith JR, Connaughton MA, Hotchkiss J,
Knobil E. On the mechanism of lactational anovulation in the rhesus monkey. *The
American journal of physiology*. 1998;274(4 Pt 1):E665-676.
28.   Maeda K, Uchida E, Tsukamura H, Ohkura N, Ohkura S, Yokoyama A. Prolactin
does not mediate the suppressive effect of the suckling stimulus on luteinizing
hormone secretion in ovariectomized lactating rats. *Endocrinologia japonica*.
1990;37(3):405-411.
29.   Smith MS. The relative contribution of suckling and prolactin to the inhibition of
gonadotropin secretion during lactation in the rat. *Biol Reprod*. 1978;19(1):77-83.
30.   Steyn FJ, Wan Y, Clarkson J, Veldhuis JD, Herbison AE, Chen C. Development of a
methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile
and adult male mice. *Endocrinology*. 2013;154(12):4939-4945.
31.   Czieselsky K, Prescott M, Porteous R, Campos P, Clarkson J, Steyn FJ, Campbell RE,
Herbison AE. Pulse and Surge Profiles of Luteinizing Hormone Secretion in the
Mouse. *Endocrinology*. 2016;157(12):4794-4802.
32.   Briski KP, Sylvester PW. Effects of sequential acute stress exposure on stress-induced
pituitary luteinizing hormone and prolactin secretion. *Life sciences*.
1987;41(10):1249-1255.

33.   Steyn FJ, Huang L, Ngo ST, Leong JW, Tan HY, Xie TY, Parlow AF, Veldhuis JD,
Waters MJ, Chen C. Development of a method for the determination of pulsatile
growth hormone secretion in mice. *Endocrinology*. 2011;152(8):3165-3171.
34.   Brown RS, Kokay IC, Phillipps HR, Yip SH, Gustafson P, Wyatt A, Larsen CM,
Knowles P, Ladyman SR, LeTissier P, Grattan DR. Conditional Deletion of the
Prolactin Receptor Reveals Functional Subpopulations of Dopamine Neurons in the
Arcuate Nucleus of the Hypothalamus. *J Neurosci*. 2016;36(35):9173-9185.
35.   Gustafson P, Ladyman SR, McFadden S, Larsen C, Khant Aung Z, Brown RSE, Bunn
SJ, Grattan DR. Prolactin receptor-mediated activation of pSTAT5 in the pregnant
mouse brain. *J Neuroendocrinol*. 2020;32(11):e12901.
36.   Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA. Regulation of
Kiss1 gene expression in the brain of the female mouse. *Endocrinology*.
2005;146(9):3686-3692.
37.   Moult PJ, Rees LH, Besser GM. Pulsatile gonadotrophin secretion in
hyperprolactinaemic amenorrhoea an the response to bromocriptine therapy. *Clin
Endocrinol (Oxf)*. 1982;16(2):153-162.
38.   Porteous R, Haden P, Hackwell E, Singline A, Herde M, Desai R, Handelsman D,
Grattan D, Herbison A. Reformulation of PULSAR for Analysis of Pulsatile LH
Secretion and a Revised Model of Estrogen-Negative Feedback in Mice.
Endocrinology. 2021;162(11).
39.   Clarkson J, d'Anglemont de Tassigny X, Colledge WH, Caraty A, Herbison AE.
Distribution of kisspeptin neurones in the adult female mouse brain. *J
Neuroendocrinol*. 2009;21(8):673-682.
40.   Clarkson J, Boon WC, Simpson ER, Herbison AE. Postnatal development of an
estradiol-kisspeptin positive feedback mechanism implicated in puberty onset.
Endocrinology. 2009;150(7):3214-3220.
41.   Kim CK, Yang SJ, Pichamoorthy N, Young NP, Kauvar I, Jennings JH, Lerner TN,
Berndt A, Lee SY, Ramakrishnan C. Simultaneous fast measurement of circuit
dynamics at multiple sites across the mammalian brain. *Nature methods*.
2016;13(4):325-328.
42.   Sherathiya VN, Schaid MD, Seiler JL, Lopez GC, Lerner TN. GuPPy, a Python
toolbox for the analysis of fiber photometry data. *Scientific Reports*.
2021;11(1):24212.
43.   Goodman RL, Karsch FJ. Pulsatile secretion of luteinizing hormone: differential
suppression by ovarian steroids. *Endocrinology*. 1980;107(5):1286-1290.
44.   Skinner DC, Evans NP, Delaleu B, Goodman RL, Bouchard P, Caraty A. The
negative feedback actions of progesterone on gonadotropin-releasing hormone
secretion are transduced by the classical progesterone receptor. *Proc Natl Acad Sci U
S A*. 1998;95(18):10978-10983.
45.   McQuillan HJ, Clarkson J, Kauff A, Han SY, Yip SH, Cheong I, Porteous R, Heather
AK, Herbison AE. Definition of the estrogen negative feedback pathway controlling
the GnRH pulse generator in female mice. *Nature Communications*. 2022;13(1):7433.
46.   Bohnet HG, Schneider HP. Prolactin as a cause of anovulation. In: Crosignani PG,
Robyn C, eds. Prolactin and human reproduction. New York: Academic Press;
1977:153-159.
47.   McNeilly. Effects of lactation on fertility. 1979.
48.   McNeilly AS, Tay CC, Glasier A. Physiological mechanisms underlying lactational
amenorrhea. *Annals of the New York Academy of Sciences*. 1994;709:145-155.
49.   Herbison AE. The Gonadotropin-Releasing Hormone Pulse Generator.
Endocrinology. 2018;159(11):3723-3736.

50.   de Roux N, Genin E, Carel JC, Matsuda F, Chaussain JL, Milgrom E.
720   Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived
721   peptide receptor GPR54. *Proc Natl Acad Sci U S A*. 2003;100(19):10972-10976.
722   51.   Seminara et al. The GPR54 gene as a regulator of puberty. 2003.
723   52.   Morishige WK, Pepe GJ, Rothchild I. Serum luteinizing hormone, prolactin and
724   progesterone levels during pregnancy in the rat. *Endocrinology*. 1973;92(5):1527-
725   1530.
726   53.   Virgo BB, Bellward GD. Serum progesterone levels in the pregnant and postpartum
727   laboratory mouse. *Endocrinology*. 1974;95(5):1486-1490.
728   54.   Pointis G, Rao B, Latreille M, Mignot T-M, Cedard L. Progesterone levels in the
729   circulating blood of the ovarian and uterine veins during gestation in the mouse.
730   *Biology of Reproduction*. 1981;24(4):801-805.
731   55.   Bridges RS. A quantitative analysis of the roles of dosage, sequence, and duration of
732   estradiol and progesterone exposure in the regulation of maternal behavior in the rat.
733   *Endocrinology*. 1984;114(3):930-940.
734   56.   Sugimoto Y, Yamasaki A, Segi E, Tsuboi K, Aze Y, Nishimura T, Oida H, Yoshida
735   N, Tanaka T, Katsuyama M. Failure of parturition in mice lacking the prostaglandin F
736   receptor. *Science*. 1997;277(5326):681-683.
737   57.   Thapa S, Short RV, Potts M. Breast feeding, birth spacing and their effects on child
738   survival. *Nature*. 1988;335(6192):679-682.
739   58.   Zakar T, Hertelendy F. Progesterone withdrawal: key to parturition. *American journal
740   of obstetrics and gynecology*. 2007;196(4):289-296.
741   59.   Rolland R, Lequin RM, Schellekens LA, Jong FHD. The role of prolactin in the
742   restoration of ovarian function during the early post-partum period in the human
743   female: I. A study during physiological lactation. *Clinical Endocrinology*.
744   1975;4(1):15-25.
745   60.   Kokay IC, Petersen SL, Grattan DR. Identification of prolactin-sensitive GABA and
746   kisspeptin neurons in regions of the rat hypothalamus involved in the control of
747   fertility. *Endocrinology*. 2011;152(2):526-535.
748   61.   Hodson DJ, Townsend J, Tortonese DJ. Characterization of the Effects of Prolactin in
749   Gonadotroph Target Cells1. *Biology of Reproduction*. 2010;83(6):1046-1055.
750   62.   Tortonese DJ, Brooks J, Ingleton PM, McNeilly AS. Detection of prolactin receptor
751   gene expression in the sheep pituitary gland and visualization of the specific
752   translation of the signal in gonadotrophs. *Endocrinology*. 1998;139(12):5215-5223.
753   63.   Henderson HL, Townsend J, Tortonese DJ. Direct effects of prolactin and dopamine
754   on the gonadotroph response to GnRH. *J Endocrinol*. 2008;197(2):343-350.
755   64.   Woodside C, Jans JE. Role of the nutritional status of the litter and length and
756   frequency of mother-litter contact bouts in prolonging lactational diestrus in rats.
757   *Horm Behav*. 1995;29(2):154-176.
758   65.   Liu X, Herbison AE. Dopamine regulation of gonadotropin-releasing hormone neuron
759   excitability in male and female mice. *Endocrinology*. 2013;154(1):340-350.
760   66.   Greer ME, Moraczewski T, Rakoff JS. Prevalence of hyperprolactinemia in
761   anovulatory women. *Obstetrics and gynecology*. 1980;56(1):65-69.
762   67.   Evans W, Cronin M, Thorner M. Hypoganadism in hyperprolactinemia: proposed
763   mechanisms. 1982.
764   68.   Koike K, Miyake A, Aono T, Sakumoto T, Ohmichi M, Yamaguchi M, Tanizawa O.
765   Effect of prolactin on the secretion of hypothalamic GnRH and pituitary
766   gonadotropins. *Hormone research*. 1991;35 Suppl 1:5-12.
767   69.   Patel SS, Bamigboye V. Hyperprolactinaemia. *Journal of obstetrics and gynaecology
768   : the journal of the Institute of Obstetrics and Gynaecology*. 2007;27(5):455-459.

70. Cohen-Becker IR, Selmanoff M, Wise PM. Hyperprolactinemia alters the frequency
and amplitude of pulsatile luteinizing hormone secretion in the ovariectomized rat.
Neuroendocrinology. 1986;42(4):328-333.
71. Sarkar DK, Yen SS. Hyperprolactinemia decreases the luteinizing hormone-releasing
hormone concentration in pituitary portal plasma: a possible role for beta-endorphin
as a mediator. *Endocrinology*. 1985;116(5):2080-2084.
72. Fox SR, Hoefer MT, Bartke A, Smith MS. Suppression of pulsatile LH secretion,
pituitary GnRH receptor content and pituitary responsiveness to GnRH by
hyperprolactinemia in the male rat. *Neuroendocrinology*. 1987;46(4):350-359.
73. Park SK, Keenan MW, Selmanoff M. Graded hyperprolactinemia first suppresses LH
pulse frequency and then pulse amplitude in castrated male rats. *Neuroendocrinology*.
1993;58(4):448-453.
74. Park SK, Selmanoff M. Dose-dependent suppression of postcastration luteinizing
hormone secretion exerted by exogenous prolactin administration in male rats: a
model for studying hyperprolactinemic hypogonadism. *Neuroendocrinology*.
1991;53(4):404-410.
75. Grattan DR, Jasoni CL, Liu X, Anderson GM, Herbison AE. Prolactin regulation of
gonadotropin-releasing hormone neurons to suppress luteinizing hormone secretion in
mice. *Endocrinology*. 2007;148(9):4344-4351.
788
789

 Figures

Figure 1. *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} mice do not undergo the normal period of lactational

infertility and the lactation-induced suppression of kisspeptin immunoreactivity is

absent. (A) Kisspeptin immunoreactivity shown in representative photomicrographs from

diestrus, nulligravid, non-lactating (NL; left) and lactating (right) *Prlr*$^{lox/lox}$ control and

Prlr$^{lox/lox}$/*Camk2a*^{Cre} mice (from anteroventral periventricular nucleus (AVPV) region of

RP3V). (B) Representative photomicrographs showing mid arcuate nucleus (mARC) of

lactating *Prlr*$^{lox/lox}$ (top) and lactating *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} mice (bottom). (C) Total kisspeptin

cell number for the RP3V (NL *Prlr*$^{lox/lox}$ (n = 6) versus lactating *Prlr*$^{lox/lox}$ control (n = 8) p =

0.0100, NL *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} (n = 5) versus lactating *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} (n = 8) p =

0.6409). Two-way ANOVA followed by Tukey's multiple comparisons test. (D)

Quantification of kisspeptin fibre density in the arcuate nucleus (Fiji software, measured in

percentage voxels per region of interest), showing total kisspeptin fibre density in the arcuate

nucleus (lactating *Prlr*$^{lox/lox}$ control n = 8, lactating *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} n = 7, p = 0.0020,

unpaired two-tailed t test). (E) Lactating *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} mice (blue, n = 8) resume

estrous cycles significantly earlier (100% within 6-10 days of lactation) than lactating

Prlr$^{lox/lox}$ controls (grey, n = 10) (p = <0.0001, Log-rank (Mantel-Cox) test). Scale bar image

and insert = 50μm. Values are shown as mean ± SEM.

Figure 2. Prolactin action in the brain during lactation is necessary for the suppression of pulsatile LH secretion. Examples of pulsatile LH levels in the blood from lactating *Prlr*[lox/lox] control and lactating *Prlr*[lox/lox]/*Camk2a*[Cre] mice that have either been treated with vehicle (sesame oil, s.c., grey, veh) or 4mg/kg mifepristone (in sesame oil, s.c., blue, mif). Graphs show LH pulse frequency (B; interaction p = 0.2807, genotype p = 0.0024, state p = 0.8558), and mean LH levels (C; interaction p = 0.8697, genotype p = 0.0031, state p = 0.8586). Lactating vehicle-treated *Prlr*[lox/lox] (n = 8), lactating mifepristone-treated *Prlr*[lox/lox] (n = 8), lactating vehicle-treated *Prlr*[lox/lox]/*Camk2a*[Cre] (n = 8), lactating mifepristone-treated *Prlr*[lox/lox]/*Camk2a*[Cre] (n = 7). Asterisks indicate LH pulse peaks as detected by PULSAR Otago analysis. Two-way ANOVA followed by Tukey's multiple comparisons test. Values are shown as mean ± SEM.

Figure 3. Longitudinal recordings of arcuate kisspeptin neuron GCaMP population

activity throughout different reproductive states in the same *Kiss1*^{Cre} mice.

Representative neuronal activity from three *Kiss1*[Cre] mice throughout the NL (diestrus,

nulligravid, non-lactating), pregnant, lactating, and post-weaning states. The time points

monitored in order were: diestrus NL, day 4 pregnancy, day 14 pregnancy, day 18/19

pregnancy (overnight), day 7 lactation, day 14 lactation, day 18 lactation, 24 hours after

weaning (day 22 postpartum), return to normal cycling following weaning (return to estrous

cycles), and 10 days post ovariectomy (OVX). Asterisks indicate SEs. Note: change in y axes

scale on all 3 OVX timepoints and mouse shown in dataset (B) from day 4 of pregnancy

onwards.

Figure 4. Synchronised Ca²⁺ events (SE) are consistently correlated with pulsatile LH

secretion across different reproductive states in *Kiss1*^Cre mice. (A) When fibre photometry

was paired with serial blood sampling for pulsatile LH secretion, the relationship between

SEs and LH pulses was examined. Each time an SE was seen during a recording with blood

sampling, pulsatile secretion of LH was also observed, with 100% correlation (p = <0.0001,

chi-squared test; 73 out of 73 SEs observed led to an LH pulse). Representative examples of

paired photometry and blood sampling are shown from the NL (diestrus, nulligravid, non-

lactating) state, from day 7 lactation, and from day 14 lactation. (B) Quantitative analysis of

SE frequency per hour across different reproductive states in $Kiss1^{Cre}$ mice (p = 0.0012,

mixed effect analysis (fixed type III) with Tukey's multiple comparisons tests). (C)

Quantitative analysis of relative SE amplitude of normalised ΔF/F across different

reproductive states (p = 0.0118, mixed effect analysis (fixed type III) with Tukey's multiple

comparisons tests). Black asterisks indicate SEs, red asterisks indicate LH pulse peaks as

detected by PULSAR Otago analysis. Values shown as mean ± SEM.

Figure 5. Activity of arcuate kisspeptin neurons on day 18 and 19 of pregnancy in

Kiss1^Cre **mice.** Fibre photometry recordings of mice on the evening of day 18 of pregnancy

(1800 hours) to the morning of day 19 of pregnancy (0800 hours) show low amplitude SEs.

3-hour section blown up for ease of viewing. (C) No difference was seen between frequency

of SEs (per 60 minutes) in the NL (diestrus, nulligravid, non-lactating) versus day 18/19

pregnancy (p = 0.4063, paired two-tailed t test), however a significant decrease in relative SE

amplitude is seen (D; p = 0.0141, paired two-tailed t test). Asterisks indicate SEs. Dotted line

in insert of (A) and (B) indicates 3 standard deviations (3SD). Grey shaded region = lights

off.

Figure 6. *Prlr*^{lox/lox}/*Kiss1*^{Cre} mice do not undergo the normal period of lactational infertility and show early reactivation of arcuate kisspeptin neurons prior to estrus in lactation. (A) $Prlr^{lox/lox}$/$Kiss1^{Cre}$ mice resume estrous cycles significantly earlier (63% by day 10 of lactation, n = 32, blue) than $Prlr^{lox/lox}$ controls (4% by day 19 of lactation, n = 30, grey) (p = <0.0001, Log-rank (Mantel-Cox) test). (B) Representative fibre photometry traces from day 5 of lactation from either a $Kiss1^{Cre}$ control mouse or $Prlr^{lox/lox}$/$Kiss1^{Cre}$ mice. Mice with Prlr knocked out of arcuate kisspeptin neurons ($Prlr^{lox/lox}$/$Kiss1^{Cre}$) show SEs early in lactation. Asterisks indicate SEs.

 Supplementary data

Supplementary Figure 1. Proportion of kisspeptin neurons showing *Prlr* deletion using

RNAscope in either *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} or *Prlr*$^{lox/lox}$/*Kiss1*^{Cre} mice. Representative

photomicrographs showing RNAscope labelling for *Prlr* (blue) and *Kiss1* (red) in the rostral

periventricular region of the third ventricle (RP3V, A) or arcuate nucleus (ARC, C, E), in

either intact (*Prlr*$^{lox/lox}$ control n = 5, *Prlr*$^{lox/lox}$/*Camk2a*^{Cre}, n = 5, A) or ovariectomised (OVX;

Prlr$^{lox/lox}$ control n = 4, *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} n = 5, C; *Prlr*$^{lox/lox}$ control n = 6,

Prlr$^{lox/lox}$/*Kiss1*^{Cre} n = 7, E) mice. Compared to *Prlr*$^{lox/lox}$ control mice, *Prlr*$^{lox/lox}$/*Camk2a*^{Cre}

mice show a significant decrease in the percentage of *Kiss1*-expressing cells co-expressing

Prlr in both the RP3V (B; p = <0.0001) and arcuate nucleus (D; p = 0.0009) (unpaired two-

tailed t tests). (F) A significant decrease in the percentage of *Kiss1*-expressing cells co-

expressing *Prlr* was seen in *Prlr*$^{lox/lox}$/*Kiss1*^{Cre} compared to *Prlr*$^{lox/lox}$ controls (p = <0.0001,

unpaired two-tailed t test). (G) No correlation was found between percentage of *Kiss1* cells

co-expressing with *Prlr* and the day of estrus return during lactation (p = 0.1912, simple

linear regression). Solid black arrows = doubled labelled cells expressing both *Kiss1* and

Prlr; black outlined arrows = *Kiss1* cells with sparse or no co-labelling for *Prlr*. Scale bar =

150μm, insert = 60μm. Values are shown as mean ± SEM.

890
891

Supplementary Figure 2. Mifepristone dose response and effect on litter weight gain.

(A) Mifepristone dose response trial showing dose of 4mg/kg was sufficient to terminate

pregnancy in all mice (p = 0.072, chi-squared test, n = 6 both groups). (B) Mifepristone or

vehicle treatment had no effect on litter weight gain from day 3 to day 5 of lactation

(interaction of time x genotype & treatment p = 0.5322; time p = <0.0001; genotype and

treatment p = 0.8811; subject p = <0.0001; two-way ANOVA). Values are shown as mean ±

SEM.

Supplementary Figure 3: Pulsatile LH secretion profiles of *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice

and *Prlr*^{lox/lox} controls following vehicle or mifepristone treatment. Individual LH pulse

data from lactating *Prlr*^{lox/lox} control and *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice treated with either vehicle

(sesame oil, s.c., grey) or 4mg/kg mifepristone (in sesame oil, s.c., blue). Lactating vehicle-

treated *Prlr*^{lox/lox} (n = 8), lactating mifepristone-treated *Prlr*^{lox/lox} (n = 8), lactating vehicle-

treated *Prlr*^{lox/lox}/*Camk2a*^{Cre} (n = 8), lactating mifepristone-treated *Prlr*^{lox/lox}/*Camk2a*^{Cre} (n =

7). Asterisks indicate LH pulse peaks as detected by PULSAR Otago analysis.

Supplementary Figure 3: Pulsatile LH secretion profiles of *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} mice

and *Prlr*$^{lox/lox}$ controls following vehicle or mifepristone treatment. Individual LH pulse

data from lactating *Prlr*$^{lox/lox}$ control and *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} mice treated with either vehicle

(sesame oil, s.c., grey) or 4mg/kg mifepristone (in sesame oil, s.c., blue). Lactating vehicle-

treated *Prlr*$^{lox/lox}$ (n = 8), lactating mifepristone-treated *Prlr*$^{lox/lox}$ (n = 8), lactating vehicle-

treated *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} (n = 8), lactating mifepristone-treated *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} (n =

7). Asterisks indicate LH pulse peaks as detected by PULSAR Otago analysis.

Supplementary Figure 4. Gestational and maternal phenotyping of *Prlr*^{lox/lox}/*Camk2a*^{Cre}

and *Prlr*^{lox/lox}/*Kiss1*^{Cre} mice and their respective *Prlr*^{lox/lox} controls. (A-H) Data show;

gestational weight gain (A; *Prlr*^{lox/lox} n = 11, *Prlr*^{lox/lox}/*Camk2a*^{Cre} n = 11; E; *Prlr*^{lox/lox} n = 31,

Prlr^{lox/lox}/*Kiss1*^{Cre} n =27), gestation length (B; *Prlr*^{lox/lox} n = 11, *Prlr*^{lox/lox}/*Camk2a*^{Cre} n = 11; F;

Prlr^{lox/lox} n = 31, *Prlr*^{lox/lox}/*Kiss1*^{Cre} n =27), number of live pups on day 3 of lactation (C;

Prlr^{lox/lox} n = 11; *Prlr*^{lox/lox}/*Camk2a*^{Cre} n = 11; G; *Prlr*^{lox/lox} n = 31, *Prlr*^{lox/lox}/*Kiss1*^{Cre} n =27),

and litter weight gain between day 3-8 of lactation (D; *Prlr*^{lox/lox} n = 8; *Prlr*^{lox/lox}/*Camk2a*^{Cre} n

= 8) or day 3-20 of lactation (H; *Prlr*^{lox/lox} n = 22, *Prlr*^{lox/lox}/*Kiss1*^{Cre} n = 20). There were no

differences in any of these parameters (A, C, E, G, unpaired two-tailed t test; B, F, Mann

Whitney test; D, H, repeated measures mixed effect analysis, fixed effects (type III) with

Šídák's multiple comparisons test. Grey = *Prlr*^{lox/lox}; blue in top row A-D =

Prlr^{lox/lox}/*Camk2a*^{Cre}, blue in bottom row E-H = *Prlr*^{lox/lox}/*Kiss1*^{Cre}. N numbers of

Prlr^{lox/lox}/*Kiss1*^{Cre} mice and respective *Prlr*^{lox/lox} controls changed due to: *Prlr*^{lox/lox} (n = 5),

and *Prlr*^{lox/lox}/*Kiss1*^{Cre} (n = 5) mice euthanised prior to day 20 lactation due to COVID-19

lockdown; *Prlr*^{lox/lox}/*Kiss1*^{Cre} (n = 2) showed estrus and therefore underwent blood sampling

(with respective *Prlr*^{lox/lox} controls (n =2)). All these mice were euthanised 2 hours following

blood sampling; a *Prlr*^{lox/lox} dams' (n =1) litter losing weight and being excluded from

subsequent analysis. Values are shown as mean ± SEM.

931
Supplementary Figure 5. Miniature synchronised event (SE)-like activity on day 14 of

pregnancy in _Kiss1_^{Cre} mice does not result in pulsatile LH secretion. (A) Paired fibre

photometry and blood sampling from representative mouse on day 14 of pregnancy showing

miniature SE-like activity have no significant effect on pulsatile LH secretion (red).

Supplementary Table 1. Statistics table. Abbreviations for tables below: DF = degrees of

freedom; mc = multiple comparison; CI = 95% confidence interval; MW U = Mann Whitney

U; MEA = Mixed effect analysis (fixed effects (type III)), RM = repeated measures. Supp =

supplementary data figure.

Fig.	Description	Statistical analysis				
1C	RP3V Kisspeptin immunoreactivity in *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice		**p**	**DF**	**CI**	**R^2**
	Interaction	Two-way ANOVA	0.0034	1		
	Genotype	Two-way ANOVA	0.0251	1		
	State	Two-way ANOVA	0.1380	1		
	Prlr^{lox/lox}:NL vs. *Prlr*^{lox/lox}:Lactating	Tukey's mc	0.0100	23	5.175 to 44.95	
	Prlr^{lox/lox}:NL vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:NL	Tukey's mc	0.9429	23	-17.79 to 26.81	
	Prlr^{lox/lox}:NL vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Lactating	Tukey's mc	0.9228	23	-24.38 to 15.39	
	Prlr^{lox/lox}:Lactating vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:NL	Tukey's mc	0.0565	23	-41.55 to 0.4412	
	Prlr^{lox/lox}:Lactating vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Lactating	Tukey's mc	0.0010	23	-47.97 to -11.15	
	Prlr^{lox/lox}/*Camk2a*^{Cre}:NL vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Lactating	Tukey's mc	0.6409	23	-30.00 to 11.99	
1D	Arcuate Kisspeptin immunoreactivity in *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice	Unpaired two-tailed t test	0.0020	14	0.6692 to 2.424	0.5052
1E	*Prlr*^{lox/lox}/*Camk2a*^{Cre} do not show lactational diestrus		**p**	**X^2**	**DF**	
	Percentage of mice showing estrus	Log rank (Mantel-Cox) test	<0.0001	42.37	1	
2B	Effect of mifepristone treatment on LH pulse frequency in *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice		**p**	**DF**	**CI**	
	Interaction	Two-way ANOVA	0.2807	1		
	Genotype	Two-way ANOVA	0.0024	1		
	Treatment	Two-way ANOVA	0.8558	1		
	Prlr^{lox/lox}:Veh vs. *Prlr*^{lox/lox}:Mif	Tukey's mc	0.7922	27	-0.9804 to 1.980	
	Prlr^{lox/lox}:Veh vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Veh	Tukey's mc	0.3861	27	-2.355 to 0.6054	
	Prlr^{lox/lox}:Veh vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Mif	Tukey's mc	0.1487	27	-2.765 to 0.3002	
	Prlr^{lox/lox}:Mif vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Veh	Tukey's mc	0.0758	27	-2.765 to 0.3002	
	Prlr^{lox/lox}:Mif vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Mif	Tukey's mc	0.0223	27	-2.765 to 0.3002	
	Prlr^{lox/lox}/*Camk2a*^{Cre}:Veh vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Mif	Tukey's mc	0.9188	27	-1.890 to 1.175	
2C	Effect of mifepristone treatment on mean LH in *Prlr*^{lox/lox}/*Camk2a*^{Cre} mice		**p**	**DF**		
	Interaction	Two-way ANOVA	0.8697	1		
	Genotype	Two-way ANOVA	0.0031	1		
	Treatment	Two-way ANOVA	0.8586	1		
	Prlr^{lox/lox}:Veh vs. *Prlr*^{lox/lox}:Mif	Tukey's mc	0.9947	26	-0.1021 to 0.1220	
	Prlr^{lox/lox}:Veh vs. *Prlr*^{lox/lox}/*Camk2a*^{Cre}:Veh	Tukey's mc	0.1528	26	-0.2015 to 0.02263	

Fig.	Description	Statistical analysis				
	$Prlr^{lox/lox}$:Veh vs. $Prlr^{lox/lox}/Camk2a^{Cre}$:Mif	Tukey's mc	0.1765	26	-0.2047 to 0.02672	
	$Prlr^{lox/lox}$:Mif vs. $Prlr^{lox/lox}/Camk2a^{Cre}$:Veh	Tukey's mc	0.0803	26	-0.2076 to 0.008853	
	$Prlr^{lox/lox}$:Mif vs. $Prlr^{lox/lox}/Camk2a^{Cre}$:Mif	Tukey's mc	0.0974	26	-0.2110 to 0.01306	
	$Prlr^{lox/lox}/Camk2a^{Cre}$:Veh vs. $Prlr^{lox/lox}/Camk2a^{Cre}$:Mif	Tukey's mc	>0.9999	26	-0.1116 to 0.1125	
4B	Kisspeptin activity across different reproductive states		**p**	**DF**	**CI**	
	SE frequency across states	MEA	0.0012			
	NLvs. Day 4 pregnancy	Tukey's mc	0.0828	5	-0.1356 to 2.052	
	NL vs. Day 14 pregnancy	Tukey's mc	0.0586	5	-0.05175 to 2.552	
	NL vs. Day 18/19 pregnancy	Tukey's mc	0.9837	4	-5.518 to 3.932	
	NL vs. Day 7 lactation	Tukey's mc	0.0912	4	-0.2517 to 2.752	
	NL vs. Day 14 lactation	Tukey's mc	0.6584	3	-1.598 to 2.931	
	NL vs. Day 18 lactation	Tukey's mc	0.9638	4	-1.661 to 2.461	
	NL vs. 24h after weaning	Tukey's mc	>0.9999	3	-3.097 to 2.847	
	NL vs. Return to estrous cycles	Tukey's mc	0.9993	4	-3.016 to 2.449	
	NL vs. OVX	Tukey's mc	0.0009	5	-4.955 to -2.100	
	Day 4 pregnancy vs. Day 14 pregnancy	Tukey's mc	0.5635	5	-0.3793 to 0.9626	
	Day 4 pregnancy vs. Day 18/19 pregnancy	Tukey's mc	0.6960	4	-6.958 to 3.455	
	Day 4 pregnancy vs. Day 7 lactation	Tukey's mc	0.6396	4	-0.5180 to 1.101	
	Day 4 pregnancy vs. Day 14 lactation	Tukey's mc	0.4286	3	-1.026 to 0.4430	
	Day 4 pregnancy vs. Day 18 lactation	Tukey's mc	0.2948	4	-1.579 to 0.4622	
	Day 4 pregnancy vs. 24h after weaning	Tukey's mc	0.1309	3	-2.649 to 0.4820	
	Day 4 vs. Return to estrous cycles	Tukey's mc	0.1986	4	-3.185 to 0.7012	
	Day 4 pregnancy vs. OVX	Tukey's mc	<0.0001	5	-5.683 to -3.290	
	Day 14 pregnancy vs. Day 18/19 pregnancy	Tukey's mc	0.5272	4	-7.022 to 2.936	
	Day 14 pregnancy vs. Day 7 lactation	Tukey's mc				
	Day 14 pregnancy vs. Day 14 lactation	Tukey's mc	0.0343	3	-1.092 to -0.07436	
	Day 14 pregnancy vs. Day 18 lactation	Tukey's mc	0.0092	4	-1.380 to -0.3202	
	Day 14 pregnancy vs. 24h after weaning	Tukey's mc	0.0072	3	-2.072 to -0.6781	
	Day 14 pregnancy vs. Return to estrous cycles	Tukey's mc	0.0210	4	-2.731 to -0.3355	
	Day 14 pregnancy vs. OVX	Tukey's mc	<0.0001	5	-5.877 to -3.679	
	Day 18/19 pregnancy vs. Day 7 lactation	Tukey's mc	0.6053	3	-4.432 to 8.518	
	Day 18/19 pregnancy vs. Day 14 lactation	Tukey's mc	0.9434	2	-11.74 to 14.66	
	Day 18/19 pregnancy vs. Day 18 lactation	Tukey's mc	0.9458	3	-5.853 to 8.239	
	Day 18/19 pregnancy vs. 24h after weaning	Tukey's mc	0.9807	2	-6.951 to 8.287	
	Day 18/19 pregnancy vs. Return to estrous cycles	Tukey's mc	0.9952	3	-4.164 to 5.183	
	Day 18/19 pregnancy vs. OVX	Tukey's mc	0.2556	4	-7.450 to 1.980	
	Day 7 lactation vs. Day 14 lactation	Tukey's mc	0.0382	3	-1.112 to -0.05439	
	Day 7 lactation vs. Day 18 lactation	Tukey's mc	0.0325	3	-1.578 to -0.1222	
	Day 7 lactation vs. 24h after weaning	Tukey's mc	0.0412	2	-2.620 to -0.1304	
	Day 7 lactation vs. Return to estrous cycles	Tukey's mc	0.0514	3	-3.083 to 0.01609	
	Day 7 lactation vs. OVX	Tukey's mc	0.0002	4	-5.887 to -3.669	

Fig.	Description	Statistical analysis				
	Day 14 lactation vs. Day 18 lactation	Tukey's mc	0.8130	3	-1.405 to 0.8715	
	Day 14 lactation vs. 24h after weaning	Tukey's mc	0.1642	2	-2.285 to 0.7020	
	Day 14 lactation vs. Return to estrous cycles	Tukey's mc	0.2736	2	-3.366 to 1.466	
	Day 14 lactation vs. OVX	Tukey's mc	0.0026	3	-5.734 to -2.655	
	Day 18 lactation vs. 24h after weaning	Tukey's mc	0.1662	3	-1.362 to 0.3122	
	Day 18 lactation vs. Return to estrous cycles	Tukey's mc	0.1946	3	-1.849 to 0.4823	
	Day 18 lactation vs. OVX	Tukey's mc	0.0016	4	-5.475 to -2.380	
	24h after weaning vs. Return to estrous cycles	Tukey's mc	0.9643	3	-1.180 to 0.8630	
	24h after weaning vs. OVX	Tukey's mc	0.0080	3	-5.192 to -1.614	
	Return to estrous cycles vs OVX	Tukey's mc	0.0113	4	-5.382 to -1.107	
4C	Kisspeptin activity across different reproductive states		**p**	**DF**	**CI**	
	SE amplitude across states	MEA	0.0118			
	NL vs. Day 4	Tukey's mc	0.7336	3	-0.9745 to 1.654	
	NL vs. Day 18/19	Tukey's mc	0.2608	3	-0.6762 to 2.086	
	NL vs. Day 14	Tukey's mc	0.0851	3	-0.1362 to 1.365	
	NL vs. Day 18	Tukey's mc	0.1766	4	-0.2966 to 1.467	
	NL vs. 24h after weaning	Tukey's mc	0.4700	3	-0.9997 to 2.178	
	NL vs. Return to estrous cycles	Tukey's mc	0.4514	4	-0.5997 to 1.546	
	NL vs. OVX	Tukey's mc	0.9938	5	-0.9886 to 0.7359	
	Day 4 vs. Day 18/19	Tukey's mc	0.7336	1	-5.882 to 6.612	
	Day 4 vs. Day 14	Tukey's mc	0.7858	2	-1.431 to 1.981	
	Day 4 vs. Day 18	Tukey's mc	0.6250	2	-0.9138 to 1.405	
	Day 4 vs. 24h after weaning	Tukey's mc	0.2399	1	-0.8660 to 1.365	
	Day 4 vs. Return to estrous cycles	Tukey's mc	0.9217	2	-1.026 to 1.293	
	Day 4 vs. OVX	Tukey's mc	0.5530	3	-1.870 to 0.9381	
	Day 18/19 vs. Day 14	Tukey's mc	0.7069	2	-0.5763 to 0.3960	
	Day 18/19 vs. Day 18	Tukey's mc	0.1985	3	-0.3267 to 0.08725	
	Day 18/19 vs. 24h after weaning	Tukey's mc	0.6222	2	-0.6603 to 0.4286	
	Day 18/19 vs. Return to estrous cycles	Tukey's mc	0.2384	2	-0.7754 to 0.3120	
	Day 18/19 vs. OVX	Tukey's mc	0.1494	3	-2.103 to 0.4405	
	Day 14 vs. Day 18	Tukey's mc	0.9929	3	-0.3118 to 0.2526	
	Day 14 vs. 24h after weaning	Tukey's mc	0.5747	2	-0.1375 to 0.08624	
	Day 14 vs. Return to estrous cycles	Tukey's mc	0.4210	2	-0.6226 to 0.3396	
	Day 14 vs. OVX	Tukey's mc	0.3306	3	-2.375 to 0.8931	
	Day 18 vs. 24h after weaning	Tukey's mc	>0.9999	3	-0.3977 to 0.4056	
	Day 18 vs. Return to estrous cycles	Tukey's mc	0.6644	3	-0.5038 to 0.2799	
	Day 18 vs. OVX	Tukey's mc	0.1809	4	-1.793 to 0.3702	
	24h after weaning vs. Return to estrous cycles		0.5101	3	-0.4457 to 0.2140	
	24h after weaning vs. OVX	Tukey's mc	0.2874	3	-2.185 to 0.7540	
	Return to estrous cycles vs. OVX	Tukey's mc	0.2096	4	-1.563 to 0.3637	
5C	NL vs day 18/19 pregnancy frequency of SE		**p**	**R^2**	**CI**	
		Paired two-tailed t test	0.4063	0.2364	-2.646 to 4.945	
5D	NL vs day 18/19 pregnancy relative SE amplitude		**p**	**R^2**	**CI**	
		Paired two-tailed t test	0.0141	0.8989	-1.606 to -0.3816	
6A	*Prlr*$^{lox/lox}$/*Kiss1*^{Cre} do not show lactational diestrus		**p**	**X^2**	**DF**	
	Percentage of mice showing estrus	Log rank (Mantel-Cox) test	<0.0001	38.02	1	
Supp 1	Proportion of kisspeptin neurons showing *Prlr* deletion using RNAscope		**p**	**R^2**	**CI**	
	B: *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} RP3V	Unpaired two-tailed t test	<0.0001	0.8740	-55.02 to 28.69	
	D: *Prlr*$^{lox/lox}$/*Camk2a*^{Cre} ARC	Unpaired two-tailed t test	0.0009	0.8123	-69.72 to 8.860	

Fig.	Description	Statistical analysis				
	F: *Prlr*[lox/lox]/*Kiss1*[Cre] ARC	Unpaired two-tailed t test	<0.0001	0.9300	-64.27 to -44.46	
	G: correlation of day of estrus and average density of *Prlr* on *Kiss1*-expressing cells (per animal)	Simple linear regression	0.1912	0.3135		
Supp 2A	Mifepristone functional dose response trial		**p**	**X²**	**Z**	
		Chi-squared test	0.0072	6	2.449	
Supp 2B	Mifepristone has no effect on litter weight gain		**p**	**DF**		
	Time x Genotype and treatment	Two-way RM ANOVA	0.5322	6		
	Time	Two-way RM ANOVA	<0.0001	2		
	Genotype and treatment	Two-way RM ANOVA	0.8811	3		
	Subject	Two-way RM ANOVA	<0.0001	22		
Supp 3A-D	*Prlr*[lox/lox]/*Camk2a*[Cre] maternal and gestational phenotyping		**p**	**R²**	**CI**	**MW U**
	A: gestational weight gain	Unpaired two-tailed t test	0.1291	0.1113	-3.076 to 0.4219	
	B: gestation length	Mann Whitney test	0.2682			45.50
	C: number of live pups	Unpaired two-tailed t test	0.7479	0.005282	-4.037 to 2.946	
	D: litter weight gain					
	Time	MEA	<0.0001			
	Genotype	MEA	0.3833			
	Time x genotype	MEA	0.3282			
	Day 3 lactation	Šídák's mc	>0.9999		-3.740 to 4.115	
	Day 4 lactation	Šídák's mc	0.9993		-3.365 to 4.490	
	Day 5 lactation	Šídák's mc	0.9863		-2.965 to 4.890	
	Day 6 lactation	Šídák's mc	0.9525		-2.690 to 5.165	
	Day 7 lactation	Šídák's mc	0.6938		-1.953 to 5.903	
	Day 8 lactation	Šídák's mc	0.619		-1.821 to 6.112	
Supp 3E-H	*Prlr*[lox/lox]/*Kiss1*[Cre] maternal and gestational phenotyping		**p**	**R²**	**CI**	**MW U**
	E: gestational weight gain	Unpaired two-tailed t test	0.6361	0.003694	-0.9985 to 1.622	
	F: gestation length	Mann Whitney test	0.7670			471.5
	G: number of live pups	Unpaired two-tailed t test	0.4522	0.009296	-0.5930 to 1.315	
	H: litter weight gain					
	Time x Genotype	Two-way RM ANOVA	0.6404			
	Time	Two-way RM ANOVA	<0.0001			
	Genotype	Two-way RM ANOVA	0.9014			
	Subject	Two-way RM ANOVA	<0.0001			
	Day 3 lactation	Šídák's mc	>0.9999		-5.407 to 3.653	
	Day 4 lactation	Šídák's mc	>0.9999		-5.204 to 3.857	
	Day 5 lactation	Šídák's mc	>0.9999		-5.159 to 3.902	
	Day 6 lactation	Šídák's mc	>0.9999		-5.105 to 3.956	
	Day 7 lactation	Šídák's mc	>0.9999		-4.976 to 4.084	
	Day 8 lactation	Šídák's mc	>0.9999		-4.226 to 4.834	
	Day 9 lactation	Šídák's mc	>0.9999		-4.757 to 4.304	
	Day 10 lactation	Šídák's mc	>0.9999		-4.650 to 4.410	
	Day 11 lactation	Šídák's mc	>0.9999		-5.154 to 3.907	
	Day 12 lactation	Šídák's mc	>0.9999		-5.242 to 3.819	
	Day 13 lactation	Šídák's mc	>0.9999		-4.936 to 4.125	
	Day 14 lactation	Šídák's mc	>0.9999		-4.979 to 4.081	
	Day 15 lactation	Šídák's mc	>0.9999		-4.776 to 4.284	
	Day 16 lactation	Šídák's mc	>0.9999		-4.152 to 4.909	
	Day 17 lactation	Šídák's mc	>0.9999		-4.216 to 4.845	
	Day 18 lactation	Šídák's mc	0.9912		-2.717 to 6.343	
	Day 19 lactation	Šídák's mc	>0.9999		-4.182 to 4.878	

Fig.	Description	Statistical analysis				
	Day 20 lactation	Šídák's mc	>0.9999		-4.752 to 4.309	

942