**Title:** Optogenetic silencing hippocampal inputs to the retrosplenial cortex causes a prolonged disruption of working memory.

**Authors:** Bárbara Pinto-Correia [1], Patricia Caldeira[1], and Miguel Remondes[1,2] *

**Affiliations:**

[1] Instituto de Medicina Molecular João Lobo Antunes,

   Faculdade de Medicina

   Universidade de Lisboa

   Lisbon, 1649-028, Portugal

[2] Faculdade de Medicina Veterinária

   Universidade Lusófona

   Lisbon, 1749-024, Portugal

**\*Correspondence and Lead Contact email:** mremondes@medicina.ulisboa.pt

**Summary**

Working memory allows us to keep information in memory for the time needed to perform a given task. Such fundamental cognitive ability relies on a neural circuit including the retrosplenial cortex (RSC), connected to several cortical areas, functionally and anatomically, namely primary visual areas, and higher cognitive areas such as the cingulate, midcingulate, and subicular cortices. RSC bears intimate anatomical and functional connections with the hippocampus, and has been implicated in integrating and translating spatial-temporal contextual information between ego- and allocentric reference frames, to compute predictions about goals in goal-directed behaviors. The relative contribution of the hippocampus and retrosplenial cortex in working memory-guided behaviors remains unclear due to the lack of studies reversibly interfering with synapses connecting the two regions during such behaviors. We here used eArch3.0, a hyperpolarizing proton pump, to silence hippocampal axon terminals in RSC while animals perform a standard delayed non-match to place task. We found that such

manipulation impairs memory retrieval, significantly decreasing performance and hastening decision-making. Furthermore, we found that such impairment outlasts light-activation of the opsin, its effects being noticed up to 3 subsequent trials.

**Introduction**

The ability to guide behavior according to knowledge stored previously in memory lies at the heart of adaptive behavior, and is essential for survival. A crucial aspect of adaptive behavior is the capacity to make decisions based on the spatial structure of the surrounding context, something globally known as spatial cognition or spatial goal-directed behavior. Such cognitive function starts with the selection of relevant information from a flood of sensory stimuli, and ends with its storage in cortico-hippocampal circuits (Morris et al., 1982; Nadel, 1990; Scoville et al., 1957) ultimately as a cognitive map manifested in a vast repertoire of neural activity, selective to multiple ego- (self-referenced) and allocentric (world-referenced) spatial variables, the seminal example of which is the existence of hippocampal place-selective neurons (O'Keefe and Dostrovsky, 1971). To allow for the planning of the movements necessary to reach an intended goal, all the while ensuring that body actions are consistent with the context acted upon, a mechanism must exist that reinstates (retrieves) allocentric contextual representations, stored in the hippocampus (HIPP) as spatial memories, aligned to the animal's viewpoint, e.g. from an egocentric perspective (Bicanski and Burgess, 2018; Burgess, 2008; Gomez et al., 2009, 2013; Michelmann et al., 2020). Such a mechanism, largely hypothetical, involves the strengthening of synapses between neurons across stations of a complex network including the subiculum, and anterior thalamus, providing head-direction signals (Chen et al., 1994a; Stackman and Taube, 1998; Taube et al., 1990), entorhinal cortex "grid" cells encoding all places in a given environment (Hafting et al., 2005), and posterior parietal cortex, responsible for integrating primary sensory information into multimodal egocentric ("self-referenced") representations of environmental boundaries and paths (Andersen et al., 1997; Ariani et al., 2015; Chen et al., 1994b; Clark et al., 2018; Kimura et al., 2004; Nakamura, 1999). In such circuit, the retrosplenial cortex (RSC) emerges as a cornerstone, encoding contextual variables in both ego- and allocentric reference frames, translating such

information between the two frames, and mediating the realignments of ego- and allocentric spatial information, according to behavioral needs. All the while, such mechanism would represent future spatial goals, and effectively anchor spatial memory-dependent decisions (Alexander and Nitz, 2015; Alexander et al., 2020; Burgess, 2008; Carstensen et al., 2021; Gomez et al., 2009, 2013; Michelmann et al., 2020; Nitzan et al., 2020).

Despite this remarkable work, we are yet to determine which neural pathways convey allocentric-referenced information stored in the HIPP, through hypothetical recall and egocentric re-referencing in RSC, onto the anterior midline cingulate (CG) and medial pre-frontal cortical areas, responsible for planning motor behavior according to previous events, present needs, and motivation (Akam et al., 2020; Brown and Braver, 2005; Cowen et al., 2012; Jin et al., 2009; Miller et al., 2019; Sober and Sabes, 2003). We have recently found that HIPP axons target the divisions of the medial mesocortex: CG, midcingulate and RSC, according to a gradient, in which RSC receives a combination of anatomically and functionally distinct inhibitory and excitatory functional synapses (Ferreira-Fernandes et al., 2019), whose manipulation results in specific behavioral deficits (Yamawaki et al., 2019). We hypothesize that these direct connections between hippocampal CA1 and RSC convey recently encoded and recalled memories to the relevant medial mesocortical regions.

To test this hypothesis we used the Delayed Non-Match to Place task (DNMP), a working memory (WM) task that requires animals to retrieve the memory of a previous decision, and of the time it occurred, to disambiguate it from similar, previously rewarded, decisions (Aggleton et al., 1986; Maguire, 2001; Shaw and Aggleton, 1995; Shaw et al., 2013; Wilson et al., 2013). We then used optogenetics to reversibly interrupt synaptic transmission between HIPP and RSC neurons during the retrieval phase of the DNMP task. Optogenetics uses light-activated membrane-bound molecules with the capacity to sustainedly, and reversibly, change neuronal membrane potential, through various molecular mechanisms, namely opening ion channels or pumping ions across the membrane (L Fenno, 2010), to excite or silence groups of neurons with behavior-specific temporal accuracy. As far as it is known, optogenetics is the only molecular technique allowing for reversible perturbation of neural activity within

the time constants of the cognitive functions at play during working memory (Yamamoto et al., 2014).

To test the hypothesis that HIPP inputs onto RSC carry contextual information necessary for DNMP performance, and are thus necessary for spatial working memory, we used the optogenetic neural silencer eArch3.0 expressed in hippocampal neurons via local injection of a viral vector, AAV5-hSyn1-eArch3.0, under a pan-neuronal promoter, or its "empty" counterpart with only YFP, in adult male Long Evans rats previously trained in DNMP to 75% performance. We then ran multiple 28-trial DNMP sessions, in which we randomly interleaved two types of trials: no-illumination (NI) or illumination during the Test, or retrieval, phase (TI), with green (525 nm) light delivered via 2 diagonal, intra-cerebral lambda fiberoptic implants, longitudinally staggered such as to deliver light to the full extent of RSC. In each animal we measured three performance variables: individual trial (error/correct) and global performance (% correct trials), response latency to the choice point (CP), and time spent at the CP. We then used multivariable Generalized Linear Mixed Model (GLMM) using the above categorical and continuous variables, to investigate the impact of silencing hippocampal synapses onto RSC neurons in WM-dependent behavior.

If indeed the HIPP input onto RSC neurons anchors the decisions during "Test" runs to the memory of the "Sample" runs, we predict that by silencing HIPP inputs in RSC we will prevent the transfer of information previously encoded in HIPP, and thus hinder the mechanism allowing RSC to inform the correct trajectory during the "Test" run, resulting in lower performance, specifically in eArch+ animals.

**Results**

**Injection of AAV-5-hSyn1-eArch3.0 results in significant expression of a fluorescent tag in RSC hippocampal terminals**

Besides abundant expression of both eArch and CTRL transgenes in hippocampal CA1, we found highly significant expression in the RSC layers as previously reported (Figure 1A, and S1) (Ferreira-Fernandes et al., 2019). We could observe that HIPP axon terminals are distributed with high density in the superficial RSC layers II-III, as well as in deeper layer V, and sparsely on all other layers, forming beads-on-a-string structures, typical of axon terminals establishing synapses. On these RSC regions we surgically

implanted two fiberoptic stubs diagonally to target both hemispheres, at a relative reverse angle in distinct AP coordinates (Figure 1B, Methods).

**DNMT sessions in which no light was delivered resulted in no differences in behavioral performance between eArch+ and CTRL animals.**

Freely moving behavior is conditioned by the relations of body shape and movement, with surrounding context structure and dynamics. As such, to make sure all behavioral alterations observed during light-delivery and opsin activation are due solely to this mechanism, it is imperative that pre-illumination DNMP testing is run under the same conditions as those used while we silence RSC-HIPP axon terminals, except for light delivery. The DNMP task includes a sample (encoding) phase, a delay (memory retention) period, and a choice (retrieval) phase, on a T-shaped maze apparatus. During the sample run, animals are rewarded for running the only available arm, with the alternative arm being blocked. This is followed by the delay period, and the "Test" (retrieval) run, during which both arms are made available, and animals are rewarded for choosing the arm previously blocked (i.e., "not matching" the arm sampled previously).

We thus ran the DNMP protocol in two groups of animals expressing either AAV-5-hSyn1-eArch3.0 (eArch+) or an empty construct (CTRL) AAV-5-hSyn1-YFP, with fiberoptic stubs surgically implanted, and patch cords attached to the fiberoptic stubs, no light being delivered (Figure 2 A-B). Under such conditions we found no differences between eArch+ and CTRL animals in either of the three sessions ran (Figure 2C, Table 1, GLMM, p=0.212, Bonferroni *post-hoc* test).

**eArch activation via green light illumination resulted in significantly lower global performance in eArch+, but not CTRL, animals.**

After the "baseline" phase above, we started the "illumination" phase protocol, in which "Test-illuminated" trials (TI), with light delivered through the fiberoptic implants in the Test epoch of the DNMP protocol, were randomly interleaved with "non-illuminated" trials (NI), where no light was delivered in the same ~25 min session (Fig. 2B). In these sessions we observed a significant difference between eArch+ and CTRL animals, with an overall lower performance in eArch+ animals (Fig. 3A, Table 2, GLMM, p=0.006, Bonferroni *post-hoc* test). In addition, we found that the ratio of correct trials in the TI

and NI trial groups (Figure 3B, Tables 2 and 3), were both significantly different from baseline levels (p<0.001), and from their CTRL counterparts (TI, p=0.02, NI, p=0.008), whose performance did not differ from baseline (p=0.252, p=0.184). The fact that more errors were committed in both TI and NI trials of an illuminated session, but only in eArch+ animals, being absent from CTRL animals, implies two things: first, the effect we observe is not the result of light alone acting upon neurons, an effect described earlier in striatal medium spiny (Owen et al., 2019), and CG neurons (Stujenske et al., 2015); second, the effects of eArch light activation of hippocampal inputs onto RSC neurons *outlasted illumination*, affecting subsequent NI trials. We were surprised by this result, as it indicates that the effects of eArch activation on either, or both, the neurophysiology of HIPP-RSC synapses and the ensuing circuit mechanisms involved in DNMP, last longer than would be expected. To directly address this possibility, we analyzed the probability of animals committing errors after illuminated trials. We observed that, overall, illuminated trials were followed by a decreased probability of correct trials on eArch+, but not CTRL, animals (Fig. 3C, Table 4, p=0.015). This is consistent with the effects of eArch activation outlasting illumination by enough time such as to affect subsequent trials.

The effect of light delivery in subsequent non-illuminated trials in sessions where the two types of trials are interleaved within 1-2 minutes, as in the present study, has at least one precedent study in (Robinson et al., 2017). In that study, trials in which a silencer opsin is activated by light are characterized by increased error probability, and abnormal neural activity with decreased temporal selectivity and coding. Thus, both neural activity and behavior of interleaved non-illuminated trials are affected by the activation of the opsin in illuminated trials, leading authors to hypothesize that such persistent disruption of temporal coding in light-off trials is caused by synaptic plasticity events largely specific to the neural activity taking place in the (previous) epoch where light is delivered.

Our findings also imply that the notion of millisecond time-resolved neuron silencing using eArch is not applicable to synaptic transmission and, by consequence, to the ensuing behavioral effects we now investigate.

**Contrary to CTRL animals, eArch+ animals do not correct course following an incorrect decision, in illuminated sessions.**

A fundamental principle of learning theory, and a common empirical finding in reward-based learning experiments, is that following an error in which reward is withheld, animals will correct their course of action. Conversely, following a correct decision, which is rewarded, animals maintain their course of action. This behavior is presumably informed by the association of the previous action with resulting outcome, which they store in memory for the next trial (Mizumori and Jo, 2013). Under this assumption, we predict that in eArch+, but not CTRL, animals, illuminated trials will be followed by persistently defective HIPP-RSC synaptic function, presumably preventing subsequent decisions from anchoring themselves to the previous action-outcome association, even in the absence of light delivery, resulting in a higher probability of errors. To test such prediction, we analyzed the probability of an error trial following error, in illuminated sessions, and found that it is significantly higher in eArch+ animals than in CTRL animals, regardless of whether the present trial is itself illuminated (Figure 4 A, Table 5, $p<0.001$, Bonferroni *post-hoc* test). Importantly, this difference is absent in baseline sessions, in which no illumination occurs (Figure 4B, Table 6, $p=0.172$, Bonferroni *post-hoc* test). This is consistent with our prediction, and suggests that silencing HIPP terminals in RSC interrupts the flow of information from the previous trial, preventing subsequent error-correction for which it is necessary.

**How long does the effect of eArch silencing last?**

The silencing effects of light-mediated eArch activation on hippocampal axon terminals onto RSC neurons outlasted illumination, affecting subsequent trials, and lowering working memory performance. We thus sought to determine how long would such an effect last. According to existing literature, once expressed in CA3-CA1 synaptic terminals, illumination of Arch with green-yellow light results in its activation into a powerful outward-moving proton pump, effectively hyperpolarizing neurons, independent of local membrane voltage (Krol et al., 2019; Miao et al., 2015). Besides this effect, eArch was recently found to reduce the amplitude and slope of fEPSP from right after the onset to over 2 minutes after illumination offset, via a pH increase that is independent of hyperpolarization, and to cause a sustained decrease in synaptic vesicle

release (El-Gaby et al., 2016). Such sustained alkalinization, specific of synaptic boutons, lasts for several minutes post-activation of eArch, and has been replicated on striatal-cortical synapses as well (Mahn et al., 2016). This previous data reporting the effects of eArch activation led us to further investigate, in our experimental conditions, how far in time would eArch behavioral effects last. We thus asked the following question: once one trial is illuminated, how many subsequent trials will be affected? To answer this question, we computed the probability of a correct trial on trial t+1, t+2, t+3, and t+4, following every illuminated trial. Our previous data shows that, overall, non-illuminated trials suffer the effects of illumination, resulting in more errors. However, by selecting the second previous and third previous trial as a trigger for our analysis, we will be biasing our analysis to include a few sequences of one or two interleaved non-illuminated trials, thus possibly attenuating the silencing effect. As depicted in Figure 5 (see also Table 7), the probability of a trial being correct is lower in eArch+, compared to CTRL animals, whenever TI happens in the previous (Figure 5A T-1, p=0.01, Bonferroni *post-hoc* test), the second previous (T-2, p=0.033), and the third previous trial (T-3, p=0.036), such difference becoming non-significant on condition T-4, when the fourth previous trial was illuminated (T-4, p=0.078). Considering that the ITI is circa 20 sec, we can reasonably estimate that some non-illuminated trials occurred between this and the n-previous illuminated one, and that, despite this, the effects of eArch activation are still present, hence after ~3 min. To try and analyze these effects in more detail, without decreasing the sample size to non-interpretable numbers, we quantified performance specifically following sequences of two consecutively non-illuminated trials. We found that after such sequences there is still a significant impairment in eArch+ animals, in agreement with the previous analysis and with the notion that the effects of eArch silencing outlast its activation by light (Figure 5B, p=0.023, Mann-Whitney U).

**Silencing HIPP terminals in RSC hastens decision processing at the Choice Point**
In previous work we have examined the role CG neurons might play during WM, by silencing CG and measuring performance in this same DNMP task, and found that silencing CG interferes with performance, with animals spending less time in the decision processes in DNMP trials (Cruz et al., 2023). We interpreted this decreased latency as hastiness, with animals taking erroneous decisions based on incomplete

cognitive processing. We asked the exact same question in the present work, and quantified latency to the choice point, and latency at the choice point, overall, in error vs correct trials (Figure 6). We found that, overall, CTRL animals spend more time at the choice point than eArch+ animals (Figure 6A, Table 8, p=0.02, Fixed Effect Omnibus Test). Furthermore, this is true for Correct but not Error trials both under NI and TI (Figure 6B, respectively p=0.008 and p=0.045, Simple Effects Parameter Estimates), suggesting that in CTRL animals correct decisions involve more time taken at the choice point, contrary to error trials in which no such difference is found.

**Discussion**

Working memory relies on the coordinated activity of multiple brain regions, allowing relevant information to be stored in cortico-hippocampal circuits in retrievable form as memory, to anchor subsequent behavioral decisions as needed. Such coordination has repeatedly been show to correlate with WM, both during behavior and outside active behavioral decisions (Jadhav et al., 2016; Remondes and Wilson, 2013, 2015), including sleep (Feliciano-Ramos et al., 2023).  In principle, direct synaptic connections between hippocampal and cortical neurons should be interrogated first as the main support of such coordination during behavior. Here, we investigated the necessity of direct hippocampal inputs into retrosplenial cortex (Ferreira-Fernandes et al., 2019), a region generally involved in the encoding of spatial-temporal context, particularly in transferring contextual information between ego- and allocentric reference frames, in the service of decisions (Alexander and Nitz, 2015; Nitzan et al., 2022), during a spatial WM task. We used eArch3.0 expressed in hippocampal neurons via the viral vector AAV5-hSyn1-eArch3.0, under a pan-neuronal promoter, or its "empty" counterpart with only YFP as control, to bilaterally silence hippocampal terminals located at RSC in rats pre-trained to 75% criterium in a DNMP task. We found that silencing HIPP terminals by delivering green light (525 nm) to RSC in eArch+ animals significantly decreased working memory performance. There were no changes in performance during baseline sessions, in which no light is ever delivered through the fiberoptic implants, with eArch+ animals performing at the same level as CTRL. However, once TI sessions start, with randomly interleaved trials with (TI trials) and without light delivery (NI trials), there is a

significantly lower performance exclusively in eArch+ animals. Interestingly, such decreased performance is seen in eArch+ animals, regardless of whether the one trial analyzed is illuminated. Taken as groups, NI and TI trials in eArch+ animals subject to illumination sessions exhibit global performance levels that are lower than both baseline and CTRL animals´ trials (TI and NI). This indicates that the effect of hippocampal terminal silencing in RSC during TI trials affected interleaved NI trials, thus spreading beyond illumination. To explore this effect further, we quantified the probability that any given trial is correct if preceded by a TI trial, and found that it is significantly lower, confirming the previous observation. Furthermore, we asked whether our manipulation would prevent error-corrective behavior, in which animals use action-outcome information after an error trial (i.e., absence of reward), to correct course in the subsequent trial. We found that, contrary to CTRL, eArch+ animals exhibit increased errors even following incorrect responses, contrary to what happens during baseline sessions where no light is ever delivered, and error rates are generally low for both eArch+ and CTRL animals.

A previous study (Robinson et al., 2017) has recently reported that optogenetic silencing entorhinal-hippocampus circuits during WM trials affects performance and neural activity in subsequent trials. This observation suggests that such persistent disruption is caused by synaptic plasticity events related to preceding neural activity affected by the optogenetic manipulation. Furthermore, a recent study recording the effect of eArch activation in CA3-CA1 synaptic responses *in vitro* shows that, besides being a powerful outward-moving proton pump that hyperpolarizes neurons independent of local membrane voltage (Krol et al., 2019; Miao et al., 2015), this manipulation significantly reduces the amplitude and slope of recorded fEPSP. Such reduction persisted from right after the onset of eArch activation to over 2 minutes after illumination offset. This effect is achieved by a pH increase independent of hyperpolarization, leading to a sustained decrease in synaptic vesicle release (El-Gaby et al., 2016). Such sustained alkalinization, specific of synaptic boutons, lasts for several minutes post-activation of eArch, and has been replicated on another study using striatal-cortical synapses (Mahn et al., 2016). Taken together with our data, these findings imply that the temporal resolution of optogenetic neuron silencing, with neurons

effectively unable of generating action potentials, cannot be generalized to the distinct realm of synaptic transmission. We wondered how long the influence of optogenetic silencing of HIPP terminals in RSC lasted during a given session. Since TI and NI trials are randomly interleaved, two and three consecutive trials of the same type (TI or NI) are rare, so we decided to quantify the probability of any trial being correct given that its $n$ preceding trial was illuminated, with $n=1, 2, 3, …$to ask whether, and for how long, performance differences between eArch+ and CTRL animals were present post TI. We found that such influence persists for three trials after illumination, with eArch+ animals still showing significantly lower performance than CTRL by the third trial post-TI, even if the two previous trials are not illuminated. Finally, as we have seen before with optogenetic silencing of CG neurons during DNMP (Cruz et al., 2023), eArch+ animals spend less time at the choice point compared to CTRL. This behavior is in line with hastier, more erroneous, decisions. In CTRL animals specifically, correct trials are characterized by an increased delay at the choice point, as if the correct decision process took place here.

In sum, by reversibly silencing the HIPP synaptic inputs to RSC we have shown that this input is necessary for working memory performance, possibly because it conveys memorized contextual and episodic information about past decisions, namely action-outcome associations, onto medial mesocortical executive decision centers.

**Acknowledgements**

We are indebted to IMM's Bioimaging, Comparative Pathology, and Rodent facilities for critical help. We want to thank all the members of the Remondes Lab for fruitful discussions and help.

Funding reference: UID/BIM/50005/2022, project funded by Fundação para a Ciência e a Tecnologia (FCT)/ Ministério da Ciência, Tecnologia e Ensino Superior (MCTES) through Fundos do Orçamento de Estado.

FCT granted a PhD Fellowship (PD/BD/128107/2016, COVID/BD/151601/2021) to BP-C, and an Exploratory Grant (IF/00201/2013), an IMM Director's Fund Award, an Investigator FCT Position at IMM-JLA (IF/00201/2013), a Research Grant (PTDC/MED-NEU/29325/2017), and the 2022.00811.CEECIND Principal Investigator contract to MR .

**Author contributions**

B P-C and MR designed the experiments, B P-C and PC conducted the experiments and analyzed the data, and MR wrote the manuscript.

**Declaration of Interests**

The authors declare no competing interests.

**FIGURE LEGENDS**

**Figure 1. Viral vector expression and fiberoptic placement (**A) Anatomical distribution of eYFP fluorescence expression in distinct AP coordinates of the retrosplenial cortex after injection in the hippocampus (HIPP). (B) Injection spots in the HIPP (C) Fiberoptic placement two fiberoptic stubs at AP reversed 30 $^{o}$ angles reaching the two hemispheres at two distinct AP levels of RSC.

**Figure 2. Delayed Non-Match to Trajectory protocol and baseline performance** (A) DNMP protocol. Rats explore the unblocked arm in a sample run, then wait in the starting area for a 15 sec delay. Once back on the track, with both arms unblocked, rats receive a reward if they choose the arm opposite the one explored in the sample run. (B) Training schedule, after reaching 75% performance, 3 baseline (no light delivered) sessions are followed by 10 sessions with illuminated trials interleaved with non-illuminated ones. (C) No differences in performance were observed in the baseline sessions (GLMM, p=0.212, *Bonferroni post-hoc test*)

**Figure 3. Optogenetic silencing of HIPP terminals in RSC causes a significant and prolonged decrease in performance** (A) Illumination of RSC during DNMP leads to a significantly lower performance in eArch+, compared to empty-vector CTRL animals (CTRL: 0.81±0.03, ARCH: 0.73±0.08, GLMM, p=0.006, Bonferroni *post-hoc* test). (B) In eArchT+ animals, we found no differences between performance in TI and NI trials taken as groups, both significantly lower than both baseline (p<0.001), and the corresponding trial groups in CTRL animals (TI, p=0.02, NI, p=0.008). In CTRL animals no differences were found among the three groups (p=0.252 and 0.184) (C) Illumination on any given trial resulted in lower performance in the subsequent trial in eArch+, but not CTRL, animals (p=0.015), regardless of whether the illuminated trial is a correct or an incorrect trial. All summary data depicted is mean+/-SD.

**Figure 4. Optogenetic silencing of HIPP terminals in RSC causes persistent error increases**. (A) eArch+ animals exhibit persistent errors (p<0.001, Bonferroni *post-hoc* test). (B) Persistent errors are absent in baseline (no light delivered) sessions (p=0.172, Bonferroni *post-hoc* test). All summary data depicted is mean+/-SD.

**Figure 5. Behavioral impairments resulting from optogenetic silencing of HIPP terminals in RSC outlast illumination** (A) eArch+ animals exhibit decreased

performance once silencing occurred at T-1 (p=0.01, Bonferroni *post-hoc* test), T-2 (p=0.033), T-3 (p=0.036), not T-4 (p=0.078). (B) Two consecutively non-illuminated trials after illumination still result in decreased performance on the subsequent trial (p=0.023, Mann-Whitney U). All summary data depicted is mean+/-SD.

**Figure 6. Animals spend more time at the choice point in Correct trials unless HIPP terminals in RSC are silenced** (A) CTRL animals spend more time at the choice point than eArch+ animals (p=0.02, Fixed Effect Omnibus Test). (B) This is true specifically of Correct trials both in NI and TI (respectively p=0.008 and p=0.045, Simple Effects Parameter Estimates). All summary data depicted is mean+/-SD.

## LEAD CONTACT AND MATERIALS AVAILABILITY

Further information and requests for resources and reagents should be directed to and will be fulfilled by Lead Contact, Miguel Remondes, DVM, PhD (mremondes@medicina.ulisboa.pt). This study did not generate new unique reagents.

## METHODS

### RATS

We used 16 (8 eArch and 8 CTRL) male Long-Evans rats aged 4-6 months and weighing 430-600g (Charles River Laboratories), housed individually and kept under a 12hr light/dark cycle, gradually deprived to 80-85% of their free-feeding body weight. All procedures were performed in accordance with EU guidelines, approved by the Portuguese National Authority for Animal Health, as well as by the animal well-being office at Instituto de Medicina Molecular João Lobo Antunes.

### BEHAVIORAL APPARATUS AND EQUIPMENT

Experiments were performed on a black acrylic T-maze comprising a central 170 cm long arm, a 25x30cm start platform, and a horizontal arm (88cm long) extending equally on both sides with two reward ports at their ends (2 cm diameter) where chocolate milk reward was delivered. A black plastic resting box (57x39x42 cm) was located near the start region to place the rats during the delay and inter-trial interval (ITI) periods. A cable tray above the maze supported the video recording and optogenetic stimulation systems. Both were linked to a computer situated in an adjacent room outside the

behavior room. The Bonsai software (Lopes et al., 2015) was used to process and control video recording, position tracking, reward delivery, and optogenetic stimulation onset.

To record the behavioral sessions, a Flea 3 PointGrey™ camera was used, capturing at a rate of 30fps. The optogenetic stimulation system used was the PlexBright system (Plexon, Dallas, TX, USA), comprising several components. The system consisted of an LD-1 Single Channel LED Driver set at 240 mA, connected to a Dual LED Commutator placed above the maze. This commutator was linked to two green 525nm LED modules, each attached to a 2-meter-long 0.66 NA patch cable with a FC ferrule tip and stainless-steel cladding. This patch tip was connected to an implanted lambda fiberoptic stub (200/230μm, 0.66 NA) with a 2mm active length and 5 mm implant length (Optogenix™) via a ceramic sleeve. Additionally, a red 5 mm LED was used to track the rats' xy position coordinates. This LED was connected to the tip of the patch cable, close to the fiberoptic stub implants.

**SURGICAL PROCEDURE: VIRAL INJECTIONS AND FIBEROPTIC STUB IMPLANTS**

Animals were bilaterally injected with the viral vector AAV5.hSyn-eArch3.0-eYFP and the respective control AAV5.hSyn-eYFP (University of North Carolina at Chapel Hill Vector Core). Viral titers were 4.4 x $10^{12}$ vg/ml and 3.3 x $10^{12}$ vg/ml, respectively. Before surgery, animals were placed in an induction chamber saturated with 5% isoflurane until absence of the righting reflex. Once the animals were deeply anesthetized, they were weighed and transferred to a heating pad. An intraperitoneal injection of a ketamine (37.5mg/kg) and medetomidine (0.5 mg/kg) cocktail (½ dose) was administered, as well as a subcutaneous injection of carprofen (5mg/kg, ½ dose), and Ringer's lactate every hour to maintain hydration levels. The heads were shaved, lidocaine applied to the ears, and eyes covered with a protection gel (Lubrithal) to prevent drying of the cornea. The animals were installed in the stereotaxic frame secured by ear bars. Anesthesia was maintained with 1–3% isoflurane in oxygen until the end of the procedure. Once animals were placed on the frame, the surgical site was scrubbed with 10% povidone iodine alternated with 70% ethanol three times before an incision was made to expose the skull. Bregma and lambda were carefully identified and marked on the skull surface.

These two landmarks were used to level the skull dorsoventrally. Once the skull was leveled, the craniotomy coordinates were marked on the bone. Before drilling, the bone surface was treated with Baytril to minimize infection.

A high-power microdrill tool was used to drill the location of four anchoring screws, on the lateral edges of the parietal bones, and to perform three craniotomies: two single-hole craniotomies (-5, ±4,), one on each hemisphere, for the microinjection spots; and one craniotomy for the fiberoptic stubs covering both hemispheres. The latter was drilled in oval shape covering the fiberoptic stub implantation coordinates (-3,+1), (-5,-1). Loose bone pieces were carefully removed with tweezers. To prevent the brain from swelling while the next steps were performed, the craniotomies were covered with hemostatic sponge previously soaked in saline solution.

The viral construct was injected using a Narishige© manual microinjection control system operating a glass micropipette plunger, and both held on the stereotaxic frame. The micropipette was filled with mineral oil and placed in the microinjector, after which the viral suspension was aspirated (0.8μl). Using bregma as a reference, the pipette was carefully positioned over the microinjection coordinates and slowly lowered until reaching the targets: -5, ±4, -2.6 and -5, ±4, -2.8, where 0.4μl of viral suspension was injected after a 5 min resting period, at a rate of 100 nl/min. Once each injection was done, the micropipette was left in place for 10min before being slowly retracted. The craniotomies were then covered with hemostatic sponge previously soaked in saline solution and left in place to serve as a barrier between the brain and the dental acrylic (Kerr© TAB 2000 fast setting) applied afterwards.

In the fiberoptic stub craniotomy the dura was removed to expose the brain surface over each implant coordinate (-3,+1)(-5,-1). The fiberoptic stubs were secured to the stereotaxic frame, positioned in a 30° angle, and lowered across the two hemispheres to their final target position (-3,+1,-2.86)(-5,-1,-2,86). Following placement of the first fiberoptic stub it was secured using dental acrylic cement (Kerr TAB 2000 fast setting) connecting it to the anchoring screws, while the opposite hemisphere was protected with hemostatic sponge. The same procedure was followed for the second fiberoptic stub.

Once the dental acrylic was cured, the skin was sutured using absorbable sutures, and 10% povidone iodine solution was applied around the wound. Carprofen (5mg/kg, ½ dose) was administered subcutaneously for analgesia. Animals were then placed in a heated cage during the first 24h post-surgery for recovery. During this period, ad libitum food, nutritional gel, and water were provided. For the first 7 days, buprenorphine (0.3 mg/kg), a postoperative analgesic, was administered dissolved in chocolate milk.

## DELAYED NON-MATCHING TO PLACE TASK AND PROTOCOL

During the week before surgery, rats underwent daily handling and were administered chocolate milk via a 1ml syringe for habituation. A week post-surgery, the process was repeated, and rats were habituated to the maze. To facilitate this habituation, chocolate milk drops were scattered across the maze and in the reward ports. Once the rats were drinking chocolate milk from the reward ports, they were habituated to do the same with the patch cords attached. Following habituation, the DNMP training sessions started.

Each session consisted of 28 trials, and each trial comprises three epochs: (1) the sample run, (2) the delay, (3) and the test run. (1) In the Sample Run, one arm was blocked, thereby the rat was forced to visit the only available arm, where a reward awaited; (2) when the rat consumed the reward, it was picked up and placed in the resting box for 15s. During this delay, the arms and the stem near the choice point were cleaned with 70% ethanol; (3) in the Test Run the block from epoch 1 was removed and rats were free to choose  which arm to visit (left or right), and were rewarded for choosing the arm opposite to the Sample arm (i.e., non-matching the sample). After the Test Run, rats were returned to the box for a 20s ITI and a new trial started.

The Sample Run arms were balanced (50% left and 50% right), and pseudo-randomized to avoid more than 3 consecutive runs to the same arm. Rats were scored based on the number of correct and incorrect choices in the Test Run. Throughout the sessions, the position of the experimenter in the room remained constant. Rats were trained until reaching a performance of at least 75% of correct choices in 3 consecutive sessions, after which we introduced optogenetic inhibition.

**OPTOGENETIC INHIBITION PROTOCOL**

The protocol consisted in 10 consecutive sessions of 28 trials, of which 14 were illuminated in the Test epoch (TI) and 14 were not illuminated (NI). This interleaved approach was chosen to prevent compensatory mechanisms. The illuminated trials had the arms pseudo-randomized, ensuring a balanced distribution of left and right sample arms (50% each), and restricted to allow a maximum of 3 consecutive trials. The onset of illumination was triggered to the position of the animal at 120 cm into the central arm and ended when the rat reached the chosen arm. Each rat completed a total of 280 trials, half of which were subjected to optogenetic inhibition. Upon concluding the experiment, the rats were sacrificed, the implant was extracted, and the fiberoptic stubs were connected to the optogenetic system to verify the emission of light through the optic fiber.

**HISTOLOGY**

At the end of the experiments, animals were sacrificed via an isoflurane overdose followed by an intraperitoneal injection of sodium pentobarbital (800 mg/kg). Rats were then transcardially perfused with 250 ml of PBS followed by 500 ml of 10% formalin solution. The brains were dissected and kept in a falcon tube containing 10% formalin solution, covered with aluminum foil, and left at room temperature for 24h. Afterwards, the brains were transferred to a 15% sucrose solution and kept at 4°C overnight. Once sunk, the brains were embedded in gelatin, frozen in 2-Methylbutane liquid nitrogen, and sectioned in 50µm coronal slices using a cryostat (model CM3050 S, Leica). The fixed brain slices were degelatinized and incubated in DAPI 1:700 (Sigma) for 20 min. These slices were then mounted, cover slipped in Mowiol, and left to dry for 24h. The expression of the eArch3.0 viral construct in the hippocampus was confirmed using an AxioZoom V16 fluorescence stereo microscope (Zeiss). The expression of the eArch3.0 viral construct in the RSC was confirmed using a LSM 880 confocal point-scanning microscope with Airyscan (Zeiss). CTRL animals underwent the exact same procedure.

**DATA AND STATISTICAL ANALYSES**

Datasets were manually assembled containing specific details of each run. Within this dataset, each session's runs were detailed with the following specifications: run number, run type (sample or test), illumination condition (TI or NI), and outcome (correct or

incorrect trial). Additionally, the rats' xy positions were tracked via a LED, recorded, and timestamped using Bonsai software (Lopes et al., 2015). Custom Python scripts were employed to compute individual runs from these data, ensuring the accuracy of individual runs while excluding any with position data interference. Performance was computed for each experimental session including baseline sessions, consisting in the three sessions before starting optogenetic inhibition. Performance for illuminated and non-illuminated conditions was calculated as the ratio of correct trials over the total number of illuminated or non-illuminated trials, respectively. Probability computations were conducted for trial accuracy given the previous trial's illumination ($trial_{n-1}$ to $trial_{n-4}$) and for error occurrence given the previous error trial, as follows:

$P$ (correct $trial_n$ | $trial_{n-x}$ illuminated) = $P$ ($trial_n$ is correct $\cap$ $trial_{n-x}$ is illuminated) / $P$ (illuminated $trials_{n-x}$)

$P$ (error $trial_n$ | $trial_{n-1}$ error) = $P$ ($trial_n$ is an error $\cap$ $trial_{n-1}$ is an error) / $P$ (error $trials_{n-1}$)

The time spent at the choice point was determined as the difference between the timestamps of choice point entry and exit. An additional 10cm was added to the region of interest to account for instances when the rat was stationary but engaged in head poking. Time values were Log-transformed to achieve a Gaussian distribution. Normalization to the baseline of log-transformed time in the choice point to the baseline was computed as: [log(latency) – log(baseline mean)] / [log(baseline mean)]. Outliers in time data were identified as values deviating by 2 standard deviations (SD) from the mean and were subsequently removed.

Generalized Linear Mixed Models (GLMM) were employed to analyze the results. Model selection was based on the Akaike Information Criterion (AIC), with the chosen model being the one that converged and exhibited the lowest AIC value. The normality of a distribution was assessed using the Shapiro test along with assessment of residuals using Quantile-Quantile (Q-Q) plots. All the analysis was conducted in Jamovi with the GAMLJ package (Jonathon Love, 2022).

To assess baseline performance differences between eArch and CTRL groups, we used a GLMM with a binomial distribution and logit link function, with individual subjects (i.e., rats) as the random effects variable. The same GLMM, followed by a Bonferroni post-hoc test, was conducted to analyze the ratio of correct trials between groups during

the experiment, and for assessing the trial history effect of illumination. The same tests were used to analyze the trial history effect of errors, using as random effect variables the "session number" and "run number". To compare the groups after two consecutive illuminated trials, a Mann-Whitney U test was applied. For analyzing Log time in the choice point, a GLMM with a Gaussian distribution and identity link function was applied, considering rat as the random effects variable. To determine significant differences between groups we used the Wald Chi-Squared test, and for differences between groups and illumination condition we used simple effects parameter estimates.

**References**

Aggleton, J.P., Hunt, P.R., and Rawlins, J.N.P. (1986). The effects of hippocampal lesions upon spatial and non-spatial tests of working memory. Behav. Brain Res. *19*, 133–146.

Akam, T., Rodrigues-Vaz, I., Marcelo, I., Zhang, X., Pereira, M., Oliveira, R.F., Dayan, P., and Costa, R.M. (2020). The Anterior Cingulate Cortex Predicts Future States to Mediate Model-Based Action Selection. Neuron *109*, 1–15.

Alexander, A.S., and Nitz, D.A. (2015). Retrosplenial cortex maps the conjunction of internal and external spaces. Nat. Neurosci. *18*, 1143–1151.

Alexander, A.S., Carstensen, L.C., Hinman, J.R., Raudies, F., William Chapman, G., and Hasselmo, M.E. (2020). Egocentric boundary vector tuning of the retrosplenial cortex. Sci. Adv. *6*.

Andersen, R., Snyder, L., Bradley, D., and Xing, J. (1997). Multimodal Representation of Space in the Posterior Parietal Cortex and Its Use in Planning Movements. Annu. Rev. Neurosci. *20*, 303–330.

Ariani, G., Wurm, M.F., and Lingnau, A. (2015). Decoding Internally and Externally Driven Movement Plans. J. Neurosci. *35*, 14160–14171.

Bicanski, A., and Burgess, N. (2018). A neural-level model of spatial memory and imagery. Elife *7*.

Brown, J.W., and Braver, T.S. (2005). Learned predictions of error likelihood in the anterior cingulate cortex. Science (80-. ). *307*, 1118–1121.

Burgess, N. (2008). Spatial cognition and the brain. Ann. N. Y. Acad. Sci. *1124*, 77–97.

Carstensen, L.C., Alexander, A.S., Chapman, G.W., Lee, A.J., and Hasselmo, M.E. (2021). Neural responses in retrosplenial cortex associated with environmental alterations. IScience *24*, 103377.

Chen, L.L., Lin, L.H., Green, E.J., Barnes, C.A., and McNaughton, B.L. (1994a). Head-direction cells in the rat posterior cortex - I. anatomical distribution and behavioral modulation. Exp. Brain Res. *101*, 8–23.

Chen, L.L., Lin, L.H., Green, E.J., Barnes, C.A., and McNaughton, B.L. (1994b). Head-direction cells in the rat posterior cortex - I. anatomical distribution and behavioral modulation. Exp. Brain Res. *101*, 8–23.

Clark, B.J., Simmons, C.M., Berkowitz, L.E., and Wilber, A.A. (2018). The retrosplenial-parietal network and reference frame coordination for spatial navigation. Behav. Neurosci. *132*, 416–429.

Cowen, S.L., Davis, G.A., and Nitz, D.A. (2012). Anterior cingulate neurons in the rat map anticipated effort and reward to their associated action sequences . J. Neurophysiol. *107*, 2393–2407.

Cruz, A.S., Cruz, S., and Remondes, M. (2023). Optogenetic silencing of the Anterior Cingulate Cortex impairs working memory in a delayed non-match to trajectory task. BioRxiv 2023.08.30.555490.

El-Gaby, M., Zhang, Y., Wolf, K., Schwiening, C.J., Paulsen, O., and Shipton, O.A. (2016). Archaerhodopsin Selectively and Reversibly Silences Synaptic Transmission through Altered pH. Cell Rep. *16*, 2259.

Feliciano-Ramos, P.A., Galazo, M., Penagos, H., and Wilson, M. (2023). Hippocampal memory reactivation during sleep is correlated with specific cortical states of the retrosplenial and prefrontal cortices. Learn. Mem. *30*, 221–236.

Ferreira-Fernandes, E., Pinto-Correia, B., Quintino, C., and Remondes, M. (2019). A Gradient of Hippocampal Inputs to the Medial Mesocortex. Cell Rep. *29*, 3266-3279.e3.

Gomez, A., Rousset, S., and Baciu, M. (2009). Egocentric-updating during navigation

facilitates episodic memory retrieval. Acta Psychol. (Amst). *132*, 221–227.

Gomez, A., Cerles, M., Rousset, S., Le Bas, J.-F., and Baciu, M. (2013). Ongoing egocentric spatial processing during learning of non-spatial information results in temporal-parietal activity during retrieval. Front. Psychol. *4*.

Hafting, T., Fyhn, M., Molden, S., Moser, M.B., and Moser, E.I. (2005). Microstructure of a spatial map in the entorhinal cortex. Nature *436*, 801–806.

Jadhav, S.P., Rothschild, G., Roumis, D.K., and Frank, L.M. (2016). Coordinated Excitation and Inhibition of Prefrontal Ensembles during Awake Hippocampal Sharp-Wave Ripple Events. Neuron.

Jin, D.Z., Fujii, N., and Graybiel, A.M. (2009). Neural representation of time in cortico-basal ganglia circuits. Proc. Natl. Acad. Sci. U. S. A. *106*, 19156–19161.

Kimura, A., Donishi, T., Okamoto, K., and Tamai, Y. (2004). Efferent connections of "posterodorsal" auditory area in the rat cortex: Implications for auditory spatial processing. Neuroscience *128*, 399–419.

Krol, A., Lopez-Huerta, V.G., Corey, T.E.C., Deisseroth, K., Ting, J.T., and Feng, G. (2019). Two eARCHT3.0 lines for optogenetic silencing of dopaminergic and serotonergic neurons. Front. Neural Circuits *13*, 417848.

L Fenno, O.Y.K.D. (2010). The Development and Application of Optogenetics. Annu. Rev. Neurosci. *34*, 110301101035033.

Lopes, G., Bonacchi, N., Frazão, J., Neto, J.P., Atallah, B. V., Soares, S., Moreira, L., Matias, S., Itskov, P.M., Correia, P.A., et al. (2015). Bonsai: an event-based framework for processing and controlling data streams. Front Neuroinform *9*, 7–14.

Maguire, E.A. (2001). Neuroimaging studies of autobiographical event memory. Philos. Trans. R. Soc. London. Ser. B Biol. Sci. *356*, 1441–1451.

Mahn, M., Prigge, M., Ron, S., Levy, R., and Yizhar, O. (2016). Biophysical constraints of optogenetic inhibition at presynaptic terminals. Nat. Neurosci. *19*, 554.

Miao, C., Cao, Q., Ito, H.T., Yamahachi, H., Witter, M.P., Moser, M.B., and Moser, E.I. (2015). Hippocampal Remapping after Partial Inactivation of the Medial Entorhinal

Cortex. Neuron.

Michelmann, S., Price, A.R., Aubrey, B., Doyle, W.K., Friedman, D., Dugan, P., Devinsky, O., Devore, S., Flinker, A., Hasson, U., et al. (2020). Moment-by-moment tracking of naturalistic learning and its underlying hippocampo-cortical interactions. BioRxiv.

Miller, A.M.P., Mau, W., and Smith, D.M. (2019). Retrosplenial Cortical Representations of Space and Future Goal Locations Develop with Learning. Curr. Biol. *29*, 2083-2090.e4.

Mizumori, S.J.Y., and Jo, Y.S. (2013). Homeostatic regulation of memory systems and adaptive decisions. Hippocampus *23*, 1103–1124.

Morris, R.G.M., Garrud, P., Rawlins, J.N.P., and O'Keefe, J. (1982). Place navigation impaired in rats with hippocampal lesions. Nature *297*, 681–683.

NADEL, L. (1990). Varieties of Spatial Cognition: Psychobiological Considerations. Ann. N. Y. Acad. Sci. *608*, 613–636.

Nakamura, K. (1999). No Title. 2503–2517.

Nitzan, N., McKenzie, S., Beed, P., English, D.F., Oldani, S., Tukker, J.J., Buzsáki, G., and Schmitz, D. (2020). Propagation of hippocampal ripples to the neocortex by way of a subiculum-retrosplenial pathway. Nat. Commun. *11*.

Nitzan, N., Swanson, R., Schmitz, D., and Buzsáki, G. (2022). Brain-wide interactions during hippocampal sharp wave ripples. Proc. Natl Acad. Sci. USA *119*, e2200931119.

O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. Brain Res. *34*, 171–175.

Owen, S.F., Liu, M.H., and Kreitzer, A.C. (2019). Thermal constraints on in vivo optogenetic manipulations. Nat. Neurosci. 2019 227 *22*, 1061–1065.

Remondes, M., and Wilson, M.A. (2013). Cingulate-hippocampus coherence and trajectory coding in a sequential choice task. Neuron *80*, 1277–1289.

Remondes, M., and Wilson, M.A. (2015). Slow-γ Rhythms Coordinate Cingulate Cortical Responses to Hippocampal Sharp-Wave Ripples during Wakefulness. Cell Rep. *13*,

1327–1335.

Robinson, N.T.M., Priestley, J.B., Rueckemann, J.W., Garcia, A.D., Smeglin, V.A., Marino, F.A., and Eichenbaum, H. (2017). Medial Entorhinal Cortex Selectively Supports Temporal Coding by Hippocampal Neurons. Neuron *94*, 677-688.e6.

Scoville, W.B., Milner, B., and Scoville WB, M.B. (1957). Loss of recent memory after bilateral hippocampal lesions. J Neurol Neurosurg Psychiatry *20*, 11–21.

Shaw, C., and Aggleton, J.P. (1995). Evidence for the Independence of Recognition and Recency Memory in Amnesic Subjects. Cortex *31*, 57–71.

Shaw, C.L., Watson, G.D.R., Hallock, H.L., Cline, K.M., and Griffin, A.L. (2013). The role of the medial prefrontal cortex in the acquisition, retention, and reversal of a tactile visuospatial conditional discrimination task. Behav. Brain Res. *236*, 94–101.

Sober, S.J., and Sabes, P.N. (2003). Multisensory integration during motor planning. J. Neurosci. *23*, 6982–6992.

Stackman, R.W., and Taube, J.S. (1998). Firing properties of rat lateral mammillary single units: Head direction, head pitch, and angular head velocity. J. Neurosci. *18*, 9020–9037.

Stujenske, J.M., Spellman, T., and Gordon, J.A. (2015). Modeling the Spatiotemporal Dynamics of Light and Heat Propagation for In Vivo Optogenetics. Cell Rep. *12*, 525–534.

Taube, J.S., Muller, R.U., and Ranck, J.B. (1990). Head-direction cells recorded from the postsubiculum in freely moving rats. I. Description and quantitative analysis. J. Neurosci. *10*, 420–435.

Wilson, D.I.G., Langston, R.F., Schlesiger, M.I., Wagner, M., Watanabe, S., and Ainge, J.A. (2013). Lateral entorhinal cortex is critical for novel object-context recognition. Hippocampus *23*, 352–366.

Yamamoto, J., Suh, J., Takeuchi, D., and Tonegawa, S. (2014). Successful execution of working memory linked to synchronized high-frequency gamma oscillations. Cell *157*, 845–857.

Yamawaki, N., Li, X., Lambot, L., Ren, L.Y., Radulovic, J., and Shepherd, G.M.G. (2019). Long-range inhibitory intersection of a retrosplenial thalamocortical circuit by apical tuft-targeting CA1 neurons. Nat. Neurosci. 22, 618–626.

B

C

# A

## SAMPLE RUN

## TEST RUN

**Previously visited**

## Delay Period

# B

| Learning ➡ | Baseline Sessions ➡ | Illumination Sessions |
|---|---|---|

| | 3 Sessions | 280 trials |
|---|---|---|
| | >75% performance | 1 session: 28 trials |
| | With patch cords | Interleaved Illumination |

ARCH Rats
CTRL Rats

| 1 | 2 | 3 | 4 | ... | 27 | 28 |

# C

A

B