Cholinergic agonist PNU282987 induces pro-inflammatory gene expression

via the MCOLN1-TFEB pathway in murine macrophages.

Havisha H. Honwad[1,2], Mehran Najibi[3], Balazs Koscso[4,5], Milena Bogunovic[4], Javier E.

Irazoqui[1*]

[1] Department of Microbiology and Physiological Systems and

[2] Immunology & Microbiology Program, University of Massachusetts Chan Medical School,

Worcester, USA

[3] Department of Pathology, New York University Grossman School of Medicine, New York,

USA

[4] Department of Pathology, University of Massachusetts Chan Medical School, Worcester, USA

[5] Obsidian Therapeutics, Cambridge, USA

*Correspondence: Javier.Irazoqui@umassmed.edu

ABSTRACT

Neurotransmitter signaling in macrophages is a modulatory mechanism during inflammation.

Treatment of macrophages with acetylcholine, or agonistic molecules carbachol and

PNU282987, considered to be specific for the Alpha7 nicotinic ACh receptor (α7NAChR),

represses cytokine expression in inflammatory disease models, leading to the idea that

α7NAChR stimulation could modulate inflammation in health and disease. However, their

effects on resting macrophages are largely unaddressed. This study assesses the impact of

PNU282987 on macrophage phenotypes *in vitro*, focusing on bone marrow-derived macrophages

(BMDMs) and RAW264.7 cells. We found that PNU282987 induces a specific pro-inflammatory

gene expression profile, elevating *Ifnb1, Il1b, Il6,* and *Tnf* levels. However, this did not translate

to increased cytokine secretion. Unexpectedly, both BMDM and RAW264.7 cells showed no

detectable α7 nicotinic acetylcholine receptor expression, challenging the proposed action

mechanism of PNU282987. Instead, our data suggest a non-receptor-mediated pathway, where

PNU282987 may activate the stress-responsive TFEB transcription factor via ROS generation,

implicating the MCOLN1-calcineurin pathway. Our findings reveal that the primary action of

PNU282987 in resting macrophages is through non-receptor-mediated pro-inflammatory gene

expression and TFEB activation, offering new insights into its pharmacological effects beyond

the conventional receptor-based mechanisms.

INTRODUCTION

The neurotransmitter acetylcholine (ACh) modulates inflammatory responses in what is known

as the cholinergic anti-inflammatory pathway (CAIP) (1, 2). Nicotinic ACh receptors composed

of α7 subunits (α7NAChR) expressed in macrophages have been implicated in mediating the

CAIP (3, 4), mainly through inhibition of NF-κB activity (5, 6). Exactly how α7NAChR

represses NF-κB is not completely understood, but the mechanism appears to include a

JAK2/STAT3 pathway (7). Nonetheless, α7NAChR activation has been proposed as a therapeutic

strategy for several inflammatory diseases, including sepsis (8, 9), ulcerative colitis (10, 11),

arthritis (12, 13), lung inflammation (14, 15), cystic fibrosis (16), Type 2 diabetes (17, 18), and

Parkinson's disease(19). α7NAChR agonists, such as PNU282987, have been used to suppress

pro-inflammatory responses of various cell types, including macrophages, in models for these

diseases (5, 7, 20, 21). PNU282987, a selective α7NAChR agonist (22, 23) has been shown to

protect against sepsis or LPS induced lung injury (24-26) or alleviate DSS-induced colitis

inflammation (27) by inhibiting inflammatory mediators in model systems. Nonetheless, to

therapeutically deploy α7NAChR agonists to treat inflammation, it is necessary to understand

their effects on cells in their resting state, yet these are vastly understudied. Importantly, it is

currently unknown if α7NAChR agonists trigger cellular stress, which could affect their safety

profiles.

Transcription factor EB (TFEB) is a sensitive reporter of cell stress. Known as a master

regulator of lysosome biogenesis and autophagy regulation (28, 29), TFEB was recently shown

to participate in the integrated stress response (30, 31). When activated by starvation, infection,

or oxidative stress, TFEB translocates from the cytosol into the nucleus to drive gene expression

(32-36). Although TFEB is broadly expressed in mice and humans, its downstream target genes

appear to be largely cell type-specific through unknown mechanisms (37). Importantly,

understanding of TFEB functions in macrophages is in its infancy.

To better understand the effects of α7NAChR agonists on innate immunity, we examined

PNU282987-treated murine bone marrow-derived macrophages (BMDMs) and a macrophage-

like cell line. Surprisingly, PNU282987 drove a pro-inflammatory gene expression program in

both systems, despite a lack of convincing evidence of α7NAChR expression in these cells. As

an alternative mechanism of action, we found that PNU282987 triggered TFEB activation in

both cell types, and that the pro-inflammatory response was highly TFEB-dependent. Moreover,

we detected higher levels of reactive oxygen species (ROS) in PNU282987-treated cells

compared to controls, suggesting that PNU282987 might activate TFEB by oxidative stress.

Further analysis of the previously described ROS-MCOLN1-calcineurin pathway of TFEB

activation (38) was consistent with this alternative mechanism of action.

MATERIALS AND METHODS

Mice

Mouse colony maintenance and procedures were performed in accordance with the Guidelines

for the Care and Use of Laboratory Animals, published by the US National Institutes of Health,

and were approved by the Institutional Animal Care and Use Committee (IACUC) of the

University of Massachusetts Chan Medical School. Wild type mice, Strain: C57BL/6J (Stock #:

000664) and *Chrna7*$^{-/-}$ mice, Strain: B6. 129S7-*Chrna7*^{tm1Bay}/J (Stock #: 003232) mice were

ordered from The Jackson Laboratory. *Tfeb*$^{fl/fl}$ mice were a gift from Dr. Andrea Ballabio,

TIGEM, Naples, Italy (29). Tfebfl/fl *LysM-Cre*$^{\pm}$ mice were bred in house (38). BMDM

genotyping for wild type and *Chrna7*$^{-/-}$ BMDMs was performed by following steps described in

Protocol 28408: Standard PCR Assay - Chrna7<tm1Bay>-Alternate 1 for Jackson lab stock No:

003232.

Isolation and differentiation of primary murine BMDMs

Bone marrow cells were isolated from separated femurs and tibias from euthanized 8-12-week-

old mice as previously described (38). Isolated bone marrow cells from C57BL/6, *Chrna7*$^{-/-}$,

Tfeb$^{fl/fl}$ and *Tfeb*$^{\Delta LysM}$ mice were cultured in BMDM differentiation medium: DMEM containing

4.5 g/L glucose, L-glutamine (Corning, 10-013-CV), sodium pyruvate supplemented with 20%

conditioned medium from L929 cells, 10% FBS, 1% AA for 7 days. Cells were seeded in non-

tissue culture treated Petri dishes (Fisherbrand, FB0875712) and incubated at 37°C, in a

humidified chamber with 5% CO_2. Cells were seeded for experiments after 7 days of

differentiation.

Cell culture and transfection

G418 sulfate (Life Technologies, 10131) was used for selection and stable cells were selected by

FACS 10 days after selection. RAW264.7 and *Tfeb^-/- Tfe3^-/-* RAW264.7 cells, a gift from Dr.

Rosa Puertollano, NHLBI (39) and iBMDMs (38) were grown in Dulbecco's Modified Eagle's

Medium (DMEM) containing 4.5 g/L glucose, L-glutamine (Corning, 10-013-CV), sodium

pyruvate supplemented with 10% Fetal Bovine Serum (FBS; Gibco, 16000044) and 1%

Antibiotic-Antimycotic (Gibco, 15240062). Cells were grown at conditions of 37°C, in a

humidified chamber with 5% CO_2 and passaged 4 to 10 times.

Drugs and Reagents

PNU282987 (Sigma-Aldrich, P6499) powder form was carefully dissolved in DMSO (Sigma-

Aldrich, D2438) - sterile, filtered, cell culture grade, Bio Performance certified, meets EP and

USP specifications thus, devoid of endotoxin or pro-inflammatory contaminants; Carbachol

(Sigma-Aldrich, 212385); MG624 Cayman Chemical Company, 24298); LPS from *S. enterica*

serotype Typhimurium (Sigma-Aldrich, L6143-1 MG); Torin-1 (Tocris, 427); FK-506 (Cayman

Chemical Company, 10,007,965); BAPTA-AM (Cayman Chemical Company, 15,551); kb-NB-

142-70 (Cayman Chemical Company, 18,002); α-bungarotoxin Alexa Fluor 647 conjugate

(Thermo Fisher Scientific, B35450)

Immunofluorescence and fluorescence imaging

Cells were seeded in 96-Well Optical Bottom Plates (Thermo Scientific, 165305) and incubated

overnight to attain cell confluency of 0.06×10^6 for experiments. Following experiments with

iBMDM GFP-TFEB cells, fixation step was performed using 4% paraformaldehyde (PFA)

(Sigma Aldrich, 158127) and cells were incubated with Hoechst stain (Anaspec, AS-83218) for

20 min at room temperature as nuclear staining. For immunofluorescence, cells were washed 3

times with PBS (Gibco Life Technologies,10010) for 5 min, permeabilized with 0.1% Triton X in

PBS at on shaker at room temperature for 5 min, washed 3 times with PBS and blocked with 5%

bovine serum albumin (Sigma Aldrich, A9647) in PBS for 1 h. After washing with PBS (3

times), cells were incubated with primary antibody for 2 h Anti-TFEB Antibody (Abcam,

ab2636). After washing 3 times in PBS, cells were incubated with the fluorescently labelled

secondary antibody (Thermo Fisher Scientific, A11058) along with Hoechst stain (Anaspec, AS-

83,218) for 1 h at room temperature. For LysoTracker staining, cells were incubated with 100

nM LysoTracker DND-99 (Thermo Fisher, L7528) added to cell culture media for 30 min before

fixation step, according to manufacturer's instructions. Image acquisition for calculating TFEB

N:C ratio and for Lysotracker mean fluorescence intensity (MFI) was performed using Lionheart

FX automated microscope (Biotek) and Gen5 software (version 3.13).

Determination of N:C ratio

Nucleus to cytoplasm (N:C) ratio for TFEB was measured using CellPofiler (40), following our

previously described protocol (38). In brief, we used the 3_channels_pipeline.cppipe. which

uses nuclear stain (Hoechst) as primary object or "seed" region for building image segmentation

masks to distinguish individual cells by marking boundaries, define subcellular regions and

generate specific fluorescent intensity values. The pipeline calculated N:C ratio as mean

fluorescent intensity of nucleus relative to mean fluorescence intensity of the cytosol.

RT-qPCR

Cells were lysed using Trizol Reagent (1 mL per 1.5 ml microcentrifuge tube). 100 μL

chloroform was added and samples were centrifuged at 12,000 rpm to separate the RNA

containing aqueous phase. The aqueous layer was transferred to Purelink RNA mini kit column

and total RNA was extracted according to manufacturer's instructions. cDNA was prepared using

iScript gDNA clear cDNA synthesis kit (Bio-Rad, 1725035). Diluted cDNA was used for RT-

qPCR, performed using SYBR Freen Super mix (Bio-Rad) and ViiA7 Reat-Time qPCR system

(Applied Bioscience). CT values were obtained for test genes and relative expression was plotted

following normalization to reference gene *Gapdh*.

RNA sequencing and differential expression analysis

BMDMs differentiated from WT, *Chrna7*$^{-/-}$, *Tfeb*$^{fl/fl}$, *Tfeb*$^{\Delta LysM}$ mice were treated with DMSO

control or PNU282987 (50 µL) for 2 h. Total RNA was extracted from samples as described

above using Purelink RNA mini kit (Thermo Fisher Scientific, 12183018A). Whole cell RNA

from 4 replicates per condition were sent to Azenta/ Genewiz for polyA selection, library

preparation and sequencing using Illumina Nova-Seq or HiSeq platform, 2x150bp, ~350M PE

reads. Azenta/Genewiz provided raw data in the form of reads in FASTQ format. FASTQ files

were analyzed using the RNA-seq pipeline (Version 1.6.0) on ViaFoundry software (V1.6.5)

(41). Using the pipeline, FASTQ paired end reads were trimmed, quality filtered by FastQC,

followed by 3' adapter sequence removal and aligned to *Mus musculus* reference genome mm10

using STAR and RSEM was used to estimate gene and isoform levels. DEBrowser (v1.28.0) was

used for gene count normalization, data analyses, generate visual graphs of differentially

expressed genes like volcano plots. Differentially expressed gene sets were considered

significant for fold change $\geq$ 1.5; Padj $\leq$ 0.01 and were used as input for g:Profiler (42).

Comparison of DE genes with public GEO RNA-Seq datasets was performed using Enrichr, a

web-based tool used for enrichment analysis that compares gene lists with gene set libraries of

known biological function (43). Venn diagrams were made using DeepVenn (44).

ELISA

Secreted levels of TNF-α (DY410-05), IL-1β (DY401-05), IL-6 (DY406-05), IFN-β (DY8234-05), IL-10 (DY417-05), TNFSF15 (DY-1896) were measured using kits purchased from R&D Systems. Briefly, 96-well plates were coated with capture antibody and incubated at room temperature overnight. Plates were washed with Tris buffered saline (TBS) with 0.5% Tween 20 (Bethyl Laboratories, E106). Wells were blocked for 1 h with Reagent Diluent - TBS containing 1% bovine serum albumin (BSA) (Bethyl Laboratories, E106). After washing, samples or standards diluted in Reagent Diluent were added and plates were incubated for 2 h at room temperature. Detection antibody was added following washing and plates were incubated for 2 h. After washing, Streptavidin-horseradish peroxidase (Strep-HRP) was added for 20 min. TMB One Component HRP Microwell substrate (Bethyl Lab, E102) was added, and reaction was stopped using ELISA Stop Solution (Bethly Laboratories, E115) Optical density (OD) was measured immediately at 450 nm as well as 540 nm using a SpectraMax Plus 384 Microplate Reader with Softmax Pro software. OD measured at 540 nm was subtracted from 450 nm to correct for optical imperfections as per manufacturer's guidelines. Concentrations of the samples (pg/μL) were interpolated from respective standard curve using GraphPad Prism 10.

Cell viability assay

Cell viability was determined using CellTiter-Glo 2.0 Assay (Promega, G7570), according to the manufacturer's instructions. Luminescence was measured using Biotek Synergy H4 Plate Reader Gen5 (version 3.14).

ROS/Superoxide detection assay

Cellular ROS/Superoxide Detection Assay Kit (Abcam, ab139476) was used according to manufacturer's instructions. Mean fluorescence intensity (MFI) was visualized on a Lionheart microscope (Biotek) and measured using Gen5 Data Analysis Software (Version 3.13).

siRNA knockdown

siRNA reagents were purchased from Dharmacon RNAi Technologies: siGENOME NonTargeting siRNA #1 (D-001210-01-05), siGENOME Mouse *Ppp3cb* (19,056) siRNA (M-063545-00-0005), siGENOME Mouse *Ppp3r1* (19,058) siRNA (M-040744-01-0005), siGENOME Mouse *Mcoln1* (94178) siRNA (M-044469-00-515 0005). For transfection, we used Lipofectamine RNAiMAX (Thermo Scientific, 13,778,030), according to manufacturer's instructions.

Western Blotting

Cell samples were washed 3 times with PBS, harvested, and lysed with 1X SDS sample buffer Blue Loading Pack (Cell Signaling Technology, 7722) at 100 μl per well of 6-well plate. Lysates were heated at 100°C for 5 min and then centrifuged for 5 min. The supernatant was collected and sonicated, gel electrophoresis was performed using Novex 4–20% Tris-Glycine Mini Gels (Invitrogen, XP04200BOX), and were then transferred onto nitrocellulose (Life Technologies, LC2009). After washing with TBS (Life Technologies, 28,358) for 5 min, membranes were soaked in blocking buffer containing 1X TBS with 5% BSA for 1 h at room temperature. After 3 washes with TBS-Tween 20 (Life Technologies, 28,360), membranes were incubated overnight at 4°C with primary antibodies and gentle agitation. Next, membranes were washed thrice with TBS-Tween 20 and incubated with HRP-conjugated secondary antibody (Cell Signaling Technology, 7074 1:2000) for 1 h at room temperature with gentle agitation. Membranes were

then washed with TBSTween20 and incubated with LumiGLO® (Cell Signaling Technology,

7003) for 1 min and imaged using Gel box imager (Syngene) GeneSys software. Primary

antibodies and dilutions were as follows: ACTB/β-actin antibody (Cell Signaling Technology,

4967; 1:1000), α7nAChR antibody (Abcam ab216485, 1:500)

α-bungarotoxin staining and flow cytometry

α-bungarotoxin, Alexa Fluor 647 Conjugate (Thermo Fisher Scientific, B35450) was used

according to manufacturer's guidelines to stain murine peritoneal macrophages and muscularis

macrophages, isolated according to previously described methods (45) by flow cytometry-based

cell sorting (46). Data were analyzed by FlowJo.

Mouse whole brain lysate for Western Blot

In brief, Frozen brain tissue from adult 9-month-old C57Bl/6J mice were homogenized in Cell

Lysis Buffer (EPX-99999-000) and centrifuged at 4°C for 20 min at 18000 x g to obtain brain

lysate.

Determination of bactericidal activity

Escherichia coli DH5α was purchased (ThermoFisher Scientific,18265017). *Staphylococcus*

aureus NCTC8325 is a gift from Fred Ausubel, MGH Research Institute, USA. Bacteria were

grown overnight at 37°C in TSB (Sigma-Aldrich, T8907) supplemented with 10 μg/ml nalidixic

acid for *S. aureus*, or LB medium (Fisher Scientific, BP1425-500) for E. coli DH5α. Cultures

were diluted 1:50 in the respective media the next day and grown at 37°C to mid-exponential

phase. WT and *Tfe3 Tfeb* dKO RAW 264.7 cells were infected with *S. aureus* (MOI = 10) and

DH5α *E. coli* (MOI = 50) and incubated at 37°C for 30 mins as previously described (38).

Lysates were serially diluted, aliquots were plated on TSA (Difco, BD, 236,950)-nalidixic acid

agar (*S. aureus*) or LB agar (*E. coli*) and incubated overnight at 37°C, and colonies were counted

for colony-forming units/ml calculation.

Statistical analysis

Quantitative data comparisons were analyzed using statistical tests, performed as indicated in

each Figure Legend. $P \leq 0.05$ were judged to be significant. For normally distributed data and

comparisons with a single condition, we used the two-sample two-sided t-test. For multiple

comparisons, we used one-way ANOVA followed by Tukey's post-hoc test and Two-way

ANOVA and Sidak's multiple comparison tests.

RESULTS

PNU282987 induces bone marrow-derived macrophages to express pro-inflammatory

genes

To understand the effects of PNU282987 treatment on macrophage phenotype, we

exposed *in vitro*-differentiated bone marrow-derived macrophages (BMDMs) to 50 μM

PNU282987 for 2 h. We chose an early timepoint of exposure to focus on the initial primary

response triggered by this treatment and to avoid secondary effects due to long-term treatment.

Immediately following 2 h exposure, we collected RNA and performed RNA-seq to identify

differentially expressed genes between PNU282987-treated cells and DMSO-treated control.

PNU282987-treated cells showed 572 induced genes and 119 repressed genes (**Fig. 1A, Table**

S1). The induced genes were enriched for various KEGG pathways, most of which were related

to pro-inflammatory signaling (*e.g.* "TNF signaling pathway", "Cytokine-cytokine receptor

interaction", "NF-κB signaling pathway", and "Lipid and atherosclerosis", **Fig. 1B** and **Table**

S2). Consistently, complementary Gene Ontology (GO) and Reactome enrichment analyses also

showed enrichment of pro-inflammatory categories (**Table S2**). In contrast, the repressed genes

were not significantly enriched for any KEGG pathways or GO categories. These results

suggested that PNU282987 induced a pro-inflammatory gene expression profile in BMDMs.

Among the induced genes, *Ifnb1, Il1b, Il6, Tnf* and *Il10* stood out. These genes encode

the precursors for cytokines IFN-β1, IL-1β, IL-6, TNF-α and IL-10 respectively, which are key

regulators of inflammation. To test if the induced mRNA expression resulted in enhanced

cytokine production, we performed ELISA of culture supernatants from BMDMs treated with

PNU282987 for 12 h to allow for protein production and secretion. This PNU282987 treatment

did not induce the secretion of any of the cytokines (**Fig. 1C-G**). PNU282987 also did not cause

measurable lactate dehydrogenase (LDH) release (**Fig. 1H**), ruling out cell death as a reason for

the lack of cytokine secretion.

In contrast, PNU282987 had a marked effect on cytokine secretion of LPS-primed

macrophages. After stimulation with LPS for 12 h, secretion of IL-6, TNF-α, and IFN-β1 was

elevated (**Fig. 1C-E**). Subsequent PNU282987 treatment after LPS stimulation strongly inhibited

IL-6 secretion, but not that of TNF-α and IFN-β1 (**Fig. 1C-E**). PNU282987 treatment for 2 h

prior to LPS stimulation again only inhibited IL-6 (**Fig. 1C-E**). These results showed that

PNU282987 inhibits LPS-stimulated IL-6 secretion under both regimens, consistent with prior

reports of anti-inflammatory activity for PNU282987 in LPS-stimulated macrophages (24, 47).

Taken together, the data reveal that PNU282987 elicits an early pro-inflammatory gene

expression profile in macrophages that does not result in enhanced cytokine secretion. In the

context of LPS stimulation, PNU282987 prophylactically or therapeutically reduced LPS-

induced IL-6.

To verify these results in an alternative *in vitro* model, we used RAW264.7 cells, a

murine macrophage-like cell line. Similar to our observations with wild type BMDMs,

PNU282987 treatment for 2 h induced a clear pro-inflammatory gene signature (**Fig. S1A, B and**

Table S3). The induced pro-inflammatory genes included *Ifnb1, Il1b, Il6,* and *Tnf,* as was the

case with BMDMs. Also consistent with our BMDM results, PNU282987 treatment did not drive

secretion of IL-6, TNF-α, IFN-β1, or IL-1β after 12 h (**Fig. S1C-F**). Thus, in RAW264.7 cells as

in BMDMs, PNU282987 treatment induced an early pro-inflammatory gene expression profile

that did not result in secretion of pro-inflammatory cytokines.

In RAW264.7 cells, LPS stimulation elicited IL-6 and TNF-α secretion (**Fig. S1C-F**).

Prior PNU282987 treatment partially inhibited IL-6, TNF-α, and IFN-β1 secretion, and

PNU282987 treatment after LPS stimulation did not significantly change levels of IL-6, TNF-α

and IFN-β1. Over the same time scale, PNU282987 induced the same 240 genes in BMDMs and

RAW264.7 cells (**Fig. S1G**). These overlapping genes were enriched for pro-inflammatory

genes, especially those related to TNF signaling (**Fig. S1H**). This is likely a direct effect, since

PNU282987 did not induce TNF-α secretion. Taken together, these results establish that

PNU282987 induces pro-inflammatory gene expression programs in two distinct *in vitro*

macrophage models.

BMDMs and RAW264.7 cells do not express detectable levels of acetylcholine receptors

The proposed mechanistic target of PNU282987 is the α7 nicotinic acetylcholine receptor

(α7NAChR) (23). However, direct evidence of α7NAChR expression in *in vitro* and *in vivo*

macrophage models is limited. To measure expression of α7NAChR, we performed RT-qPCR

using previously published (47, 48) as well as newly designed oligonucleotide primers, and

measures *Chrna7* gene expression RNA samples from wild type and *Chrna7*[-/-] mice (49). These

approaches showed robust *Chrna7* expression in wild type brain lysate controls (**Fig. 2A**). In

contrast, the same approach did not show *Chrna7* expression in wild type BMDMs and did not

show significant differences with *Chrna7$^{-/-}$* BMDMs (**Fig. 2B**). Similarly, we did not detect

significant expression of nicotinic receptor subunits (α3, α4, α5, α6, α9, α10, β3, and β4) or

muscarinic receptors (M1, M2, M3, M4, M5) in either background (**Fig. 2C**). We detected

variable low expression of β2 nicotinic receptor subunit which showed significant difference

between wild type and *Chrna7$^{-/-}$* BMDMs (**Fig. 2C**). Thus, at the transcript level we did not

obtain convincing evidence of α7NAChR expression in BMDMs.

We used immunoblotting to detect α7NAChR in whole cell lysates using an anti-

α7NAChR antibody that was previously used for the same application (50, 51). We used *Chrna7$^{-/-}$*

$^{/-}$ BMDMs as negative controls, genotyped to ensure their identity (**Fig. S2A**). Immunoblots

showed several bands, but no differences between wild type and *Chrna7$^{-/-}$* (**Fig. S2B**). Thus, we

did not obtain convincing evidence of α7NAChR protein expression.

Fluorescently labelled α-bungarotoxin has been used to detect α7NAChR surface

expression in THP-1 macrophage cells, monocyte derived macrophages and murine intestinal

macrophages (52-54). α-bungarotoxin staining of primary murine peritoneal macrophages and

intestinal macrophage populations isolated from small and large bowel of wild type mice (45)

yielded negative results for α7NAChR expression (**Fig. S3-S6**). Additionally, we were unable to

detect α7NAChR expression by immunofluorescence staining using α7NAChR antibody (not

shown). Thus, by three orthogonal approaches and in multiple macrophage cell types, we were

unable to detect the expression of α7NAChR. Moreover, Immgen (**Fig. S7A**) and the Human

Protein Atlas (**Fig. S8A**) report that both mRNA and protein levels are undetectable in murine

and human macrophages. These data called into question the mechanism of action of

PNU282987.

***Chrna7^{-/-}* BMDMs are defective in the transcriptional response to PNU282987**

Given our lack of evidence of α7NAChR expression, we tested the role of α7NAChR

genetically. Mice deleted for *Chrna7* were generated previously (49). We used RNA-seq to

measure the PNU282987-induced response in *Chrna7^{-/-}* BMDMs. Compared to vehicle control,

PNU282987 treatment induced the expression of just 12 genes in *Chrna7^{-/-}* BMDMs (including

Tnf, Fnip2, Gng4, Rragd) (**Fig. S2C, Table S4**). Thus, the response to PNU282987 was blunted

in *Chrna7^{-/-}* BMDMs.

Compared to wild type BMDMs, PNU282987-treated *Chrna7^{-/-}* BMDMs did not induce

564 genes that were induced in wild type BMDMs. Only 8 genes were induced in both

backgrounds (**Fig. 2D** and **Table S5**). Direct comparison of the transcriptomes of PNU282987-

treated wild type and *Chrna7^{-/-}* BMDMs showed that the vast majority of the innate immune

signaling and pro-inflammatory gene changes were only induced in the wild type (**Fig. 2E, F** and

Table S6). On the other hand, PNU282987-treated *Chrna7^{-/-}* BMDMs differentially expressed

genes related to Ras and Rap1 signaling and the lysosome (**Table S6**). These results suggested

that *Chrna7^{-/-}* BMDMs lack the PNU282987-induced pro-inflammatory transcriptional response.

To better understand why *Chrna7^{-/-}* BMDMs exhibited such a strongly defective response

to PNU282987, we examined gene expression comparing DMSO-treated control BMDMs given

the uncertainty of α7NAChR mediating PNU282987 induced response. Expression of genes

encoding macrophage markers CD11a, CD11c, CD14 and MHC Class II showed significant

differences between wild type and *Chrna7^{-/-}* BMDMs, while other macrophage marker genes

(genes encoding for CD15, CD16, CD64, CD32, F4/80, CD33, CD68, CD80, and CD86) did not

show significant differences (**Fig. S2D, Table S7**). These results ruled out a broad defect in

macrophage *in vitro* differentiation.

Compared to wild type at baseline, *Chrna7$^{-/-}$* BMDMs showed decreased expression of

many immune-related genes, such as *Il6, Il1a, Il1b, Cxcl2*, and *Nox1* (Fig. S2E, F, G and **Table**

S8). These results showed that *Chrna7$^{-/-}$* BMDMs show significant differences in the expression

of a large number of immunity-related genes at baseline, which in itself may cause an altered

response to PNU282987.

PNU282987 activates innate immunity and stress-response transcription factor TFEB

Given the lack of α7NAChR expression on BMDMs and the broad defect in innate

immune gene expression in *Chrna7$^{-/-}$* BMDMs, we considered an alternative hypothesis for the

pro-inflammatory response to PNU282987: namely, the induction of cellular stress. To detect

cellular stress, we examined transcription factor TFEB, which participates in the integrated stress

response, as shown in mouse embryonic fibroblasts and retinal pigment epithelial cells (30).

TFEB resides in the cytosol in an inactive state. Under stressful conditions, including nutrient

deprivation (29, 32), oxidative stress (55, 56), and pathogen infection(35, 38), TFEB translocates

to the nucleus and drives the expression of autophagy, lysosomal, and immune genes. Thus,

nuclear accumulation of TFEB can serve as a sensitive indicator of cell stress.

To determine TFEB translocation, we measured the ratio of nuclear and cytosolic TFEB

staining (N/C ratio) by immunofluorescence microscopy in cells that were treated with

PNU282987 and the commonly used α7NAChR inhibitor, MG624 (57, 58). Resting wild type

BMDMs showed low nuclear staining (**Fig. 3A, G**). Treatment with PNU282987 modestly but

significantly increased TFEB nuclear localization (**Fig. 3B, G**), which was slightly but

significantly inhibited by MG624 pretreatment (**Fig. 3D, G**).

To probe the specificity of this response, we used carbachol, which is a broad-spectrum

cholinergic agonist (59-61). Carbachol treatment more strongly increased TFEB nuclear

localization (**Fig. 3E, H**). Moreover, MG624 pretreatment completely suppressed TFEB

translocation (**Fig. 3F, H**). MG624 treatment, alone or in combination, suppressed TFEB

expression, judging by the overall staining intensity of MG624-treated cells compared to DMSO

controls (**Fig. 3A, C, D, F**)**.** Thus, although MG624 is considered a specific α7NAChR inhibitor,

it also reduces TFEB expression.

To verify the biological significance of TFEB nuclear translocation, we measured

lysosomes by Lysotracker staining (28). Treatment with PNU282987 and carbachol increased

Lysotracker staining compared to vehicle alone **(Fig. 3I-Q)**, which correlated with the TFEB

N/C ratios of each treatment (**Fig. 3G, H**). Moreover, MG624 suppressed Lysotracker staining

(**Fig. 3L, O, Q**), in correlation with the reduced TFEB expression. These data suggest that the

induced nuclear TFEB is active.

Phagocytosis functions upstream of TFEB to drive its nuclear localization (62). To test if

PNU282987 might activate phagocytosis as a mechanism for TFEB activation, we measured

phagocytosis and bactericidal killing of *Escherichia coli* and of *Staphylococcus aureus,* which is

more resistant to bactericide (63, 64). In both infection models, PNU282987 and carbachol pre-

treatment had no effect on either bacterial uptake or bactericidal killing by wild type BMDMs

compared to vehicle controls (**Fig. S9A, B**). Therefore, TFEB activation by PNU282987 is

unlikely a result of increased phagocytic activity and nor did PNU282987 pre-dispose

macrophages to kill bacteria more efficiently.

PNU282987 might be toxic, which could activate TFEB in stressed cells. To measure

toxicity of PNU282987, we treated BMDMs with PNU282987 and measured cell viability. Over

a 128-fold concentration range, PNU282987 or carbachol did not cause significant cell death in

wild type (**Fig. S9C, D**) and *Chrna7*[-/-] BMDMs (**Fig. S9E, F**). These results are consistent with a

previous study using murine peritoneal macrophages (24) and a recent report using RAW264.7

cells (47). These data ruled out toxic stress as a mechanism of TFEB activation by PNU282987.

PNU282987 induces ROS, which are required for TFEB activation

We and others recently showed that oxidative stress and reactive oxygen species can

activate TFEB in macrophages and other cell types (36, 38, 55). Therefore, PNU282987 could

plausibly activate TFEB via ROS. To test if PNU282987 induces ROS production, we measured

ROS levels in PNU282987-treated BMDMs. PNU282987 induced both the accumulation of ROS

overall and H_2O_2 specifically (**Fig. 4A, B**). Surprisingly, MG624 suppressed ROS and H_2O_2

accumulation induced by PNU282987(**Fig. 4A, B**). MG624 alone also depressed H_2O_2 levels

compared to vehicle controls, suggesting that this compound may be redox active (**Fig. 4B**).

These results showed that PNU282987 induces ROS in BMDMs, which could be a mechanism

for TFEB activation.

At baseline, *Chrna7*[-/-] BMDMs showed significantly higher total ROS and H_2O_2

compared to wild type cells (**Fig. 4C, D**). The ROS and H_2O_2 levels in *Chrna7*[-/-] BMDMs were

similar to those in PNU282987-treated wild type BMDMs, and treatment with PNU282987 did

not lead to further increases (**Fig. 4E, F**). Taken together, these results showed that PNU282987

induces ROS and MG624 inhibits them, providing a plausible hypothesis for the activation of

TFEB by PNU282987 through ROS and its inhibition by MG624. To directly test this

hypothesis, we attempted to disrupt TFEB nuclear localization with antioxidants NAC and

NACA. However, both agents induced TFEB nuclear localization on their own, precluding this

test (not shown).

TFEB activation by PNU282987 requires MCOLN1, calcineurin, and protein kinase D.

As an alternative way to test the mechanism of activation of TFEB, we examined the

ROS-MCOLN1-calcineurin pathway. In previous studies, we showed that phagocytosis activates

TFEB through a pathway that involves ROS, Ca^{2+} release, and calcineurin, in that sequence (38).

To examine the role of intracellular Ca^{2+} in PNU282987-induced TFEB translocation, we pre-

treated immortalized BMDMs expressing GFP-TFEB (38) with cell-permeable Ca^{2+}-chelating

agent BAPTA-AM prior to PNU282987 treatment. BAPTA-AM pretreatment suppressed the

PNU282987-induced nuclear localization of GFP-TFEB (**Fig. 5C, E**), suggesting that

intracellular Ca^{2+} is required for PNU282987-induced TFEB nuclear localization. Consistently,

silencing of *Trpml1,* which encodes the main lysosomal Ca^{2+} transporter MCOLN1, also

prevented GFP-TFEB translocation induced by PNU282987 (**Fig. 5K-M**), suggesting that Ca^{2+}

export from the lysosome by MCOLN1 is required.

Similarly, pre-treatment with FK506, which inhibits calcineurin (65), prevented GFP-

TFEB nuclear translocation caused by PNU282987 (**Fig. 5D, E**). This result suggested that

calcineurin is required for PNU282987 to induce TFEB translocation. In support of this

conclusion, silencing of *Ppp3cb* and *Ppp3r1,* respectively encoding catalytic and regulatory

subunits of calcineurin, partially suppressed PNU282987-induced GFP-TFEB translocation (**Fig.**

5F-I, M). These results strongly suggest that calcineurin is required for PNU282987-induced

TFEB activation.

In our previous research, we found that protein kinase D1 is required for TFEB activation

during bacterial infection and phagocytosis (38, 66). To test whether protein kinase D might be

involved in the PNU282987-induced activation of TFEB, we used the inhibitor kb-NB142-70

(67). Pretreatment of BMDMs with kb-NB142-70 suppressed TFEB translocation caused by

PNU282987 (**Fig. S9G-K**), thus implicating protein kinase D in this mechanism of activation as

well. Taken altogether, these results are consistent with PNU282987-induced activation of TFEB

through the ROS-MCOLN1-calcineurin pathway.

TFEB mediates a large fraction of the transcriptional response to PNU282987

To define the extent to which TFEB activation drives the transcriptional response to

PNU282987, we performed RNA-seq of conditionally TFEB-deleted BMDMs (38). The

expression of a set of 23 macrophage differentiation markers was not significantly different

between $Tfeb^{\text{fl/fl}}$ and $Tfeb^{\Delta \text{LysM}}$ BMDMs, showing that *in vitro* macrophage differentiation was

not affected by *Tfeb* deletion (**Fig. S10A**).

At baseline, TFEB deletion decreased the expression of 665 genes compared to $Tfeb^{\text{fl/fl}}$

controls (**Fig. S10B and Table S9**). Surprisingly, this gene set was not enriched for autophagy

nor lysosomal genes, as would have been expected based on data from other cell types (68).

Instead, this TFEB-dependent gene set was enriched in functional categories related to "PI3K-

Akt signaling", "ECM-receptor interaction", "Proteoglycans in cancer", and the extracellular

matrix (**Fig. S10C, Table S10**). TFEB deletion also resulted in higher expression of 138 genes

(**Table S9**), but this gene set did not exhibit significant enrichment of functional categories.

These results suggest that TFEB promotes the expression of extracellular matrix genes in resting

macrophages.

In PNU282987-treated cells compared to wild type, deletion of TFEB resulted in

decreased expression of TNF signaling genes in addition to the baseline defect in the expression

of genes related to extracellular matrix and PI3K-AKT (**Fig. 6A, B**). The genes that were more

highly expressed in TFEB-deleted cells were not significantly enriched for any informative

functional category. Thus, TFEB promotes the expression of TNF signaling genes in response to

PNU282987.

Of 118 genes that were upregulated by PNU282987 in wild type controls, only 66 were

also upregulated in TFEB-deleted BMDMs (**Fig. 6C**). 19 additional genes were induced only in

TFEB-deleted cells. None of these gene sets, *i.e.* TFEB-dependent, -independent, or

compensatory, were enriched for any relevant functional category. Thus, TFEB is required for

about 40% of the transcriptional response that is triggered in wild type BMDMs by PNU282987.

From these analyses, gene *Tnfsf15* stood out as the sole PNU282987-induced gene

encoding a soluble signaling mediator. Tumor necrosis factor superfamily-15 (TNFSF15), also

called TL1A, is a ligand for death receptor 3 (DR3), which activates signaling pathways involved

in the activation, proliferation, apoptosis, and pro-inflammatory responses in immune cells,

including monocytes and macrophages (69). Therefore, it was possible that PNU282987 induced

TNFSF15 production as well as its transcription, leading to an activated macrophage state *in*

vitro. To test that hypothesis, we performed ELISA using supernatants from wild type and

Tfeb[fl/fl] BMDMs. Treatment with PNU282987 did not increase soluble TNFSF15 in these cells

(**Fig. S10D**), suggesting that TNFSF15 is not likely to be a mediator of PNU282987-induced

pro-inflammatory gene expression.

To further test these conclusions, we turned to RAW264.7 cells that were deleted for *Tfeb*

and *Tfe3,* which encodes a TFEB paralog that heterodimerizes with TFEB but can also function

independently as a homodimer (39, 70). In this system, PNU282987-treated *Tfeb Tfe3* double

knockout cells showed defective expression of 550 genes compared to wild type controls (**Fig.**

S11A, Table S11). These genes were enriched in pro-inflammatory pathways, notably TNF and

NOD receptor signaling, as well as lysosomal genes (**Fig. S11B, Table S12**), showing that TFEB

and TFE3 are important for the pro-inflammatory gene expression program that is triggered by

PNU282987. The genes that were more highly expressed in the *Tfeb Tfe3* knockout cells did not

show enrichment of any relevant functional categories. These data strongly support the

conclusion that PNU282987 induces a pro-inflammatory gene expression signature in a TFEB

and TFE3-dependent manner.

Recall that PNU282987 did not induce cytokine secretion in RAW264.7 cells, despite

their transcriptional induction (**Fig. S11C-F**). Surprisingly, deletion of *Tfeb* and *Tfe3* completely

abrogated IL-6 and IFN-β1 production by LPS (but not that of TNF-α, **Fig. S11C-F**). Moreover,

PNU282987 induced a 3-fold increase in LDH release in the *Tfeb Tfe3* knockout cells (**Fig.**

S11G), suggesting that TFEB and TFE3 may mitigate PNU282987 cellular toxicity.

However, at baseline *Tfeb Tfe3* double knockout RAW264.7 cells showed decreased

genes expression of macrophage differentiation markers CD11b, CD15, CD16, CD32, CD64,

CD33, and CD86, as well as TLR2 and TLR4 compared to wild type (**Fig. S11H, Table S13**),

suggesting that the *Tfeb Tfe3* double knockout RAW264.7 cells may show an impaired

macrophage phenotype compared to the wild type control. As with the wild type RAW264.7

cells, the double knockout cells did not detectably express cholinergic receptors, except

α10NAChR, for which the transcript was detected at the same level as in wild type cells (**Fig.**

S11I). *Tfeb*^{fl/fl} and *Tfeb*^{ΔLysM} BMDMs did not show expression of any cholinergic receptor,

similar to wild type BMDMs (**Fig. 2C, S11I**). Taken together, these data show that TFEB (and

TFE3) are important mediators of the PNU282987-induced pro-inflammatory gene transcription

response in macrophages.

DISCUSSION

We describe the early phenotypic effects of PNU282987 treatment on macrophage gene

expression *in vitro*. PNU282987 triggered a large number of pro-inflammatory genes in naïve

primary macrophages and macrophage cell lines. We show evidence that these pro-inflammatory

expression profiles do not result in increased cytokine secretion, at least at the exposure times

that we examined. Because we did not find conclusive evidence of cholinergic receptor

expression in primary macrophages of diverse genetic backgrounds nor in the RAW264.7 cell

line, we hypothesize that PNU282987 may instead exert this activity by cellular stress.

Consistent with this hypothesis, we observe TFEB activation in PNU282987-treated cells, and

deletion of TFEB (and TFE3) significantly impaired PNU282987-induced gene expression

changes. Furthermore, we provide evidence that PNU282987induces the accumulation of ROS in

treated macrophages, and that they trigger TFEB activation through the previously described

ROS-MCOLN1-calcineurin Ca^{2+} signaling pathway. Thus, our findings reveal at least three key

insights.

The first key insight is that secreted pro-inflammatory cytokines (e.g. IFN-β1, IL-6 and

TNF-α) did not increase following PNU282987 treatment, despite their gene induction. It is

possible that longer exposure times (>12 h) might result in induced release. However, we

focused on primary effects of PNU282987 treatment as opposed to the effects of long-term

exposure to avoid confounding secondary responses. Alternatively, the cytokines may be

regulated post-transcriptionally. This interpretation has precedent: transcription of inflammatory

mediators such as TNF-α and IL-6 is uncoupled from secretion by post-transcriptional regulation

mechanisms at the level of mRNA stability or mRNA translation (71). Our RNA-seq analysis

showed induction of *Zfp36l1* in wild type and *Tfeb*[fl/fl] BMDMs by PNU282987. *Zfp36l1* encodes

RNA-binding protein Tristetraprolin (TTP), which belongs to the ZFP36 family of proteins

involved in post-transcriptional regulation of many inflammatory genes (72, 73). Whether

Zfp36l1 contributes to uncoupling TNF-α or IL-6 gene expression from secretion under these

conditions requires further investigation. Consistently, a 5 min pre-treatment with ACh was

sufficient to repress LPS-induced TNF-α, IL-1β, and IL-6 secretion in human monocyte-derived

macrophages, without significant effects on their transcripts (1). Therefore, it is likely that

PNU282987-induced cytokine transcription does not result in enhanced secretion due to post-

transcriptional regulation.

The induction of pro-inflammatory gene expression by PNU282987 appears to run

contrary to the large body of data supporting a cholinergic anti-inflammatory pathway (CAIP).

Studies show that stimulation with ACh or α7NAChR agonists, such as nicotine or choline,

dampen inflammation by suppressing NF-κB in LPS-stimulated human monocyte-derived

macrophages or RAW264.7 macrophages (9, 74). PNU282987 treatment alleviated diverse

inflammatory conditions in mice, such as sepsis-associated lung injury (25), psoriasis-like

inflammation (51), or experimental colitis (27) by inhibiting NF- κB. A unifying theme in these

studies is that PNU282987 was administered to already activated macrophages. Because we were

interested in the primary effects in naïve macrophages, this represents a significant difference

with our approach. Nonetheless, we replicate the inhibitory effect of PNU29287 on LPS-induced

secretion of TNF-α and IL-6. However, the lack of detectable α7NAChR expression in our

systems casts doubt on an α7NAChR-mediated mechanism of action. Published work implicated

JAK2 in the mechanism of cholinergic anti-inflammatory effects through NF-κB repression (7,

75); yet other reports showed that JAK2 can be activated by ROS (76, 77), supporting an

alternative explanation. Moreover, nicotine triggers ROS in murine BMDMs (78) and

RAW264.7 macrophages (79), lending further support to a ROS-mediated mechanism.

The second key insight is the surprising lack of cholinergic receptor expression by

macrophages. Early reports measured the expression of α7NAChR in primary human

macrophages using a nested RT-PCR technique and western blot with α7NAChR antibody

specific to rat and human α7 subunit (3) and in mouse alveolar macrophages using

immunofluorescence (80). However, these studies did not show the specificity of the antibodies

with *Chrna7*[-/-] controls. Another study reported α7NAChR expression in IFN-γ-polarized and IL-

4-polarized BMDMs using RT-PCR (not RT-qPCR), but naïve macrophages were not tested (81).

In yet another study, α7NAChR mRNA expression by RT-PCR was reported in thioglycolate-

elicited peritoneal macrophages that were stimulated with LPS for 24 h (48), or in murine bone

marrow cells, with high variability among littermates (82). Undetectable or highly variable

expression of *Chrna7* transcript among mononuclear phagocytes, as well as levels of *Chrna7*

mRNA not correlating with expression of functional receptors have been noted (83). Large-scale

studies, including our own RNA-seq experiments, do not detect *Chrna7* expression either in

BMDMs or various tissue-resident macrophages, even after pro-inflammatory activation (84). In

our hands, the same oligonucleotide primers that were reported (47, 48, 85) were unable to detect

Chrna7 transcript by highly sensitive RT-qPCR (**Table S14**). Thus, we are unable to conclude

that murine naïve bone marrow-derived macrophages express *Chrna7* transcript.

555   In addition, we used α-bungarotoxin to detect surface expression of α7NAChR by flow

cytometry, similar to previous studies that detected α7NAChR expression on murine intestinal

muscularis macrophages (53) and human primary macrophages (54) but were not able to detect

α7NAChR protein expression in murine peritoneal or muscularis macrophages. The Human

Protein Atlas and ImmGen databases do not support α7NAChR expression in macrophages of

various origins (**Fig. S7, S8**). Thus, there is a lack of unequivocal evidence of α7NAChR

expression in macrophages.

562   It is possible that specific stimuli induce α7nAChR expression. α7NAChR expression by

RT-qPCR was reported to increase in human monocyte-derived macrophages following

treatment with apolipoproteins (86). However, α7NAChR expression was undetectable in

untreated controls, by RT-qPCR, fluorescence staining, and immunoblot. Using the same

antibody, we also failed to detect α7NAChR expression in naïve BMDMs. Another study

reported α7NAChR expression in J774A.1 cells and murine peritoneal macrophages following

LPS treatment but reported no expression in unstimulated macrophages by FITC-labelled

bungarotoxin fluorescence imaging and immunoblot (87). In contrast, a recent study reported no

differences in *Chrna7* mRNA expression of α7nAChR between LPS treated and untreated

RAW264.7 macrophages by RT-qPCR , but the levels of expression were very low (47). We were

unable to detect expression after testing the same primer pairs. Additionally, we found no α-

bungarotoxin staining in murine colonic macrophages from *Salmonella enterica*-infected animals

(**Fig. S6A, B**). Thus, it is unclear whether LPS or other stimuli induce *Chrna7* in murine

macrophages.

Chrna7[-/-] macrophages were severely impaired in their response to PNU282986. These

results are difficult to reconcile with the lack of α7NAChR expression. One possibility is that the

whole-animal deletion might impair the bone marrow compartment and affect *in vitro* properties

(but not differentiation *per se*, see **Fig. S2C**) of BMDMs. To address this possibility, we

attempted a conditional deletion strategy to delete *Chrna7* in monocytes (88) but were

unsuccessful in generating the strain. Alternatively, genetic background may affect the behavior

of the knockout cells. The *Chrna7* knockout was constructed by deleting *Chrna7* exons 8-10 in

the 129/SvEv background and introduced in the C57BL/6 background (49). It is possible that

genomic elements surrounding the transgene or retention of 129S genome affect BMDM

responses to chemical probes such as PNU282987 (89). Whichever the case, we cannot exclude a

contribution of the *Chrna7* locus to the macrophage response to PNU282987 despite its lack of

expression. For instance, our data show that the *Chrna7* knockout cells exhibit a constitutively

higher level of ROS than wild type, which may desensitize them to PNU282987 through

mechanisms that do not hinge on *Chrna7* expression.

Given the lack of conclusive evidence showing cholinergic receptor expression, it is

surprising that MG624 inhibited the nuclear translocation of TFEB and lysosomal biogenesis by

PNU29287. We observed decreased staining of TFEB in MG624-treated cells, suggesting that

MG624 independently causes TFEB repression, which could explain its suppression of

lysosomal biogenesis caused by PNU282987. It is unclear if this mechanism of TFEB repression

is transcriptional or post-transcriptional.

Another key insight is that TFEB was both activated by PNU282987 and biologically

significant for the transcriptional response to it. TFEB was shown to be activated by PNU282987

in murine neuronal N2a cells (90) and in pancreatic acinar cells (91), suggesting that this activity

of PNU282987 may extend beyond macrophages. We find that the mechanism of TFEB

activation in macrophages involves the ROS-responsive MCOLN1-Ca^{2+}-calcineurin pathway

that we described previously (38). Additional upstream regulators of TFEB, such as mTORC1

(92), PKC-α (66) and PKC-β (93), LRRK2 (94), ERK2 (29), p38 MAPK(95), and AKT (96)

could also be involved, but further investigation is required to assess their relevance in the

context of PNU282987-induced TFEB activation.

Tfeb Tfe3 dKO RAW264.7 cells exhibited a stronger defect than *Tfeb*$^{\Delta Lysm}$ BMDMs,

which could be due to the partial redundancy between TFEB and TFE3. We also detected

defective expression of several macrophage markers in the *Tfeb Tfe3* dKO RAW264.7 cells;

nonetheless, these cells remain capable of secreting LPS-induced TNF-α (**Fig. S11D**), and of

phagocytosis and bactericidal killing (38), ruling out a dramatic defect in macrophage

differentiation. However, this may also reflect biological differences between primary cells and

the cell line. Previous work suggested that TFEB was required for human THP-1 cell

differentiation induced by PMA (97); in contrast, we did not detect a defect in BMDM

differentiation in *Tfeb*-deleted murine cells. Furthermore, we previously showed that PMA

activates TFEB *via* protein kinase D1 (66). Thus, PMA may activate a TFEB-dependent

differentiation path in THP-1 cells that is not fully representative of *in vivo* differentiation or M-

CSF-driven *in vitro* differentiation, as in our model system.

The mechanism of PNU282987-induced ROS is currently unknown. Cellular sources of

endogenous ROS include mitochondria, but how PNU282987 affects mitochondrial function in

macrophages is not known. Moreover, the mechanism by which MG624 reduces TFEB

expression is also unknown. TFEB is degraded by ubiquitination and the proteasome in HeLa

cells (98) and neuron-like SH-SY5Y cells or HEK293 cells (99), but whether MG624 affects

TFEB ubiquitination or proteasomal degradation is unknown. Nonetheless, our findings provide

new insight into the mechanisms of action of two widely used chemical agents that are broadly

believed to be highly specific and shed new light on the central role of TFEB in the regulation of

transcriptional phenotypes in macrophages. Based on our results, the interpretation of

experiments that use PNU29287, carbachol, MG624, and potentially other cholinergic agents,

must take into account this alternative mechanism of action.

FIGURE LEGENDS

Figure 1. PNU282987 induces pro-inflammatory genes in BMDMs

A. RNA-seq of BMDMs treated with PNU282987 (50 µM, 2 h) or DMSO. Volcano plot of

differentially expressed genes in PNU282987 treated cells relative to DMSO treated control (fold

change (FC) $\geq$ 1.5; Padj. $\leq$ 0.01). Red indicates upregulated (UP), blue denotes downregulated

(DOWN) genes.

B. Gene ontology (GO) over-representation and KEGG pathway enrichment analysis of

PNU282987 induced genes.

C-G. Measurement of secreted cytokines by ELISA from wild type (WT) BMDMs under the

following conditions: untreated (unt), DMSO, PNU282987 (50 µM, 12 h), LPS (100 ng/mL, 12

h), LPS (100 ng/mL, 2 h) + PNU282987 (50 µM, 12 h), PNU282987 (50 µM, 2 h) + LPS (100

ng/mL, 12 h). Cytokines: IL-6 (**C**), TNF-α (**D**), IFN-β1 (**E**), IL-1β (**F**), and IL-10 (**G**).

H. Lactate dehydrogenase (LDH) release measured in culture supernatants.

n = 3 biological replicates, *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$, One-way

ANOVA followed by Tukey's post-hoc test.

Figure 2. BMDMs from *Chrna7$^{-/-}$* mice are defective in the transcriptional response to PNU282987

A. RT-qPCR, relative expression of *Chrna7*, *Chnrb2,* and *Il6* normalized to *Gapdh,* of mouse whole brain lysate following PBS or LPS (2 mg/kg, 24 h) intraperitoneal injections (3 biological replicates).

B. RT-qPCR of *Chrna7* in WT and *Chrna7$^{-/-}$* BMDMs relative to *Gapdh* (4 biological replicates, ns: not significant ($p \geq 0.05$, 2-sample 2-sided unpaired *t* test).

C. RT-qPCR of transcripts encoding various nicotinic and muscarinic receptor subunit genes in WT and *Chrna7$^{-/-}$* BMDMs (4 biological replicates, Two-way ANOVA, multiple comparisons, ****$p \leq 0.0001$)

D. Venn diagram showing the overlap between PNU282987-induced genes in WT BMDMs compared to PNU282987-induced genes in *Chrna7$^{-/-}$* BMDMs.

E. Volcano plot showing differentially expressed genes in PNU282987-treated *Chrna7$^{-/-}$* BMDMs compared to wild type. Red, upregulated genes in PNU282987 treated *Chrna7$^{-/-}$* BMDMs; blue, upregulated genes in PNU282987 wild type BMDMs (fold change (FC) ≥ 1.5; Padj. ≤ 0.01).

F. GO over-representation and KEGG pathway analysis for genes upregulated in PNU282987-treated WT BMDMs.

Figure 3. PNU282987 activates TFEB in BMDMs

A-F. Representative immunofluorescence micrographs of WT BMDMs stained with anti-TFEB antibody, treated with DMSO control (**A**), PNU282987 (50 µM, 2 h) (**B**), MG624 (1 µM, 2 h) (**C**), MG624 (1 µM, 2 h pre-treatment) + PNU282987 (50 µM, 2 h) (**D**), Carbachol (50 µM, 2 h) (**E**), MG624 (1 µM, 2 h pre-treatment) + Carbachol (50 µM, 2 h) (**F**).

G, H. Nucleus to cytoplasmic (N:C) ratio of TFEB in BMDMs, measured using CellProfiler software (400 cells per condition, **$p \leq 0.01$, ****$p \leq 0.0001$, One-way ANOVA followed by Tukey's post-hoc test).

I-P. LysoTracker staining (red) of GFP-TFEB iBMDMs treated with DMSO control (**I**), PNU282987 (50 µM, 2 h) (**J**), Carbachol (50 µM, 2 h) (**M**) or MG624 (1 µM, 2 h) (**K**) MG624 (1 µM, 2 h pre-treatment) + PNU282987 (50 µM, 2 h) (**L**), MG624 (1 µM, 2 h pre-treatment) + Carbachol (50 µM, 2 h) (**O**) with LPS (100 ng/mL, 2 h) (**N**) and Torin-1 (1.25 uM, 1 h) (**P**).

(**Q**) LysoTracker mean fluorescence intensity per cell (MFI/cell) measured with CellProfiler. Images and quantification of 3 biological replicates (400 cells per condition, ****$p \leq 0.0001$, One-way ANOVA followed by Tukey's post-hoc test).

Figure 4. PNU282987 induces ROS production

A, B. Total ROS (**A**) and (**B**) superoxide in WT BMDMs treated with DMSO, PNU282987 (50 µM, 2 h), or following pre-treatment with MG624 (1 µM, 2 h). Pyocyanin (200 µM, 30 min) and NAC (5 mM, 30 min), positive and negative control, respectively. (400 cells per condition, *$p \leq 0.05$, ****$p \leq 0.0001$, One-way ANOVA followed by Tukey's post-hoc test).

C, D. Total ROS (**C**) and superoxide (**D**) shown as mean fluorescence intensity in WT and *Chrna7*[-/-] BMDMs (200 cells per condition, ***$p \leq 0.0001$, 2-sided 2-sample unpaired *t* test).

E, F. Total ROS (**E**) and superoxide (**F**) measured in WT and *Chrna7*[-/-] BMDMs following treatment with PNU282987 (50 µM, 2 h), NAC (5 mM, 30 min) and Pyocyanin (200 µM, 30 min). 400 cells per condition, ****p ≤ 0.0001, One-way ANOVA followed by Tukey's post-hoc test.

Figure 5. TFEB activation by PNU282987 requires MCOLN1 and calcineurin

A-D. Representative fluorescence micrographs of GFP-TFEB iBMDMs treated with DMSO (**A**), PNU282987 (50 µM, 2 h) (**B**), and PNU282987 (50 µM, 2 h) following pre-treatment with BAPTA-AM (10 µM, 3 h) (**C**) or FK506 (5 µM, 6 h) (**D**).

E. Quantification of TFEB N:C ratio using CellProfiler (3 biological replicates, 350 cells per condition, ****p ≤ 0.0001, One-way ANOVA followed by Tukey's post-hoc test).

F-L. Representative fluorescence micrographs of GFP-TFEB iBMDMs treated with siRNA, scrambled (*Scr*) (**F, G**) or directed against *Ppp3cb* (**H**), *Ppp3r1* (**I**), and *Trpml1* (**K, L**) for 48 h prior to treatment with PNU282987 (50 µM, 2 h) or serum starvation (**J, L**).

M. Quantification of GFP-TFEB N:C ratio by CellProfiler (3 biological replicates, n=300 cells) **** p ≤ 0.0001 (one-way ANOVA followed by Tukey's post-hoc test).

Figure 6. TFEB mediates 40% of the transcriptional response to PNU282987 in BMDMs

A. Volcano plot showing differentially expressed genes in PNU282987-induced *Tfeb*[ΔLysM] BMDMs compared to PNU282987-induced *Tfeb*[fl/fl] BMDMs. Red, upregulated genes in PNU282987-treated *Tfeb*[ΔLysM] BMDMs and blue, upregulated genes in PNU282987-treated *Tfeb*[fl/fl] BMDMs (fold change (FC) ≥ 1.5; Padj. ≤ 0.01).

B. Gene ontology (GO) over-representation and (KEGG) pathway enrichment analysis of PNU282987 induced genes in *Tfeb*[fl/fl] BMDMs.

C. Venn diagram showing the overlap between PNU282987-induced genes in *Tfeb*^{fl/fl} BMDMs

and PNU282987-induced genes in *Tfeb*^{ΔLysM} BMDMs.

REFERENCES

1.   Borovikova LV, Ivanova S, Zhang M, Yang H, Botchkina GI, Watkins LR, et al. Vagus

nerve stimulation attenuates the systemic inflammatory response to endotoxin. Nature.

2000;405(6785):458-62.

2.   Kelly MJ, Breathnach C, Tracey KJ, Donnelly SC. Manipulation of the inflammatory

reflex as a therapeutic strategy. Cell Reports Medicine. 2022;3(7).

3.   Wang H, Yu M, Ochani M, Amella CA, Tanovic M, Susarla S, et al. Nicotinic

acetylcholine receptor α7 subunit is an essential regulator of inflammation. Nature.

2003;421(6921):384-8.

4.   Rosas-Ballina M, Olofsson PS, Ochani M, Valdés-Ferrer SI, Levine YA, Reardon C, et al.

Acetylcholine-synthesizing T cells relay neural signals in a vagus nerve circuit. Science.

2011;334(6052):98-101.

5.   Baez-Pagan CA, Delgado-Velez M, Lasalde-Dominicci JA. Activation of the Macrophage

alpha7 Nicotinic Acetylcholine Receptor and Control of Inflammation. J Neuroimmune

Pharmacol. 2015;10(3):468-76.

6.   Yoshikawa H, Kurokawa M, Ozaki N, Nara K, Atou K, Takada E, et al. Nicotine inhibits

the production of proinflammatory mediators in human monocytes by suppression of I-κB

phosphorylation and nuclear factor-κB transcriptional activity through nicotinic acetylcholine

receptor α7. Clinical & Experimental Immunology. 2006;146(1):116-23.

7.   de Jonge WJ, van der Zanden EP, The FO, Bijlsma MF, van Westerloo DJ, Bennink RJ, et

al. Stimulation of the vagus nerve attenuates macrophage activation by activating the Jak2-

STAT3 signaling pathway. Nat Immunol. 2005;6(8):844-51.

8.   Van Westerloo DJ, Giebelen IA, Florquin S, Daalhuisen J, Bruno MJ, de Vos AF, et al.

The cholinergic anti-inflammatory pathway regulates the host response during septic peritonitis.

Journal of Infectious Diseases. 2005;191(12):2138-48.

9.   Wang H, Liao H, Ochani M, Justiniani M, Lin X, Yang L, et al. Cholinergic agonists

inhibit HMGB1 release and improve survival in experimental sepsis. Nature medicine.

2004;10(11):1216-21.

10.   Ghia JE, Blennerhassett P, Kumar–Ondiveeran H, Verdu EF, Collins SM. The vagus

nerve: a tonic inhibitory influence associated with inflammatory bowel disease in a murine

model. Gastroenterology. 2006;131(4):1122-30.

11.   Bai A, Guo Y, Lu N. The effect of the cholinergic anti-inflammatory pathway on

experimental colitis. Scandinavian journal of immunology. 2007;66(5):538-45.

12.   Westman M, Engström M, Catrina A, Lampa J. Cell specific synovial expression of

nicotinic alpha 7 acetylcholine receptor in rheumatoid arthritis and psoriatic arthritis.

Scandinavian journal of immunology. 2009;70(2):136-40.

13.   van Maanen MA, Lebre MC, van der Poll T, LaRosa GJ, Elbaum D, Vervoordeldonk MJ,

et al. Stimulation of nicotinic acetylcholine receptors attenuates collagen-induced arthritis in

mice. Arthritis & Rheumatism. 2009;60(1):114-22.

14.   Gwilt CR, Donnelly LE, Rogers DF. The non-neuronal cholinergic system in the airways:

an unappreciated regulatory role in pulmonary inflammation? Pharmacology & therapeutics.

2007;115(2):208-22.

15. Su X, Matthay MA, Malik AB. Requisite role of the cholinergic α7 nicotinic

acetylcholine receptor pathway in suppressing gram-negative sepsis-induced acute lung

inflammatory injury. The journal of immunology. 2010;184(1):401-10.

16. Greene CM, Ramsay H, Wells RJ, O'Neill SJ, McElvaney NG. Inhibition of Toll-Like

Receptor 2-Mediated Interleukin-8 Production in Cystic Fibrosis Airway Epithelial Cells via the

α7-Nicotinic Acetylcholine Receptor. Mediators of Inflammation. 2010;2010.

17. Marrero MB, Lucas R, Salet C, Hauser TA, Mazurov A, Lippiello PM, et al. An α7

nicotinic acetylcholine receptor-selective agonist reduces weight gain and metabolic changes in a

mouse model of diabetes. Journal of Pharmacology and Experimental Therapeutics.

2010;332(1):173-80.

18. Pavlov VA, Tracey KJ. The vagus nerve and the inflammatory reflex—linking immunity

and metabolism. Nature Reviews Endocrinology. 2012;8(12):743-54.

19. Bencherif M, Lippiello PM, Lucas R, Marrero MB. Alpha7 nicotinic receptors as novel

therapeutic targets for inflammation-based diseases. Cell Mol Life Sci. 2011;68(6):931-49.

20. Youssef ME, Moustafa Y, Abdelrazek H. Molecular mechanisms of α7-nAchR-mediated

anti-inflammatory effects. Indian Journal of Physiology and Pharmacology. 2021;64(3):158-73.

21. Lu B, Kwan K, Levine YA, Olofsson PS, Yang H, Li J, et al. α7 nicotinic acetylcholine

receptor signaling inhibits inflammasome activation by preventing mitochondrial DNA release.

Molecular medicine. 2014;20:350-8.

22. Bodnar AL, Cortes-Burgos LA, Cook KK, Dinh DM, Groppi VE, Hajos M, et al.

Discovery and structure− activity relationship of quinuclidine benzamides as agonists of α7

nicotinic acetylcholine receptors. Journal of medicinal chemistry. 2005;48(4):905-8.

23.   Hajos M, Hurst R, Hoffmann W, Krause M, Wall T, Higdon N, et al. The selective α7

nicotinic acetylcholine receptor agonist PNU-282987 [N-[(3R)-1-azabicyclo [2.2. 2] oct-3-yl]-4-

chlorobenzamide hydrochloride] enhances GABAergic synaptic activity in brain slices and

restores auditory gating deficits in anesthetized rats. Journal of Pharmacology and Experimental

Therapeutics. 2005;312(3):1213-22.

24.   Shao Z, Li Q, Wang S, Chen Z. Protective effects of PNU-282987 on sepsis-induced

acute lung injury in mice. Molecular Medicine Reports. 2019;19(5):3791-8.

25.   Pinheiro NM, Santana FPR, Almeida RR, Guerreiro M, Martins MA, Caperuto LC, et al.

Acute lung injury is reduced by the α7nAChR agonist PNU-282987 through changes in the

macrophage profile. The FASEB Journal. 2017;31(1):320-32.

26.   Zhao X, Yu Z, Lv Z, Meng L, Xu J, Yuan S, et al. Activation of alpha-7 nicotinic

acetylcholine receptors (α7nAchR) promotes the protective autophagy in LPS-induced acute

lung injury (ALI) in vitro and in vivo. Inflammation. 2019;42:2236-45.

27.   Pu W, Su Z, Wazir J, Zhao C, Wei L, Wang R, et al. Protective effect of α7 nicotinic

acetylcholine receptor activation on experimental colitis and its mechanism. Molecular

Medicine. 2022;28(1):104.

28.   Sardiello M, Palmieri M, Di Ronza A, Medina DL, Valenza M, Gennarino VA, et al. A

gene network regulating lysosomal biogenesis and function. Science. 2009;325(5939):473-7.

29.   Settembre C, Di Malta C, Polito VA, Arencibia MG, Vetrini F, Erdin S, et al. TFEB links

autophagy to lysosomal biogenesis. science. 2011;332(6036):1429-33.

30.   Martina JA, Diab HI, Brady OA, Puertollano R. TFEB and TFE 3 are novel components

of the integrated stress response. The EMBO journal. 2016;35(5):479-95.

31.   Nabar NR, Kehrl JH. Focus: Infectious Diseases: The Transcription Factor EB Links

Cellular Stress to the Immune Response. The Yale Journal of Biology and Medicine.

2017;90(2):301.

32.   Roczniak-Ferguson A, Petit CS, Froehlich F, Qian S, Ky J, Angarola B, et al. The

transcription factor TFEB links mTORC1 signaling to transcriptional control of lysosome

homeostasis. Science signaling. 2012;5(228):ra42-ra.

33.   Settembre C, Zoncu R, Medina DL, Vetrini F, Erdin S, Erdin S, et al. A lysosome-to-

nucleus signalling mechanism senses and regulates the lysosome via mTOR and TFEB. The

EMBO journal. 2012;31(5):1095-108.

34.   Nezich CL, Wang C, Fogel AI, Youle RJ. MiT/TFE transcription factors are activated

during mitophagy downstream of Parkin and Atg5. Journal of Cell Biology. 2015;210(3):435-50.

35.   Visvikis O, Ihuegbu N, Labed SA, Luhachack LG, Alves A-MF, Wollenberg AC, et al.

Innate host defense requires TFEB-mediated transcription of cytoprotective and antimicrobial

genes. Immunity. 2014;40(6):896-909.

36.   Zhang X, Cheng X, Yu L, Yang J, Calvo R, Patnaik S, et al. MCOLN1 is a ROS sensor in

lysosomes that regulates autophagy. Nature communications. 2016;7(1):12109.

37.   Tan A, Prasad R, Lee C, Jho E-h. Past, present, and future perspectives of transcription

factor EB (TFEB): mechanisms of regulation and association with disease. Cell Death &

Differentiation. 2022;29(8):1433-49.

38.   Najibi M, Honwad HH, Moreau JA, Becker SM, Irazoqui JE. A novel Nox/Phox-Cd38-

Naadp-Tfeb axis important for macrophage activation during bacterial phagocytosis. Autophagy.

2022;18(1):124-41.

39.   Pastore N, Brady OA, Diab HI, Martina JA, Sun L, Huynh T, et al. TFEB and TFE3

cooperate in the regulation of the innate immune response in activated macrophages. Autophagy.

2016;12(8):1240-58.

40.   McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, et al.

CellProfiler 3.0: Next-generation image processing for biology. PLoS biology.

2018;16(7):e2005970.

41.   Yukselen O, Turkyilmaz O, Ozturk AR, Garber M, Kucukural A. DolphinNext: a

distributed data processing platform for high throughput genomics. BMC genomics. 2020;21:1-

16.

42.   Reimand J, Arak T, Adler P, Kolberg L, Reisberg S, Peterson H, et al. g: Profiler—a web

server for functional interpretation of gene lists (2016 update). Nucleic acids research.

2016;44(W1):W83-W9.

43.   Evangelista JE, Xie Z, Marino GB, Nguyen N, Clarke Daniel JB, Ma'ayan A. Enrichr-

KG: bridging enrichment analysis across multiple libraries. Nucleic Acids Research.

2023;51(W1):W168-W79.

44.   Hulsen T. DeepVenn--a web application for the creation of area-proportional Venn

diagrams using the deep learning framework Tensorflow. js. arXiv preprint arXiv:221004597.

2022.

45.   Koscsó B, Gowda K, Schell TD, Bogunovic M. Purification of dendritic cell and

macrophage subsets from the normal mouse small intestine. Journal of immunological methods.

2015;421:1-13.

46.   Koscsó B, Bogunovic M. Analysis and purification of mouse intestinal dendritic cell and

macrophage subsets by flow cytometry. Current Protocols in Immunology. 2016;114(1):14.39. 1-

14.39. 14.

47.   Tan Y, Chu Z, Shan H, Zhangsun D, Zhu X, Luo S. Inflammation regulation via an

agonist and antagonists of α7 nicotinic acetylcholine receptors in RAW264. 7 macrophages.

Marine Drugs. 2022;20(3):200.

48.   Kawashima K, Yoshikawa K, Fujii YX, Moriwaki Y, Misawa H. Expression and function

of genes encoding cholinergic components in murine immune cells. Life sciences. 2007;80(24-

25):2314-9.

49.   Orr-Urtreger A, Göldner FM, Saeki M, Lorenzo I, Goldberg L, De Biasi M, et al. Mice

deficient in the α7 neuronal nicotinic acetylcholine receptor lack α-bungarotoxin binding sites

and hippocampal fast nicotinic currents. Journal of Neuroscience. 1997;17(23):9165-71.

50.   Lei W, Zhao C, Sun J, Jin Y, Duan Z. Activation of α7nAChR preserves intestinal barrier

integrity by enhancing the HO-1/STAT3 signaling to inhibit NF-κB activation in mice.

Biomedicine & Pharmacotherapy. 2022;149:112733.

51.   Chen Y, Lian P, Peng Z, Wazir J, Ma C, Wei L, et al. Alpha-7 nicotinic acetylcholine

receptor agonist alleviates psoriasis-like inflammation through inhibition of the STAT3 and NF-

κB signaling pathway. Cell Death Discovery. 2022;8(1):141.

52.   Nemethova A, Michel K, Gomez-Pinilla PJ, Boeckxstaens GE, Schemann M. Nicotine

attenuates activation of tissue resident macrophages in the mouse stomach through the β2

nicotinic acetylcholine receptor. PLoS One. 2013;8(11):e79264.

53.   Matteoli G, Gomez-Pinilla PJ, Nemethova A, Di Giovangiulio M, Cailotto C, van Bree

SH, et al. A distinct vagal anti-inflammatory pathway modulates intestinal muscularis resident

macrophages independent of the spleen. Gut. 2014;63(6):938-48.

54.   Siniavin AE, Streltsova MA, Kudryavtsev DS, Shelukhina IV, Utkin YN, Tsetlin VI.

Activation of α7 nicotinic acetylcholine receptor upregulates HLA-DR and macrophage

receptors: Potential role in adaptive immunity and in preventing immunosuppression.

Biomolecules. 2020;10(4):507.

55.   Martina JA, Puertollano R. Protein phosphatase 2A stimulates activation of TFEB and

TFE3 transcription factors in response to oxidative stress. Journal of Biological Chemistry.

2018;293(32):12525-34.

56.   Martina JA, Guerrero-Gómez D, Gómez-Orte E, Antonio Barcena J, Cabello J, Miranda-

Vizuete A, et al. A conserved cysteine-based redox mechanism sustains TFEB/HLH-30 activity

under persistent stress. The EMBO Journal. 2021;40(3):e105793.

57.   Gotti C, Balestra B, Moretti M, Rovati G, Maggi L, Rossoni G, et al. 4-Oxystilbene

compounds are selective ligands for neuronal nicotinic αBungarotoxin receptors. British journal

of pharmacology. 1998;124(6):1197-206.

58.   Tanibuchi Y, Wu J, Toyohara J, Fujita Y, Iyo M, Hashimoto K. Characterization of [3H]

CHIBA-1001 binding to α7 nicotinic acetylcholine receptors in the brain from rat, monkey, and

human. Brain research. 2010;1348:200-8.

59.   Li Y-z, Liu X-h, Rong F, Hu S, Sheng Z-y. Carbachol inhibits TNF-α-induced endothelial

barrier dysfunction through alpha 7 nicotinic receptors. Acta Pharmacologica Sinica.

2010;31(10):1389-94.

60.   Zhou G, Hu S, Lv Y, Song Q, Zou X, Sheng Z. Carbachol alleviates rat cytokine release

and organ dysfunction induced by lipopolysaccharide. Journal of Trauma and Acute Care

Surgery. 2011;71(1):157-62.

61.   Halder N, Lal G. Cholinergic system and its therapeutic importance in inflammation and

autoimmunity. Frontiers in Immunology. 2021;12:660342.

62.   Gray MA, Choy CH, Dayam RM, Ospina-Escobar E, Somerville A, Xiao X, et al.

Phagocytosis enhances lysosomal and bactericidal properties by activating the transcription

factor TFEB. Current Biology. 2016;26(15):1955-64.

63.   Bantawa K, Sah SN, Subba Limbu D, Subba P, Ghimire A. Antibiotic resistance patterns

of Staphylococcus aureus, Escherichia coli, Salmonella, Shigella and Vibrio isolated from

chicken, pork, buffalo and goat meat in eastern Nepal. BMC research notes. 2019;12(1):1-6.

64.   van den Honert MS, Gouws PA, Hoffman LC. A preliminary study: Antibiotic resistance

of Escherichia coli and Staphylococcus aureus from the meat and feces of various South African

wildlife species. Food Science of Animal Resources. 2021;41(1):135.

65.   Liu J, Farmer Jr JD, Lane WS, Friedman J, Weissman I, Schreiber SL. Calcineurin is a

common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. Cell.

1991;66(4):807-15.

66.   Najibi M, Labed SA, Visvikis O, Irazoqui JE. An evolutionarily conserved PLC-PKD-

TFEB pathway for host defense. Cell reports. 2016;15(8):1728-42.

67.   LaValle CR, Bravo-Altamirano K, Giridhar KV, Chen J, Sharlow E, Lazo JS, et al. Novel

protein kinase D inhibitors cause potent arrest in prostate cancer cell growth and motility. BMC

chemical biology. 2010;10:1-16.

68.   Carey K, Paulus G, Wang L, Balce D, Luo J, Bergman P, et al. TFEB transcriptional

responses reveal negative feedback by BHLHE40 and BHLHE41. Cell Rep 33: 108371. 2020.

69.   Xu W-D, Li R, Huang A-F. Role of TL1A in inflammatory autoimmune diseases: a

comprehensive review. Frontiers in Immunology. 2022;13:891328.

70.   Martina JA, Diab HI, Lishu L, Jeong-A L, Patange S, Raben N, et al. The nutrient-

responsive transcription factor TFE3 promotes autophagy, lysosomal biogenesis, and clearance

of cellular debris. Science signaling. 2014;7(309):ra9-ra.

71.   Stumpo DJ, Lai WS, Blackshear PJ. Inflammation: cytokines and RNA-based regulation.

Wiley Interdisciplinary Reviews: RNA. 2010;1(1):60-80.

72.   Makita S, Takatori H, Nakajima H. Post-transcriptional regulation of immune responses

and inflammatory diseases by RNA-binding ZFP36 family proteins. Frontiers in immunology.

2021;12:711633.

73.   Jalonen U, Nieminen R, Vuolteenaho K, Kankaanranta H, Moilanen E, editors. Down-

regulation of tristetraprolin expression results in enhanced IL-12 and MIP-2 production and

reduced MIP-3alpha synthesis in activated macrophages. SCANDINAVIAN JOURNAL OF

IMMUNOLOGY; 2007: BLACKWELL PUBLISHING 9600 GARSINGTON RD, OXFORD

OX4 2DQ, OXON, ENGLAND.

74.   Parrish WR, Rosas-Ballina M, Gallowitsch-Puerta M, Ochani M, Ochani K, Yang L-H, et

al. Modulation of TNF release by choline requires α7 subunit nicotinic acetylcholine receptor-

mediated signaling. Molecular medicine. 2008;14(9):567-74.

75.   Peña G, Cai B, Liu J, van der Zanden EP, Deitch EA, de Jonge WJ, et al.

Unphosphorylated STAT3 modulates alpha7 nicotinic receptor signaling and cytokine production

in sepsis. European journal of immunology. 2010;40(9):2580-9.

76.   Simon AR, Rai U, Fanburg BL, Cochran BH. Activation of the JAK-STAT pathway by

reactive oxygen species. American Journal of Physiology-Cell Physiology. 1998;275(6):C1640-

C52.

77.   Liu J, Han X, Zhang T, Tian K, Li Z, Luo F. Reactive oxygen species (ROS) scavenging

biomaterials for anti-inflammatory diseases: from mechanism to therapy. Journal of Hematology

& Oncology. 2023;16(1):116.

78.   Mao C, Li D, Zhou E, Zhang J, Wang C, Xue C. Nicotine exacerbates atherosclerosis

through a macrophage-mediated endothelial injury pathway. Aging (Albany NY).

2021;13(5):7627-43.

79.   Tsoyi K, Jang HJ, Kim JW, Chang HK, Lee YS, Pae H-O, et al. Stimulation of alpha7

nicotinic acetylcholine receptor by nicotine attenuates inflammatory response in macrophages

and improves survival in experimental model of sepsis through heme oxygenase-1 induction.

Antioxidants & redox signaling. 2011;14(11):2057-70.

80.   Su X, Lee JW, Matthay ZA, Mednick G, Uchida T, Fang X, et al. Activation of the $\alpha 7$

nAChR reduces acid-induced acute lung injury in mice and rats. American journal of respiratory

cell and molecular biology. 2007;37(2):186-92.

81.   Lee RH, Vazquez G. Evidence for a prosurvival role of alpha-7 nicotinic acetylcholine

receptor in alternatively (M2)-activated macrophages. Physiol Rep. 2013;1(7):e00189.

82.   St-Pierre S, Jiang W, Roy P, Champigny C, LeBlanc É, Morley BJ, et al. Nicotinic

acetylcholine receptors modulate bone marrow-derived pro-inflammatory monocyte production

and survival. PLoS One. 2016;11(2):e0150230.

83.   Richter K, Grau V. Signaling of nicotinic acetylcholine receptors in mononuclear

phagocytes. Pharmacol Res. 2023;191:106727.

84.   Lavin Y, Winter D, Blecher-Gonen R, David E, Keren-Shaul H, Merad M, et al. Tissue-

resident macrophage enhancer landscapes are shaped by the local microenvironment. Cell.

2014;159(6):1312-26.

85.   Fujii T, Mashimo M, Moriwaki Y, Misawa H, Ono S, Horiguchi K, et al. Expression and

Function of the Cholinergic System in Immune Cells. Front Immunol. 2017;8:1085.

86.   Lin Y-K, Yeh C-T, Kuo K-T, Fong I-H, Yadav VK, Kounis NG, et al. Apolipoprotein

(a)/Lipoprotein (a)-Induced oxidative-inflammatory α7-nAChR/p38 MAPK/IL-6/RhoA-GTP

signaling Axis and M1 macrophage polarization modulate inflammation-associated development

of coronary artery spasm. Oxidative medicine and cellular longevity. 2022;2022.

87.   Khan MA, Farkhondeh M, Crombie J, Jacobson L, Kaneki M, Martyn JA.

Lipopolysaccharide upregulates alpha7 acetylcholine receptors: stimulation with GTS-21

mitigates growth arrest of macrophages and improves survival in burned mice. Shock.

2012;38(2):213-9.

88.   Hernandez CM, Cortez I, Gu Z, Colón-Sáez JO, Lamb PW, Wakamiya M, et al. Research

tool: validation of floxed α7 nicotinic acetylcholine receptor conditional knockout mice using in

vitro and in vivo approaches. The Journal of physiology. 2014;592(15):3201-14.

89.   Weiss FD, Alvarez Y, Sahu A, Shakeri F, Lee HE, Gellner A-K, et al. Retention of ES

cell-derived 129S genome drives NLRP1 hypersensitivity and transcriptional deregulation in

Nlrp3-/-mice. bioRxiv. 2024:2024.01. 03.573991.

90.   Ito T, Inden M, Ueda T, Asaka Y, Kurita H, Hozumi I. The neuroprotective effects of

activated alpha7 nicotinic acetylcholine receptor against mutant copper-zinc superoxide

dismutase 1-mediated toxicity. Sci Rep. 2020;10(1):22157.

91.   Li B, Wu J, Bao J, Han X, Shen S, Ye X, et al. Activation of alpha7nACh receptor

protects against acute pancreatitis through enhancing TFEB-regulated autophagy. Biochim

Biophys Acta Mol Basis Dis. 2020;1866(12):165971.

92.   Martina JA, Chen Y, Gucek M, Puertollano R. MTORC1 functions as a transcriptional

regulator of autophagy by preventing nuclear transport of TFEB. Autophagy. 2012;8(6):903-14.

93.   Ferron M, Settembre C, Shimazu J, Lacombe J, Kato S, Rawlings DJ, et al. A RANKL–

PKCβ–TFEB signaling cascade is necessary for lysosomal biogenesis in osteoclasts. Genes &

development. 2013;27(8):955-69.

94.   Nabar NR, Heijjer CN, Shi C-S, Hwang I-Y, Ganesan S, Karlsson MC, et al. LRRK2 is

required for CD38-mediated NAADP-Ca2+ signaling and the downstream activation of TFEB

(transcription factor EB) in immune cells. Autophagy. 2022;18(1):204-22.

95.   Hsu CL, Lee EX, Gordon KL, Paz EA, Shen W-C, Ohnishi K, et al. MAP4K3 mediates

amino acid-dependent regulation of autophagy via phosphorylation of TFEB. Nature

Communications. 2018;9(1):942.

96.   Palmieri M, Pal R, Nelvagal HR, Lotfi P, Stinnett GR, Seymour ML, et al. mTORC1-

independent TFEB activation via Akt inhibition promotes cellular clearance in neurodegenerative

storage diseases. Nature communications. 2017;8(1):14338.

97.   Martina JA, Jeong E, Puertollano R. p38 MAPK-dependent phosphorylation of TFEB

promotes monocyte-to-macrophage differentiation. EMBO reports. 2023;24(2):e55472.

98.   Sha Y, Rao L, Settembre C, Ballabio A, Eissa NT. STUB 1 regulates TFEB-induced

autophagy–lysosome pathway. The EMBO journal. 2017;36(17):2544-52.

99.   Li C, Wang X, Li X, Qiu K, Jiao F, Liu Y, et al. Proteasome inhibition activates

autophagy-lysosome pathway associated with TFEB dephosphorylation and nuclear

translocation. Frontiers in cell and developmental biology. 2019;7:170.

A WT BMDMs (PNU282987 vs DMSO)

B UP in WT BMDMs (PNU282987 vs DMSO)

C IL-6
D TNF-α
E IFN-β1
F IL-1β
G IL-10

H LDH

A PNU282987 treatment (*Tfeb*^fl/fl vs *Tfeb*^ΔLysM BMDMs)

B UP in *Tfeb*^fl/fl

C