Transient destabilization of whole brain dynamics induced by DMT

Juan Ignacio Piccinini[1*], Yonatan Sanz Perl[1,2], Carla Pallavicini[1, 3], Gustavo Deco[2,4,5,6,] Morten Kringelbach[7,8,9], David Nutt[10], Robin Carhart-Harris[10, 11], Christopher Timmermann[10], Enzo Tagliazucchi[1,12*]

[1]Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, and CONICET - Universidad de Buenos Aires, Instituto de Física Aplicada e Interdisciplinaria (INFINA). Buenos Aires, Argentina.

[2]Center for Brain and Cognition, Computational Neuroscience Group, Department of Information and Communication Technologies, Universitat Pompeu Fabra, Barcelona, Spain

[3]Université Paris Cité, CNRS, Integrative Neuroscience and Cognition Center, Paris, France.

[4]Institució Catalana de la Recerca i Estudis Avançats (ICREA), Barcelona, Spain

[5]Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[6]School of Psychological Sciences, Monash University, Melbourne, Clayton VIC, Australia

[7]Centre for Eudaimonia and Human Flourishing, University of Oxford, Oxford, UK

[8]Department of Psychiatry, University of Oxford, Oxford, UK

[9]Center for Music in the Brain, Department of Clinical Medicine, Aarhus University, Denmark

[10]Centre for Psychedelic Research, Division of Psychiatry, Department of Brain Sciences, Imperial College London, London, UK

[11]Departments of Neurology and Psychiatry, University of California San Francisco, San Francisco, California, USA

[12]Latin American Brain Health Institute (BrainLat), Universidad Adolfo Ibañez, Santiago, Chile

* Corresponding authors: Juan Piccinini (piccijuan@gmail.com), Enzo Tagliazucchi (tagliazucchi.enzo@googlemail.com)

Abstract

The transition towards the brain state induced by psychedelic drugs is frequently neglected in favor

of a static description of their acute effects. We used a time-dependent whole-brain model to

reproduce large-scale brain dynamics measured with fMRI from 15 volunteers under 20 mg

intravenous N,N-Dimethyltryptamine (DMT), a short-acting psychedelic. To capture its transient

effects, we parametrized the proximity to a global bifurcation using a pharmacokinetic equation.

Simulated perturbations revealed a transient of heightened reactivity concentrated in fronto-parietal

regions and visual cortices, correlated with serotonin 5HT2a receptor density, the primary target of

psychedelics. These advances suggest a mechanism to explain key features of the psychedelic state

and also predicts that the temporal evolution of these features aligns with pharmacokinetics. Our

results contribute to understanding how psychedelics introduce a transient where minimal

perturbations can achieve a maximal effect, shedding light on how short psychedelic episodes may

extend an overarching influence over time.

14
15

Introduction

Psychedelic drugs offer an opportunity to investigate how changes in the brain across multiple

spatiotemporal scales interact with human consciousness and cognition [1]. At the molecular level,

psychedelics bind to the serotonin 5HT2a receptor [2], recruiting specific intracellular signaling

pathways that are different from those implicated in the action of non-psychedelic 5HT2A agonists

[3,4]. The subjective effects of psychedelics may also depend on other pharmacological and non-

pharmacological factors [5], the latter including the context of drug intake and the mindset of the user

[6]. At the systems level, psychedelics increase global network integration measured with functional

magnetic resonance imaging (fMRI) [7-12], and evidence from multiple modalities links their effects to

increased entropy and complexity of spontaneous brain activity fluctuations [13-19]. An integrative

understanding of psychedelic action would require the identification of causal mechanisms behind these empirical observations, from molecules to subjective experience [1].

In recent years, generative whole-brain activity models have been increasingly adopted to test potential mechanisms underlying neuroimaging data, including successful applications to the specific case of psychedelic compounds psilocybin and lysergic acid diethylamide (LSD) [20]. In these two case studies, biophysical models consisting of local excitatory and inhibitory populations with excitatory long-range connections were used to provide evidence supporting psychedelic-induced modulation of 5HT2a synaptic scaling [21-23]. Herzog and colleagues implemented a similar model to show that 5HT2a receptor stimulation is consistent with increased brain-wide entropy [13], in agreement with the theoretical model of psychedelic action proposed by Carhart-Harris [24]. A complementary approach is to consider phenomenological models to investigate changes in global brain dynamics in line with insights and metrics from complexity science [25]. This approach was followed to demonstrate that LSD increases the complexity of spontaneous brain activity as assessed via fMRI [26], an observation consistent with previous studies showing an expanded repertoire of brain states with facilitated transitions between them [19,27], constituting a potential mechanism for the psychedelic-induced enhancement of neural flexibility [28]. Moreover, the dynamical characterization of psychedelic action reveals a scenario opposite to that of unconsciousness, where a reduction in the repertoire of brain states and an increase in their stability against perturbations is observed [25,29,30] as hypothesized in previous theoretical work [24]

To date, whole-brain models have been used to investigate the mechanisms underlying the steady state effects of LSD and psilocybin, two classic psychedelic drugs. However, it is currently unknown whether the putative mechanisms of psychedelic action identified by these models also extend to the transitions from pre-dose baseline and the acute effects. Time-dependent models capable of fitting dynamics across transitions are necessary to distinguish between the

neurochemical modulation of brain activity, aligned with the drug pharmacokinetics, and the indirect effects of factors linked to the interplay between expectations, subjective effects and the short-term impact of the experience on the emotional state of the subjects [1]. Moreover, developing computational models sensitive to slow temporal fluctuations in brain activity may contribute to determine whether the pharmacology of certain psychedelics is multi-phasic, as has been proposed for the case of LSD [31]. The short-lasting psychedelic effects of intravenous DMT are ideally suited for this purpose, as they are highly dynamic and give way to the recovery of the baseline state after around 30 minutes [32]. Making progress in this direction, recent studies have found that specific EEG features of the DMT state are correlated with the serum concentration of the drug [17,33]. However, the dynamical principles behind the transition to altered consciousness remain to be addressed by modeling efforts.

In this work, we adopted a model previously used to investigate the effects of psychedelic drugs [26], introducing time dependency in the parameter governing the proximity to dynamical criticality, i.e. a phase transition between qualitatively different dynamics (chaos/noise vs. statistical regularities/oscillations, as shown by Piccinini et al. 2021). To model the relatively short-lasting effects induced by DMT, the temporal evolution of the model bifurcation parameter was constrained to represent a simplified version of DMT pharmacokinetics [34]. The optimization of the free parameters underlying this temporal evolution allowed us to reproduce the empirical functional connectivity dynamics (FCD) [35], and subsequently the optimal peak intensity and latency for DMT vs. a placebo control condition. Finally, we investigated the stability of the simulated dynamics against external perturbations, thus assessing potential correlations between the local regional reactivity and the density of 5HT2a receptors, the main pharmacological target of DMT [2].

Results

Figure 1 provides an outline of the methods and the overall procedure. Following previous work, the local dynamics in the whole-brain model were given by Stuart-Landau nonlinear oscillators, with a bifurcation at a=0 [35]. For a>0 the model presents stable oscillations, while a<0 results in stable spirals which extinguish the oscillation amplitude until the dynamics are dominated by the additive noise term, η(t). Close to the bifurcation (i.e. dynamic criticality), the noise introduces spontaneous switches between both regimes, resulting in oscillations with complex amplitude fluctuations. To model the time-dependent changes introduced by DMT, we put forward an assumed equation for the bifurcation parameter a(t) given by the gamma function shown in Fig. 1. Parameters λ and β determine the peak amplitude and its latency, respectively. For an adequate choice of parameters, this function rises rapidly and then presents a slow decay, constituting an approximate description of the pharmacokinetics of bolus intravenous DMT administration [34]. The local Stuart-Landau nonlinear oscillators were coupled using an averaged structural connectivity matrix inferred using diffusion tensor imaging (DTI) from healthy individuals and scaled by the global coupling parameter G. Since we aimed to capture the temporal evolution introduced by the short-acting effects of DMT, we optimized the gamma function parameters λ and β to fit the reproduction of the mean FCD computed from 15 participants, which contains in its i, j entry the similarity between functional connectivity (FC) matrices computed over short windows starting at time points i and j (see Methods). Note that in contrast to previous applications of this model to resting state data [21,35-37], we did not optimize the statistical similarity between the empirical and simulated FCD matrices; instead, we employed the Frobenius distance for their comparison, as we were interested in capturing the temporal evolution introduced by DMT.

To determine the optimal gamma function parameters to reproduce the whole-brain dynamics represented by the FCD, we performed an exhaustive exploration of all pairwise combinations of λ

and β within a range of values compatible with the expected DMT pharmacokinetics (see Methods).

The results of this exploration are shown in Fig. 2A for the placebo (left) and DMT conditions

(right), with lower values indicating a better model fit. For the placebo condition, the optimal values

were located within a large region of comparatively low amplitude and variable latency, evidencing

a weak temporal dependency without a clearly defined peak of maximal intensity. Conversely, for

DMT this region was displaced and reduced to encompass shorter and less variable latencies,

together with larger amplitude values. Figure 2B represents the empirical (left column) and optimal

simulated (right column) FCD matrices for the placebo (first row) and DMT conditions (second

row). Note that the two diagonal blocks of the FCD matrices separate the baseline (see Fig.1,

"Functional connectivity dynamics") and post-administration periods. It can be seen that the model

is capable of approximating this temporal structure, as well as the overall intensity of the matrix

element values.

After determining the optimal λ and β for each independent run of the simulation, we explored the

corresponding $a(t)$ plots displayed in Fig. 3A for placebo and DMT, with thicker lines indicating

the mean across all simulations. This figure shows that dynamics start from a baseline of sustained

oscillations at $a=0.07$. After DMT infusion, we observed a sharp and rapid decrease of $a(t)$ which

displaced whole-brain dynamics towards the bifurcation at $a=0$. After peaking after ≈ 5 minutes, the

parameter $a(t)$ gradually recovered towards baseline values at the end of the scanning session. In

contrast, results for the placebo condition showed a lower peak amplitude combined with longer

latencies, to the point where the peaks were not reached during the scanning session. As a

consequence, $a(t)$ for the placebo condition approximated the constant baseline value with

comparatively smaller temporal variation. Figure 3B summarizes the differences between the

dynamics of $a(t)$ found for placebo and DMT in the two-dimensional space spanned by λ and β,

where each point represents optimal values for an independent run of the simulation, and the larger

full circles indicate the mean across all simulations. For the DMT condition, we could observe that

λ and β were clustered in the upper left corner, indicating comparatively low latencies and high

amplitude peaks (λ=159.3 ± 7, β=284 ± 37, 95% confidence interval [CI]). Conversely, the placebo

condition resulted in low amplitude values and more variable latencies, skewed towards larger

values relative to DMT (λ=65.6 ± 9, β=588 ± 69, 95% CI). Both clusters of values could be clearly

separated, supporting the finding of qualitatively different a(t) dynamics between conditions (t-test

p < 0.0001 for both parameters)

Figure 1. *Overview of the whole-brain model and the choice of optimization target.* The upper left panel

("Local model") presents the dynamics of a single node, consisting of a Stuart-Landau nonlinear oscillator

with bifurcation parameter a, and the imaginary and real part of complex variable z vs. a, showing a transition

between stable spirals and limit cycles at a=0. The bottom left panel ("Time dependency") presents the

temporal parametrization of the bifurcation parameter a(t), given by a gamma function with parameters λ

(peak amplitude) and β (peak latency). The upper right panel ("Inter-areal coupling") illustrates how the

nodes were coupled following the structural connectivity given by DTI scaled by G to reproduce the empirical

FCD. Finally, the bottom right panel ("Functional connectivity dynamics") shows the computation of the

FCD matrix, which contains in its i, j entry the similarity between FC matrices computed over short windows

starting at time points i and j. The diagonal block encased in red indicates the baseline period before the

administration of DMT.

Figure 2. *Exploration of parameter space and optimal Functional Connectivity Dynamics.* **A)** Averaged

parameter space (n = 50) reveals different peak amplitude (λ) and latency (β) values for placebo vs. DMT.

Optimal performance of DMT is restricted to a narrower region **B)** Empirical and optimal simulated FCD

(columns) for the placebo and DMT conditions (rows) average over n = 15 subjects (simulations).

Figure 3. *The temporal evolution of the bifurcation parameter, $a(t)$, distinguishes between DMT and the*

placebo condition. **(A)** a(t), defined as a gamma function, for n = 50 independent runs of parameter

optimization, compared between both conditions. Plots with thicker lines indicate the curves associated with

mean value parameters. **(B)** Two-dimensional representation of the optimal gamma function parameters, λ

and β, both for placebo and DMT. Individual points indicate the outcome of the n =50 independent runs of

parameter optimization, while the larger full circles represent the average across simulation runs.

Comparisons for the means of both parameters were made with two-sided t-tests with p-values < 0.0001 for

both parameters.

After determining the optimal parameters to fit the FCD, we used the resulting models to investigate the time-

dependent effects of delivering external perturbations. This analysis was motivated in previous reports

indicating that psychedelics increase the reactivity to external stimuli and facilitate transitions between brain

metastable states [26,27]. Based on these results, we hypothesized that DMT would result in a transient episode

of more sensitive brain dynamics, arising due to a destabilization of whole-brain dynamics, i.e. due to the

increased proximity to dynamic criticality.

To test our hypothesis, we introduced a periodic external perturbation at the frequency of the

endogenous oscillations, which was previously shown to maximize the effect on the ongoing brain

activity [38,39]. We determined the reactivity or sensitivity to the perturbation at each time point, noted

here by $\chi(t)$, computing the derivative of the induced FCD changes with a numerical

differentiation algorithm (see methods for more details). To facilitate the interpretation of the

results, we introduced this perturbation at nodes located within six resting state networks (RSN) [40]

known to encompass different functional brain systems: primary visual (Vis), extrastriate (ES),

auditory (Aud), sensorimotor (SM), default mode (DM) and executive control (EC) cortices. Figure

4A shows $\chi(t)$ for each RSN, both for the placebo (top) and DMT (bottom) conditions. For the

latter, it is clear that $\chi(t)$ is aligned with the expected evolution of drug effect intensity, with the

largest reactivity obtained at the extrastriate visual cortex. In contrast, the time dependency of $\chi(t)$

was considerably less marked for the placebo group. Figure 4B displays the peak of $\chi(t)$, i.e. χ_{max},

for each RSN and for three different external perturbation intensities (F_{ext}), comparing the DMT

and placebo conditions. Consistent with the results shown in Fig. 4A, the peak values were

systematically lower in placebo vs. DMT for all intensities, indicating higher reactivity to external

perturbations during the acute effects of the drug.

Figure 4. *Time-dependent effect of simulated perturbations indicate higher reactivity for DMT vs. placebo.*

(A) Reactivity $\chi(t)$ normalized by the number of nodes in the corresponding RSN, for placebo (top) and DMT

(bottom). DMT curves peak around 4 minutes after the dose, restoring baseline values at the end of the session.

Placebo shows a lower peak amplitude and longer latencies remaining constant during the whole study. Shaded

regions of each line denote the standard deviation of the bootstrapped reactivities ($n_{boot} = 200$). **(B)** Plots of the

peak $\chi(t)$ across time (χ_{max}) for each RSN and three different external perturbation intensities (F_{ext}).

Comparisons between conditions (for a given perturbation intensity) and between successive perturbation

intensities (for a given condition) were made with two-sided t-tests, giving p-values < 0.0001 for all

comparisons. Executive control (EC) network was an exception when comparing $F_{ext} = 0.005$ and $F_{ext} = 0.01$ (p

$= 0.23$).

Finally, we investigated the correlation between the 5HT2a receptor density (see Methods) and the

peak reactivity (χ_{max}) across RSN. Based on the known pharmacological action of DMT [2], we

expected a positive correlation between these two variables, i.e. that RSN with higher 5HT2a

receptor density would show higher χ_{max} values, and vice-versa. Figure 5A illustrates the spatial

configuration of the RSN, while the mean 5HT2a receptor density per RSN is shown in Fig. 5B. To

assess if the difference in χ_{max} between conditions can be attributed to the local density of

serotonin receptors, we first computed the difference between the reactivity curves of DMT and

placebo, resulting in $\Delta\chi(t)$. Figure 5C shows the peak of $\Delta\chi(t)$, $\Delta\chi_{max}$, vs. the 5HT2a receptor

density of each RSN, together with the best least-squares linear fit. To estimate the significance of

this correlation, we conducted a bootstrap procedure which resulted in the distribution of correlation

coefficients (ρ) presented in the inset of this panel, with a mean value of $\rho = 0.9059 \pm 0.0003$,

95% CI. Figure 5D displays the mean ρ ($\Delta\chi_{max}$ vs. 5HT2a receptor density) across a range of

external perturbation intensities (F_{ext}). This plot indicates that low stimulation amplitudes exert an

effect that is comparatively independent of the local 5HT2a receptor density. However, as the

perturbation intensity increases, the receptor density becomes more relevant, peaking at $\rho > 0.9$.

Finally, as the intensity keeps increasing, the reactivity decouples again from the receptor density.

Figure 5. *The peak differential reactivity to external perturbations ($\Delta\chi_{max}$) between DMT and placebo correlates with the local 5HT2a receptor density.* **(A)** Spatial configuration of the six RSN that constrained the simulated delivery of the perturbation (Vis: primary visual, ES: extrastriate, Aud: auditory, SM: sensorimotor, DM: default mode, EC: executive control), adapted from [40]. **(B)** Mean 5HT2a receptor density per RSN. **(C)** Peak of $\Delta\chi(t)$, $\Delta\chi_{max}$, vs. the 5HT2a receptor density of each RSN (mean $\pm$ SD) together with the best least-squares linear fit. The inset shows the distribution of correlation coefficients (ρ) obtained using a bootstrap procedure (n = 1000). **(D)** Correlation coefficient ρ vs. the external perturbation intensity, F_{ext}.

Discussion

This study represents a first step towards the computational modeling of time-dependent psychedelic effects. Our main finding is that DMT destabilizes (i.e. brings closer to the global bifurcation point, or dynamic criticality) whole-brain dynamics, and that the extent of this destabilization is compatible with the characteristic pharmacokinetics of the drug, here constrained by a gamma function [34]. Conversely, an inactive placebo resulted in extreme parameter values that

flattened this function, thus approximating a constant value over time. A consequence of this loss of

stability is the heightened sensitivity to external perturbations [38], paralleling the dynamics of the

bifurcation parameter and thus being maximal when the perturbation is applied in nodes belonging

to RSN with high 5HT2a receptor density. This increased sensitivity may affect how the brain

responds to incoming sensory stimuli under the effects of the drug, and may also impact on its

capacity to amplify endogenous events linked to ongoing mentation and cognitive processing. Note

our emphasis on *endogenous events* in this regard, as a converse reduced sensitivity to perturbation

may be true in relation to certain external perturbations, such as sensory evoked potentials ([41]).

Indeed, recent work showed that external stimulation delivered during the psychedelic state has

stronger effects relative to a control condition, which include changes in the intensity of the

experience as well as a substantial modification of entropy-based metrics of neural activity [42]

Previous studies have used similar models to investigate the mechanisms underlying psychedelic-

induced changes in fMRI resting state activity [21-23,26]. However, these studies were concerned

primarily with the steady state effects of the drugs, thus neglecting the analysis of periods when

their intensity changes over time, such as the transitions between the baseline and the acute state.

These transitions are difficult to capture for the oral administration of LSD or psilocybin, which

results in a slower onset and a less marked transition towards the psychedelic state [43,44]. In contrast,

the intravenous administration of DMT results in a fast-onset and relatively brief psychedelic

episode that peaks in subjective intensity only a few minutes after the infusion [32] as indeed occurred

in our previous work [9,17]. Our current study shows that the whole-brain FCD dynamics induced by

DMT on fMRI data recapitulate this temporal evolution, which was not observed for the placebo. It

is important to note that even if the intensity of the DMT effects is not reflected on the amplitude of

the time-domain fMRI signals, whole-brain FCD contains sufficient information to determine the

temporal evolution of the DMT experience as shown by the optimal fit of the model to the FCD

data.

The transient destabilization of brain dynamics induced by DMT is consistent with multiple

experimental results and theoretical accounts of psychedelic action in the human brain [24]. Drawing

from the theory of bifurcations in dynamical systems, as the model approximates the bifurcation

point, both the complexity and entropy of simulated brain activity are expected to increase [13-19],

together with an expansion in the repertoire of possible metastable states (Tagliazucchi, Carhart-

Harris et al. 2014). Also, near the bifurcation point, the sensitivity of the system to external

perturbations is maximized [26,38], predicting a larger response to a perturbation within the

psychedelic state. A prolonged period of recovery from perturbation is known as *critical slowing*

and is considered to be one of several signatures of critical states, such as a global dynamic

bifurcation [45]. Using non-invasive transcranial magnetic stimulation (TMS) combined with

electroencephalography (EEG), Ort and colleagues could not find changes in cortical reactivity

induced by psilocybin; however, they informed changes in spectral content and subjective

experience linked to the stimulation [46]. This result is partially consistent with our prediction,

especially when taking into account the differences between TMS and a periodic perturbation

delivered at the natural frequency of the endogenous oscillations. Empirically, this perturbation

could be achieved by non-invasive methods such as transcranial alternating current stimulation

(tACS) [47]; however, to date this form of stimulation has not been investigated in participants

undergoing the effects of psychedelic drugs.

Our results draw a parallel between the effects of DMT on whole-brain dynamics and the theory of

dynamical and statistical criticality. The main difference between conditions was the behavior of the

bifurcation parameter across time, resulting in values closer to the critical value coincident with the

peak effects of DMT. This finding, which is consistent with prior work implying increased

signatures of criticality in the psychedelic state [45,48,49], can be interpreted through the critical brain

hypothesis, stating that the major features of brain dynamics can be explained by proximity to a

second order phase transition [50,51]. Near this critical point, the system exhibits divergent

susceptibility (i.e. reactivity), allowing minor perturbations to propagate throughout the entire

system [52]. Extending this analogy, we can postulate that 5HT2a receptor activation shifts the

network state towards criticality, which is consistent with multiple other findings related to the

brain dynamics under the effects of psychedelic drugs, as discussed by [25]. We are mindful however,

that certain specific stimuli consonant with the ongoing state, such as music [53], may have a bigger

psychological and, presumably, neurobiological [54] impact under psychedelics. This matter may

point to the special value of utilizing naturalistic stimuli with good ecological validity in the context

of psychedelic research.

We also report that the sensitivity to external perturbations covariates with the density of 5HT2a

receptors. Since we normalized this value by the total number of nodes in each RSN, the

explanation for this correlation should be found in the influence of 5HT2a in the organization of

functional and structural connectivity. This serotonin receptor subtype is implicated in large-scale

heterogeneities of the human cerebral cortex, constituting an important factor to distinguish

between unimodal and the higher level integrative transmodal cortex [55]. Moreover, by bringing the

global dynamics closer to dynamical criticality, 5HT2a receptor activation may facilitate the

switching between inter-areal coupling that may underly psychedelic-induced neural [56,57] and

possibly cognitive flexibility [28], and also open a transient window where the increased functional

diversity may exert an effect in the underlying structural connectivity via plasticity effects [58].

Eventually, this could contribute to consolidating the long-term effects associated with brief

psychedelic episodes [59]. For high values of the external perturbation amplitude, the reactivity

decoupled from the 5HT2a receptor density, likely indicating the saturation of the induced effects

on whole-brain dynamics.

Beyond the current application, the temporal parametrization of whole-brain models may find other

uses to test whether a certain temporal process underlies the recorded neuroimaging data, and to

inform the potential mechanisms behind the empirical observations. The signal preprocessing steps

applied to resting state fMRI data typically include high pass filters which are necessary to attenuate

the effects of scanner drift and head movement [60]. Therefore, the fMRI time series may not directly

reflect the dynamics of the phenomenon under study, unless the experiment is designed to

maximize the signal-to-noise ratio, e.g. with tasks or stimulation structured in according to an

adequate block design [61]. However, it may not be possible to control the onset and length of the

temporal process under study, as in the current application to the study of a transient

pharmacologically-induced event. This could also occur in the study of seizures [62], sleep-related

transients and graphoelements (e.g. spindles, K-complexes) [63], spontaneous mentation or ongoing

cognition [64], and in other endogenous phenomena with an interesting temporal structure beyond

experimental control. In these cases, the use of time-dependent whole-brain models may be useful

to detect the fingerprints of the event in the FCD, by comparing different temporal parametrizations

by fitting and contrasting their corresponding goodness of fit.

One important limitation of our approach is that the temporal parametrization a priori constrains the

model and its capacity to detect the underlying temporal dynamics. By choosing a gamma function,

our approach only allowed us to test whether the FCD dynamics under DMT could be better

explained by this specific temporal evolution of the bifurcation parameter relative to the placebo

condition. However, since we did not specify a prior the direction of the effect, this approach

allowed us to test whether DMT could bring the dynamics closer to the global bifurcation and

therefore towards a point of decreased stability. The choice of the gamma function can be justified

by the prior literature on the time course of DMT-induced effects, as well as its effects on EEG

activity and their correlation with the drug pharmacokinetics [9,16,17,32]. While more complex models

could be implemented to adequately describe these dynamics [33], our choice has the merit of

constituting a qualitative description of the transient effects of DMT without incurring in a

proliferation of free parameters required for more neurobiological realism. Moreover, the

comparatively poor temporal resolution of fMRI may limit the potential gain of including a more

nuanced description of pharmacodynamics. This limitation could be addressed by reproducing our

work with data from faster imaging modalities (such as EEG and MEG).

In summary, our work shifts the focus from the reproduction of the steady-state effects of

psychedelics towards disentangling the dynamics of the fast-onset and short-lived intravenous DMT

experience. The future implementation of more realistic biophysical models could contribute to our

understanding of how the interaction between drugs, neurotransmitters and receptors is capable of

initiating a cascade of neural events which ultimately results in a global brain state associated with

profound alterations in consciousness and cognitive processing with potentially lasting

consequences [59]. The combination of these models with a more nuanced description of drug

pharmacokinetics and pharmacodynamics could also contribute to explain the potential multiphasic

effects of some psychedelic compounds, and to test potential mechanisms behind their long-lasting

effects, which are a key aspect of their possible therapeutic use in patients with depression and other

psychiatric disorders.

Materials and Methods

Study participants and experimental design

This is a re-analysis of a previously published EEG-fMRI dataset acquired from healthy participants

under the acute effects of DMT [9]. The original publication can be referenced for in-depth

methodological details.

The study followed a single-blind, placebo-controlled design, with all participants providing written

informed consent. The experimental protocol received approval from the National Research Ethics

Committee London—Brent and the Health Research Authority, conducted in adherence to the

Declaration of Helsinki (2000), International Committee on Harmonization Good Clinical Practices

guidelines, and the National Health Service Research Governance Framework.

Volunteers complete two visits at the Imperial College Clinical Imaging Facility, spaced two weeks

apart. On each testing day, participants were subject to separate scanning sessions. In the initial

session, they received intravenous administration of either placebo (10 mL sterile saline) or 20 mg

DMT (fumarate form dissolved in 10 mL sterile saline). This was done in a counter-balanced order,

with half receiving placebo and the rest receiving DMT. The first session comprised continuous

resting-state scans lasting 28 minutes, with DMT/placebo administered at the end of the 8th minute,

and scanning concluding 20 minutes after injection. Participants lay in the scanner with closed eyes

aided by an eye mask, while fMRI data was recorded. In the second session, participants were cued

to verbally rate the subjective intensity of drug effects every minute in real-time. Only fMRI data

from the first scanning session was used for the present analysis.

A cohort of 20 participants completed all study visits, consisting of 7 females with a mean age of

33.5 years and a standard deviation of 7.9. For the present study data from only 15 of the subjects

were used, the rest being discarded due to strong head movement artifacts inside the scanner (see

below for further information on the exclusion criterion).

fMRI acquisition and preprocessing

Images were acquired using a 3T MR scanner (Siemens Magnetom Verio syngo MR B17) with a

12-channel head coil for compatibility with EEG acquisition. Functional imaging was performed

using a T2*-weighted BOLD-sensitive gradient echo planar imaging sequence [repetition time (TR)

= 2000ms, echo time (TE) = 30 ms, acquisition time (TA) = 28.06 mins, flip angle (FA) = 80°,

voxel size = $3.0 \times 3.0 \times 3.0$mm3, 35 slices, interslice distance = 0 mm].

The preprocessing steps involved despiking, slice time correction, motion correction, brain

extraction, rigid body registration to anatomical scans, nonlinear registration to 2mm MNI brain,

denoising (via Independent Component Analysis), and scrubbing (using an framewise displacement

[FD] threshold of 0.4, with scrubbed volumes replaced by the mean of surrounding volumes) [65].

Further steps included spatial smoothing (FWHM) of 6 mm, band-pass filtering between 0.01 and

0.08 Hz, linear and quadratic detrending, and regressing out nine nuisance regressors. These

regressors, all band-pass filtered, consisted of six motion-related parameters (3 translations, 3

rotations) and three anatomically related parameters (ventricles, draining veins, local white matter).

Finally, time series were extracted for each Automated Anatomical Labeling (AAL) [66] template

region by averaging the time series from all voxels within the corresponding region.

Out of 20 participants, five were excluded from group analyses due to excessive head movement

during the 8-minute post-DMT period (>20% of scrubbed volumes with an framewise displacement

threshold [FD] of 0.4).

Anatomical connectivity matrix

The anatomical connectivity matrix was obtained applying diffusion tensor imaging (DTI) on

diffusion-weighted imaging (DWI) data obtained from 16 healthy right-handed participants (11 men

and 5 women, mean age: 24.75 ± 2.54 years), recruited online at Aarhus University, Denmark.

Participants with psychiatric or neurological disorders (or a history thereof) were excluded. For

detailed MRI acquisition parameters, we refer readers to a previous publication. Anatomical

connectivity networks were constructed in three steps. Firstly, whole-brain network regions were

defined using the AAL atlas. Secondly, connections between nodes in the whole-brain network

(edges) were estimated through probabilistic tractography applied to individual participant DTI

data. Lastly, results were averaged across participants.

DTI preprocessing utilized the probtrackx tool from the FSL diffusion imaging toolbox (Fdt)

(www.fsl.fmrib.ox.ac.uk/fsl/fslwiki/FDT) with default parameters. Subsequently, local probability

distributions of fiber directions were estimated at each voxel. The connectivity probability from

seed voxel i to another voxel j was defined as the proportion of fibers passing through voxel i that

reached voxel j, with 5000 streamlines sampled per voxel. This process extended from the voxel to

the region level, where in a region of interest consisting of n voxels, $5000 \times n$ fibers were sampled.

The connectivity probability from region i to region j was calculated as the number of sampled

fibers in region i that connected the two regions, divided by $5000 \times n$, where n represents the

number of voxels in region i. The resulting anatomical connectivity matrices were then thresholded

at 0.1% (equivalent to a minimum of five streamlines), yielding the anatomical connectivity

matrices C_{ij} used in the model.

Whole-brain computational model

We simulated brain activity measured with fMRI at the whole-brain level by using a Stuard-Landau

oscillator (i.e. Hopf normal form) for the local dynamics (see Fig. 1, "Local model"). This

phenomenological model aims to directly simulate recorded brain signals. The emergent global

dynamics are simulated by including mutual interactions between brain areas according to the

anatomical connectivity matrix $\boldsymbol{C_{ij}}$ obtained from DTI (see Fig. 1, "Inter-areal coupling"). The full

model consists of 90 coupled nodes representing the 90 cortical and subcortical brain areas from

AAL parcellation, with the following temporal evolution for region $\boldsymbol{n}$:

1
$$\frac{dx_n}{dt} = (a(t) - x_n{}^2 - y_n{}^2)x_n - \omega y_n + G \sum_{p=1}^{90} C_{np}(x_p - x_n) + \gamma\eta_n(t)$$

2
$$\frac{dy_n}{dt} = (a(t) - x_n{}^2 - y_n{}^2)x_n + \omega_n x_n + G \sum_{p=1}^{90} C_{np}(y_p - y_n) + \gamma\eta_n(t)$$

3
(1)

Here, η_n represents additive Gaussian noise with standard deviation γ (set to 0.05), C_{np} are the

matrix elements of the anatomical connectivity matrix, G is a factor that scales the anatomical

connectivity (fixed at $G = 0.5$, as determined previously elsewhere (Ipina et al., 2020), ω_n is the

peak frequency of node n in the 0.01-0.08 Hz band (determined using Fourier analysis applied to

the empirical time series and averaged across participants), and variable x_n represents the empirical

brain activity signal of node n, which was used to compute the simulated FC and FCD matrices.

The local dynamics present a supercritical bifurcation at a=0, so that if a>0 the system engages in a

stable limit cycle (i.e. oscillations) with angular frequency ω_n, and when a<0 the local dynamics

are attracted to a stable fixed point representing a low activity state dominated by noise. Close to the

bifurcation, the additive noise can induce a switching behavior between regimes, resulting

oscillations with a complex envelope [35].

To model the short-lasting effects of DMT, we introduce a time-dependent bifurcation parameter

a(t) following a gamma function that represents a simplified description of drug pharmacokinetics

[34]:

21
$$a(t) = \lambda\left(\frac{te^{-t/\beta}}{N}\right)$$

22
(2)

where N is a constant that normalizes the term between brackets, λ is a scale parameter determining

the amplitude of the peak, and β the parameter that controls the rate of decay of the function, and

therefore is related to the latency of the peak (see Fig. 1, "Time dependency").

Model fitting to empirical data

To characterize the time-dependent structure of brain dynamics, we computed the FCD matrices

(see Fig. 1, "Functional connectivity dynamics") [36]. For this purpose, each full-length signal of 28

min was split up into M = 82 sliding windows of 60 s each, with an overlap of 40 s between

successive windows. For each sliding window centered at time t, the functional connectivity matrix

FC(t) was computed. The FCD matrix is a M $\times$ M symmetric matrix whose t_1, t_2 entry is defined

by the Pearson correlation coefficient between the upper triangular parts of the two matrices $FC(t_1)$

and $FC(t_2)$. These matrices were computed for each of the fifteen participants and simulations by

exhaustively exploring the model parameters related to the temporal evolution of the bifurcation

parameter, λ and β. To compare the FCD matrices taking into account their temporal structure, we

used the Frobenius distance between the elements of the empirical and simulated matrices.

Model fitting to baseline data

To set the model to the baseline conditions before the administration of the drug, we conducted a

search for the best value of the bifurcation parameter by fitting the FCD submatrix of the first 22

temporal windows. This corresponds to the first 8 minutes previous to the injection of DMT. The

fitting was done for both conditions (DMT and placebo) and, as expected, resulted in the same

value of a = 0.07.

Fitting the temporal evolution of the bifurcation parameter

We fixed the value of the bifurcation parameter to the baseline value of a=0.07 for the first 8

minutes of the simulation and introduced the time dependency afterwards, given by the difference

between the baseline value and the gamma function $\lambda \left(\frac{te^{-t/\beta}}{N} \right)$, with t=0 here indicating the time of

the drug injection. Next, we performed an exhaustive exploration of the parameter space spanned by

λ and β, searching for the optimal combination of parameters. For this purpose, we explored a grid

given by $\lambda = [0, 200]$ and $\beta = [20, 900]$ in steps of 5 and 20 units, respectively. For each parameter

combination, we computed the FCD 15 times (i.e. once per participant) randomly changing the

initial conditions of the model. The resulting FCD matrices were averaged and compared with the

empirical FCD using the Frobenius distance. This procedure was repeated 50 times for each pair of

parameters.

Simulated perturbations

Next, we modeled an oscillatory perturbation and investigated the response of the whole-brain

model fitted as explained in the previous section. The perturbation was given by an external

additive periodic forcing term applied to node $\mathbf{n}$, $\mathbf{F_n} = \mathbf{F_{ext}}(\cos(\boldsymbol{\omega_n} t) + i \sin(\boldsymbol{\omega_n} t))$, where the

real part of the perturbation is added to the equation for $\frac{dx_n}{dt}$ in Eq. 1 and the imaginary part to the

equation for $\frac{dy_n}{dt}$. In the equation for $\mathbf{F_n}$, $\boldsymbol{\omega_n}$ is the natural frequency of the time series

corresponding to node $\mathbf{n}$ (same as in Eq. 1).

To facilitate the interpretation of the results, we applied this perturbation to nodes located within six

different resting state networks (RSN) identified using independent component analysis (ICA) by

Beckmann, et al. [40]. To account for the time variation of the reactivity, we sampled equally spaced

values of $a(t)$, here noted a_p, with p indexing the time sample. Then, for every one of those values,

we performed a simulation keeping a_p constant until the end of the simulation, i.e. for every

simulation we set a=0.07 for the first 8 minutes and then the constant value, a_p until the end of the

simulation. The perturbation was applied after the first 8 minutes corresponding to the baseline,

varying the amplitude F_{ext} from 0 to 0.015 in 00125 steps. The maximum value of the range was

chosen as it saturated the correlation between 5HT2a receptor density and local reactivity, as shown

in Fig. 5. For each combination of RSN, F_{ext} and a_p we computed the resulting FCD and its

distance to the empirical condition, and then assessed the impact of the perturbation as explained in

the following section.

Measure of reactivity to perturbations

We interpret the whole-brain model reactivity as the sensitivity of brain activity to the external

periodic stimulation. Following an analogy with the concept of susceptibility in statistical physics,

we defined the reactivity as the following derivative:

$$\chi(t) \; = \; \frac{\partial M}{\partial F_{ext}}$$

12   (3)

where M denotes the Frobenius distance between the simulated and empirical FCD matrices.

Therefore $\chi(t)$ quantifies how a certain RSN in a particular moment responds to an external

perturbation in terms of its impact on the Frobenius metric. This magnitude was computed using a

second order finite difference method. We evaluated $\chi(t)$ relative to its value at t=0. Furthermore,

given that the number of nodes differs across the RSN, we computed the $\chi(t)$ normalized by the

number of nodes of each RSN.

Bootstrap method

To determine the peak of $\chi(t)$ across the duration of the simulation, we used a bootstrapping

procedure to obtain a distribution of peak values which allowed us to compute a confidence interval

for the resulting average value. The bootstrap procedures were conducted by randomly drawing

samples (with replacement) from the distribution of values, creating different $\chi(t)$ curves in each

iteration and measuring its maximum value. The size of the sampled subset was equal to that of the

original distribution. This procedure was repeated 200 times, generating a bootstrap distribution of

the desired magnitude. When calculating the correlations between local 5HT2a receptor density and

the maximum reactivity per RSN, the bootstrap was done 1000 times to generate the histograms.

Statistical tests and Confidence Intervals

95% Confidence Intervals were calculated using the standard error of the mean (SEM) and a z-score

of 1.96. Metric comparisons were achieved using a two-sided paired t-test assuming non-equal

variance between pair of metrics. To determine the peak location and significance of $\chi(t)$ across the

duration of the simulation, we implemented a bootstrap procedure (see section "*Bootstrap*

method").

Receptor density maps

The receptor density maps used were estimated using PET tracer studies obtained by Hansen and

colleagues [67]. All PET images were registered to the MNI-ICBM 152 nonlinear 2009 (version c,

asymmetric) template and subsequently parcellated to the 90 region AAL atlas (Tzourio-Mazoyer et

al., 2002). For more details on the acquisitions and limitations of the data set see the original

publication [67].

References

1   Kwan, A. C., Olson, D. E., Preller, K. H. & Roth, B. L. The neural basis of psychedelic action.
       Nat Neurosci **25**, 1407-1419, doi:10.1038/s41593-022-01177-4 (2022).

2   Nichols, D. E. Psychedelics. *Pharmacol Rev* **68**, 264-355, doi:10.1124/pr.115.011478
       (2016).

Wallach, J. *et al.* Identification of 5-HT(2A) receptor signaling pathways associated with psychedelic potential. *Nat Commun* **14**, 8221, doi:10.1038/s41467-023-44016-1 (2023).

Gonzalez-Maeso, J. *et al.* Hallucinogens recruit specific cortical 5-HT(2A) receptor-mediated signaling pathways to affect behavior. *Neuron* **53**, 439-452, doi:10.1016/j.neuron.2007.01.008 (2007).

Zamberlan, F. *et al.* The Varieties of the Psychedelic Experience: A Preliminary Study of the Association Between the Reported Subjective Effects and the Binding Affinity Profiles of Substituted Phenethylamines and Tryptamines. *Front Integr Neurosci* **12**, 54, doi:10.3389/fnint.2018.00054 (2018).

Carhart-Harris, R. L. *et al.* Psychedelics and the essential importance of context. *J Psychopharmacol* **32**, 725-731, doi:10.1177/0269881118754710 (2018).

Tagliazucchi, E. *et al.* Increased Global Functional Connectivity Correlates with LSD-Induced Ego Dissolution. *Curr Biol* **26**, 1043-1050, doi:10.1016/j.cub.2016.02.010 (2016).

Carhart-Harris, R. L. *et al.* Neural correlates of the LSD experience revealed by multimodal neuroimaging. *Proc Natl Acad Sci U S A* **113**, 4853-4858, doi:10.1073/pnas.1518377113 (2016).

Timmermann, C. *et al.* Human brain effects of DMT assessed via EEG-fMRI. *Proc Natl Acad Sci U S A* **120**, e2218949120, doi:10.1073/pnas.2218949120 (2023).

Preller, K. H. *et al.* Changes in global and thalamic brain connectivity in LSD-induced altered states of consciousness are attributable to the 5-HT2A receptor. *Elife* **7**, doi:10.7554/eLife.35082 (2018).

Luppi, A. I. *et al.* LSD alters dynamic integration and segregation in the human brain. *Neuroimage* **227**, 117653, doi:10.1016/j.neuroimage.2020.117653 (2021).

Bedford, P. *et al.* The effect of lysergic acid diethylamide (LSD) on whole-brain functional and effective connectivity. *Neuropsychopharmacology* **48**, 1175-1183, doi:10.1038/s41386-023-01574-8 (2023).

Herzog, R. *et al.* A whole-brain model of the neural entropy increase elicited by psychedelic drugs. *Sci Rep* **13**, 6244, doi:10.1038/s41598-023-32649-7 (2023).

Schartner, M. M., Carhart-Harris, R. L., Barrett, A. B., Seth, A. K. & Muthukumaraswamy, S. D. Increased spontaneous MEG signal diversity for psychoactive doses of ketamine, LSD and psilocybin. *Sci Rep* **7**, 46421, doi:10.1038/srep46421 (2017).

Viol, A., Palhano-Fontes, F., Onias, H., de Araujo, D. B. & Viswanathan, G. M. Shannon entropy of brain functional complex networks under the influence of the psychedelic Ayahuasca. *Sci Rep* **7**, 7388, doi:10.1038/s41598-017-06854-0 (2017).

Pallavicini, C. *et al.* Neural and subjective effects of inhaled N,N-dimethyltryptamine in natural settings. *J Psychopharmacol* **35**, 406-420, doi:10.1177/0269881120981384 (2021).

Timmermann, C. *et al.* Neural correlates of the DMT experience assessed with multivariate EEG. *Sci Rep* **9**, 16324, doi:10.1038/s41598-019-51974-4 (2019).

Lebedev, A. V. *et al.* LSD-induced entropic brain activity predicts subsequent personality change. *Hum Brain Mapp* **37**, 3203-3213, doi:10.1002/hbm.23234 (2016).

Tagliazucchi, E., Carhart-Harris, R., Leech, R., Nutt, D. & Chialvo, D. R. Enhanced repertoire of brain dynamical states during the psychedelic experience. *Hum Brain Mapp* **35**, 5442-5456, doi:10.1002/hbm.22562 (2014).

Cofre, R. *et al.* Whole-Brain Models to Explore Altered States of Consciousness from the Bottom Up. *Brain Sci* **10**, doi:10.3390/brainsci10090626 (2020).

Deco, G. *et al.* Whole-Brain Multimodal Neuroimaging Model Using Serotonin Receptor Maps Explains Non-linear Functional Effects of LSD. *Curr Biol* **28**, 3065-3074 e3066, doi:10.1016/j.cub.2018.07.083 (2018).

Kringelbach, M. L. *et al.* Dynamic coupling of whole-brain neuronal and neurotransmitter systems. *Proc Natl Acad Sci U S A* **117**, 9566-9576, doi:10.1073/pnas.1921475117 (2020).

Burt, J. B. *et al.* Transcriptomics-informed large-scale cortical model captures topography of pharmacological neuroimaging effects of LSD. *Elife* **10**, doi:10.7554/eLife.69320 (2021).

Carhart-Harris, R. L. *et al.* The entropic brain: a theory of conscious states informed by neuroimaging research with psychedelic drugs. *Front Hum Neurosci* **8**, 20, doi:10.3389/fnhum.2014.00020 (2014).

Girn, M. *et al.* A complex systems perspective on psychedelic brain action. *Trends Cogn Sci* **27**, 433-445, doi:10.1016/j.tics.2023.01.003 (2023).

Jobst, B. M. *et al.* Increased sensitivity to strong perturbations in a whole-brain model of LSD. *Neuroimage* **230**, 117809, doi:10.1016/j.neuroimage.2021.117809 (2021).

Singleton, S. P. *et al.* Receptor-informed network control theory links LSD and psilocybin to a flattening of the brain's control energy landscape. *Nat Commun* **13**, 5812, doi:10.1038/s41467-022-33578-1 (2022).

Doss, M. K. *et al.* Psilocybin therapy increases cognitive and neural flexibility in patients with major depressive disorder. *Transl Psychiatry* **11**, 574, doi:10.1038/s41398-021-01706-y (2021).

Demertzi, A. *et al.* Human consciousness is supported by dynamic complex patterns of brain signal coordination. *Sci Adv* **5**, eaat7603, doi:10.1126/sciadv.aat7603 (2019).

Solovey, G. *et al.* Loss of Consciousness Is Associated with Stabilization of Cortical Activity. *J Neurosci* **35**, 10866-10877, doi:10.1523/JNEUROSCI.4895-14.2015 (2015).

Marona-Lewicka, D., Thisted, R. A. & Nichols, D. E. Distinct temporal phases in the behavioral pharmacology of LSD: dopamine D2 receptor-mediated effects in the rat and implications for psychosis. *Psychopharmacology (Berl)* **180**, 427-435, doi:10.1007/s00213-005-2183-9 (2005).

Strassman, R. J. Human psychopharmacology of N,N-dimethyltryptamine. *Behav Brain Res* **73**, 121-124, doi:10.1016/0166-4328(96)00081-2 (1996).

Eckernas, E., Timmermann, C., Carhart-Harris, R., Roshammar, D. & Ashton, M. N,N-dimethyltryptamine affects electroencephalography response in a concentration-dependent

manner-A pharmacokinetic/pharmacodynamic analysis. *CPT Pharmacometrics Syst Pharmacol* **12**, 474-486, doi:10.1002/psp4.12933 (2023).

Salway, R. & Wakefield, J. Gamma generalized linear models for pharmacokinetic data. *Biometrics* **64**, 620-626, doi:10.1111/j.1541-0420.2007.00897.x (2008).

Deco, G., Kringelbach, M. L., Jirsa, V. K. & Ritter, P. The dynamics of resting fluctuations in the brain: metastability and its dynamical cortical core. *Sci Rep* **7**, 3095, doi:10.1038/s41598-017-03073-5 (2017).

Hansen, E. C., Battaglia, D., Spiegler, A., Deco, G. & Jirsa, V. K. Functional connectivity dynamics: modeling the switching behavior of the resting state. *Neuroimage* **105**, 525-535, doi:10.1016/j.neuroimage.2014.11.001 (2015).

Piccinini, J. *et al.* Noise-driven multistability vs deterministic chaos in phenomenological semi-empirical models of whole-brain activity. *Chaos* **31**, 023127, doi:10.1063/5.0025543 (2021).

Jobst, B. M. *et al.* Increased Stability and Breakdown of Brain Effective Connectivity During Slow-Wave Sleep: Mechanistic Insights from Whole-Brain Computational Modelling. *Sci Rep* **7**, 4634, doi:10.1038/s41598-017-04522-x (2017).

Ipina, I. P. *et al.* Modeling regional changes in dynamic stability during sleep and wakefulness. *Neuroimage* **215**, 116833, doi:10.1016/j.neuroimage.2020.116833 (2020).

Beckmann, C. F., DeLuca, M., Devlin, J. T. & Smith, S. M. Investigations into resting-state connectivity using independent component analysis. *Philos Trans R Soc Lond B Biol Sci* **360**, 1001-1013, doi:10.1098/rstb.2005.1634 (2005).

Murray, C. H. *et al.* Low doses of LSD reduce broadband oscillatory power and modulate event-related potentials in healthy adults. *Psychopharmacology (Berl)*, doi:10.1007/s00213-021-05991-9 (2021).

Pedro A.M. Mediano, F. E. R., Christopher Timmermann, Leor Roseman, David J. Nutt, Amanda Feilding, Mendel Kaelen, Morten L. Kringelbach, Adam B. Barrett, Anil K. Seth,

Suresh Muthukumaraswamy, Daniel Bor, View Robin L. Carhart-Harris. Effects of external stimulation on psychedelic state neurodynamics. *ACS Chem Neurosci* (2024).

Dolder, P. C. *et al.* Pharmacokinetics and Pharmacodynamics of Lysergic Acid Diethylamide in Healthy Subjects. *Clin Pharmacokinet* **56**, 1219-1230, doi:10.1007/s40262-017-0513-9 (2017).

Holze, F., Becker, A. M., Kolaczynska, K. E., Duthaler, U. & Liechti, M. E. Pharmacokinetics and Pharmacodynamics of Oral Psilocybin Administration in Healthy Participants. *Clin Pharmacol Ther* **113**, 822-831, doi:10.1002/cpt.2821 (2023).

Toker, D. *et al.* Consciousness is supported by near-critical slow cortical electrodynamics. *Proc Natl Acad Sci U S A* **119**, doi:10.1073/pnas.2024455119 (2022).

Ort, A. *et al.* TMS-EEG and resting-state EEG applied to altered states of consciousness: oscillations, complexity, and phenomenology. *iScience* **26**, 106589, doi:10.1016/j.isci.2023.106589 (2023).

Helfrich, R. F. *et al.* Entrainment of brain oscillations by transcranial alternating current stimulation. *Curr Biol* **24**, 333-339, doi:10.1016/j.cub.2013.12.041 (2014).

Atasoy, S. *et al.* Connectome-harmonic decomposition of human brain activity reveals dynamical repertoire re-organization under LSD. *Sci Rep* **7**, 17661, doi:10.1038/s41598-017-17546-0 (2017).

Varley, T. F., Carhart-Harris, R., Roseman, L., Menon, D. K. & Stamatakis, E. A. Serotonergic psychedelics LSD & psilocybin increase the fractal dimension of cortical brain activity in spatial and temporal domains. *Neuroimage* **220**, 117049, doi:10.1016/j.neuroimage.2020.117049 (2020).

Chialvo, D. R. Emergent complex neural dynamics. *Nature Physics* **6**, 744-750 (2010).

Cocchi, L., Gollo, L. L., Zalesky, A. & Breakspear, M. Criticality in the brain: A synthesis of neurobiology, models and cognition. *Prog Neurobiol* **158**, 132-152, doi:10.1016/j.pneurobio.2017.07.002 (2017).

Tian, Y., Tan, Z., Hou, H., Li, G., Cheng, A., Qiu, Y., ... & Sun, P. Theoretical foundations of studying criticality in the brain. *Network Neuroscience* **6(4)**, 1148-1185 (2022).

Kaelen, M. *et al.* LSD enhances the emotional response to music. *Psychopharmacology (Berl)* **232**, 3607-3614, doi:10.1007/s00213-015-4014-y (2015).

Kaelen, M. *et al.* LSD modulates music-induced imagery via changes in parahippocampal connectivity. *Eur Neuropsychopharmacol* **26**, 1099-1109, doi:10.1016/j.euroneuro.2016.03.018 (2016).

Luppi, A. I. *et al.* A role for the serotonin 2A receptor in the expansion and functioning of human transmodal cortex. *Brain*, doi:10.1093/brain/awad311 (2023).

Roseman, L., Leech, R., Feilding, A., Nutt, D. J. & Carhart-Harris, R. L. The effects of psilocybin and MDMA on between-network resting state functional connectivity in healthy volunteers. *Front Hum Neurosci* **8**, 204, doi:10.3389/fnhum.2014.00204 (2014).

Petri, G. *et al.* Homological scaffolds of brain functional networks. *J R Soc Interface* **11**, 20140873, doi:10.1098/rsif.2014.0873 (2014).

Rocha, R. P. *et al.* Recovery of neural dynamics criticality in personalized whole-brain models of stroke. *Nat Commun* **13**, 3683, doi:10.1038/s41467-022-30892-6 (2022).

Aday, J. S., Mitzkovitz, C. M., Bloesch, E. K., Davoli, C. C. & Davis, A. K. Long-term effects of psychedelic drugs: A systematic review. *Neurosci Biobehav Rev* **113**, 179-189, doi:10.1016/j.neubiorev.2020.03.017 (2020).

Cordes, D. *et al.* Frequencies contributing to functional connectivity in the cerebral cortex in "resting-state" data. *AJNR Am J Neuroradiol* **22**, 1326-1333 (2001).

Liu, J., Duffy, B. A., Bernal-Casas, D., Fang, Z. & Lee, J. H. Comparison of fMRI analysis methods for heterogeneous BOLD responses in block design studies. *Neuroimage* **147**, 390-408, doi:10.1016/j.neuroimage.2016.12.045 (2017).

Chaudhary, U. J., Duncan, J. S. & Lemieux, L. Mapping hemodynamic correlates of seizures using fMRI: A review. *Hum Brain Mapp* **34**, 447-466, doi:10.1002/hbm.21448 (2013).

Duyn, J. H. EEG-fMRI Methods for the Study of Brain Networks during Sleep. *Front Neurol* **3**, 100, doi:10.3389/fneur.2012.00100 (2012).

Brechet, L. *et al.* Capturing the spatiotemporal dynamics of self-generated, task-initiated thoughts with EEG and fMRI. *Neuroimage* **194**, 82-92, doi:10.1016/j.neuroimage.2019.03.029 (2019).

Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* **59**, 2142-2154, doi:10.1016/j.neuroimage.2011.10.018 (2012).

Tzourio-Mazoyer, N. *et al.* Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* **15**, 273-289, doi:10.1006/nimg.2001.0978 (2002).

Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional organization of the human neocortex. *Nat Neurosci* **25**, 1569-1581, doi:10.1038/s41593-022-01186-3 (2022).

Acknowledgments

E.T. is supported by PICT-2019–02294 (Agencia I+D+i, Argentina), ANID/FONDECYT Regular 1220995 (Chile) and by PIP 1122021010 0800CO (2022-2024) (CONICET, Argentina). The collection of the empirical data used in this manuscript was funded via a donation by Patrick Vernon, mediated by the Beckley Foundation. C.T. is funded by a donation by Anton Bilton to the Centre for Psychedelic Research.

Author Contributions

J.I.P, Y.S-P, C.P and E.T designed the research. J.I.P conducted the research. J.I.P, Y.S-P and E.T analyzed and interpreted the results. J.I.P created the code and made the figures. D.N, R.C-H and C.T provided the curated data. J.I.P and E.T wrote the manuscript. C.P provided analytic support. Y.S-P and E.T supervised the research. All authors edited the manuscript.

Competing interests

The authors declare not competing interest.

Data and materials availability

All data and code needed to evaluate the conclusions in this paper are available at https://github.com/juanpiccinini/DMT-whole-brain.