Scarcity of fixed carbon transfer in a model microbial phototroph-heterotroph interaction

Short Title: Scarcity of microbial carbon transfer

Sunnyjoy Dupuis[1,2], Usha F. Lingappa[2], Xavier Mayali[3], Eve S. Sindermann[2], Jordan L.

Chastain[2,4], Peter K. Weber[3], Rhona Stuart[3], Sabeeha S. Merchant[1,2,5,6]

[1]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720,

USA

[2]California Institute for Quantitative Biosciences (QB3), University of California, Berkeley, CA

94720, USA

[3]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore,

CA 94550, USA

[4]College of Chemistry, University of California, Berkeley, CA 94720, USA

[5]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA

[6]Division of Environmental Genomics and Systems Biology, Lawrence Berkeley National

Laboratory, Berkeley, CA, USA

Correspondence: Sabeeha S. Merchant, sabeeha@berkeley.edu, +1 510 664-5154,

408B Stanley Hall, Berkeley, CA 94720-3220

Study Funding: This work was supported by The Gordon and Betty Moore Foundation

Symbiosis in Aquatic Systems Initiative Investigator Award GBML9203

(https://doi.org/10.37807/GBMF9203), and the Lawrence Livermore National Laboratory

μBiospheres Scientific Focus Area award SCW1039 from the DOE OBER Genomic Sciences

Program. S.D. acknowledges support from NIH T32 Genetic Dissection of Cells and Organisms

training grant 1T32GM132022-01.

ABSTRACT

Although the green alga *Chlamydomonas reinhardtii* has long served as a reference organism, few studies have interrogated its role as a primary producer in microbial interactions. Here, we quantitatively investigated *C. reinhardtii's* capacity to support a heterotrophic microbe using the established coculture system with *Mesorhizobium japonicum*, a vitamin B_{12}-producing α-proteobacterium. Using stable isotope probing and nanoscale secondary ion mass spectrometry (nanoSIMS), we tracked the flow of photosynthetic fixed carbon and consequent bacterial biomass synthesis under continuous and diurnal light with single-cell resolution. We found that more ^{13}C fixed by the alga was taken up by bacterial cells under continuous light, invalidating the hypothesis that the alga's fermentative degradation of starch reserves during the night would boost *M. japonicum* heterotrophy. $^{15}NH_4$ assimilation rates and changes in cell size revealed that *M. japonicum* cells reduced new biomass synthesis in coculture with the alga but continued to divide – a hallmark of nutrient limitation often referred to as reductive division. Despite this sign of starvation, the bacterium still synthesized vitamin B_{12} and supported the growth of a B_{12}-dependent *C. reinhardtii* mutant. Finally, we showed that bacterial proliferation could be supported solely by the algal lysis that occurred in coculture, highlighting the role of necromass in carbon cycling. Collectively, these results reveal the scarcity of fixed carbon in this microbial trophic relationship (particularly under environmentally relevant light regimes), demonstrate B_{12} exchange even during bacterial starvation, and underscore the importance of quantitative approaches for assessing metabolic coupling in algal-bacterial interactions.

KEY WORDS

rhizobia; chlorophyte; photoautotroph; diel; cobalamin; co-cultivation; symbiosis; SIP

INTRODUCTION

Photosynthetic microbes are responsible for a large fraction of the primary productivity

on our planet [1]. These organisms supply the microbial world with raw materials for life,

thereby shaping communities and driving carbon cycling [2, 3]. Through exudation, lysis, and

predation, photosynthetic microbes provide a variety of reduced carbon substrates with various

diffusion rates and accessibility to heterotrophic bacteria. The primary consumers recycle carbon

back to inorganic forms, closing the "microbial loop" that constitutes a major component of

biogeochemical carbon cycling [2, 4, 5]. Heterotrophic bacteria can also influence algal

productivity, for example, by changing the availability of critical micronutrients for algal growth

and metabolism [6–8].

The green alga *Chlamydomonas reinhardtii* has served as a reference organism for

discoveries in photosynthesis, cilia-based motility, and the eukaryotic cell cycle [9]. Extensive

work on axenic (bacteria-free) cultures has provided detailed knowledge of the pathways and

regulons for photosynthesis and primary metabolism. This chlorophyte has a flexible metabolic

repertoire, with the capacity to grow by photoautotrophy alone or by acetate assimilation. Under

diurnal cycles, *C. reinhardtii*'s metabolism is shaped by programmed waves of gene expression

that accommodate diurnal changes in light and oxygen availability in natural environments [10,

11]. Photosynthesis fuels cell growth and starch synthesis during the day, and starch reserves are

glycolytically degraded at night, providing a source of energy until the next dawn. Although

extensive knowledge of its metabolism makes *C. reinhardtii* an attractive model organism to

study microbial trophic interactions, relatively little is known about how *C. reinhardtii*

contributes to its environment and influences its neighbors in the microbial world.

In its natural soil context, *C. reinhardtii* is surrounded by other microbes that exchange

nutrients, compete for resources, and communicate with one another. *C. reinhardtii* can enrich

for particular heterotrophic bacteria from bulk soil, demonstrating that the alga's growth can

influence the metabolisms and behaviors of various bacteria [12, 13]. Studies of

Chlamydomonas-bacteria interactions have primarily focused on impacts to the alga's behavior,

in part due to technical challenges in probing bacterial physiology in mixed cultures. Bacteria are

known to supply vitamins and fixed nitrogen, facilitate iron assimilation, and impart thermal

tolerance to algal cultures [6, 7, 14–20]. Coculture of *C. reinhardtii* with various heterotrophic

bacteria (e.g. *Bradyrhizobium japonicum*, *Pseudomonas* spp.) has been shown to stimulate algal

H_2 production, starch accumulation, and viability under sulfur deprivation by consuming O_2 and

alleviating accumulation of fermentation byproducts from the medium [21]. Bacterial secondary

metabolites can also influence *C. reinhardtii*'s cell cycle and aggregation [22, 23]. In addition,

interactions between *C. reinhardtii* and vitamin B_{12}-producing rhizobia like *Mesorhizobium*

japonicum (previously referred to as *Mesorhizobium loti*), *Sinorhizobium meliloti*, and

Rhizobium leguminosarum have been used to study the evolution of vitamin dependency across

the algal tree of life [6, 15, 24–28]. *C. reinhardtii* does not require vitamin B_{12} for growth,

because it has retained a functional cobalamin-independent isoform of methionine synthase,

METE1. However, when vitamin B_{12} is supplied, *METE1* is repressed, and the vitamin is used as

a cofactor for the cobalamin-dependent isoform, METH1. When *C. reinhardtii* was grown in the

laboratory with a continuous supply of the vitamin, the alga acquired deleterious mutations in the

repressed *METE1* gene, yielding B_{12}-requiring strains (*mete1*) in less than 500 generations [25].

The interaction between the *C. reinhardtii mete1* mutant and *M. japonicum* is the most

well-studied *C. reinhardtii* coculture system. It has been suggested to be facultatively

mutualistic: the mutant alga receives the vitamin cofactor, and the bacterium is believed to

receive reduced carbon compounds that support heterotrophic growth. However, the mechanism

and magnitude of the flux of carbon transfer from *C. reinhardtii* to *M. japonicum* are unknown.

Quantitative investigations of algal-bacterial interactions have been challenging due to technical

barriers for studying eukaryotes and prokaryotes simultaneously. We have addressed this

challenge by using spatiotemporal stable isotope tracing via nanoscale secondary ion mass

spectrometry (nanoSIMS) to probe this simple model interaction – a technique which has

typically been used for studying complex polycultures from the field. We quantified the transfer

of fixed carbon from wild-type *C. reinhardtii* to *M. japonicum* in continuous and diurnal light.

Although some photosynthetically fixed carbon was taken up by the bacterial population, *M.*

japonicum growth was severely limited, resulting in reductive bacterial division. Thus, *C.*

reinhardtii primary productivity yields a low carrying capacity for this bacterium, particularly

under diurnal light. Nonetheless, *M. japonicum* provided adequate B_{12} to enable growth of the *C.*

reinhardtii mete1 strain even when the bacterium itself exhibited signs of starvation. These

results highlight that in natural contexts, bacteria may not proliferate to the high cell densities

that are possible in nutrient-rich media often used in the laboratory, but they can still greatly

influence the metabolism and evolution of their neighbors in the microbial world.

MATERIALS AND METHODS

Strains and culture conditions

Three strains of *C. reinhardtii* were used in this study. The *mete1* strain generated

through experimental evolution on 1000 ng/l vitamin B_{12} [25] and its parental wild-type strain

were provided by Alison G. Smith (University of Cambridge, UK). According to its haplotype,

this strain is closely related to strain S24- (SI Fig. 1) [29]. For experiments testing the impact of

diurnal light, we used the cell-wall reduced strain CC-5390. This strain is amenable to

synchronization under diurnal cycles, providing a system with exceptional signal-to-noise for

resolving diurnal patterns [10]. *M. japonicum* strain MAFF303099 was also provided by Alison

G. Smith.

All experiments were performed in amended HSM minimal medium [30] with a modified

trace element solution [31], 2 mM $MgSO_4 \cdot 7H_2O$, 10 µg/l biotin, and 10 µM $CoCl_2$ unless

otherwise specified (Supplementary Data File S2: Media composition). Cultures were grown in

either 250 ml Erlenmeyer flasks containing 100 ml medium and bubbled with filter-sterilized air

(provided by an aquarium pump), 6-well plates with 3 ml medium, or 250 ml beakers with 50 ml

medium. Algal cultures and cocultures were incubated at 28°C, agitated at 110–130 rpm, and

illuminated with 200 µmol photons/m^2/s cool white light (Sylvania Dulux L55WFT55DL/841)

continuously or under diurnal 12-h-light/12-h-dark cycles.

C. reinhardtii cells were precultivated axenically in photoautotrophic conditions for at

least 3 passages prior to experiments. The *mete1* strain was precultivated with 200–250 ng/l

vitamin B_{12} (cyanocobalamin). For experiments comparing coculture in continuous light versus

diurnal light, *C. reinhardtii* CC-5390 cells were precultivated in flat-panel photobioreactors

(Photon System Instruments, Drásov, Czechia) as previously described under the respective light

regime to achieve adequate synchrony of the diurnal populations [10].

Bacterial cells were precultivated in 14 ml polystyrene round-bottom tubes with 3 ml

amended HSM containing 0.2% sucrose, agitated at 200 rpm, and incubated at 28°C. When

larger volumes were required for inoculation, cells were precultivated instead in 250 ml

Erlenmeyer flasks containing 100 ml medium, continuously bubbled with air, and agitated at

110–130 rpm. Bacterial cells were collected by centrifugation at 8000 x*g* and washed three times

in 0.85% NaCl to remove sucrose before use as inoculum. Optical density at 600 nm (OD_{600}) was

used to estimate the cell density of washed cell suspensions used for inoculation.

Cell number, size, and cytotoxicity measurements

C. reinhardtii cell density and size were determined using a hemocytometer and ImageJ

or a Beckman Coulter Multisizer 3 with a 50 μm orifice (Beckman Coulter, CA, USA). Bacterial

cell density was determined by counting colony forming units (CFU) in 10 μl spots of serial

dilutions on Typtone Yeast (TY) agar medium (Supplementary Data File S2: Media

composition) after 4 d of incubation at 30°C in the dark.

Algal cellular integrity was estimated using CellTox Green Cytotoxicity Assay (Promega,

WI, USA). *C. reinhardtii* sample densities were adjusted to $1\text{x}10^6$ cells/ml. Triplicate killed

control samples were prepared by heating cells at 90°C for 10 min. Fluorescence was measured

in a black-walled 96-well plate using a SpectraMax iD3 plate reader (Molecular Devices, CA,

USA). Background fluorescence of a medium blank was subtracted from all values. Results were

reported relative to killed control samples.

Samples for fluorescence microscopy were fixed overnight in 4% paraformaldehyde at

4°C in the dark. Fixed samples were then collected by centrifugation at 10,000 xg, washed in

phosphate buffered saline (PBS) (pH 7), and stored in 1:1 ethanol:PBS at -20°C. For imaging,

samples were spotted on 0.22 μm GTTP type isopore polycarbonate filters and washed with

sterile water to remove residual salts. Prepared filters were mounted on slides with 5 μg/ml

DAPI in Citifluor antifade mounting medium buffered with PBS and imaged using a Zeiss

Axioimager M2 microscope with a DAPI filter set (Ex 350/50, Em 460/50). Bacterial cell

lengths were measured manually using ImageJ Version 2.1.0/1.53c by drawing a line from one

pole of the cell to the other along the longer axis.

Total organic carbon measurements

Total nonpurgeable organic carbon content (NPOC) of cells was determined using a

TOC-L Shimadzu Total Organic Carbon Analyzer (Shimadzu, Kyoto, Japan). 10–40 ml culture

was collected by centrifugation at 8,000 xg for 2 min, and the supernatant was collected and then

stored at -20°C until analysis. Spent media were diluted 2-fold and HCl was added to 27 mM.

All samples were sparged to remove inorganic carbon. Some organic carbon may be purged from

the sample by this method, so we report "nonpurgeable" organic carbon.

Stable isotope labelling and nanoSIMS

$^{15}NH_4Cl$ (99% ^{15}N, Cambridge Isotope Laboratories, MA, USA) was provided directly in

the culture medium, replacing 50% of the NH_4Cl in HSM. 13 h after inoculation, $^{13}CO_2$ was

provided to cultures by bubbling air through a solution of $NaH^{13}CO_3$ (99% ^{13}C, Cambridge

Isotope Laboratories) and then into the cultures as previously described [32]. Killed controls

were generated by treating cells with 4% paraformaldehyde for 20 min in the dark. Coculture

experiments were otherwise conducted as described above.

For nanoSIMS analyses, samples were fixed and deposited on filters as described above

for fluorescence microscopy. Dry filter pieces were then mounted on 1 inch round stubs using

carbon tabs (Ted Pella, Inc., CA, USA) and gold coated. Data were collected on a NanoSIMS 50

(CAMECA, Gennevilliers, France) using a 1.5 pA Cs^+ primary ion beam. The secondary ion

masses of $^{12}C_2^-$, $^{12}C^{13}C^-$, $^{12}C^{14}N^-$, $^{12}C^{15}N^-$, and $^{32}S^-$ were collected. Raw nanoSIMS images were

processed using L'image (L. Nittler, Carnegie Institution of Washington, D.C., USA). Individual

cell regions of interest (ROIs) were manually circled using size and high $^{12}C^{14}N$ signal to

identify bacterial cells and high $^{12}C^{13}C$ signal along with the secondary electron image to identify

algal cells. Isotope enrichment in atom percent enrichment (APE) for $^{12}C^{13}C^-/^{12}C_2^-$ and $^{12}C^{15}N^-$

$/^{12}C^{14}N^-$ were calculated using standard ratios of 0.02247 and 0.00367, respectively. Enrichment

data are also presented as net assimilation (N_{net} and C_{net}). The fraction of newly synthesized

biomass relative to total biomass was calculated against an unlabeled biomass standard as

previously described [33]:

$$X_{net} = \frac{F_{cell} - F_{unlabelled}}{F_{substrate} - F_{unlabelled}}$$

Where F is the abundance of the heavy isotope (^{15}N or ^{13}C) over the total ($^{14}N+^{15}N$ or $^{12}C+^{13}C$,

respectively). For ^{15}N, net assimilation was calculated relative to the known isotopic composition

of the initial source substrate added to the medium. For ^{13}C, net bacterial assimilation was

calculated relative to the average algal biomass isotopic composition in the respective

experiment (continuous light or diurnal light) as a rough estimation of the source substrate.

Vitamin B$_{12}$ bioassay

Vitamin B$_{12}$ concentration was determined using the *Escherichia coli* B$_{12}$ bioassay

described previously [34, 35]. The bioassay uses two *E. coli* strains: the Δ*metE*Δ*metH* control

strain requires methionine for growth, and the Δ*metE* strain can grow when provided either

methionine or B$_{12}$. The amount of B$_{12}$ in a sample is determined by subtracting the growth of the

Δ*metE*Δ*metH* control strain from the growth of the Δ*metE* strain after 24 h incubation in the

sample and then comparing to a vitamin B$_{12}$ standard curve.

Culture samples and vitamin B$_{12}$ standards were boiled at 100°C for 10 min. Cell debris

was removed by centrifugation at 16,000 x*g* for 2 min, and the supernatants were snap frozen in

liquid N$_2$ and stored at -80°C until analysis. *E. coli* strains were precultivated twice in

polystyrene round-bottom tubes in 2 ml M9 medium (Supplementary Data File S2: Media

composition) with 0.2% glucose and 1 mg/ml methionine at 37°C and 250 rpm agitation for 24

h. *E. coli* inocula were then collected by centrifugation at 10,000 x*g* for 1 min and washed three

times with M9 medium prior to use in the bioassay. Samples and standards were thawed at room

temperature and diluted in 2X M9 medium with 0.4% glucose and water to a final volume of 2

ml in polystyrene round-bottom tubes, to which either the $\Delta metE$ strain or the $\Delta metE \Delta metH$

control strain was added to a starting density of 0.01 OD_{600}. The bioassay was then incubated at

37°C and agitated at 250 rpm for 24 h, and growth was measured by OD_{600}. Sample B_{12}

concentration was determined relative to a standard curve.

Spent medium and cell lysate experiments

Wild-type *C. reinhardtii* was grown for 48 h in continuous light until density reached 1–

4×10^6 cells/ml. Cells were collected by centrifugation in 500 ml polypropylene bottles for 4 min

at 11,900 xg and 4°C. The supernatant was passed through a 0.22 μm filter to generate sterile

spent medium. Cell pellets were resuspended in fresh medium to the original culture volume and

then sonicated using a Fisherbrand Model 505 Sonic Dismembrator (Fisher Scientific, PA, USA)

on ice for 45 min with alternating 10 s at 30% intensity and 15 s "off" steps. The supernatant of

the resulting cell lysate was then passed through a 0.22 μm filter to generate sterile soluble cell

lysate. Spent medium, soluble cell lysate, and parallel live *C. reinhardtii* cultures were then

inoculated with *M. japonicum* as described above, and growth was monitored over another 48 h.

To determine the bacterial growth supported by substrates in *C. reinhardtii* cell lysate

when these substrates are present at a relevant concentration, wild-type *C. reinhardtii* was grown

for 48 h in continuous light, and spent medium and soluble cell lysate were prepared as described

above. Then, the NPOC of the two culture fractions was measured, and the cell lysate was

diluted to the same NPOC concentration as the spent medium using fresh medium. 3 ml of the

diluted cell lysate was inoculated with *M. japonicum* as described above in parallel with spent

medium, fresh medium, and fresh medium with 150 μg/ml sucrose in polystyrene round-bottom

tubes. Cultures were incubated at 30°C and agitated at 200 rpm, and growth was monitored for

roughly 48 h.

RESULTS

***M. japonicum* growth is severely limited in coculture with *C. reinhardtii*, especially under**

diurnal light

Previous work suggested that *C. reinhardtii* supports heterotrophic growth of the

rhizobial bacterium *M. japonicum* when cocultured in minimal medium given light and

atmospheric CO_2 [25, 26, 28, 36]. As a first step in characterizing the transfer of fixed carbon in

this relationship, we compared the growth of *M. japonicum* when incubated alone or together

with wild-type *C. reinhardtii* in minimal medium with either no exogenous carbon source or

with 150 µg/ml sucrose (Fig. 1). Coculture with the bacterium did not have a discernable impact

on *C. reinhardtii* growth or physiology (Fig. 1B, SI Fig. 2, $p > 0.05$). *M. japonicum* grew

considerably with sucrose and only marginally without it, but we found that coculture with *C.*

reinhardtii did not significantly increase the maximum bacterial density in either condition (Fig.

1C, $p > 0.05$). Although the maximum density of *M. japonicum* in coculture without sucrose

($\sim 3 \times 10^7$–5×10^8 CFU/ml) was similar to what has been observed previously [28], we found that

the bacterium was able to achieve a comparable density without the alga, in contrast to previous

reports.

As the growth of *M. japonicum* was not significantly stimulated by wild-type *C.*

reinhardtii, it seemed that the alga did not supply adequate substrates for the bacterium's

heterotrophic metabolism under these conditions. This could be because *C. reinhardtii* released

fixed carbon compounds in insufficient quantities, or because *M. japonicum* is unable to

metabolize the specific compounds generated by *C. reinhardtii*. We had initially illuminated

cocultures with continuous light to maximize algal carbon fixation and growth rate, but we

hypothesized that the alga's nighttime metabolism could stimulate additional growth of *M.*

japonicum. During the day, the alga fixes carbon to build biomass and stores photosynthate as

starch [10]. After sufficient growth during the light phase, cells initiate DNA synthesis and

mitosis in the dark, followed by a G0 phase during which starch reserves are degraded by

glycolysis and genes for fermentation are expressed. Fermentation "waste products" released by

algae in the night could serve as a readily available source of fixed carbon for cooccurring

bacteria.

To test the impact of different light regimes on the interaction, we used the readily

synchronized *C. reinhardtii* strain CC-5390. Synchronized populations allow excellent temporal

separation of cells in distinct metabolic phases of the cell cycle, and could reveal a diurnal

pattern in bacterial growth resulting from the algal partner's diurnal metabolism. We compared

cocultures and monocultures under continuous and diurnal light over several days with high

temporal resolution. *C. reinhardtii*'s doubling time is slower under diurnal light (cells only

divide once per day), but we designed the experiment to achieve the same final amount of algal

biomass under the two light regimes. *C. reinhardtii* strain CC-5390 density reached roughly

4×10^6 cells/ml under all conditions (Fig. 2A, 2B). This was roughly 2X higher than the density

reached by the wild-type strain used in Fig. 1, although the strains appeared to reach stationary

phase in the same amount of time under continuous light (72 h). Algal growth was again

unaffected by the bacterium. In contrast, we found that *M. japonicum* density increased

significantly in the presence of CC-5390 under both light regimes (Fig. 2C–2E). CC-5390 may

support more growth of the bacterium than the wild-type strain because it is more prone to cell

lysis and releases more nonpurgeable organic carbon (NPOC) into the spent medium (SI Fig. 3).

However, we could not discern a clear temporal pattern in bacterial proliferation over the diurnal

cycles, and contrary to our hypothesis, we found that continuously illuminated cocultures

supported 3-fold more bacterial cells than did diurnally illuminated cocultures (Fig. 2E, $p =$

0.04). Spent medium from continuous light-grown *C. reinhardtii* had significantly more NPOC

than spent medium from diurnal light-grown *C. reinhardtii* of the same culture density,

suggesting that more dissolved organic carbon was available to the bacterium under continuous

light (Fig. 2F, $p < 0.05$). Algal cell lysis was similar under the two light regimes, so differences

in algal spent medium NPOC suggest that *C. reinhardtii* exudes more reduced carbon in

continuous light than in diurnal light (SI Fig. 4, $p > 0.05$). Finally, the presence of *M. japonicum*

did not significantly reduce the concentration of NPOC in the spent medium (Fig. 2F, Student's

t-test, $p > 0.05$). This could mean that either *M. japonicum* did not take up significant amounts of

organic carbon from the medium or that it could not fully oxidize it to CO_2, and instead released

reduced carbon waste products back into the extracellular milieu.

Stable isotope probing reveals mismatch between bacterial biomass synthesis and division

in coculture

To determine the amount of carbon fixed by *C. reinhardtii* that is taken up by *M.*

japonicum, we conducted stable isotope probing experiments using $^{13}CO_2$. Cocultures of CC-

5390 and *M. japonicum* grown under continuous and diurnal light as in Fig. 2 were bubbled with

air containing $^{13}CO_2$. $^{15}NH_4Cl$ was added to the medium as an independent tracer of biomass

synthesis, complementing measurements of cell density. We assayed label uptake with single-

cell resolution using nanoSIMS.

We found substantial ^{13}C enrichment of the algal cells, which remained at a similar

concentration throughout the timepoints examined (Fig. 3A, SI Fig. 5, SI Fig. 6). In addition, we

observed ^{13}C enrichment of bacterial cells, which increased with time in cocultures under both

light regimes (Fig. 3A, SI Fig. 5, SI Fig. 6, $p < 0.05$). Killed controls and monocultures of *M.*

japonicum on various substrates demonstrated that this enrichment was not a result of

nonspecific background or direct CO_2 incorporation by the bacterium (i.e. through anapleurotic

carboxylation reactions) (SI Fig. 6, SI Fig. 7). Thus, a meaningful amount of carbon derived

from algal photoautotrophy was in fact incorporated into *M. japonicum* biomass in the

cocultures.

We compared the extent of carbon transfer under continuous and diurnal light by

calculating the fraction of bacterial carbon that was derived from algal carbon, C_{net}, using the

average ^{13}C content of algal biomass over the course of each experiment as a rough measure of

the source carbon isotopic composition. We found that significantly more algal ^{13}C was

transferred to bacterial cells under continuous light over 48 h (0.17 median C_{net} in continuous

light, 0.09 median C_{net} in diurnal light) (Fig. 3B, $p = 6 \times 10^{-6}$). This was consistent with our

observation that spent medium NPOC and *M. japonicum* cell density were greater in cocultures

exposed to continuous light than those exposed to diurnal light (Fig. 2, SI Fig. 8A).

Both the ^{13}C and ^{15}N enrichment in *M. japonicum* cells were lower than we had expected

from the increase in bacterial CFUs observed in the experiment (>3-fold increase in CFUs during

the first 24 h of coculture under continuous light), suggesting a mismatch between the change in

cell density and the bacterial biomass synthesized (Fig. 3C, SI Fig. 8A). We quantified the

magnitude of this mismatch using the ^{15}N label, for which the source isotopic composition was

better constrained, as opposed to that of the ^{13}C label which was delivered in the gas phase in an

open system. Using the source ^{15}N composition and the measured isotopic composition of

bacterial cells at each timepoint, we calculated the fraction of newly synthesized biomass

generated from the source substrate over the course of the experiment: N_{net} [33]. We compared

this to the expected N_{net} that would be observed if each doubling in CFUs in the culture occurred

upon a doubling in bacterial biomass (0.99 N_{net} for cells grown on sucrose after 8–13 doublings,

0.63–0.97 N_{net} for cells co-cultivated with *C. reinhardtii* after 1–5 doublings) (SI Fig. 8B).

Indeed, the measured N_{net} of *M. japonicum* cells in the cocultures was much lower than the

expected values at all measured timepoints (0.18–0.34 N_{net}), whereas the measured values

matched the expected values during logarithmic-phase growth on sucrose (Fig. 4A). This showed

that in coculture with *C. reinhardtii*, *M. japonicum* cells reduced new biomass synthesis by 24 h

and yet continued to divide – a hallmark of bacterial starvation often referred to as reductive

division [37–39]. As nutrients available for a bacterium's growth decrease, the cell volume

added prior to cell division is reduced, leading to smaller daughter cells [40–43].

If the deviation of the observed N_{net} from the expected N_{net} indeed reflected reductive

division, bacterial cells in the coculture would be smaller than cells in the logarithmic phase of

growth on sucrose, which are typically 2 μm in length. When we measured the size of the

bacterial cells, we found that *M. japonicum* cells were indeed smaller in the cocultures: mean cell

length was 1.4–1.5 μm after 24 h in coculture, whereas it was 2.3 μm after 24 h of growth on

sucrose (Fig. 4B, 4C). When sucrose was presumably depleted from the medium over the

subsequent day in the monoculture, mean cell length decreased to a similar extent, demonstrating

that reductive division occurs upon carbon limitation in *M. japonicum*.

Taken together, these data show that photosynthetically fixed carbon was indeed

transferred from *C. reinhardtii* to *M. japonicum* under both continuous and diurnal light, and that

although this amount was not sufficient for balanced logarithmic-phase growth of the bacterium,

it allowed for a small amount of bacterial proliferation.

Reductively dividing bacteria still deliver sufficient vitamin B$_{12}$ to support growth of the

B$_{12}$-requiring *mete1* strain

As reductive division had not previously been reported in the *C. reinhardtii-M.*

japonicum interaction, we wondered whether the phenomenon also occurs during coculture with

the B$_{12}$-requiring *mete1* strain of *C. reinhardtii*, and if so, whether the bacterium still synthesizes

adequate B$_{12}$ to support the mutant's growth. We compared the change in bacterial cell size after

growth on 150 µg/ml sucrose to changes during coculture with either wild-type *C. reinhardtii* or

mete1. In all conditions, we observed a 37–50% reduction in *M. japonicum* cell length over 54 h

of cultivation (Fig. 5A, 5B). This suggested that bacterial carbon limitation occurs to a similar

extent during coculture with the *mete1* strain as with the wild type.

To test whether reductively dividing *M. japonicum* still delivers vitamin B$_{12}$ to *C.*

reinhardtii, we compared the growth of the *C. reinhardtii mete1* strain in coculture and

monoculture without exogenous B$_{12}$, and we monitored changes in vitamin B$_{12}$ concentration

overtime using a bioassay [34, 35]. *mete1* growth was limited by B$_{12}$ relative to the wild-type *C.*

reinhardtii, but the B$_{12}$-requiring strain grew significantly more in the presence of *M. japonicum*

despite reductive bacterial division (Fig. 5C, 5E, $p < 0.05$). Furthermore, we found that *M.*

japonicum produced significant vitamin B$_{12}$ in coculture, but only after 24 h (Fig. 5F, $p < 0.05$),

even though bacterial biomass synthesis and cell size have decreased substantially by that time

(Fig. 4). This demonstrates that bacterial B$_{12}$ synthesis and sharing occur despite the carbon

limitation experienced during the trophic interaction with the alga.

Although there were roughly 10X more *C. reinhardtii* cells present in the wild-type

coculture than the B$_{12}$-limited mutant coculture, there was no difference in *M. japonicum* density

between these two conditions (Fig. 5D, $p > 0.05$). Thus, greater algal density does not translate

into greater *M. japonicum* growth, as noted previously [28]. We observed this phenomenon in

cocultures inoculated at several starting relative abundances (SI Fig. 10). This suggests that

carbon transfer from the alga to the bacterium could be indirect.

***C. reinhardtii* lysis can explain observed *M. japonicum* proliferation in coculture**

To distinguish the source of the fixed carbon that supports *M. japonicum* proliferation in

coculture, we compared *M. japonicum* growth with live *C. reinhardtii* to growth in *C. reinhardtii*

spent medium and in the soluble fraction of *C. reinhardtii* cell lysate. Spent medium supported

the same number of *M. japonicum* cells as did coculture, demonstrating that the observed

increase does not require a direct interaction with the alga (Fig. 6A). Cell lysate from *C.*

reinhardtii supported almost an order of magnitude more *M. japonicum* proliferation. Therefore,

we hypothesized that the fixed carbon transferred in our cocultures could be sourced from a few

lysed *C. reinhardtii* cells rather than from photosynthate exuded by live algal cells.

Having determined the increase in viable *M. japonicum* cells supported by lysate from a

known density of *C. reinhardtii* cells (1.7×10^8 CFU/ml on average after 48 h in lysate from

1.6×10^6 *C. reinhardtii* cells/ml culture), we estimated the theoretical increase in *M. japonicum*

CFUs supported by a single lysed *C. reinhardtii* cell to be 100 ± 43 CFU in 48 h. From that, we

estimated how many *C. reinhardtii* cells would have to lyse in the coculture to enable the *M.*

japonicum proliferation observed: roughly 1.1×10^5 *C. reinhardtii* cells/ml, which corresponds to

1.8% of the *C. reinhardtii* populations in those cocultures. To test the actual degree of algal lysis

in our cultures, we measured algal cellular integrity using the CellTox Green Cytotoxicity Assay.

This assay uses a membrane-impermeable DNA dye, resulting in fluorescence from cells whose

plasma membrane has been compromised or from extracellular DNA [44]. We compared live

culture fluorescence to killed control samples and found that 1–3% of the wild-type *C.*

reinhardtii population had compromised cellular integrity (Fig. 6B). Thus, the fraction of dead or

dying *C. reinhardtii* cells measured by this assay is comparable to that needed to explain the

observed *M. japonicum* proliferation.

Finally, we reasoned that if *M. japonicum* is primarily growing on algal cell lysate, then

cell lysate should support the same amount of bacterial proliferation as spent medium when

present at the same organic carbon content. Thus, we compared *M. japonicum* growth in spent

medium and diluted cell lysate from wild-type *C. reinhardtii* when both were present at roughly

15 µg/ml NPOC. Indeed, algal cell lysate alone supported the same *M. japonicum* growth as

spent medium when present at a relevant concentration: bacterial cell density was not

significantly different between these two conditions at most timepoints examined (Fig. 6C, $p >$

0.05). The slight increase in *M. japonicum* density in the diluted cell lysate condition could be

attributed to increased availability of micronutrients in the fresh medium used to dilute the algal

cell lysate. Taken together, these data are consistent with the interpretation that *M. japonicum*

growth in coculture is enabled by substrates that are released through *C. reinhardtii* cell lysis

rather than by substrates exuded by live *C. reinhardtii* cells.

DISCUSSION

Most studies of symbioses between *C. reinhardtii* and heterotrophic bacteria to date have

focused on impacts on the alga, leaving open the question of how this model phototroph might

influence bacterial growth and physiology. Here, we have investigated *C. reinhardtii's* role as a

primary producer in the established coculture system with *M. japonicum*. We have tied carbon

cycling in the interaction to alterations in bacterial growth and physiology through quantitative,

single-cell analyses. Under continuous light, we found that the alga increased the number of

viable bacterial cells, albeit marginally so. We tested whether nighttime metabolism might

increase the alga's propensity for carbon sharing, but we found that diurnal illumination actually

decreased the productivity of the partnership. Using nanoSIMS to visualize ^{13}C enrichment with

high spatial resolution, we quantitatively demonstrated transfer of carbon fixed by the model alga

to the bacterium. Roughly 17% of the carbon in *M. japonicum* cells was derived from *C.*

reinhardtii over 36 h under continuous light, whereas only ~9% of the *M. japonicum* carbon

came from *C. reinhardtii* over the same amount of time under diurnal light. Bulk measurements

showed that algal cultures of similar densities had more organic carbon in their spent medium

after continuous illumination than after diurnal illumination. Together, these results suggest that

the increased metabolic rate realized under continuous light may be more beneficial for *M.*

japonicum (and potentially other bacteria) than fermentative catabolism of starch presumed to

occur during the night. It is also possible that although *C. reinhardtii* expresses genes for

fermentative metabolism in the night, fermentative metabolites may not actually accumulate to

high enough levels in culture unless cells experience anoxia.

With *C. reinhardtii* as the sole source of fixed carbon, *M. japonicum* exhibited signs of

starvation, even under continuous light, highlighting the meagerness of the microbial trophic

relationship. The quantitative stable isotope probing experiments revealed that during coculture

with *C. reinhardtii*, *M. japonicum* synthesized significantly less biomass than expected given the

increase in viable bacterial cells. By direct measurement, we showed that *M. japonicum* divided

into small daughter cells within 24 h of carbon limitation in coculture, a physiological change

often referred to as reductive division. It is well known that as nutrients become limiting,

bacterial size at birth and the amount of cell volume each bacterium adds prior to division

decrease [40–43]. Studies on freshwater and marine bacteria have documented up to 90%

reduction in cell volume during carbon limitation [45–48]. This phenomenon is often considered

an adaptive strategy by which diverse bacteria survive starvation [37–39, 49]. By increasing cell

number and reducing cell volume, bacteria increase the probability that a member of the

population will encounter sufficient nutrients again and increase their surface area-to-volume

ratio to improve nutrient uptake. Reductively dividing cells also exhibit changes in stress

resistance and macromolecule synthesis. However, we found that despite its apparent starvation,

M. japonicum provided enough vitamin B_{12} to support a B_{12}-dependent *C. reinhardtii* mutant.

Vitamin supply enables growth of auxotrophs, influences gene expression and metabolism, and

can even give rise to vitamin dependency in algae [25, 50]. Thus, our results suggest that even

under austere conditions with meager supply of growth substrates, bacteria can still influence

algal physiology and evolution.

We also found that a direct interaction with *C. reinhardtii* was not required to achieve the

M. japonicum proliferation observed in coculture: growth in algal spent medium yielded the

same maximum density of viable *M. japonicum* cells as did coculture. Furthermore,

mechanically lysed *C. reinhardtii* cells yielded considerably higher *M. japonicum* densities. This

observation led us to hypothesize that perhaps this bacterium does not metabolize the exudates of

live *C. reinhardtii* cells at all. Instead, perhaps the small amount of bacterial growth observed in

coculture and in spent medium is due to the lysis of a small fraction of algal cells. It has long

been appreciated that even during rapid exponential growth, microbial cultures experience a

small death rate [51]. Furthermore, cell lysis is an important mechanism of nutrient transfer in

natural environments, and microbial necromass is a major component of the soil organic carbon

pool [52, 53]. For the experimental conditions examined here, we showed that ~2% of the *C.*

reinhardtii population in our cultures are present as necromass. Based on the *M. japonicum*

growth yields achieved in mechanically lysed *C. reinhardtii* cells, we calculated that this is

almost exactly the amount of algal necromass required to explain the *M. japonicum* proliferation

that occurred in coculture. When present at the same organic carbon concentration, algal cell

lysate supported the same amount of *M. japonicum* growth as algal spent medium. Therefore, we

propose that lysis is the primary mechanism of trophic transfer in this coculture system, rather

than exudation or direct exchange.

Given that very little growth of *M. japonicum* was observed from an interaction with *C.*

reinhardtii – even under conditions optimized for algal productivity – our results may suggest

that this model alga exhibits a low carrying capacity for heterotrophic bacteria in the

environment. Similar bacterial densities have been reported for other *C. reinhardtii* cocultures in

minimal media, including those with *Methylobacterium aquaticum* [54, 55]. However, it is

possible that other bacteria, such as those that may have co-evolved with *C. reinhardtii* in its

native environment, may possess greater capacity to metabolize compounds that *C. reinhardtii*

releases during growth. In addition, other bacteria may achieve greater access to growth

substrates through chemotaxis, interspecies signaling and direct exchange, or algicidal activity

[3, 56–59]. The mode of cell lysis (i.e. mechanical vs. viral) could also greatly impact the algal-

derived nutrients released to cooccurring heterotrophs [60]. Nonetheless, stable isotope probing

experiments on other algal-bacterial interactions and on seawater microbial communities have

reported similar levels of trophic carbon transfer to those we observed here [56, 61, 62]. This

suggests that although *C. reinhardtii* and *M. japonicum* may not commonly encounter one

another in nature, this relationship may be representative of naturally occurring microbial

phototroph-heterotroph interactions. Studying this model system has revealed important aspects

of microbial ecology, and our quantitative approach can serve as a roadmap for the future work

necessary to understand *C. reinhardtii*'s full potential as a primary producer in the microbial

world.

ACKNOWLEDGMENTS

This work was supported by The Gordon and Betty Moore Foundation Symbiosis in Aquatic

Systems Initiative Investigator Award GBML9203 (https://doi.org/10.37807/GBMF9203), and

the Lawrence Livermore National Laboratory μBiospheres Scientific Focus Area award

SCW1039 from the DOE OBER Genomic Sciences Program. S.D. acknowledges support from

NIH T32 Genetic Dissection of Cells and Organisms training grant 1T32GM132022-01. We

thank Dr. Alison G. Smith and her group at the University of Cambridge for providing strains

and for their generous discussions with us. We thank Dr. Michiko E. Taga at UC Berkeley for

her guidance on the work. We thank Dr. Denise Schichnes and the UC Berkeley Biological

Imaging Facility for fluorescence microscopy support. We also thank Christina Ramon at

Lawrence Livermore National Laboratory and Charles Perrino at UC Berkeley for analytical

support.

COMPETING INTERESTS

The authors declare no competing financial interests.

DATA AVAILABILITY

All data underlying this article are available in Supplementary Data File S1.

REFERENCES

1.   Field CB, Behrenfeld MJ, Randerson JT, Falkowski P. Primary production of the

biosphere: Integrating terrestrial and oceanic components. *Science (1979)* 1998; **281**: 237–

240.

2.   Azam F, Fenchel T, Field J, Gray J, Meyer-Reil L, Thingstad F. The ecological role of

water-column microbes in the sea. *Mar Ecol Prog Ser* 1983; **10**: 257–263.

3.   Seymour JR, Amin SA, Raina J-B, Stocker R. Zooming in on the phycosphere: the

506   ecological interface for phytoplankton–bacteria relationships. *Nat Microbiol* 2017; **2**:

507   17065.

508   4.   Pomeroy LR. The ocean's food web, a changing paradigm. *Bioscience* 1974; **24**: 499–504.

509   5.   Azam F, Malfatti F. Microbial structuring of marine ecosystems. *Nat Rev Microbiol* 2007;

510   **5**: 782–791.

511   6.   Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. Algae acquire vitamin

512   B_{12} through a symbiotic relationship with bacteria. *Nature* 2005; **438**: 90–93.

513   7.   Amin SA, Green DH, Hart MC, Küpper FC, Sunda WG, Carrano CJ. Photolysis of iron–

514   siderophore chelates promotes bacterial–algal mutualism. *Proceedings of the National*

515   *Academy of Sciences* 2009; **106**: 17071–17076.

516   8.   Hom EFY, Aiyar P, Schaeme D, Mittag M, Sasso S. A chemical perspective on

517   microalgal-microbial interactions. *Trends Plant Sci* 2015; **20**: 689–693.

518   9.   Salomé PA, Merchant SS. A series of fortunate events: Introducing Chlamydomonas as a

519   reference organism. *Plant Cell* 2019; **31**: 1682–1707.

520   10.   Strenkert D, Schmollinger S, Gallaher SD, Salomé PA, Purvine SO, Nicora CD, et al.

521   Multiomics resolution of molecular events during a day in the life of Chlamydomonas.

522   *Proceedings of the National Academy of Sciences* 2019; **116**: 2374–2383.

523   11.   Zones JM, Blaby IK, Merchant SS, Umen JG. High-resolution profiling of a synchronized

524   diurnal transcriptome from *Chlamydomonas reinhardtii* reveals continuous cell and

525   metabolic differentiation. *Plant Cell* 2015; **27**: 2743–2769.

526   12.   Durán P, Flores-Uribe J, Wippel K, Zhang P, Guan R, Melkonian B, et al. Shared features

527   and reciprocal complementation of the Chlamydomonas and Arabidopsis microbiota. *Nat*

528   *Commun* 2022; **13**: 406.

529   13.   Kim B-H, Ramanan R, Cho D-H, Oh H-M, Kim H-S. Role of Rhizobium, a plant growth

530   promoting bacterium, in enhancing algal biomass through mutualistic interaction. *Biomass*

531   *Bioenergy* 2014; **69**: 95–105.

532   14.   Lőrincz Z, Preininger É, Kósa A, Pónyi T, Nyitrai P, Sarkadi L, et al. Artificial tripartite

533   symbiosis involving a green alga (Chlamydomonas), a bacterium (Azotobacter) and a

534   fungus (Alternaria): Morphological and physiological characterization. *Folia Microbiol*

535   *(Praha)* 2010; **55**: 393–400.

536   15.   Xie B, Bishop S, Stessman D, Wright D, Spalding MH, Halverson LJ. *Chlamydomonas*

537   *reinhardtii* thermal tolerance enhancement mediated by a mutualistic interaction with

538   vitamin B_{12}-producing bacteria. *ISME J* 2013; **7**: 1544–1555.

539   16.   Amin SA, Parker MS, Armbrust EV. Interactions between diatoms and bacteria.

540   *Microbiology and Molecular Biology Reviews* 2012; **76**: 667–684.

541   17.   Gobler C, Norman C, Panzeca C, Taylor G, Sañudo-Wilhelmy S. Effect of B-vitamins

542   (B_1, B_{12}) and inorganic nutrients on algal bloom dynamics in a coastal ecosystem. *Aquatic*

543   *Microbial Ecology* 2007; **49**: 181–194.

544   18.   Sañudo-Wilhelmy SA, Gobler CJ, Okbamichael M, Taylor GT. Regulation of

545   phytoplankton dynamics by vitamin B_{12}. *Geophys Res Lett* 2006; **33**: L04604.

546   19.   Helliwell KE. The roles of B vitamins in phytoplankton nutrition: new perspectives and

547   prospects. *New Phytologist* 2017; **216**: 62–68.

548   20.   Keshtacher-Liebso E, Hadar Y, Chen Y. Oligotrophic bacteria enhance algal growth under

549   iron-deficient conditions. *Appl Environ Microbiol* 1995; **61**: 2439–41.

550   21.   Fakhimi N, Gonzalez-Ballester D, Fernández E, Galván A, Dubini A. Algae-bacteria

551   consortia as a strategy to enhance H_2 production. *Cells* 2020; **9**: 1353.

552   22.   Krespach MKC, Stroe MC, Flak M, Komor AJ, Nietzsche S, Sasso S, et al. Bacterial

553   marginolactones trigger formation of algal gloeocapsoids, protective aggregates on the

554   verge of multicellularity. *Proceedings of the National Academy of Sciences* 2021; **118**.

555   23.   Windler M, Stuart R, Deutzmann JS, Mayali X, Navid A, D'haeseleer P, et al. Bacterial

556   exometabolites influence *Chlamydomonas* cell cycle and double algal productivity. *FEMS*

557   *Microbiol Ecol* 2022; **98**.

558   24.   Kazamia E, Czesnick H, Nguyen TT Van, Croft MT, Sherwood E, Sasso S, et al.

559   Mutualistic interactions between vitamin B_{12}-dependent algae and heterotrophic bacteria

560   exhibit regulation. *Environ Microbiol* 2012; **14**: 1466–1476.

561   25.   Helliwell KE, Collins S, Kazamia E, Purton S, Wheeler GL, Smith AG. Fundamental shift

562   in vitamin B_{12} eco-physiology of a model alga demonstrated by experimental evolution.

563   *ISME J* 2015; **9**: 1446–1455.

564   26.   Bunbury F, Helliwell KE, Mehrshahi P, Davey MP, Salmon DL, Holzer A, et al.

565   Responses of a newly evolved auxotroph of Chlamydomonas to B_{12} deprivation. *Plant*

566   *Physiol* 2020; **183**: 167–178.

567   27.   Helliwell KE, Scaife MA, Sasso S, Araujo APU, Purton S, Smith AG. Unraveling vitamin

568   B_{12} -responsive gene regulation in algae. *Plant Physiol* 2014; **165**: 388–397.

569   28.   Bunbury F, Deery E, Sayer AP, Bhardwaj V, Harrison EL, Warren MJ, et al. Exploring

570   the onset of B_{12}-based mutualisms using a recently evolved *Chlamydomonas* auxotroph

571   and B_{12}-producing bacteria. *Environ Microbiol* 2022; **24**: 3134–3147.

572   29.   Gallaher SD, Fitz-Gibbon ST, Glaesener AG, Pellegrini M, Merchant SS.

573   Chlamydomonas genome resource for laboratory strains reveals a mosaic of sequence

574   variation, identifies true strain histories, and enables strain-specific studies. *Plant Cell*

575   2015; **27**: 2335–2352.

576   30.   Sueoka N. Mitotic replication of deoxynucleic acid in *Chlamydomonas reinhardi*.

577   *Proceedings of the National Academy of Sciences* 1960; **46**: 83–91.

578   31.   Kropat J, Hong-Hermesdorf A, Casero D, Ent P, Castruita M, Pellegrini M, et al. A

579   revised mineral nutrient supplement increases biomass and growth rate in

580   *Chlamydomonas reinhardtii*. *The Plant Journal* 2011; **66**: 770–780.

581   32.   Lingappa UF, Dupuis S, Mayali X, Merchant SS. Isotopically labelled inorganic carbon

582   delivered to algal cultures via bubbler bottle. *protocols.io* 2023.

583   33.   Dekas AE, Parada AE, Mayali X, Fuhrman JA, Wollard J, Weber PK, et al.

584   Characterizing chemoautotrophy and heterotrophy in marine archaea and bacteria with

585   single-cell multi-isotope nanoSIP. *Front Microbiol* 2019; **10**.

586   34.   Mok KC, Hallberg ZF, Taga ME. Purification and detection of vitamin B_{12} analogs. 2022.

587   pp 61–85.

588   35.   Dupuis S, Schmollinger S, Merchant SS. Quantitative detection of vitamin B_{12} in algae by

589   bioassay and ICP-MS/MS. *protocols.io* 2022.

36.   Laeverenz Schlogelhofer H, Peaudecerf FJ, Bunbury F, Whitehouse MJ, Foster RA, Smith

AG, et al. Combining SIMS and mechanistic modelling to reveal nutrient kinetics in an

algal-bacterial mutualism. *PLoS One* 2021; **16**: e0251643.

37.   Thorne SH, Williams HD. Adaptation to nutrient starvation in *Rhizobium leguminosarum*

bv. phaseoli: analysis of survival, stress resistance, and changes in macromolecular

synthesis during entry to and exit from stationary phase. *J Bacteriol* 1997; **179**: 6894–

6901.

38.   Shimaya T, Okura R, Wakamoto Y, Takeuchi KA. Scale invariance of cell size

fluctuations in starving bacteria. *Commun Phys* 2021; **4**: 238.

39.   Bergkessel M, Delavaine L. Diversity in starvation survival strategies and outcomes

among heterotrophic proteobacteria. *Microb Physiol* 2021; **31**: 146–162.

40.   Schaechter M, Maaløe O, Kjeldgaard NO. Dependency on medium and temperature of

cell size and chemical composition during balanced growth of *Salmonella typhimurium*. *J*

Gen Microbiol 1958; **19**: 592–606.

41.   Vadia S, Levin PA. Growth rate and cell size: a re-examination of the growth law. *Curr*

Opin Microbiol 2015; **24**: 96–103.

42.   Campos M, Surovtsev IV, Kato S, Paintdakhi A, Beltran B, Ebmeier SE, et al. A constant

size extension drives bacterial cell size homeostasis. *Cell* 2014; **159**: 1433–1446.

43.   Taheri-Araghi S, Bradde S, Sauls JT, Hill NS, Levin PA, Paulsson J, et al. Cell-size

control and homeostasis in bacteria. *Current Biology* 2015; **25**: 385–391.

44.   Haire TC, Bell C, Cutshaw K, Swiger B, Winkelmann K, Palmer AG. Robust microplate-

based methods for culturing and *in vivo* phenotypic screening of *Chlamydomonas*

reinhardtii. *Front Plant Sci* 2018; **9**.

45. Vrede K, Heldal M, Norland S, Bratbak G. Elemental composition (C, N, P) and cell

volume of exponentially growing and nutrient-limited bacterioplankton. *Appl Environ*

Microbiol 2002; **68**: 2965–71.

46. Troussellier M, Bouvy M, Courties C, Dupuy C. Variation of carbon content among

bacterial species under starvation condition. *Aquatic Microbial Ecology* 1997; **13**: 113–

119.

47. Holmquist L, Kjelleberg S. Changes in viability, respiratory activity and morphology of

the marine Vibrio sp. strain S14 during starvation of individual nutrients and subsequent

recovery. *FEMS Microbiol Ecol* 1993; **12**: 215–223.

48. Mårdén P, Tunlid A, Malmcrona-Friberg K, Odham G, Kjelleberg S. Physiological and

morphological changes during short term starvation of marine bacterial islates. *Arch*

Microbiol 1985; **142**: 326–332.

49. Nyström T, Larsson C, Gustafsson L. Bacterial defense against aging: role of the

Escherichia coli ArcA regulator in gene expression, readjusted energy flux and survival

during stasis. *EMBO J* 1996; **15**: 3219–28.

50. Sokolovskaya OM, Shelton AN, Taga ME. Sharing vitamins: Cobamides unveil microbial

interactions. *Science* 2020; **369**.

51. Kelly CD, Rahn O. The growth rate of individual bacterial cells. *J Bacteriol* 1932; **23**:

147–153.

52. Gude S, Pherribo GJ, Taga ME. Emergence of metabolite provisioning as a by-product of

evolved biological functions. *mSystems* 2020; **5**.

53. Sokol NW, Slessarev E, Marschmann GL, Nicolas A, Blazewicz SJ, Brodie EL, et al. Life

and death in the soil microbiome: how ecological processes influence biogeochemistry.

Nat Rev Microbiol 2022; **20**: 415–430.

54. Ban S, Lin W, Wu F, Luo J. Algal-bacterial cooperation improves algal photolysis-

mediated hydrogen production. *Bioresour Technol* 2018; **251**: 350–357.

55. Calatrava V, Hom EFY, Llamas Á, Fernández E, Galván A. OK, thanks! A new

mutualism between Chlamydomonas and methylobacteria facilitates growth on amino

acids and peptides. *FEMS Microbiol Lett* 2018; **365**.

56. Samo TJ, Kimbrel JA, Nilson DJ, Pett-Ridge J, Weber PK, Mayali X. Attachment

between heterotrophic bacteria and microalgae influences symbiotic microscale

interactions. *Environ Microbiol* 2018; **20**: 4385–4400.

57. Jackrel SL, Yang JW, Schmidt KC, Denef VJ. Host specificity of microbiome assembly

and its fitness effects in phytoplankton. *ISME J* 2021; **15**: 774–788.

58. Aiyar P, Schaeme D, García-Altares M, Carrasco Flores D, Dathe H, Hertweck C, et al.

Antagonistic bacteria disrupt calcium homeostasis and immobilize algal cells. *Nat*

Commun 2017; **8**: 1756.

59. Raina J-B, Giardina M, Brumley DR, Clode PL, Pernice M, Guagliardo P, et al.

Chemotaxis increases metabolic exchanges between marine picophytoplankton and

heterotrophic bacteria. *Nat Microbiol* 2023; **8**: 510–521.

60. Pherribo GJ, Taga ME. Bacteriophage-mediated lysis supports robust growth of amino

acid auxotrophs. *ISME J* 2023; **17**: 1785–1788.

61.   Arandia-Gorostidi N, Weber PK, Alonso-Sáez L, Morán XAG, Mayali X. Elevated

temperature increases carbon and nitrogen fluxes between phytoplankton and

heterotrophic bacteria through physical attachment. *ISME J* 2017; **11**: 641–650.

62.   de-Bashan LE, Mayali X, Bebout BM, Weber PK, Detweiler AM, Hernandez J-P, et al.

Establishment of stable synthetic mutualism without co-evolution between microalgae and

bacteria demonstrated by mutual transfer of metabolites (NanoSIMS isotopic imaging)

and persistent physical association (Fluorescent *in situ* hybridization). *Algal Res* 2016; **15**:

179–186.

FIGURES

Figure 1: Wild-type *C. reinhardtii* does not significantly improve growth of *M. japonicum*.

Triplicate cultures (*n* = 3) were inoculated with or without 150 µg/ml sucrose and maintained under continuous light, bubbled with air, and shaken at 110 rpm. (A) Representative image of cultures 72 h after inoculation. (B) Wild-type *C. reinhardtii* cell density in cultures over time after inoculation. (C) *M. japonicum* cell density in cultures over time after inoculation. The maximum bacterial density in coculture was not significantly different from the maximum bacterial density in monoculture with or without sucrose by two-tailed Student's *t*-tests (*p* > 0.05).

Figure 2: Coculture light regime impacts organic carbon release and *M. japonicum* growth.

Quadruplicate (*n* = 4) monocultures (colors) and cocultures (black and white) were bubbled with air and shaken at 125 rpm. Cultures were inoculated to reach the same final algal cell density at the end of the experiment. (A, B) Growth of cell wall reduced strain CC-5390 in continuous light or diurnal light (12-h-light/12-h-dark), respectively, with and without *M. japonicum*. The grey backgrounds indicate sample timepoints that occurred during the dark phase. Under diurnal light, CC-5390 exhibits synchronous growth, doubling roughly once per day upon nightfall. (C, D) *M. japonicum* growth under the continuous and diurnal light regimes, respectively, with and without CC-5390. (E) Maximum bacterial density achieved under the various trophic conditions tested. (F) NPOC in spent medium from CC-5390 and cocultures sampled at the indicated times from the continuous and diurnal light regimes in (A) and (B), respectively. Lower-case letters indicate

significantly different groups by paired two-tailed Student's *t*-test ($p < 0.05$). Error bars represent

the standard deviation from the mean.

Figure 3: Stable isotope probing demonstrates carbon flux in continuous and diurnal light.
Duplicate cocultures ($n = 2$) of CC-5390 and *M. japonicum* were grown as described in Fig. 2
but with 50% of their NH_4^+ provided as ^{15}N and with $^{13}CO_2$ added to the air starting 13 h after
inoculation. Isotope enrichment was measured using nanoSIMS at the indicated times after
inoculation. The grey backgrounds indicate sample timepoints that occurred during the dark
phase. (A) ^{13}C atom percent enrichment (APE) data of $n \geq 5$ individual cells grown under
continuous light or diurnal light. The dashed lines represent the maximum ^{13}C APE that occurred
in *M. japonicum* monoculture controls provided sucrose as a result of nonspecific background or
direct CO_2 incorporation during heterotrophic growth; see SI Fig. 6 for more details. Asterisks
indicate significant differences over time in a given culture by two-tailed Student's *t*-test ($p <
0.05$). (B) C_{net} of *M. japonicum* cells ($n \geq 10$), calculated relative to the algal biomass as the
source. Median values are indicated, and asterisks indicate significant differences by two-tailed
Student's *t*-test ($p < 0.05$). (C) ^{15}N APE data of $n \geq 5$ individual cells grown under continuous

light or diurnal light. Asterisks indicate significant differences over time in a given culture by

two-tailed Student's *t*-test ($p < 0.05$).

Figure 4: *M. japonicum* undergoes reductive division upon depletion of sucrose and during algal coculture. Duplicate cultures ($n = 2$) (monoculture with 150 µg/ml sucrose or coculture under continuous light or diurnal light) were grown as described in Fig. 2 but with 50% of their NH_4^+ provided as ^{15}N and with $^{13}CO_2$ added to the air starting 13 h after inoculation. Isotope enrichment was measured using nanoSIMS and cells were imaged by fluorescence microscopy at the indicated times after inoculation. The grey backgrounds indicate sample timepoints that occurred during the dark phase. (A) N_{net} of *M. japonicum* cells ($n \geq 10$) (circles) relative to the

calculated expected N_{net} (blue bars) if each doubling in CFU/ml represented a doubling in biomass (SI Fig. 8). Black lines represent mean values. (B) Representative fluorescence microscopy images of cells stained with DAPI. (C) Cell length of $n \geq 50$ *M. japonicum* cells per culture type ($n \geq 25$ cells from each of the duplicate cultures) measured using ImageJ. Mean cell length is indicated, and asterisks indicate significant differences by two-tailed Student's *t*-test ($p < 0.05$).

Figure 5: *M. japonicum* delivers vitamin B$_{12}$ to *C. reinhardtii* despite reductive division.

Triplicate cultures ($n = 3$) were maintained under continuous light, shaken at 130 rpm, and assayed at the indicated times after inoculation. (A) Representative fluorescence microscopy images of *M. japonicum* with 150 µg/ml sucrose, in coculture with *C. reinhardtii* wild type, and in coculture with *C. reinhardtii mete1*. Cells were stained with DAPI, and *C. reinhardtii* cells are indicated with white arrowheads. (B) Length of *M. japonicum* cells ($n = 50$, 10–20 cells from each of the three triplicate cultures) measured using ImageJ. Mean cell length is indicated. (C) Representative image of cultures 54 h after inoculation. (D) *M. japonicum* density in monoculture with 150 µg/ml sucrose (pink bars) or in coculture (black bars) with wild-type (circles) and *mete1 C. reinhardtii* (triangles), estimated as CFU/ml. (E) Cell density of *C.*

reinhardtii wild type (circles) or *mete1* (triangles) in coculture (black bars) and of *mete1* alone without added vitamin B_{12} (green bars). (F) Concentration of vitamin B_{12} in cultures measured using an *E. coli* bioassay. Error bars represent the standard deviation from the mean, and asterisks indicate significant differences by two-tailed Student's *t*-test ($p < 0.05$).

Figure 6: A small degree of algal lysis can explain the *M. japonicum* growth that occurs in coculture. Triplicate cultures (*n* = 3) of wild-type *C. reinhardtii* were grown under continuous light and shaken at 125 rpm for 24 h, reaching 1–4x10^6 cells/ml. These cultures were then split and used to generate parallel live algal cultures, algal spent media, and algal cell lysate, each then inoculated with *M. japonicum* in triplicate (*n* = 3). (A) *M. japonicum* density 45 h after inoculation in monoculture with no reduced carbon source, coculture with live *C. reinhardtii* cells, *C. reinhardtii* spent medium, or *C. reinhardtii* cell lysate. Asterisks indicate significant differences by two-tailed Student's *t*-test (*p* < 0.05). (B) The approximate degree of *C. reinhardtii* cell lysis required to explain *M. japonicum* growth in coculture based on *M. japonicum* growth yield in *C. reinhardtii* cell lysate (blue bar), and the degree of *C. reinhardtii* cell lysis measured in the cultures used for the experiment in panel (A) by CellTox Green fluorescence, reported relative to a 100% killed control. The expected value and the mean value are indicated. (C) *M. japonicum* density in triplicate cultures (*n* = 3) over time after inoculation in monoculture with or without 150 µg/ml sucrose, in *C. reinhardtii* spent medium, or in *C.*

reinhardtii cell lysate when diluted to the same NPOC concentration as the spent medium. The

asterisk indicates a significant difference in density between spent medium and diluted cell lysate

40 h after inoculation by Student's *t*-test ($p < 0.05$); density was not significantly different

between these two conditions at the other timepoints examined ($p > 0.05$). Error bars represent

the standard deviation from the mean.