Silver nanoparticles can be sampled by ultrafiltration probe but elution into & recovery from

plasma and DPBS differs *in vitro*

Marije Risselada[1*], Robyn R. McCain[2], Miriam G Bates[1], Makensie L Anderson[1]

* Corresponding author

E-mail: mrissela@purdue.edu

[1]Department of Veterinary Clinical Sciences, College of Veterinary Medicine, Purdue

University, West Lafayette, Indiana, United States of America

[2]Center for Clinical Translational Research, College of Veterinary Medicine, Purdue University,

West Lafayette, Indiana, United States of America

Abstract

We compared 1) the influence of elution fluid on rate, pattern, and completeness of silver nanoparticle (AgNP) elution, and 2) ultrafiltration (UF) probe and direct sampling *in vitro*. Six specimens (2.5ml of 0.02mg/ml 10nm AgNP and 5.0ml of 30% poloxamer 407) contained in a dialysis tube (12-14kDa pores) were placed in 100ml Dulbecco's Phosphate Buffered Saline (DPBS) (n=3) or canine plasma (n=3) for 96h on a stirred hot plate (37°C and 600rpm) and sampled 20 times. Six pipette and UF probe samples were taken of a 0.001mg AgNP/ml DPBS or plasma solution. Inductively coupled plasma mass spectrometry was used to analyze Ag. Stock plasma contained Ag. At 96h, 5/6 dialysis tubes had not fully released AgNP. One peak in hourly Ag increase was present in DPBS (10-13h), and two peaks in plasma (6-8h and 10-13h). The hourly Ag increase in plasma decreased earlier than in DPBS. UF probe sampling was possible in both DPBS and plasma and resulted in higher Ag concentrations but with more variation than pipette samples. While *in vitro* use of DPBS might be more cost effective, plasma should be considered due to difference in elution and recovery.

Introduction

With the increase in antimicrobial resistance in veterinary medicine [1,2], novel strategies and antimicrobials to combat infections are (re)gaining favor. One strategy might be to deliver a high local dose of antibiotics [3-7]. Another might be to explore the use of non-antibiotic antimicrobials, such as silver (Ag), especially silver nanoparticles (AgNP). They have received considerable interest for their use against micro-organisms [8,9] and in wound care [10]. Prior silver elution studies have been performed into phosphate-buffered saline (PBS) from poloxamer 407 [11], into deionized water baths (from silver oxide films) [12], or aqueous samples with varying NaCl concentrations (from solid coated specimens) [13]. Plasma might be a closer *in vitro* fluid approximation to an *in vivo* environment than PBS and could mimic protein interactions that might be encountered. The elution into human plasma from an antibiotic containing polymethyl methacrylate (PMMA) construct as well as the effect plasma on PMMA properties was reported [14]. However, plasma has not been evaluated for suitability of Ag release studies, and whether the benefit could outweigh the increased cost of commercial canine plasma (~$2,400/500ml; Innovative research, Novi, MI) over PBS (~$30/500ml; Gibco, Thermofisher Scientific, Billings, MT).

The ultimate goal of sustained release compounds assessed by elution studies is their *in vivo* application. Ultrafiltration (UF) probes have been used successfully in dogs [15,16] and are used to sample compounds present in the tissue, without the need for more invasive sampling methods, such as tissue biopsies [17]. Establishing the feasibility of sampling Ag with UF probes would be preferable prior to their use to investigate the *in vivo* pharmacokinetic profile of Ag containing local delivery products. However, the possibility of sampling AgNP via UF probe has

not been assessed, nor has a comparison of canine UF probes and direct sampling been reported

for AgNP.

We aimed to 1) determine and compare the influence of elution fluid on rate, pattern, and

completeness of AgNP elution *in vitro*, and 2) compare UF probe sampling with direct samples.

We hypothesized that 1) elution of AgNP into Dulbecco's PBS (DPBS) will be like elution in

canine plasma, and that 2) UF probe sampling will be constant over time and yield similar results

as direct sampling.

Materials and Methods

An observational elution study and repeat sampling study were performed separately. Canine Ultrafiltration Probes (BASi Instruments, West-Lafayette, IN) were assembled per manufacturer instructions [18].

Elution

Six elution specimens were prepared. Each contained 2.5ml of a 0.02mg/ml 10nm silver nanoparticles (AgNP) in aqueous buffer with sodium citrate as stabilizer (Sigma Aldrich, Saint Louis, MO) mixed with 5.0ml 30% poloxamer 407 (Pluronic® F-127, Sigma Aldrich) (1:2 ratio with a total Ag content of 0.05mg/specimen). The specimens were prepared in individual 12ml syringes <2 hours before use and stored refrigerated and shielded from light. Individual 150ml crystallization dishes (Synthware glass, Pleasant Prairie, WI) with 100ml fluid were prewarmed for 2 hours prior to starting the elution study. Three dishes had Dulbecco's phosphate-buffered saline without Calcium chloride or magnesium chloride added (DPBS, Gibco, Thermofisher Scientific, Billings, MT) and three had canine plasma with dipotassium EDTA (K2-EDTA) (IGCNPLAK2E500ml Canine Plasma lot 41273, Innovative research, Novi, MI). A 10 multi-position hotplate with magnetic stir bars was used (RT 10, IKA Magnetic Stirrers, Wilmington, NC) with settings at 37°C and 600 rpm throughout the experiment. Room temperature was set at 68°F. Specimens were created using a 10cm long strip of 1 inch dialysis tubing with 12-14kDa pores (Carolina Biological Supply Co, Burlington, NC) [19]). The distal free end was folded up length wise and secured in folded position using 2 large surgical stainless steel hemoclips (Hemoclips® Teleflex, Morrisville, NC) in opposite direction 11]. The Ag-poloxamer mix was then placed in the tube, and the proximal free end of the tube folded and closed in similar fashion. All specimens were created in one batch and then placed in the prewarmed DPBS or

canine plasma within 5 minutes after assembly. The t=0 sample was taken immediately after all specimens were submerged in the same order of assembly and placement. Twenty samples (0.15ml each) were taken over 96 hours with a decrease in frequency (0, 1, 2, 3, 4, 5, 6, 8, 10, 13, 17, 22, 27, 34, 42, 48, 58, 66, 72, 96 hours) with the sampling order consistent throughout. At 96 hours, an additional sample (by needle aspiration through the tube) of the fluid contained within the dialysis tube was taken.

Ultrafiltration probe sampling

Two solutions were prepared immediately prior to sampling in a 50ml conical centrifuge tube (Thermofisher Scientific) using graduated pipettes (Thermofisher Scientific) and a 3ml syringe (Covidien, Mansfield, MA). Each specimen contained 0.03mg of AgNP (1.5ml) total in 28.5ml of either DPBS or canine plasma for a fluid concentration of 0.001mg/ml of AgNP. Ultrafiltration probes were tested for patency and sampling using DPBS prior to use. Full submerging of the probe near the lowest point of the tube was ensured, and positioning was checked after each manipulation. Vacutainers without additives (Vacuette Blood collection tube, 3.0ml, no additive, Greiner Bio-One, Sigma Aldrich, St Louis, MO) were used to collect the probe samples and an additional 10ml of air was removed to increase negative pressure in each vacutainer to increase sampling speed. The initial sample of the specimen was discarded to avoid risk of dilution by plain DPBS. Repeat samples using both a UF probe and direct sampling were taken at 0, 5, 10, 15, 30 and 60 minutes.

Samples of stock AgNP, DPBS and plasma were taken prior to the start of the study. All samples were stored at -78°C until batch analysis.

Sample and data analysis

The quantity of Ag in each sample was determined via inductively coupled plasma mass

spectrometry (ICP-MS) as previously described [11]. DPBS samples were diluted in 2% HNO3

and plasma samples were digested over night at 70C in the 1:1 mixture of 70% HNO3 and H2O2

and analyzed using ICP-MS (Perkin Elmer NexION 300D) to determine the concentration of

silver within each sample. The short-term precision was less than 3% relative standard deviation

(RSD), and the long-term stability was <4% relative standard deviation over 4 hours. Isotope-

ratio precision was less than 0.08% relative standard deviation. The AgNP detection limit was

0.001ng/ml, and quantification limit at 0.002ng/ml, and all samples below this limit were

recorded as 0ng/ml. Silver concentrations are expressed in parts per billion (ppb), with

1ppb=1ng/ml. Hourly increase in Ag was calculated as the difference between the measured

values at two subsequent time points divided by the hours between time points (expressed as

ppb/hr). The data of the 3 elution specimens will be expressed graphically as mean±SD. The

values of the repeat sampling experiment will be reported as individual results and a mean±SD

(RSD) in an observational manner without further statistical analysis. The RSD was calculated

by dividing the SD by the mean and will be expressed as a %.

Results

Silver concentrations for stock solutions were: DPBS 0.19ppb; plasma 4.83ppb and AgNP 24,940ppb. Assembled UF probes initially did not reliably yield appropriate negative suction to obtain a sample, and probes were not re-usable for repeat experiments. Additional negative pressure applied to the vacutainers together with sealing connecting points with glue (3g Tube, The Gorilla Glue Company, Cincinnati, OH) allowed single use sampling.

Elution

No leakage of any dialysis tubes was observed under gentle pressure immediately after assembly. Five out of 6 dialysis tubes were fully filled at 96 hours, the sixth was not fully filled when retrieved. The five fully filled specimens still contained more Ag than the surrounding fluid (Table 1), and release of Ag was not complete at 96 hours. A burst release of Ag was seen both into DPBS and plasma in the first 13 (DPBS) and 8 (plasma) hours, with the baseline amount in plasma higher than in DPBS (Fig 1). The highest hourly increase in Ag was between 10-13 hours in both fluids, with a clear single peak in DPBS (20.66ppb/hr) whereas two peaks were present for plasma: between 6-8 hours (12.31ppb/hr) and between 10-13 hours (13.85ppb/hr). The hourly increase of Ag in DPBS tapered between 27-84 hours but continued throughout. The increase in Ag in plasma decreased earlier, between 8-60 hours, and measured Ag decreased thereafter. The amount of Ag measured at 96 hours in fluid and the specimens was higher in DPBS than in plasma (Fig 2).

Fig 1: A. Elution of Ag into either DPBS (light grey) or plasma (dark grey), expressed as ppb over 96 hours. The elution of Ag into DBPS continued, leading to a higher fluid concentration, whereas the increase of Ag in plasma tapered between 24 – 60 hours and Ag decreased after 60 hours. B. The increase in Ag is expressed as Ag (ppb)/hour measured over the timeframe prior to

the sampling time. A burst elution pattern is evident for both elution fluids. C. Elution of Ag into

either DPBS or plasma, expressed as ppb over the first 22 hours. D. Hourly increase in Ag over

the first 22 hours.

Fig 2: A. Ag amount (ppb) in elution fluid and remainder of the specimen at 96 hours.

Measured Ag content in both the fluid and specimen remnant were less in plasma than DPBS. B.

Pipette and ultrafiltration (UF) probe sampling of 1,247ppb Ag in DPBS and plasma solution.

Both methods, in both mediums underestimated the Ag content. The UF probe samples had a

wider variation in sampled Ag. The dotted line indicates 1,247ppb Ag.

Table 1 Silver (Ag) at 96 hours expressed as parts per billion (ppb).

Ag in DPBS		Ag in Plasma	
Fluid	Specimen	Fluid	Specimen
2137.97*	203.32*	25	32.34
293.18	512.65	20.02	68.29
248.75	528.26	23.36	58.90

149   Each tube contained 6660ppb Ag at the start of the experiment. *denotes the sample that was not
150   fully filled and the surrounding fluid.
151
152

Ultrafiltration probe sampling

Anticipated Ag content based on the stock (24,940ppb Ag (AgNP stock) diluted in 30ml) was

1,247ppb Ag of the diluted fluid. Ultrafiltration probes were able to collect Ag in both DPBS and

plasma, with Ag content in fluid obtained via UF probe sampling higher than the corresponding

pipette samples (Fig 2). However, the UF probe-obtained samples had more variation and values

differed between sampling times and decreased over time (Table 2). Samples obtained by both

methods had a measured Ag lower than anticipated, with the underestimation greater in DPBS

than plasma.

Table 2: Pipette and ultrafiltration (UF) probe sampling of a planned 1,247ppb Ag in DPBS and
plasma solution.

Repeat sampling of Ag (ppb)	DPBS		Plasma	
Time (min)	UF probe	pipette	UF probe	pipette
0	502.67	1.19	791.28	116.19
5	48.31	11.31	946.58	99.18
10	33.88	0.93	370.54	225.08
15	15.47	3.18	117.41	74.07
30	2.33	0.80	1111.30	116.02
60	3.95	6.93	785.83	89.27
mean±SD (RSD)	101.10±197.53 (195%)	4.06±4.25 (105%)	687.2±372.2 (54%)	119.9±53.9 (45%)

163   **Table 2**. Probes were allowed to sample for 4 minutes to obtain enough sample volume. Direct
164   samples were taken using a pipettor at the time point. UF probe samples were taken by attaching
165   a vacutainer to the probe for 4 minutes at the start time. DPBS= Dulbecco's Phosphate Buffered
166   Saline, UF= ultrafiltration. The mean is provided with both standard deviation (SD) and relative
167   standard deviation (RSD).
168

Discussion

Elution into DPBS yielded different results than elution into plasma, with a longer sustained initial hourly increase in fluid Ag and more variation in Ag measurements with a resultant less smooth curve. A burst release pattern was evident for both DBPS and plasma, which corresponds with an earlier study where ~88% of the Ag released from poloxamer 407 within the first 24 hours [11]. While the pore size of the dialysis tube should allow AgNPs to cross freely, the Ag concentration within the dialysis tube was still higher at 96 hours than the surrounding fluid, indicating incomplete release of the specimen in both elution media. Silver could be measured in DPBS and plasma by UF probe sampling.

The higher initial values of Ag found in plasma might be explained by Ag measured in stock plasma vs DPBS solution. It was then followed by an increase of Ag content similar in shape to the DPBS curve. However, this initial starting value would mask the initial release of Ag at the time when release will be highest and is a limitation of using plasma as the elution medium. We chose to report the measured values as-is instead of detracting the value obtained from stock plasma, as we only had a single measurement for plasma rather than repeat samples and felt that correcting could inadvertently introduce error as well.

Plasma with EDTA was chosen as it was the cheapest commercial option available. Ethylenediaminetetraacetic acid (EDTA) chelates silver [20], and its presence in plasma theoretically could help extract Ag from the specimen contained within the dialysis tube. However, chelation of Ag into stable complexes also might hinder analyses, although measuring of excreted EDTA-metal complexes before and during chelation therapy has been described [20,21]. In addition, nitric acid digestion has been used to separate heavy metals from EDTA

[22]. However, depending on the drug for which the elution is performed a different anti-

coagulation agent might be preferred.

We used a 1:2 ratio of AgNP:poloxamer in this study to maximize the Ag content while still

maintaining full gelatination properties of the poloxamer. In a prior elution study [4] a 1:4 ratio

of AgNP:poloxamer was used with full gelatination, whereas specimens did not fully gel in a 1:1

ratio in a different study [23] but did when we tested at a 1:2 ratio (unpublished data). The

specimen composition as well as methodology (a single, larger amount of elution fluid with

sampling as opposed to full exchange of smaller amounts of elution fluid) between this study and

the prior study might account for the earlier tapering of elution and the flattening of the curve.

The fluid exchange would increase the gradient and therefore drive migration of Ag across the

dialysis tube membrane. The fluid quantity was chosen to allow a model with a larger amount of

fluid without full exchange was chosen to allow continuous stirring, the ability to submerge a

large specimen, and for cost of plasma (~$2,400 per 500ml).

While aluminum foil was wrapped tightly around the opening of the dishes, fluid loss, in

addition to sampling loss due to warming of the fluid was possible. Aluminum foil was chosen as

the initial intent was to perform all sampling with UF probes in addition to direct sampling and

using snap-on or screw-on lids would have damaged the probes. The study was converted to

direct sampling only at the last minute due to concerns of probe functioning and continued

patency during the initial assembly, and fluid was already in the dishes. The evaporative fluid

loss could account for the increase in Ag concentration of the fluid at the end of the study. No

attempt was made to estimate fluid losses during the study, as measuring at each time point

would have necessitated interrupting the elution and manipulating the specimen and might cause

additional fluid loss. However, measuring the volume at 96 hours could have been performed.

Fluid loss during a continuous release elution model is a known limitation of the model chosen

[19], however removing and replacing all fluid at each time point might artificially keep a larger

gradient intact than a continuous model would. In addition, a continuous model might more

closely resemble the *in vivo* decrease in gradient between the slow-release compound and its

environment.

The concentration of Ag measured in the samples obtained both by UF probe and pipette

sampling decreased over time. This could be explained by the stagnant nature of the fluids within

the tube, although care was taken to sample from the lower most area of the fluid in case of

sedimentation. Elution experiments ideally are performed with constant stirring [19], or agitation

prior to sampling [11]. We opted to not agitate the tubes to keep the probes fully submerged at

all times and given that the probe would remain in the same place *in vivo*. The tubes for pipette

sampling were similarly not agitated to not introduce a variable between the two methods, and

samples were obtained from the lowest area in the tube. The higher variation seen in samples

obtained by UF probes is interesting and indicates that care must be taken with *in vivo* sampling

to account for variability, for example by taking multiple samples.

Additional limitations could be inconsistency in mixing the specimens and solutions as well

as sampling or measurement errors. We included 3 repeats of the elution study to minimize the

effect of both inconsistency and sampling. Six repeat samples were used in the static sampling

study, however inconsistency between the solutions could still exist.

Conclusions

Elution into plasma and UF probe sampling of Ag in plasma are possible. While use of DPBS

might be more cost effective, plasma should be considered for *in vitro* silver elution studies due

to difference in elution and recovery. Silver nanoparticles in poloxamer 407 might release more

slowly than other drugs that have been studied under similar conditions. Both UF probe sampling

and direct sampling underestimate the actual Ag concentration in the fluid. Ultrafiltration probe

sampling of Ag is possible and UF probes could be used to measure *in vivo* local tissue

concentrations, but the possibility of variations in the sampled fluid should be kept in mind.

Acknowledgements

The authors thank the Nanomedicines Characterization Core Facility (NCore) at the Center

for Nanotechnology in Drug Delivery (CNDD) and Dr Marina Sokolsky for their assistance with

sample and data analysis.

References

References

1. Perrenten V, Kadlec K, Schwarz S, Andersson UG, Finn M, Greko C, et al. Clonal spread of

methicillin-resistant staphylococcus pseudintermedius in Europe and North America: an

international multi-centre study. J Antimicrob Chemo 2010; 65: 1145-1154.

2. Cohn LA, Middleton JR. A veterinary perspective on methicillin resistant staphylococci. J Vet

Emerg Crit Care 2010; 20: 31-45.

3. Ham K, Griffon D, Seddighi M, Johnson AL. Clinical application of tobramycin-impregnated

calcium sulfate beads in six dogs (2002-2004). J Am Anim Hosp Assoc 2008; 44: 320-326.

4. Peterson LC, Kim SE, Lewis DD, et al. Calcium sulfate antibiotic-impregnated bead

implantation for deep surgical site infection associated with orthopedic surgery in small animals.

Vet Surg 2021; 50: 748–757.

5. Lee SJ, Frederick SW, Cross AR. Use of an amikacin-infused collagen sponge concurrent with

implant removal for treatment of tibial plateau leveling osteotomy surgical site infection in 31

cases. Vet Surg 2019; 48: 700-706.

6. Reed TP, Thomas LA, Weeren FR, et al. A novel dextran polymer hydrogel local

antimicrobial therapy in dogs: A pilot study. Can Vet J. 2016; 57: 189-195.

7. Smith JJ, Chou P-Y, Filliquist B, et al. Number of previous surgeries and antibiotic resistance

decreases the success of local administration

of antibiotic-impregnated poloxamer 407 hydrogel when managing orthopedic surgical site

infections in dogs. J Am Vet Med Assoc 2023; 261: 1186-1192.

8. Li W-R, Xie X-B, Shi Q-S, Zeng H-Y, OU-Yang Y-S, Chen Y-B. Antibacterial activity and

mechanism of silver nanoparticles on Escherichia coli. Appl Microbiol Biotechnol.

2010;85:1115-1122.92. Senthilkumar P, Rashmitha S, Veera P, Ignatious C, SaiPriya C, Samrot

A. Antibacterial Activity of Neem Extract and its Green Synthesized Silver Nanoparticles

against Pseudomonas aeruginosa. J Pure Appl Microbiol. 2018;12:969-974.

10. Augustine R, Hasan A, Yadu Nath VK, Thomas J, Augustine A, Kalarikkal N, et al. Mater

Sci Mater Med. 2018;29:163.

11. Peterson J, Risselada M. In vitro elution of silver nanoparticles from three carrier media. Am

J Vet Res. 2022;83(8):1-7.

12. Goderecci SS, Kaiser E, Yanakas M, Norris Z, Scaturro J, Oszust R, et al. Silver Oxide

Coatings with High Silver-Ion Elution Rates and Characterization of Bactericidal Activity.

Molecules. 2017 Sep 7;22(9):1487. doi: 10.3390/molecules22091487.

13. Kent RD, Vikesland PJ. Controlled evaluation of silver nanoparticle dissolution using atomic

force microscopy. Environ Sci Technol. 2012 Jul 3;46(13):6977-84. doi: 10.1021/es203475a.

Epub 2012 Jan 13.

14. Schmidt-Malan SM, Greenwood-Quaintance KE, Berglund LJ, Mandrekar J, Patel R.

Oritavancin polymethylmethacrylate (PMMA)-compressive strength testing and in vitro elution.

J Orthop Surg Res. 2019;14(1):43. doi: 10.1186/s13018-019-1080-6.

15. Maaland MG, Guardabassi L, Papich MG. Minocycline pharmacokinetics and

pharmacodynamics in dogs: dosage recommendations for treatment of meticillin-resistant

Staphylococcus pseudintermedius infections. Dermatol. 2014;25(3):182-e47. doi:

10.1111/vde.12130.

16. Messenger KM, Wofford JA, Papich MG. Carprofen pharmacokinetics in plasma and in

control and inflamed canine tissue fluid using in vivo ultrafiltration. J Vet Pharmacol Ther.

2016;39(1):32-9. doi: 10.1111/jvp.12233. Epub 2015 May 10.

17. Maxwell EA, Phillips H, Clark-Price SC, Vieson MD, Selmic LE, Schaeffer DJ, et al.

Pharmacokinetics of platinum and safety evaluation of carboplatin-impregnated calcium sulfate

hemihydrate beads after implantation in healthy cats. Vet Surg 2020;49:748-757.

18. https://www.basinc.com/ultrafiltration-probes-and-accessories

19. Risselada M, Marcellin-Little DJ, Messenger KM, Griffith E, Davidson GS, Papich MG.

Assessment of in vitro release of carboplatin from six carrier media. Am J Vet Res.

2016;17(12):1381-1386.

20. Fulgenzi A and Ferrero ME. EDTA Chelation Therapy for the Treatment of Neurotoxicity.

Int J Mol Sci. 2019;20(5): 1019.

21. Robotti E, Quasso F, Mandredi M, Gosetti F, Mazzucco E, Isidoro C, et al. Determination by

ICP-MS and multivariate data analysis of elemental urine excretion profile during the EDTA

chelation therapy: A case study. J Trace Elem Biol. 2020;62:126608. doi:

10.1016/j.jtemb.2020.126608.

22.https://patentimages.storage.googleapis.com/5c/43/f6/ac08bfd470cea7/US20080038169A1.pd

f. Accessed 2023-10-27

23. Bates MG, Risselada M, Peña-Hernandez DC, Hendrix K, Moore GE. Antibacterial activity

of silver nanoparticles against Escherichia coli and methicillin-resistant Staphylococcus

pseudintermedius is affected by incorporation into carriers for sustained release. Am J Vet Res.

2024 Jan 13:1-11. doi: 10.2460/ajvr.23.10.0229. Online ahead of print.

Supporting information

314   S1: Excel file with original data

Figure 1ABCD

A

B

Figure 2AB