Silver nanoparticles can be sampled by ultrafiltration probe but elution into & recovery from

plasma and DPBS differs *in vitro*

Marije Risselada[1*], Robyn R McCain[2], Miriam G Bates[1], Makensie L Anderson[1]

* Corresponding author

E-mail: mrissela@purdue.edu

[1]Department of Veterinary Clinical Sciences, College of Veterinary Medicine, Purdue

University, West Lafayette, Indiana, United States of America

[2]Center for Clinical Translational Research, College of Veterinary Medicine, Purdue University,

West Lafayette, Indiana, United States of America

Abstract

We compared 1) the influence of elution fluid on rate, pattern, and completeness of silver

nanoparticle (AgNP) elution, and 2) ultrafiltration (UF) probe and direct sampling *in vitro*. Six

specimens (2.5ml of 0.02mg/ml 10nm AgNP and 5.0ml of 30% poloxamer 407) contained in a

dialysis tube (12-14kDa pores) were placed in 100ml Dulbecco's Phosphate Buffered Saline

(DPBS) (n=3) or canine plasma (n=3) for 96h on a stirred hot plate (37°C and 600rpm) and

sampled 20 times. Six pipette and UF probe samples were taken of a 0.001mg AgNP/ml DPBS

or plasma solution. Inductively coupled plasma mass spectrometry was used to analyze Ag.

Stock plasma contained Ag. At 96h, 5/6 dialysis tubes had not fully released AgNP. One peak in

hourly Ag increase was present in DPBS (10-13h), and two peaks in plasma (6-8h and 10-13h).

The hourly Ag increase in plasma decreased earlier than in DPBS. UF probe sampling was

possible in both DPBS and plasma and resulted in higher Ag concentrations but with more

variation than pipette samples. While *in vitro* use of DPBS might be more cost effective, plasma

should be considered due to difference in elution and recovery.

Introduction

With the increase in antimicrobial resistance in veterinary medicine [1,2], novel strategies and antimicrobials to combat infections are (re)gaining favor. One strategy might be to deliver a high local dose of antibiotics [3-7]. Another might be to explore the use of non-antibiotic antimicrobials, such as silver (Ag), especially silver nanoparticles (AgNP). They have received considerable interest for their use against micro-organisms [8,9] and in wound care [10]. Poloxamer 407 is a versatile reverse gelatinating polymer that is safe for implantable use [11,12], and has found use for delivery of antifungals [12], chemotherapeutics [11,13], and antibiotics [14]. It also might have inherent antimicrobial properties of its own [15]. Prior silver elution studies have been performed into phosphate-buffered saline (PBS) from poloxamer 407 [16], into deionized water baths (from silver oxide films) [17], or aqueous samples with varying NaCl concentrations (from solid coated specimens) [18]. Plasma might be a closer *in vitro* fluid approximation to an *in vivo* environment than PBS and could mimic protein interactions that might be encountered. The elution into human plasma from an antibiotic containing polymethyl methacrylate (PMMA) construct as well as the effect plasma on PMMA properties was reported [19]. However, plasma has not been evaluated for suitability of Ag release studies, and whether the benefit could outweigh the increased cost of commercial canine plasma (~$2,400/500ml; Innovative research, Novi, MI) over PBS (~$30/500ml; Gibco, Thermofisher Scientific, Billings, MT).

The ultimate goal of sustained release compounds assessed by elution studies is their *in vivo* application. Ultrafiltration (UF) probes have been used successfully in dogs [20,21] and are used to sample compounds present in the tissue, without the need for more invasive sampling methods, such as tissue biopsies [22]. Establishing the feasibility of sampling Ag with UF probes

would be preferable prior to their use to investigate the *in vivo* pharmacokinetic profile of Ag

containing local delivery products. However, the possibility of sampling AgNP via UF probe has

not been assessed, nor has a comparison of canine UF probes and direct sampling been reported

for AgNP.

We aimed to 1) determine and compare the influence of elution fluid on rate, pattern, and

completeness of AgNP elution *in vitro*, and 2) compare UF probe sampling with direct samples.

We hypothesized that 1) elution of AgNP into Dulbecco's PBS (DPBS) will be similar to elution

in canine plasma, and that 2) UF probe sampling will be constant over time and yield similar

results as direct sampling.

Materials and Methods

An observational elution study and repeat sampling study were performed separately. Canine Ultrafiltration Probes (BASi Instruments, West-Lafayette, IN) were assembled as per manufacturer instructions [23].

Elution

A commercial 0.02mg/ml 10nm silver nanoparticles (AgNP) in aqueous buffer with sodium citrate as stabilizer (Sigma Aldrich, Saint Louis, MO) was used for the study. Six elution specimens were prepared. Each contained 2.5ml of the 0.02 mg/ml 10nm AgNP stock solution mixed with 5.0ml 30% poloxamer 407 (Pluronic® F-127, Sigma Aldrich) (1:2 ratio with a total Ag content of 0.05mg/specimen). The specimens were prepared in individual 12ml syringes <2 hours before use and stored refrigerated and shielded from light. Individual 150ml crystallization dishes (Synthware glass, Pleasant Prairie, WI) with 100ml fluid were prewarmed for 2 hours prior to starting the elution study. Three dishes had Dulbecco's phosphate-buffered saline without Calcium chloride or magnesium chloride added (DPBS, Gibco, Thermofisher Scientific, Billings, MT) and three had canine plasma with dipotassium EDTA (K2-EDTA) (IGCNPLAK2E500ml Canine Plasma lot 41273, Innovative research, Novi, MI). A 10 multi-position hotplate with magnetic stir bars was used (RT 10, IKA Magnetic Stirrers, Wilmington, NC) with settings at 37°C and 600 rpm throughout the experiment. Room temperature was set at 68°F. Specimens were created using a 10cm long strip of 1 inch dialysis tubing with 12-14kDa pores (Carolina Biological Supply Co, Burlington, NC) [24]). The distal free end was folded up length wise and secured in folded position using 2 large surgical stainless steel hemoclips (Hemoclips® Teleflex, Morrisville, NC) in opposite direction [16]. The 1:2 AgNP-poloxamer mix was then placed in the tube, and the proximal free end of the tube folded and closed in

similar fashion. All specimens were created in one batch and then placed in the prewarmed

DPBS or canine plasma within 5 minutes after assembly. The t=0 sample was taken immediately

after all specimens were submerged in the same order of assembly and placement. Twenty

samples (0.15ml each) were taken over 96 hours with a decrease in frequency (0, 1, 2, 3, 4, 5, 6,

8, 10, 13, 17, 22, 27, 34, 42, 48, 58, 66, 72, 96 hours) with the sampling order consistent

throughout. At 96 hours, an additional sample (by needle aspiration through the tube) of the fluid

contained within the dialysis tube was taken.

Ultrafiltration probe sampling

Two specimens were prepared immediately prior to sampling in a 50ml conical centrifuge

tube (Thermofisher Scientific) using graduated pipettes (Thermofisher Scientific) and a 3ml

syringe (Covidien, Mansfield, MA). Each specimen contained 0.03mg of AgNP (1.5ml of the

commercial 0.02mg AgNP stock solution) total in 28.5ml of either DPBS (n=1) or canine plasma

(n=1) for a targeted fluid concentration of 0.001mg/ml of AgNP. Ultrafiltration probes were

tested for patency and sampling using DPBS prior to use for the study.

Full submerging of the probe near the lowest point of the tube was ensured, and positioning

was checked after each manipulation. Vacutainers without additives (Vacuette Blood collection

tube, 3.0ml, no additive, Greiner Bio-One, Sigma Aldrich, St Louis, MO) were used to collect

the probe samples and an additional 10ml of air was removed to increase negative pressure in

each vacutainer to increase sampling speed. The initial sample of the study specimen was

discarded to avoid risk of dilution by plain DPBS. Repeat samples using both a UF probe and

direct sampling using a pipette from the same area as the membrane of the UF probe were taken

at 0, 5, 10, 15, 30 and 60 minutes from the plasma and DPBS specimen.

Samples of commercial stock AgNP solution, DPBS and canine plasma were taken prior to

the start of the study. All samples were stored at -78°C until batch analysis.

Sample and data analysis

The quantity of Ag in each sample was determined via inductively coupled plasma mass

spectrometry (ICP-MS) as previously described [16]. DPBS samples were diluted in 2% HNO3

and plasma samples were digested over night at 70°C in the 1:1 mixture of 70% HNO3 and

H2O2 and analyzed using ICP-MS (Perkin Elmer NexION 300D) to determine the concentration

of silver within each sample. The short-term precision was less than 3% relative standard

deviation (RSD), and the long-term stability was <4% relative standard deviation over 4 hours.

Isotope-ratio precision was less than 0.08% relative standard deviation. The Ag detection limit

was 0.001ng/ml, and quantification limit at 0.002ng/ml, and all samples below this limit were

recorded as 0ng/ml. Silver concentrations were expressed in parts per billion (ppb), with

1ppb=1ng/ml. Hourly increase in Ag was calculated as the difference between the measured

values at two subsequent time points divided by the hours between time points (expressed as

ppb/hr). The data of the 3 elution specimens will be expressed graphically as mean±SD. The

values of the repeat sampling experiment will be reported as individual results and a mean±SD

(RSD) for both specimens in an observational manner without further statistical analysis. The

RSD was calculated by dividing the SD by the mean and will be expressed as a %.

Results

Silver concentrations for stock solutions used in this study were: DPBS stock 0.19ppb Ag; plasma stock 4.83ppb Ag and the commercial 0.02mg/ml AgNP solution contained 24,940ppb Ag. Assembled UF probes initially did not reliably yield appropriate negative suction to obtain a sample, and probes were not re-usable for repeat experiments. Additional negative pressure applied to the vacutainers together with sealing connecting points with glue (3g Tube, The Gorilla Glue Company, Cincinnati, OH) allowed single use sampling.

Elution

No leakage of any dialysis tubes was observed under gentle pressure immediately after assembly. Five out of 6 dialysis tubes were fully filled at 96 hours, the sixth was not fully filled when retrieved. The five fully filled specimens still contained more Ag than the surrounding fluid at 96 hours (Table 1), and release of Ag was not complete at 96 hours. A burst release of Ag was seen both into DPBS and plasma in the first 13 hours (DPBS, light grey) and 8 hours (plasma, dark grey), with the baseline amount in plasma higher than in DPBS (Fig 1AB). The increase of Ag measured in plasma decreased earlier, between 8-60 hours (dark grey, Fig 1AB), and the measured Ag decreased thereafter. The hourly increase of Ag in DPBS (light grey) tapered between 27-84 hours but continued throughout the study (Fig 1C). The highest hourly increase in Ag was between 10-13 hours in both DPBS (light grey) and plasma (dark grey) (Fig 1D). A clear single peak in DPBS (20.66ppb/hr) whereas two peaks were present for plasma: between 6-8 hours (12.31ppb/hr) and between 10-13 hours (13.85ppb/hr) (Fig 1D). The amount of Ag measured at 96 hours was higher in the remaining Ag:poloxamer mix contained within the tube than the surrounding fluid (both DPBS and plasma) (Fig 2). The Ag concentration measured

at 96 hours in DPBS was higher (both specimen and fluid) than in plasma (Fig 2). The mean±SD

total amount of Ag removed via sampling for the three DPBS fluid set ups was 330±51ng Ag

and for the three plasma fluid set ups 198.1±71ng Ag.

Fig 1: A. Elution of Ag into either DPBS (light grey) or plasma (dark grey), expressed as ppb

Ag over 96 hours. The elution of Ag into DBPS continued longer, leading to a higher fluid

concentration, whereas the increase of Ag in plasma tapered between 24 - 60 hours and Ag

decreased after 60 hours. B. The increase in Ag is expressed as Ag (ppb)/hour measured over the

timeframe prior to the sampling time. A burst elution pattern is evident for both elution fluids. C.

Elution of Ag into either DPBS or plasma, expressed as ppb Ag over the first 22 hours. D.

Hourly increase in Ag over the first 22 hours. The measured values of Ag in plasma varied more

for each time point and between time points as evidenced by the larger bars and the shape of the

curve.

Fig 2: Ag amount (ppb) in elution fluid (DPBS or plasma) and remainder of the

Ag:poloxamer specimen at 96 hours. DPBS elution fluid is shown in light grey, with the

remaining specimen in shaded light grey. The plasma elution fluid is shown in dark gray.

Measured Ag content in both the fluid and Ag:poloxamer specimen remnant at 96 hours were

less in plasma than DPBS, with the specimens still higher in Ag content than the surrounding

DPBS or plasma.

Table 1: Silver (Ag) at 96 hours expressed as parts per billion (ppb) for the remaining elution
fluid outside of the dialysis tube and the remaining specimen contained within the dialysis tube.

Ag in DPBS		Ag in Plasma	
Fluid	**Specimen**	**Fluid**	**Specimen**
2137.97*	203.32*	25	32.34
293.18	512.65	20.02	68.29
248.75	528.26	23.36	58.90

170 Each tube contained 6660ppb Ag at the start of the experiment. *denotes the specimen that was
171 not fully filled and the surrounding elution fluid. Silver at 96 hours was analyzed once for six
172 specimens (3 each in DPBS and plasma) and elution fluid.

Ultrafiltration probe sampling

Anticipated Ag content based on the stock (24,940ppb Ag (AgNP stock) diluted in 30ml) was

1,247ppb Ag of the diluted fluid. Ultrafiltration probes were able to collect Ag in both DPBS and

plasma, with Ag content in fluid obtained via UF probe sampling higher than the corresponding

pipette samples (Fig 3). However, the UF probe-obtained samples had more variation and values

differed between sampling times and decreased over time (Table 2). Samples obtained by both

methods had a measured Ag lower than anticipated, with the underestimation greater in DPBS

than plasma.

Fig 3: Pipette and ultrafiltration (UF) probe sampling of 1,247ppb Ag in DPBS (light grey)

and plasma (dark grey) solution. Pipette sampling is represented by a solid sphere (DPBS) or

square (plasma) symbol, while the corresponding UF sampling data is shown by a solid sphere or

square. Both methods underestimated the Ag content in both mediums. The samples obtained via

UF probe had a wider variation in measured Ag. The dotted line indicates 1,247ppb Ag.

Table 2: Repeat pipette and ultrafiltration (UF) probe sampling of a planned solution of 1,247ppb
Ag in DPBS and plasma solution (one specimen each).

Repeat sampling of Ag (ppb)	AgNP in DPBS specimen (n=1)		AgNP in plasma specimen (n=1)	
Time (min)	UF probe	pipette	UF probe	pipette
0	502.67	1.19	791.28	116.19
5	48.31	11.31	946.58	99.18
10	33.88	0.93	370.54	225.08
15	15.47	3.18	117.41	74.07
30	2.33	0.80	1111.30	116.02
60	3.95	6.93	785.83	89.27
mean±SD (RSD)	101.10±197.53 (195%)	4.06±4.25 (105%)	687.2±372.2 (54%)	119.9±53.9 (45%)

191   Probes were allowed to sample for 4 minutes to obtain enough sample volume. Direct samples
192   were taken using a pipettor at the time point. UF probe samples were taken by attaching a
193   vacutainer to the probe for 4 minutes at the start time. DPBS= Dulbecco's Phosphate Buffered
194   Saline, UF= ultrafiltration. The mean is provided with both standard deviation (SD) and relative
195   standard deviation (RSD) for each sampling method over time from the same fluid.
196

Discussion

Elution into DPBS yielded different results than elution into plasma, with a longer sustained initial hourly increase in fluid Ag and more variation in Ag measurements with a resultant less smooth curve. A burst release pattern was evident for both DBPS and plasma as elution fluid. The burst release corresponds with an earlier study where ~88% of the Ag was released from poloxamer 407 within the first 24 hours [16]. While the pore size of the dialysis tube should allow AgNPs to cross freely, the Ag concentration within the dialysis tube was still higher at 96 hours than the surrounding fluid (both DPBS and plasma), indicating incomplete release of the specimen in both elution media. Silver could be measured in DPBS and plasma by UF probe sampling.

The higher initial values of Ag found in plasma might be explained by Ag being present in stock plasma while no Ag was found in the DPBS stock solution. The higher initial value in plasma was then followed by an increase in Ag content similar in shape to the DPBS curve. However, the presence of Ag, and the initial starting value, would mask the initial release of Ag at the time when the release will be highest and is a limitation of using plasma as the elution fluid. We chose to report the measured values as-is instead of detracting the initial Ag measured obtained from stock plasma, as we only had a single measurement for the stock plasma rather than repeat samples and thus felt that correcting could inadvertently introduce error as well.

Plasma with EDTA was chosen as it was the cheapest commercial option available. Ethylenediaminetetraacetic acid (EDTA) chelates silver [25], and its presence in plasma theoretically could help extract Ag from the specimen contained within the dialysis tube. However, chelation of Ag into stable complexes also might hinder analyses, although measuring of excreted EDTA-metal complexes before and during chelation therapy has been described

[25,26]. In addition, nitric acid digestion has been used to separate heavy metals from EDTA

[27]. However, depending on the compound for which the elution is performed and its chelation

abilities with EDTA plasma with a different anti-coagulation agent might be preferred.

We used a 1:2 ratio of AgNP:poloxamer in this study to maximize the Ag content while still

maintaining the full gelatinating properties of the poloxamer. In a prior elution study, a 1:4 ratio

of AgNP:poloxamer was used and the specimens fully gelatinated [4]. However, specimens did

not fully gel in a 1:1 ratio in a different study [15] but did when we tested at a 1:2 ratio

(unpublished data). The specimen composition, as well as methodology (a single, larger amount

of elution fluid with sampling as opposed to full exchange of smaller amounts of elution fluid)

between this study and the prior study might account for the earlier tapering of elution and the

flattening of the curve. The fluid exchange would increase the gradient and therefore drive

migration of Ag across the dialysis tube membrane. The fluid quantity was chosen to allow a

model with a larger amount of fluid without full exchange to allow continuous stirring, the

ability to submerge a large specimen, and due to the cost of canine plasma (~$2,400 per 500ml).

Removing a sample at each time point (0.15ml each for a total of 3 ml) could impact the Ag

concentration by both removing fluid as well as Ag. The total amount of Ag removed over the

entirety of the 96 hours was 330ng for the DPBS fluid set ups and 158ng for the plasma set ups.

Ideally the removed amount of Ag would have been added back in a corrected calculating using

the currently present volume of the elution fluid, however, the fluid present at each time point

was not measured, and using an approximation might lead to a flawed correction, and more error

than not correcting.

The dialysis tube model was chosen to avoid the poloxamer from dissolving immediately

upon placing the AgNP:poloxamer mix in the elution DPBS or plasma fluid. Prior studies with

the same commercial AgNP and dialysis tubing [16] yielded appropriate migration of Ag

through the membrane. Given that no plasma was placed within the tube, there were no concerns

of Ag complexing with proteins, leading to inability to migrate across the membrane due to pore

size (12-14kDa).

While aluminum foil was wrapped tightly around the opening of the dishes, fluid loss, in

addition to sampling loss due to warming of the fluid was possible. Aluminum foil was chosen as

the initial intent was to perform all sampling with UF probes in addition to direct sampling and

using snap-on or screw-on lids would have damaged the probes. The study was converted to

direct sampling only at the last minute due to concerns of probe functioning and continued

patency during the initial assembly, and fluid was already in the dishes being prewarmed. The

evaporative fluid loss could account for the increase in Ag concentration of the fluid at the end of

the study. No attempt was made to estimate fluid losses during the study, as measuring at each

time point would have necessitated interrupting the elution and manipulating the specimen and

might cause additional fluid loss. However, measuring the volume at 96 hours could have been

performed. Fluid loss during a continuous release elution model is a known limitation of the

model chosen [24], however removing and replacing all fluid at each time point might artificially

keep a larger gradient intact than a continuous model would. In addition, a continuous model

might more closely resemble the *in vivo* decrease in gradient between the slow-release

compound and its environment.

The concentration of Ag measured in the samples obtained both by UF probe and pipette

sampling decreased over time. This could be explained by the stagnant nature of the fluids within

the tube, although care was taken to sample from the lowest area of the fluid in case of

sedimentation. Elution experiments ideally are performed with constant stirring [24], or agitation

prior to sampling [16]. We opted to not agitate the tubes in order to keep the probes fully

submerged at all times. In addition, the probe would remain in the same place *in vivo* as well,

and a stationary fluid was felt to be appropriate given that no concentration gradient was present.

Pipette samples were obtained from the area in the tube in the location of filtration membranes of

the UF as to not introduce a variable between the two methods. The higher variation seen in

samples obtained by UF probes is interesting and indicates that care must be taken with *in vivo*

sampling to account for variability, for example by taking multiple samples at each time point.

Additional limitations could be inconsistency in mixing the specimens and solutions as well

as sampling or measurement errors. We included 3 repeats of the elution study to minimize the

effect of both inconsistency and sampling. Six repeat samples were used in the static sampling

study, however inconsistency between the solutions could still exist.

Conclusions

Elution into plasma and UF probe sampling of Ag in plasma are possible. While use of DPBS might be more cost effective, plasma should be considered for *in vitro* silver elution studies due to difference in elution and recovery. Silver nanoparticles in poloxamer 407 might release more slowly than other drugs that have been studied under similar conditions. Both UF probe sampling and direct pipette sampling underestimate the actual Ag concentration in the fluid. Ultrafiltration probe sampling of Ag is possible and UF probes could be used to measure *in vivo* local tissue concentrations, but the possibility of variations in the sampled fluid should be kept in mind.

Acknowledgements

The authors thank the Nanomedicines Characterization Core Facility (NCore) at the Center for Nanotechnology in Drug Delivery (CNDD) and Dr Marina Sokolsky for their assistance with sample and data analysis.

References

1. Perrenten V, Kadlec K, Schwarz S, Andersson UG, Finn M, Greko C, et al. Clonal spread of methicillin-resistant staphylococcus pseudintermedius in Europe and North America: an international multi-centre study. J Antimicrob Chemo 2010; 65: 1145-1154.

2. Cohn LA, Middleton JR. A veterinary perspective on methicillin resistant staphylococci. J Vet Emerg Crit Care 2010; 20: 31-45.

3. Ham K, Griffon D, Seddighi M, Johnson AL. Clinical application of tobramycin-impregnated calcium sulfate beads in six dogs (2002-2004). J Am Anim Hosp Assoc 2008; 44: 320-326.

4. Peterson LC, Kim SE, Lewis DD, et al. Calcium sulfate antibiotic-impregnated bead implantation for deep surgical site infection associated with orthopedic surgery in small animals. Vet Surg 2021; 50: 748–757.

5. Lee SJ, Frederick SW, Cross AR. Use of an amikacin-infused collagen sponge concurrent with implant removal for treatment of tibial plateau leveling osteotomy surgical site infection in 31 cases. Vet Surg 2019; 48: 700-706.

6. Reed TP, Thomas LA, Weeren FR, et al. A novel dextran polymer hydrogel local antimicrobial therapy in dogs: A pilot study. Can Vet J. 2016; 57: 189-195.

7. Smith JJ, Chou P-Y, Filliquist B, et al. Number of previous surgeries and antibiotic resistance decreases the success of local administration of antibiotic-impregnated poloxamer 407 hydrogel when managing orthopedic surgical site infections in dogs. J Am Vet Med Assoc 2023; 261: 1186-1192.

8. Li W-R, Xie X-B, Shi Q-S, Zeng H-Y, OU-Yang Y-S, Chen Y-B. Antibacterial activity and mechanism of silver nanoparticles on Escherichia coli. Appl Microbiol Biotechnol. 2010;85:1115-1122.92. Senthilkumar P, Rashmitha S, Veera P, Ignatious C, SaiPriya C, Samrot

A. Antibacterial Activity of Neem Extract and its Green Synthesized Silver Nanoparticles against Pseudomonas aeruginosa. J Pure Appl Microbiol. 2018;12:969-974.

10. Augustine R, Hasan A, Yadu Nath VK, Thomas J, Augustine A, Kalarikkal N, et al. Mater Sci Mater Med. 2018;29:163.

11. Mathews KG, Linder KE, Davidson GS, et al. Assessment of clotrimazole gels for in vitro stability and in vivo retention in the frontal sinus of dogs. Am J Vet Res 2009; 70: 640–647.

12. Risselada M, Linder KE, Griffith E, et al. Pharmacokinetics and toxicity of subcutaneous administration of carboplatin in Poloxamer 407 in a rodent model. PloS one 2017;12:e0186018.

13. Risselada M, Tuohy JL, Law M, et al. Local administration of carboplatin in poloxamer 407 after an ulnar osteosarcoma removal in a dog. J Am Anim Hosp Assoc 2020;56(6):325-330.

14. Risselada M, Spies KE, Kim SY. Amikacin in 30% poloxamer 407 is a versatile local therapy with no adverse effects in 29 dogs. J. Am Vet Med Assoc. Under review.

15. Bates MG, Risselada M, Peña-Hernandez DC, Hendrix K, Moore GE. Antibacterial activity of silver nanoparticles against Escherichia coli and methicillin-resistant Staphylococcus pseudintermedius is affected by incorporation into carriers for sustained release. Am J Vet Res. 2024 Jan 13:1-11. doi: 10.2460/ajvr.23.10.0229. Online ahead of print.

16. Peterson J, Risselada M. In vitro elution of silver nanoparticles from three carrier media. Am J Vet Res. 2022;83(8):1-7.

17. Goderecci SS, Kaiser E, Yanakas M, Norris Z, Scaturro J, Oszust R, et al. Silver Oxide Coatings with High Silver-Ion Elution Rates and Characterization of Bactericidal Activity. Molecules. 2017 Sep 7;22(9):1487. doi: 10.3390/molecules22091487.

18. Kent RD, Vikesland PJ. Controlled evaluation of silver nanoparticle dissolution using atomic

force microscopy. Environ Sci Technol. 2012 Jul 3;46(13):6977-84. doi: 10.1021/es203475a.

Epub 2012 Jan 13.

19. Schmidt-Malan SM, Greenwood-Quaintance KE, Berglund LJ, Mandrekar J, Patel R.

Oritavancin polymethylmethacrylate (PMMA)-compressive strength testing and in vitro elution.

J Orthop Surg Res. 2019;14(1):43. doi: 10.1186/s13018-019-1080-6.

20. Maaland MG, Guardabassi L, Papich MG. Minocycline pharmacokinetics and

pharmacodynamics in dogs: dosage recommendations for treatment of meticillin-resistant

Staphylococcus pseudintermedius infections. Dermatol. 2014;25(3):182-e47. doi:

10.1111/vde.12130.

21. Messenger KM, Wofford JA, Papich MG. Carprofen pharmacokinetics in plasma and in

control and inflamed canine tissue fluid using in vivo ultrafiltration. J Vet Pharmacol Ther.

2016;39(1):32-9. doi: 10.1111/jvp.12233. Epub 2015 May 10.

22. Maxwell EA, Phillips H, Clark-Price SC, Vieson MD, Selmic LE, Schaeffer DJ, et al.

Pharmacokinetics of platinum and safety evaluation of carboplatin-impregnated calcium sulfate

hemihydrate beads after implantation in healthy cats. Vet Surg 2020;49:748-757.

23. https://www.basinc.com/ultrafiltration-probes-and-accessories

24. Risselada M, Marcellin-Little DJ, Messenger KM, Griffith E, Davidson GS, Papich MG.

Assessment of in vitro release of carboplatin from six carrier media. Am J Vet Res.

2016;17(12):1381-1386.

25. Fulgenzi A and Ferrero ME. EDTA Chelation Therapy for the Treatment of Neurotoxicity.

Int J Mol Sci. 2019;20(5): 1019.

26. Robotti E, Quasso F, Mandredi M, Gosetti F, Mazzucco E, Isidoro C, et al. Determination by

ICP-MS and multivariate data analysis of elemental urine excretion profile during the EDTA

chelation therapy: A case study. J Trace Elem Biol. 2020;62:126608. doi:

10.1016/j.jtemb.2020.126608.

27.https://patentimages.storage.googleapis.com/5c/43/f6/ac08bfd470cea7/US20080038169A1.pd

f. Accessed 2023-10-27

Supporting information

S1: Excel file with original data