Oncogenic RAS signaling suppresses ferroptosis via transcriptional upregulation of
GCH1

Jonathan K. M. Lim[1*], Frauke Stölting[1#], Tal Levy[2,3#], Leonie Thewes[4], Daniel Picard[1,5,6] Sofya
Tishina[7,8], Hai-Feng Zhang[9,10], Oksana Lewandowska[1,11], Tobias Reiff[11], Marc Remke[1,5,6],
Johannes Brägelmann[7,12,13], Filippo Beleggia[7,12,14], Carsten Berndt[4], Silvia von Karstedt[7,8,13],
Guido Reifenberger[1,6], Barak Rotblat[2,3], Gabriel Leprivier[1*]

[1]Institute of Neuropathology, University Hospital Düsseldorf and Medical Faculty, Heinrich
Heine University, Düsseldorf, Germany

[2]Department of Life Sciences, Faculty of Natural Sciences, Ben-Gurion University of the
Negev, Beer-Sheva, Israel

[3]The National Institute for Biotechnology in the Negev, Ben-Gurion University of the Negev,
Beer-Sheva, Israel

[4]Department of Neurology, University Hospital Düsseldorf and Medical Faculty, Heinrich-
Heine University Düsseldorf, Düsseldorf, Germany

[5]Department of Pediatric Oncology, Hematology, and Clinical Immunology, University Hospital
Düsseldorf and Medical Faculty, Heinrich Heine University, Düsseldorf, Germany

[6]German Cancer Consortium (DKTK) partner site Essen/Düsseldorf, Düsseldorf, Germany

[7]Department of Translational Genomics, University Hospital Cologne and Faculty of Medicine,
University of Cologne, Cologne, Germany

[8]CECAD Cluster of Excellence, University of Cologne, Cologne, Germany

[9]Department of Pathology and Laboratory Medicine, University of British Columbia,
Vancouver, BC, Canada

[10]Department of Molecular Oncology, BC Cancer Agency, Vancouver, BC, Canada

[11]Institute of Genetics, Heinrich-Heine-University, Düsseldorf, Germany

[12]Mildred Scheel School of Oncology Aachen Bonn Cologne Düsseldorf (MSSO ABCD),
Faculty of Medicine and University Hospital Cologne, University of Cologne, Cologne,
Germany

[13]Center for Molecular Medicine Cologne (CMMC), Faculty of Medicine and University Hospital
Cologne, University of Cologne, Cologne, Germany

[14]Department I of Internal Medicine, Faculty of Medicine and University Hospital Cologne,
University of Cologne, Cologne, Germany

*[*]Correspondence: jonathan.lim@med.uni-duesseldorf.de, gabriel.leprivier@med.uni-
duesseldorf.de

[#]Equal contribution

Abstract

Ferroptosis is an iron-dependent form of regulated cell death arising from excessive lipid
peroxidation. While seminal work described that oncogenic RAS transformation drives
synthetic lethal vulnerability to archetypal ferroptosis inducers including erastin (eradicator of
RAS and ST-expressing cells) and RSL3 (Ras selective lethal 3), more recent work suggest
that oncogenic RAS signaling may confer ferroptosis resistance. Thus, the impact of
oncogenic RAS on the cellular response to ferroptosis is still unclear. Here, we provide unifying
evidence across multiple cellular models that oncogenic RAS signaling suppresses
ferroptosis. Using integrated proteo- and transcriptomic analyses, we uncovered that

oncogenic RAS signaling upregulates the ferroptosis suppressor GTP cyclohydrolase I
(GCH1) via transcriptional induction by the transcription factor ETS1 downstream of the RAS-
MAPK signaling cascade. Targeted repression of Gch1 or of the tetrahydrobiopterin (BH4)
synthesis pathway, which is mediated by GCH1, was sufficient to sensitize oncogenic RAS
transformed cells to ferroptosis in 2D and 3D cell models, highlighting a mechanism through
which RAS promotes resistance to ferroptosis induction. Furthermore, we found that GCH1
expression is clinically relevant and correlates with RAS signaling activation in human cancers.
Overall, this study redefines oncogenic RAS signaling to be a ferroptosis suppressor, and
identifies GCH1 as a mediator of this effect and a potential vulnerability for targeting RAS
driven cancers.

Introduction

Ferroptosis is an iron-dependent, non-apoptotic form of cell death resulting from the excessive
accumulation of lipid hydroperoxides. Cellular defenses against ferroptosis are mediated by
the combined action of the cystine/glutamate antiporter xCT, which dictates the rates of
glutathione (GSH) biosynthesis by importing cystine, as well as the GSH-dependent
selenoenzyme glutathione peroxidase 4 (GPX4), which directly detoxifies lipid hydroperoxides
(1–4). Independent of the xCT-GPX4 axis, other major regulators of ferroptosis include
ferroptosis suppressor protein 1 (FSP1), which mediates ubiquinol (CoQH2) recycling, and
GTP cyclohydrolase 1 (GCH1), which mediates tetrahydrobiopterin (BH4) synthesis, both of
which are lipophilic radical-trapping antioxidants with the propensity to reduce toxic lipid
hydroperoxides (5–9). Ferroptosis has been implicated in cancer and is proposed to act as a
barrier against tumorigenesis. Indeed, the tumor suppressors p53 and BAP1 have been
individually linked to ferroptosis induction through their ability to repress the expression of
SLC7A11 (the gene encoding xCT) (10, 11). Intriguingly, ferroptosis was initially described in
synthetic lethal screens used to identify compounds that preferentially kill mutant RAS[G12V]
transformed cells but not normal isogenic counterparts – termed RAS-selective lethal (RSL)
compounds, which include erastin (xCT inhibitor) and RSL3 (GPX4 inhibitor) (12–14). These
initial studies suggest that oncogenic RAS sensitizes cells to ferroptosis, which was supported
by some studies using newly-discovered ferroptosis inducers (FINs)(3, 15).

RAS is a small GTPase encoded by the *HRAS*, *KRAS*, and *NRAS* proto-oncogenes (16–18).
Physiologically, RAS is responsible for transmitting growth factors signals from receptor
tyrosine kinases through to the activation of the prototypical RAF-MEK-ERK and PI3K-AKT
pathways, which promote proliferation and survival, respectively (17, 18). *RAS* genes are
frequently mutated in human cancer, particularly in pancreatic, colorectal, and lung
adenocarcinomas (18). Upon mutation, RAS acquires oncogenic properties that mediate
tumor initiation and maintenance. Furthermore, oncogenic RAS confers resistance to
apoptosis through multiple signaling cascades, rendering it difficult to target RAS-mutant
tumors (19–21). Therefore, exploiting ferroptosis induction as a means to circumvent such
resistance represents an attractive alternative approach to treating RAS-mutant cancers.

Nonetheless, the impact of oncogenic RAS on ferroptosis is still unclear. Despite the initial
discovery of ferroptosis as an oncogenic RAS-selective lethality in the cellular systems
mentioned, previous work by us and others demonstrated that oncogenic RAS in fact
enhances intracellular anti-oxidant capacity. In particular, we demonstrated that oncogenic
RAS signaling transcriptionally upregulates *SCL7A11* to promote glutathione biosynthesis and
confer resistance to oxidative stress (22). Furthermore, we recently reported that oncogenic
RAS induces FSP1 transcription to protect cells against RSL3-induced ferroptosis (23). Thus,
the *bona fide* role of oncogenic RAS in modulating the response to ferroptosis needs to be
clarified, as this holds significant future ramifications for the clinical application of FINs as a
cancer therapy.

Here, we provide substantial evidence across multiple cellular models that oncogenic RAS
expression suppresses ferroptosis induction by multiple classes of FINs, strongly supporting
that oncogenic RAS signaling mediates the protection of cells, rather than their sensitization

to ferroptosis induction. Transcriptome and proteomic profiling further reveal that among
various ferroptosis modulators, *GCH1* expression is highly upregulated upon KRAS^{G12V}
transformation. Indeed, we provide strong evidence that *GCH1* transcription is under the
regulation of the transcription factor ETS1 downstream of the RAS-driven RAF-MEK-ERK
pathway. Functionally, we uncovered that GCH1 and BH4 synthesis contributes to oncogenic
KRAS^{G12V}-mediated ferroptosis resistance and reduction of lipid peroxidation (LP) in vitro and
in vivo. This has clinical relevance as we report that *GCH1* expression is increased in RAS
mutant tumors. Collectively, these findings offer 1) a renewed perspective for the role of
oncogenic RAS as a predictor of ferroptosis resistance, 2) the mechanism by which this
occurs, and 3) the translational potential of inhibiting GCH1/BH4 as a novel strategy for the
sensitization of RAS mutant tumors to ferroptosis.

Results

Oncogenic RAS signaling activation confers resistance to ferroptosis

The role of oncogenic RAS in modulating cellular susceptibility to ferroptosis is still
controversial. To provide unifying evidence as to the impact of oncogenic RAS on ferroptosis,
we sought to interrogate the response of multiple isogenic cellular RAS systems to different
classes of ferroptosis inducers (FINs). First, we found that mouse NIH 3T3 fibroblasts
transformed with mutant KRAS^{G12V} (NIH 3T3 KRAS^{G12V}) were completely resistant to the class
I FINs erastin and sulfasalazine (SSZ), class II FINs ML162 and ML210, and class IV FIN
FINO2, while non-transformed counterparts NIH 3T3 MSCV showed varying levels of cell
death following a 16-hour treatment with each of these compounds (Fig. 1A and S1A). This is
consistent with our previous findings that mouse embryonic fibroblasts display enhanced
resistance to the class II FIN RSL3 following endogenous level expression of oncogenic KRAS
(23). Co-treatment with the ferroptosis-selective antioxidant ferrostatin-1 (Fer-1) abolished cell
death induced in non-transformed fibroblasts by the different FINs, confirming the induction of
ferroptosis (Fig. 1A and Fig. S1A). Strikingly, NIH 3T3 KRAS^{G12V} were completely resistant
even when exposed to high concentrations of erastin (up to 20 μM), while non-transformed
cells were susceptible across a range of erastin treatment (Fig. S1A). Second, we confirmed
these initial observations using a derivative of a widely-utilized mutant RAS overexpression
system, namely human BJ fibroblasts immortalized with the catalytic subunit of human
telomerase (hTERT), SV40 large T and small T antigen (LT and ST respectively) and
transformed by ectopic expression of oncogenic HRAS^{G12V} (BJ-hTERT/LT/ST/HRAS^{G12V}, also
known as BJeLR cells) (2, 3, 24). These models previously led to the identification of RSL3
and erastin as selective lethal agents against cells expressing both mutant RAS and small T
antigen (ST) (12–14). To determine the impact of mutant RAS in these models, but
independently of ST, we subjected BJ-hTERT/LT/HRAS^{G12V} and the corresponding isogenic
BJ-hTERT/LT control cells to FINs treatment. Interestingly, BJ-hTERT/LT/HRAS^{G12V} showed
enhanced resistance to RSL3, ML162, ML210 and erastin, as compared to BJ-TERT/LT (Fig.
1B), in contrast to previous findings obtained with BJeLR compared to BJ-hTERT (12–14).
While these results further support that oncogenic RAS confers resistance against ferroptosis
induction, they also highlight the contribution of ST to ferroptosis sensitivity (12). In agreement
with increased resistance to multiple FINs and our previously reported data, BJ-
hTERT/LT/HRAS^{G12V} cells expressed higher levels of SLC7A11 and FSP1 than BJ-hTERT/LT
(Fig. S1, B and C) (22, 23). Third, to extend the validity of our findings in human cell lines, we
assessed the impact of KRAS depletion on susceptibility to ferroptosis, using shRNA-
mediated knockdown of KRAS in mutant KRAS^{G12V}-expressing human pancreatic
adenocarcinoma cell line CFPAC1. Notably, KRAS knockdown rendered CFPAC1 cells more
sensitive to RSL3-induced cell death, which was again rescued by Fer-1 (Fig. 1C and D).

Finally, to determine the generalizability of oncogenic RAS-mediated resistance to ferroptosis,
we leveraged cancer cell line drug sensitivity data from the Cancer Therapeutic Response
Portal v2 (CTRPv2), available from Broad Institute's DepMap portal (25, 26). We observed a
correlation across over 700 cancer cell lines between *KRAS* mutation and resistance (i.e.,

higher median area under the curve [AUC] values) to ML162 (Fig. 1E), while this was not
observed for RSL3, ML210, and erastin (Fig. S1, D to F). Given that the RAS pathway can be
active even in cancer cells harboring *KRAS wild-type* (*KRAS^wt*), due to various genetic events,
we analyzed RAS pathway activity in the same dataset using a validated transcriptional
signature of 84 genes (RAS84) optimized to capture RAS oncogenic activity (27). Remarkably,
we found that higher RAS pathway activation positively correlated with resistance to ML162,
RSL3, ML210, and erastin (Fig. 1, F to I). Taken together, these findings provide strong
evidence that oncogenic RAS signaling activation functionally protects cells from ferroptosis
and correlates with ferroptosis resistance across a large number of cancer cells.

Oncogenic RAS signaling leads to upregulation of GCH1

The fact that RAS pathway activation correlates with ferroptosis resistance across many
cancer entities suggests that it may be mediated by additional mechanisms besides FSP1 and
xCT upregulation. Therefore, we next sought to identify novel downstream mediators of
oncogenic RAS signaling that could protect cells against ferroptosis. To this end, we
performed RNA sequencing of NIH 3T3 KRAS^G12V and control cells and integrated this data
with our previous proteomics analysis performed on the same cell lines (28). These screens
identified 2022 transcripts and 448 proteins that were significantly altered following KRAS^G12V-
expression, among which were 193 overlapping transcripts/proteins (Fig. S2A). Consistent
with our prior investigations, transcriptome profiling identified *Slc7a11* and *Aifm2* (encoding
Fsp1), but not *Gpx4*, as upregulated by oncogenic RAS signaling (Fig. 2A). Notably, the gene
encoding GTP cyclohydrolase I, *Gch1*, was among the most significantly upregulated genes
in NIH 3T3 KRAS^G12V cells (Fig. 2A). Indeed, among the 193 overlapping transcripts/proteins
significantly altered in KRAS^G12V-expressing cells, Gch1 represented the 5th most upregulated
transcript and 10th most upregulated protein (Table S1). Importantly, GCH1 was recently
identified as a suppressor of ferroptosis (7, 8). Indeed, GCH1 is the rate-limiting enzyme in
tetrahydrobiopterin (BH4) synthesis, a radical-trapping antioxidant with potent activity against
LP (Fig. 2B) (8). For this reason, we decided to focus on the regulation of *GCH1* by oncogenic
RAS. The upregulation of *Gch1* in NIH 3T3 KRAS^G12V versus control cells was confirmed using
qRT-PCR (Fig. 2C), as well as at the protein level using mass spectrometry-based proteomics
(Fig. 2D). In agreement, human BJ-hTERT/LT/HRAS^G12V cells showed a similar increase in
GCH1 expression relative to BJ-hTERT/LT isogenic counterparts (Fig. 2E). In human cancer
cells harboring *KRAS* mutations, including CFPAC1 and non-small cell lung carcinoma cell
line H460, shRNA-mediated knockdown of KRAS conversely led to decreases in *GCH1*
expression (Fig. 2, F and G and Fig. S2B). Similarly, treatment of mutant KRAS^G12C-expressing
pancreatic cancer cell line MIA PaCa-2 with the KRAS^G12C inhibitor AMG 510 decreased *GCH1*
expression (Figure 2H).

Additionally, by analyzing expression data of IMR90 human fibroblasts we found elevated
GCH1 expression upon stable expression of HRAS^G12V, as compared to control pBabe, but
not upon constitutively active AKT1 (Figure 2I) (29). This suggests that *GCH1* might be
upregulated downstream of RAS-MAPK signaling but not PI3K-AKT signaling. To further
ascertain this, we treated NIH 3T3 KRAS^G12V cells with the MEK inhibitor PD184352 or the
AKT inhibitor MK-2206 and found that the former but not the latter led to decreased *Gch1*
expression, further supporting that the regulation of *GCH1* downstream of oncogenic RAS is
mediated by MAPK (RAF-MEK-ERK) signaling (Figure 2J). Moreover, MEK inhibition led to a
sustained depletion of *Gch1* expression for up to 72 hours (Fig. S2C). Altogether, the data
suggests that oncogenic RAS signaling leads to upregulation of *GCH1* expression in a MAPK-
dependent manner.

GCH1 mediates oncogenic RAS-driven ferroptosis resistance

196 We next investigated the role of GCH1 in mediating ferroptosis suppression downstream of
197 oncogenic RAS signaling. For these studies, we first employed ferroptosis sensitive non-
198 transformed NIH 3T3 cells to determine whether supplementation with BH4, the end product
199 of the biosynthesis pathway downstream of GCH1, could phenocopy ferroptosis resistance

mediated by KRAS-mutant expression in these cells. Strikingly, cells supplemented with BH4, or its oxidized form dihydrobiopterin (BH2), showed near-complete insensitivity to erastin or FINO2, similar to NIH 3T3 KRAS^{G12V} cells (Fig. 3, A and B). While NIH 3T3 MSCV exhibited higher amounts of LP as NIH 3T3 KRAS^{G12V} cells when treated with erastin or FINO2, consistent with the ferroptosis-related phenotypes observed in these cells, the levels of LP were significantly reduced upon supplementation of NIH 3T3 MSCV with BH2 (Fig. 3, C and D). This was not the case with BH4 treatment (Fig. 3, C and D), suggesting a possible difference in the mechanism of ferroptosis protection between BH2 and BH4, or decreased BH4 efficiency since the oxidation of BH4 upon contact with culture medium may yield multiple oxidation products. Furthermore, BH4 or BH2 supplementation abrogated ferroptosis sensitivity of KRAS-knockdown CFPAC1 cells under RSL3 treatment (Fig. 3E), linking mutant KRAS-mediated ferroptosis resistance to the activity of the BH4 synthesis pathway. Finally, to ascertain of the contribution of GCH1 to KRAS^{G12V}-mediated resistance to ferroptosis, we knocked down *Gch1* expression in NIH 3T3 KRAS^{G12V} cells using CRISPR interference (CRISPRi). This was sufficient to restore the sensitivity of NIH 3T3 KRAS^{G12V} to ferroptosis induced by erastin or FINO2 (Fig. 3F and G, Fig. S3A). Taken together, these findings suggest that oncogenic RAS signaling leads to ferroptosis resistance by co-opting GCH1 function.

Targeting the BH4 synthesis pathway sensitizes oncogenic RAS to ferroptosis in vivo

Next, we sought to determine the contribution of GCH1 to oncogenic RAS-mediated tumorigenic potential or the ability of GCH1 inhibition to synergize with ferroptosis induction in vivo. To this end, we first assessed the ability of NIH 3T3 KRAS^{G12V} cells to grow under anchorage-independent conditions following CRISPRi-mediated *Gch1* knockdown using soft agar colony formation assays. While knockdown of *Gch1* expression did not observably decrease colony growth of KRAS-transformed cells under normal conditions, it resulted in significantly decreased tumorigenic potential of these cells under conditions of ferroptosis induction, suggesting that GCH1 inhibition may also synergize with ferroptosis induction in vivo (Fig. 4A and Fig. S3C). Similarly, this effect was observed in KRAS-transformed cells treated with SPRi3, an inhibitor against sepiapterin reductase (SPR), the enzyme mediating the final step of BH4 synthesis (Fig. 2B), under ferroptosis-induced conditions (Fig. 4B). In agreement, SPRi3 treatment synergized with RSL3-mediated ferroptosis induction in reducing the size of pancreatic organoids derived from the LsL-KRAS^{G12D} transgenic mouse model of pancreatic cancer (Fig. 4C) (30).

GCH1 expression is clinically relevant and correlates with oncogenic RAS activation in human cancers

It was previously reported that *GCH1* expression contributes to lung cancer, breast cancer, and glioblastoma development and that its overexpression is linked to poor prognosis of glioblastoma patients (31–33). Given the potential of GCH1 inhibition as a therapeutic strategy for the sensitization of RAS driven tumors to ferroptosis induction, we now analyzed the clinical relevance of *GCH1* expression in human cancers harboring KRAS mutation or constitutive RAS signaling activation through analysis of The Cancer Genome Atlas (TCGA) datasets. Notably, a pan-cancer analysis revealed that *GCH1* expression is higher in *KRAS-mutant* (*KRAS^{mut}*) tumors than in *KRAS^{wt}* tumors (Fig. 5A). We then extended the same analysis of *GCH1* expression to several tumor types of which a significant proportion of tumors are known to harbour activating mutations in *KRAS*, namely pancreatic adenocarcinoma (PAAD), colon adenocarcinoma (COAD) and lung adenocarcinoma (LUAD). Such analyses revealed that *GCH1* is overexpressed in PAAD, COAD, and LUAD relative to corresponding non-tumor tissue (Fig. 5, B, C, and D). Further, *GCH1* overexpression was confirmed in two orthogonal PAAD datasets (Fig. S4A and B) (34, 35). Consistent with the above-findings, we observed a positive correlation between *GCH1* levels and RAS signaling activation (based on RAS84 signature) in COAD, but also in several other human cancers including rectal adenocarcinoma (READ), uterine carcinosarcoma (UCS), skin cutaneous melanoma (SKCM), low-grade glioma (LGG), and GBM (Fig. 5E-J). Of note, *GCH1* expression was recently reported to be elevated

in glioblastoma but the mechanisms by which this occurs remains unclear (31). Together with
previous reports that genetic alterations in components of the RAS-MAPK cascade are
frequently observed in glioblastoma, our evidence suggests that *GCH1* overexpression is
associated with oncogenic RAS signaling activation in this tumor type (36). In agreement,
over-representation analysis of genes positively co-expressed with *GCH1* in glioblastoma
exhibited significant enrichment of genes known to be upregulated upon KRAS activation (Fig.
5K). Moreover, *Gch1* levels are higher in tumors derived from a genetically engineered mouse
model of high-grade glioma, constituted from *Nf1*-deletion (which promotes RAS pathway
activation), as compared to matched normal tissue (Fig. 5L), but this was not observed in a *c-*
Myc-driven mouse model of liver carcinoma (Fig. S4C) (37). Together, our findings reveal the
clinical relevance of *GCH1* regulation by the oncogenic RAS pathway in human cancers.

Transcriptional upregulation of *GCH1* by oncogenic RAS signaling is mediated by ETS1

We next sought to determine the mechanism by which oncogenic RAS signaling regulates
GCH1 expression. To this end, we conducted gene set enrichment analysis (GSEA) of genes
displaying positive co-expression with *Gch1* from the above RNA sequencing data of NIH 3T3
KRAS^{G12V} and control cells. Notably, genes having at least one binding site for the transcription
factor Ets-1 showed significant enrichment, and this gene set was among the top five
transcription factor target gene sets showing enrichment (Fig. 6A, Fig. S5A). Given the fact
that ETS1 is a known substrate of ERK downstream of the RAS-MAPK signaling pathway, we
postulated that the regulation of *GCH1* by oncogenic RAS is mediated by ETS1, which to our
knowledge has not been reported (Fig. 6B). In support of this, siRNA-mediated knockdown of
Ets-1 expression in NIH 3T3 KRAS^{G12V} cells significantly decreased *Gch1* expression (Fig.
6C). Similarly, knockdown of *ETS1* expression in H460 cells using two distinct siRNAs led to
a decrease in *GCH1* expression (Fig. 6D). Next, to ascertain whether ETS1 controls the
induction of *GCH1*, we performed dual-luciferase reporter assays in HEK293T cells using a
reporter construct consisting of a transcriptionally active region (TAR) of *GCH1* directly
upstream of *Firefly luciferase*. Sequence analysis of this region revealed the presence of three
ETS1 binding sites, based on the consensus 5'-GGAA/T-3' (Fig. 6E). While ectopic ETS1
expression was sufficient to activate the TAR of *GCH1* in a dose-dependent manner,
mutations in each of three putative ETS1 binding sites within this region abrogated this effect,
with concurrent mutations in all three sites having the strongest effect (Fig. 6, F and G). In
support of this, chromatin immunoprecipitation (ChIP)-qRT-PCR in NIH 3T3 KRAS^{G12V} cells
and ChIP-sequencing analysis of publicly-available UCSC Encode data provided strong
evidence that endogenous ETS1 occupies the human and mouse TAR region of *GCH1* (Fig.
6, H and I). Consistent with our functional analyses above, pan-cancer data from the Clinical
Proteomic Tumor Analysis Consortium (CPTAC) indicate that *GCH1* expression levels are
positively correlated with ETS1 protein expression (Fig. 6J), whereas publicly-available PAAD,
COAD, and LUAD datasets similarly show a positive correlation between *GCH1* and *ETS1*
mRNA expression (Fig. 6, K to M) (35, 38, 39). Together, these data support that *GCH1* is
transcriptionally upregulated by the MEK-ERK-ETS1 pathway downstream of oncogenic RAS.

Discussion

There is a growing interest in the potential of inducing ferroptosis in tumor cells as an
anticancer strategy. However, ferroptosis sensitivity varies substantially among cancer cells
due to distinct tissue lineages, mutation status of oncogenes or tumor suppressor genes, as
well as the acquisition of drug-tolerant or differentiated cell states (3, 10, 40–42). Thus,
understanding the mechanisms that contribute to LP and ferroptosis will aid the development
of effective approaches to overcome ferroptosis resistance. In this study, we demonstrate that
oncogenic activation of RAS signaling, one of the most mutated pathways in cancer, is a major
determinant of ferroptosis resistance. Notably, ferroptosis originally emerged as an alternative
form of necrosis induced by erastin and RSL3, identified from synthetic lethal compound
screens for their selective cytotoxicity towards engineered mutant HRAS-overexpressing cells
known as BJeLR cells (12–14). Following these seminal findings, the identification of novel

classes of FINs have often been accompanied by validation in the BJeLR model, over time substantiating a paradigm that oncogenic RAS-transformed cells are ostensibly sensitive to ferroptosis inducers, even though the association between KRAS mutational status and ferroptosis sensitivity could never be confirmed in comprehensive human cancer cell line screens (3, 43). On the contrary, our findings here, together with our previous study suggest rather that oncogenic RAS signaling renders cells resistant to ferroptosis, as demonstrated using multiple functional models (23). One of the models used here, BJ-hTERT/LT/HRAS[V12] cells, are similar to BJeLR but allowed us to interrogate the effect of mutant RAS independently of ST, which was initially used as an additional transforming factor. Indeed, in the absence of ST, mutant HRAS[V12] expression led to a paradoxical resistance to FINs suggesting that the presence of ST may confound the contribution of RAS signaling in modulating the sensitivity of cells to ferroptosis. Notably, one of the mechanisms by which ST promotes malignant transformation is by negatively regulating the protein phosphatase 2A (PP2A) family (24). Thus, the mechanisms by which ST or PP2A possibly modulate ferroptosis response warrant further investigation.

It was previously suggested that the expression level of mutant RAS, whether expressed from an endogenous locus or ectopically overexpressed, will have differential effects on ferroptosis sensitivity of the cell, depending ultimately on the net outcome of how cellular processes related to ferroptosis, including metabolism, redox and iron homeostasis, are balanced (23, 44, 45). This could plausibly explain why oncogenic RAS sensitizes cells to ferroptosis in some contexts, while in others protect cells from ferroptosis. However, our previous findings in endogenous KRAS cellular models and current findings in ectopic cellular KRAS and HRAS models, all demonstrably protect cells from FINs. An independent study similarly observed that rhabdomyosarcoma cells were protected against ferroptosis upon ectopic expression of various oncogenic RAS isoforms (46). In support of this, our survey of a large array of cancer cell lines (from the CCLE database) also revealed a significant association between oncogenic RAS signaling activation and ferroptosis resistance. Altogether, our evidence from various contexts are in accordance with a role of oncogenic RAS signaling as a ferroptosis suppressor, and highlights that the accepted view of ferroptosis as a synthetic lethal vulnerability of RAS-driven tumors needs to be revisited.

A major finding of our work is that oncogenic activation of the RAS signaling pathway induces GCH1 transcription to promote ferroptosis suppression. GCH1 is the rate-limiting enzyme in de novo synthesis of cellular BH4, converting GTP to dihydroneopterin phosphate (NH2TP), which is further processed to 6-pyruvoyltetrahydropterin (6-PTP) via 6-pyruvoyltetrahydropterin synthase (PTS), and finally producing BH4 via SPR (47). In turn, BH4 is an essential cofactor for a number of enzymes including nitric oxide synthases (NOSs) and tyrosine hydroxylase, critical for the synthesis of nitric oxide (NO) and dopamine respectively. More recently, BH4 itself was described to be a potent antioxidant with the ability to trap lipid radicals and prevent the propagation of LP chain reaction, as well as to selectively prevent oxidative degradation of polyunsaturated fatty acyl chain-containing phospholipids that drive ferroptosis execution (7, 8). Consistently, we found that *GCH1*-depletion re-sensitized KRAS-mutant cells to ferroptosis induction, while BH4 supplementation potently rescued this effect. In our cellular models, a direct radical-trapping effect of BH4 is likely, but an indirect effect of BH4 from its cofactor role cannot be completely ruled out. Intriguingly, BH4 reduction leads to uncoupling of NOSs, resulting in superoxide generation, which can promote ferroptosis. Conversely, NO, even though considered a radical species, paradoxically has been described to suppress ferroptosis by terminating LP propagation (48). Further, evidence suggest that GCH1 overexpression leads to an accumulation of the antioxidant CoQH2, since BH4 is also a cofactor for phenylalanine to tyrosine conversion, which can generate 4-OH-benzoate, a precursor for CoQH2 synthesis (7). It is therefore conceivable that BH4 generation downstream of oncogenic RAS signaling could also protect cells against ferroptosis via NOS- or CoQH2-dependent mechanisms. Of note, although the enhancement of GCH1-mediated BH4 synthesis is sufficient to suppress ferroptosis, oxidized BH4 (BH2) can also be

regenerated via the function of dihydrofolate reductase (DHFR) (8). Thus, whether DHFR serves a compensatory role for BH4 maintenance under GCH1-depleted conditions, and whether simultaneously targeting BH4 synthesis and DHFR using pharmacological inhibitors may be a viable therapeutic strategy against RAS-driven cancers, is likely to be a subject of future interest.

Interestingly, GCH1 was reported to play a role in promoting tumorigenesis. In particular, GCH1 was shown to facilitate the growth of glioblastoma in vivo, possibly by preserving the redox balance (31). In line with that, Cronin *et al.* recently demonstrated that GCH1 contributes to the development of non-small cell lung cancer NSCLC in a genetically-engineered mouse model driven by KRAS (32). In both studies, whether the pro-tumorigenic function of GCH1 is linked to the control of ferroptosis is unclear. While our data do not show that GCH1 is a determinant of tumorigenicity of KRAS-transformed fibroblasts in vitro and in vivo, this may be due to the lack of contribution from the tumor microenvironment or the absence of a fully functional immune system. Yet, we provide evidence that targeting GCH1 or SPR sensitizes cells to ferroptosis induction in KRAS-driven cancer models in vitro and in tumor organoids. Taken together, while targeting GCH1 alone could be a viable therapeutic strategy to target RAS driven tumors, our findings suggest that combining inhibitors of the BH4 pathway with FINs, such as IKE, may represent a more effective therapeutic approach to target these tumor types. Alternatively, combining RAS pathway inhibitors with FINs may prove to be an attractive therapeutic strategy since we and others have shown that repressing RAS signaling may sensitize cells to FINs, particularly in light of well-documented resistance to RAS inhibitors (49, 50).

In summary, this study uncovers that oncogenic RAS signaling transcriptionally regulates the GCH1-BH4 axis to mediate ferroptosis resistance. Along with our previous work, we thus demonstrate that oncogenic RAS signaling induces an extensive transcriptional program promoting the upregulation of three major arms of ferroptosis defense which also includes the xCT-glutathione axis and the FSP1-CoQH2 axis (22, 23). Consistently, our findings redefine the role of oncogenic RAS signaling to be a ferroptosis suppressor, which may help guide future strategies to maximize the clinical utility of FINs as next-generation therapeutic modalities against refractory and recurrent RAS-mutant tumors.

Materials and Methods

Cell culture

NIH 3T3 cells expressing mutant KRAS[G12V] and control MSCV were generated and cultured as previously described (22). BJ cells expressing human telomerase (hTERT), SV40 large T (LT) and mutant HRAS[G12V] (hTERT/LT/HRAS[G12V]) and control hTERT/LT were a kind gift from William C. Hahn (Dana-Farber Cancer Institute, Boston, U. S. A). H460 cells were a kind gift from Julian Downward (Francis Crick Institute, London, UK). MIA PaCa-2 cells were a kind gift from Barbara Grüner (German Cancer Consortium DKTK, Essen, Germany). HEK293T and CFPAC1 cells were purchased from American Type Culture Collection (Rockville, MD). Unless otherwise specified, all cell lines were maintained in DMEM (Gibco) supplemented with 10% fetal calf serum (FCS; Sigma Aldrich). All cells were kept at 37 °C with 5% CO2 and tested for mycoplasma at regular intervals.

Murine pancreatic tumor organoids treatment

Pancreatic tumor organoids were generated as previously described (23). Organoids were seeded at 100 cells/well, with or without SPRi3 (50 µM) treatment and allowed to grow for one week following which RSL3 (1 µM) treatments were added for an additional 48 hours. Organoid sizes were imaged and quantified using the BZ-H4M/Measurement Application Software (Keyence).

Full Materials and Methods available in Supplemental information

Acknowledgements

Research in the lab of G.L. was funded by the Deutsche Forschungsgemeinschaft (grant no. LE 3751/2-1), the German Cancer Aid (grant no. 70112624), the Elterninitiative Düsseldorf e.V. (grant no. 701910003), the Dr. Rolf M. Schwiete Stiftung (grant no. 2020-018) and the Research Commission of the Medical Faculty, Heinrich Heine University Düsseldorf (grant no. 2016-056 and 2020-044). J.K.M.L was supported by the Deutsche Forschungsgemeinschaft (Walter Benjamin Fellowship no. LI3844/1-1), the Dr. Rolf M. Schwiete Stiftung (grant no. 2023-029) and the Research Commission of the Medical Faculty, Heinrich Heine University Düsseldorf (grant no. 2022-12). Research in the lab of S.v.K was funded by: a collaborative research center grant on cell death (CRC1403, project ID 414786233), predictability in evolution (CRC1310, project ID 325931972), small cell lung cancer (CRC1399, project ID 413326622), B-cell lymphomas (CRC1530, project ID 455784452), through a priority program on ferroptosis (SPP2306, project ID 461704389) all funded by the Deutsche Forschungsgemeinschaft, an eMed consortium grant by the BMBF (InCa-01ZX1901A), via CANTAR which is funded through the program " Netzwerke 2021", an initiative of the Ministry of Culture and Science of the State of Northrhine Westphalia, Germany and a project grant (A06) funded by the center for molecular medicine cologne (CMMC). Research in the lab of F.B. was funded by the Deutsche Forschungsgemeinschaft (SFB1399 grant no. 41332662), the German Cancer Aid (grant no. 70113041) and the Mildred Scheel Nachwuchszentrum (grant no. 70113307). J.B. was funded by: a collaborative research center grant on small cell lung cancer (CRC1399 grant no. 413326622) by the Deutsche Forschungsgemeinschaft, via CANTAR which is funded through the program " Netzwerke 2021", an initiative of the Ministry of Culture and Science of the State of Northrhine Westphalia, Germany, a career advancement grant by the Center for Molecular Medicine Cologne (CMMC, CAP18) and the Mildred Scheel Nachwuchszentrum (grant no. 70113307). C.B. was funded by the Deutsche Forschungsgemeinschaft via GRK2578 (project ID 417677437).

Author contributions

Conceptualization, J.K.M.L. and G.L.; Methodology, J.K.M.L., F.S., T.L., L.T., D.P., S.T., H.Z., O.L., J.B., F.B.; Resources, T.R., M.R., C.B., S.v.K., G.R., B.R.; Writing – Original Draft, J.K.M.L and G.L.; Writing – Review & Editing, J.K.M.L., F.S., S.T., H.Z., T.R., F.B., S.v.K., G.R., B.R., and G.L.; Visualization, J.K.M.L. and G.L.; Supervision, J.K.M.L and G.L.; Funding Acquisition, J.K.M.L., B.R., G.R. and G.L.

Competing interests

J.B. has received a research grant by Bayer AG outside of the presented work. The other authors declare no potential competing interests.

References

1. S. J. Dixon, *et al.*, Ferroptosis: an iron-dependent form of nonapoptotic cell death. *Cell* **149**, 1060–1072 (2012).

2. S. J. Dixon, *et al.*, Pharmacological inhibition of cystine-glutamate exchange induces endoplasmic reticulum stress and ferroptosis. *Elife* **3**, e02523 (2014).

3. W. S. Yang, *et al.*, Regulation of ferroptotic cancer cell death by GPX4. *Cell* **156**, 317–331 (2014).

4. J. P. Friedmann Angeli, *et al.*, Inactivation of the ferroptosis regulator Gpx4 triggers acute renal failure in mice. *Nat Cell Biol* **16**, 1180–1191 (2014).

5. S. Doll, *et al.*, FSP1 is a glutathione-independent ferroptosis suppressor. *Nature* **575**, 693–698 (2019).

6. K. Bersuker, *et al.*, The CoQ oxidoreductase FSP1 acts parallel to GPX4 to inhibit
ferroptosis. *Nature* **575**, 688–692 (2019).

7. V. A. N. Kraft, *et al.*, GTP Cyclohydrolase 1/Tetrahydrobiopterin Counteract Ferroptosis
through Lipid Remodeling. *ACS Cent Sci* **6**, 41–53 (2020).

8. M. Soula, *et al.*, Metabolic determinants of cancer cell sensitivity to canonical ferroptosis
inducers. *Nat Chem Biol* **16**, 1351–1360 (2020).

9. H.-F. Zhang, R. I. Klein Geltink, S. J. Parker, P. H. Sorensen, Transsulfuration, minor
player or crucial for cysteine homeostasis in cancer. *Trends Cell Biol* **32**, 800–814
(2022).

10. L. Jiang, *et al.*, Ferroptosis as a p53-mediated activity during tumour suppression.
Nature **520**, 57–62 (2015).

11. Y. Zhang, *et al.*, BAP1 links metabolic regulation of ferroptosis to tumour suppression.
Nat Cell Biol **20**, 1181–1192 (2018).

12. S. Dolma, S. L. Lessnick, W. C. Hahn, B. R. Stockwell, Identification of genotype-
selective antitumor agents using synthetic lethal chemical screening in engineered
human tumor cells. *Cancer Cell* **3**, 285–296 (2003).

13. N. Yagoda, *et al.*, RAS-RAF-MEK-dependent oxidative cell death involving voltage-
dependent anion channels. *Nature* **447**, 864–868 (2007).

14. W. S. Yang, B. R. Stockwell, Synthetic lethal screening identifies compounds activating
iron-dependent, nonapoptotic cell death in oncogenic-RAS-harboring cancer cells. *Chem
Biol* **15**, 234–245 (2008).

15. I. Poursaitidis, *et al.*, Oncogene-Selective Sensitivity to Synchronous Cell Death
following Modulation of the Amino Acid Nutrient Cystine. *Cell Rep* **18**, 2547–2556
(2017).

16. J. L. Bos, ras oncogenes in human cancer: a review. *Cancer Res* **49**, 4682–4689 (1989).

17. J. Downward, Targeting RAS signalling pathways in cancer therapy. *Nat Rev Cancer* **3**,
11–22 (2003).

18. Y. Pylayeva-Gupta, E. Grabocka, D. Bar-Sagi, RAS oncogenes: weaving a tumorigenic
web. *Nat Rev Cancer* **11**, 761–774 (2011).

19. A. D. Cox, C. J. Der, The dark side of Ras: regulation of apoptosis. *Oncogene* **22**, 8999–
9006 (2003).

20. S. von Karstedt, *et al.*, Cancer cell-autonomous TRAIL-R signaling promotes KRAS-
driven cancer progression, invasion, and metastasis. *Cancer Cell* **27**, 561–573 (2015).

21. F. J. H. Hoogwater, *et al.*, Oncogenic K-Ras turns death receptors into metastasis-
promoting receptors in human and mouse colorectal cancer cells. *Gastroenterology* **138**,
2357–2367 (2010).

22. J. K. M. Lim, *et al.*, Cystine/glutamate antiporter xCT (SLC7A11) facilitates oncogenic
RAS transformation by preserving intracellular redox balance. *Proc Natl Acad Sci U S A*
116, 9433–9442 (2019).

23. F. Müller, *et al.*, Elevated FSP1 protects KRAS-mutated cells from ferroptosis during
tumor initiation. *Cell Death Differ* **30**, 442–456 (2023).

24. W. C. Hahn, *et al.*, Enumeration of the simian virus 40 early region elements necessary
for human cell transformation. *Mol Cell Biol* **22**, 2111–2123 (2002).

25. A. Tsherniak, *et al.*, Defining a Cancer Dependency Map. *Cell* **170**, 564-576.e16 (2017).

26. F. M. Behan, *et al.*, Prioritization of cancer therapeutic targets using CRISPR-Cas9
screens. *Nature* **568**, 511–516 (2019).

27. P. East, *et al.*, RAS oncogenic activity predicts response to chemotherapy and outcome
in lung adenocarcinoma. *Nat Commun* **13**, 5632 (2022).

28. H.-F. Zhang, *et al.*, Proteomic Screens for Suppressors of Anoikis Identify IL1RAP as a
Promising Surface Target in Ewing Sarcoma. *Cancer Discov* **11**, 2884–2903 (2021).

29. A. L. Kennedy, *et al.*, Activation of the PIK3CA/AKT pathway suppresses senescence
induced by an activated RAS oncogene to promote tumorigenesis. *Mol Cell* **42**, 36–49
(2011).

30. S. R. Hingorani, *et al.*, Preinvasive and invasive ductal pancreatic cancer and its early
detection in the mouse. *Cancer Cell* **4**, 437–450 (2003).

31. A. N. Tran, *et al.*, Reactive species balance via GTP cyclohydrolase I regulates
glioblastoma growth and tumor initiating cell maintenance. *Neuro Oncol* **20**, 1055–1067
(2018).

32. S. J. F. Cronin, *et al.*, Phenotypic drug screen uncovers the metabolic GCH1/BH4
pathway as key regulator of EGFR/KRAS-mediated neuropathic pain and lung cancer.
Sci Transl Med **14**, eabj1531 (2022).

33. Z. Wang, *et al.*, GTP Cyclohydrolase Drives Breast Cancer Development and Promotes
EMT in an Enzyme-Independent Manner. *Cancer Res* **83**, 3400–3413 (2023).

34. L. Badea, V. Herlea, S. O. Dima, T. Dumitrascu, I. Popescu, Combined gene expression
analysis of whole-tissue and microdissected pancreatic ductal adenocarcinoma identifies
genes specifically overexpressed in tumor epithelia. *Hepatogastroenterology* **55**, 2016–
2027 (2008).

35. R. Janky, *et al.*, Prognostic relevance of molecular subtypes and master regulators in
pancreatic ductal adenocarcinoma. *BMC Cancer* **16**, 632 (2016).

36. M. Ceccarelli, *et al.*, Molecular Profiling Reveals Biologically Discrete Subsets and
Pathways of Progression in Diffuse Glioma. *Cell* **164**, 550–563 (2016).

37. A. Méndez-Lucas, *et al.*, Identifying strategies to target the metabolic flexibility of
tumours. *Nat Metab* **2**, 335–350 (2020).

38. S. Cherradi, P. Martineau, C. Gongora, M. Del Rio, Claudin gene expression profiles and
clinical value in colorectal tumors classified according to their molecular subtype. *Cancer*
Manag Res **11**, 1337–1348 (2019).

39. A. L. Tarca, *et al.*, Strengths and limitations of microarray-based phenotype prediction:
lessons learned from the IMPROVER Diagnostic Signature Challenge. *Bioinformatics*
29, 2892–2899 (2013).

40. M. J. Hangauer, *et al.*, Drug-tolerant persister cancer cells are vulnerable to GPX4
inhibition. *Nature* **551**, 247–250 (2017).

41. V. S. Viswanathan, *et al.*, Dependency of a therapy-resistant state of cancer cells on a
lipid peroxidase pathway. *Nature* **547**, 453–457 (2017).

42. J. Tsoi, *et al.*, Multi-stage Differentiation Defines Melanoma Subtypes with Differential
Vulnerability to Drug-Induced Iron-Dependent Oxidative Stress. *Cancer Cell* **33**, 890-
904.e5 (2018).

43. R. P. Abrams, W. L. Carroll, K. A. Woerpel, Five-Membered Ring Peroxide Selectively
Initiates Ferroptosis in Cancer Cells. *ACS Chem Biol* **11**, 1305–1312 (2016).

44. J. K. M. Lim, G. Leprivier, The impact of oncogenic RAS on redox balance and
implications for cancer development. *Cell Death Dis* **10**, 955 (2019).

45. G. M. DeNicola, *et al.*, Oncogene-induced Nrf2 transcription promotes ROS
detoxification and tumorigenesis. *Nature* **475**, 106–109 (2011).

46. C. Schott, U. Graab, N. Cuvelier, H. Hahn, S. Fulda, Oncogenic RAS Mutants Confer
Resistance of RMS13 Rhabdomyosarcoma Cells to Oxidative Stress-Induced
Ferroptotic Cell Death. *Front Oncol* **5**, 131 (2015).

47. B. Thöny, G. Auerbach, N. Blau, Tetrahydrobiopterin biosynthesis, regeneration and
functions. *Biochem J* **347 Pt 1**, 1–16 (2000).

48. T. Homma, *et al.*, Nitric oxide protects against ferroptosis by aborting the lipid
peroxidation chain reaction. *Nitric Oxide* **115**, 34–43 (2021).

49. V. Bhatt, *et al.*, Inhibition of autophagy and MEK promotes ferroptosis in Lkb1-deficient
Kras-driven lung tumors. *Cell Death Dis* **14**, 61 (2023).

50. M. M. Awad, *et al.*, Acquired Resistance to KRASG12C Inhibition in Cancer. *N Engl J*
Med **384**, 2382–2393 (2021).

51. H. Thorvaldsdóttir, J. T. Robinson, J. P. Mesirov, Integrative Genomics Viewer (IGV):
high-performance genomics data visualization and exploration. *Brief Bioinform* **14**, 178–
192 (2013).

Fig. 1. Oncogenic activation of RAS signaling suppresses ferroptosis. (A) Cell death
quantification of NIH 3T3 cells expressing mutant KRAS[G12V] or control MSCV upon treatment
with ML162 (2 µM), ML210 (10 µM), FINO2 (20 µM), sulfasalazine (SSZ; 0.7 mM) or vehicle
for 24 h in the presence or absence of ferrostatin-1 (Fer-1; 5 µM), analyzed by propidium
iodide (PI) staining and flow cytometry. (B) Cell death quantification of BJ cells expressing
human telomerase (hTERT), SV40 large T (LT) and mutant HRAS[G12V] (hTERT/LT/HRAS[G12V])
or control hTERT/LT upon treatment with RSL3 (1 µM), ML162 (0.5 µM), ML210 (1 µM),
erastin (ERA; 5 µM) or vehicle for 24 h in the presence or absence of Fer-1 (5 µM), analyzed
as in (A). (C) Cell death quantification of shKRAS knockdown and scrambled (scr) control
CFPAC1 cells upon treatment with indicated concentrations of RSL3 for 24 h in the presence
or absence of Fer-1 (5 µM), analyzed as in (A) and (B). (D) Relative *KRAS* mRNA expression

in shKRAS knockdown and scr control CFPAC1 cells analyzed by qRT-PCR. (E) Correlation between *KRAS* mutation status and sensitivity to ML162 in cancer cells. *wt, wild-type. mut, mutant*. (F, G, H, and I) Correlation between *RAS* signature score and sensitivity to ML162, RSL3, ML210, and erastin in cancer cells. *RAS*^low and *RAS*^high cells are divided according to median AUC. (E-I) Data are obtained from the Cancer Therapeutic Response Portal v2 (CTRPv2) database, accessed via the DepMap portal. Higher AUC values indicate increased resistance to compound treatment. Data represent mean ± SD of three independent experiments. Unpaired, two-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Fig. 2. Oncogenic RAS signaling transcriptionally upregulates GCH1. (A) Volcano plots comparing transcriptomic (RNA sequencing) and proteomic profiles of NIH 3T3 cells expressing mutant KRAS^G12V versus control MSCV. (B) Schematic depicting GCH1-mediated tetrahydrobiopterin (BH4) biosynthesis. (C) Relative *Gch1* mRNA expression in NIH 3T3 cells expressing mutant KRAS^G12V and control MSCV analyzed by qRT-PCR. (D) Relative Gch1 peptide level in NIH 3T3 cells expressing mutant KRAS^G12V and control MSCV analyzed by mass spectrometry. (E) Relative *GCH1* mRNA expression in BJ cells expressing human telomerase (hTERT), SV40 large T (LT) and mutant HRAS^G12V (hTERT/LT/HRAS^G12V) or control hTERT/LT analyzed by qRT-PCR. (F) Relative *GCH1* mRNA expression in shKRAS knockdown and scrambled (scr) control CFPAC1 cells analyzed by qRT-PCR. (G) Relative *GCH1* mRNA expression in shKRAS knockdown and scr control H460 cells analyzed by qRT-PCR. (H) Relative *GCH1* mRNA expression analyzed by qRT-PCR and phosphorylated ERK (p-ERK) levels analyzed by Western Blot, in MIA PaCa-2 cells upon treatment with AMG510 (10 µM) or vehicle for the indicated times. (I) Relative *GCH1* mRNA expression in IMR90 cells expressing myristoylated AKT1 (myrAKT1), mutant HRAS^G12V, or control pBabe, analyzed by microarray. Data are obtained from GSE45276, accessed via the R2 platform (http://r2.amc.nl) (29). (J) Relative *Gch1* mRNA expression analyzed by qRT-PCR, and expression of indicated proteins analyzed by Western Blot in NIH 3T3 cells expressing mutant KRAS^G12V (NIH 3T3 KRAS^G12V) upon treatment with MEK inhibitor PD184352 (10 µM; MEKi), AKT inhibitor MK2206 (10 µM; AKTi) or vehicle for the indicated times. Data represent mean ± SD of three independent experiments. Unpaired, two-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Fig. 3. GCH1 mediates oncogenic RAS-driven ferroptosis suppression in 2D models. (A) Cell death quantification of NIH 3T3 cells expressing mutant KRAS^G12V or control MSCV upon treatment with indicated concentrations of erastin (Era) or vehicle for 24 h in the presence or absence of dihydrobiopterin (BH2; 50 µM), tetrahydrobiopterin (BH4; 50 µM), or ferrostatin-1 (Fer-1; 5 µM), analyzed by propidium iodide (PI) staining and flow cytometry. (B) Cell death quantification of NIH 3T3 cells expressing mutant KRAS^G12V or control MSCV upon treatment with indicated concentrations of FINO2 or vehicle for 24 h in the presence or absence of BH2 (50 µM), BH4 (50 µM), or Fer-1 (5 µM), analyzed as in (A). (C) Lipid peroxidation quantification of NIH 3T3 cells expressing mutant KRAS^G12V or control MSCV upon treatment with indicated concentrations of Era or vehicle for 6 h in the presence or absence of BH2 (50 µM), BH4 (50 µM), or Fer-1 (5 µM), analyzed by C11-BODIPY(581/591) staining and flow cytometry. (D) Lipid peroxidation quantification of NIH 3T3 cells expressing mutant KRAS^G12V or control MSCV upon treatment with indicated concentrations of FINO2 or vehicle for 6 h in the presence or absence of dihydrobiopterin BH2 (50 µM), BH4 (50 µM), or Fer-1 (5 µM), analyzed as in (C). (E) Cell death quantification of shKRAS knockdown CFPAC1 cells upon treatment with RSL3 (5 µM) for 24 h in the presence or absence of BH2 (50 µM) or BH4 (50 µM), analyzed as in (A) and (B). (F) Relative *Gch1* mRNA expression in CRISPR interference (CRISPRi)-mediated Gch1 knockdown (sgGch1-1 and sgGch1-2) and EGFP knockdown (sgEGFP) control NIH 3T3 cells expressing mutant KRAS^G12V (NIH 3T3 KRAS^G12V), analyzed by qRT-PCR. (G) Cell death quantification of sgGch1-1, sgGch1-2, and sgEGFP NIH 3T3 KRAS^G12V cells upon treatment with indicated concentrations of Era or vehicle for 24 h in the presence or absence of BH2 (50 µM) or BH4 (50 µM), or Fer-1 (5 µM), analyzed as in (A) and

622 (B). Data represent mean ± SD of three independent experiments. Unpaired, one-tailed *t test*;
623 *P < 0.05, **P < 0.01, ***P < 0.001. ns, not significant.

Fig. 4. GCH1 mediates oncogenic RAS-driven ferroptosis suppression in 3D models. (A)
Colony formation of CRISPR interference (CRISPRi)-mediated Gch1 knockdown (sgGch1-1
and sgGch1-2) and control (sgEGFP) NIH 3T3 cells expressing mutant KRAS^{G12V} (NIH 3T3
KRAS^{G12V}) grown in soft agar for 21 days and treated with the indicated concentrations of
RSL3. Colonies were imaged and quantified using ImageJ, and normalized to control. (B)
Colony formation of NIH 3T3 KRAS^{G12V} cells grown in soft agar for 21 days and treated with
SPRi3 or RSL3, alone or in combination. Colonies were imaged and quantified using ImageJ,
and normalized to control. (C) Pancreatic organoids were seeded with or without SPRi3 (50
μM) and allowed to grow for one week following which RSL3 (1 μM) treatments were added
for 48 hours. Organoids were imaged and quantified using the BZ-H4M/Measurement
Application Software (Keyence). Data represent mean ± SD of three independent
experiments. Unpaired, one-tailed *t test*; *P < 0.05, **P < 0.01, ***P < 0.001. ns, not significant.

Fig. 5. GCH1 expression is clinically relevant and correlates with oncogenic RAS activation in
human cancers. (A) Pan-cancer analysis of *GCH1* expression in tumor samples stratified
according to *KRAS* mutation status. *wt, wild-type. mut, mutant*. Data are obtained from The
Cancer Genome Atlas (TCGA), analyzed using the cBioPortal platform. (B, C, and D) *GCH1*
expression in tumor samples of pancreatic adenocarcinoma (PAAD), colon adenocarcinoma
(COAD), and lung adenocarcinoma (LUAD) from TCGA, in comparison to corresponding non-
tumor tissue from TCGA and/or Genotype-Tissue Expression (GTEx) database. Data are
analyzed using the UCSC Xena portal. (E, F, G, H, I, and J) Correlation of *GCH1* expression
and *RAS* signature score in tumor samples of COAD, rectal adenocarcinoma (READ), uterine
carcinosarcoma (UCS), skin cutaneous melanoma (SKCM), low-grade glioma (LGG), and
glioblastoma (GBM). Data are obtained from TCGA, accessed via the DepMap portal.
Correlations were quantified using Spearman's rank correlation coefficient. (K) Over-
representation analysis for Human Molecular Signatures Database (MSigDB) Hallmark gene
sets of genes positively co-expressed with *GCH1* in tumor samples of GBM from TCGA,
analyzed using the R2 platform. (L) *Gch1* expression in tumor samples from a mouse model
of high-grade glioma with *p53fl/fl;Nf1fl/fl* driven by Ascl1 Cre-ERTM, in comparison to non-tumor
tissue, obtained from GSE57036, accessed via the R2 platform. Data represent mean ± SD
of three independent experiments. Unpaired, two-tailed *t test*; *P < 0.05, **P < 0.01, ***P <
0.001. ns, not significant.

Fig. 6. Transcriptional upregulation of GCH1 by oncogenic RAS signaling is mediated by
ETS1. (A) Gene set enrichment analysis (GSEA) of NIH 3T3 cells expressing mutant
KRAS^{G12V} (NIH 3T3 KRAS^{G12V}) versus control MSCV showing enrichment for ETS1-regulated
genes. (B) Schematic depicting regulation of ETS1 downstream of RAS-MAPK signaling
pathway. (C) Relative *Gch1* mRNA expression analyzed by qRT-PCR and Ets-1 levels
analyzed by Western Blot in siRNA-mediated Ets-1 knockdown (siEts-1) and scrambled
(siScr) control NIH 3T3 KRAS^{G12V} cells. (D) Relative *GCH1* mRNA expression analyzed by
qRT-PCR and ETS1 levels analyzed by Western Blot in si-ETS1 knockdown (siETS1-1 and
siETS1-2) and siScr control H460 cells. (E) Schematic depicting the transcriptionally active
region (TAR) of *GCH1* and ETS1 binding sites. (F) Activity measurements of a reporter
construct consisting of the TAR of *GCH1* directly upstream of *Firefly luciferase* (FLuc; *GCH1-
Luc*), normalized to control *Renilla luciferase* (RLuc), upon co-expression with the indicated
amounts of *ETS1*-expressing plasmid in HEK293T cells, and ETS1 expression levels
analyzed by Western Blot. *A. U., Arbitrary units*. (G) Activity measurements of *GCH1-Luc* or
GCH1-Luc with mutations in individual *ETS1* binding sites (*GCH1^{mut}-Luc*-1, 2, or 3) or all ETS1
binding sites (*GCH1^{mut}-Luc*-all) normalized to RLuc, upon co-expression with 400 ng of *ETS1*-
expressing plasmid in HEK293T cells, and ETS1 expression levels analyzed by Western Blot.
(H) ChIP-qPCR analysis showing occupancy of Ets-1 at the *Gch1* locus in NIH 3T3 KRAS^{G12V}
cells. (I) ChIP-sequencing data showing ETS1, H3K4me3, and H3K27ac occupancy profiles
at the *GCH1* locus, in HepG2 cells. Data are obtained from UCSC Encode database, analyzed

using Integrative Genomics Viewer (IGV) (51). (J) Correlation between *GCH1* expression and ETS1 protein levels across multiple tumor entities from the Clinical Proteomic Tumor Analysis Consortium (CPTAC) and TCGA, analyzed using the cBioPortal platform. (K, L, and M) Correlation of *GCH1* expression and *ETS1* expression in tumor samples of PAAD, COAD, and lung adenocarcinoma (LUAD) obtained from GSE62165, GSE72970, and GSE43580 respectively, accessed via the R2 platform (35, 38, 39). Correlations were quantified using Spearman's rank correlation coefficient. Data represent mean ± SD of three independent experiments. Unpaired, two-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Fig. S1. (A) Cell death quantification of NIH 3T3 cells expressing mutant KRAS[G12V] or control MSCV upon treatment with the indicated concentrations of erastin (Era) or vehicle for 24 h in the presence or absence of ferrostatin-1 (Fer-1; 5 µM), analyzed by propidium iodide (PI) staining and flow cytometry. (B and C) Relative *GCH1* and *xCT* mRNA expression in BJ cells expressing human telomerase (hTERT), SV40 large T (LT) and mutant HRAS[G12V] (hTERT/LT/HRAS[G12V]) or control hTERT/LT analyzed by qRT-PCR. Data represent the mean ± SEM. (D, E, and F) Correlation between *KRAS* mutation status and sensitivity to RSL3, ML210, and erastin in cancer cells. *wt, wild-type. mut, mutant*. Data are obtained from the CTRP database, accessed via the DepMap portal. Data represent mean ± SD of three independent experiments. Unpaired, two-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Fig. S2. (A) *left,* Venn diagram depicting significantly altered transcripts and proteins (fold change > 2 or < -2; p < 0.05) in NIH 3T3 cells expressing mutant KRAS[G12V] relative to control MSCV based on RNA sequencing and proteomic profiling. *right,* Heatmap showing concordance in transcript and protein levels of 193 overlapping transcripts/proteins significantly altered in KRAS[G12V]-expressing cells. (B) Relative *KRAS* mRNA expression in shKRAS knockdown and scrambled (scr) control H460 cells analyzed by qRT-PCR. (C) Relative *Gch1* mRNA expression in NIH 3T3 cells expressing mutant KRAS[G12V] (NIH 3T3 KRAS[G12V]) upon treatment with MEK inhibitor PD184352 (10 µM; MEKi) or vehicle for the indicated times, analyzed by qRT-PCR. Data represent mean ± SD of three independent experiments. Unpaired, two-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Fig. S3. (A) Cell death quantification of sgGch1-1, sgGch1-2, and sgEGFP NIH 3T3 KRAS[G12V] cells upon treatment with indicated concentrations of FINO2 or vehicle for 24 h in the presence or absence of dihydrobiopterin (BH2; 50 µM) or tetrahydrobiopterin (BH4; 50 µM), or ferrostatin-1 (Fer-1; 5 µM), analyzed by propidium iodide (PI) staining and flow cytometry. (B) Ranking of correlation scores between GCH1 expression and sensitivity to CTRP compounds (median AUC). Data are obtained from the Cancer Therapeutic Response Portal v2 (CTRPv2) database, accessed via the DepMap portal. (C) Colony formation of CRISPR interference (CRISPRi)-mediated Gch1 knockdown (sgGch1-1 and sgGch1-2) and control (sgEGFP) NIH 3T3 cells expressing mutant KRAS[G12V] (NIH 3T3 KRAS[G12V]) grown in soft agar for 21 days. Colonies were imaged and quantified using ImageJ, and normalized to control. Data represent mean ± SD of three independent experiments. Unpaired, one-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Fig. S4. (A and B) *GCH1* expression in tumor samples of PAAD in comparison to non-tumor tissue, obtained from GSE15471 and GSE62165, accessed via the R2 platform (34, 35) (C) *Gch1* expression in tumor samples from a mouse model of liver cancer driven by human c-MYC overexpression, in comparison to non-tumor tissue, obtained from GSE129013, accessed via the NCBI Gene Expression Omnibus (GEO) platform (37). Data represent mean ± SD of three independent experiments. Unpaired, two-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Fig. S5. (A) Gene set enrichment analysis (GSEA) of NIH 3T3 cells expressing mutant KRAS[G12V] (NIH 3T3 KRAS[G12V]) versus control MSCV showing enrichment for target genes regulated by indicated transcription factors. (B) Correlation of *GCH1* expression and *ETS1* expression in tumor samples of pancreatic adenocarcinoma (PAAD) obtained from GSE15471

727   accessed via the R2 platform (34). Correlations were quantified using Spearman's rank
728   correlation coefficient. Data represent mean ± SD of three independent experiments.
729   Unpaired, two-tailed *t test*; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$. ns, not significant.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Figure 1

A

B

H460

C

NIH 3T3 KRAS^{G12V}

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5