Title

Quantifying the thermal tolerance traits of individual corals reveals widespread variation across the Great Barrier Reef

Running title

Variation in GBR corals heat tolerance

Authors

Hugo Denis[1,2,3], Line K. Bay[4], Véronique J.L. Mocellin[4], Melissa S. Naugle[3], Gaël Lecellier[1,5], Steven W. Purcell[3], Véronique Berteaux-Lecellier[1], Emily J. Howells[3]

Affiliations

1 UMR250/9220 ENTROPIE (IRD-CNRS-UR-IFREMER-UNC), Promenade Roger-Laroque, Noumea cedex, New Caledonia, France.

2 ED 129, SU Sorbonne Université, 4, Place Jussieu, 75252 Paris, France

3 National Marine Science Centre, Southern Cross University, Coffs Harbour, NSW, Australia

4 Australian Institute of Marine Science, Townsville, QLD, Australia

5 Institut de Sciences Exactes et Appliquées (ISEA) EA7484, 145, Avenue James Cook – BP R4 98 851 Nouméa, New Caledonia

Contact info

Hugo DENIS

hugo.denis@ird.fr

Keywords

coral bleaching, coral reefs, acute heat stress, intraspecific variation, phenotype-environment association, global warming.

Abstract

The capacity of reef-building corals to adapt to global warming depends upon standing heritable variation in tolerance traits upon which selection can act. While heat tolerance is known to differ markedly among coral taxa and reefs, limited knowledge exists on within-species variation among spatial scales ranging from meters to hundreds of kilometers. Here, we performed standardized acute heat stress assays to quantify the thermal tolerance of 768 colonies of *Acropora spathulata* from 14 reefs spanning 1060 km (9.5° latitude) of the Great Barrier Reef (GBR). Colony thermal tolerance was determined by the rate of decline in physiological traits under +3 °C, +6 °C and +9 °C above the local maximum monthly mean (MMM). Photochemical efficiency and chlorophyll retention traits revealed considerable variation in thermal thresholds among individual colonies both among reefs (~6 °C) and within reefs (~3 °C). Although tolerance rankings of colonies varied between traits, the most heat tolerant corals (i.e. within the top 25% of each trait) were found at virtually all reefs across the GBR, indicating widespread phenotypic variation. Reef-scale environmental predictors explained 12–62% of trait variation and heat tolerance was the greatest at reefs exposed to high thermal averages and recent thermal stress likely reflecting local adaptation and stress pre-acclimatization. We also found evidence that heat tolerance was further influenced by thermal fluctuations, ocean currents and cloud cover. Importantly, heat tolerance relative to local summer temperatures was the greatest in the southern GBR (~9.5 °C above MMM), suggestive of higher potential to adapt to future warming. Together, these results can be used to identify naturally tolerant coral populations and individuals for conservation and restoration applications.

Introduction

Global warming modifies species thermal environments (Masson-Delmotte et al., 2022) and directly impacts organism fitness (Angilletta et al., 2003; Tracy & Christian, 1986). The persistence of species and the ecosystem services they may provide (e.g., food and water supply, regulation of natural hazards, support for tourism; Costanza et al., 2014) therefore depends on their response to rising and extreme temperatures. Yet, considerable intraspecific variation in thermal tolerance exists in marine ecosystems (Bennett et al., 2019) as some species have colonized large geographical areas spanning contrasted thermal environments (Reusch, 2014). Species can adjust to local temperatures through physiological acclimatization of individuals within their lifespan through phenotypic plasticity, and/or selection for particular phenotypes within populations leading to changes in genetic variation across generations (Somero, 2010). Documenting phenotypic differences along thermal gradients can thus inform conservation and restoration efforts (Anthony et al., 2020; Bowden-Kerby, 2022) by improving predictions of species adaptive potential (Bennett et al., 2019), climate refugia (Lachs, Humanes, Mumby, et al., 2023) or extinctions (Sinervo et al., 2010). As such, thermal tolerance variation has been documented across terrestrial and marine taxa including plants (Reusch, 2014; Smith et al., 2017), reptiles (Herrando-Pérez et al., 2019), arthropods (Franken et al., 2018), birds (Charmantier et al., 2008; Noakes et al., 2016), marine invertebrates (Morley et al., 2012) and fishes (Fangue et al., 2006; McKenzie et al., 2021).

Reef-building (scleractinian) corals are among the organisms most at risk from global warming as they occupy narrow thermal niches in environments where current summer temperatures often approach their upper thermal limits and increasingly exceed them (Hughes et al., 2017). The energetic requirements of corals rely on a finely-tuned symbiotic relationship with photosynthetic dinoflagellates (family Symbiodiniaceae; LaJeunesse et al., 2018; Muscatine & Porter, 1977; Stat et al., 2006) that can be severely impaired under heat

stress (Hoegh-Guldberg et al., 2007). The loss of symbiont cells and/or their photosynthetic pigments caused by symbiophagy or symbiont exocytosis (Oakley & Davy, 2018; Roberty & Plumier, 2022) is termed 'coral bleaching'. Increasingly frequent mass bleaching events have been observed over the past 4 decades (Hughes, 2020; Hughes et al., 2021) and are predicted to have impacted > 70% of the world's coral reefs (Virgen-Urcelay & Donner, 2023), leading to widespread coral mortality (Bruno & Selig, 2007; De'ath et al., 2012; Gardner et al., 2003; Riegl et al., 2018). Understanding the extent of variation in coral heat tolerance across phylogeny, space, and time is therefore crucial to predict the future of these ecosystems (Cornwall et al., 2023; Dixon et al., 2022; Matz et al., 2020; Sully et al., 2022).

Intraspecific variation in coral heat tolerance exists at local scales (e.g., Cunning et al., 2016; Drury, Bean, et al., 2022; Drury, Marhoefer et al., 2021; Martin, et al., 2022; Jin et al., 2016; McWilliam et al., 2022; Sampayo et al., 2008) and is an indicator of adaptive capacity and metapopulation persistence probability (Holstein et al., 2022; Howells et al., 2022; Parkinson et al., 2015). Populations with low intraspecific variation may suffer near extirpation at local scales (Álvarez-Noriega et al., 2023; Hughes et al., 2018; Riegl et al., 2018). On the contrary, high intraspecific variation in physiological traits can buffer populations against the loss of genetic and functional diversity (McWilliam et al., 2022). Coral thermal tolerance traits are shaped by thermal history (Hughes et al., 2019; Schoepf et al., 2021; Thompson & Van Woesik, 2009) and other environmental conditions (e.g., irradiance (Gonzalez-Espinosa & Donner, 2021), hydrodynamic regimes (Page et al., 2019), nutrients (Wiedenmann et al., 2013) and water oxygen content (Alderdice et al., 2022). Coral colonies can adjust to these variable conditions through phenotypic plasticity (DeCarlo et al., 2019; Hackerott et al., 2021), heritable genetic effects (Bairos-Novak et al., 2021; Dixon et al., 2015; Howells et al., 2021; Jin et al., 2016; Kirk et al., 2018; Rose et al., 2018; Rose et al., 2021), differences in symbiont communities (Berkelmans & Van Oppen, 2006; Howells et al., 2012; Sampayo et al., 2008) or a combination of these three factors (Howells et al., 2016; Thomas et al., 2018).

Yet, further research is required to understand the relative importance of these drivers — especially micro- and macro-scale environmental conditions — across species and regions.

Quantifying intraspecific variation in thermal thresholds during natural marine heatwaves is challenged by differences in heat stress magnitude and duration over space and time and the technical constraints of surveying large spatial areas during a single heatwave event. Consequently, controlled experiments have been widely used to characterize intraspecific variation in thermal tolerance among reefs (Cornwell et al., 2021; Evensen et al., 2022; Marzonie et al., 2022; Shaw et al., 2016; Voolstra et al., 2021), habitats (Klepac & Barshis, 2022; Marhoefer et al., 2021; Thomas et al., 2018; Voolstra et al., 2020), and individuals (Humanes et al., 2022; Kavousi et al., 2020; Walker, Nestor, et al., 2022). Recently, acute heat stress assays (< 48 h) have been developed to measure individual thermal thresholds in a standardized, high-throughput and cost-effective manner (Barshis et al., 2013; Cornwell et al., 2021; Cunning et al., 2021; Evensen et al., 2021; Evensen et al., 2022; Klepac et al., 2023; Marzonie et al., 2022; Morikawa & Palumbi, 2019; Voolstra et al., 2021; Voolstra et al., 2020), yielding consistent results across weeks (Walker, Cornwell, et al., 2022), seasons (Cunning et al., 2021) and with long-term heat stress (Evensen et al., 2021; Voolstra et al., 2020; but see Klepac et al., 2023). In these experiments, coral heat tolerance has been mostly inferred from photosynthetic efficiency (F_v/F_m), a common proxy of bleaching onset (Jones et al., 2000; Schoepf et al., 2015). However, the role of symbiont photochemical damage in bleaching initiation has been increasingly challenged (Gardner et al., 2017; Krueger et al., 2015) and can be offset by host protective mechanisms (Baird et al., 2009; Oakley & Davy, 2018; Roberty & Plumier, 2022), stressing the importance of relating F_v/F_m measurements to subsequent coral bleaching responses (i.e., the loss of symbiont cells and/or chlorophyll). To date, acute heat stress assays have investigated the response from a few colonies at many reefs or many colonies at a single reef, but have not yet evaluated intraspecific variation across multiple spatial scales.

To address the aforementioned knowledge gaps, we used acute heat stress assays to measure photochemical efficiency and bleaching traits in 768 colonies of *Acropora spathulata* from 14 reefs spanning 1,060 km (~9.5 degrees of latitude) of the Great Barrier Reef (GBR). *A. spathulata* is an abundant corymbose species found on reef flats and upper slopes that builds ecologically important structural complexity on the GBR (Cooper et al., 2011; Wallace et al., 2012). We identify reefs where heat tolerant individuals are most likely to be found and use multiple environmental datasets (i.e., satellite observations, numerical models, in-situ loggers) to determine environmental predictors of tolerance traits. The large sample size of this study provides GBR-wide information to inform conventional management via spatial and temporal protection and novel interventions, including seeding heat-tolerant corals onto reefs.

Material & Methods

Study sites, experimental design and setup

Study sites and sample collection

We quantified thermal tolerance traits of 768 colonies of *A. spathulata* across 14 reefs of the GBR (Figure 1a, Table S1). At each reef, 40–60 colonies were sampled on SCUBA from one or two sites (covering 50–4200 m^2) between 28 February and 26 March 2022 (GBRMPA permit G21/45166.1) over a single or two consecutive days. For each colony, 12 nubbins (~8 cm) were collected for standardized acute heat stress experiments and *in-situ* metadata was recorded. This included colony-level GPS coordinates linked to *in-situ* photographic records (Lukoschek et al., 2016), time-corrected depth (-0.54 ± 0.76 m relative to the lowest astronomical tide) and visual pigmentation (Coral Watch Health Chart scores at 0.5 intervals; Siebeck et al., 2006). Colonies showing signs of disease were rarely encountered but intentionally excluded to avoid physiological bias unrelated to temperature effects.

Figure 1. Location of sample sites and acute heat stress design to quantify variation in the heat tolerance of the reef-building coral *Acropora spathulata* on the Great Barrier Reef. **a.** Map of the 14 reefs sampled across a latitudinal and thermal gradient. Isobaths of 0, 200, 1,000 and 2,000 m were obtained from the naturalearth 10 m resolution dataset (Kelso & Patterson, 2010) and are represented in shades of blue. Coral reef locations were obtained from the UNEP database (UNEP-WCMC, 2010). Sites labels are colored according to their maximum monthly mean (MMM) (dark red: > 28.5 °C, blue: < 27.5 °C, Table S1). **b.** A total of 768 coral colonies were measured in standardized acute heat stress experiments. 12 nubbins per colony were randomly divided among 3 tanks for each of 4 temperature treatments relative to the local MMM of each reef. **c.** Standardized target temperature profiles to elicit increasing levels of heat stress, following (Evensen et al., 2021). Arrows indicate the timing of physiological measurements of the maximum quantum yield of photosystem II (F_v/F_m) and hyperspectral imagery (Hyp). **d.** Representative image of *A. spathulata* at Fitzroy Reef (© Mila Grinblat 2022).

Acute heat stress assay design

Acute heat stress assays were conducted on board a research vessel the day following collection. The experimental design consisted of three replicate tanks for each of four temperature treatments of 0 °C (control), 3 °C, 6 °C, and 9 °C above the local Maximum Monthly Mean (MMM, 1985–1990+1993 climatology as defined by NOOA; Liu et al., 2003)) for each reef site. These treatments were designed to elicit increasing levels of thermal stress following Voolstra et al. (2020). Immediately after collection, 12 nubbins from each coral colony were mounted on separate experimental racks with unique identifiers and held

overnight at ambient seawater temperature. The following morning, one nubbin from each colony was randomly assigned to each of the 12 tanks (Figure 1b). Assays started at 11:00 am and consisted of 3 hours ramp-up from ambient temperature (within 1 °C of the MMM) to target temperature, 3 hours hold at the target temperature, 2 hours ramp down to MMM, and a final ~12-h hold at MMM (Figure 1c). During the experiments, temperatures in individual tanks were recorded at a 1-min intervals using Hobo loggers (Onset) and closely matched their target profiles (mean$_{\Delta T}$ = 0.36 °C, Table S2) with the exception of one site (St Crispin: mean$_{\Delta T}$ = 1.61 °C).

Automated experimental system

A portable experimental system ("Seasim-in-a-box", National Sea Simulator, Australian Institute of Marine Science; Fig S1) was modified from Marzonie et al. (2022) to consist of replicate 50-L acrylic tanks outfitted with insulating water jackets and supplied with 0.8 L.min^1 unfiltered seawater directly pumped from the ocean (Iwaki MX400). Each treatment sumps, with recirculating water supply to the insulating water jackets, were brought from ambient to target temperatures by heating elements (Omega 2kW titanium). In parallel, four distinct flow-through seawater were heated by convection via their titanium coil (Wateco 56″ titanium heat exchanger) placed in the sumps, and pumped to the replicate experimental tanks using submersible pumps (Reefe RP2400LV). Tanks were equipped with independent white/blue LED lighting (300 W) to supply ~ 300 µmol photons m^{-2}. s^{-1} with a diurnal light regime timed to *in-situ* conditions (11:13). Each tank was equipped with circulators (Tunze 6015), independent temperature probes (TC Direct PT-100) and PAR sensors (Skye Quantum PAR – SKL26250) to control temperature and lighting. The monitoring and control of temperature and lighting were achieved by a system consisting of a Programmable Logic Controller (Siemens Simatic 571500), a Weidmuller UR20 Remote IO Signal Input and Outputs and a Human Machine Interface (HMI TP700 Comfort panel)

Phenotypic traits

Photochemical efficiency

The maximum quantum yield of photosystem II (F_v/F_m) of coral photosymbionts (Symbiodiniaceae) was used as an initial rapid, non-invasive measure of heat tolerance. A decline in F_v/F_m reflects early physiological impairment in corals and has been reported to respond consistently between acute- and long-term heat exposures (Evensen et al., 2021; Voolstra et al., 2020; but see Klepac et al., 2023). Measurements were taken 2 hours after the end of temperature ramp-down (> 1-hour dark adaptation) using an Imaging PAM chlorophyll fluorometer (IMAG-K7, Walz Germany) with the following settings: Measuring Light = 2 (freq = 1), Saturation Pulse = 7 (Int = 30s), Damp = 1, Gain = 1. For each fragment, three measurements were extracted from non-overlapping areas and underwent several quality check and filtration steps (Supplementary materials and methods). After filtration, a total of 769 colonies and 8,453 fragments were retained in further analyses.

Hyperspectral image-based assessment of chlorophyll content

Non-invasive estimates of chlorophyll content were used as a second metric of heat tolerance. The loss of chlorophyll underpins visual bleaching scores commonly used in acute heat stress experiments and has been shown to correspond with mortality risk (Walker, Cornwell, et al., 2022). Total chlorophyll content was assessed in fragments from reflectance measurements taken 11 hours after the end of the temperature ramp-down using a hyperspectral camera (Resonon, Pika XC2) with the following settings: Integration time = 39.2ms, Gain = 0, Frame rate = 22 fps. MATLAB software was used to compute the normalized difference vegetation index (NDVI, Supplementary material and methods) from reflectance measurements, where NDVI = $(R_{720}-R_{670})/(R_{720}+R_{670})$ and R_x is the reflectance at x nm. NDVI is used in a wide range of organisms including scleractinian corals (Rocha et al., 2013; Wijgerde et al., 2014) and has been validated as an accurate proxy of chlorophyll content in the soft coral *Sarcophton* cf. *glaucum* (Leal et al., 2015). We repeated this

validation in *A. spathulata* using spectrophotometric determination from tissue extractions (Ritchie, 2008) with a strong relationship between NDVI and log-transformed chlorophyll-a (R^2=0.74, Supplementary material and methods, Figure S2).

Phenotypic data analysis

For each trait — the maximum quantum yield (F_v/F_m) and chlorophyll content (NDVI) — we calculated two colony-level metrics of heat tolerance. The temperature that resulted in a 50% decline in response (i.e., the median effective dose, or ED50) which represents an absolute thermal threshold, and the performance retention under extreme heat (+9 °C), which represents how coral traits decline relative to their local reef MMM.

The response of raw F_v/F_m and NDVI to temperature across experiments was assessed using a linear mixed effect model incorporating fixed effects of site, treatment, fragment size (estimated from pixel numbers in RGB images) and random intercepts for each colony and tank using package *lme4* (Bates et al., 2014), including 95% bootstrap confidence intervals for fixed effects (Table S3). Tanks were first treated as a block factor in the analysis. Despite small variations in temperatures among replicate tanks, tank had a minor effect on fragment performances under heat stress (explained 1.44% of F_v/F_m random variance σ_i^2 and 0.8% of NDVI σ_i^2, linear mixed effect model, Table S3). Likewise, fragment size effect appears to be negligible for F_v/F_m (B = 1.4 × 10^{-6}, df = 7473, 95% CI: -4.8 × 10^{-7}–3.0 × 10^{-6}) and very weak for NDVI (B = 8.5 × 10^{-6}, df = 8988, 95% CI:6.14 × 10^{-6}–1.0 × 10^{-5}).

ED50 thermal thresholds

For each colony, the decrease in F_v/F_m and NDVI was modeled over temperature treatments using dose response curves in the R package *drc* (Ritz et al., 2015). Colonies with <2 fragments per treatment after previous quality control filtering (Supplementary material and methods) were excluded to ensure robust estimates of individual responses across treatments (excluded colonies: F_v/F_m = 103, NDVI = 4). All models were fitted using the *drm* function based on the mean hold temperature recorded within each tank for each

experiment, with constraints set on parameters (Table S4a). The best model (Weibull type II with 3 parameters) was chosen as the one with the lowest Akaike's information criterion score (AIC) for most reef sites using the *mselect* function (6/14 and 11/14 reefs for F_v/F_m and NDVI, respectively; Table S4b). Because using the same model is a prerequisite to compare ED50s, we used this model for all colonies even though some curves showed a better fit to log-logistic or quadratic models. The model equation is

$$f(x; b, d, e) = d(1 - \exp(-\exp(b(\log(x) - \log(e)))))$$

where *b* is steepness of the curve, *d* the upper asymptote (lower asymptote set to 0) and *e* the inflexion point. An example of dose response curves can be found in Figure S3. Dose response curves were filtered to remove individuals that showed poor fit to the data, notably when the decline in phenotypic traits was minor or absent up to +6 °C or +9 °C resulting in ED50s with wide confidence intervals (Supplementary material and methods). After filtration 652 and 725 colony-level ED50 estimates were retained for F_v/F_m and NDVI respectively.

Performance retention under thermal stress

Performance retention under extreme heat was computed as

$$Xi_{MMM+9/MMM} = (Xi_{MMM+9}/Xi_{MMM}) \, x \, ((Tmean_{MMM+9C} - Tmean_{MMM})/9)$$

where Xi is the trait (F_v/F_m or NDVI) averaged across fragments of individual i in the MMM and MMM+9 treatments, respectively, and $Tmean_{MMM+9C}$ and $Tmean_{MMM}$ are the average hold temperatures for these treatments. The second term accounts for small differences between target and effective temperatures across tanks and experiments. Performance retention can be computed even for colonies that experienced minor decline in phenotypic traits under +9 °C which prevented fitting a dose-response curve. After filtration of colonies with < 2 fragments in the MMM and MMM+9C treatments, 730 and 765 colony-level performance retention estimates were calculated for F_v/F_m and NDVI respectively.

Heat tolerance variation analysis

The variation in heat tolerance (ED50 and performance retention) among and within sites was investigated using an analysis of variance (ANOVA). Homogeneity of variance between sites was assessed using Levene's test on residuals from groups medians (Table S5b) and Tukey's HSD post-hoc tests for pairwise comparisons (Table S5c). Since variance was unequal across sites, we used one-way Welch's ANOVA to compare site means and Games-Howell post-hoc tests with adjusted *p*-values at α = 0.05 for pairwise comparisons (Table S5d). Following Cunning et al. (2021), we assessed within-site variation in heat tolerance by computing ED50 adjusted as the grand mean of the total distribution plus residuals from site. Finally, we investigated the similarity of colony thermal tolerance rankings between F_v/F_m and NDVI traits using Spearman's rank correlation as assumption of normality was violated. For both ED50 and performance retention, we computed both GBR-wide and site-level correlations (St Crispin Reef excluded; Table S6).

Environmental data acquisition and analysis

To examine environmental conditions experienced by *A. spathulata* across the GBR, we retrieved 10 environmental variables characterized by 24 predictors computed from *in-situ* loggers, numerical models, and satellite observations (Table S7). Reef thermal history was characterized using the NOAA Coral Watch v3.1 sea surface temperature (SST) satellite product (a.k.a CoralTemp; NOAA Coral Reef Watch, 2018, updated daily, 5km, 1985–present) for thermal predictors based on historical climatology (MMM, Degree Heating Weeks (DHW)) as it has the largest temporal coverage. For contemporary thermal predictors (e.g., annual average and thermal range, 2014–present) we used the eReefs GBR1 Hydro model (CSIRO, 2020a) as it delivers temperature predictions at a finer spatial resolution (~1 km, hourly) and higher accuracy for the last decade (ME = 0.4 °C–0.9 °C; Steven et al., 2019). Temperature time-series from both sources were validated using *in-situ* records (Supplementary material and methods).

Global averages (2010–2019) of environmental variables that may play a secondary role in heat tolerance (e.g., chlorophyll, turbidity, current velocity) were retrieved from eReefs GBR4 model (CSIRO, 2020b; hourly, 4 km). In addition, we included the average cloud fraction (% of pixel masked by clouds) and cloud fraction anomaly for the 30 days prior to collection (Gonzalez-Espinosa & Donner, 2021) retrieved from NASA-MODIS 1°, monthly database (Savtchenko et al., 2004).

Our environmental dataset also included variables measured at the individual colony level. Depths obtained from dive computers were adjusted to tide level at the time of collection (BOM, 2022) (standardized to Lowest Astronomical Tide, Figure S7). All predictors derived from eReefs were computed for the depth of each colony through linear interpolation from the model vertical levels. As the GBR experienced a marine heatwave during the sampling campaign (0.47–4.67 Degree Heating Weeks), we included pigmentation levels prior to collection as an explanatory variable to account for heat stress experienced in the weeks preceding the experiment (Figure S8).

Principal Component Analysis (PCA) was performed on environmental predictors for the 768 colonies to visualize the distribution of sites and colonies along environmental gradients.

Phenotype by environment analysis

The influence of environmental predictors on heat tolerance traits (ED50 and performance retention in F_v/F_m and NDVI) was evaluated using random forest ensemble learning and ridge regression on a set of 12 low to moderately correlated predictors (pairwise absolute Pearson correlation = 0.01–0.69; Figure S9, Table S9; St Crispin Reef excluded). Because any association of heat tolerance traits with MMM and DHW at the time of collection may reflect different mechanisms (e.g., adaptation vs acclimatization), both were retained in the dataset despite their strong correlation ($R = 0.78$).

Random forest models were built using *cforest* function from *party* R package (Hothorn et al., 2015). For each of the 4 heat tolerance metrics, separate random forests were grown to

1,000 trees (ntree) with 5 environmental predictors tried at each split (mtry). Predictor importance and model accuracy were assessed through repeated 10-fold stratified (across sites) cross-validation (70/30 split, Supplementary material and methods). The importance of each predictor was estimated by computing the marginal increase in out-of-bag sample MSE (Mean Inc MSE) when training the model with the predictors randomly shuffled.

Ridge regression was performed for each phenotypic trait using the *glmnet* R package (Friedman et al., 2021). The optimal tuning parameter λ of the penalty term was selected using k-fold cross-validation (λ.1se; Figure S11 and Figure S12). Model accuracy (R^2) and environmental predictor coefficients were estimated using bootstrap resampling (repeated 100-fold stratified cross validation, Figure S13 and Table S10).

All statistical analysis were conducted using R v4.0.4 (R CoreTeam, 2020) and figures created using package *ggplot2* (Wickham, 2011) and Inkscape v1.2 (Bah, 2011). Scripts and data used for the analyses are available online: https://github.com/hde08/GBR-HeatTolVar-GCB/tree/main.

Results

Under acute heat stress, the majority of *A. spathulata* colonies on the GBR showed minimal decline in photochemical efficiency (F_v/F_m) and chlorophyll content (NDVI) at +3 °C (≤ 0.5%) and +6 °C (≤ 2.5%) above their local MMM. However, under the more extreme temperature of +9 °C, these traits declined by 2–80% (F_v/F_m) and 10–98% (NDVI), driving differences in thermal threshold (ED50) and performance retention metrics among reefs and individuals (Figure 2).

Reef-level variation in coral heat tolerance

The GBR-wide medians of colony thermal thresholds were 37.1 ± 1.0 °C for F_v/F_m ED50 (median ± SD) and 36.1 ± 0.8 °C for NDVI ED50, with reef accounting for 34% and 25% of

their respective total variation (ω^2 effect sizes, Table S5b). Mean reef-level F_v/F_m ED50s differed by 1.8 °C and generally decreased with latitude (Figure 2a). For example, F_v/F_m ED50 thresholds in the southern GBR (Heron Island and Fitzroy Reef; 36.5–36.6 °C) were 0.8 °C lower than all reefs in the central and northern GBR (36.8–38.3 °C; *p.adj* < .0001; Table S5d). However, several reefs had higher (Kelso Reef, Pelorus Island, No Name Reef) or lower (Fitzroy Island, North Direction Island) ED50s than nearby reefs at similar latitudes (*p.adj* < .05). Conversely, NDVI ED50 thresholds were less variable among reefs (by 1.3 °C) and were lower at Martin Reef in the northern GBR than all other reefs (*p.adj* < .01; Figure 2c). At St Crispin Reef, both F_v/F_m ED50 and NDVI ED50 were 0.8–2.6 °C lower than all other reefs but are likely to be negatively biased due to experimental treatments not reaching target temperatures (Table S2).

GBR-wide performance retention (performance under +9 °C relative to local conditions) was 46 ± 12% for F_v/F_m and 25 ± 19% for NDVI with reef accounting for 44% and 59% of their respective total variation (ω^2 effect sizes, Table S5b). Performance retention was 2–41% (F_v/F_m) and 60–65% (NDVI) higher at southern reefs than the rest of the GBR (Figure 2b and Figure 2d), thus following an opposite trend to F_v/F_m ED50.

Figure 2. Variation in *Acropora spathulata* thermal tolerance among 14 reefs on the Great Barrier Reef (GBR) under acute experimental heat stress. The decline in maximum quantum yield (F_v/F_m) and total chlorophyll content (NDVI) was used to measure ED50 temperatures (a, c) and performance retention under extreme heat (i.e., +9 °C/MMM; b, d). Left panels display boxplots of the median (solid line) and first and third quartiles (box) for each reef colored by their MMM. Right panels show the distribution of values, with the GBR-wide median and standard deviation. Reefs are ordered (left to right) by increasing latitude. * Values for St Crispin Reef are negatively biased due to experimental treatments not reaching target temperatures.

Colony-level variation in coral heat tolerance

At the colony-level, thermal thresholds varied by up to 6°C, approximately 3 times the reef-level variation (1.8 °C, Figure S4). Within reefs, there was a 3.3 °C and 3.0 °C average range in colony-level ED50s for F_v/F_m and NDVI, respectively. The magnitude of this variation differed among reefs (p < 0.001, Levene's test; Table S5b) but no specific reef exhibited

higher or lower variation for both traits. Colony-level variation in performance retention under +9 °C was also higher than reef-level variation (by 48% for F_v/F_m and by 17% for NDVI). Although colony-level ED50s were prone to uncertainty (median 95% CI range = 2.51 ± 3.34 °C and 2.99 ± 5.56 °C, Supplementary material and methods), congruent patterns between ED50 and performance retention support high variation in acute heat tolerance among and within reefs of the GBR.

Importantly, heat tolerant colonies were widely distributed across the GBR. For example, colonies that were ranked within the top 25% of F_v/F_m ED50 and NDVI ED50 occurred at virtually all reefs (Figure 3a, b). However, the most tolerant colonies were generally more abundant at central and northern reefs for F_v/F_m and southern reefs for NDVI. Additionally, several reefs exhibited a high proportion of tolerant individuals for both traits (e.g., Mackay Reef, Moore Reef, and Pelorus Island; Figure 3).

Response to acute heat stress differs between traits

The ranking of reef- and colony-level responses under heat stress differed between phenotypic traits. Overall, there was no GBR-wide correlation between F_v/F_m and NDVI ED50s (ρ = 0.00, Figure 3c), but a moderate positive correlation between their performance retention at +9 °C (ρ = 0.43, Figure 3d). At the reef-level, there was a moderate positive correlation between F_v/F_m ED50 and NDVI ED50 at Chicken and No Name (ρ > 0.3) and a weak positive correlation at Davies, Heron, Mackay, Martin and Moore (ρ = 0.1–0.3, Table S6). Similarly, there was a weak to moderate positive correlation between the performance retention of F_v/F_m and NDVI at all reefs (ρ = 0.1–0.3) except Chicken (ρ = 0.02), North Direction (ρ = 0.84) and Pelorus Island (ρ = -0.22; Table S6)

Figure 3. Spatial distribution of *Acropora spathulata* acute heat tolerance on the Great Barrier Reef measured by **a.** maximum quantum yield (F_v/F_m) and **b.** chlorophyll content (NDVI). Pie charts at each reef represent the proportion of individual colony ED50s falling between each quartile of the global distribution (blue: ED50 < Q1, yellow: Q1 < ED50 < Q2, orange: Q2 < ED50 < Q3, red: Q3 <ED50). Pie charts radii are proportional to average retained performance under extreme heat (i.e., +9 °C/MMM), normalized by the global range for plotting purposes. Correlation between **c.** F_v/F_m and NDVI ED50s and **d.** retained performance under extreme heat (+9 °C/MMM). Each point represents a distinct colony (n = 652 and 730 for c and d, respectively) and Spearman's-ρ and *p*-values are indicated.

Environmental drivers of acute heat tolerance metrics

The Principal Component Analysis on 24 quantitative environmental predictors divided the 768 colonies into three major clusters (Figure 4). The first component (46.1% explained variance) separated cooler southern reefs from warmer central and northern sites and was mainly driven by MMM (1.7 °C range) and DHW at the time of collection. The second

component (15.1% explained variance) separated northern and central sites along an inshore-offshore gradient that was mainly driven by turbidity, sea surface current velocities, and temperature variation (annual range and the rate of change from spring to summer). Within clusters, colonies grouped by reefs as most predictors were retrieved at the site level and were minimally separated by their water depth or pigmentation. Pigmentation scores prior to collection differed between individuals and sites but had no clear association with DHW at the time of collection (Figure S8).

Environmental predictors explained a moderate to large proportion of variation in heat stress responses of *A. spathulata* across the GBR (R^2 = 0.12–0.62). The predictive accuracy of random forest (RF) and ridge regression (RR) was stronger when heat tolerance was expressed for both F_v/F_m and NDVI as performance retention (R^2 = 0.27–0.62) than ED50 thresholds (R^2 = 0.12–0.31). Top environmental predictors slightly differed among traits and metrics but variation in heat tolerance was primarily associated with site-level thermal history (Figure 5). MMM (Figure 5b) and DHW at time of collection (Figure 5d) were predominant metrics associated with response to acute heat stress (i.e., top RF predictors for 3 of the 4 heat tolerance metrics, Table S10a) while being strongly correlated to each other (R = 0.78). The direction of associations differed between traits with MMM and DHW at the time of collection being positively associated with F_v/F_m ED50 (β = 0.06–0.07), but negatively with performance retention in F_v/F_m and NDVI and NDVI ED50 (β = -0.03–-0.06; Table S10b). Secondary thermal history associations occurred between the annual range in temperature and both performance retention metrics (F_v/F_m β = -0.03, NDVI β =-0.04; Figure 5c), and between the frequency of DHW > 4 and both ED50 thresholds (F_v/F_m ED50 β = 0.04, NDVI ED50 β = -0.06; Figure 5e). Particularly, the frequency of DHW > 4 in the year before collection showed a strong negative association with NDVI ED50 (β = -0.060, top predictor in RF), but no significant association with any other trait metric.

To a lesser extent, the heat tolerance of *A. spathulata* was associated with water flow, shading, and depth. F_v/F_m tolerance metrics were generally higher at reefs with high sea

surface current velocity (F_v/F_m ED50 β = 0.042, +9/MMM β = 0.037) and lower at sites with positive cloud anomalies in the 30 days prior to collection (F_v/F_m ED50 β = -0.017, +9/MMM β = -0.031; Figure 5a). Turbidity showed little to no relationship with heat tolerance metrics. At the colony level, increasing water depth was positively related to F_v/F_m metrics (F_v/F_m ED50: β = 0.042, +9/MMM: β = 0.004), but negatively related to NDVI metrics (NDVI ED50: β = -0.028, +9/MMM: β = -0.015). In addition, colonies with higher levels of pigmentation at the time of collection tended to have higher NDVI ED50 thresholds (β = 0.028), but there was no relationship with other tolerance metrics.

Figure 4. Principal component analysis (PCA) conducted on 24 quantitative environmental predictors of *Acropora spathulata* heat tolerance metrics (*n* = 768 colonies sampled across the Great Barrier Reef, colored by their reef of origin). Loadings projected on the first two components are depicted with arrows, and labels are shown for moderately correlated predictors (|R| < 0.7) used in further analyses. CF30a = Cloud fraction anomaly for 30 days prior collection, DHW_collection = Degree Heating Weeks (DHW) at the day of collection, DHW_freq_sup4 = Frequency of DHW > 4, DHW_freq_sup4_1year = Frequency of DHW > 4 in the year prior collection, MMM = Maximum Monthly Mean (1985–1990+1993 climatology), ROTC_ss = Rate of temperature change in spring/summer, TSA_DHW_stdev = standard

deviation of DHW. Numbers in parentheses represent the proportion of variance explained by each principal component.

Figure 5 Environmental drivers of acute heat tolerance of *Acropora spathulata* on the Great Barrier Reef. Heat tolerance was measured as the maximum quantum yield of photosystem II (F_v/F_m) and chlorophyll content (NDVI) expressed as ED50 temperature thresholds and performance retention under extreme heat (+9 °C). Relationships between environmental predictors (defined in Table S7) and heat tolerance metrics were assessed using ridge regression and random forest models. **a.** Heatmap of associations evaluated with random forest variable importance (10-fold stratified cross validation, circle size) and ridge regression standardized coefficients (multiple bootstrap resampling $n = 100$, circle color). The variance explained by each model (random forest/ridge R^2) is shown above each tolerance trait metric. **b-e**. Thermal history maps of (**b**) maximum monthly mean temperature as defined by NOOA (Liu et al., 2003) (**c**) annual temperature range (2014–2022), (**d**) peak degree heating weeks during sampling and (**e**) annual frequency of degree heating weeks above 4 since 1985. White dots indicate reef sites sampled in this study.

Discussion

The large geographic scope of this study (14 reefs across 9.5 degrees of latitude) and sample size (n = 768) revealed extensive intraspecific variation in the heat tolerance of reef-building corals both within and among reefs on the Great Barrier Reef (GBR). This work builds upon previous studies quantifying reef-level variation in acute heat stress tolerance (0.9–1.9 °C; Evensen et al., 2022; Marzonie et al., 2022; 1.8 °C in our study). We documented the highest variation in colony-level tolerance within reefs (3–3.3 °C) to date, exceeding Floridan nurseries (2.5 °C; Cunning et al., 2016). This resulted in >6 °C variation in colony ED50 thresholds where tolerant colonies — based on multiple traits and metrics — were widely distributed across the GBR. Environmental predictors of heat-tolerance variation support adaptation and/or acclimatization of populations to local conditions while unexplained differences among colonies may be due to host and/or symbiont adaptive genetic variation within reefs.

Acute heat tolerance metrics linked to historical and recent environmental conditions

All heat tolerance metrics were most strongly related to thermal history supporting adaptation of corals to their local thermal regimes (Bay & Palumbi, 2014; Howells et al., 2013; Hughes et al., 2019; Sully et al., 2019; Thompson & Van Woesik, 2009). These results also corroborate ecological experiments that found 4 °C difference in *A. spathulata* larvae survival thresholds between Lizard Island (14.7 °S) and One Tree Island (23.5 °S) i.e., the same latitudinal range as our study (Woolsey et al., 2015). F_v/F_m ED50 thresholds increased at reefs exposed to higher thermal averages (MMM), annual thermal variation (AR) and exposure to marine heat waves (frequency of DHW > 4 and standard deviation of DHW). DHW at the time of sampling was also strongly positively associated with F_v/F_m ED50, suggesting that acclimatization of the holobiont to higher temperatures possibly through upregulation of photoprotective mechanisms can increase the heat tolerance trait F_v/F_m ED50 as reported by Cunning et al. (2021). Conversely, there were weaker unexpected negative

associations between all thermal history metrics and NDVI ED50. This may reflect accumulated heat stress leading to early health impairment of the holobiont and lower chlorophyll retention, which is supported by the strong negative relationship of NDVI ED50 with frequency of DHW >4 in the 12 months prior to sampling. This is similar to F_v/F_m ED50 findings of Marzonie et al. (2022) in the Coral Sea, suggesting that some reef populations may be impacted over long time periods after heat exposure. Differences in the association with thermal history among traits could thus occur from a decoupling between host and symbiont baseline conditions and/or stress responses (discussed below; Baird et al., 2009; Gardner et al., 2017; Krueger et al., 2015). Nevertheless, several reefs in our study (No Name, North Direction, Mackay) exhibited a high proportion of tolerant individuals for both F_v/F_m and NDVI ED50, showing that some colonies and populations may perform better under heat stress across multiple traits (Lachs, Humanes, Pygas, et al., 2023). These populations will be more likely to play a key role in the GBR resilience in a warming climate and thus their mapping in space and time could provide valuable information for spatial planning of Australia's marine parks into the future.

The heat tolerance metrics of *A. spathulata* measured here were further associated with predictors that influence water movement, light, and within-reef temperature. Strong water movement can delay photochemical damage under thermal stress through coral surface cooling (Fabricius, 2006), increases in respiration and metabolic transfers, and removal of toxins (Nakamura et al., 2005; Page et al., 2019). Here we observed a positive association between surface current velocity and F_v/F_m metrics which suggests a carryover of these effects to our experiments. Symbiont responses to experimental heat stress may also have been driven by solar irradiance, a factor known to mediate bleaching responses under high seawater temperatures (Lesser & Farrell, 2004). Indeed F_v/F_m ED50 thresholds were reduced at reefs with a high positive cloud fraction anomaly prior to the experiments and may indicate that symbionts acclimated to lower light levels (Anthony & Hoegh-Guldberg, 2003) experienced greater irradiance-induced photosynthetic damage in the experimental system (Roberty et al.,

2015). Overall, these results demonstrate that empirically derived thermal tolerance metrics are shaped by long-term adaptation as well as acclimatization to recent conditions.

Standing within-reef variation in heat tolerance supports adaptive potential to climate change

There was greater variation in the acute heat tolerance of *A. spathulata* colonies within reefs than among reef-level averages (ED50 ~3.3 vs 1.8 °C and ~3.0 vs 1.6 °C for F_v/F_m and NDVI respectively), despite most colonies being sampled from a narrow depth range at a single site less than 600 m^2. Importantly, we found that heat-tolerant corals (defined as the top 25 percentile from the GBR ED50 distributions) occurred at nearly every reef, including the cooler southern GBR. These results align with previous results on *A. cervicornis* in the Florida Keys (Cunning et al., 2021), *A. hyacinthus* in Palau (Cornwell et al., 2021) and *A. hyacinthus* in the GBR (Naugle et al., 2024). Within-reef variation remained mostly unexplained by colony-level predictors (depth and pigmentation) and could be elucidated by incorporating genomic data not yet available here. Individual differences in heat tolerance may be underpinned by genomic variation within the host (Drury, Bean, et al., 2022; Jin et al., 2016), symbionts (Berkelmans & Van Oppen, 2006; Jones et al., 2008; Palacio-Castro et al., 2023) and/or their interactions (Kavousi et al., 2020). For example, incorporating the proportion of *Durusdinium* cells and host polygenic scores based on putative heat-adaptive loci increased the predictive accuracy of GBR *A. millepora* bleaching models by ~22% (Fuller et al., 2020). In addition, deciphering the independent role of environmental drivers requires a better characterization across microhabitats (Ainsworth et al., 2021) and depth (Wyatt et al., 2023) as well as optimization of sampling strategies across environmental gradients to reduce the collinearity of predictors (Selmoni et al., 2020). The large standing variation in thermal tolerance at the reef scale reported in this study has important implications for the conservation of this species on the GBR. It may notably support *A. spathulata* adaptation to climate change through selection of pre-existing heat tolerance alleles (Matz et al., 2020). Furthermore, rather than translocating

individuals from warmer reefs (e.g., Howells et al., 2013) restoration projects can target local heat tolerant individuals when their phenotypes are known or by sampling a diversity of genets.

Coral populations from the Southern GBR may live further from their upper thermal limit

Acute thermal thresholds (ED50) for *A. spathulata* populations on the GBR were 9.5–9.6 °C above their local MMM on southern reefs compared to 7.5–8.8 °C on central and northern reefs. This is consistent with higher performance retention under extreme heat on southern reefs and latitudinal differences in the thermal thresholds of *A. spathulata* larvae (Woolsey et al., 2015). Consequently, under comparable levels of thermal stress, southern populations are expected to be less susceptible to bleaching than central and northern populations. In line with this expectation, southern GBR reefs have experienced less frequent and intense bleaching events than central and northern reefs (Hughes et al., 2021) with 90% less bleaching-related mortality during the 1985–2012 period (De'ath et al., 2012). The lower frequency and magnitude of bleaching events on southern reefs might also allow more time for adaptation to occur through the migration of heat-tolerance associated alleles from warmer northern reefs (Matz et al., 2020).

Conversely, the low cumulative heat stress experienced by those regions may result in coral assemblages that are naïve to high heat stress (Hughes et al., 2021). As such, mild but highly prevalent coral bleaching occurred in the southern GBR during 2016–2017 (21% of colonies; Kennedy et al., 2017) and 2020 (48% of colonies; Nolan et al., 2021; 86% of colonies; Page et al., 2023) and occurred under lower cumulative heat stress (DHW) than in central and northern GBR (Hughes et al., 2021). Further investigations are thus required to understand how recurrent warming disturbances and gene flow with northern regions will shape the evolution of these cooler reefs.

Divergent effects of acute heat stress on thermal tolerance traits

Here we found a moderate correlation between F_v/F_m and NDVI performance retention under extreme heat. This supports the use of photochemical apparatus integrity as an early proxy of coral bleaching (Evensen et al., 2021; Jones et al., 2000; Voolstra et al., 2020) and the importance of symbiont tolerance in driving holobiont heat stress response. For instance, some symbiont species or genotypes may produce more photoprotective pigments or antioxidant compounds, alleviating oxidative stress for the host (Oakley & Davy, 2018; Roberty & Plumier, 2022). Conversely, we found no correlation between ED50s of F_v/F_m and NDVI traits. The only two other studies using F_v/F_m and chlorophyll-related ED50 thresholds found congruent reef-level variation between the two traits for some species (e.g., *Pocillopora verrucosa*; Evensen et al., 2022 and *Acropora hyacinthus*; Naugle et al., 2024) but not others (e.g., *Porites lobata*; Evensen et al., 2022). In both studies, visual chlorophyll scores showed higher noise and lower differences in reef ED50s which aligns with *A. spathulata* pairwise reef differences in NDVI ED50s (0.39 °C) being almost half that of F_v/F_m ED50 (0.67 °C). This may also be due to differences in the timing of measurement between traits as their decline occurs at different rates (Gardner et al., 2017). In *A. tenuis*, F_v/F_m values were found to be stable 0–24 hours after the end of a similar acute heat stress, while visual pigmentation and chlorophyll decreased up to 24 hours after the end of temperature ramp-down (Nielsen et al., 2022). Therefore delaying the timing of NDVI measurements in our study may have revealed higher divergence in chlorophyll retention between reefs. Finally, oxidative stress and photosynthetic damage can occur under moderate heat stress without a decrease in chlorophyll content or symbiont density (Middlebrook et al., 2012). Interactions between algal oxidative stress and physiological reactions such as antioxidant mechanisms (Roberty & Plumier, 2022), immune response from the host (Baird et al., 2009) or nutrient cycling between symbiotic partners (Rädecker et al., 2021) may indeed lead to differential bleaching outcome under the same level of symbiont photodamage.

Divergent responses between traits highlight the importance of defining heat tolerance and selecting phenotypic traits accordingly. Chlorophyll content may be an appropriate proxy for visual holobiont bleaching and mortality as its decline after 16 hours of acute heat stress was found to persist and reflect mortality in the next month (Walker, Cornwell, et al., 2022). However, photosynthetic efficiency may be a better proxy to detect sub-lethal effects as corals can experience physiological stress of reduced tissue biomass and symbiont loss long before changes in pigmentation are visually detectable (Fitt et al., 2001; Gardner et al., 2017). Finally, which phenotypic trait under acute experimental stress would best predict heat tolerance in natural conditions remains unclear and could vary with marine heatwave magnitude and duration as well as the biological level at which they are assessed (Klepac et al., 2023). As no single trait or metric can fully capture the complexity of the coral holobiont heat stress response, it is important to measure multiple traits whenever achievable, including relevant host traits (e.g., antioxidant capacity).

Conclusion

Using standardized acute heat stress assays, we found that heat tolerant corals were widely distributed across the GBR, including cooler, southern reefs. This suggests potential for adaptive responses to climate change, if heat tolerance traits have a heritable basis (Bairos-Novak et al., 2021; Howells et al., 2022). Rankings of coral heat tolerance can guide human interventions and their interpretation and application may depend on the specific traits and metrics measured. Corals with high absolute thermal tolerance can help to decipher heat-stress protective mechanisms (Cleves et al., 2020; Drury, Bean, et al., 2022) and be used as material for assisted gene flow through translocation (e.g., Drury & Lirman; Morikawa & Palumbi, 2019; Quigley et al., 2019), selective breeding (Howells et al., 2021; Quigley et al., 2020) and cryopreservation (Hagedorn et al., 2021). On the contrary, corals with high thermal tolerance relative to local conditions may be better targets for local propagation and breeding, decreasing the risk of carrying pathogens and phenotype-environment mismatches (Sellares Blasco et al., 2021). Nonetheless, natural adaptive processes and human interventions must

be accompanied by reduced anthropogenic emissions through ambitious national and international commitments to secure the persistence of species and populations across the entire GBR.

References

Ainsworth, T. D., Leggat, W., Silliman, B. R., Lantz, C. A., Bergman, J. L., Fordyce, A. J., Page, C. E., Renzi, J. J., Morton, J., & Eakin, C. M. (2021). Rebuilding relationships on coral reefs: Coral bleaching knowledge-sharing to aid adaptation planning for reef users: Bleaching emergence on reefs demonstrates the need to consider reef scale and accessibility when preparing for, and responding to, coral bleaching. *BioEssays*, 2100048.

Alderdice, R., Perna, G., Cárdenas, A., Hume, B. C., Wolf, M., Kühl, M., Pernice, M., Suggett, D. J., & Voolstra, C. R. (2022). Deoxygenation lowers the thermal threshold of coral bleaching. *Scientific reports*, *12*(1), 1-14.

Álvarez-Noriega, M., Marrable, I., Noonan, S. H., Barneche, D. R., & Ortiz, J. C. (2023). Highly conserved thermal performance strategies may limit adaptive potential in corals. *Proceedings of the Royal Society B*, *290*(1990), 20221703.

Angilletta, M. J., Wilson, R. S., Navas, C. A., & James, R. S. (2003). Tradeoffs and the evolution of thermal reaction norms. *Trends in Ecology & Evolution*, *18*(5), 234-240.

Anthony, K. R., & Hoegh-Guldberg, O. (2003). Kinetics of photoacclimation in corals. *Oecologia*, *134*, 23-31.

Anthony, K. R. N., Helmstedt, K. J., Bay, L. K., Fidelman, P., Hussey, K. E., Lundgren, P., Mead, D., McLeod, I. M., Mumby, P. J., Newlands, M., Schaffelke, B., Wilson, K. A., & Hardisty, P. E. (2020). Interventions to help coral reefs under global change-A complex decision challenge. *PloS one*, *15*(8), e0236399. https://doi.org/10.1371/journal.pone.0236399

Bah, T. (2011). *Inkscape: guide to a vector drawing program*. Prentice hall press.

Baird, A. H., Bhagooli, R., Ralph, P. J., & Takahashi, S. (2009). Coral bleaching: the role of the host. *Trends in Ecology & Evolution*, *24*(1), 16-20.

Bairos-Novak, K. A.-O., Hoogenboom, M. A.-O., van Oppen, M. A.-O., & Connolly, S. A.-O. (2021). Coral adaptation to climate change: Meta-analysis reveals high heritability across multiple traits. LID - 10.1111/gcb.15829 [doi]. *Global Change Biology*(27), 5694-5710.

Barshis, D. J., Ladner, J. T., Oliver, T. A., Seneca, F. O., Traylor-Knowles, N., & Palumbi, S. R. (2013). Genomic basis for coral resilience to climate change. *Proceedings of the National Academy of Sciences*, *110*(4), 1387-1392.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *ArXiv preprint arXiv:1406.5823*.

Bay, R. A., & Palumbi, S. R. (2014). Multilocus adaptation associated with heat resistance in reef-building corals. *Current biology*, *24*(24), 2952-2956.

Bennett, S., Duarte, C. M., Marbà, N., & Wernberg, T. (2019). Integrating within-species variation in thermal physiology into climate change ecology. *Philosophical Transactions of the Royal Society B*, *374*(1778), 20180550.

Berkelmans, R., & Van Oppen, M. J. (2006). The role of zooxanthellae in the thermal tolerance of corals: a 'nugget of hope'for coral reefs in an era of climate change. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1599), 2305-2312.

BOM. (2022). *Tide Predictions for Australia, South Pacific and Antarctica*. Australian government Bureau of Meteorology. Retrieved 19/04/2022 from http://www.bom.gov.au/australia/tides/

Bowden-Kerby, A. (2022). Coral-Focused Climate Change Adaptation and Restoration Based on Accelerating Natural Processes: Launching the "Reefs of Hope" Paradigm. *Oceans*, *4*(1), 13-26.

Bruno, J. F., & Selig, E. R. (2007). Regional decline of coral cover in the Indo-Pacific: timing, extent, and subregional comparisons. *PloS one*, *2*(8), e711.

Charmantier, A., McCleery, R. H., Cole, L. R., Perrins, C., Kruuk, L. E., & Sheldon, B. C. (2008). Adaptive phenotypic plasticity in response to climate change in a wild bird population. *Science*, *320*(5877), 800-803.

Cleves, P. A., Tinoco, A. I., Bradford, J., Perrin, D., Bay, L. K., & Pringle, J. R. (2020). Reduced thermal tolerance in a coral carrying CRISPR-induced mutations in the gene for a heat-shock transcription factor. *Proceedings of the National Academy of Sciences*, *117*(46), 28899-28905.

Cooper, T. F., Berkelmans, R., Ulstrup, K. E., Weeks, S., Radford, B., Jones, A. M., Doyle, J., Canto, M., O'Leary, R. A., & van Oppen, M. J. (2011). Environmental factors controlling the distribution of Symbiodinium harboured by the coral *Acropora millepora* on the Great Barrier Reef. *PloS one*, *6*(10), e25536.

Cornwall, C. E., Comeau, S., Donner, S. D., Perry, C., Dunne, J., van Hooidonk, R., Ryan, S., & Logan, C. A. (2023). Coral adaptive capacity insufficient to halt global transition of coral reefs into net erosion under climate change. *Global Change Biology*.

Cornwell, B., Armstrong, K., Walker, N. S., Lippert, M., Nestor, V., Golbuu, Y., & Palumbi, S. R. (2021). Widespread variation in heat tolerance and symbiont load are associated with growth tradeoffs in the coral *Acropora hyacinthus* in Palau. *eLife*, *10*, e64790.

Costanza, R., De Groot, R., Sutton, P., Van der Ploeg, S., Anderson, S. J., Kubiszewski, I., Farber, S., & Turner, R. K. (2014). Changes in the global value of ecosystem services. *Global environmental change*, *26*, 152-158.

CSIRO. (2020a). *eReefs GBR 1 hydrodynamic model version 2.* Data set accessed 2022-07-13 at https://extraction.ereefs.aims.gov.au/.

CSIRO. (2020b). *eReefs GBR 4 hydrodynamic model version 2.* Data set accessed 2022-04-22 at https://extraction.ereefs.aims.gov.au/.

Cunning, R., Parker, K. E., Johnson-Sapp, K., Karp, R. F., Wen, A. D., Williamson, O. M., Bartels, E., D'Alessandro, M., Gilliam, D. S., & Hanson, G. (2021). Census of heat tolerance among Florida's threatened staghorn corals finds resilient individuals throughout existing nursery populations. *Proceedings of the Royal Society B*, *288*(1961), 20211613.

Cunning, R., Ritson-Williams, R., & Gates, R. D. (2016). Patterns of bleaching and recovery of Montipora capitata in Kāne ʻohe Bay, Hawai ʻi, USA. *Marine Ecology Progress Series*, *551*, 131-139.

De'ath, G., Fabricius, K. E., Sweatman, H., & Puotinen, M. (2012). The 27–year decline of coral cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences*, *109*(44), 17995-17999.

DeCarlo, T. M., Harrison, H. B., Gajdzik, L., Alaguarda, D., Rodolfo-Metalpa, R., D'Olivo, J., Liu, G., Patalwala, D., & McCulloch, M. T. (2019). Acclimatization of massive reef-building corals to consecutive heatwaves. *Proceedings of the Royal Society B*, *286*(1898), 20190235.

Dixon, A. M., Forster, P. M., Heron, S. F., Stoner, A. M. K., & Beger, M. (2022). Future loss of local-scale thermal refugia in coral reef ecosystems. *PLOS Climate*, *1*(2), e0000004. https://doi.org/10.1371/journal.pclm.0000004

Dixon, G. B., Davies, S. W., Aglyamova, G. V., Meyer, E., Bay, L. K., & Matz, M. V. (2015). Genomic determinants of coral heat tolerance across latitudes. *Science*, *348*(6242), 1460-1462.

Drury, C., Bean, N. K., Harris, C. I., Hancock, J. R., Huckeba, J., Roach, T. N., Quinn, R. A., & Gates, R. D. (2022). Intrapopulation adaptive variance supports thermal tolerance in a reef-building coral. *Communications biology*, *5*(1), 1-10.

Drury, C., & Lirman, D. Genotype by environment interactions in coral bleaching. *Proceedings of the Royal Society B*, *288*(1946), 20210177.

Drury, C., Martin, R. E., Knapp, D. E., Heckler, J., Levy, J., Gates, R. D., & Asner, G. P. (2022). Ecosystem-scale mapping of coral species and thermal tolerance. *Frontiers in Ecology and the Environment*.

Evensen, N. R., Fine, M., Perna, G., Voolstra, C. R., & Barshis, D. J. (2021). Remarkably high and consistent tolerance of a Red Sea coral to acute and chronic thermal stress exposures. *Limnology and Oceanography*.

Evensen, N. R., Voolstra, C. R., Fine, M., Perna, G., Buitrago-López, C., Cárdenas, A., Banc-Prandi, G., Rowe, K., & Barshis, D. J. (2022). Empirically derived thermal thresholds of four coral species along the Red Sea using a portable and standardized experimental approach. *Coral Reefs*, 1-14.

Fabricius, K. E. (2006). Effects of irradiance, flow, and colony pigmentation on the temperature microenvironment around corals: Implications for coral bleaching? *Limnology and Oceanography*, *51*(1), 30-37.

Fangue, N. A., Hofmeister, M., & Schulte, P. M. (2006). Intraspecific variation in thermal tolerance and heat shock protein gene expression in common killifish, *Fundulus heteroclitus*. *Journal of Experimental Biology*, *209*(15), 2859-2872.

Fitt, W. K., Brown, B. E., Warner, M. E., & Dunne, R. P. (2001). Coral bleaching: interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral Reefs*, *20*, 51-65.

Franken, O., Huizinga, M., Ellers, J., & Berg, M. P. (2018). Heated communities: large inter- and intraspecific variation in heat tolerance across trophic levels of a soil arthropod community. *Oecologia*, *186*, 311-322.

Friedman, J., Hastie, T., Tibshirani, R., Narasimhan, B., Tay, K., Simon, N., & Qian, J. (2021). Package 'glmnet'. *CRAN R Repositary*.

Fuller, Z. L., Mocellin, V. J., Morris, L. A., Cantin, N., Shepherd, J., Sarre, L., Peng, J., Liao, Y., Pickrell, J., & Andolfatto, P. (2020). Population genetics of the coral *Acropora millepora*: Toward genomic prediction of bleaching. *Science*, *369*(6501).

Gardner, S. G., Raina, J.-B., Nitschke, M. R., Nielsen, D. A., Stat, M., Motti, C. A., Ralph, P. J., & Petrou, K. (2017). A multi-trait systems approach reveals a response cascade to bleaching in corals. *BMC biology*, *15*(1), 1-14.

Gardner, T. A., Côté, I. M., Gill, J. A., Grant, A., & Watkinson, A. R. (2003). Long-term region-wide declines in Caribbean corals. *Science*, *301*(5635), 958-960.

Gonzalez-Espinosa, P. C., & Donner, S. D. (2021). Cloudiness reduces the bleaching response of coral reefs exposed to heat stress. *Global Change Biology*.

Hackerott, S., Martell, H. A., & Eirin-Lopez, J. M. (2021). Coral environmental memory: causes, mechanisms, and consequences for future reefs. *Trends in Ecology & Evolution*.

Hagedorn, M., Page, C. A., O'Neil, K. L., Flores, D. M., Tichy, L., Conn, T., Chamberland, V. F., Lager, C., Zuchowicz, N., & Lohr, K. (2021). Assisted gene flow using cryopreserved sperm in critically endangered coral. *Proceedings of the National Academy of Sciences*, *118*(38).

Herrando-Pérez, S., Ferri-Yáñez, F., Monasterio, C., Beukema, W., Gomes, V., Belliure, J., Chown, S. L., Vieites, D. R., & Araújo, M. B. (2019). Intraspecific variation in lizard heat tolerance alters estimates of climate impact. *Journal of Animal Ecology*, *88*(2), 247-257.

Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield, P., Gomez, E., Harvell, C. D., Sale, P. F., Edwards, A. J., & Caldeira, K. (2007). Coral reefs under rapid climate change and ocean acidification. *science*, *318*(5857), 1737-1742.

Holstein, D. M., Smith, T. B., van Hooidonk, R., & Paris, C. B. (2022). Predicting coral metapopulation decline in a changing thermal environment. *Coral Reefs*, 1-12.

Hothorn, T., Hornik, K., Strobl, C., Zeileis, A., & Hothorn, M. T. (2015). Package 'party'. *Package Reference Manual for Party Version 0.9-998, 16*, 37.

Howells, E., Beltran, V., Larsen, N., Bay, L., Willis, B., & Van Oppen, M. (2012). Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nature Climate Change, 2*(2), 116-120.

Howells, E. J., Abrego, D., Liew, Y. J., Burt, J. A., Meyer, E., & Aranda, M. (2021). Enhancing the heat tolerance of reef-building corals to future warming. *Science Advances, 7*(34), eabg6070.

Howells, E. J., Abrego, D., Meyer, E., Kirk, N. L., & Burt, J. A. (2016). Host adaptation and unexpected symbiont partners enable reef-building corals to tolerate extreme temperatures. *Global Change Biology, 22*(8), 2702-2714.

Howells, E. J., Bay, L. K., & Bay, R. A. (2022). Identifying, Monitoring, and Managing Adaptive Genetic Variation in Reef-Building Corals under Rapid Climate Warming. *Coral Reef Conservation and Restoration in the Omics Age*, 55-70.

Howells, E. J., Berkelmans, R., van Oppen, M. J., Willis, B. L., & Bay, L. K. (2013). Historical thermal regimes define limits to coral acclimatization. *Ecology, 94*(5), 1078-1088.

Hughes, T. (2020). Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. Science 359, 80–83 (2018). *Scientific reports, 10*, 20346.

Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D., Baird, A. H., Babcock, R. C., Beger, M., Bellwood, D. R., & Berkelmans, R. (2017). Global warming and recurrent mass bleaching of corals. *Nature, 543*(7645), 373-377.

Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Dietzel, A., Eakin, C. M., Heron, S. F., Hoey, A. S., Hoogenboom, M. O., & Liu, G. (2018). Global warming transforms coral reef assemblages. *Nature, 556*(7702), 492-496.

Hughes, T. P., Kerry, J. T., Connolly, S. R., Álvarez-Romero, J. G., Eakin, C. M., Heron, S. F., Gonzalez, M. A., & Moneghetti, J. (2021). Emergent properties in the responses of tropical corals to recurrent climate extremes. *Current biology, 31*(23), 5393-5399. e5393.

Hughes, T. P., Kerry, J. T., Connolly, S. R., Baird, A. H., Eakin, C. M., Heron, S. F., Hoey, A. S., Hoogenboom, M. O., Jacobson, M., & Liu, G. (2019). Ecological memory modifies the cumulative impact of recurrent climate extremes. *Nature Climate Change, 9*(1), 40-43.

Humanes, A., Lachs, L., Beauchamp, E. A., Bythell, J. C., Edwards, A. J., Golbuu, Y., Martinez, H. M., Palmowski, P., Treumann, A., & van der Steeg, E. (2022). Within-population variability in coral heat tolerance indicates climate adaptation potential. *Proceedings of the Royal Society B, 289*(1981), 20220872.

Jin, Y. K., Lundgren, P., Lutz, A., Raina, J.-B., Howells, E. J., Paley, A. S., Willis, B. L., & Van Oppen, M. J. (2016). Genetic markers for antioxidant capacity in a reef-building coral. *Science Advances, 2*(5), e1500842.

Jones, A. M., Berkelmans, R., van Oppen, M. J., Mieog, J. C., & Sinclair, W. (2008). A community change in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field evidence of acclimatization. *Proceedings of the Royal Society B: Biological Sciences, 275*(1641), 1359-1365.

Jones, R. J., Ward, S., Amri, A. Y., & Hoegh-Guldberg, O. (2000). Changes in quantum efficiency of Photosystem II of symbiotic dinoflagellates of corals after heat stress, and of bleached corals sampled after the 1998 Great Barrier Reef mass bleaching event. *Marine and Freshwater Research, 51*(1), 63-71.

Kavousi, J., Denis, V., Sharp, V., Reimer, J. D., Nakamura, T., & Parkinson, J. E. (2020). Unique combinations of coral host and algal symbiont genotypes reflect intraspecific variation in heat stress responses among colonies of the reef-building coral, *Montipora digitata. Marine Biology, 167*, 1-15.

Kelso, N. V., & Patterson, T. (2010). Introducing natural earth data-naturalearthdata. com. *Geographia Technica, 5*(82-89), 25.

Kennedy, E. V., Ordoñez, A., & Diaz-Pulido, G. (2017). Coral bleaching in the southern inshore Great Barrier Reef: a case study from the Keppel Islands. *Marine and Freshwater Research*, *69*(1), 191-197.

Kirk, N. L., Howells, E. J., Abrego, D., Burt, J. A., & Meyer, E. (2018). Genomic and transcriptomic signals of thermal tolerance in heat-tolerant corals (*Platygyra daedalea*) of the Arabian/Persian Gulf. *Molecular Ecology*, *27*(24), 5180-5194.

Klepac, C. N., & Barshis, D. J. (2022). High-resolution in situ thermal metrics coupled with acute heat stress experiments reveal differential coral bleaching susceptibility. *Coral Reefs*, 1-13.

Klepac, C. N., Petrik, C., Karabelas, E., Owens, J., Hall, E. R., & Muller, E. M. (2023). Assessing Acute Thermal Assays as a Rapid Screening Tool for Coral Restoration. *Research Square*. https://doi.org/10.21203/rs.3.rs-3140553/v1

Krueger, T., Hawkins, T. D., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., Leggat, W., Fisher, P. L., & Davy, S. K. (2015). Differential coral bleaching—Contrasting the activity and response of enzymatic antioxidants in symbiotic partners under thermal stress. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *190*, 15-25.

Lachs, L., Humanes, A., Mumby, P. J., Donner, S. D., Bythell, J. C., Beauchamp, E., Bukurou, L., Buzzoni, D., de la Torre Cerro, R., & East, H. K. (2023). Unexpectedly high coral heat tolerance at thermal refugia. *BioRxiv*, 2023.2006. 2016.545328.

Lachs, L., Humanes, A., Pygas, D. R., Bythell, J. C., Mumby, P. J., Ferrari, R., Figueira, W. F., Beauchamp, E., East, H. K., & Edwards, A. J. (2023). No apparent trade-offs associated with heat tolerance in a reef-building coral. *Communications biology*, *6*(1), 400.

LaJeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra, C. R., & Santos, S. R. (2018). Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Current biology*, *28*(16), 2570-2580. e2576.

Leal, M. C., Jesus, B., Ezequiel, J., Calado, R., Rocha, R. J. M., Cartaxana, P., & Serôdio, J. (2015). Concurrent imaging of chlorophyll fluorescence, Chlorophyll a content and green fluorescent proteins-like proteins of symbiotic cnidarians. *Marine ecology*, *36*(3), 572-584.

Lesser, M. P., & Farrell, J. H. (2004). Exposure to solar radiation increases damage to both host tissues and algal symbionts of corals during thermal stress. *Coral Reefs*, *23*, 367-377.

Liu, G., Strong, A. E., & Skirving, W. (2003). Remote sensing of sea surface temperatures during 2002 Barrier Reef coral bleaching. *Eos, Transactions American Geophysical Union*, *84*(15), 137-141.

Lukoschek, V., Riginos, C., & van Oppen, M. J. (2016). Congruent patterns of connectivity can inform management for broadcast spawning corals on the Great Barrier Reef. *Molecular Ecology*, *25*(13), 3065-3080.

Marhoefer, S. R., Zenger, K. R., Strugnell, J. M., Logan, M., van Oppen, M. J., Kenkel, C. D., & Bay, L. K. (2021). Signatures of adaptation and acclimatization to reef flat and slope habitats in the coral Pocillopora damicornis. *Frontiers in Marine Science*, *8*, 704709.

Marzonie, M. R., Bay, L. K., Bourne, D. G., Hoey, A. S., Matthews, S., Nielsen, J. J., & Harrison, H. B. (2022). The effects of marine heatwaves on acute heat tolerance in corals. *Global Change Biology*.

Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., & Shukla, P. R. (2022). *Global Warming of 1.5° C: IPCC Special Report on Impacts of Global Warming of 1.5° C above Pre-industrial Levels in Context of Strengthening Response to Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*. Cambridge University Press.

Matz, M. V., Treml, E. A., & Haller, B. C. (2020). Estimating the potential for coral adaptation to global warming across the Indo-West Pacific. *Global Change Biology, 26*(6), 3473-3481.

McKenzie, D. J., Zhang, Y., Eliason, E. J., Schulte, P. M., Claireaux, G., Blasco, F. R., Nati, J. J., & Farrell, A. P. (2021). Intraspecific variation in tolerance of warming in fishes. *Journal of fish biology, 98*(6), 1536-1555.

McWilliam, M., Madin, J. S., Chase, T. J., Hoogenboom, M. O., & Bridge, T. C. (2022). Intraspecific variation reshapes coral assemblages under elevated temperature and acidity. *Ecology letters.*

Middlebrook, R., Anthony, K. R., Hoegh-Guldberg, O., & Dove, S. (2012). Thermal priming affects symbiont photosynthesis but does not alter bleaching susceptibility in Acropora millepora. *Journal of Experimental Marine Biology and Ecology, 432*, 64-72.

Morikawa, M. K., & Palumbi, S. R. (2019). Using naturally occurring climate resilient corals to construct bleaching-resistant nurseries. *Proceedings of the National Academy of Sciences, 116*(21), 10586-10591.

Morley, S., Martin, S., Bates, A., Clark, M., Ericson, J., Lamare, M., & Peck, L. (2012). Spatial and temporal variation in the heat tolerance limits of two abundant Southern Ocean invertebrates. *Marine Ecology Progress Series, 450*, 81-92.

Muscatine, L., & Porter, J. W. (1977). Reef corals: mutualistic symbioses adapted to nutrient-poor environments. *Bioscience, 27*(7), 454-460.

Nakamura, T., Van Woesik, R., & Yamasaki, H. (2005). Photoinhibition of photosynthesis is reduced by water flow in the reef-building coral Acropora digitifera. *Marine Ecology Progress Series, 301*, 109-118.

Naugle, M. S., Denis, H., Mocellin, V. J. L., Laffy, P. W., Popovic, I., Bay. L. K., Howells, E. J. (2024). Environmental, host, and symbiont drivers of heat tolerance in a species complex of reef-building corals. *PREPRINT (Version 1) Available at Biorxiv.*

Nielsen, J., Matthews, G., Frith, K., Harrison, H., Marzonie, M., Slaughter, K., Suggett, D., & Bay, L. (2022). Experimental considerations of acute heat stress assays to quantify coral thermal tolerance.

NOAA Coral Reef Watch. (2018, updated daily). *NOAA Coral Reef Watch Version 3.1, Daily Global 5-km Satellite Sea Surface Temperature Product (a.k.a CoralTemp), Jun. 1, 1985-July. 3, 2022. College Park, Maryland, USA: NOAA Coral Reef Watch. Data set accessed 2022-07-04 at*
https://coralreefwatch.noaa.gov/product/5km/index_5km_sst.php.

Noakes, M. J., Wolf, B. O., & McKechnie, A. E. (2016). Seasonal and geographical variation in heat tolerance and evaporative cooling capacity in a passerine bird. *Journal of Experimental Biology, 219*(6), 859-869.

Nolan, M. K., Schmidt-Roach, S., Davis, A. R., Aranda, M., & Howells, E. J. (2021). Widespread bleaching in the One Tree Island lagoon (Southern Great Barrier Reef) during record-breaking temperatures in 2020. *Environmental monitoring and assessment, 193*(9), 1-8.

Oakley, C., & Davy, S. (2018). Cell biology of coral bleaching. In *Coral bleaching* (pp. 189-211). Springer.

Page, C. A., Giuliano, C., Bay, L. K., & Randall, C. J. (2023). High survival following bleaching underscores the resilience of a frequently disturbed region of the Great Barrier Reef. *Ecosphere, 14*(2), e4280.

Page, C. E., Leggat, W., Heron, S. F., Choukroun, S. M., Lloyd, J., & Ainsworth, T. D. (2019). Seeking resistance in coral reef ecosystems: the interplay of biophysical factors and bleaching resistance under a changing climate: the interplay of a reef's biophysical factors can mitigate the coral bleaching response. *BioEssays, 41*(7), 1800226.

Palacio-Castro, A. M., Smith, T. B., Brandtneris, V., Snyder, G. A., van Hooidonk, R., Maté, J. L., Manzello, D., Glynn, P. W., Fong, P., & Baker, A. C. (2023). Increased dominance of heat-tolerant symbionts creates resilient coral reefs in near-term ocean warming. *Proceedings of the National Academy of Sciences, 120*(8), e2202388120.

Parkinson, J. E., Banaszak, A. T., Altman, N. S., LaJeunesse, T. C., & Baums, I. B. (2015). Intraspecific diversity among partners drives functional variation in coral symbioses. *Scientific reports*, *5*(1), 15667.

Quigley, K., Randall, C., Van Oppen, M., & Bay, L. (2020). Assessing the role of historical temperature regime and algal symbionts on the heat tolerance of coral juveniles. *Biology open*, *9*(1), bio047316.

Quigley, K. M., Bay, L. K., & van Oppen, M. J. (2019). The active spread of adaptive variation for reef resilience. *Ecology and Evolution*, *9*(19), 11122-11135.

Rädecker, N., Pogoreutz, C., Gegner, H. M., Cárdenas, A., Roth, F., Bougoure, J., Guagliardo, P., Wild, C., Pernice, M., & Raina, J.-B. (2021). Heat stress destabilizes symbiotic nutrient cycling in corals. *Proceedings of the National Academy of Sciences*, *118*(5).

Reusch, T. B. (2014). Climate change in the oceans: evolutionary versus phenotypically plastic responses of marine animals and plants. *Evolutionary Applications*, *7*(1), 104-122.

Riegl, B., Johnston, M., Purkis, S., Howells, E., Burt, J., Steiner, S. C., Sheppard, C. R., & Bauman, A. (2018). Population collapse dynamics in *Acropora downingi*, an Arabian/Persian Gulf ecosystem-engineering coral, linked to rising temperature. *Global Change Biology*, *24*(6), 2447-2462.

Ritchie, R. J. (2008). Universal chlorophyll equations for estimating chlorophylls a, b, c, and d and total chlorophylls in natural assemblages of photosynthetic organisms using acetone, methanol, or ethanol solvents. *Photosynthetica*, *46*(1), 115-126.

Ritz, C., Baty, F., Streibig, J. C., & Gerhard, D. (2015). Dose-response analysis using R. *PloS one*, *10*(12), e0146021.

Roberty, S., Fransolet, D., Cardol, P., Plumier, J.-C., & Franck, F. (2015). Imbalance between oxygen photoreduction and antioxidant capacities in Symbiodinium cells exposed to combined heat and high light stress. *Coral Reefs*, *34*, 1063-1073.

Roberty, S., & Plumier, J.-C. (2022). Bleaching physiology: who's the 'weakest link'—host vs. symbiont? *Emerging Topics in Life Sciences*, *6*(1), 17-32.

Rocha, R. J., Pimentel, T., Serôdio, J., Rosa, R., & Calado, R. (2013). Comparative performance of light emitting plasma (LEP) and light emitting diode (LED) in ex situ aquaculture of scleractinian corals. *Aquaculture*, *402*, 38-45.

Rose, N. H., Bay, R. A., Morikawa, M. K., & Palumbi, S. R. (2018). Polygenic evolution drives species divergence and climate adaptation in corals. *Evolution*, *72*(1), 82-94.

Rose, N. H., Bay, R. A., Morikawa, M. K., Thomas, L., Sheets, E. A., & Palumbi, S. R. (2021). Genomic analysis of distinct bleaching tolerances among cryptic coral species. *Proceedings of the Royal Society B*, *288*(1960), 20210678.

Sampayo, E., Ridgway, T., Bongaerts, P., & Hoegh-Guldberg, O. (2008). Bleaching susceptibility and mortality of corals are determined by fine-scale differences in symbiont type. *Proceedings of the National Academy of Sciences*, *105*(30), 10444-10449.

Savtchenko, A., Ouzounov, D., Ahmad, S., Acker, J., Leptoukh, G., Koziana, J., & Nickless, D. (2004). Terra and Aqua MODIS products available from NASA GES DAAC. *Advances in Space Research*, *34*(4), 710-714.

Schoepf, V., Sanderson, H., & Larcombe, E. (2021). Coral heat tolerance under variable temperatures: Effects of different variability regimes and past environmental history vs. current exposure. *Limnology and Oceanography*.

Schoepf, V., Stat, M., Falter, J. L., & McCulloch, M. T. (2015). Limits to the thermal tolerance of corals adapted to a highly fluctuating, naturally extreme temperature environment. *Scientific reports*, *5*(1), 1-14.

Sellares Blasco, R. I., Villalpando, M. F., Guendulain-Garcia, S. D., & Cróquer, A. (2021). Assisted coral reproduction in the Dominican Republic: a successful story to replicate in the Caribbean. *Frontiers in Marine Science*, *8*, 927.

Selmoni, O., Vajana, E., Guillaume, A., Rochat, E., & Joost, S. (2020). Sampling strategy optimization to increase statistical power in landscape genomics: A simulation-based approach. *Molecular ecology resources*, *20*(1), 154-169.

Shaw, E. C., Carpenter, R. C., Lantz, C. A., & Edmunds, P. J. (2016). Intraspecific variability in the response to ocean warming and acidification in the scleractinian coral Acropora pulchra. *Marine Biology*, *163*, 1-11.

Siebeck, U., Marshall, N., Klüter, A., & Hoegh-Guldberg, O. (2006). Monitoring coral bleaching using a colour reference card. *Coral Reefs*, *25*, 453-460.

Sinervo, B., Mendez-De-La-Cruz, F., Miles, D. B., Heulin, B., Bastiaans, E., Villagrán-Santa Cruz, M., Lara-Resendiz, R., Martínez-Méndez, N., Calderón-Espinosa, M. L., & Meza-Lázaro, R. N. (2010). Erosion of lizard diversity by climate change and altered thermal niches. *Science*, *328*(5980), 894-899.

Smith, A. B., Alsdurf, J., Knapp, M., Baer, S. G., & Johnson, L. C. (2017). Phenotypic distribution models corroborate species distribution models: A shift in the role and prevalence of a dominant prairie grass in response to climate change. *Global Change Biology*, *23*(10), 4365-4375.

Somero, G. (2010). The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine 'winners' and 'losers'. *Journal of Experimental Biology*, *213*(6), 912-920.

Stat, M., Carter, D., & Hoegh-Guldberg, O. (2006). The evolutionary history of Symbiodinium and scleractinian hosts—symbiosis, diversity, and the effect of climate change. *Perspectives in Plant Ecology, Evolution and Systematics*, *8*(1), 23-43.

Steven, A. D., Baird, M. E., Brinkman, R., Car, N. J., Cox, S. J., Herzfeld, M., Hodge, J., Jones, E., King, E., & Margvelashvili, N. (2019). eReefs: An operational information system for managing the Great Barrier Reef. *Journal of Operational Oceanography*, *12*(sup2), S12-S28.

Sully, S., Burkepile, D. E., Donovan, M., Hodgson, G., & Van Woesik, R. (2019). A global analysis of coral bleaching over the past two decades. *Nature communications*, *10*(1), 1-5.

Sully, S., Hodgson, G., & van Woesik, R. (2022). Present and future bright and dark spots for coral reefs through climate change. *Global Change Biology*.

Team, R. C. (2020). R: A language and environment for statistical computing (Version 4.0. 3)[Computer software]. R Foundation for Statistical Computing. In.

Thomas, L., Rose, N. H., Bay, R. A., López, E. H., Morikawa, M. K., Ruiz-Jones, L., & Palumbi, S. R. (2018). Mechanisms of thermal tolerance in reef-building corals across a fine-grained environmental mosaic: lessons from Ofu, American Samoa. *Frontiers in Marine Science*, *4*, 434.

Thompson, D. M., & Van Woesik, R. (2009). Corals escape bleaching in regions that recently and historically experienced frequent thermal stress. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1669), 2893-2901.

Tracy, C. R., & Christian, K. A. (1986). Ecological relations among space, time, and thermal niche axes. *Ecology*, *67*(3), 609-615.

UNEP-WCMC, W.-C. (2010). WRI, & TNC.(2010). *Global distribution of warm-water coral reefs, compiled from multiple sources including the Millennium Coral Reef Mapping Project. Version*, *1*.

Virgen-Urcelay, A., & Donner, S. D. (2023). Increase in the extent of mass coral bleaching over the past half-century, based on an updated global database. *PloS one*, *18*(2), e0281719.

Voolstra, Valenzuela, J. J., Turkarslan, S., Cárdenas, A., Hume, B. C., Perna, G., Buitrago-López, C., Rowe, K., Orellana, M. V., & Baliga, N. S. (2021). Contrasting heat stress response patterns of coral holobionts across the Red Sea suggest distinct mechanisms of thermal tolerance. *Molecular Ecology*, *30*(18), 4466-4480.

Voolstra, C. R., Buitrago-López, C., Perna, G., Cárdenas, A., Hume, B. C., Rädecker, N., & Barshis, D. J. (2020). Standardized short-term acute heat stress assays resolve

historical differences in coral thermotolerance across microhabitat reef sites. *Global Change Biology*, *26*(8), 4328-4343.

Walker, N. S., Cornwell, B. H., Nestor, V., Armstrong, K. C., Golbuu, Y., & Palumbi, S. R. (2022). Persistence of phenotypic responses to short-term heat stress in the tabletop coral *Acropora hyacinthus*. *PloS one*, *17*(9), e0269206.

Walker, N. S., Nestor, V., Golbuu, Y., & Palumbi, S. R. (2022). Coral bleaching resistance variation is linked to differential mortality and skeletal growth during recovery. *Evolutionary Applications*, 16(12), 504-517.

Wallace, C. C., Done, B. J., & Muir, P. R. (2012). Revision and catalogue of worldwide staghorn corals Acropora and Isopora (Scleractinia: Acroporidae) in the Museum of Tropical Queensland.

Wickham, H. (2011). ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics*, *3*(2), 180-185.

Wiedenmann, J., D'Angelo, C., Smith, E. G., Hunt, A. N., Legiret, F.-E., Postle, A. D., & Achterberg, E. P. (2013). Nutrient enrichment can increase the susceptibility of reef corals to bleaching. *Nature Climate Change*, *3*(2), 160-164.

Wijgerde, T., van Melis, A., Silva, C. I., Leal, M. C., Vogels, L., Mutter, C., & Osinga, R. (2014). Red light represses the photophysiology of the scleractinian coral *Stylophora pistillata*. *PloS one*, *9*(3), e92781.

Woolsey, E., Keith, S., Byrne, M., Schmidt-Roach, S., & Baird, A. (2015). Latitudinal variation in thermal tolerance thresholds of early life stages of corals. *Coral Reefs*, *34*(2), 471-478.

Wyatt, A. S., Leichter, J. J., Washburn, L., Kui, L., Edmunds, P. J., & Burgess, S. C. (2023). Hidden heatwaves and severe coral bleaching linked to mesoscale eddies and thermocline dynamics. *Nature communications*, *14*(1), 1-17.

Acknowledgments

We thank the Traditional Owners of the Great Barrier Reef for their free prior and informed consent to enter and undertake coral sampling in their Sea Country. We pay our respect to their Elders, past, present, and future. We thank Andrea Severati, Steven Green, and Justin Hochen from the Australian Institute of Marine Science for technical support and building the acute heat stress experimental system. We also thank Samantha Goyen, Riverside Marine Cruise crew members, and numerous AIMS staff and volunteers for their support while collecting and experimenting at sea. We are thankful to Eric Lawrey and Eduardo Klein for their support with environmental data collection as well as Jonathan Kok and Eric Hochberg for developing tools and pipelines for hyperspectral imaging data extraction.

Authors contributions

L.K.B. and E.J.H. initiated the project and designed the experiment. H.D., L.K.B., V.M., M.N. and E.H. collected fragments and performed acute heat stress experiments. H.D. and V.M.

measured total chlorophyll content in the laboratory. H.D. extracted image-based phenotypic measurements and retrieved environmental data together with M.N. H.D. performed the data analysis, supported by E.J.H., G.L. and V.B.L. H.D. wrote the first version of the manuscript, which was critically revised by all other authors. All authors read and approved the final manuscript.

Data availability statement

Scripts and data used for the analyses are available online: https://github.com/hde08/GBR-HeatTolVar-GCB/tree/main; publicly archived on https://zenodo.org/records/xxx.

Funding information

This work was supported by the Reef Restoration and Adaptation Program funded by the partnership between the Australian Government's Reef Trust and the Great Barrier Reef Foundation.

Competing interests

The authors declare no competing interests.