Title

Thermal tolerance traits of individual corals are widely distributed across the Great Barrier

Reef

Abstract

Adaptation of reef-building corals to global warming depends upon standing heritable

variation in tolerance traits upon which selection can act. Yet limited knowledge exists on

heat tolerance variation among conspecific individuals separated by meters to hundreds of

kilometers. Here, we performed standardized acute heat stress assays to quantify the

thermal tolerance traits of 768 colonies of *Acropora spathulata* from 14 reefs spanning 1060

km (9.5° latitude) of the Great Barrier Reef. Thermal thresholds for photochemical efficiency

and chlorophyll retention varied considerably among individual colonies both among reefs

(~6 °C) and within reefs (~3 °C). Although tolerance rankings of colonies varied between

traits, the most heat tolerant corals (i.e. top 25% of each trait) were found at virtually all reefs,

indicating widespread phenotypic variation. Reef-scale environmental predictors explained

12–62% of trait variation. Corals exposed to high thermal averages and recent thermal stress

exhibited the greatest photochemical performance, likely reflecting local adaptation and

stress pre-acclimatization, and the lowest chlorophyll retention suggesting stress pre-

sensitization. Importantly, heat tolerance relative to local summer temperatures was the

greatest on southern reefs suggestive of higher adaptive potential. These results can be

used to identify naturally tolerant coral populations and individuals for conservation and

restoration applications.

Introduction
25

Global warming modifies species thermal environments[1] and directly impacts organism

fitness[2]. The persistence of species and the ecosystem services they provide therefore

depends on their response to rising and extreme temperatures. Yet, considerable

intraspecific variation in thermal tolerance exists in marine ecosystems[3] as some species

have colonized large geographical areas spanning contrasted thermal environments[4].

Individuals can adjust to local temperatures within their lifespan through physiological

acclimatization and selection for particular phenotypes within populations can lead to genetic

adaptation across generations[5]. Documenting phenotypic differences along thermal gradients

can thus inform conservation and restoration efforts[6] by improving predictions of species

adaptive potential[3], climate refugia[7] or extinctions[8].

Reef-building (scleractinian) corals are among the organisms most at risk from global

warming as they occupy narrow thermal niches in environments where current summer

temperatures often approach and increasingly exceed their upper thermal limits[9]. The

energetic requirements of corals rely on a finely-tuned symbiotic relationship with

photosynthetic dinoflagellates (family Symbiodiniaceae)[10] that can be severely impaired

under heat stress[11], triggering a loss of symbiont cells and/or their photosynthetic pigments

known as coral bleaching[12,13]. Increasingly frequent mass bleaching events have been

observed over the past 4 decades[14,15] and are predicted to have impacted > 70% of the

world's coral reefs[16], leading to widespread coral mortality[17,18]. Understanding the extent of

variation in coral heat tolerance across phylogeny, space, and time is therefore crucial to

predict the future of these ecosystems[19-22].

Intraspecific variation in coral heat tolerance exists at local scales[23-29] and is an indicator of

adaptive capacity and metapopulation persistence probability[30,31]. Populations with low

intraspecific variation may suffer near extirpation at local scales[18,32,33]. On the contrary, high

intraspecific variation in physiological traits can buffer populations against the loss of genetic

and functional diversity[25]. Coral thermal tolerance traits are shaped by thermal history[34-36]

and other environmental conditions (e.g., irradiance[37], hydrodynamic regimes[38], nutrients[39]

and water oxygen content[40]). Coral colonies can adjust to these variable conditions through

phenotypic plasticity[41,42], heritable genetic effects[28,43,44], differences in symbiont

communities[27,45] or a combination of these three factors[46,47]. Yet, further research is required

to understand the relative importance of these drivers across environments, species and

regions.

Quantifying intraspecific variation in thermal thresholds during natural marine heatwaves is

challenged by differences in heat stress magnitude and duration over space and time and

the technical constraints of surveying large spatial areas during a single heatwave event.

Consequently, controlled experiments have been widely used to characterize intraspecific

variation in thermal tolerance among reefs[48-52], habitats[29,47,53], and individuals[54,55]. Recently,

acute heat stress assays (< 48 h) have been developed to measure individual thermal

thresholds in a standardized, high-throughput and cost-effective manner[48-51,53,56-60], yielding

consistent results across weeks[61], seasons[56] and with long-term heat stress[53,57,but see[60]). In

these experiments, coral heat tolerance has been mostly inferred from photosynthetic

efficiency (F_v/F_m), a common proxy of bleaching onset[62]. However, the role of symbiont

photochemical damage in bleaching initiation has been increasingly challenged[63,64] and can

be offset by host protective mechanisms[12,13,65] stressing the importance of relating F_v/F_m

measurements to subsequent coral bleaching responses (i.e., the loss of symbiont cells

and/or chlorophyll). To date, acute heat stress assays have investigated the response from a

few colonies at many reefs or many colonies at a single reef, but have not yet evaluated

intraspecific variation across multiple spatial scales.

To address the aforementioned knowledge gaps, we used acute heat stress assays to

measure photochemical efficiency and bleaching traits in 768 colonies of *Acropora*

spathulata from 14 reefs spanning 1,060 km (~9.5 degrees of latitude) of the Great Barrier

Reef (GBR; Figure 1a). *A. spathulata* is an abundant corymbose species found on reef flats

and upper slopes that builds ecologically important structural complexity on the GBR (Figure

1d) [66]. We identify reefs where heat tolerant individuals are most likely to be found and use

multiple environmental datasets (i.e., satellite observations, numerical models, in-situ

loggers) to determine environmental predictors of tolerance traits. The large sample size of

this study provides GBR-wide information to inform conventional management via spatial and

temporal protection and novel interventions, including seeding heat-tolerant corals onto

reefs.

Results

Acute heat stress assays were performed using standardized temperature treatments to elicit

increasing levels of heat stress of 0 °C (control), 3 °C, 6 °C, and 9 °C above the local

Maximum Monthly Mean (MMM)[67] (Figure 1b). Following exposure of replicate fragments of

A. spathulata colonies to each treatment, two phenotypic traits were measured: the

maximum photochemical efficiency of photosystem II (F_v/F_m) and hyperspectral imagery

estimate of chlorophyll content (normalized difference vegetation index; NDVI; Figure 1c).

The decline of each colony trait was modeled over temperature treatments using dose

response curves and was used to calculate two colony-level metrics of heat tolerance: the

temperature that resulted in a 50% decline in response (i.e., the median effective dose, or

ED50) which represents an absolute thermal threshold, and the performance under extreme

heat (+9 °C), which represents how coral traits decline relative to their local reef MMM.

The majority of *A. spathulata* colonies on the GBR showed minimal decline in photochemical

efficiency (F_v/F_m) and chlorophyll content (NDVI) at +3 °C (≤ 0.5%) and +6 °C (≤ 2.5%) above

their local MMM. However, under the more extreme temperature of +9 °C, these traits

declined by 2–80% (F_v/F_m) and 10–98% (NDVI), driving differences in thermal threshold

(ED50) and performance retention metrics among reefs and individuals (Figure 2).

Figure 1. Location of sample sites and acute heat stress design to quantify variation in the heat tolerance of the reef-building coral *Acropora spathulata* on the Great Barrier Reef. **a.** Map of the 14 reefs sampled across a latitudinal and thermal gradient. Isobaths of 0, 200, 1,000 and 2,000 m were obtained from the naturalearth 10 m resolution dataset and are represented in shades of blue. Coral reef locations were obtained from the UNEP database[68]. Sites labels are colored according to their maximum monthly mean (MMM) (dark red: > 28.5 °C, blue: < 27.5 °C, Table S1). **b.** A total of 768 coral colonies were measured in standardized acute heat stress experiments. 12 nubbins per colony were randomly divided among 3 tanks for each of 4 temperature treatments relative to the local MMM of each reef. **c.** Standardized target temperature profiles to elicit increasing levels of heat stress, following Evensen *et al.* [57]. Arrows indicate the timing of physiological measurements of the maximum quantum yield of photosystem II (F_v/F_m) and hyperspectral imagery (Hyp). **d.** Representative image of *A. spathulata* at Fitzroy Reef (© Mila Grinblat 2022).

Reef-level variation in coral heat tolerance

The GBR-wide medians of colony thermal thresholds were 37.1 ± 1.0 °C for F_v/F_m ED50 (median ± SD) and 36.1 ± 0.8 °C for NDVI ED50, with reef accounting for 34% and 25% of their respective total variation (ANOVA ω^2 effect sizes, Table S5b). Mean reef-level F_v/F_m ED50s differed by 1.8 °C and generally decreased with latitude (Figure 2a). For example, F_v/F_m ED50 thresholds in the southern GBR (Heron Island and Fitzroy Reef; 36.5–36.6 °C) were 0.8 °C lower than all reefs in the central and northern GBR (36.8–38.3 °C; *p.adj* < .0001; Table S5d). However, several reefs had higher (Kelso Reef, Pelorus Island, No Name Reef) or lower (Fitzroy Island, North Direction Island) ED50s than nearby reefs at similar

latitudes (*p.adj* < .05). Conversely, NDVI ED50 thresholds were less variable among reefs

(by 1.3 °C) and were lower at Martin Reef in the northern GBR than all other reefs (*p.adj* <

.01; Figure 2c). At St Crispin Reef, both F_v/F_m ED50 and NDVI ED50 were 0.8–2.6 °C lower

than all other reefs but are likely to be negatively biased due to experimental treatments not

reaching target temperatures (Table S2).

GBR-wide performance retention (performance under +9 °C relative to local conditions) was

46 ± 12% for F_v/F_m and 25 ± 19% for NDVI with reef accounting for 44% and 59% of their

respective total variation (ω² effect sizes, Table S5b). Performance retention was 2–41%

(F_v/F_m) and 60–65% (NDVI) higher at southern reefs than the rest of the GBR (Figure 2b and

Figure 2d), thus following an opposite trend to F_v/F_m ED50.

Figure 2. Variation in *Acropora spathulata* thermal tolerance among 14 reefs on the Great Barrier Reef (GBR) under acute experimental heat stress. The decline in maximum quantum yield (F_v/F_m) and total chlorophyll content (NDVI) was used to measure ED50 temperatures (a, c) and performance retention under extreme heat (i.e., +9 °C/MMM; b, d). Left panels display boxplots of the median (solid line) and first and third quartiles (box) for each reef colored by their MMM. Right panels show the distribution of values, with the GBR-wide median and standard deviation. Reefs are ordered (left to right) by increasing latitude. * Values for St Crispin Reef are negatively biased due to experimental treatments not reaching target temperatures.

Colony-level variation in coral heat tolerance

At the colony-level, thermal thresholds varied by up to 6°C, approximately 3 times the reef-level variation (1.8 °C). Within reefs, there was a 3.3 °C and 3.0 °C average range in colony-level ED50s for F_v/F_m and NDVI, respectively (Figure S1). The magnitude of this variation differed among reefs ($p < 0.001$, Levene's test; Table S5b) but no specific reef exhibited higher or lower variation for both traits. Colony-level variation in performance retention under +9 °C was also higher than reef-level variation (by 48% for F_v/F_m and by 17% for NDVI). Although colony-level ED50s were prone to uncertainty (median 95% CI range = 2.51 ± 3.34 °C and 2.99 ± 5.56 °C, Supplementary material and methods), congruent patterns between ED50 and performance retention support high variation in acute heat tolerance among and within reefs of the GBR.

Importantly, heat tolerant colonies were widely distributed across the GBR. For example, colonies that were ranked within the top 25% of F_v/F_m ED50 and NDVI ED50 occurred at virtually all reefs (Figure 3a, b). However, the most tolerant colonies were generally more abundant at central and northern reefs for F_v/F_m and southern reefs for NDVI. Additionally, several reefs exhibited a high proportion of tolerant individuals for both traits (e.g., Mackay Reef, Moore Reef, and Pelorus Island; Figure 3).

Response to acute heat stress differs between traits

The ranking of reef- and colony-level responses under heat stress differed between phenotypic traits. Overall, there was no GBR-wide correlation between F_v/F_m and NDVI

ED50s (ρ = 0.00, Figure 3c), but a moderate positive correlation between their performance

retention at +9 °C (ρ = 0.43, Figure 3d). At the reef-level, there was a moderate positive

correlation between F_v/F_m ED50 and NDVI ED50 at Chicken and No Name (ρ > 0.3) and a

weak positive correlation at Davies, Heron, Mackay, Martin and Moore (ρ = 0.1–0.3, Table

S6). Similarly, there was a weak to moderate positive correlation between the performance

retention of F_v/F_m and NDVI at all reefs (ρ = 0.1–0.3) except Chicken (ρ = 0.02), North

Direction (ρ = 0.84) and Pelorus Island (ρ = -0.22; Table S6).

Figure 3. Spatial distribution of *Acropora spathulata* acute heat tolerance on the Great
Barrier Reef measured by **a.** maximum quantum yield (F_v/F_m) and **b.** chlorophyll content
(NDVI). Pie charts at each reef represent the proportion of individual colony ED50s falling
between each quartile of the global distribution (blue: ED50 < Q1, yellow: Q1 < ED50 < Q2,
orange: Q2 < ED50 < Q3, red: Q3 <ED50). Pie charts radii are proportional to average
retained performance under extreme heat (i.e., +9 °C/MMM), normalized by the global range
for plotting purposes. Correlation between **c.** F_v/F_m and NDVI ED50s and **d**. retained

performance under extreme heat (+9 °C/MMM). Each point represents a distinct colony (n =

652 and 730 for c and d, respectively) and Spearman's-ρ and *p*-values are indicated.

Environmental drivers of acute heat tolerance metrics

To examine environmental conditions experienced by *A. spathulata* across the GBR, we

retrieved 10 environmental variables characterized by 24 quantitative predictors computed

from *in-situ* loggers, numerical models, and satellite observations (Table S7). A Principal

Component Analysis on environmental predictors divided the 768 colonies into three major

clusters (Figure S2). The first component (46.1% explained variance) separated cooler

southern reefs from warmer central and northern sites and was mainly driven by MMM (1.7

°C range) and Degree Heating Weeks (DHW) at the time of collection. The second

component (15.1% explained variance) separated northern and central sites along an

inshore-offshore gradient that was mainly driven by turbidity, sea surface current velocities,

and temperature variation (annual range and the rate of change from spring to summer).

Within clusters, colonies grouped by reefs as most predictors were retrieved at the site level

and were minimally separated by their water depth or pigmentation. Pigmentation scores

prior to collection differed between individuals and sites but had no clear association with

DHW at the time of collection (Figure S3).

Environmental drivers of heat tolerance traits (ED50 and performance retention in F_v/F_m and

NDVI) were evaluated using random forest ensemble learning and ridge regression on 12

low to moderately correlated predictors (pairwise absolute Pearson correlation = 0.01–0.69).

Together, these predictors explained a moderate to large proportion of variation in heat

stress responses of *A. spathulata* across the GBR (R^2 = 0.12–0.62). The predictive accuracy

of random forest (RF) and ridge regression (RR) was stronger when heat tolerance was

expressed for both F_v/F_m and NDVI as performance retention (R^2 = 0.27–0.62) than ED50

thresholds (R^2 = 0.12–0.31). Top environmental predictors slightly differed among traits and

metrics but variation in heat tolerance was primarily associated with site-level thermal history

(Figure 4). MMM (Figure 4b) and DHW at time of collection (Figure 4d) were predominant

metrics associated with response to acute heat stress (i.e., top RF predictors for 3 of the 4

heat tolerance metrics, Table S10a) while being strongly correlated to each other ($R = 0.78$).

The direction of associations differed between traits with MMM and DHW at the time of

collection being positively associated with F_v/F_m ED50 ($\beta = 0.06–0.07$), but negatively with

performance retention in F_v/F_m and NDVI and NDVI ED50 ($\beta = -0.03–-0.06$; Table S10b).

Secondary thermal history associations occurred between the annual range in temperature

and both performance retention metrics (F_v/F_m $\beta = -0.03$, NDVI $\beta = -0.04$; Figure 4c), and

between the frequency of DHW > 4 and both ED50 thresholds (F_v/F_m ED50 $\beta = 0.04$, NDVI

ED50 $\beta = -0.06$; Figure 4e). Particularly, the frequency of DHW > 4 in the year before

collection showed a strong negative association with NDVI ED50 ($\beta = -0.060$, top predictor in

RF), but no significant association with any other trait metric.

To a lesser extent, the heat tolerance of *A. spathulata* was associated with water flow,

shading, and depth. F_v/F_m tolerance metrics were generally higher at reefs with high sea

surface current velocity (F_v/F_m ED50 $\beta = 0.042$, +9/MMM $\beta = 0.037$) and lower at sites with

positive cloud anomalies in the 30 days prior to collection (F_v/F_m ED50 $\beta = -0.017$, +9/MMM β

= -0.031; Figure 4a). Turbidity showed little to no relationship with heat tolerance metrics. At

the colony level, increasing water depth was positively related to F_v/F_m metrics (F_v/F_m ED50:

$\beta = 0.042$, +9/MMM: $\beta = 0.004$), but negatively related to NDVI metrics (NDVI ED50: $\beta = -$

0.028, +9/MMM: $\beta = -0.015$). In addition, colonies with higher levels of pigmentation at the

time of collection tended to have higher NDVI ED50 thresholds ($\beta = 0.028$), but there was no

relationship with other tolerance metrics.

Figure 4. Environmental drivers of acute heat tolerance of *Acropora spathulata* on the Great Barrier Reef. Heat tolerance was measured as the maximum quantum yield of photosystem II (F_v/F_m) and chlorophyll content (NDVI) expressed as ED50 temperature thresholds and performance retention under extreme heat (+9 °C). Relationships between environmental predictors (defined in Table S7) and heat tolerance metrics were assessed using ridge regression and random forest models. **a.** Heatmap of associations evaluated with random forest variable importance (10-fold stratified cross validation, circle size) and ridge regression standardized coefficients (multiple bootstrap resampling $n = 100$, circle color). The variance explained by each model (random forest/ridge R^2) is shown above each tolerance trait metric. **b-e**. Thermal history maps of (**b**) maximum monthly mean temperature as defined by NOOA[67] (**c**) annual temperature range (2014–2022), (**d**) peak degree heating weeks during sampling and (**e**) annual frequency of degree heating weeks above 4 since 1985. White dots indicate reef sites sampled in this study.

Discussion

The large geographic scope of this study (14 reefs across 9.5 degrees of latitude) and sample size ($n = 768$) revealed extensive intraspecific variation in the heat tolerance of reef-building corals across the Great Barrier Reef (GBR). This resulted in more than 6 °C variation in colony thermal thresholds where tolerant colonies — based on multiple traits and metrics — were widely distributed among and within reefs. Environmental predictors of heat-tolerance variation support adaptation and/or acclimatization of populations to local

conditions while unexplained differences among colonies may be due to host and/or

symbiont adaptive genetic variation within reefs.

Acute heat tolerance metrics linked to historical and recent environmental conditions

All heat tolerance metrics were most strongly related to thermal history supporting adaptation

of corals to their local thermal regimes[35,36,69,70]. These results also corroborate ecological

experiments that found 4 °C difference in *A. spathulata* larvae survival thresholds between

Lizard Island (14.7 °S) and One Tree Island (23.5 °S) i.e., the same latitudinal range as our

study[71]. F_v/F_m ED50 thresholds increased at reefs exposed to higher thermal averages

(MMM), annual thermal variation (AR) and exposure to marine heat waves (frequency of

DHW > 4 and standard deviation of DHW). DHW at the time of sampling was also strongly

positively associated with F_v/F_m ED50, suggesting that acclimatization of the holobiont to

higher temperatures possibly through upregulation of photoprotective mechanisms can

increase the heat tolerance trait F_v/F_m ED50 as reported by Cunning *et al.* [56]. Conversely,

there were weaker unexpected negative associations between all thermal history metrics and

NDVI ED50. This may reflect accumulated heat stress leading to early health impairment of

the holobiont and lower chlorophyll retention, which is supported by the strong negative

relationship of NDVI ED50 with frequency of DHW >4 in the 12 months prior to sampling.

This is similar to F_v/F_m ED50 findings of Marzonie *et al.* [49] in the Coral Sea, suggesting that

some reef populations may be impacted over long time periods after heat exposure.

Differences in the association with thermal history among traits could thus occur from a

decoupling between host and symbiont baseline conditions and/or stress responses

(discussed below)[63,64,65]. Nevertheless, several reefs in our study (No Name, North Direction,

Mackay) exhibited a high proportion of tolerant individuals for both F_v/F_m and NDVI ED50,

showing that some colonies and populations may perform better under heat stress across

multiple traits[72]. These populations will be more likely to play a key role in the GBR resilience

in a warming climate and thus their mapping in space and time could provide valuable

information for spatial planning of Australia's marine parks into the future.

The heat tolerance metrics of *A. spathulata* measured here were further associated with

predictors that influence water movement, light, and within-reef temperature. Strong water

movement can delay photochemical damage under thermal stress through coral surface

cooling, increases in respiration and metabolic transfers and removal of toxins[38,73] and the

positive association between surface current velocity and F_v/F_m metrics suggests a carryover

of these effects to our experiments. Solar irradiance is also a factor known to mediate

bleaching responses under high seawater temperatures[74] and F_v/F_m ED50 thresholds were

indeed reduced at reefs with a high cloud cover prior to the experiments. This may indicate

that symbionts acclimated to lower light levels[75] experienced greater irradiance-induced

photosynthetic damage in the experimental system[76]. Overall, these results demonstrate that

empirically derived thermal tolerance metrics are not only shaped by long-term adaptation

but also acclimatization to recent conditions.

Standing within-reef variation in heat tolerance supports adaptive potential to climate change

There was greater variation in the acute heat tolerance of *A. spathulata* colonies within reefs

than among reef-level averages (ED50 ~3.3 vs 1.8 °C and ~3.0 vs 1.6 °C for F_v/F_m and NDVI

respectively), despite most colonies being sampled from a narrow depth range at a single

site less than 600 m^2. Importantly, we found that heat-tolerant corals (defined as the top 25

percentile from the GBR ED50 distributions) occurred at nearly every reef, including the

cooler southern GBR. These results align with previous results on *A. cervicornis* in the

Florida Keys[56], *A. hyacinthus* in Palau[51] and *A. hyacinthus* in the GBR[77]. Within-reef variation

remained mostly unexplained by colony-level predictors (depth and pigmentation) and could

be elucidated by incorporating genomic data not yet available here. Individual differences in

heat tolerance may be underpinned by genomic variation within the host[23,28], symbionts[45,78]

and/or their interactions[55]. For example, incorporating the proportion of *Durusdinium* cells

and host polygenic scores based on putative heat-adaptive loci increased the predictive

accuracy of GBR *A. millepora* bleaching models by ~22%[79]. In addition, deciphering the

independent role of environmental drivers requires a better characterization across

microhabitats[80] and depth[81] as well as optimization of sampling strategies across

environmental gradients to reduce the collinearity of predictors[82]. The large standing variation

in thermal tolerance at the reef scale reported in this study has important implications for the

conservation of this species on the GBR. It may notably support *A. spathulata* adaptation to

climate change through selection of pre-existing heat tolerance alleles[20]. Furthermore, rather

than translocating individuals from warmer reefs (e.g., Howells *et al.* [69]) restoration projects

can target local heat tolerant individuals when their phenotypes are known or by sampling a

diversity of genets.

Coral populations from the Southern GBR may live further from their upper thermal limit

Acute thermal thresholds (ED50) for *A. spathulata* populations on the GBR were 9.5–9.6 °C

above their local MMM on southern reefs compared to 7.5–8.8 °C on central and northern

reefs. This is consistent with higher performance retention under extreme heat on southern

reefs and latitudinal differences in the thermal thresholds of *A. spathulata* larvae[83].

Consequently, under comparable levels of thermal stress, southern populations are expected

to be less susceptible to bleaching than central and northern populations. In line with this

expectation, southern GBR reefs have experienced less frequent and intense bleaching

events than central and northern reefs[15] with 90% less bleaching-related mortality during the

1985–2012 period[17]. The lower frequency and magnitude of bleaching events on southern

reefs might also allow more time for adaptation to occur through the migration of heat-

tolerance associated alleles from warmer northern reefs[20].

Conversely, the low cumulative heat stress experienced by those regions may result in coral

assemblages that are naïve to high heat stress[15]. As such, mild but highly prevalent coral

bleaching occurred in the southern GBR during 2016–2017 (21% of colonies; Kennedy et al.

[84]) and 2020 (86% of colonies; Page et al. [85]; 48% of colonies; Nolan et al. [86]) and occurred

under lower cumulative heat stress (DHW) than in central and northern GBR[15]. Further

investigations are thus required to understand how recurrent warming disturbances and gene

flow with northern regions will shape the evolution of these cooler reefs.

Divergent effects of acute heat stress on thermal tolerance traits

Here we found a moderate correlation between F_v/F_m and NDVI performance retention under

extreme heat. This supports the use of photochemical apparatus integrity as an early proxy

of coral bleaching[53,57,62] and the importance of symbiont tolerance in bleaching response. For

instance, some symbiont species or genotypes may produce more photoprotective pigments

or antioxidant compounds, alleviating oxidative stress for the host [12,13]. Conversely, we found

no correlation between ED50s of F_v/F_m and NDVI traits. The only two other studies using

F_v/F_m and chlorophyll-related ED50 thresholds found congruent reef-level variation between

the two traits for some species (e.g., Pocillopora verrucosa[48] and *Acropora hyacinthus[77])* but

not others (e.g., Porites lobata[48]). In both studies, visual chlorophyll scores showed higher

noise and lower differences in reef ED50s which aligns with *A. spathulata* pairwise reef

differences in NDVI ED50s (0.39 °C) being almost half that of F_v/F_m ED50 (0.67 °C). This

may also be due to differences in the timing of measurement between traits as their decline

occurs at different rates[63]. In *A. tenuis*, F_v/F_m values were found to be stable 0–24 hours after

the end of an acute heat stress, while chlorophyll decreased up to 24 hours after the end of

temperature ramp-down[87]. Therefore delaying the timing of NDVI measurements in our study

may have revealed higher divergence in chlorophyll retention between reefs. Finally, algal

oxidative stress and photosynthetic damage can be alleviated by antioxidant mechanisms[13]

or immune response from the host[65] and thus occur without a decrease in chlorophyll content

or symbiont density[88].

Divergent responses between traits highlight the importance of defining heat tolerance and

selecting phenotypic traits accordingly. Chlorophyll content may be an appropriate proxy for

holobiont bleaching and mortality as its decline during an acute heat stress can persist and

reflect mortality in the next month[61]. However, photosynthetic efficiency may be a better

proxy to detect sub-lethal effects as corals can experience physiological stress of reduced

tissue biomass and symbiont loss long before changes in pigmentation are visually

detectable[63]. Finally, which phenotypic trait under acute experimental stress would best

predict heat tolerance in natural conditions remains unclear and could vary with marine

heatwave magnitude and duration as well as the biological level at which they are

assessed[60]. Our results demonstrate that no single trait or metric can fully capture the

complexity of the coral holobiont heat stress response, and highlight the importance of

measuring multiple traits whenever achievable, including relevant host traits (e.g., antioxidant

capacity).

Conclusion

Using standardized acute heat stress assays, we found that heat tolerant corals were widely

distributed across the GBR, including cooler, southern reefs. This suggests potential for

adaptive responses to climate change, if heat tolerance traits have a heritable basis[31,44].

Rankings of coral heat tolerance can guide human interventions and their interpretation and

application may depend on the specific traits and metrics measured. Corals with high

absolute thermal tolerance can help to decipher heat-stress protective mechanisms[23,89] and

be used as material for assisted gene flow through translocation [59,90] and selective

breeding[43]. On the contrary, corals with high thermal tolerance relative to local conditions

may be better targets for local propagation and breeding, decreasing the risk of carrying

pathogens and phenotype-environment mismatches[91]. Nonetheless, natural adaptive

processes and human interventions must be accompanied by reduced anthropogenic

emissions through ambitious national and international commitments to secure the

persistence of species and populations across the entire GBR.

Methods

Study sites, experimental design and setup

Study sites and sample collection

We quantified thermal tolerance traits of 768 colonies of *A. spathulata* across 14 reefs of the

GBR (Figure 1a, Table S1). At each reef, 40–60 colonies were sampled on SCUBA from one

or two sites (covering 50–4200 m^2) between 28 February and 26 March 2022 (GBRMPA

permit G21/45166.1) over a single or two consecutive days. For each colony, 12 nubbins (~8

cm) were collected for standardized acute heat stress experiments and *in-situ* metadata was

recorded. This included colony-level GPS coordinates linked to *in-situ* photographic

records[92], time-corrected depth (-0.54 ± 0.76 m relative to the lowest astronomical tide) and

visual pigmentation (Coral Watch Health Chart scores at 0.5 intervals)[93]. Colonies showing

signs of disease were rarely encountered but intentionally excluded to avoid physiological

bias unrelated to temperature effects.

Acute heat stress assay design

A portable automated experimental system ("Seasim-in-a-box", National Sea Simulator, XXX;

Figure S4) was modified from Marzonie *et al.* [49] to conduct acute heat stress assays onboard

a research vessel the day following collection (Supplementary materials and methods). The

experimental design consisted of three replicate tanks for each of four temperature

treatments of 0 °C (control), 3 °C, 6 °C, and 9 °C above the local Maximum Monthly Mean

(MMM, 1985–1990+1993 climatology as defined by NOOA[67]) for each reef site. These

treatments were designed to elicit increasing levels of thermal stress following Voolstra *et al.*

[53]. Immediately after collection, 12 nubbins from each coral colony were mounted on

separate experimental racks with unique identifiers and held overnight at ambient seawater

temperature. The following morning, one nubbin from each colony was randomly assigned to

each of the 12 tanks (Figure 1b). Assays started at 11:00 am and consisted of 3 hours ramp-

up from ambient temperature (within 1 °C of the MMM) to target temperature, 3 hours hold at

the target temperature, 2 hours ramp down to MMM, and a final ~12-h hold at MMM (Figure

1c). During the experiments, temperatures in individual tanks were recorded at a 1-min

intervals using Hobo loggers (Onset) and closely matched their target profiles (mean$_{\Delta T}$ = 0.36

°C, Table S2) with the exception of one site (St Crispin: mean$_{\Delta T}$ = 1.61 °C).

Phenotypic traits

Photochemical efficiency

The maximum quantum yield of photosystem II (F_v/F_m) of coral photosymbionts

(Symbiodiniaceae) was used as an initial rapid, non-invasive measure of heat tolerance. A

decline in F_v/F_m reflects early physiological impairment in corals and has been reported to

respond consistently between acute- and long-term heat exposures[53,57]; but see Klepac *et al.*

[60]. Measurements were taken 2 hours after the end of temperature ramp-down (> 1-hour dark

adaptation) using an Imaging PAM chlorophyll fluorometer (IMAG-K7, Walz Germany) with

the following settings: Measuring Light = 2 (freq = 1), Saturation Pulse = 7 (Int = 30s), Damp

= 1, Gain = 1. For each fragment, three measurements were extracted from non-overlapping

areas and underwent several quality check and filtration steps (Supplementary materials and

methods). After filtration, a total of 769 colonies and 8,453 fragments were retained in further

analyses.

Hyperspectral image-based assessment of chlorophyll content

Non-invasive estimates of chlorophyll content were used as a second metric of heat

tolerance. The loss of chlorophyll underpins visual bleaching scores commonly used in acute

heat stress experiments and has been shown to correspond with mortality risk[61]. Total

chlorophyll content was assessed in fragments from reflectance measurements taken 11

hours after the end of the temperature ramp-down using a hyperspectral camera (Resonon,

Pika XC2) with the following settings: Integration time = 39.2ms, Gain = 0, Frame rate = 22

fps. MATLAB software was used to compute the normalized difference vegetation index

(NDVI, Supplementary material and methods) from reflectance measurements, where NDVI

$= (R_{720}-R_{670})/(R_{720}+R_{670})$ and R_x is the reflectance at x nm. NDVI is a proxy for chlorophyll

used in a wide range of organisms including scleractinian corals[94] and has been validated in

the soft coral *Sarcophton* cf. *glaucum*[95]. We repeated this validation in *A. spathulata* using

spectrophotometric determination from tissue extractions[96] with a strong relationship between

NDVI and log-transformed chlorophyll-a (R^2=0.74, Supplementary material and methods,

Figure S5).

Phenotypic data analysis

The response of raw F_v/F_m and NDVI to temperature across experiments was assessed

using a linear mixed effect model incorporating fixed effects of site, treatment, fragment size

(estimated from pixel numbers in RGB images) and random intercepts for each colony and

tank using package *lme4*[97], including 95% bootstrap confidence intervals for fixed effects

(Table S3). Tanks were first treated as a block factor in the analysis. Despite small variations

in temperatures among replicate tanks, tank had a minor effect on fragment performances

under heat stress (explained 1.44% of F_v/F_m random variance σ_i^2 and 0.8% of NDVI σ_i^2,

linear mixed effect model, Table S3). Likewise, fragment size effect appears to be negligible

for F_v/F_m (B = 1.4×10^{-6}, df = 7473, 95% CI: -4.8×10^{-7}–3.0×10^{-6}) and very weak for NDVI

(B = 8.5×10^{-6}, df = 8988, 95% CI:6.14×10^{-6}–1.0×10^{-5}).

ED50 thermal thresholds

For each colony, the decrease in F_v/F_m and NDVI was modeled over temperature treatments

using dose response curves in the R package *drc*[98]. Colonies with <2 fragments per

treatment after previous quality control filtering (Supplementary material and methods) were

excluded to ensure robust estimates of individual responses across treatments (excluded

colonies: F_v/F_m = 103, NDVI = 4). All models were fitted using the *drm* function based on the

mean hold temperature recorded within each tank for each experiment, with constraints set

on parameters (Table S4a). The best model (Weibull type II with 3 parameters) was chosen

as the one with the lowest Akaike's information criterion score (AIC) for most reef sites using

the *mselect* function (6/14 and 11/14 reefs for F_v/F_m and NDVI, respectively; Table S4b).

Because using the same model is a prerequisite to compare ED50s, we used this model for

all colonies even though some curves showed a better fit to log-logistic or quadratic models.

The model equation is

$$f(x; b, d, e) = d(1 - \exp\left(-\exp\left(b(\log(x) - \log(e))\right)\right))$$

where *b* is steepness of the curve, *d* the upper asymptote (lower asymptote set to 0) and *e*

the inflexion point. An example of dose response curves can be found in Figure S6. Dose

response curves were filtered to remove individuals that showed poor fit to the data, notably

when the decline in phenotypic traits was minor or absent up to +6 °C or +9 °C resulting in

ED50s with wide confidence intervals (Supplementary material and methods). After filtration

652 and 725 colony-level ED50 estimates were retained for F_v/F_m and NDVI respectively.

480   Performance retention under thermal stress

481   Performance retention under extreme heat was computed as

$$Xi_{MMM+9/MMM} = (Xi_{MMM+9}/Xi_{MMM})\, x\, ((Tmean_{MMM+9C} - Tmean_{MMM})/9)$$

where Xi is the trait (F_v/F_m or NDVI) averaged across fragments of individual i in the MMM

and MMM+9 treatments, respectively, and $Tmean_{MMM+9C}$ and $Tmean_{MMM}$ are the average hold

temperatures for these treatments. The second term accounts for small differences between

target and effective temperatures across tanks and experiments. Performance retention can

be computed even for colonies that experienced minor decline in phenotypic traits under +9

°C which prevented fitting a dose-response curve. After filtration of colonies with < 2

fragments in the MMM and MMM+9C treatments, 730 and 765 colony-level performance

retention estimates were calculated for F_v/F_m and NDVI respectively.

Heat tolerance variation analysis

The variation in heat tolerance (ED50 and performance retention) among and within sites

was investigated using an analysis of variance (ANOVA). Homogeneity of variance between

sites was assessed using Levene's test on residuals from groups medians (Table S5b) and

Tukey's HSD post-hoc tests for pairwise comparisons (Table S5c). Since variance was

unequal across sites, we used one-way Welch's ANOVA to compare site means and Games-

Howell post-hoc tests with adjusted p-values at α = 0.05 for pairwise comparisons (Table

S5d). Following Cunning et $al.$ [56], we assessed within-site variation in heat tolerance by

computing ED50 adjusted as the grand mean of the total distribution plus residuals from site.

Finally, we investigated the similarity of colony thermal tolerance rankings between F_v/F_m and

NDVI traits using Spearman's rank correlation as assumption of normality was violated. For

both ED50 and performance retention, we computed both GBR-wide and site-level

correlations (St Crispin Reef excluded; Table S6).

Environmental data acquisition and analysis

To examine environmental conditions experienced by *A. spathulata* across the GBR, we

retrieved 10 environmental variables characterized by 24 predictors computed from *in-situ*

loggers, numerical models, and satellite observations (Table S7). Reef thermal history was

characterized using the NOAA Coral Watch v3.1 sea surface temperature (SST) satellite

product (1985–present; 5km)[99] for thermal predictors based on historical climatology (MMM,

DHW) as it has the largest temporal coverage. For contemporary thermal predictors (e.g.,

annual average and thermal range, 2014–present) we used the eReefs GBR1 Hydro

model[100] as it delivers temperature predictions at a finer spatial resolution (~1 km, hourly)

and higher accuracy for the last decade (ME = 0.4–0.9 °C)[101]. Temperatures from both

sources were validated using *in-situ* records (Supplementary material and methods).

Global averages (2010–2019) of environmental variables that may play a secondary role in

heat tolerance (e.g., chlorophyll, turbidity, current velocity) were retrieved from eReefs GBR4

model (hourly; 4 km)[102]. In addition, we included the average cloud fraction (% of pixel

masked by clouds) and cloud fraction anomaly for the 30 days prior to collection[37] retrieved

from NASA-MODIS 1°, monthly database[103].

Our environmental dataset also included variables measured at the individual colony level.

Depths obtained from dive computers were adjusted to tide level at the time of collection

(standardized to Lowest Astronomical Tide, Figure S7). Predictors derived from eReefs were

computed at each colony depth through vertical interpolation from the model z-levels. As the

GBR experienced a marine heatwave during the sampling campaign (0.47–4.67 DHW), we

included pigmentation levels prior to collection as an explanatory variable to account for heat

stress experienced in the weeks preceding the experiment (Figure S3).

Principal Component Analysis (PCA) was performed on environmental predictors for the 768

colonies to visualize the distribution of sites and colonies along environmental gradients.

Phenotype by environment analysis

The influence of environmental predictors on heat tolerance traits (ED50 and performance

retention in F_v/F_m and NDVI) was evaluated using random forest ensemble learning and ridge

regression on a set of 12 low to moderately correlated predictors (pairwise absolute Pearson

correlation = 0.01–0.69; Figure S8, Table S9; St Crispin Reef excluded). Because any

association of heat tolerance traits with MMM and DHW at the time of collection may reflect

different mechanisms (e.g., adaptation vs acclimatization), both were retained in the dataset

despite their strong correlation ($R = 0.78$).

Random forest models were built using *cforest* function from *party* R package[104]. For each of

the 4 heat tolerance metrics, separate random forests were grown to 1,000 trees (ntree) with

5 environmental predictors tried at each split (mtry). Predictor importance and model

accuracy were assessed through repeated 10-fold stratified (across sites) cross-validation

(70/30 split, Supplementary material and methods). The importance of each predictor was

estimated by computing the marginal increase in out-of-bag sample MSE (Mean Inc MSE)

when training the model with the predictors randomly shuffled.

Ridge regression was performed for each phenotypic trait using the *glmnet* R package[105].

The optimal tuning parameter λ of the penalty term was selected using k-fold cross-validation

(λ.1se; Figure S9 and Figure S10). Model accuracy (R^2) and environmental predictor

coefficients were estimated using bootstrap resampling (repeated 100-fold stratified cross

validation, Figure S11 and Table S10).

All statistical analysis were conducted using R v4.0.4 and figures created using package

ggplot2 and Inkscape v1.2.

References

1.   Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., and Shukla, P.R.
(2022). Global Warming of 1.5° C: IPCC Special Report on Impacts of Global
Warming of 1.5° C above Pre-industrial Levels in Context of Strengthening Response
to Climate Change, Sustainable Development, and Efforts to Eradicate Poverty.
Cambridge University Press.
2.   Angilletta, M.J., Wilson, R.S., Navas, C.A., and James, R.S. (2003). Tradeoffs and
the evolution of thermal reaction norms. Trends in Ecology & Evolution *18*, 234-240.
3.   Bennett, S., Duarte, C.M., Marbà, N., and Wernberg, T. (2019). Integrating within-
species variation in thermal physiology into climate change ecology. Philosophical
Transactions of the Royal Society B *374*, 20180550.
4.   Reusch, T.B. (2014). Climate change in the oceans: evolutionary versus
phenotypically plastic responses of marine animals and plants. Evolutionary
applications *7*, 104-122.
5.   Somero, G. (2010). The physiology of climate change: how potentials for
acclimatization and genetic adaptation will determine 'winners' and 'losers'. Journal of
Experimental Biology *213*, 912-920.
6.   Anthony, K.R.N., Helmstedt, K.J., Bay, L.K., Fidelman, P., Hussey, K.E., Lundgren,
P., Mead, D., McLeod, I.M., Mumby, P.J., Newlands, M., et al. (2020). Interventions to
help coral reefs under global change-A complex decision challenge. PLoS One *15*,
e0236399. 10.1371/journal.pone.0236399.
7.   Lachs, L., Humanes, A., Mumby, P.J., Donner, S.D., Bythell, J.C., Beauchamp, E.,
Bukurou, L., Buzzoni, D., de la Torre Cerro, R., and East, H.K. (2023). Unexpectedly
high coral heat tolerance at thermal refugia. bioRxiv, 2023.2006. 2016.545328.
8.   Sinervo, B., Mendez-De-La-Cruz, F., Miles, D.B., Heulin, B., Bastiaans, E., Villagrán-
Santa Cruz, M., Lara-Resendiz, R., Martínez-Méndez, N., Calderón-Espinosa, M.L.,
and Meza-Lázaro, R.N. (2010). Erosion of lizard diversity by climate change and
altered thermal niches. Science *328*, 894-899.
9.   Hughes, T.P., Kerry, J.T., Álvarez-Noriega, M., Álvarez-Romero, J.G., Anderson,
K.D., Baird, A.H., Babcock, R.C., Beger, M., Bellwood, D.R., and Berkelmans, R.
(2017). Global warming and recurrent mass bleaching of corals. Nature *543*, 373-377.
10.   LaJeunesse, T.C., Parkinson, J.E., Gabrielson, P.W., Jeong, H.J., Reimer, J.D.,
Voolstra, C.R., and Santos, S.R. (2018). Systematic revision of Symbiodiniaceae
highlights the antiquity and diversity of coral endosymbionts. Current Biology *28*,
2570-2580. e2576.

11. Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P.,
Gomez, E., Harvell, C.D., Sale, P.F., Edwards, A.J., and Caldeira, K. (2007). Coral
reefs under rapid climate change and ocean acidification. Science *318*, 1737-1742.
12. Oakley, C., and Davy, S. (2018). Cell biology of coral bleaching. In Coral bleaching,
(Springer), pp. 189-211.
13. Roberty, S., and Plumier, J.-C. (2022). Bleaching physiology: who's the 'weakest
link'—host vs. symbiont? Emerging Topics in Life Sciences *6*, 17-32.
14. Hughes, T. (2020). Spatial and temporal patterns of mass bleaching of corals in the
Anthropocene. Science 359, 80–83 (2018). Scientific Reports *10*, 20346.
15. Hughes, T.P., Kerry, J.T., Connolly, S.R., Álvarez-Romero, J.G., Eakin, C.M., Heron,
S.F., Gonzalez, M.A., and Moneghetti, J. (2021). Emergent properties in the
responses of tropical corals to recurrent climate extremes. Current Biology *31*, 5393-
5399. e5393.
16. Virgen-Urcelay, A., and Donner, S.D. (2023). Increase in the extent of mass coral
bleaching over the past half-century, based on an updated global database. PloS
One *18*, e0281719.
17. De'ath, G., Fabricius, K.E., Sweatman, H., and Puotinen, M. (2012). The 27–year
decline of coral cover on the Great Barrier Reef and its causes. Proceedings of the
National Academy of Sciences *109*, 17995-17999.
18. Riegl, B., Johnston, M., Purkis, S., Howells, E., Burt, J., Steiner, S.C., Sheppard,
C.R., and Bauman, A. (2018). Population collapse dynamics in Acropora downingi, an
Arabian/Persian Gulf ecosystem-engineering coral, linked to rising temperature.
Global change biology *24*, 2447-2462.
19. Dixon, A.M., Forster, P.M., Heron, S.F., Stoner, A.M.K., and Beger, M. (2022). Future
loss of local-scale thermal refugia in coral reef ecosystems. PLOS Climate *1*,
e0000004. 10.1371/journal.pclm.0000004.
20. Matz, M.V., Treml, E.A., and Haller, B.C. (2020). Estimating the potential for coral
adaptation to global warming across the Indo-West Pacific. Global Change Biology
26, 3473-3481.
21. Sully, S., Hodgson, G., and van Woesik, R. (2022). Present and future bright and
dark spots for coral reefs through climate change. Global Change Biology.
22. Cornwall, C.E., Comeau, S., Donner, S.D., Perry, C., Dunne, J., van Hooidonk, R.,
Ryan, S., and Logan, C.A. (2023). Coral adaptive capacity insufficient to halt global
transition of coral reefs into net erosion under climate change. Global Change
Biology.
23. Drury, C., Bean, N.K., Harris, C.I., Hancock, J.R., Huckeba, J., Roach, T.N., Quinn,
R.A., and Gates, R.D. (2022). Intrapopulation adaptive variance supports thermal
tolerance in a reef-building coral. Communications Biology *5*, 1-10.
24. Drury, C., Martin, R.E., Knapp, D.E., Heckler, J., Levy, J., Gates, R.D., and Asner,
G.P. (2022). Ecosystem-scale mapping of coral species and thermal tolerance.
Frontiers in Ecology and the Environment.
25. McWilliam, M., Madin, J.S., Chase, T.J., Hoogenboom, M.O., and Bridge, T.C.
(2022). Intraspecific variation reshapes coral assemblages under elevated
temperature and acidity. Ecology Letters.
26. Cunning, R., Ritson-Williams, R., and Gates, R.D. (2016). Patterns of bleaching and
recovery of Montipora capitata in Kāne 'ohe Bay, Hawai 'i, USA. Marine Ecology
Progress Series *551*, 131-139.
27. Sampayo, E., Ridgway, T., Bongaerts, P., and Hoegh-Guldberg, O. (2008). Bleaching
susceptibility and mortality of corals are determined by fine-scale differences in
symbiont type. Proceedings of the National Academy of Sciences *105*, 10444-10449.
28. Jin, Y.K., Lundgren, P., Lutz, A., Raina, J.-B., Howells, E.J., Paley, A.S., Willis, B.L.,
and Van Oppen, M.J. (2016). Genetic markers for antioxidant capacity in a reef-
building coral. Science Advances *2*, e1500842.
29. Marhoefer, S.R., Zenger, K.R., Strugnell, J.M., Logan, M., van Oppen, M.J., Kenkel,
C.D., and Bay, L.K. (2021). Signatures of adaptation and acclimatization to reef flat

and slope habitats in the coral Pocillopora damicornis. Frontiers in Marine Science *8*,
704709.

30.   Holstein, D.M., Smith, T.B., van Hooidonk, R., and Paris, C.B. (2022). Predicting coral
metapopulation decline in a changing thermal environment. Coral Reefs, 1-12.

31.   Howells, E.J., Bay, L.K., and Bay, R.A. (2022). Identifying, Monitoring, and Managing
Adaptive Genetic Variation in Reef-Building Corals under Rapid Climate Warming.
Coral Reef Conservation and Restoration in the Omics Age, 55-70.

32.   Hughes, T.P., Kerry, J.T., Baird, A.H., Connolly, S.R., Dietzel, A., Eakin, C.M., Heron,
S.F., Hoey, A.S., Hoogenboom, M.O., and Liu, G. (2018). Global warming transforms
coral reef assemblages. Nature *556*, 492-496.

33.   Álvarez-Noriega, M., Marrable, I., Noonan, S.H., Barneche, D.R., and Ortiz, J.C.
(2023). Highly conserved thermal performance strategies may limit adaptive potential
in corals. Proceedings of the Royal Society B *290*, 20221703.

34.   Schoepf, V., Sanderson, H., and Larcombe, E. (2021). Coral heat tolerance under
variable temperatures: Effects of different variability regimes and past environmental
history vs. current exposure. Limnology and Oceanography.

35.   Thompson, D.M., and Van Woesik, R. (2009). Corals escape bleaching in regions
that recently and historically experienced frequent thermal stress. Proceedings of the
Royal Society B: Biological Sciences *276*, 2893-2901.

36.   Hughes, T.P., Kerry, J.T., Connolly, S.R., Baird, A.H., Eakin, C.M., Heron, S.F., Hoey,
A.S., Hoogenboom, M.O., Jacobson, M., and Liu, G. (2019). Ecological memory
modifies the cumulative impact of recurrent climate extremes. Nature Climate Change
9, 40-43.

37.   Gonzalez-Espinosa, P.C., and Donner, S.D. (2021). Cloudiness reduces the
bleaching response of coral reefs exposed to heat stress. Global Change Biology.

38.   Page, C.E., Leggat, W., Heron, S.F., Choukroun, S.M., Lloyd, J., and Ainsworth, T.D.
(2019). Seeking resistance in coral reef ecosystems: the interplay of biophysical
factors and bleaching resistance under a changing climate: the interplay of a reef's
biophysical factors can mitigate the coral bleaching response. BioEssays *41*,
1800226.

39.   Wiedenmann, J., D'Angelo, C., Smith, E.G., Hunt, A.N., Legiret, F.-E., Postle, A.D.,
and Achterberg, E.P. (2013). Nutrient enrichment can increase the susceptibility of
reef corals to bleaching. Nature Climate Change *3*, 160-164.

40.   Alderdice, R., Perna, G., Cárdenas, A., Hume, B.C., Wolf, M., Kühl, M., Pernice, M.,
Suggett, D.J., and Voolstra, C.R. (2022). Deoxygenation lowers the thermal threshold
of coral bleaching. Scientific Reports *12*, 1-14.

41.   DeCarlo, T.M., Harrison, H.B., Gajdzik, L., Alaguarda, D., Rodolfo-Metalpa, R.,
D'Olivo, J., Liu, G., Patalwala, D., and McCulloch, M.T. (2019). Acclimatization of
massive reef-building corals to consecutive heatwaves. Proceedings of the Royal
Society B *286*, 20190235.

42.   Hackerott, S., Martell, H.A., and Eirin-Lopez, J.M. (2021). Coral environmental
memory: causes, mechanisms, and consequences for future reefs. Trends in Ecology
& Evolution.

43.   Howells, E.J., Abrego, D., Liew, Y.J., Burt, J.A., Meyer, E., and Aranda, M. (2021).
Enhancing the heat tolerance of reef-building corals to future warming. Science
advances *7*, eabg6070.

44.   Bairos-Novak, K.A.-O., Hoogenboom, M.A.-O., van Oppen, M.A.-O., and Connolly,
S.A.-O. (2021). Coral adaptation to climate change: Meta-analysis reveals high
heritability across multiple traits. Global Change Biology *27*, 5694-5710.

45.   Berkelmans, R., and Van Oppen, M.J. (2006). The role of zooxanthellae in the
thermal tolerance of corals: a 'nugget of hope'for coral reefs in an era of climate
change. Proceedings of the Royal Society B: Biological Sciences *273*, 2305-2312.

46.   Howells, E.J., Abrego, D., Meyer, E., Kirk, N.L., and Burt, J.A. (2016). Host adaptation
and unexpected symbiont partners enable reef-building corals to tolerate extreme
temperatures. Global change biology *22*, 2702-2714.

47. Thomas, L., Rose, N.H., Bay, R.A., López, E.H., Morikawa, M.K., Ruiz-Jones, L., and
Palumbi, S.R. (2018). Mechanisms of thermal tolerance in reef-building corals across
a fine-grained environmental mosaic: lessons from Ofu, American Samoa. Frontiers
in Marine Science *4*, 434.
48. Evensen, N.R., Voolstra, C.R., Fine, M., Perna, G., Buitrago-López, C., Cárdenas, A.,
Banc-Prandi, G., Rowe, K., and Barshis, D.J. (2022). Empirically derived thermal
thresholds of four coral species along the Red Sea using a portable and standardized
experimental approach. Coral Reefs, 1-14.
49. Marzonie, M.R., Bay, L.K., Bourne, D.G., Hoey, A.S., Matthews, S., Nielsen, J.J., and
Harrison, H.B. (2022). The effects of marine heatwaves on acute heat tolerance in
corals. Global Change Biology.
50. Voolstra, Valenzuela, J.J., Turkarslan, S., Cárdenas, A., Hume, B.C., Perna, G.,
Buitrago-López, C., Rowe, K., Orellana, M.V., and Baliga, N.S. (2021). Contrasting
heat stress response patterns of coral holobionts across the Red Sea suggest distinct
mechanisms of thermal tolerance. Molecular ecology *30*, 4466-4480.
51. Cornwell, B., Armstrong, K., Walker, N.S., Lippert, M., Nestor, V., Golbuu, Y., and
Palumbi, S.R. (2021). Widespread variation in heat tolerance and symbiont load are
associated with growth tradeoffs in the coral Acropora hyacinthus in Palau. Elife *10*,
e64790.
52. Shaw, E.C., Carpenter, R.C., Lantz, C.A., and Edmunds, P.J. (2016). Intraspecific
variability in the response to ocean warming and acidification in the scleractinian coral
Acropora pulchra. Marine Biology *163*, 1-11.
53. Voolstra, C.R., Buitrago-López, C., Perna, G., Cárdenas, A., Hume, B.C., Rädecker,
N., and Barshis, D.J. (2020). Standardized short-term acute heat stress assays
resolve historical differences in coral thermotolerance across microhabitat reef sites.
Global Change Biology *26*, 4328-4343.
54. Humanes, A., Lachs, L., Beauchamp, E.A., Bythell, J.C., Edwards, A.J., Golbuu, Y.,
Martinez, H.M., Palmowski, P., Treumann, A., and van der Steeg, E. (2022). Within-
population variability in coral heat tolerance indicates climate adaptation potential.
Proceedings of the Royal Society B *289*, 20220872.
55. Kavousi, J., Denis, V., Sharp, V., Reimer, J.D., Nakamura, T., and Parkinson, J.E.
(2020). Unique combinations of coral host and algal symbiont genotypes reflect
intraspecific variation in heat stress responses among colonies of the reef-building
coral, Montipora digitata. Marine Biology *167*, 1-15.
56. Cunning, R., Parker, K.E., Johnson-Sapp, K., Karp, R.F., Wen, A.D., Williamson,
O.M., Bartels, E., D'Alessandro, M., Gilliam, D.S., and Hanson, G. (2021). Census of
heat tolerance among Florida's threatened staghorn corals finds resilient individuals
throughout existing nursery populations. Proceedings of the Royal Society B *288*,
20211613.
57. Evensen, N.R., Fine, M., Perna, G., Voolstra, C.R., and Barshis, D.J. (2021).
Remarkably high and consistent tolerance of a Red Sea coral to acute and chronic
thermal stress exposures. Limnology and Oceanography.
58. Barshis, D.J., Ladner, J.T., Oliver, T.A., Seneca, F.O., Traylor-Knowles, N., and
Palumbi, S.R. (2013). Genomic basis for coral resilience to climate change.
Proceedings of the National Academy of Sciences *110*, 1387-1392.
59. Morikawa, M.K., and Palumbi, S.R. (2019). Using naturally occurring climate resilient
corals to construct bleaching-resistant nurseries. Proceedings of the National
Academy of Sciences *116*, 10586-10591.
60. Klepac, C.N., Petrik, C., Karabelas, E., Owens, J., Hall, E.R., and Muller, E.M. (2023).
Assessing Acute Thermal Assays as a Rapid Screening Tool for Coral Restoration.
Research Square. https://doi.org/10.21203/rs.3.rs-3140553/v1.
61. Walker, N.S., Cornwell, B.H., Nestor, V., Armstrong, K.C., Golbuu, Y., and Palumbi,
S.R. (2022). Persistence of phenotypic responses to short-term heat stress in the
tabletop coral Acropora hyacinthus. PLOS ONE *17*, e0269206.

62. Jones, R.J., Ward, S., Amri, A.Y., and Hoegh-Guldberg, O. (2000). Changes in
quantum efficiency of Photosystem II of symbiotic dinoflagellates of corals after heat
stress, and of bleached corals sampled after the 1998 Great Barrier Reef mass
bleaching event. Marine and Freshwater Research *51*, 63-71.
63. Gardner, S.G., Raina, J.-B., Nitschke, M.R., Nielsen, D.A., Stat, M., Motti, C.A.,
Ralph, P.J., and Petrou, K. (2017). A multi-trait systems approach reveals a response
cascade to bleaching in corals. BMC biology *15*, 1-14.
64. Krueger, T., Hawkins, T.D., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O.,
Leggat, W., Fisher, P.L., and Davy, S.K. (2015). Differential coral bleaching—
Contrasting the activity and response of enzymatic antioxidants in symbiotic partners
under thermal stress. Comparative Biochemistry and Physiology Part A: Molecular &
Integrative Physiology *190*, 15-25.
65. Baird, A.H., Bhagooli, R., Ralph, P.J., and Takahashi, S. (2009). Coral bleaching: the
role of the host. Trends in Ecology & Evolution *24*, 16-20.
66. Wallace, C.C., Done, B.J., and Muir, P.R. (2012). Revision and catalogue of
worldwide staghorn corals Acropora and Isopora (Scleractinia: Acroporidae) in the
Museum of Tropical Queensland.
67. Liu, G., Strong, A.E., and Skirving, W. (2003). Remote sensing of sea surface
temperatures during 2002 Barrier Reef coral bleaching. Eos, Transactions American
Geophysical Union *84*, 137-141.
68. UNEP-WCMC, W.-C. (2010). WRI, & TNC.(2010). Global distribution of warm-water
coral reefs, compiled from multiple sources including the Millennium Coral Reef
Mapping Project. Version *1*.
69. Howells, E.J., Berkelmans, R., van Oppen, M.J., Willis, B.L., and Bay, L.K. (2013).
Historical thermal regimes define limits to coral acclimatization. Ecology *94*, 1078-
1088.
70. Sully, S., Burkepile, D.E., Donovan, M., Hodgson, G., and Van Woesik, R. (2019). A
global analysis of coral bleaching over the past two decades. Nature communications
10, 1-5.
71. Woolsey, E.S., Byrne, M., and Baird, A.H. (2013). The effects of temperature on
embryonic development and larval survival in two scleractinian corals. Marine
Ecology Progress Series *493*, 179-184.
72. Lachs, L., Humanes, A., Pygas, D.R., Bythell, J.C., Mumby, P.J., Ferrari, R., Figueira,
W.F., Beauchamp, E., East, H.K., and Edwards, A.J. (2023). No apparent trade-offs
associated with heat tolerance in a reef-building coral. Communications Biology *6*,
400.
73. Nakamura, T., Van Woesik, R., and Yamasaki, H. (2005). Photoinhibition of
photosynthesis is reduced by water flow in the reef-building coral Acropora digitifera.
Marine Ecology Progress Series *301*, 109-118.
74. Lesser, M.P., and Farrell, J.H. (2004). Exposure to solar radiation increases damage
to both host tissues and algal symbionts of corals during thermal stress. Coral reefs
23, 367-377.
75. Anthony, K.R., and Hoegh-Guldberg, O. (2003). Kinetics of photoacclimation in
corals. Oecologia *134*, 23-31.
76. Roberty, S., Fransolet, D., Cardol, P., Plumier, J.-C., and Franck, F. (2015).
Imbalance between oxygen photoreduction and antioxidant capacities in
Symbiodinium cells exposed to combined heat and high light stress. Coral Reefs *34*,
1063-1073.
77. Naugle, M. S., Denis, H., Mocellin, V. J. L., Laffy, P. W., Popovic, I., Bay. L. K.,
Howells, E. J. (2024). Environmental, host, and symbiont drivers of heat tolerance in
a species complex of reef-building corals. *BioRxiv.* [Preprint.]. Available from:
https://doi.org/10.1101/2024.01.31.575130.
78. Palacio-Castro, A.M., Smith, T.B., Brandtneris, V., Snyder, G.A., van Hooidonk, R.,
Maté, J.L., Manzello, D., Glynn, P.W., Fong, P., and Baker, A.C. (2023). Increased

dominance of heat-tolerant symbionts creates resilient coral reefs in near-term ocean
warming. Proceedings of the National Academy of Sciences *120*, e2202388120.

79. Fuller, Z.L., Mocellin, V.J., Morris, L.A., Cantin, N., Shepherd, J., Sarre, L., Peng, J.,
Liao, Y., Pickrell, J., and Andolfatto, P. (2020). Population genetics of the coral
Acropora millepora: Toward genomic prediction of bleaching. Science *369*.

80. Ainsworth, T.D., Leggat, W., Silliman, B.R., Lantz, C.A., Bergman, J.L., Fordyce, A.J.,
Page, C.E., Renzi, J.J., Morton, J., and Eakin, C.M. (2021). Rebuilding relationships
on coral reefs: Coral bleaching knowledge-sharing to aid adaptation planning for reef
users: Bleaching emergence on reefs demonstrates the need to consider reef scale
and accessibility when preparing for, and responding to, coral bleaching. BioEssays,
2100048.

81. Wyatt, A.S., Leichter, J.J., Washburn, L., Kui, L., Edmunds, P.J., and Burgess, S.C.
(2023). Hidden heatwaves and severe coral bleaching linked to mesoscale eddies
and thermocline dynamics. Nature Communications *14*, 1-17.

82. Selmoni, O., Vajana, E., Guillaume, A., Rochat, E., and Joost, S. (2020). Sampling
strategy optimization to increase statistical power in landscape genomics: A
simulation-based approach. Molecular ecology resources *20*, 154-169.

83. Woolsey, E., Keith, S., Byrne, M., Schmidt-Roach, S., and Baird, A. (2015).
Latitudinal variation in thermal tolerance thresholds of early life stages of corals. Coral
Reefs *34*, 471-478.

84. Kennedy, E.V., Ordoñez, A., and Diaz-Pulido, G. (2017). Coral bleaching in the
southern inshore Great Barrier Reef: a case study from the Keppel Islands. Marine
and Freshwater Research *69*, 191-197.

85. Page, C.A., Giuliano, C., Bay, L.K., and Randall, C.J. (2023). High survival following
bleaching underscores the resilience of a frequently disturbed region of the Great
Barrier Reef. Ecosphere *14*, e4280.

86. Nolan, M.K., Schmidt-Roach, S., Davis, A.R., Aranda, M., and Howells, E.J. (2021).
Widespread bleaching in the One Tree Island lagoon (Southern Great Barrier Reef)
during record-breaking temperatures in 2020. Environmental monitoring and
assessment *193*, 1-8.

87. Nielsen, J., Matthews, G., Frith, K., Harrison, H., Marzonie, M., Slaughter, K.,
Suggett, D., and Bay, L. (2022). Experimental considerations of acute heat stress
assays to quantify coral thermal tolerance. Scientific reports *12*, 16831.

88. Middlebrook, R., Anthony, K.R., Hoegh-Guldberg, O., and Dove, S. (2012). Thermal
priming affects symbiont photosynthesis but does not alter bleaching susceptibility in
Acropora millepora. Journal of Experimental Marine Biology and Ecology *432*, 64-72.

89. Cleves, P.A., Tinoco, A.I., Bradford, J., Perrin, D., Bay, L.K., and Pringle, J.R. (2020).
Reduced thermal tolerance in a coral carrying CRISPR-induced mutations in the gene
for a heat-shock transcription factor. Proceedings of the National Academy of
Sciences *117*, 28899-28905.

90. Quigley, K.M., Bay, L.K., and van Oppen, M.J. (2019). The active spread of adaptive
variation for reef resilience. Ecology and evolution *9*, 11122-11135.

91. Sellares Blasco, R.I., Villalpando, M.F., Guendulain-Garcia, S.D., and Cróquer, A.
(2021). Assisted coral reproduction in the Dominican Republic: a successful story to
replicate in the Caribbean. Frontiers in Marine Science *8*, 927.

92. Lukoschek, V., Riginos, C., and van Oppen, M.J. (2016). Congruent patterns of
connectivity can inform management for broadcast spawning corals on the Great
Barrier Reef. Molecular ecology *25*, 3065-3080.

93. Siebeck, U., Marshall, N., Klüter, A., and Hoegh-Guldberg, O. (2006). Monitoring
coral bleaching using a colour reference card. Coral Reefs *25*, 453-460.

94. Wijgerde, T., van Melis, A., Silva, C.I., Leal, M.C., Vogels, L., Mutter, C., and Osinga,
R. (2014). Red light represses the photophysiology of the scleractinian coral
Stylophora pistillata. PLoS one *9*, e92781.

95. Leal, M.C., Jesus, B., Ezequiel, J., Calado, R., Rocha, R.J.M., Cartaxana, P., and
Serôdio, J. (2015). Concurrent imaging of chlorophyll fluorescence, Chlorophyll a

content and green fluorescent proteins-like proteins of symbiotic cnidarians. Marine
Ecology *36*, 572-584.
96. Ritchie, R.J. (2008). Universal chlorophyll equations for estimating chlorophylls a, b,
c, and d and total chlorophylls in natural assemblages of photosynthetic organisms
using acetone, methanol, or ethanol solvents. Photosynthetica *46*, 115-126.
97. Bates, D., Mächler, M., Bolker, B., and Walker, S. (2014). Fitting linear mixed-effects
models using lme4. arXiv preprint arXiv:1406.5823.
98. Ritz, C., Baty, F., Streibig, J.C., and Gerhard, D. (2015). Dose-response analysis
using R. PloS one *10*, e0146021.
99. NOAA Coral Reef Watch (2018, updated daily). *NOAA Coral Reef Watch Version*
3.1, Daily Global 5-km Satellite Sea Surface Temperature Product (a.k.a CoralTemp),
Jun. 1, 1985-July. 3, 2022. College Park, Maryland, USA: NOAA Coral Reef Watch.
Data set accessed 2022-07-04 at
https://coralreefwatch.noaa.gov/product/5km/index_5km_sst.php.
100. CSIRO (2020). eReefs GBR 1 hydrodynamic model version 2. Data set accessed
2022-07-13 at https://extraction.ereefs.aims.gov.au/.
101. Steven, A.D., Baird, M.E., Brinkman, R., Car, N.J., Cox, S.J., Herzfeld, M., Hodge, J.,
Jones, E., King, E., and Margvelashvili, N. (2019). eReefs: An operational information
system for managing the Great Barrier Reef. Journal of Operational Oceanography
12, S12-S28.
102. CSIRO (2020). eReefs GBR 4 hydrodynamic model version 2. Data set accessed
2022-04-22 at https://extraction.ereefs.aims.gov.au/.
103. Savtchenko, A., Ouzounov, D., Ahmad, S., Acker, J., Leptoukh, G., Koziana, J., and
Nickless, D. (2004). Terra and Aqua MODIS products available from NASA GES
DAAC. Advances in Space Research *34*, 710-714.
104. Hothorn, T., Hornik, K., Strobl, C., Zeileis, A., and Hothorn, M.T. (2015). Package
'party'. Package Reference Manual for Party Version 0.9-998 *16*, 37.
105. Friedman, J., Hastie, T., Tibshirani, R., Narasimhan, B., Tay, K., Simon, N., and Qian,
J. (2021). Package 'glmnet'. CRAN R Repositary.

Data and code availability statement

890 Scripts and data used for the analyses are available online: https://github.com/hde08/Aspat-

891 GBR-HeatTolVar/tree/main. Data is publicly archived on [Anonymous Repository].