***In vivo* CRISPR screens identify novel virulence genes among proteins of**

unassigned subcellular localization in *Toxoplasma*

Yuta Tachibana[1,2], Miwa Sasai[1,2,3], and Masahiro Yamamoto[1,2,3]*

[1]Department of Immunoparasitology, Research Institute for Microbial Diseases,

[2]Laboratory of Immunoparasitology, WPI Immunology Frontier Research Center,

[3]Department of Immunoparasitology, Center for Infectious Disease Education and

Research, Osaka University, Suita, Osaka 565-0871, Japan

Running title: *In vivo* CRISPR screens on hyperLOPIT-unassigned genes

Key words: *Toxoplasma, CRISPR screen, hyperLOPIT-unassigned virulence factors*

*Correspondence:

Masahiro Yamamoto

(E-mail: myamamoto@biken.osaka-u.ac.jp)

ABSTRACT

      The research field to identify and characterize virulence genes in *Toxoplasma gondii* has been dramatically advanced by a series of *in vivo* CRISPR screens. Although subcellular localizations of thousands of proteins were predicted by the spatial proteomic method called hyperLOPIT, those of more than 1000 proteins remained unassigned and their essentiality in virulence was also unknown. In this study, we generated two small-scale gRNA libraries targeting approximately 600 hyperLOPIT-unassigned proteins and performed *in vivo* CRISPR screens. As a result, we identified several *in vivo* fitness-conferring genes that were previously unreported. We further characterized two candidates, TgGTPase and TgRimM, which are localized in the cytoplasm and the apicoplast, respectively. Both genes are essential for parasite virulence and widely conserved in the phylum Apicomplexa. Collectively, our current study provides a resource for estimating the *in vivo* essentiality of *Toxoplasma* proteins with previously unknown localizations.

IMPORTANCE

      Toxoplasma gondii is a protozoan parasite that causes severe infection in immunocompromised patients or newborns. *Toxoplasma* possesses more than 8000 genes; however, the genes that determine parasite virulence were not fully identified. The apicomplexan parasites, including *Toxoplasma*, developed unique organelles that do not exist in other model organisms; thus, determining the subcellular location of parasite proteins is important for understanding their functions. Here, we used *in vivo* CRISPR genetic screens that enabled us to investigate hundreds of genes in *Toxoplasma* during mouse infection. We screened approximately 600 parasite proteins with previously unknown subcellular localizations. We identified many novel genes that confer parasite fitness in mice. Among the top hits, we characterized two essential virulence genes, TgGTPase and TgRimM, which are widely conserved in the phylum Apicomplexa. Our findings will contribute to understanding how apicomplexans adapt to the host environment and cause disease.

INTRODUCTION

An obligate intracellular parasite, *Toxoplasma gondii*, is an important human and animal pathogen that causes life-threatening toxoplasmosis (1). *Toxoplasma* has been the model apicomplexan parasite owing to its easiness of *in vitro* cultivation, genetic tractability, *in vivo* disease model in mice, and various biological assays to investigate the function of the genes (2). As unicellular eukaryotes that adapt to intracellular parasitism, apicomplexans have developed unique secretory organelles (e.g., micronemes, rhoptries, dense granules), which are absent in other well-studied eukaryotic cell systems (3). These secretory organelles are key features of the apicomplexans and discharge various types of proteins that are essential for invasion, replication, and egress (3). Likewise, the apicoplast is a unique non-photosynthetic plastid-like organelle and essential for apicomplexans' metabolism (fatty acid, heme, iron-sulfur cluster, and isoprenoid synthesis) (4). Therefore, uncovering the subcellular localizations of parasite proteins is crucial to understand the protein functions and parasite pathogenesis mechanism.

Although the parasite protein localization is typically revealed by cell biological methods such as classical immunofluorescence assay (IFA), immuno-electron microscopy, and, recently, ultrastructure expansion microscopy (U-ExM) (5, 6), the parasite spatial proteome has been largely unknown. Recent studies elegantly addressed the issues in *Toxoplasma* (7) and *Cryptosporidium* (8) by a spatial proteomic method called hyperLOPIT (9), yielding plenty of knowledge on parasite subcellular proteome. In a previous study using hyperLOPIT on *Toxoplasma* (7), approximately 3800 proteins were identified in the tachyzoite stage; however, about 1,200 of them are still classified as unassigned subcellular localizations. Moreover, the functions of these genes mostly remained to be elucidated.

In addition to classical forward and reverse genetics, *in vitro* or *in vivo* CRISPR genetic screens have been used to assess the essentiality of the parasite genes in the lytic cycles or virulence (10-15). We previously performed *in vivo* CRISPR screens using small-scaled guide RNA (gRNA) libraries based on hyperLOPIT-assigned localizations (e.g., rhoptries, dense granules, nucleus, endoplasmic reticulum (ER), Golgi apparatus, cytosol, apicoplast, and mitochondrion) and identified novel secreted and non-secreted virulence factors (14). However, the essentiality of hyperLOPIT-unassigned proteins in

virulence remains mostly unknown.

In this study, we generated two small-scaled gRNA libraries targeting genes
encoding hyperLOPIT-unassigned proteins and performed *in vivo* CRISPR screens in
mice. We identified several novel genes that contribute to the parasite's *in vivo* fitness.
Among the top hits, we further focused on two candidates, TgGTPase (TGGT1_277840)
and TgRimM (TGGT1_321310). TgGTPase is a guanosine triphosphatase (GTPase) that
localizes in the parasite cytosol, and TgRimM is a novel apicoplast-resident protein
possessing a ribosome maturation factor M (RimM) domain and a C2H2-type zinc finger
domain. Both genes are widely conserved among the apicomplexans, and gene deletions
led to the complete loss of virulence in mice, suggesting that TgGTPase and TgRimM are
essential for the parasite adaptation to the *in vivo* environment. Overall, our current study
provides a clue for the research community to estimate the *in vivo* essentiality of proteins
with previously unknown localizations.

RESULTS

***In vivo* CRISPR screening targeting proteins of unassigned subcellular localization**

To identify genes necessary for *Toxoplasma* survival in mice, we utilized a
previously established *in vivo* CRISPR screen platform using type I strain and C57BL/6
mice (14). We newly generated two gRNA libraries (Unassigned_1 and Unassigned_2
libraries), each targeting 295 genes classified as unassigned localization in hyperLOPIT
(7) and control genes for *in vitro* and *in vivo* selection. Among approximately 1200
hyperLOPIT-unassigned proteins, we excluded genes with previously published *in vitro*
fitness scores in human foreskin fibroblasts (HFF) less than -1.5 to reduce the size of
the gRNA library (10). Transfected RHΔhxgprt parasites were selected with
pyrimethamine in Vero cells for four passages to generate a pooled mutant population
(*in vitro* sample). The parasite mutant pools were used to infect C57BL/6 mice with 10^7
parasites each by injection into the footpad. After seven days post infection, parasites
were recovered from the spleen and expanded in Vero cells for one passage (*in vivo*
sample). The gRNA sequences were amplified by PCR from the input library and the
genomic DNA extracted from the parasites of *in vitro* and *in vivo* samples. The gRNAs
were sequenced by next-generation sequencing to determine their relative abundance.

We calculated *in vitro* and *in vivo* fitness scores of each gene as the average $\log_2$ fold
change of guide abundances between conditions. We analyzed the screen results from
both sublibraries to assess the reproducibility **(Tables S1 and S2)**. The *in vitro* essential
and dispensable controls were successfully separated, with lower scores for the essential
genes and higher scores for the dispensable genes (**Figure S1A**). The correlations
between our *in vitro* fitness scores and the genome-wide *in vitro* fitness scores in HFFs
were high (r = 0.78 and 0.76, respectively) (**Figure S1B**) (10). High reproducibility of
in vivo fitness scores was observed between independent infections in mice (r =
0.75±0.05 and 0.68±0.07, respectively) (**Figure S1C**). These data demonstrated that our
in vitro and *in vivo* CRISPR screens using Unassigned_1 and Unassigned_2 libraries
were highly reproducible.
To identify genes that confer parasite *in vivo* fitness during infection in mice,
we compared *in vitro* and *in vivo* fitness scores (**Figure 1A, Tables S1 and S2**). To
rank the candidate genes, we calculated the distance of each gene from the regression
line (**Figure 1B**). As reported previously (14), the *in vivo* essential controls ROP18 and
GRA23 showed negative *in vivo* fitness scores and ranked highly (**Figure 1B**). This
analysis highlighted both previously identified and unidentified genes. For instance, the
top hits from Unassigned_1 library contain hypothetical protein (TGGT1_211850),
thioredoxin (Txn: TGGT1_224060), NUDIX hydrolase (Nudix: TGGT1_282190),
CAAX metallo endopeptidase (Caax: TGGT1_221170), vacuolar iron transporter (VIT:
TGGT1_266800), DNA damage inducible protein 1 (DDI1: TGGT1_304680),
TgGTPase (TGGT1_277840), GLT2 (TGGT1_239752), apical annuli methyltransferase
(AAMT: TGGT1_310070), and mRNA cleavage factor family protein (mRNACF:
TGGT1_221190). It has been reported that VIT mainly localizes to the plant-like
vacuolar compartment (PLVAC) and regulates parasite iron metabolism (16). DDI1 is
one of the components of the parasite ubiquitin-proteasome system and is localized in
the cytoplasm and nucleus (17). TgGTPase was identified as an interactor of the Nd
complex, which facilitates rhoptry exocytosis (18). GLT2 is a glucosyltransferase and
required for parasite disaccharide metabolism (19). AAMT was identified as an apical
annuli component (20). The top hits from Unassigned_2 library also contain several
hypothetical proteins (TGGT1_300220, TGGT1_312840, TGGT1_277870,

TGGT1_250790, TGGT1_221640, and TGGT1_273905), ribosome maturation factor

RimM domain-containing protein (TgRimM: TGGT1_321310), TBC6

(TGGT1_237280), DnaJ domain-containing protein (Dnaj: TGGT1_203380), CPSF A

subunit region protein (CPSF: TGGT1_ 267710), and rRNA-processing protein

EFG1(EFG1: TGGT1_240660). Although TGGT1_300220 was also shown to be one of

the top hits in another *in vivo* CRISPR screen (15), no further investigation was

performed. TBC6 is one of the TBC-domain containing proteins and is localized in the

ER or cytoplasmic vesicles (21). TGGT1_273905 localizes to the cytoplasm (22).

Although some candidates (VIT and DDI1) were previously reported to be required for

parasite virulence (16, 17), none of the other top hits are assessed regarding virulence so

far.

Identification of novel genes that contribute to parasite virulence in mice

We generated individual knockout strains to examine whether these top-hit

genes affect parasite virulence (**Figures S2A and S2B**). WT mice were locally infected

with these mutant strains into the footpad, and the survival rates were monitored. We

chose footpad infection rather than intraperitoneal infection because our previous study

revealed that the former is more suitable for assessing virulence in using RH strain (14)

(**Figures 2A and 2B**). Among the candidates from Unassigned_1 library, all mice

survived in the infection of ΔTGGT1_211850, ΔDDI1, ΔTgGTPase, or ΔAAMT

parasites. Other tested mutants also showed reduced virulence (ΔGLT2, ΔCaax,

ΔmRNACF, ΔTxn, ΔNudix, and ΔVIT). Among the candidates from Unassigned_2

library, all mice survived in the infection of ΔTGGT1_300220, ΔTgRimM, ΔEFG1,

ΔCPSF, or ΔTGGT1_277870 parasites. ΔTGGT1_250790, ΔDnaj, ΔTGGT1_312840,

or ΔTGGT1_273905 parasites showed reduced virulence. WT mice infected with

ΔTBC6, and ΔTGGT1_221640 succumbed in a time course similar to those infected

with WT parasites. Collectively, these data demonstrated that our *in vivo* CRISPR

screens targeting hyperLOPIT-unassigned proteins highlighted known and novel *in vivo*

fitness-conferring genes that affect virulence.

TgGTPase is a widely conserved GTPase that contributes to virulence

Among the top-ranking candidates from Unassigned_1 library, we chose to

characterize TgGTPase further (**Figure 3A and Tables S1**). TgGTPase is a GTPase that

is broadly conserved across the phylum Apicomplexa (**Figures 3B and 3C**). To study

the function of TgGTPase, we complemented the ΔTgGTPase strain with C-terminal

HA-tagged TgGTPase expressed from the tubulin promoter that we will refer to as

"ΔTgGTPase::TgGTPase-HA" (**Figure 3D**). Western blot showed the band of

TgGTPase-HA at around 40 kDa (**Figure 3E**). The HA-tagged TgGTPase showed a

cytoplasmic localization by IFA (**Figure 3F**). To validate the essentiality of TgGTPase

on pathogenesis, we infected C57BL/6 mice intraperitoneally with 10^3 tachyzoites of

the WT, ΔTgGTPase, and ΔTgGTPase::TgGTPase-HA parasites and assessed mouse

survival (**Figure 3G**). All the mice intraperitoneally infected with ΔTgGTPase parasites

survived. By contrast, ΔTgGTPase::TgGTPase-HA parasites fully restored virulence as

well as WT parasites. Taken together, these results suggest that TgGTPase is a

cytoplasmic protein essential for virulence.

TgRimM is a novel apicoplast-resident protein possessing C2H2 zinc finger that

contributes to virulence in mice

Among the top-ranking candidates from Unassigned_2 library, we chose to

characterize TgRimM further (**Figure 4A and Tables S2**). TgRimM possesses a signal

peptide, a ribosome maturation factor RimM domain, and a single C2H2-type zinc finger

domain (**Figure 4B**) (23, 24). We searched for TgRimM homologs and found that they

are broadly conserved across the phylum Apicomplexa except for *Cryptosporidium* spp.

(**Figure 4C**). To characterize TgRimM, we complemented the ΔTgRimM strain with C-

terminal HA-tagged TgRimM expressed from the tubulin promoter that we will refer to

as "ΔTgRimM::TgRimM-HA" (**Figure 4D**). The HA-tagged TgRimM showed an

apicoplast localization co-stained with streptavidin (an apicoplast marker) by IFA (**Figure**

4E). Western blot against TgRimM-HA showed the main band at around 75 kDa,

probably a processed form of TgRimM (**Figure 4F**). To check the effect of TgRimM on

virulence, we infected C57BL/6 mice intraperitoneally with 10^3 tachyzoites of the WT,

ΔTgRimM, and ΔTgRimM::TgRimM-HA parasites and assessed the mouse survival. All

the mice intraperitoneally infected with ΔTgRimM parasites survived. In contrast,

ΔTgRimM::TgRimM-HA parasites restored virulence (**Figure 4G**). Next, we assessed

the importance of the C2H2-type zinc finger domain in TgRimM. We generated a

complemented strain lacking the C2H2 zinc finger (ΔTgRimM::TgRimM$^{\Delta ZF}$-HA)

(**Figure 4C**). Deleting the zinc finger domain did not alter the apicoplast localization of

TgRimM$^{\Delta ZF}$-HA (**Figure S3**). However, ΔTgRimM::TgRimM$^{\Delta ZF}$-HA could not restore

virulence (**Figure 4G**). Thus, these results suggest that TgRimM is essential for virulence

in mice and that the C2H2 zinc finger is indispensable for TgRimM function in the

apicoplast.

DISCUSSION

Despite much effort from the research community, many apicomplexan proteins

still need to be annotated owing to their uniqueness and absence in other model organisms.

The CRISPR screens enabled the researchers to investigate hundreds to thousands of

genes in *Toxoplasma* at once *in vitro* culture or *in vivo* infection to identify essential genes

for parasite growth, metabolism, drug resistance, differentiation, host-pathogen

interaction, and virulence (10-15, 25-33). Most *in vivo* CRISPR screens in *Toxoplasma*

have been focused on identifying secreted virulence effectors such as rhoptry bulb

proteins (ROPs) and dense granule proteins (GRAs) (11-13). However, recent studies are

in progress to search for non-secreted virulence factors by *in vivo* CRISPR screens (14,

15). In this study, we focused on the hyperLOPIT-unassigned proteins and performed *in*

vivo CRISPR screens to identify novel virulence factors in *Toxoplasma*. As a result, we

identified many candidate genes that were required for the parasite *in vivo* fitness in mice

and further characterized TgGTPase and TgRimM among them.

TgGTPase was originally reported as an interactor of the Nd complex, which

plays an important role in rhoptry secretion (18). However, considering that our study

revealed its distribution in the cytosol, TgGTPase might have broader functions than

rhoptry secretion. TgGTPase homologs are widely conserved across the phylum

Apicomplexa, suggesting that this GTPase is functionally important across the species in

the phylum. In the transposon mutagenesis screen in *Plasmodium falciparum* (34),

PfGTPase (PF3D7_1348100) possesses lower mutagenesis scores (Mutagenesis Index

Score = 0.316 and Mutagenesis Fitness Score = -2.437), suggesting that PfGTPase is

essential for optimal growth of asexual blood-stages *in vitro*. Future studies on TgGTPase
homologs in *Plasmodium* or *Cryptosporidium* will reveal their conserved function in the
phylum.

We identified TgRimM as an apicoplast-resident protein. A homolog in *P.*
falciparum (PfRimM: PF3D7_1032000) is also predicted to be localized in the apicoplast
by signal peptide prediction (35). In the transposon mutagenesis screen in *P. falciparum*
(34), PfRimM also possesses very low mutagenesis scores (Mutagenesis Index Score =
0.158 and Mutagenesis Fitness Score = -2.8), indicating that PfRimM is essential for
growth of asexual blood-stages *in vitro*. Assuming that RimM homologs in the phylum
Apicomplexa (ApiRimM) are located in the apicoplast, the lack of RimM homologs in
Cryptosporidium is reasonable because *Cryptosporidium* spp. are known to have lost the
apicoplast (36-38). RimM domain-containing protein was reported to regulate 16S rRNA
processing in bacteria (39). It is reported that a RimM domain-containing protein in
Arabidopsis thaliana (AtRimM) is located in the chloroplasts and contributes to rRNA
maturation and proteostasis (40). Considering that the apicoplast evolved from secondary
endosymbiosis (41) and that the zinc finger domain in TgRimM is essential for its
function in our study, ApiRimM might facilitate rRNA maturation and proteostasis in the
apicoplast. Previous studies on *Toxoplasma* apicoplast have shown that apicoplast
function is essential for the parasite *in vitro* survival and growth (42-44). Moreover,
apicoplast-assigned proteins in hyperLOPIT showed a small number of *in vitro*
dispensable proteins (7), also supporting its bare essentiality. In sharp contrast, other than
TgRimM, we previously identified two additional apicoplast-resident proteins (EF-P:
TGGT1_258380 and hypothetical protein: TGGT1_204350) that are dispensable for *in*
vitro fitness but essential for *in vivo* fitness in CRISPR screens (14). The presence of this
type of apicoplast proteins that are dispensable *in vitro* but indispensable *in vivo* might
imply unknown characteristics of the apicoplast. Considering that the apicoplast is a
potent drug target against toxoplasmosis and malaria (45), the presence of such apicoplast
proteins that are dispensable *in vitro* but indispensable *in vivo* might expand the range of
anti-toxoplasmosis and -malarial drug targets in the future. In combination with the
database of hyperLOPIT, *in vivo* CRISPR screens succeeded in identifying such "new
class" apicoplast proteins essential for virulence.

Limitations of our current study are that, although approximately 1,200 proteins

are classified as unassigned subcellular localizations in hyperLOPIT, our sublibraries

cover only half of them. Therefore, the remaining proteins still need to be addressed in

the future. Also, although we identified previously unreported virulence genes in this

study, their precise functions and localizations, other than TgGTPase and TgRimM,

mostly remained unaddressed. Future works will aim to characterize these remaining

proteins further.

In conclusion, our *in vivo* CRISPR screen platform provides a resource for the

community to identify virulence genes encoding proteins with previously unknown

localizations and functions.

MATERIALS AND METHODS

***Toxoplasma* strains**

RHΔhxgprt (14), RHΔhxgprtΔku80 (46) and its derivatives of *Toxoplasma* were

maintained in Vero cells and passaged every 3 days using RPMI (Nacalai Tesque)

supplemented with 2 % heat-inactivated fetal bovine serum (FBS; JRH Bioscience), 100

U/ml penicillin and 0.1 mg/ml streptomycin (Nacalai Tesque) in incubators at 37°C and

5% CO_2.

Host cell culture

Vero cells were maintained in RPMI (Nacalai Tesque) supplemented with 10 %

heat-inactivated FBS, 100 U/ml penicillin and 0.1 mg/ml streptomycin (Nacalai Tesque)

in incubators at 37°C and 5% CO_2. Human foreskin fibroblasts (HFFs) were maintained

in DMEM (Nacalai Tesque) supplemented with 10% heat-inactivated FBS, 100 U/ml

penicillin and 0.1 mg/ml streptomycin (Nacalai Tesque) in incubators at 37°C and 5%

CO_2.

Mice

C57BL/6NCrSlc (C57BL/6N) mice were purchased from SLC. All experiments

were conducted in 8-10-week-old female mice. All animal experiments were conducted

with the approval of the Animal Research Committee of Research Institute for Microbial

Diseases in Osaka University.

300 _In vitro_ and _in vivo_ pooled CRISPR screens

The gRNA sequences of Unassigned_1 and Unassigned_2 sublibrary were

selected from the genome-wide gRNA library (10). The selected gRNA sequences were

cloned into the modified pU6-Universal vector by cloning a T2A, DHFR, T2A, and RFP

in frame with Cas9, where the expression of gRNA and Cas9-T2A-DHFR-T2A-RFP

cassettes was independently transcribed. The insertion of the selected gRNA sequences

into the   vector was performed by VectorBuilder. The gRNA library (200 μg) was

linearized with NotI and transfected into approximately $1-2\times10^8$ RHΔhxgprt parasites

divided between four separate cuvettes. Then, transfected parasites were grown in 4×150-

mm dishes with confluent Vero cell monolayers. Pyrimethamine (Sigma) was added 24

h post transfection. All the parasites were passaged every 3 days until Passage 3 without

filtration. After 2 days (Passage 4), the parasites were syringe lysed, filtered, and counted

for genomic DNA preparation or for mouse infection. For genomic DNA preparation, at

least 1×10^8 parasites were pelleted and stored at -80 °C. For mouse infection, the parasites

were resuspended in PBS at a concentration of 2.5×10^8 parasites/ml. Then, 1×10^7

parasites in 40 μl PBS were injected into the footpad of each anesthetized mouse. Parasite

viability was determined by plaque assay. At 7 days post infection, the spleens were

collected and crushed by a plunger and passed through a cell strainer to make single cell

suspensions. Then, the suspensions were pelleted and added to 2×150-mm dishes per

spleen with confluent Vero cell monolayers. After 2-4 days, when the parasites

completely lysed out, they were filtered and counted. At least 1×10^8 parasites were

pelleted and stored at -80 °C. Parasite genomic DNA was extracted using the DNeasy

Blood and Tissue kit (Qiagen) according to the manufacturer's instructions. Integrated

gRNA sequences were PCR-amplified and barcoded with KOD FX Neo (TOYOBO)

using Primer 1 and Primer 2 (**Table S3**). Genomic DNA (1μg) was used for the template.

The resulting libraries were sequenced on a DNBSEQ-G400RS (MGI) using Primer 3

and Primer 4 (**Table S3**).

Bioinformatic analysis of the CRISPR screen

Following demultiplexing, gRNA sequencing reads were aligned to the gRNA library. The abundance of each gRNA was calculated and normalized to the total number of aligned reads (47). For *in vitro* analysis, the $\log_2$ fold-change between the P4 sample and the library was calculated for each gRNA. The fitness score for each gene was calculated as the mean $\log_2$ fold change for the top five scoring guides. For *in vivo* analysis, the $\log_2$ fold-change between each *in vivo* sample and the P4 sample was calculated for each gRNA as described above. The median fitness score across mouse replicates was used as the *in vivo* fitness score (14). For a given gene, gRNAs were compared using Wilcoxon rank sum test between P4 vs WT mice. The p values for each test were adjusted using the Benjamini-Hochberg method. The distance of each gene from the regression line was calculated as below.

$$distance\bigl(ax + by + c = 0, (x_0, y_0)\bigr) = \frac{ax_0 + b\,y_0 + c}{\sqrt{a^2 + b^2}}$$

All analyses were conducted by R (v4.1.1) with package stats (v3.6.2) and visualized by ggplot2 (v3.4.0).

Plasmid construction for knockout *Toxoplasma*

For construction of the CRISPR/Cas9 plasmids for targeting a gene of interest (GOI), two oligonucleotide primers (GOI_gRNA1_F and GOI_gRNA1_R, GOI_gRNA2_F and GOI_gRNA2_R) containing gRNA sequence were annealed and cloned into BsaI site of the pU6-Universal vector (Addgene #52694). To generate a construct for deleting the entire coding region of GOI, flanking regions of 60 bp of 5' and 3' outside the gRNAs were used to surround the genes. Forward and reverse primers with homology to the floxed HXGPRT cassette and to the 5' and 3' coding sequence of the GOI was used (GOI_homology_F and GOI_ homology _R). The primer sequences used for the genetic disruption are shown in **Table S3**.

Generation of gene knockout *Toxoplasma*

RHΔhxgprtΔku80 were filtered and resuspended in Cytomix (10mM KPO$_4$, 120mM KCl, 0.15 mM CaCl$_2$, 5mM MgCl$_2$, 25 mM HEPES, 2 mM EDTA). Parasites were mixed with 50 µg of each gRNA1 and gRNA2 CRISPR plasmids with the PCR-

amplified targeting fragment for each GOI, and supplemented with 2 mM ATP, 5mM

GSH. Parasites were electroporated by GENE PULSER II (Bio-Rad Laboratories).

Transfected parasites were selected by 25 µg/ml mycophenolic acid (MPA) (Sigma) and

50 µg/ml Xanthine (Wako) to obtain stably resistant clones. Then, parasites were

subjected to limiting dilution in 96-well plates to isolate single clones. To confirm the

disruption of the gene, we analyzed the mRNA expression by quantitative RT-PCR.

Complementation of TgGTPase and TgRimM

To complement the ΔTgGTPase and ΔTgRimM parasites, TgGTPase, TgRimM,

and TgRimM$^{\Delta ZF}$ cDNA was generated by PCR amplification from cDNA of the RH strain.

The PCR products were inserted into the pUPRT plasmid vector, which possesses cDNA

of interest with SPOT and HA epitope tag driven by the tubulin promoter. Indicated

parasites were transfected with the pUPRT complementation vector and gRNA targeting

the UPRT locus, then selected by FUDR (Wako) (48). Isolated clones were examined for

protein expression by western blotting and IFA.

TgGTPase and TgRimM phylogenetic analysis

376   ToxoDB (23), VEuPathDB (49) and BLAST were searched for TgGTPase or

377   TgRimM homologs across the phylum Apicomplexa. Clustal Omega was used to align

378   obtained homologs (50). Phylogenetic trees were visualized by ggtree (v3.2.1) (51).

Immunofluorescence assay

HFFs were grown on coverslips, infected with parasites for 24-30 h, and fixed

in PBS containing 3.7% paraformaldehyde (PFA) for 10 min at room temperature. Cells

were permeabilized with PBS containing 0.1% Triton-X for 10 min and then blocked with

in PBS containing 3% BSA for 1 h at room temperature. Then, the coverslips were

incubated with the primary antibodies for 1 h, followed by incubation with appropriate

secondary antibodies, streptavidin and DAPI for 30 min. The coverslips were mounted

using PermaFluor (Thermo Scientific). Images were acquired by confocal laser

microscopy (Olympus FV3000 IX83). Primary antibodies used were mouse anti-HA.11

(BioLegend, 901514), rabbit anti-GAP45 (52), and Alexa Fluor 594 streptavidin

(Invitrogen, S11227).

Assessment of *in vivo* virulence in mice

Mice were infected with 10^3 tachyzoites in 200 µl (intra-peritoneum) or 40 µl

(intra-footpad) of PBS. Parasite viability was determined by plaque assay. The mouse

health condition and survival were monitored daily until 30 days (intra-peritoneum) or 40

days (intra-footpad) post infection, respectively.

Quantitative RT-PCR

Total RNA was extracted by RNeasy kit (QIAGEN), and cDNA was synthesized

by Verso reverse transcription (Thermo Fisher Scientic) according to the manufacturer's

instructions. Quantitative RT-PCR was performed with a CFX Connect real-time PCR

system (Bio-Rad Laboratories) and a Go-Taq real-time PCR system (Promega). The data

were analyzed by the ΔΔCT method and normalized to ACT1 in each sample. The primer

sequences are listed in **Table S3**.

Western blotting

Cells were lysed in lysis buffer (1% NP-40, 150 mM NaCl, 20 mM Tris-HCl, pH

7.5) containing a protease inhibitor cocktail (Nacalai Tesque). The cell lysates were

separated by SDS-PAGE and transferred to polyvinylidene difluoride membranes

(Immobilon-P; Millipore). Membranes were blocked in 5% skim milk in 0.05% Tween-

20 in PBS for 1 h at room temperature, followed by incubation with primary antibodies

for 1 h at room temperature. Blots were then incubated with appropriate secondary

antibodies for 1 h at room temperature. Primary antibodies used were mouse anti-HA

(MBL, M180-7) and rabbit anti-ROP18 (this study).

Generation of custom anti-ROP18 antibody

Custom anti-ROP18 antibody (rabbit polyclonal) was generated against a synthetic C-

terminal peptide of ROP18 (AQNFEQQEHLHTE). The epitope identification, peptide

synthesis, rabbit immunization and serum collection were conducted by Cosmo Bio. The
specificity of anti-ROP18 antibody was validated for western blotting.

Quantification and statistical analysis

Information about the number of biological replicates and the type of statistical
tests used can be found in the figure legends. All statistical analyses except for the
survival rates were performed using R (4.1.1, https://www.r-project.org/). For correlation
analysis, Pearson's correlation was used. Data with p values < 0.05 were considered
statistically significant. The statistical analysis of survival rates was performed by the log-
rank test using the GraphPad Prism9 software.

Data availability

The CRISPR screen data of two sublibraries have been deposited to the NCBI
GEO. The GEO accession numbers are **GSE253884** and **GSE253885**. Any additional
information required to reanalyze the data reported in this paper is available from the
corresponding author upon request.

ACKNOWLEGDEMENT

We thank M. Enomoto and N. Yamagishi (Osaka University) for secretarial and
technical assistance. We acknowledge the NGS core facility at the Research Institute for
Microbial Diseases of Osaka University for the sequencing. We thank Dr. D. Soldati-
Favre for anti-GAP45 antibody. This study was supported by Fusion Oriented Research
for Disruptive Science and Technology (JPMJFR206D), and Moonshot research &
development (JPMJMS2025) from Japan Science and Technology Agency, the Research
Program on Emerging and Re-emerging Infectious Diseases (JP23fk0108682) from the
Agency for Medical Research and Development (AMED), a Grant-in-Aid for
Transformative Research Area (B) (Establishment of PLAMP as a new concept to
determine self and nonself for obligatory intracellular pathogens; 20B304), for Scientific
Research (A) (19H00970), for JSPS Research Fellow (23KJ1469) from the Ministry of
Education, Culture, Sports, Science and Technology, program from Joint Usage and Joint
Research Programs of the Institute of Advanced Medical Sciences, Tokushima University,

Takeda Science Foundation, Mochida Memorial Foundation, Astellas Foundation for

Research on Metabolic Disorders, Naito Foundation, the Chemo-Sero-Therapeutic

Research Institute, Research Foundation for Microbial Diseases of Osaka University,

BIKEN Taniguchi Scholarship, and Joint Research Program of Research Center for

Global and Local Infectious Diseases of Oita University (2021B06).

AUTHOR CONTRIBUTIONS

Conceptualization, Y.T. and M.Y.; Methodology, Y.T. and M.Y.; Software and Formal

Analysis, Y.T.; Investigation, Y.T. and M.Y.; Writing – Original Draft, Y.T. and M.Y.;

Writing – Review & Editing, Y.T., M.S. and M.Y.; Funding Acquisition, M.Y.;

Supervision, M.Y.

FIGURE LEGENDS

Figure 1. *In vivo* CRISPR screening targeting proteins with hyperLOPIT-

unassigned localization.

(A) Scatterplots showing in vitro and in vivo fitness scores from Unnassigned_1 (left)

and Unassigned_2 (right) libraries, respectively. ROP18 and GRA23 are labeled in

orange as control genes.

(B) Ranking tables for top 15 *in vivo* fitness-conferring genes in each library ordered by

the distance from the regression line.

Figure 2. Validation of candidate genes that affect parasite virulence.

(A) C57BL/6 mice were inoculated into the footpad with 10^3 tachyzoites of the indicated

knockout strains from the unassigned_1 library. WT (n = 6), ΔTGGT1_211850 (n = 5),

ΔDDI1 (n = 5), ΔTgGTPase (n = 5), ΔAAMT (n = 5), ΔGLT2 (n = 5), ΔCaax (n = 5),

ΔmRNACF (n = 5), ΔTxn (n = 5), ΔNudix (n = 5), ΔVIT (n = 5).

(B) C57BL/6 mice were inoculated into the footpad with 10^3 tachyzoites of the indicated

knockout strains from the unassigned_2 library. WT (n = 6), ΔTGGT1_300220 (n = 5),

ΔTgRimM (n = 5), ΔEFG1 (n = 5), ΔCPSF (n = 5), ΔTGGT1_277870 (n = 5),

ΔTGGT1_250790 (n = 5), ΔDnaj (n = 5), ΔTGGT1_312840 (n = 5), ΔTGGT1_273905

(n = 5), ΔTBC6 (n = 5), ΔTGGT1_221640 (n = 5). N.S., not significant (log-rank test).

Figure 3. TgGTPase is a cytosolic GTPase that contributes to parasite virulence.

(A) Rank-ordered plot of genes from Unassigned_1 library.

(B) Schematic of putative domains in TgGTPase.

(C) Phylogenetic tree of selected TgGTPase homologs across the apicomplexan phylum.

(D) Schematic of TgGTPase knockout and complementation of TgGTPase-HA.

(E) Western blot analysis of TgGTPase-HA.

(F) Immunofluorescence images of ΔTgGTPase::TgGTPase-HA showing cytosolic localization of TgGTPase-HA (green). GAP45 (magenta) stains the inner membrane complex of the parasites.

(G) Survival curves of C57BL/6 mice intraperitoneally infected with 10^3 tachyzoites of WT (n = 5), ΔTgGTPase (n = 5), and ΔTgGTPase::TgGTPase-HA (n = 5).

Data are representative of three independent experiments (E and F).

Figure 4. TgRimM is an apicoplast-resident protein that is essential for parasite virulence.

(A) Rank-ordered plot of genes from Unassigned_2 library.

(B) Schematic of putative domains in TgRimM. SP is predicted by SignalP 6.0.

(C) Phylogenetic tree of selected TgRimM homologs across the apicomplexan phylum.

(D) Schematic of TgRimM knockout and complementation of wild-type and mutant TgRimM.

(E) Immunofluorescence images of ΔTgRimM::TgRimM-HA showing TgRimM-HA (green) is co-stained with streptavidin (magenta), an apicoplast marker.

(F) Western blot analysis of TgRimM-HA.

(G) Survival curves of C57BL/6 mice intraperitoneally infected with 10^3 tachyzoites of WT (n = 5), ΔTgRimM (n = 5), ΔTgRimM::TgRimM-HA (n = 5), or ΔTgRimM::TgRimM$^{\Delta ZF}$-HA (n = 5).

Data are representative of three independent experiments (E and F).

SUPPLEMENTAL MATERIAL

Supplementary Figure 1. Assessing the reproducibility of CRISPR screens.

Related to Figure 1.

(A) Rank-ordered plots for *in vitro* fitness scores of 4th passage.

(B) Correlation between our *in vitro* fitness scores and the *in vitro* fitness scores in HFF.

(C) Overlay of *in vivo* fitness scores for each mouse. Pearson's correlation coefficients

are shown as mean ± SD.

Supplementary Figure 2. Generating gene knockout parasites.

Related to Figure 2.

(A) Schematic of gene knockout strategy.

(B) Quantitative RT-PCR validations for indicated parasites.

Supplementary Figure 3. TgRimM$^{\Delta ZF}$-HA localized in the apicoplast.

Related to Figure 4.

Immunofluorescence images of ΔTgRimM::TgRimM$^{\Delta ZF}$-HA showing TgRimM$^{\Delta ZF}$-HA

(green) is co-stained with streptavidin (magenta), an apicoplast marker. Data are

representative of three independent experiments.

Table S1.

Summary of *in vivo* CRISPR screen using Unassigned_1 library, gRNA sequences, raw

count data, and *in vivo* fitness scores for each mouse.

Table S2.

Summary of *in vivo* CRISPR screen using Unassigned_2 library, gRNA sequences, raw

count data, and *in vivo* fitness scores for each mouse.

Table S3.

Lists of the primers used in this study.

REFERENCES

1.   Montoya JG, Liesenfeld O. 2004. Toxoplasmosis. Lancet 363:1965-76.

2.   Kim K, Weiss LM. 2004. Toxoplasma gondii: the model apicomplexan. Int J Parasitol

34:423-32.

3.   Lebrun M, Carruthers VB, Cesbron-Delauw M-F. 2020. Toxoplasma secretory proteins

and their roles in parasite cell cycle and infection, p 607-704, Toxoplasma gondii. Elsevier.

4.   Kloehn J, Lacour CE, Soldati-Favre D. 2021. The metabolic pathways and transporters of

the plastid organelle in Apicomplexa. Curr Opin Microbiol 63:250-258.

5.   Gambarotto D, Zwettler FU, Le Guennec M, Schmidt-Cernohorska M, Fortun D, Borgers

S, Heine J, Schloetel JG, Reuss M, Unser M, Boyden ES, Sauer M, Hamel V, Guichard P.

2019. Imaging cellular ultrastructures using expansion microscopy (U-ExM). Nat

Methods 16:71-74.

6.   Tosetti N, Dos Santos Pacheco N, Bertiaux E, Maco B, Bournonville L, Hamel V, Guichard

P, Soldati-Favre D. 2020. Essential function of the alveolin network in the subpellicular

microtubules and conoid assembly in Toxoplasma gondii. Elife 9.

7.   Barylyuk K, Koreny L, Ke H, Butterworth S, Crook OM, Lassadi I, Gupta V, Tromer E,

Mourier T, Stevens TJ, Breckels LM, Pain A, Lilley KS, Waller RF. 2020. A Comprehensive

Subcellular Atlas of the Toxoplasma Proteome via hyperLOPIT Provides Spatial Context

for Protein Functions. Cell Host Microbe 28:752-766 e9.

8.   Guerin A, Strelau KM, Barylyuk K, Wallbank BA, Berry L, Crook OM, Lilley KS, Waller

RF, Striepen B. 2023. Cryptosporidium uses multiple distinct secretory organelles to

interact with and modify its host cell. Cell Host Microbe 31:650-664 e6.

9.   Christoforou A, Mulvey CM, Breckels LM, Geladaki A, Hurrell T, Hayward PC, Naake T,

Gatto L, Viner R, Martinez Arias A, Lilley KS. 2016. A draft map of the mouse pluripotent

stem cell spatial proteome. Nat Commun 7:8992.

10.   Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, Thiru P, Saeij JPJ,

Carruthers VB, Niles JC, Lourido S. 2016. A Genome-wide CRISPR Screen in Toxoplasma

Identifies Essential Apicomplexan Genes. Cell 166:1423-1435 e12.

11.   Young J, Dominicus C, Wagener J, Butterworth S, Ye X, Kelly G, Ordan M, Saunders B,

Instrell R, Howell M, Stewart A, Treeck M. 2019. A CRISPR platform for targeted in vivo

screens identifies Toxoplasma gondii virulence factors in mice. Nat Commun 10:3963.

12.   Sangare LO, Olafsson EB, Wang Y, Yang N, Julien L, Camejo A, Pesavento P, Sidik SM,

Lourido S, Barragan A, Saeij JPJ. 2019. In Vivo CRISPR Screen Identifies TgWIP as a

Toxoplasma Modulator of Dendritic Cell Migration. Cell Host Microbe 26:478-492 e8.

13.   Butterworth S, Torelli F, Lockyer EJ, Wagener J, Song OR, Broncel M, Russell MRG,

Moreira-Souza ACA, Young JC, Treeck M. 2022. Toxoplasma gondii virulence factor

ROP1 reduces parasite susceptibility to murine and human innate immune restriction.

PLoS Pathog 18:e1011021.

14.   Tachibana Y, Hashizaki E, Sasai M, Yamamoto M. 2023. Host genetics highlights IFN-

gamma-dependent Toxoplasma genes encoding secreted and non-secreted virulence

factors in in vivo CRISPR screens. Cell Rep 42:112592.

15.   Giuliano CJ, Wei KJ, Harling FM, Waldman BS, Farringer MA, Boydston EA, Lan TCT,

Thomas RW, Herneisen AL, Sanderlin AG, Coppens I, Dvorin JD, Lourido S. 2023.
Functional profiling of the Toxoplasma genome during acute mouse infection. bioRxiv
doi:10.1101/2023.03.05.531216.

16. Aghabi D, Sloan M, Gill G, Hartmann E, Antipova O, Dou Z, Guerra AJ, Carruthers VB,
Harding CR. 2023. The vacuolar iron transporter mediates iron detoxification in
Toxoplasma gondii. Nat Commun 14:3659.

17. Zhang H, Liu J, Ying Z, Li S, Wu Y, Liu Q. 2020. Toxoplasma gondii UBL-UBA shuttle
proteins contribute to the degradation of ubiquitinylated proteins and are important for
synchronous cell division and virulence. FASEB J 34:13711-13725.

18. Aquilini E, Cova MM, Mageswaran SK, Dos Santos Pacheco N, Sparvoli D, Penarete-
Vargas DM, Najm R, Graindorge A, Suarez C, Maynadier M, Berry-Sterkers L, Urbach S,
Fahy PR, Guerin AN, Striepen B, Dubremetz JF, Chang YW, Turkewitz AP, Lebrun M.
2021. An Alveolata secretory machinery adapted to parasite host cell invasion. Nat
Microbiol 6:425-434.

19. Gas-Pascual E, Ichikawa HT, Sheikh MO, Serji MI, Deng B, Mandalasi M, Bandini G,
Samuelson J, Wells L, West CM. 2019. CRISPR/Cas9 and glycomics tools for Toxoplasma
glycobiology. J Biol Chem 294:1104-1125.

20. Engelberg K, Chen CT, Bechtel T, Sanchez Guzman V, Drozda AA, Chavan S, Weerapana
E, Gubbels MJ. 2020. The apical annuli of Toxoplasma gondii are composed of coiled-coil
and signalling proteins embedded in the inner membrane complex sutures. Cell Microbiol
22:e13112.

21. Quan JJ, Nikolov LA, Sha J, Wohlschlegel JA, Bradley PJ. 2023. Toxoplasma gondii
encodes an array of TBC-domain containing proteins including an essential regulator that
targets Rab2 in the secretory pathway. bioRxiv doi:10.1101/2023.05.28.542599.

22. Roumegous C, Abou Hammoud A, Fuster D, Dupuy JW, Blancard C, Salin B, Robinson
DR, Renesto P, Tardieux I, Frenal K. 2022. Identification of new components of the basal
pole of Toxoplasma gondii provides novel insights into its molecular organization and
functions. Front Cell Infect Microbiol 12:1010038.

23. Harb OS, Roos DS. 2020. ToxoDB: Functional Genomics Resource for Toxoplasma and
Related Organisms. Methods Mol Biol 2071:27-47.

24. Teufel F, Almagro Armenteros JJ, Johansen AR, Gislason MH, Pihl SI, Tsirigos KD,
Winther O, Brunak S, von Heijne G, Nielsen H. 2022. SignalP 6.0 predicts all five types
of signal peptides using protein language models. Nat Biotechnol 40:1023-1025.

25. Butterworth S, Kordova K, Chandrasekaran S, Thomas KK, Torelli F, Lockyer EJ, Edwards
A, Goldstone R, Koshy AA, Treeck M. 2023. High-throughput identification of
Toxoplasma gondii effector proteins that target host cell transcription. Cell Host Microbe

31:1748-1762 e8.

26.   Chen Y, Liu Q, Xue JX, Zhang MY, Geng XL, Wang Q, Jiang W. 2021. Genome-Wide
618   CRISPR/Cas9 Screen Identifies New Genes Critical for Defense Against Oxidant Stress
619   in Toxoplasma gondii. Front Microbiol 12:670705.

620   27.   Harding CR, Sidik SM, Petrova B, Gnadig NF, Okombo J, Herneisen AL, Ward KE,
621   Markus BM, Boydston EA, Fidock DA, Lourido S. 2020. Genetic screens reveal a central
622   role for heme metabolism in artemisinin susceptibility. Nat Commun 11:4813.

623   28.   Krishnamurthy S, Maru P, Wang Y, Bitew MA, Mukhopadhyay D, Yamaryo-Botte Y,
624   Paredes-Santos TC, Sangare LO, Swale C, Botte CY, Saeij JPJ. 2023. CRISPR Screens
625   Identify Toxoplasma Genes That Determine Parasite Fitness in Interferon Gamma-
626   Stimulated Human Cells. mBio 14:e0006023.

627   29.   Krishnan A, Kloehn J, Lunghi M, Chiappino-Pepe A, Waldman BS, Nicolas D, Varesio E,
628   Hehl A, Lourido S, Hatzimanikatis V, Soldati-Favre D. 2020. Functional and
629   Computational Genomics Reveal Unprecedented Flexibility in Stage-Specific Toxoplasma
630   Metabolism. Cell Host Microbe 27:290-306 e11.

631   30.   Lockyer EJ, Torelli F, Butterworth S, Song OR, Howell S, Weston A, East P, Treeck M.
632   2023. A heterotrimeric complex of Toxoplasma proteins promotes parasite survival in
633   interferon gamma-stimulated human cells. PLoS Biol 21:e3002202.

634   31.   Paredes-Santos TC, Bitew MA, Swale C, Rodriguez F, Krishnamurthy S, Wang Y, Maru P,
635   Sangare LO, Saeij JPJ. 2023. Genome-wide CRISPR screen identifies genes synthetically
636   lethal with GRA17, a nutrient channel encoding gene in Toxoplasma. PLoS Pathog
637   19:e1011543.

638   32.   Waldman BS, Schwarz D, Wadsworth MH, 2nd, Saeij JP, Shalek AK, Lourido S. 2020.
639   Identification of a Master Regulator of Differentiation in Toxoplasma. Cell 180:359-372
640   e16.

641   33.   Wang Y, Sangare LO, Paredes-Santos TC, Hassan MA, Krishnamurthy S, Furuta AM,
642   Markus BM, Lourido S, Saeij JPJ. 2020. Genome-wide screens identify Toxoplasma gondii
643   determinants of parasite fitness in IFNgamma-activated murine macrophages. Nat
644   Commun 11:5258.

645   34.   Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, Udenze K, Bronner IF,
646   Casandra D, Mayho M, Brown J, Li S, Swanson J, Rayner JC, Jiang RHY, Adams JH. 2018.
647   Uncovering the essential genes of the human malaria parasite Plasmodium falciparum by
648   saturation mutagenesis. Science 360.

649   35.   Cilingir G, Broschat SL, Lau AO. 2012. ApicoAP: the first computational model for
650   identifying apicoplast-targeted proteins in multiple species of Apicomplexa. PLoS One
651   7:e36598.

652 36. Abrahamsen MS, Templeton TJ, Enomoto S, Abrahante JE, Zhu G, Lancto CA, Deng M,
653   Liu C, Widmer G, Tzipori S, Buck GA, Xu P, Bankier AT, Dear PH, Konfortov BA, Spriggs
654   HF, Iyer L, Anantharaman V, Aravind L, Kapur V. 2004. Complete genome sequence of
655   the apicomplexan, Cryptosporidium parvum. Science 304:441-5.

656 37. Xu P, Widmer G, Wang Y, Ozaki LS, Alves JM, Serrano MG, Puiu D, Manque P, Akiyoshi
657   D, Mackey AJ, Pearson WR, Dear PH, Bankier AT, Peterson DL, Abrahamsen MS, Kapur
658   V, Tzipori S, Buck GA. 2004. The genome of Cryptosporidium hominis. Nature 431:1107-
659   12.

660 38. Zhu G, Marchewka MJ, Keithly JS. 2000. Cryptosporidium parvum appears to lack a
661   plastid genome. Microbiology (Reading) 146 (Pt 2):315-321.

662 39. Bylund GO, Wipemo LC, Lundberg LA, Wikstrom PM. 1998. RimM and RbfA are
663   essential for efficient processing of 16S rRNA in Escherichia coli. J Bacteriol 180:73-82.

664 40. Liu K, Lee KP, Duan J, Kim EY, Singh RM, Di M, Meng Z, Kim C. 2023. Cooperative role
665   of AtRsmD and AtRimM proteins in modification and maturation of 16S rRNA in plastids.
666   Plant J 114:310-324.

667 41. van Dooren GG, Striepen B. 2013. The algal past and parasite present of the apicoplast.
668   Annu Rev Microbiol 67:271-89.

669 42. Leveque MF, Berry L, Cipriano MJ, Nguyen HM, Striepen B, Besteiro S. 2015.
670   Autophagy-Related Protein ATG8 Has a Noncanonical Function for Apicoplast
671   Inheritance in Toxoplasma gondii. mBio 6:e01446-15.

672 43. Niu Z, Ye S, Liu J, Lyu M, Xue L, Li M, Lyu C, Zhao J, Shen B. 2022. Two apicoplast
673   dwelling glycolytic enzymes provide key substrates for metabolic pathways in the
674   apicoplast and are critical for Toxoplasma growth. PLoS Pathog 18:e1011009.

675 44. Sanchez SG, Bassot E, Cerutti A, Mai Nguyen H, Aida A, Blanchard N, Besteiro S. 2023.
676   The apicoplast is important for the viability and persistence of Toxoplasma gondii
677   bradyzoites. Proc Natl Acad Sci U S A 120:e2309043120.

678 45. McFadden GI, Yeh E. 2017. The apicoplast: now you see it, now you don't. Int J Parasitol
679   47:137-144.

680 46. Huynh MH, Carruthers VB. 2009. Tagging of endogenous genes in a Toxoplasma gondii
681   strain lacking Ku80. Eukaryot Cell 8:530-9.

682 47. Sidik SM, Huet D, Lourido S. 2018. CRISPR-Cas9-based genome-wide screening of
683   Toxoplasma gondii. Nat Protoc 13:307-323.

684 48. Shen B, Brown KM, Lee TD, Sibley LD. 2014. Efficient gene disruption in diverse strains
685   of Toxoplasma gondii using CRISPR/CAS9. mBio 5:e01114-14.

686 49. Alvarez-Jarreta J, Amos B, Aurrecoechea C, Bah S, Barba M, Barreto A, Basenko EY,
687   Belnap R, Blevins A, Bohme U, Brestelli J, Brown S, Callan D, Campbell LI, Christophides

GK, Crouch K, Davison HR, DeBarry JD, Demko R, Doherty R, Duan Y, Dundore W,
Dyer S, Falke D, Fischer S, Gajria B, Galdi D, Giraldo-Calderon GI, Harb OS, Harper E,
Helb D, Howington C, Hu S, Humphrey J, Iodice J, Jones A, Judkins J, Kelly SA, Kissinger
JC, Kittur N, Kwon DK, Lamoureux K, Li W, Lodha D, MacCallum RM, Maslen G,
McDowell MA, Myers J, Nural MV, Roos DS, et al. 2024. VEuPathDB: the eukaryotic
pathogen, vector and host bioinformatics resource center in 2023. Nucleic Acids Res
52:D808-D816.

50.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Soding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of
high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol
7:539.

51.   Yu G. 2020. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc
Bioinformatics 69:e96.

52.   Plattner F, Yarovinsky F, Romero S, Didry D, Carlier MF, Sher A, Soldati-Favre D. 2008.
Toxoplasma profilin is essential for host cell invasion and TLR11-dependent induction of
an interleukin-12 response. Cell Host Microbe 3:77-87.

Figure 1. *In vivo* CRISPR screening targeting proteins with hyperLOPIT-unassigned localization.

(A) Scatterplots showing *in vitro* and *in vivo* fitness scores from Unnassigned_1 (left) and Unnassigned_2 (right) libraries, respectively. ROP18 and GRA23 are labeled in

orange as control genes. (B) Ranking tables for top 15 *in vivo* fitness-conferring genes in each library ordered by the distance from the regression line.

Figure 2 _Tachibana et al.

Figure 2. Validation of candidate genes that affect parasite virulence.

(A) C57BL/6 mice were inoculated into the footpad with 10^3 tachyzoites of the indicated knockout strains from the unassigned_1 library. WT (n = 6), ΔTGGT1_211850 (n = 5), ΔDDI1 (n = 5), ΔTgGTPase (n = 5), ΔAAMT (n = 5), ΔGLT2 (n = 5), ΔCaax (n = 5), ΔmRNACF (n = 5), ΔTxn (n = 5), ΔNudix (n = 5), ΔVIT (n = 5).

(B) C57BL/6 mice were inoculated into the footpad with 10^3 tachyzoites of the indicated knockout strains from the unassigned_2 library. WT (n = 6), ΔTGGT1_300220 (n = 5), ΔTgRimM (n = 5), ΔEFG1 (n = 5), ΔCPSF (n = 5), ΔTGGT1_277870 (n = 5), ΔTGGT1_250790 (n = 5), ΔDnaj (n = 5), ΔTGGT1_312840 (n = 5), ΔTGGT1_273905 (n = 5), ΔTBC6 (n = 5), ΔTGGT1_221640 (n = 5). N.S., not significant (log-rank test).

Figure 3 Tachibana et al.

Figure 3. TgGTPase is a cytosolic GTPase that contributes to parasite virulence.

(A) Rank-ordered plot of genes from Unassigned_1 library. (B) Schematic of putative domains in TgGTPase. (C) Phylogenetic tree of selected TgGTPase homologs across the apicomplexan phylum. (D) Schematic of TgGTPase knockout and complementation of TgGTPase-HA. (E) Western blot analysis of TgGTPase-HA. (F) Immunofluorescence images of ΔTgGTPase::TgGTPase-HA showing cytosolic localization of TgGTPase-HA (green). GAP45 (magenta) stains the inner membrane complex of the parasites. (G) Survival curves of C57BL/6 mice intraperitoneally infected with 10^3 tachyzoites of WT (n = 5), ΔTgGTPase (n = 5), and ΔTgGTPase::TgGTPase-HA (n = 5). Data are representative of three independent experiments (E and F).

Figure 4. TgRimM is an apicoplast-resident protein that is essential for parasite virulence.

(A) Rank-ordered plot of genes from Unassigned_2 library. (B) Schematic of putative domains in TgRimM. SP is predicted by SignalP 6.0. (C) Phylogenetic tree of selected TgRimM homologs across the apicomplexan phylum. (D) Schematic of TgRimM knockout and complementation of wild-type and mutant TgRimM. (E) Immunofluorescence images of ΔTgRimM::TgRimM-HA showing TgRimM-HA (green) is co-stained with streptavidin (magenta), an apicoplast marker. (F) Western blot analysis of TgRimM-HA. (G) Survival curves of C57BL/6 mice intraperitoneally infected with 10^3 tachyzoites of WT (n = 5), ΔTgRimM (n = 5), ΔTgRimM::TgRimM-HA (n = 5), or ΔTgRimM::TgRimM$^{\Delta ZF}$-HA (n = 5). Data are representative of three independent experiments (E and F).