Complex interactions in the life cycle of a simple parasite shape the

evolution of virulence

Luís M. Silva [1,2] *

Jacob C. Koella [1]

[1] Institute of Biology, University of Neuchâtel, Rue Emilie-Argand 11, 2000 Neuchâtel,

Switzerland

[2] Department of Zoology, University of British Columbia, 6270 University Boulevard,
Vancouver, BC V6T 1Z4, Canada

* Corresponding author: luis.silva@ubc.ca

ORCID

LMS: 0000-0002-8203-4006

22 **Classification:** Biological Sciences, Evolution

24 **Keywords**: pathogen evolution, virulence, transmission, exploitation, microsporidia

Significance statement

Classical expectations on parasite evolution assume a trade-off between transmission rate and harm to the host (virulence), favoring parasites with intermediate virulence. However, recent studies challenge this idea, suggesting it is too simplistic. Here, by selecting the parasite *Vavraia culicis* for early or late transmission (or shorter and longer time within the host) in the host *Anopheles gambiae* we demonstrate that understanding virulence evolution requires considering the entirety and the type of transmission cycle and not merely the within-host stage. These findings highlight the importance of including all transmission stages in parasite evolution studies and demonstrate how certain social conditions might unintentionally select more virulent parasites.

Abstract

Evolutionary expectations about the virulence of parasites (i.e., the parasite-induced mortality rate of the host) often focus solely on the within-host transmission stage, overlooking the time spent between hosts and variations in transmission cycles. Moreover, parasite growth rate within the host is closely linked to virulence. We here suggest that a simplified view of transmission and parasite evolution makes it difficult to predict how virulence will evolve. We illustrate our ideas with a parasite with a simple life cycle, the microsporidian *Vavraia culicis*, which infects the mosquito *Anopheles gambiae*. We selected the parasite over six host generations for early or late host transmission, corresponding to shorter or longer time within the host. Selecting for late transmission increased their exploitation of the host, resulting in higher virulence and a shorter life cycle with rapid infective spore production, comparatively to selection for early transmission. In response, hosts infected with late-selected spores shortened their life cycle and shifted to earlier reproduction. Notably, higher virulence was negatively correlated with the endurance of infective spores outside the host, demonstrated in a parallel study using these parasite lines. These findings emphasize the importance of considering the entire transmission cycle in studies of parasite evolution and raise concerns about how host density and social settings might influence virulence evolution.

Introduction

Virulence is generally defined as the degree to which a parasite reduces the fitness of its host[1], commonly measured as the cost in host survival. It is the consequence of complex interactions between host and parasite strategies[2]. On one hand, the host can influence virulence through its ability to clear the parasite (its resistance) or to limit its effects and associated costs (its tolerance)[3,4]. On the other hand, the parasite's influence on virulence can be decomposed into processes that are linked to using its host's resources and growing (exploitation) or mechanisms that damage the host independently of the parasite's growth, such as toxins (per parasite pathogenicity)[2,5,6].

Despite the complexity of this interplay[7], most ideas about the evolution of virulence are based on

simple notions: that virulence is governed by the parasite and that its evolution is constrained by its

trade-off with the rate of transmission. Thus, to maximize a parasite's total transmission (that is, the

rate of transmission times the period of infection), evolutionary pressure should favor a higher rate of

transmission until the advantage of faster transmission is balanced by a shorter infection period due to

greater virulence. Therefore, evolution should lead to an intermediate level of virulence[8,9]. While this

and variations of this idea have been extensively discussed and studied for the last 50 years[8–10],

empirical studies have given mixed support[11,12]. This is partly due to an overly simplistic description

of transmission, the key parasite fitness measure and a vital trait for infection development and

spread. As a parasite fitness trait, transmission is rather complex, since it encompasses stages within

and among hosts, each with their own set of factors that can interact and trade-off against each other[13].

Nevertheless, several of these factors have been studied and indeed whether transmission is vertical or

horizontal[14], continuous or not[15], or whether a parasite is directly transmitted, has environmental

stages[13], or is vector-borne[16] strongly affects predictions about the evolution of virulence. However, a

strongly overlooked factor is the timing of transmission, that is the time it takes for the parasite to

grow, develop, and potentially evolve within its host before being transmitted.

According to the trade-off model earlier transmission should lead to higher virulence[17], though details

of the timing of transmission and pathogenicity can lead to more complex predictions. One study that

measured the effect of time to transmission was performed by Cooper and colleagues on nuclear

polyhedrosis virus (NPV) infections in the moth *Lymantria dispar*[18]. They selected the virus for early

or late transmission and tracked over generations the virulence and viral growth at two-time points of

the moth development. Selection for early transmission led to greater virulence but lower viral burden

than selection for late transmission. While this result appears to agree with the virulence-transmission

trade-off theory, it is difficult to interpret, for the experiment selected not only for early or late

transmission but also inadvertently for rapid growth. This situation is made more complex by the fact

that a longer time within the host before transmission might provide an opportunity for competing

parasites to evolve[19,20]. Such competition is thought to be responsible for the evolution of higher

virulence of the microsporidian *Glugoides intestinalis* when its host, *Daphnia magna*, is allowed to

live longer[21]. Additionally, part of the parasite's impact may be due to the phenotypic responses of the

host, which are governed by trade-offs between early and late reproduction.

Lastly, most studies that aspire to understand parasite evolution tend to focus on one transmission

stage, either within or among hosts, and neglect the link between the two. This raises major concerns

for epidemiology and the understanding of parasite and virulence evolution[13]. As expected from life

history theory, a high fitness in one environment is likely to trade-off with success in other

environments. This is particularly crucial for parasites, as their fitness is dependent on the success in

the first host, survival in the environment they are transmitted to (e.g., water, soil, surfaces), and

ability to infect a new host. Hence, it is important to relate and integrate the diverse environments a

parasite is exposed to, to fully understand its fitness and evolution and the focus on merely one of
them might greatly bias our predictions regarding its evolution. For instance, a parasite that has a high
fitness within one host, might not be as well equipped to survive the outside host environment, and
vice-versa. Such trade-offs have been hinted at by some authors but in poorly controlled conditions.
Taking this into account, we first aimed to test whether selecting for the parasite's time to
transmission affects its virulence. Next, in a common garden experiment, we examined if changes in
virulence due to parasite evolution influenced infection dynamics and the response of non-evolved
hosts. We then integrated our results into the modern virulence decomposition framework, which
allowed us to separate the parasite's impact on the host into growth-dependent (exploitation) and
growth-independent (per parasite pathogenicity) costs. Finally, we connected these findings to a
parallel study with the same parasite lines, which assessed differences in parasite endurance and
infection ability after exposure to various temperatures and durations outside of the host.
For this, we used the mosquito *Anopheles gambiae* and its microsporidian parasite *Vavraia culicis*,
which is a common generalist parasite of mosquitoes with low costs to the host[22,23] and whose life
cycle allows us to easily control and manipulate both time-to and mode-of transmission. With this
approach, we hoped to shed light on the role and importance of considering time to transmission, as
well as within and among hosts' evolution in evolutionary and epidemiological studies. Results from
this study are of universal interest for the understanding of parasite evolution and disease spread but
also particularly relevant in vector-borne disease evolution where most vector-transmitted parasites
require long-lived hosts or vectors to complete their life cycle.
118

Results

In this study we selected the microsporidian parasite *V. culicis* for early or late transmission (and
shorter or longer time within the host) in the host *Anopheles gambiae* for six generations (Fig. 1a).
Then, we assessed changes in parasite development within the host, as well as the host response to the
parasite, when infected by the early or late-selected parasite, or upon infection with the reference
stock parasite that did not undergo selection. Virulence of the infection by the different parasites was
also measured and decomposed into exploitation (cost dependent due to parasite growth) and per
parasite pathogenicity (cost independent of parasite growth). It is noteworthy to emphasize virulence
in this study is measured as a cost in host survival.
Concerning the parasite, we quantified its: i) cost in host survival, or virulence (Fig. 1bc); ii) cost in
host fecundity (Fig. 1d); iii) spore rate and dynamics (Fig. 3); and iv) decomposed the parasite
contribution for virulence through the quantification of cost dependent and independent of growth in
host survival (Fig. 4). Regarding the host, we measured: i) its fecundity shift in response to the
different parasites (Fig. 1d); and ii) developmental time and mortality during the larval and pupal

stage upon infection (Fig. 2). Altogether, with this set of results, we aimed to understand the drivers

of parasite and virulence evolution within the host stage and how they contribute to host infectivity.

Selection for late transmission increases virulence and shifts host fecundity

The selection regime had a cost in host longevity, with the individuals infected with spores selected

for late transmission living less than individuals infected with early-selected or unselected spores ($\chi 2$

$= 138.82$, $df = 2$, $p < 0.001$; Fig. 1b). The comparison of the maximum hazard (a proxy for virulence)

from the replicates of each regime gave a similar result, with the infection from late-selected spores

leading to an increase in maximum hazard and therefore virulence ($\chi 2 = 13.239$, $df = 1$, $p < 0.001$;

Fig. 1c). Both of these results suggest that selection for later transmission, and consequently longer

time within the host, selects for more virulent parasites which greatly reduce the lifespan of the host,

in comparison with infection by early-selected or unselected spores (Fig. 1bc). See SI Appendix Table

S1 for further details on these analyses.

When accounting for the cost of infection in fecundity, we observed this was strongly affected by the

selection regime as well ($df = 2$, F= 5.914, $p = 0.003$; Fig. 1d). However, in this case, infection by

both early- and late-selected parasites led to a lower cost for the host in comparison to infection by the

unselected reference parasite. Nevertheless, the cost of infection in host fecundity was also affected

by day ($df = 1$, F= 20.280, $p < 0.001$; Fig. 1d), with the cost increasing from day 10 to 15 of host

adulthood. Although there was no significant effect of an interaction between the selection regime and

assay day, a multiple comparisons test was still performed to measure differences between each

treatment for each day. Hence, we observed that the cost in fecundity in individuals infected with late-

selected parasites on day 10 was less severe than either that of individuals with late parasites on day

15 of adulthood ($df = 258$, t-ratio = 4.164, $p<0.001$) or unselected parasites on day 10 ($df = 258$, t-

ratio = 3.533, $p<0.001$) and 15 ($df = 258$, t-ratio = 4.916, $p=0.006$). In sum, these results suggest

infection by selected parasites had a lower cost in fecundity in comparison to infection by the

unselected one. Also, the cost in fecundity increased with infection progression, in particular for

individuals infected with late-selected parasites, where we observed an increase in fecundity for the

first day, suggesting a shift in fecundity by the host. Further statistical details on SI Appendix Table

S1.

Selection shortened host and parasite life cycles

Afterward, we focused on parasite-driven changes in host development (Fig. 2). First, we compared

larvae mortality between selected and unselected parasites, and then between early and late-selected

parasites. We did not observe any statistical difference for either of the comparisons, showing parasite

selection does not affect larvae mortality (Fig. 2a). However, at the pupal stage, we observed that a

host infected by an early-selected parasite was more likely to die in this stage, comparatively to hosts

infected by late-selected parasites ($\chi 2 = 7.890$, $df = 1$, $p = 0.005$; Fig. 2b). Nevertheless, selection for

late transmission favored parasites that shortened the host time to pupation, when compared to early transmitted ones ($\chi2 = 14.267$, $df = 1$, $p < 0.001$; Fig. 2c). Although the last result provides evidence of a parasite-driven shortening of host developmental time, such outcome had no consequence in adult host body size ($\chi2 = 0.054$, $df = 1$, $p = 0.816$; Fig. 2d). We believe the shortening of the host cycle is considerably small and therefore, does not substantially affect the host growth. Further details can be found in the SI Appendix Table S2, and replicate variation is presented in Figure S1.

We then turned our focus to changes in infection parameters, such as spore rate (proxy for when the infective stages are produced) and their respective number produced per day (severity of infection and individual infectiousness). First, the spore rate increased with time ($\chi2 = 114.375$, $df = 1$, $p < 0.001$) as it is expected. However, this increase was stronger for parasites selected for late transmission and weaker for unselected parasites (interaction between selection regime and day: $\chi2 = 24.854$, $df = 2$, $p < 0.001$) (Fig. 3a), meaning late-selected spores shortened their life cycle and started producing infective spores much sooner than the other treatments. The number of spores produced per day was also affected by the selection regime ($\chi2 = 41.857$, $df = 2$, $p < 0.001$), day ($\chi2 = 158.934$, $df = 1$, $p < 0.001$) and their interaction ($\chi2 = 10.993$, $df = 2$, $p = 0.004$). Thus, the average number of spores produced increased with the progression of infection and did so more strongly if the parasites had been selected for late transmission (Fig. 3b). Additional details are in SI Appendix Table S3.

An increase in host exploitation underpins virulence evolution

We then aimed to relate the latter changes in infection dynamics to the parasite's respective virulence. For this we decomposed virulence into two components: a) exploitation, i.e. cost in host survival due to parasite growth; b) per parasite pathogenicity, i.e., cost in host survival independent of parasite growth. This decomposition allows us to understand if parasite evolution is due to changes in the parasite growth rate, or pathogenicity, such as the acquisition of virulence factors or toxin production. In our case, exploitation (i.e., cost due to parasite growth) was affected by the selection regime ($df = 2$, $F = 55.466$, $p < 0.001$), but per parasite pathogenicity was not ($df = 1$, $F = 0.302$, $p = 0.602$) (Fig. 4ab; See SI Appendix Table S4 for statistical comparisons). These findings suggest that, although we did not detect a significant effect of selection for early transmission on virulence, it did increase the parasite's ability to exploit the host, though to a lesser degree than selected for late transmission. Considering *V. culicis* growth slows down around days 15 and 18 of adulthood, almost resembling a plateau, we used spore density as a proxy of individual infectiousness and tested for a relationship between infectiousness and virulence in this model system. It is important to mention that this sort of rationale is valid for the parasite model in question, as our measure of parasite growth is the number of infective cells that will be transmitted to the environment, and consequently infect future hosts. Therefore, for this model, spore growth is equivalent to host infectiousness, one of the transmission stages discussed in Silva *et al.* (2024). Using a linear model, we detected an effect of virulence (i.e., natural log of the maximum hazard) on host infectivity (i.e., natural log of mean spore density) ($df =$

1, F =31.534, $p < 0.001$, Fig. 4c, statistical details on SI Appendix Table S4), providing evidence for

a positive correlation between host infectivity, or spore growth, and virulence in this system.

Figure 1. Longevity, fecundity and virulence evolution. (a) Experimental design, describing direct

and indirect forms of parasite selection in this study. **(b)** Host survival after infection with evolved or

reference parasites. Early and Late curves were generated with 487 and 519 females split throughout

five replicates. The stock was composed of 96 females. **(c)** Virulence, as the natural log of the

maximum hazard for each of the replicates of each selection regime ($n = 5$). The dashed line

represents the virulence of the parasite stock. See SI Appendix Table S1 for a complete statistical

analysis. **(d)** Cost of infection in fecundity, as the percentage difference in egg numbers of infected

mosquitoes comparatively to the uninfected ones, for days 10 and 15 post adult emergence. The

dashed line corresponds to the reference fecundity or uninfected individuals. The sample size was the

following: Early with 39 and 45, Late with 56 and 50, Stock with 40 and 38 for days 10 and 15,

respectively.

Figure 2. Host developmental traits. Selected regimes (i.e., early and late) were analyzed together

with and separately from the stock. **(a)** Larvae mortality across regimes. The regimes were composed

of 2068, 2011 and 2037 individuals for Early, Late and Stock. **(b)** Pupae mortality with each regime

consisting of 1955, 1873 and 1936 in the same regime order. **(c)** Early pupation, as the proportion of

individuals that pupated until day eight post eclosion. In the same regime order, the sample size was

1934, 1867 and 1929 individuals. **(d)** Body size at adult emergence using wing length of five-day-old

females as a proxy. Each regime consisted of 50 individuals. Different letters denote means that they

are significantly different from one another. See SI Appendix Table S2 for complete statistical

analysis and Figure S1 for mean and standard error for each of the replicates.

Figure 3. Infection dynamics. (a) Spore rate, as the daily proportion of females with detectable

spores for each treatment during the first ten days of adulthood. **(b)** Spore dynamics, as the mean

number of spores per female for each of the first ten days, as well as days 15 and 18 (i.e., chronic

load). For each regime and time-point, 50 females were assayed. See SI Appendix Table S3 for

complete statistical analysis.

Figure 4. Virulence decomposition. (a) Exploitation was measured as the mean spore density at days

15 and 18 (i.e., spore load divided by wing size) for every replicate of each regime. **(b)** Per parasite

pathogenicity, as the relationship between virulence and spore density. **(c)** Infectiousness-virulence

relationship for this infection model. Given that in this infection model, the spore load is expected to

decelerate in the measured chronic phase, the number of spores is not expected to increase and

therefore provide a fair estimate of the total number of spores each of the regimes can transmit. There

is a positive relationship between virulence and host infectiousness ($df = 1$, F = 31.534, p <0.001).

Different letters denote means that they are significantly different from one another. See SI Appendix

Table S4 for further details on the statistical analysis.

Discussion

We demonstrated that time to transmission affects the evolution of virulence. In particular, we showed

that contrary to the predictions of a theoretical model[17], a longer time within the host (i.e., late

transmission) was selected for greater virulence than a shorter time (i.e., early transmission).

Furthermore, we provided contrasting evidence to a study on nuclear polyhedrosis virus (NPV)

infections in the moth *Lymantria dispar*[18], in which selection of the virus for late transmission led to

lower virulence (and higher viral burden) than selection for early transmission. Below we argue and

explain why these two contrasting findings might reflect current limitations in the study of parasite

evolution. We also demonstrate that our study serves as a suitable example to reflect on classical

theories of parasite evolution and how the integration of parasite life history is vital for the

understanding of its evolution and consequently, virulence itself.

The different outcomes observed in these two studies emphasize the complexity of the evolution of

virulence[7,12,13]. In particular, it may result from the different levels at which selection operates on

parasites: i) competition between parasites within one host; ii) the ability to survive an environment

outside the host; iii) or transmission among hosts[13]. The interlinking of these, and dependent traits,

can either limit or facilitate their evolvability, which in turn has major consequences for the

understanding of virulence evolution. Indeed, if we only consider selection at the level of transmission

among hosts, as assumed by much of the theory of the evolution of virulence[8,9] and by Osnas and

Dobson (2010), selection for early transmission would lead to a higher killing rate than selection for

late transmission.

However, if we instead consider sources of selection within the host, we can expect a different

evolutionary outcome, which in turn aligns with our results.

Since we selected for early and late transmission by letting the parasite be transmitted from

mosquitoes that died early or that lived a long time, we indirectly selected for shorter and longer

periods of infection. A longer period of infection, in turn, increases the chance of mutation (an

unlikely explanation for our results) or the strength of density-dependence as the parasite load

increases, and thus increases the possibility of competition among parasite genotypes[24,25], in a

phenomenon described as clonal interference[19]. This selects for more transmissible or more rapidly

replicating parasites in our study (Fig. 3 and 4) and others[19,24]. This result is experimentally supported

by a recent study from Sheen and colleagues (2024), which suggested that in the absence of turnover

rates in poultry markets, and longer host lifespan, virulence is no longer expected to maintain at

intermediate levels but instead continuously increase as observed with circulating parasites such as

H5N1[26]. We expect this phenomenon to happen not only in poultry markets with lower turnover but

also in society and the wild, and literature has reflected this trend[25]. Longer infection gives the

immune response more time to act, giving greater selective pressure for the parasite to evade

immunity and, consequently, replicate more rapidly and be more virulent[27], thus hindering the fixation

of genotypes with low virulence[28]. In addition, the host's immune response should increase at higher

parasite burdens[6,29] Therefore, early-transmitted parasites, which have had little time to replicate, are

subjected to less selective pressure from the host's immune response. Late-transmitted parasites,

however, grow to a higher density and are subjected to a strong immune response, which then fades

with the mosquito's age[30,31], letting the selected genotypes replicate rapidly. Overall, thus, in contrast

to selection due to transmission among hosts, within-host processes are likely to select for high

virulence of parasites that can replicate for a long time and that transmit late.

A good example of this rationale is provided by the evolutionary models about obligate killers[12,32–34],

which assume that the host is a limited resource, making the parasite's growth density-dependent and

decoupled from its transmission. This makes it possible for "selfish" genotypes to grow without

counter-selection[32,35]. In our study, although *V. culicis* is not an obligate killer (see Materials and

Methods), the experimental design made it behave like one, by restricting its transmission route, and

therefore, both assumptions are likely to hold in our experiment. In particular, our protocol eliminated

many costs of transmission. Thus, for example, since spores (infective stage of our parasite) were

sampled shortly after their production, transmission was independent of their long-term quality or

infectivity, so any trade-off between the replication rate and quality of spores played no role in
transmission[12,36,37]. Indeed, the positive correlation between chronic spore load (i.e., a proxy for
infectiousness) and virulence (when results of all selection regimes and replicates are pooled) (Fig.
4c) corroborates this uncontrollable growth hypothesis. Several studies and models have supported
this hypothesis[5,38,39].
This hypothesis introduce us to the third level at which selection can act on: the ability to endure the
environment among hosts, a transmission stage common to most parasites[13,38,40]. This resting stage is
crucial for *V. culicis* because its transmission cycle encompasses death of an individual on water and
release of infective spores that will be ingested by young larvae. These spores can sit and wait for a
few hours to a several months[41], making their survival in the external environment essential for a
successful transmission. In respect to this, a follow-up study of ours with these same evolved lines
quantified the parasites' ability to endure external conditions and their resilience to different
temperatures over time [42]. Consistent with life history theory, we found that spore growth within the
host was negatively correlated with survival in the external environment. This meant that spores
selected for late transmission, which had the highest growth rate within a host, were the most
susceptible to dying in the external environment. This effect was more pronounced with longer
exposure to the outside environment and at higher temperatures. In contrast, early-selected and
unselected reference spores remained highly viable for the whole duration of the experiment, i.e., 90
days. A reduction in parasite viability in late transmitted spores led to lower infectivity in new hosts
and less severe infections with lower parasite burden compared to their counterparts. Hence, we
demonstrated that selection within the host can inadvertently impact parasite fitness outside of the
host. This is of particular significance from a social perspective, as increased host density due to
scenarios with social crowding or aggregation behaviors might shorten the time parasites spend in the
external environment. This favors parasites that prioritize evolution within the host stage, and which
invest in host exploitation and within-host growth.
Several other results in this study are noteworthy. First, adaptation to either early or late transmission
increased the parasite's prevalence of spores (Fig. 3a), growth (Fig. 3b) and exploitation of the host
(Fig. 4a). This is most likely because the parasite switched from being maintained on two mosquito
species as hosts to a parasite infecting only on *An. gambiae*. The parasite could thus specialize on a
single host without being impeded by the cost of having to also perform on a second one[43–45].
However, selection did not affect per-parasite pathogenicity for any of the regimes (i.e., damage
independent of parasite growth such as production of toxins, Fig. 4b). This is likely due to the limited
microsporidian genome[46,47] and, to our knowledge, lack of virulence factors on which selection can
act.
Second, our selection protocol had an impact on some aspects of the mosquitoes' development. For
instance, infection by late-selected parasites led to a shortening of the host life cycle through early
pupation (Fig. 2c). This result is likely due to the properties of this parasite treatment. Indeed it has

been hypothesized infection by more virulent parasites induces strong stress responses that activate

higher levels of immunity and shorten the host developmental time[6,48]. This hypothesis is supported

by our results, as the most virulent parasite (late-selected) shortened the host developmental time (Fig.

2c), while the less virulent parasite (early-selected) did not, and instead carried a cost in pupae

mortality (Fig. 2b). Nevertheless, neither of these changes was big enough to impact body size at

adult emergence (Fig. 4d).

Third, infection decreased fecundity, in particular later in life. This impact does not appear to be

linked to the density of spores, for the average number of spores did not differ between the time

points when fecundity was measured and infection by the unselected parasites gave the lowest

fecundity despite having low spore loads (Fig. 3). Rather the pattern may reflect the host's response to

infection. Indeed, as suggested by the life-history theory[49], the shorter the lifespan of infected

mosquitoes was, the greater the early investment in fecundity, while late fecundity was low and

similar in each selection regime.

Altogether, our study highlights the importance of considering the whole parasite life cycle to

understand its evolution. The findings and arguments presented here have the intention to make the

reader reflect on classical theories and its applicability into modern society and disease spread

strategies. We validate many of the concerns previously discussed in literature concerning the current

theory of virulence evolution and how they link to the oversimplification of parasite transmission, the

main parasite fitness measure. Although usefully simplistic, the current theory of virulence disregards

a considerable portion of parasite strategies, which encompass different transmission stages that

evidently cannot be neglected, even when solely studying within host aspects of infection.

Importantly, we believe this study has major consequences for epidemiology, vector-control and

disease spread strategies, regardless of the host-parasite model system in question. The latter is due to

the nature of parasites' transmission cycles, which as discussed in detail by Silva *et al.* (2024) present

the same transmission stages regardless of the taxa, and therefore evolutionary outcomes and

interactions discussed here can be adapted and further explored in most parasite taxa. Moreover, one

of the key take-home messages is the importance of the balance between within-host and among host

evolution. This has obvious consequences for disease evolution and spread, as it has become

increasingly obvious and alarming in recent pandemics, such as the SARS-CoV2, with the rise of

disproportionately high transmission associated to higher virulence in certain social settings[50–52]. Only

through the understanding of this balance, can we predict and design better disease control strategies.

Undeniably, a continued short-sighted view of parasite evolution with within-host evolution as the

sole focus of the field will hinder our understanding of the parasite evolution and promulgate the

controversy between virulence and transmission.

Lastly, we believe is imperative to make a small nod to the importance of these findings to the

ecology of this parasite. Microsporidia, such as the one in this study, have been proposed as

biological agents for malaria control[22,53] due to their ability to suppress *Plasmodium* infections in

wild- and lab-grown mosquitoes[54,55]. In line with this, is possible that the geographical distribution of
mosquitoes (and microsporidia they carry) might influence microsporidia evolution and therefore its
control potential. Surely, infection by our microsporidia is unlikely to affect mosquito population
numbers, as our late-selected treatment with shorter lifespan has great earlier fecundity (Fig. 1d), but
the same cannot be expected regarding wild mosquito density. Although, the mechanisms
underpinning *Plasmodium* suppression are still unknown, it is fair to expect microsporidia evolution
to trade-off with its ability to interfere with *Plasmodium* development to a certain extent. Hence, this
system seems a unique model to test microbe-driven vector-control mechanisms, while allowing to
study of fundamental evolutionary questions regarding infection and microbe interactions.
386

Materials and methods

Experimental model
We used the microsporidian parasite *Vavraia culicis floridensis*, provided by James Becnel (USDA,
Gainesville, FL) and the Kisumu strain of the mosquito host *Anopheles gambiae (s.s)*. Mosquitoes
were reared under standard laboratory conditions ($26 \pm 1^\circ$C, $70 \pm 5\%$ relative humidity and 12 h
light/dark). *V. culicis* is a generalist, obligatory, intracellular parasite that infects mosquito epithelial
gut cells. Mosquitoes are exposed to the microsporidian during the larval aquatic stage, where they
ingest spores in the water and food. After replication within the larvae, they start producing spores
from within the adult gut cells, re-infecting new cells and propagating the cycle until they kill the
host. Spores are transmitted when infected individuals die in water or from mother to offspring by
adhering to the surface of the eggs. Before starting the experiment, *V. culicis* was kept in large
numbers by alternatingly infecting Aedes *aegypti* and *An. gambiae* populations, to ensure it remains a
generalist parasite. The parasite was allowed to develop within these populations until host death
when *V. culicis* spores were retrieved for later infections.
401

Experimental evolution and rearing
A starting population of 300 freshly hatched *An. gambiae* larvae were grown in groups of 50 per petri
dish (120 mm diameter x 17 mm) containing 80 mL of distilled water. Larvae were fed with daily
Tetramin Baby fish food according to their age: 0.04, 0.06, 0.08, 0.16, 0.32 and 0.6 mg/larva for 0, 1,
2, 3, 4 and 5 days or older, respectively. On the second day of development, individuals were exposed
to 10,000 spores of *V. culicis* per larva. Upon pupation, individuals were transferred to cages (21 x 21
x 21 cm) and left to emerge into adults. Freshly prepared cotton pads with 6% sucrose solution were
provided throughout their adult life. At adulthood, two selection regimes were generated (Fig. 1a): i)
Early transmission, where the spores from the first 50 females to die were retrieved, mixed and used
to infect the following generation of the same regime; ii) and late transmission, where the spores from
the last 50 females to die were used. Each regime consisted of five replicate populations and was

selected for six generations prior to assessing differences due to selection at F7. To ensure

transmission only happened through parasite killing, dead individuals were removed shortly after

death, and female mosquitoes were never blood-fed and therefore unable to reproduce. The retrieved

dead females were kept in 2mL microcentrifuge tubes with 1 mL of distilled water at 4ºC for no

longer than 40 days, to maintain the viability of the spores. At the end of each generation, we pooled

all selected individuals, added a stainless-steel bead (Ø 5 mm) to each of the tubes and homogenized

the samples using a Qiagen TissueLyser LT at a frequency of 30 Hz for one minute. The

concentration of the resulting spore solution was then estimated by counting spores in a

haemocytometer under a phase-contrast microscope (400x magnification) and the dose for the next

generation was adjusted to 10,000 spores/larva. We only selected the parasite, and therefore, we used

mosquitoes from the stock population to start every generation.

Response to selection

After six generations of selection, we measured the effect of parasite selection on a set of host traits:

larvae and pupae mortality, time to pupation, body size, fecundity, longevity and infection dynamics

for each of the *V. culicis* selection regimes and replicates. We included the "stock" *V. culicis*

population as a regime as our baseline reference. All the common garden experiments were performed

in a similar design to the selection protocol unless said otherwise. Both larvae and pupae mortality, as

well as time to pupation, were assessed every 24 ± 1 h. To assess fecundity, adults were allowed to

mate in a cage (21 x 21 x 21 cm) for the first seven days of adulthood. The males were removed on

the seventh day. Two days prior to the assay day, females were allowed to blood feed for five minutes

on Tiago G. Zeferino's arm. Two days later, 100 females from each treatment and replicate were

individualized in cups (5 x 5 x 10 cm) with Whatmann filter paper (110mm) and water and allowed to

lay eggs for 48 hours. Afterwards, all females were killed, and the number of eggs counted. Body size

at adult emergence was measured using wing length at day 5 as a proxy [14]. Longevity was tracked in

adult females, which were kept in individual cups (5 x 5 x 10 cm) with access to fresh sugar solutions

and a humid filter paper. Infection dynamics was followed by sacrificing alive females at day 1, 2, 3,

4, 5, 6, 7, 8, 9, 10, 15 and 18 post adult emergence and then quantifying their spore load, as described

in the previous section.

Statistical analysis

All the analyses were performed with R version 4.3.1 in RStudio version 3034.06.2+561. The

following packages were used for visualizing and plotting the data: "ggplot2"[56], "purrr"[57], "scales"[58],

"dplyr"[59] and "tidyr"[60], as well as created with BioRender.com and edited in Microsoft PowerPoint

16.16.27. Statistical analysis and their diagnosis were conducted using: "DHARMa"[61], "car"[62],

"survival"[63], "lme4"[64], "muhaz"[65], "emmeans"[66] and "multcomp"[67].

Longevity was compared across regimes using a cox proportional hazard with the regime as the

explanatory variable and the replicate as a random factor. Then, we generated a survival curve for

each replicate of each regime and extracted its maximum hazard, as in[6], our proxy for virulence.

Differences in virulence were assessed through a linear model with the natural log of the maximum

hazard as a dependent variable, the regime as the explanatory variable and the replicate as a random

factor.

Concerning the cost of infection in fecundity, we first tested if the proportion of non-egglaying

females was the same across all regimes and days for each replicate. For this, we ran a generalized

linear model with a binomial structure and both regime and day post-adult emergence as fixed factors,

as well as their interaction, and replicate as a random factor. There was no effect of any of the

treatments and therefore, only considering females who had laid eggs, we calculated the percentual

cost of infection by subtracting the mean uninfected fecundity from each infected individual

fecundity, for day 10 and 15 of adulthood, and dividing the resulting value by the infected individual

fecundity and multiplying it by 100. Then, we assessed differences due to regime and/or days post-

adult emergence using a linear model with replicate as a random factor.

Larvae and pupae mortality were measured as the proportion of dead individuals, for each of the

stages. A generalized linear model with a binomial error structure, and the replicate as a random

factor, was used to determine the effect of regime on each of the developmental stage mortality.

Individuals pupated from day 6 to 11 post eclosion, but approximately 77%, 79% and 82% of them

pupated on day eight or nine, for Early, Late and Stock, respectively. Therefore, we split individuals

into two categories, the ones that pupated until day eight and the ones that pupated from day nine

onwards and used a generalized linear model with binomial distribution to measure differences in

early pupation. As before, the regime was considered an explanatory variable and the replicate as a

random factor.

We defined spore rate as the proportion of females with detectable spores per regime for each of the

first ten days of adulthood. Our detection threshold is estimated to be 100 spores. Mosquitoes with

detectable spores were then used to estimate spore dynamics from day one to 18 of adulthood. The

spore rate was analysed using a generalized linear model with a binomial distribution, while for spore

dynamics we used a linear model to assess differences in the log-transformed number of spores. In

both models, we had regime and day, and their interaction, as explanatory factors and the replicate as

a random factor.

To decompose virulence, we used the maximum hazard from the longevity curves and (since the

average maximum hazard day was 20) the spore density on days 15 and 18. Spore density was

calculated as the individual spore load at days 15 and 18 adjusted by individual wing length.

Virulence decomposition analysis was done as in [6]. In summary, exploitation was measured as the

natural log-transformed mean spore density from days 15 and 18 for a given regime and replicate,

while per parasite pathogenicity was given as the slope of the relationship between natural log-

transformed maximum hazard and (i.e., virulence) and exploitation. Lastly, post-hoc multiple comparisons were performed with "emmeans", using the default Tukey adjustment, across the different models.

Author contributions

LMS conceived the overall idea. JCK and LMS designed the experiments. LMS collected and analysed the data. LMS wrote the first draft of the manuscript. Both authors contributed critically to the drafts.

Acknowledgements

We thank Tiago G. Zeferino and Gwendoline Acerbi for their advice and technical support. LMS and the project were supported by SNF grant 310030_192786.

Competing Interests

The authors declare that they have no competing interests.

References

1. Read, A. F. The evolution of virulence. *Trends Microbiol.* **2**, 73–76 (1994).

2. Råberg, L. & Stjernman, M. The evolutionary ecology of infectious disease virulence. *Ecoimmunology* **548**, 78 (2012).

3. Råberg, L., Graham, A. L. & Read, A. F. Decomposing health: tolerance and resistance to parasites in animals. *Philos. Trans. R. Soc. B Biol. Sci.* **364**, 37–49 (2009).

4. Kutzer, M. A. M. & Armitage, S. A. O. Maximising fitness in the face of parasites: a review of host tolerance. *Zoology* **119**, 281–289 (2016).

5. Fraser, C., Hollingsworth, T. D., Chapman, R., de Wolf, F. & Hanage, W. P. Variation in HIV-1 set-point viral load: epidemiological analysis and an evolutionary hypothesis. *Proc. Natl. Acad. Sci.* **104**, 17441–17446 (2007).

6. Acuña-Hidalgo, B., Silva, L. M., Franz, M., Regoes, R. R. & Armitage, S. A. O. Decomposing virulence to understand bacterial clearance in persistent infections. *Nat. Commun.* **13**, 1–14 (2022).

7. Carlsson, F. & Råberg, L. The germ theory revisited: A noncentric view on infection outcome. *Proc. Natl. Acad. Sci.* **121**, e2319605121 (2024).

8. Anderson, R. M. & May, R. M. Coevolution of hosts and parasites. *Parasitology* **85**, 411–426 (1982).

9. May, R. M. & Anderson, R. M. Epidemiology and genetics in the coevolution of parasites and

hosts. *Proc. R. Soc. London. Ser. B. Biol. Sci.* **219**, 281–313 (1983).

10. Cressler, C. E., McLeod, D. V, Rozins, C., Van Den Hoogen, J. & Day, T. The adaptive

evolution of virulence: a review of theoretical predictions and empirical tests. *Parasitology*

143, 915–930 (2016).

11. Alizon, S., Hurford, A., Mideo, N. & Van Baalen, M. Virulence evolution and the trade-off

hypothesis: history, current state of affairs and the future. *J. Evol. Biol.* **22**, 245–259 (2009).

12. Alizon, S. & Michalakis, Y. Adaptive virulence evolution: the good old fitness-based

approach. *Trends Ecol. Evol.* **30**, 248–254 (2015).

13. Silva, L. M., King, K. C. & Koella, J. C. Dissecting transmission to understand parasite

evolution. (2024).

14. Agnew, P. & Koella, J. C. Virulence, parasite mode of transmission, and host fluctuating

asymmetry. *Proc. R. Soc. London. Ser. B Biol. Sci.* **264**, 9–15 (1997).

15. Godinho, D. P. *et al.* Limited host availability disrupts the genetic correlation between

virulence and transmission. *Evol. Lett.* **7**, 58–66 (2023).

16. Day, T. Parasite transmission modes and the evolution of virulence. *Evolution (N. Y).* **55**,

2389–2400 (2001).

17. Osnas, E. E. & Dobson, A. P. Evolution of virulence when transmission occurs before disease.

Biol. Lett. **6**, 505–508 (2010).

18. Cooper, V. S. *et al.* Timing of transmission and the evolution of virulence of an insect virus.

Proc. R. Soc. London. Ser. B Biol. Sci. **269**, 1161–1165 (2002).

19. Strelkowa, N. & Lässig, M. Clonal interference in the evolution of influenza. *Genetics* **192**,

671–682 (2012).

20. Pan, G. *et al.* Invertebrate host responses to microsporidia infections. *Dev. Comp. Immunol.*

83, 104–113 (2018).

21. Ebert, D. & Mangin, K. L. The influence of host demography on the evolution of virulence of

a microsporidian gut parasite. *Evolution (N. Y).* **51**, 1828–1837 (1997).

22. Lorenz, L. M. & Koella, J. C. The microsporidian parasite Vavraia culicis as a potential late

life–acting control agent of malaria. *Evol. Appl.* **4**, 783–790 (2011).

23. Lorenz, L. M. & Koella, J. C. Maternal environment shapes the life history and susceptibility

to malaria of Anopheles gambiae mosquitoes. *Malar. J.* **10**, 1–8 (2011).

24. Xue, K. S., Moncla, L. H., Bedford, T. & Bloom, J. D. Within-host evolution of human

influenza virus. *Trends Microbiol.* **26**, 781–793 (2018).

25. Schmitz, D. A., Allen, R. C. & Kümmerli, R. Negative interactions and virulence differences

drive the dynamics in multispecies bacterial infections. *Proc. R. Soc. B* **290**, 20231119 (2023).

26. Sheen, J. K., Rasambainarivo, F., Saad-Roy, C. M., Grenfell, B. T. & Metcalf, C. J. E. Markets

as drivers of selection for highly virulent poultry pathogens. *Nat. Commun.* **15**, 605 (2024).

27. Roy, B. A. & Kirchner, J. W. Evolutionary dynamics of pathogen resistance and tolerance.

Evolution (N. Y). **54**, 51–63 (2000).

28.   Fleming-Davies, A. E. *et al.* Incomplete host immunity favors the evolution of virulence in an
emergent pathogen. *Science (80-.).* **359**, 1030–1033 (2018).

29.   Moreno-García, M., Condé, R., Bello-Bedoy, R. & Lanz-Mendoza, H. The damage threshold
hypothesis and the immune strategies of insects. *Infect. Genet. Evol.* **24**, 25–33 (2014).

30.   Shaw, A. C., Joshi, S., Greenwood, H., Panda, A. & Lord, J. M. Aging of the innate immune
system. *Curr. Opin. Immunol.* **22**, 507–513 (2010).

31.   Izhar, R. & Ben-Ami, F. Host age modulates parasite infectivity, virulence and reproduction.
J. Anim. Ecol. **84**, 1018–1028 (2015).

32.   Ebert, D. & Weisser, W. W. Optimal killing for obligate killers: the evolution of life histories
and virulence of semelparous parasites. *Proc. R. Soc. London. Ser. B Biol. Sci.* **264**, 985–991
(1997).

33.   Ben-Ami, F. The virulence–transmission relationship in an obligate killer holds under diverse
epidemiological and ecological conditions, but where is the tradeoff? *Ecol. Evol.* **7**, 11157–
11166 (2017).

34.   Bérénos, C., Schmid-Hempel, P. & Mathias Wegner, K. Evolution of host resistance and
trade-offs between virulence and transmission potential in an obligately killing parasite. *J.*
Evol. Biol. **22**, 2049–2056 (2009).

35.   Levin, B. R. & Bull, J. J. Short-sighted evolution and the virulence of pathogenic
microorganisms. *Trends Microbiol.* **2**, 76–81 (1994).

36.   Dearsly, A. L., Sinden, R. E. & Self, I. A. Sexual development in malarial parasites:
gametocyte production, fertility and infectivity to the mosquito vector. *Parasitology* **100**, 359–
368 (1990).

37.   Leggett, H. C., Brown, S. P. & Reece, S. E. War and peace: social interactions in infections.
Philos. Trans. R. Soc. B Biol. Sci. **369**, 20130365 (2014).

38.   Messenger, S. L., Molineux, I. J. & Bull, J. J. Virulence evolution in a virus obeys a trade off.
Proc. R. Soc. London. Ser. B Biol. Sci. **266**, 397–404 (1999).

39.   Paul, R. E. L. *et al.* Experimental evaluation of the relationship between lethal or non-lethal
virulence and transmission success in malaria parasite infections. *BMC Evol. Biol.* **4**, 1–12
(2004).

40.   Bhardwaj, R. & Agrawal, A. Likelihood of survival of coronavirus in a respiratory droplet
deposited on a solid surface. *Phys. Fluids* **32**, (2020).

41.   Becnel, J. J. & Weiss, L. M. *Microsporidia: pathogens of opportunity*. (John Wiley & Sons
Incorporated, 2014).

42.   Silva, L. M. & Koella, J. C. Virulence evolution: thinking outside of the host. *bioRxiv* 2005–
2024 (2024).

43.   Remold, S. Understanding specialism when the jack of all trades can be the master of all.

595   *Proc. R. Soc. B Biol. Sci.* **279**, 4861–4869 (2012).

596 44. Visher, E. *et al.* The evolution of host specialization in an insect pathogen. *Evolution (N. Y).*
597   **76**, 2375–2388 (2022).

598 45. G. Zeferino, T. & C. Koella, J. Host-specific effects of a generalist parasite of mosquitoes.
599   *bioRxiv* 2002–2024 (2024).

600 46. Wadi, L. & Reinke, A. W. Evolution of microsporidia: An extremely successful group of
601   eukaryotic intracellular parasites. *PLoS Pathog.* **16**, e1008276 (2020).

602 47. Khalaf, A., Francis, O. & Blaxter, M. L. Genome evolution in intracellular parasites:
603   Microsporidia and Apicomplexa. *J. Eukaryot. Microbiol.* e13033 (2024).

604 48. Roth, O. & Kurtz, J. The stimulation of immune defence accelerates development in the red
605   flour beetle (Tribolium castaneum). *J. Evol. Biol.* **21**, 1703–1710 (2008).

606 49. Stearns, S. C. *The evolution of life histories.* **249**, (Oxford university press Oxford, 1992).

607 50. Wong, G. *et al.* MERS, SARS, and Ebola: the role of super-spreaders in infectious disease.
608   *Cell Host Microbe* **18**, 398–401 (2015).

609 51. Lloyd-Smith, J. O., Schreiber, S. J., Kopp, P. E. & Getz, W. M. Superspreading and the effect
610   of individual variation on disease emergence. *Nature* **438**, 355–359 (2005).

611 52. Stein, R. A. Super-spreaders in infectious diseases. *Int. J. Infect. Dis.* **15**, e510–e513 (2011).

612 53. Koella, J. C., Lorenz, L. & Bargielowski, I. Microsporidians as evolution-proof agents of
613   malaria control? *Adv. Parasitol.* **68**, 315–327 (2009).

614 54. Bargielowski, I. & Koella, J. C. A possible mechanism for the suppression of Plasmodium
615   berghei development in the mosquito Anopheles gambiae by the microsporidian Vavraia
616   culicis. *PLoS One* **4**, e4676 (2009).

617 55. Herren, J. K. *et al.* A microsporidian impairs Plasmodium falciparum transmission in
618   Anopheles arabiensis mosquitoes. *Nat. Commun.* **11**, 1–10 (2020).

619 56. Wickham, H. *ggplot2: elegant graphics for data analysis.* (springer, 2016).

620 57. Henry, L. & Wickham, H. purrr: Functional Programming Tools. R package version 0.3. 4.
621   (2020).

622 58. Wickham, H., Wickham, M. H. & RColorBrewer, I. Package 'scales'. (2022).

623 59. Wickham, H., François, R., Henry, L., Müller, K. & Wickham, M. H. Package 'dplyr'. *A*
624   *Gramm. Data Manip. R Packag. version* **8**, (2019).

625 60. Wickham, H. & Wickham, M. H. Package 'tidyr'. *Easily Tidy Data with'spread'and'gather*
626   *()'Functions* (2017).

627 61. Hartig, F. & Hartig, M. F. Package 'DHARMa'. *R Packag.* (2017).

628 62. Fox, J. *et al.* Package 'car'. *Vienna R Found. Stat. Comput.* **16**, (2012).

629 63. Therneau, T. M. & Lumley, T. Package 'survival'. *Surviv. Anal. Publ. CRAN* **2**, 3 (2014).

630 64. Bates, D. *et al.* Package 'lme4'. *URL http//lme4. r-forge. r-project. org* (2009).

631 65. Gentleman, R. & Winsemius, M. D. Package 'muhaz'. (2015).

632   66.   Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. Package 'emmeans'. *R Packag.*
633   *version* **1**, (2019).
634   67.   Hothorn, T. *et al.* Package 'multcomp'. *Simultaneous inference Gen. Parametr. Model. Proj.*
635   *Stat. Comput. Vienna, Austria* (2016).
636

Supplementary Information

"Complex interactions in the life cycle of a simple parasite shape the evolution of virulence"

by Luis M. Silva and Jacob C. Koella

Table S1. Longevity, fecundity and virulence evolution.

Table S2. Host developmental traits.

Table S3. Infection dynamics.

Table S4. Virulence decomposition.

Figure S1. Host developmental traits per replicate.

665 **Table S1. Longevity, fecundity and virulence evolution**

Tested effect			666
Model 1a - Host survival	$\chi 2$	*df*	*p*
Regime	138.82	2	**< 0.001**
Model 1b - Virulence	$\chi 2$	*df*	*p*
Regime	13.239	1	**< 0.001** 669
Model 1c - Cost of infection in fecundity	*df*	**F**	*p*
Regime	2	5.914	**0.063** 670
Day	1	20.279	**<0.001** 671
Replicate	4	0.695	0.596
Regime : Day	2	1.583	0.267 672

693 **Table S2.** Host developmental traits.

Tested effect	χ2	df	p
Model 2a - Larvae mortality			
Selected *vs*. Stock:			
Selection	3.564	1	0.059
Early *vs*. Late:			
Regime	3.437	1	0.064
Model 2b- Pupae mortality			
Selected *vs*. Stock:			
Selection	2.324	1	0.127
Early *vs*. Late:			
Regime	7.890	1	**0.005**
Model 2c - Early pupation			
Selected *vs*. Stock:			
Selection	0.010	1	0.920
Early *vs*. Late:			
Regime	14.267	1	**< 0.001**
Model 2d - Body size			
Selected *vs*. Stock:			
Selection	0.027	1	0.869
Early *vs*. Late:			
Regime	0.054	1	0.816

Table S3. Infection dynamics.

Tested effect			
Model 3a - Spore rate	χ^2	*df*	*p*
Regime	4.921	2	0.085
Day	114.375	1	**< 0.001**
Regime : Day	24.854	2	**< 0.001**
Model 3b - Infection dynamics	χ^2	*df*	*p*
Regime	41.857	2	**<0.001**
Day	158.934	1	**< 0.001**
Regime : Day	10.993	2	**0.004**

749   **Table S4.** Virulence decomposition.

Tested effect			
Model 4a - Exploitation	*df*	**F**	*p*
Regime	2	55.466	**< 0.001**
Multiple comparisons for Model 4a	*df*	*t*-ratio	*p*
Early - Late	12	-6.086	**0.001**
Early - Stock	12	4.401	**0.002**
Late - Stock	12	10.487	**<0.001**
Model 4b - Per parasite pathogenicity	*df*	**F**	*p*
Regime	1	0.441	0.535
Day	1	2.866	0.141
Regime : Day	1	0.300	0.606
Model 4b - Virulence-transmission	*df*	**F**	*p*
ln maximum hazard	1	9.244	**0.009**

Figure S1. Host developmental traits per replicate. (a) Larvae mortality across regimes and

replicates. Sample size as follows: Early with 357, 441, 491, 428 and 351: Late with 422, 427, 344,

423 and 395; Stock with 404, 484, 441, 401 and 307 for replicate 1 to 5 respectively. **(b)** Pupae

mortality for each regime and replicate. Missing bars represent replicates with no pupae mortality.

Sample size was: Early with 335, 419, 467, 401 and 333: Late with 395, 395, 320, 395 and 368; Stock

with 383, 455, 420, 385 and 293 for replicate 1 to 5 respectively. **(c)** Early pupation, as the proportion

of individuals that pupated until day eight post eclosion. Sample size was the following: Early with

330, 416, 465, 398 and 325: Late with 395, 395, 320, 394 and 363; Stock with 381, 452, 419, 385 and

292 for replicate 1 to 5 respectively. **(d)** Body size at adult emergence using wing length of five-day-

old females as a proxy. Each replicate consisted of 10 individuals, illustrated by a different dot shape.