Assembly of the infant gut microbiome and resistome are linked to bacterial strains in mother's milk

Pamela Ferretti[1,*], Mattea Allert[2,*], Kelsey E. Johnson[2], Marco Rossi[1], Timothy Heisel[3], Sara Gonia[3], Dan Knights[4,5], David A. Fields[6], Frank W. Albert[2], Ellen W. Demerath[2], Cheryl A. Gale[3], and Ran Blekhman[1]

[1] Section of Genetic Medicine, Department of Medicine, University of Chicago, Chicago, IL, USA
[2] Department of Genetics, Cell Biology, and Development, University of Minnesota, Minneapolis, MN, USA
[3] Department of Pediatrics, University of Minnesota, Minneapolis, MN, USA
[4] Department of Computer Science and Engineering, University of Minnesota, Minneapolis, MN, USA
[5] BioTechnology Institute, College of Biological Sciences, University of Minnesota, Minneapolis, MN, USA
[6] Department of Pediatrics, the University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA

* these authors contributed equally
Corresponding author: Ran Blekhman (blekhman@uchicago.edu)

Abstract

The establishment of the gut microbiome in early life is critical for healthy infant development. Although human milk is recommended as sole nutrition for the infant, little is known about how variation in the milk microbiome shapes the microbial communities in the infant gut. Here, we quantified the similarity between the maternal milk and the infant gut microbiome using 507 metagenomic samples collected from 195 mother-infant pairs at one, three, and six months postpartum. Microbial taxonomic overlap between milk and the infant gut was driven by *Bifidobacterium longum*, and infant microbiomes dominated by *B. longum* showed greater temporal stability than those dominated by other species. We identified numerous instances of strain sharing between milk and the infant gut, involving both commensal (e.g. *B. longum*) and pathobiont species (e.g. *K. pneumoniae*). Shared strains also included typically oral species such as *S. salivarius* and *V. parvula*, suggesting possible transmission from the infant's oral cavity to the mother's milk. At one month, the infant gut microbiome was enriched in biosynthetic pathways, suggesting that early colonisers might be more metabolically independent than later ones. Lastly, we observed significant overlap in antimicrobial resistance

gene carriage within mother-infant pairs. Together, our results suggest that the human milk microbiome has an important role in the assembly, composition, and stability of the infant gut microbiome.

Introduction

The gut microbiome composition and maturation in early life plays an important role in the development of the immune system[1,2], nutrient absorption[3,4], and metabolism regulation[5]. Its assembly is influenced by microbes acquired from the mother[6–9], among other sources[10], starting from the first day of life[11]. While the impact of the maternal gastrointestinal, vaginal, oral and cutaneous microbial species on the infant gut microbiome assembly and development have been extensively investigated[6,11–15], the role of the maternal breast milk microbiome in modulating infant gut microbes remains poorly understood.

Human breast milk represents, ideally, the sole source of nutrition for the infant in the first semester of life[16]. Nevertheless, milk remains heavily understudied, estimated to be less than 0.2% of all human-associated public metagenomic samples, based on data from 2021[17]. Maternal milk provides the infant with nutrients, such as proteins, lipids, carbohydrates (including human milk oligosaccharides, HMOs), maternal immune cells, antibodies, and live bacteria[18–20]. The milk, and the bacteria that it carries, have therefore the potential to significantly impact infant gut microbiome composition, stability, and functionality. In addition, the presence of antimicrobial resistance genes in the milk microbiome can further influence the infant's health and its gut microbiome composition[21]. The microbiome could also underlie the benefits of exclusive breastfeeding for a variety of chronic conditions including asthma[22–24], childhood obesity[25], type 1 diabetes[26], and allergic disease[27].

Despite its clear importance for infant health, research on the human milk microbiome has lagged behind studies of other human body sites. This is partly due to the nature of the sample itself: rich in human cells and fat content, and characterized by low microbial biomass[28], milk samples are challenging to process and sequence using standard approaches[28]. This is especially true for colostrum, the milk produced in the first days postpartum, as its particularly low production volume further exacerbates the low biomass challenge typical of all milk samples[29], and its collection would reduce the amount available for the infant. The few available milk microbiome cohorts are either limited by a small sample size or by lack of paired infant samples. Studies have also been hindered by the use of 16S rRNA gene sequencing[30,31], which

cannot reliably identify microbial strains[32] and does not provide insights into the functional
potential of the bacterial communities. Previous amplicon studies have identified a "core"
microbial signature of human breast milk microbiome, mostly dominated by taxa belonging to
the *Staphylococcus* and *Streptococcus* genera, followed by genera such as *Lactobacillus*,
Bifidobacterium, *Veillonella,* and *Escherichia* among others[19,33]. Bifidobacteria are of particular
interest due to their critical role in infant health[34]. They are among the most abundant taxa in the
gut of breastfed infants, and their depletion has been associated with immune and metabolic
disorders[34–36]. Bifidobacteria, in particular *B. longum subsp. infantis*, are able to digest HMOs[37].
The ability to use HMOs as a carbon source, as well as the genetic mechanisms underlying this
ability, varies across members of the *Bifidobacterium* genus[38]. Notably, the depletion of genes
required for the degradation of HMOs in bifidobacteria has been linked to systemic inflammation
and immune dysregulation in infants, highlighting the importance of bifidobacteria-mediated
HMOs degradation in infant health[39].

In this work, we investigate the relationship between the maternal milk microbiome and the
infant gut microbiome by analyzing paired breast milk and infant stool samples collected during
the first six months postpartum from a large cohort of healthy mother-infant pairs. Using high-
throughput shotgun metagenomics, we characterize microbial species composition and
temporal stability, and identify multiple instances of strain-level sharing between mothers and
infants, as well as strain persistence in the infant gut over time. We then analyze the microbial
functional potential, with particular focus on the antimicrobial resistance gene carriage of both
breast milk and infant stool samples over time.

90 Results

91 *Bifidobacterium longum* drives the limited compositional overlap between
92 the infant gut and the maternal milk microbiome
93
We collected and sequenced 507 microbiome samples from 195 mother-infant pairs recruited in
Minneapolis, USA. All infants were born at term and were exclusively breastfed at one month of
age. Breast milk was collected at one and three months postpartum (n=173), while infant stool
samples were collected at one and six months postpartum (n=334, Fig. 1A). The majority of
infants were vaginally born (75%), exclusively breastfed at six months (76%), and antibiotics
naive (67% had not received antibiotics by six months of age) (Extended Data 1 and Suppl.

Table 1, 2). Species-level taxonomic profiling was performed with MetaPhlAn4[40] (Suppl. Table
3).

Maternal milk was characterized by a significantly lower species richness than infant stool
samples (p=2.1x10^{-16} when comparing milk at one month versus infant stool at one month, and
p=1.7x10^{-5} when comparing milk at three months versus infant stool at one month, paired t-test;
Fig. 1B). Microbial diversity in both infant stool samples and maternal milk moderately increased
over time (p=0.06, and p=0.09, respectively; paired t-test). The milk microbiome was dominated
by *Bifidobacterium longum* (Fig. 1C). Other prominent members of the milk microbiome included
Bifidobacterium breve and *Bifidobacterium bifidum,* skin-associated species such as
Staphylococcus epidermidis and *Cutibacterium acnes*, and oral-associated species such as
Streptococcus salivarius (Fig. 1C). This is consistent with previous studies showing detecting
skin-associated taxa in milk in addition to bifidobacteria[41,42]. Maternal secretor status was not
significantly associated with differences in the milk microbiome alpha and beta diversity (p=0.76
paired t-test; Extended Data 2A-B).

The infant gut microbiome at one month was dominated by *B. longum*, *B. breve, B. bifidum*,
Escherichia coli, Bacteroides fragilis, Phocaeiola vulgatus, and Phocaeiola dorei, as well as by
species typically associated with the oral cavity, such as *Veillonella parvula*, *Veillonella atypica,*
and *Veillonella dispar* (Fig. 1D). *B. longum* was the most prevalent species found in both milk
and infant stool samples (55.2% and 98.2%, respectively, Fig. 1C-D and Extended Data 3).
Other taxa found in both body sites included *B. breve, B. bifidum, B. fragilis,* and *P. vulgatus*
(Fig. 1C-D). *Bifidobacterium* was the most prevalent and abundant genus in milk and stool
samples at six months (Extended Data 4A). At the family taxonomic level, the prevalence of
Bifidobacteriaceae increased over time (Extended Data 4B). Based on the dominant (most
abundant) species, the infant stool samples could be divided in four groups, hereafter referred
to as predominance groups: the first three dominated by *B. longum*, *B. breve*, and *B. bifidum*,
respectively, and a fourth group dominated by species not belonging to the *Bifidobacterium*
genus, mostly *E. coli, B. fragilis*, and *P. vulgatus* (Fig. 1D). 29.3% of infant stool samples were
dominated by *B. longum*, 13.8% by *B. breve*, 5.4% by *B. bifidum* and 51.5% by non-
bifidobacteria species (Extended Data 5). Similarly, 35% of milk samples were dominated by *B.
longum*, while 44.5% were dominated by non-bifidobacteria species.

While beta diversity analysis showed that milk and infant stool microbiome composition was
overall distinct (Fig. 1E-F), both body sites were dominated by *B. longum* (Fig. 1C-D), indicating

that *B. longum* is the main driver of the limited compositional overlap between a typical milk
microbiome and a typical infant gut microbiome at the population level. In both milk and infant
stool samples, *B. longum* subsp. *longum* (*BL. longum*) and *B. longum* subsp. *infantis* (*BL.*
infantis) frequently co-existed in the same sample (81.4% of milk and stool samples, 52.1% of
milk samples at one month, 83.4% and 95.6% of stool samples at one and six months,
respectively; Extended Data 6A-B). *BL. infantis* was present at significantly higher relative
abundance than *BL. longum* in milk samples at one and three months postpartum (p=3.2x10^{-2}
and p=4x10^{-2}, respectively; t-test) and in infant stool samples at six months of age (p=1.5x10^{-6};
t-test - Extended Data 6C). *BL. longum* was significantly enriched in infant stool samples at one
month of age as compared to *BL. infantis* (p=1.6x10^{-3}; t-test - Extended Data 6C).

Figure 1. Study design, microbial diversity, and taxonomic composition of maternal milk and infant stool microbiomes over time. (A) Study design overview for the 507 samples collected from 195 mother-infant pairs. Infant stool samples (n=334) were collected at one (1M) and six months (6M) of life. Maternal breast milk samples (n=173) were collected one and three months (3M) after delivery. (B) Shannon diversity distribution for infant stool and maternal milk samples over time. P-values calculated using two-sided paired t-test (* for p ≤ 0.05, ** for p ≤ 0.01 and **** for p ≤ 0.0001). Samples included: 122 for milk at one month, 25 for milk three months, 175 for stool one month and 159 for

stool at six months of age. Taxonomic composition of (C) the most prevalent and abundant species found in the human breast milk and (D) in the infant gut microbiome samples in relation to sample collection time point, predominance group and other relevant infant metadata. The predominance group identifies the most abundant species in each sample. Ordination plot based on Bray-Curtis distance between samples for PCoA1 and PCoA2 (E) and PCoA1 and PCoA3 (F), colored by body site of origin and sampling time. Percentage of variance explained (PVE) is reported in parentheses.

B. longum in the infant gut becomes less prevalent over time and is associated with a more stable microbiome

To investigate the stability of the microbiome in our samples, we first focused on the predominance groups described in the previous section. Among bifidobacteria, *B. longum* was the species with the highest prevalence in both breast milk (55.2%) and infant stool samples (98.2%, Fig 2A). Specifically, 50.3% of milk samples were dominated by *B. longum* (17.3% by *BL. infantis* and 28.3% by *BL. longum*), 3.5% by *B. breve,* and 1.1% by *B. bifidum* (Fig. 2A and Extended Data 7A). Across the *Bifidobacterium* genus, *B. longum* was also the most abundant, present with a mean relative abundance of 12.4% in milk samples and 21.9% in infant stool samples, Fig. 2A-B). Mothers whose breast milk was dominated by *B. longum* (in particular *BL. infantis*) exhibited the most stable milk microbiome over time. Specifically, 64.3% of milk samples in which *B. longum* was the predominant species at one month postpartum maintained this dominance at three months (Fig. 2B and Extended Data 7B). In contrast, none of the milk samples initially dominated by *B. breve* retained that predominance at the three-month time point (Fig. 2B). In infant stool samples, the percentage of samples dominated by *B. longum* increased over time, from 15.7% at one month to 46.6% at six months of age (Fig. 2B). Conversely, the percentage of samples dominated by non-bifidobacteria decreased from 68.5% to 28.1% (Fig. 2B).

In the infant, *B. longum* was found in 97.7% of the stool samples collected at 1 month of age (85.6% for *BL. infantis* and 88.5% for *BL. longum,* Fig. 2C and Extended Data 7C), while *B. breve* and *B. bifidum* were found in 83.9% and 58% of the samples in this timepoint, respectively (Fig. 2C). Overall, the prevalence of bifidobacteria in the infant gut increased from one to six months of age, particularly for those infants that were still exclusively breastfed at six months (Bonferroni adjusted $p=0.08$ for *B. longum*, $p=2.4 \times 10^{-4}$ for *B. breve*, and $p=1.8 \times 10^{-3}$ for *B. bifidum,* Fisher's exact test; Fig. 2C). A similar trend was observed in *B. longum* subspecies (Bonferroni adjusted $p=3.9 \times 10^{-3}$ for *BL. longum*, $p=1.4 \times 10^{-2}$ for *BL. infantis*, Fisher's exact test; Extended Data 7C). Consistent with the prevalence patterns, we observed significant temporal

increases in the relative abundance of several *Bifidobacterium* species in the infant gut. *B.*
longum mean relative abundance significantly increased from one to six months of age
(Bonferroni adjusted p=6.2x10^{-12}, Wilcoxon Rank Sum; Fig. 2D). Subspecies analysis showed
that *BL. infantis* increased from 3.2% mean relative abundance at one month to 23.8% at six
months (Bonferroni adjusted p=3.6x10^{-17}, Wilcoxon Rank Sum) and that *BL. longum* increased
from 8.7% mean relative abundance at one month to 9.1% at six months (Bonferroni adjusted
p=1.1x10^{-5}, Wilcoxon Rank Sum) (Extended Data 7D). Overall, *BL. longum* was more abundant
than *BL. infantis* at one month but not at six months of age (Bonferroni adjusted p=4.1x10^{-11},
Wilcoxon Rank Sum, Extended Data 7D). The trend of increasing mean relative abundance
from one to six months of age was consistently observed across other *Bifidobacterium* species,
regardless of breastfeeding practices at six months (Bonferroni-adjusted p=2.4x10$^{\square\square}$ for *B.*
breve and p=1.6x10$^{\square\square}$ for *B. bifidum*, Wilcoxon rank-sum test; Fig. 2D), albeit with substantial
inter-individual variability (Extended Data 8).

Next, we looked at the longitudinal stability of the infant gut microbiome composition at the
species-level. We found that *B. longum* (in particular *B. infantis*) was more abundant at six
months compared to one month (BH-adjusted p=2.2x10^{-10}, t-test), while *E. coli* and *V. dispar*
decreased over time (BH-adjusted p=5.3x10^{-3} and p=6.7x10^{-3}, respectively, t-test; Fig. 2E and
Extended Data 7E). We then investigated if there were species that were differentially abundant
between the two timepoints when stratifying the infants for exclusive breastfeeding. Infants that
were exclusively breastfed through six months of age showed a significant increase in *B.*
longum and *B. breve* relative abundances (BH-adjusted p=3.6x10^{-9}, p=2.2x10^{-5}, respectively,
paired t-test; Fig. 2F-G and Extended Data 9A), and a significant reduction in *C. perfringens*
(BH-adjusted p=1.3x10^{-2}, paired t-test; Fig. 2G), compared to those who were not exclusively
breastfed at six months of age. Infants that were exclusively breastfed at one month but not at
six months showed an increase in *R. gnavus* (BH-adjusted p=9.7x10^{-4}, t-test; Fig. 2G) as well as
B. longum (BH-adjusted p=1.6x10^{-4}, t-test; Fig. 2F-G) and both its subspecies (BH-adjusted
p=2.5x10^{-5} for *BL. infantis* and BH-adjusted p=2.5x10^{-2} for *BL. longum*, t-test; Extended Data
9A). We then compared the relative abundance of taxa in the gut of infants introduced to solid
foods and those exclusively breastfed at six months of age. We found no taxa with statistically
significant differential abundance between infants who were exclusively breastfed at six months
and those who were introduced to solid foods (FDR>0.05). Overall, the infant stool samples that
were dominated by *B. longum* at both one and six months of age showed the most stable
community composition over time (Extended Data 9B).

219
Figure 2. Dynamics of *Bifidobacterium* predominance, abundance, and persistence in the infant gut and maternal milk
microbiomes, stratified by breastfeeding status. (A) Prevalence of each predominance group in milk and infant stool
samples and (B) the transition of samples between predominance groups over time. Each sample is assigned one of
four predominance groups indicated by the different colors. Numbers of samples included: n=145 for milk at one
month, n=25 for milk at three months, n=175 for stool samples at one month and n=159 for stool samples at six
months of age. BLo = *B. longum*, BBr = *B. breve*, BBi = *B. bifidum.* (C) Prevalence of *Bifidobacterium* species at one
and six months of age. Prevalence at six months is further stratified by exclusive breastfeeding status (n=121 for
exclusive breastfeeding; n=34 for mixed/formula diet). Samples lacking breastfeeding information at six months were
excluded from the stratified subgroups (n=4). Bonferroni-adjusted p-values were calculated using two-sided Fisher's
exact tests. (D) Relative abundance of *Bifidobacterium* species in infants at one and six months of age. Relative
abundances at six months of age are further stratified by exclusive breastfeeding status. Relative abundances were
calculated only on samples in which the species was present. Sample size is reported in brackets. P-values
calculated using two-sided Wilcoxon rank sum, * for p ≤ 0.05, ** p ≤ 0.01, *** for p ≤ 0.001 and **** for p ≤ 0.0001.
Reported p-values are adjusted using Bonferroni method. (E) Species persistence in the infant gut across all samples
and (F) stratified by breastfeeding at six months. See Extended Data 7 for plots with separation between *BL. infantis*
and *BL. longum.* (G) Relative abundance of some of the differentially abundant species between one and six months
when divided by breastfeeding (BF) at six months. BF= exclusively breastfed. Sample size is reported in brackets. P-
values calculated with two-sided paired t-test and adjusted with BH correction. **** for P ≤ 0.0001, *** for P ≤ 0.001,
** for P ≤ 0.01 and * for P ≤ 0.05.

Strains shared between breast milk and the infant gut include bifidobacteria, pathobionts, and taxa typically found in the gut and oral microbiome

While maternal microbial transmission is well documented for other body sites, it remains unclear whether specific taxa are shared between breast milk and the infant gut, which microbes are involved, and how frequently such sharing occurs. To investigate these questions, we quantified species- and strain-level overlap between breast milk and infant stool samples to investigate potential vertical transmission events (see Methods). We found that species sharing between a mother's milk and her infant's stool samples decreased over time. On average, 10.1% of taxa found in infant stool samples at one month of age were also found in their respective mother's milk collected at one month postpartum (Fig. 3A). This percentage significantly decreased when considering infant stool samples at 6 months of age (7.2%, p=1.5x10^{-2} t-test; Fig. 3A). The most commonly shared species between milk and infant stool samples at one month postpartum was *B. longum* (shared on average in 44.3% of mother-infant pairs - 46.9% for *BL. infantis* and 41.7% for *BL. longum*), followed by *S. epidermidis* (38.7%), *B. breve* (26%), and *S. salivarius* (18%, Fig. 3B). When considering stool samples at one and six months combined, *BL. infantis* was shared in 54.4% of the mother-infant pairs, while *BL. longum* was shared in 47.2% of pairs (Fig. 3C). *B. breve* and *B. bifidum* were characterized by a significantly lower rate of sharing (30% and 11%, respectively, Figure 3C). While a positive correlation between the relative abundance of bifidobacteria in milk and the corresponding infant stool samples was observed in certain mother-infant pairs, no significant association was detected at the population level (Extended Data 10).

While informative of the potential microbial sharing between mothers and infants, species-level taxonomic profiling is not sufficient to identify transmission events, due to the genetic variability between conspecific strains[32]. We therefore performed strain-level profiling with StrainPhlAn4[40], which allowed us to reliably identify strains that are found in pairs of samples, such as a mother and her infant, defined as strain sharing. We reconstructed a total of 77 strains, of which 65 were found only in infant fecal samples and 12 were found in milk samples, due to the lower coverage. We identified twelve instances of strain-sharing between a mother's milk and her infant's stools, spanning across six mother-infant pairs and ten different taxa: *B. longum* (SGB17248), *B. bifidum* (SGB17256), *P. vulgatus* (SGB1814), *E. coli* (SGB10068), *K. pneumoniae* (SGB10115)*, K. variicola* (SGB10114), *V. parvula* (SGB6939), *R. mucilaginosa* (SGB16985), *S. salivarius* (SGB8005), and a strain belonging to the *Streptococcus* genus

(SGB8007_group, Fig. 3D and Suppl. Table 4). In particular, species like *R. mucilaginosa* and

S. salivarius are considered typical members of the infant oral microbiome, while *P. vulgatus*

and *K. pneumonia* and bifidobacteria are more commonly found in the infant gut[11]. All strains

identified in milk samples were from samples collected at one month postpartum. Strains shared

between milk and infant stool samples at one month of age were also found in stool samples

collected at six months of age in 25% of cases (Fig. 3E). Overall, strain sharing between milk

and infant stool samples was higher at one month of age compared to samples collected at six

months of age (Fig. 3E), while no strain-sharing event was detected in milk samples collected at

three months postpartum, likely due to the low microbial yield in these samples.

In addition to mother-infant sharing, strain analysis can also provide information on strain

persistence in the infant gut over the first six months of age. We found that 19.4% of the strains

identified in infant stool samples at one month persisted in the infant stools until six months of

age (Suppl. Table 4). The taxa most commonly found to be persistent in the infant gut included

E. coli (23.2% of persistence events), *B. longum* (13.4%), *B. fragilis* (8.5%), *B. vulgatus* (8.5%),

and *B. bifidum* (8.5%, Suppl. Table 4 and Fig. 3F). Vaginally delivered infants had a higher

number of persistent strains than infants that were born via C-section ($p=1.8 \times 10^{-2}$, t-test; Fig.

3G). Exclusive breastfeeding at six months of age, antibiotic intake, and maternal secretor

status were not significantly associated with differences in strain persistence in the infant gut

($p=0.08$, $p=0.25$, and $p=0.84$, respectively).

291
Figure 3. Species and strain-level sharing between maternal milk and infant stool microbiomes, and temporal
persistence of microbial strains in the infant gut. (A) Percentage of the infant stool microbiome that is shared with the
respective mother, across all mother-infant pairs. Species sharing between milk samples at one month postpartum
and infant stool samples at one month and six months of age are shown on the left and right boxplot, respectively.
Two-sided t-test, * for p ≤ 0.05. Only mother-infant pairs for which milk at one month and stool samples at one and six
months were available were included (n=108 pairs). (B) Overview of the ten most frequently shared species between
the infant stool samples at one month of age and the respective mother's milk samples at one month postpartum. *BL.*
infantis = B. longum subsp. *infantis*; *BL. longum = B. longum* subsp. *longum*. (C) Mean percentage of *Bifidobacterium*

species that were shared between milk and infant stool samples, or found in only one of the body sites, across all
mother-infant pairs. Sharing was assessed between milk at one month and infant stool at one and six months, and
milk at three months and infant stool at six months. Only pairs with at least one milk and one stool sample were
included. (D) Examples of strain sharing between milk and infant stool samples for *B. longum* (SGB17248),
Phocaeicola vulgatus (SGB1814), *Streptococcus salivarius* (SGB8005) and *Klebsiella pneumoniae* (SGB10115).
Shared strains are shown in black; persistent strains (found in the same infant at one and six months) are shown in
gray. (E) Percentage of strains shared between mother-infant pairs stratified by infant stool collection time point. (F)
Taxonomic composition of strains found to be persistent in the infant gut over time (up to six months). (G) Percentage
of strains in infant stool at one month that persisted at six months divided by delivery mode. For each infant,
persistence was calculated as the number of strains shared between the one- and six-month timepoints, divided by
the total number of strains detected at one month. Each dot is a mother-infant pair. The number of infants in each
group is reported in brackets. P-value calculated with two-sided t-test, * for p ≤ 0.05.

The maternal milk and early infant gut microbiomes are enriched in metabolic pathways for the biosynthesis of essential amino acids

Next, we investigated the functional potential of the maternal milk and infant gut microbiomes

using HUMAnN3[43], a method that profiles microbial metabolic pathways from metagenomic

data. The most prevalent pathways identified in the infant stool samples were associated with

de-novo biosynthesis of molecules, in particular of essential amino acids (such as valine,

isoleucine, threonine, and lysine) and ribonucleotides (Fig. 4A). Other abundant pathways were

associated with cell structure (i.e. peptidoglycan maturation), and energy metabolism. The

pathways involved in peptidoglycan maturation were present at higher relative abundances in

infant stool samples at one month compared to six months ($p=2.7 \times 10^{-15}$, t-test; Fig. 4A), and this

increase was particularly prominent for stool samples at one month dominated by non-

bifidobacteria species (Extended Data 11A). In milk, the most prevalent and abundant pathways

were associated with the biosynthesis of nucleotides and amino acids (such as valine and

isoleucine), cell metabolism (sucrose biosynthesis, glycogen degradation, pyruvate fermentation

and folate transformations), and cell structure (peptidoglycan maturation) (Fig. 4B). In infant

stool samples, the pathways associated with essential amino acid biosynthesis were

significantly more abundant at one month compared to six months, and this trend was

significantly more pronounced for infants that were no longer exclusively breastfed at six months

(Bonferroni-adjusted $p=5.2 \times 10^{-4}$ for infants exclusively breastfed at one and six months, and

$p=4.6 \times 10^{-12}$, for infants exclusively breastfed at one month only, t-test; Fig. 4C). In particular,

milk samples at one month dominated by *B. breve*, were associated with the highest abundance

of metabolic pathways associated with the biosynthesis of essential amino acids (Extended

Data 11B). In milk, pathways associated with the biosynthesis of essential amino acids were

significantly more abundant at three months compared to one month postpartum (Bonferroni-adjusted p=2.4x10^{-2}, t-test; Fig. 4C)

We then sought to investigate the relationship between the abundance of the metabolic pathways identified from microbes within each mother's milk with those identified in her infant's stool samples. We found a significant correlation between the metabolic pathways found in breast milk with those found in infant stool samples only for two of the mother-infant pairs for which we previously identified strain sharing events (BH-adjusted p=9x10^{-20} and p=7.6x10^{-16}, for mother-infant pair 1 and 173, respectively, Spearman's correlation; Fig. 4D and Extended data 12). When investigating specific pathways, rather than specific mother-infant pairs, we find that no pathway showed a significant correlation in its abundance in milk compared to infant stool samples (Extended Data 13).

Figure 4. Functional profiling of microbial metabolic pathways in breast milk and infant stool samples over time and across mother-infant pairs. Most abundant pathways identified in (A) breastmilk at one and three months postpartum and (B) infant gut at one and six months of life. (C) Relative abundance of pathways involved in the biosynthesis of essential amino acids across sample type, collection time point and breastfeeding (BF). Error bars represent a 95% confidence interval calculated by bootstrapping (1000 times). P-values calculated using two-sided t-test, * for p ≤ 0.05, ** p ≤ 0.01, *** for p ≤ 0.001 and **** for p ≤ 0.0001. Reported p-values are adjusted using Bonferroni method. Bars represent the mean and sample size is reported in brackets. (D) P-values of Spearman correlation between the abundance of all metabolic pathways shared between the maternal breast milk and infant stool samples for each

 mother-infant pair. P-values are calculated using a two-sided Spearman correlation test and corrected for multiple
360 testing using Benjamini Hochberg correction. Adjusted p-values are shown in $\log_{10}$ scale. Only the top 25 mother-
361 infant pairs are shown. Circle size denotes the correlation coefficient (rho), and the orange line denotes the
362 significance threshold (p-value=0.01).

The infant gut and the maternal milk microbiomes harbor a diverse landscape of antimicrobial resistance genes

The gut microbiome is a reservoir of antimicrobial resistance, yet its composition and

longitudinal variability remain poorly characterized in infants[44,45]. Even less is known about the

antimicrobial resistance genes (ARGs), collectively known as the resistome[46], present in human

breast milk, and their transmission to the infant during lactation[21,47–49]. To investigate the

resistome in the maternal milk and the infant gut, we used DeepARG[50], which leverages deep

learning to predict antimicrobial resistance genes from metagenomic data. We then compared

ARG classes across sample types and collection timepoints. Maternal milk and infant stool

differed in terms of ARGs classes composition and prevalence (Fig. 5A, Extended Data 14A). In

the milk, the most prevalent (27% on average) antimicrobial resistance class identified was

against macrolide-lincosamide-streptogramin (MLS). Overall, maternal milk was characterized

by a lower diversity of detected ARG classes compared to infant stool samples (p=2.2x10^{-16}, t-

test; Extended Data 14B-C). In both milk and infant stool samples, the diversity of ARGs classes

increased over time, although this increase was statistically significant only in the milk

(p=3.1x10^{-4} and p=0.83 for milk and stool samples respectively, paired t-test; Extended Data

14C).

The infant gut resistome was mostly dominated by resistance to tetracycline, MLS,

aminoglycoside, and beta-lactams (Fig. 5A). Infants whose stool samples were dominated by

bifidobacteria were characterized by a significantly lower carriage rate of ARGs (p=7.6x10^{-10} and

2.5x10^{-2} at one and six months respectively, t-test; Fig. 5B-C). We found no significant

difference in the resistome of infants that were born via C-section compared to those born via

vaginal delivery (p=0.34 and p=0.97 at 1 month and 6 months, respectively, t-test), between

those exposed to antibiotics and those that were antibiotics-naive (p=0.80 and p=0.14 at one

month and six months, respectively, t-test), those that were introduced to solid foods at six

months of age (0.71, t-test), nor between those exclusively breastfed at six months of age and

those fed with a mixture of breast milk and formula (p=0.35, t-test; Extended Data 14D-F).

Nevertheless, we found extensive ARGs carriage in infants with no recorded pre-, during-, and

postpartum exposure to antibiotics (Fig. 5B).

When comparing the overall resistome across all mothers and infants, we found no significant

correlation between the ARGs found in milk and the ARGs found in the infant stool samples

(R=-0.044, *p*=0.37, Spearman's test; Fig. 5D). However, the infant gut resistome at one month

was positively correlated with the gut resistome at six months of age (R=0.42, p<0.001,

Spearman's test; Fig. 5E).

Considering sharing of ARGs between milk and stool samples within each mother-infant pair,

we found that mother-infant pairs shared a significantly higher number of ARGs than what was

expected by chance (p<0.016, by permutation analysis, see Methods; Fig. 5F-G). Mother-infant

pairs 1 and 173, for which strain sharing events were identified, were the pairs with the highest

rate of shared ARGs between the mother's milk and her infant stool samples (Fig. 5F). On

average, the most commonly shared antimicrobial resistance classes were associated with

resistance to peptides, fluoroquinolone, and MLS, while the most commonly shared

antimicrobial resistance genes were *MACB* (MLS class), *ACRD* (aminoglycoside class), and

TETQ (tetracycline class) (Fig. 5H).

Figure 5. Profiling and sharing of predicted antimicrobial resistance genes between maternal milk and the infant gut microbiome. (A) Prevalence of the antimicrobial resistance genes (ARGs) classes predicted in maternal milk and the infant gut microbiome over time, calculated as percentage over the total number of detected ARGs for each sample type. (B) ARGs carriage in infant stool samples, divided by collection time point, infant metadata, predominance group and ARGs class. (C) Number of detected ARGs in infant stool samples dominated by bifidobacteria and non-bifidobacteria species at one and six months of life. P-values calculated using two-sided t-test, ** for p ≤ 0.0001, * for p<0.05. (D) Correlation between ARG carriage in maternal milk and infant stool samples, and (E) between infant stool samples at one month and six months. Each dot is a combination of a mother-infant pair and a predicted ARG class. Two-sided Spearman's R-values (correlation coefficient) and p-values are reported. (F) Percentage of ARG genes shared between at least one maternal and one infant sample, for each mother-infant pair. Mean value is indicated by the orange line. The mother-infant pairs for which strain sharing events were identified are highlighted with the arrow. (G) Distribution of the percentage of shared ARGs between at least one maternal milk and one infant stool sample on permuted mother-infant pairs (pseudo-pairs). The mean value obtained from real mother-infant pairs is indicated by the orange line. (H) Percentage of mother-infant pairs sharing antimicrobial resistance genes, divided by ARG class. For each ARG class, percentage value was calculated on the total number of mother-infant pairs in which that ARG

423 class was identified in at least one sample. The most frequently shared antimicrobial resistance genes per class are
424 highlighted in orange. MLS=Macrolides, lincosamides, streptogramines.

Discussion

Although breast milk represents a critical source of nutrition for the developing infant, little is

known about the milk microbiome, how it changes over time, its metabolic potential, and how it

shapes the infant's gut microbiome. Answering these questions requires the use of high-

resolution sequencing techniques, such as metagenomics. Here, we investigated the

composition, strain sharing, functional potential, and antimicrobial resistance of the maternal

breast milk and the infant gut microbiome in early life in predominantly exclusively breastfeeding

mother-infant dyads. In line with previous studies, the milk microbiome yielded a lower number

of reads and was characterized by a reduced microbial diversity compared to the infant gut

microbiome[11,21,28]. The milk at one month postpartum, considered largely mature[18], was

dominated by bifidobacteria, in particular *B. longum*, *B. breve,* and *B. bifidum*.

The bifidobacteria signature identified in our milk samples contrasts with some previous

amplicon-based studies, which have typically reported the milk microbiome as being dominated

by *Staphylococcus* and/or *Streptococcus*[29,51–54]. While bifidobacteria are widely recognized as

infant gut commensals[55], their presence in human milk has also been documented in prior

metagenomic studies[54,56,57] and confirmed through cultivation-based[57] and quantitative real-time

PCR approaches[58,59]. This discrepancy between *Bifidobacterium*- and

Staphylococcus/Streptococcus-dominated profiles may stem from amplification biases inherent

to amplicon sequencing, as reported in other sample types[60,61], leading to the under-

representation of bifidobacteria[62] and over-representation of *Staphylococcus*[63]. Notably, the use

of universal PCR primers not specifically optimized for bifidobacteria can lead to their under-

detection[62]. Furthermore, direct comparison of sequencing methods showed higher relative

abundances of bifidobacteria in shotgun metagenomic data compared to amplicon-based

data[54]. Additional contributing factors may include variation in DNA extraction protocols[64],

particularly the inclusion and extended duration (10 min) of the bead-beating lysis step, which

has been shown to affect the detection and relative abundance of bifidobacteria[65]. In our study,

DNA extractions for milk and infant stool samples were conducted independently by

experienced personnel, therefore minimizing the likelihood of cross-contamination during this

step. Beyond methodological considerations, the biological presence of bifidobacteria in breast

milk is also supported by the ability of *B. longum* and other bifidobacteria to survive and grow at

low-oxygen conditions[66–69]. In addition, oxygen-utilizing taxa commonly found in breast milk,

such as *Streptococcus* and *Staphylococcus,* may reduce oxygen concentrations, facilitating the

growth of bifidobacteria[33]. Finally, the presence in the milk of typical oral species, such as *S.*

salivarius, is likely due to the possible transfer of microbes from the oral cavity of the infant to

the breast milk during suckling, a process called retrograde flow[70].

While the breast milk and infant stool microbiomes were largely distinct in their microbial

composition, they were both dominated by bifidobacteria. In particular, *B. longum* was the most

prevalent and abundant species in both milk and infant stool samples, driving the limited overlap

between these two sample types. While *B. longum*, *B. bifidum,* and *B. breve* were often

coexisting and present at comparable abundances in milk, these species were largely mutually

exclusive in the infant gut, suggesting a higher level of competition between bifidobacteria

species in the infant gut compared to the milk. In line with previous findings[71], the gut

microbiome of exclusively breastfed infants was characterized by higher prevalence and

abundance of bifidobacteria compared to infants who ceased exclusive breastfeeding prior to

six months of age, suggesting that prolonged breastfeeding supports the persistence and

expansion of bifidobacteria in the infant gut microbiome. At the subspecies level, we found that

both *BL. infantis* and *BL. longum* frequently co-existed in the same sample (in 81.4% of

combined milk and infant stool samples), and that both were present at high prevalence (>75%)

in infants at one month of age. Surprisingly, *BL. infantis* was already detectable at this early time

point, in contrast to a recent study that found *BL. infantis* in infant stool samples only after 10

weeks of age[37]. Although that study's meta-analysis supported the delayed appearance of *BL.*

infantis across multiple populations, samples collected from the United States cohort were

limited to the first two weeks of life, preventing a direct comparison with our samples collected at

one month of age. However, in line with the same study[37], we found that *BL. infantis* was

significantly more abundant in infant stool samples at six months compared to one month and

compared to *BL. longum* at six months of age.

At both the species- and strain-level, microbiome sharing between the mother's milk and

the infant stool samples was more pronounced at one month postpartum than at six months,

consistent with findings from previous studies on microbiome sharing between other maternal

body sites and the infant gut[11,72]. This trend suggests that microbiome sharing between milk and

the infant gut predominantly occurs during early infancy and may also reflect the effect of time,

as a longer interval between sample collections could reduce the likelihood of detecting shared

taxa. Overall, although there was a significant correlation between *Bifidobacterium* abundance

in breast milk and infant stool in some mother-infant pairs, no statistically significant correlations

emerged at the population level (e.g. across all mother-infant pairs). This discrepancy between

individual dyads and population-level trends suggest that, while breast milk may serve as a

reservoir for taxa like *BL. infantis*, successful colonization in the infant gut is likely influenced by

additional factors (e.g., pre-existing microbial community composition, host genetics, or nutrient

availability). Consequently, the relative abundance of bifidobacteria in breast milk does not

necessarily predict their relative abundance in the infant gut, suggesting more complex

mechanisms at play beyond the initial mother-to-infant microbial transmission. At the strain

level, we found twelve cases of strain sharing between the mother's milk and her infant's gut

microbiome, in particular for the commensal species *B. longum, B. bifidum, V. parvula, P.*

vulgatus, S. salivarius, R. mucilaginosa and for the pathobiont *K. pneumoniae*. Despite *K.*

pneumoniae potential association with silent sepsis in infants[73,74] and with mastitis and milk loss

in cows[75], the infants in our cohort showed no clinical manifestations nor had any of the mothers

reported mastitis/breast inflammation at the time of their study visit and milk collection. Strain-

level analysis also confirmed that the milk and the infant gut microbiome shared the same

strains of species typically found in the oral cavity (*R. mucilaginosa* and *S. salivarius*). This

could mean that either these species, initially present in the mother's milk, reach the infant gut

by first colonizing the oral cavity, or alternatively it could be that the infant colonizes the breast

milk shortly after birth with oral microbes acquired via other routes. As the first collection time

point in this study was at one month postpartum, we were unable to discern between these two

possible scenarios.

While strain sharing between the maternal and the infant gut has been extensively

investigated[8,11], only one metagenomic study has so far, to the best of our knowledge,

successfully identified strains shared between breast milk and the infant gut microbiome[57]. Our

results provide evidence that strain sharing between the infant gut and the maternal breast milk

occurs, even if at a rate considerably lower than what was previously found between the infant

and the maternal gut[11]. The fact that we were able to reconstruct strains from only ten taxa in

our milk samples, and yet all were found to be shared between mother's milk and infant gut,

indicates that our estimate for strain sharing between breast milk and infant gut is likely an

underestimate due to the low number of microbial reads in the milk samples. This is a common

challenge with this type of sample[28]. While prebiotics (e.g. HMOs) have been widely studied for

their potential role in shaping the infant gut microbiome[37,76–78], our findings support the

transmission of milk microbial strains as a contributing factor to infant gut microbiome assembly.

Strain-level analysis also revealed rates of microbial persistence in the infant gut
comparable with what previously reported[79], with 19% of the strains found in the infant stool
samples at one month persisting to six months of age, and 25% of strains shared between the
mother's milk and the infant gut at one month being identified also at six months of age. We also
found that delivery mode impacted microbial persistence rate in the infant gut: vaginally born
infants were characterized by significantly higher strain persistence over time than those born
via C-section. This result supports previous amplicon and species-level metagenomic studies
that associated C-section with a more diverse and less stable infant gut microbiome[15,80–82].

Resistome analysis showed that both the infant gut and the maternal milk harbor a
diverse landscape of antimicrobial resistance gene classes, mostly dominated by resistance
genes for tetracycline, aminoglycoside, and macrolide-lincosamide-streptogramin, even in
infants that had no recorded exposure to pre-, intra-, and postpartum antibiotics. Tetracycline is
not prescribed in pregnancy and its use is not recommended in children younger than 8 years
due to potential permanent discoloration of teeth[83]. Still, tetracycline resistance was the most
abundant antibiotic resistance class in both milk and infant stool samples at one month.
Aminoglycoside (e.g. gentamicin and streptomycin) and macrolides (e.g. azithromycin and
erythromycin) are widely used in the perinatal period[49], therefore resistance is potentially
acquired via antibiotic exposure. Milk and infant stool samples at one month were characterized
by higher prevalence and diversity in resistance classes, compared to later time points. In
addition, the majority of ARGs found in the infant gut at one month were associated with
predominance of non-bifidobacteria species, in particular *E. coli*. The presence of antibiotic
resistance in the newborn gut has been reported in previous studies[44,45,84], with indications of
resistance transmission from the maternal gut to the infant gut via mobile genetic elements[85].
Our results show a significant overlap between the resistome of infants and that of their
mother's milk, and that this overlap was particularly pronounced when evidence of strain sharing
was found. This suggests that the infant gut resistome is likely influenced to some degree by
acquisition from breastmilk, in addition to other mechanisms, such as mobile genetic elements,
vertical strain transmission and environmental exposure, and from other maternal body sites,
such as gut[85].

This study has several limitations. Milk sampling was performed via the use of breast
pumps, which could potentially impact milk microbiome composition[86]. The low microbial load in
breast milk samples limited our ability to detect strain sharing between milk and the infant gut,

particularly in the milk samples collected at three months, suggesting that the observed sharing patterns likely represent an underestimate. Notably, the samples with the higher number of strain sharing events corresponded to mother-infant pairs 1 and 173, which ranked first and third, respectively, in the number of reads mapped to MetaPhlAn4 marker gene database. These findings suggest that higher microbial yield and deeper sequencing depth for milk samples should be a priority for future metagenomic studies. The definition of a standardized and optimized protocol for human breast milk extraction to maximize microbial yield should also be prioritized by the field. In addition, as we sampled only milk and infant stool samples, we could not confirm that the strain sharing events we identified were indeed cases of microbial transmission from the maternal milk to the infant gut, rather than cases of strain acquisition by both the mother and the infant from external sources not investigated in this study. Finally, although infants were born across multiple hospitals, which could potentially influence microbial composition, all samples were collected using the same procedures, and the taxonomic composition of the samples did not cluster based on the sampling location.

In this work, we characterized the microbiome composition, function and antimicrobial resistance potential of the breast milk of mothers and the gut microbiome of their infants during the first six months postpartum. We found evidence of strain- and antimicrobial resistance gene-sharing between mother-infant pairs. Taken together, our results indicate that the maternal breast milk plays a role in infant gut microbiome and resistome establishment, development and temporal stability. Our results represent an important step towards the strain-level characterization of the maternal milk microbiome in relation to the infant's gut and its better representation in public repositories.

Methods

Sample collection

Participants in this study were enrolled as part of the Mother and Infants Linked for health (MILk) cohort[87–91]. All mothers were enrolled prenatally at the University of Minnesota in collaboration with HealthPartners Institute (Minneapolis, MN), and provided written informed consent. The study was approved by the Institutional Review Boards of the University of Minnesota, the University of Oklahoma, and HealthPartners Institute (STUDY00009021), and conducted in accordance with all relevant guidelines and regulations. All participant data included in this study were anonymized. Sample identifiers were randomized, infant age was recorded at the time of the study visit, and no hospital, location, or other personally identifiable information is included in the supplementary materials. The MILk Study is registered at ClinicalTrials.gov (NCT03301753). Eligibility criteria included a healthy, uncomplicated

pregnancy, and intention to exclusively breastfeed. All mothers were aged between 21-45
years, were not diabetic and non-smokers, and delivered a full-term infant. No case of breast
infection or mastitis was reported during the milk sample collection. All infants were singletons
and born at term with a birth weight that was appropriate for their gestational age, and were
exclusively breastfed to at least one month of age (Supplementary Table 1,2 and Extended
Data 1). All relevant metadata were collected via the hospital's electronic medical health records
and via questionnaires during sample collection (Supplementary Table 1,2 and Extended Data
1). Ethnicity and race of the participants are included in Supplementary Table 2. Samples were
collected between 2014 and 2023. Breast milk samples were collected at one and three months
postpartum. Milk collection was performed as follows: mothers fed their infant from one or both
of their breasts, until the infant was satisfied. After two hours, the milk was collected from the
right breast using a hospital grade electric breast pump (Medela Symphony; Medela, Inc., Zug,
Switzerland), until cessation of production. Each milk sample was gently mixed, and its volume
and weight were recorded. Aliquots were stored at -80 °C within 20 minutes of collection and
kept at that temperature until RNA/DNA extraction. Infant stool samples were collected at one
and six months of age. Stool samples were either collected from diapers during a study visit or
at home by the mother. In case of collection during a study visit, the sample was immediately
frozen at -80°C, while in case of home collection the sample was stored in 2 ml cryovials with
600 µl RNALater (Ambion/Invitrogen, Carlsbad, CA), and later stored at -80°C upon arrival to
the lab at the University of Minnesota, as described also in [87–91].

DNA extraction and metagenomic sequencing

Starting volume for extraction was 1 ml for breast milk, and the weight comparable to a pea-
sized scoop for the infant feces. DNA extraction was performed with PowerSoil Pro kit
(QIAGEN, Germantown, MD). Cell lysis was carried out by bead-beating for 10 minutes,
followed by elution with 100 µl of the provided elution solution. The extracted DNA was then
stored in microfuge tubes at -80°C and subsequently used to construct libraries for
metagenomic shotgun sequencing using the Illumina Nextera XT 1⁄4 kit (Illumina, San Diego,
CA, United States). DNA extractions for milk and infant stool samples were conducted
independently by experienced personnel. Metagenomic shotgun sequencing libraries for milk
and infant stool samples were then sequenced concurrently on an Illumina NovaSeq system
(Illumina, San Diego, CA) by the University of Minnesota Genomics Center. A small subset
(n=30) of milk samples at 1 month postpartum was selected to be re-sequenced at the target
depth of 30M reads per sample to improve detection of species, strains, functional pathways
and ARGs shared between mothers and their infants.. For those 30 samples, raw sequences
from the two batches were concatenated before preprocessing and analyzed together with the
rest of the samples in the downstream pipeline described below.

Quality filtering and removal of human reads

Host DNA was removed using paired-end mapping with Bowtie2[92] version 2.2.4 against a
human reference genome hg38. Unmapped paired-end reads were filtered using SAMtools[93]
version 1.9 with following parameters "samtools view -bS, samtools view -b -f 12 -F 256,
samtools sort -n -m 5G -@ 2, samtools fastq -@ 8 -0 /dev/null -s /dev/null -n". BEDtools[94] was

then used to convert the bam files to fastq files containing the non-human paired-end reads.
Then, adapter sequences were removed and the samples were filtered and trimmed using the
default parameters of Trimmomatic[95]. Host DNA was removed using the most complete version
of the human genome as reference (T2T-CHM13, including Y chromosome, available here:
https://s3-us-west-2.amazonaws.com/human-
pangenomics/T2T/CHM13/assemblies/analysis_set/chm13v2.0.fa.gz). Mean host DNA content
in breast milk samples was 97.3%. FastQC[96] was used to analyze the quality of the
metagenomic reads. Mean PHRED of the score was 33. Samples that had less than 1000 reads
that mapped to the Metaphlan4[40] database (mpa_vJan21_CHOCOPhlAnSGB_202103) were
discarded. Further pre-processing was conducted using the fastp[97,98], where reads underwent
length filtering at a minimum of 75 bp, quality trimming, additional adapter trimming, and poly-
G/poly-X trimming with the parameters "fastp -i Input R1 -o Output R1 -I Input R2 -O Output R2 -
-detect_adapter_for_pe -r -l 75 -x". After pre-processing, milk samples yielded 2.5±1.36 million
reads per sample, while stool samples yielded 21.4±8.01 million reads per sample.

Species-level taxonomic profiling and *B. longum* subspecies detection

Species-level taxonomic profiling was based on marker genes using MetaPhlAn4[40], with the
following parameters: "--bt2_ps sensitive" text. MetaPhlAn4 infers taxonomic prevalence and
abundance by using unique marker genes for 26,970 species-level genome bins[40]. As
MetaPhlAn4 has no marker genes to identify and distinguish *B. longum* subspecies (mainly *BL.
infantis* and *BL. longum*), we used a modified version of the standard MetaPhlAn4 database.
Details of the modified database can be found in the original publication[37], while the database
can be found on GitHub (https://github.com/yassourlab/MetaPhlAn-B.infantis/). Unclassified *B.
longum* subspecies was not included in the subspecies analyses. Merged abundance table was
created using the dedicated MetaPhlAn4 utils script. Contaminants found in the blank samples
as well as additional known contaminant species were removed before downstream analysis.
Milk samples with no identifiable bacterial taxa via MetaPhlAn4 were excluded from downstream
analysis (n=26). MetaPhlAn4 profiles are available in Supplementary Table 3. Heatmaps for
species-level taxonomic composition were generated with ComplexHeatmaps[99,100], using
euclidean distance for hierarchical clustering. Correlation results shown in Extended Data 10
were calculated using the sm_statCorr() function part of smplot2, using Pearson as correlation
method.

Maternal secretor status

662 Mothers with an inactive fucosyltransferase 2 (*FUT2*) gene are classified as non secretors.
663 Secretor status for the mothers in this study was obtained from[91] and was calculated based on
664 the concentration of 2'-fucosyllactose in breast milk.

Strain-level profiling and strain sharing

Strain-level profiling was performed using StrainPhlAn4[40], based on single-nucleotide variant
calling, using default parameters. Strain-sharing events between samples were identified based
on the output of the "strain_transmission.py" script, which is part of the StrainPhlAn4 release.

Detailed step-by-step description is available in
https://github.com/biobakery/MetaPhlAn/wiki/Strain-Sharing-Inference. Briefly, the
"strain_transmission.py" script takes as input the phylogenetic tree computed by StrainPhlAn4
for each species-level genome bin (SGB), SGB ID, and metadata table containing information
on subjectID, relation (e.g. mother-infant pair), and sampling time point. Precomputed
thresholds provided in "VallesColomerM_2022_Nov19_thresholds.tsv" were used when
possible, otherwise the default threshold of 0.03 was provided to the script. The script outputs a
list of all sample pairs in which strain sharing was detected. The collated output of the
"strain_transmission.py" script is available as Supplementary Table 4. Phylogenetic trees were
visualized with iTOL[101].

Alpha and beta diversity

680 Both alpha diversity and ordination plots were computed using the "diversity" and "vegdist"
681 functions, respectively, included in the "vegan" R package (v2.6-4).

Functional profiling

Functional prediction was performed with HUMAnN3[43]. Biosynthetic potential of essential amino
acids were calculated by searching the output pathways for the following keywords: "L-histidine
biosynthesis", "L-isoleucine biosynthesis", "L-isoleucine biosynthesis", "L-lysine biosynthesis",
"L-methionine biosynthesis", "L-phenylalanine biosynthesis", "L-threonine biosynthesis", and "L-
valine biosynthesis". Profiles relative abundance plots show 95% confidence interval using
bootstrapping with 1000 repetitions using the Hmisc R library (v5.0-1). Raw functional profiles
are available in Supplementary Table 5.

Antimicrobial resistance genes prediction

Antimicrobial resistance genes were predicted from raw metagenomic short reads using
DeepARG v1.0.2[50]. DeepARG leverages deep learning models to identify over 30 antimicrobial
resistance classes. Predicted resistance is classified as "predicted" or "potential" by the models.
To reduce false positives detection, ARG predictions classified as "multidrug" or "unclassified"
were excluded from downstream analyses and a minimum 95% identity threshold was applied.
Raw DeepARG profiles are available in Supplementary Table 6. Correlation results shown in
Figure 5D-E were calculated using the sm_statCorr() function part of smplot2, using Spearman
as correlation method. Permutation analysis in Figure 5G was performed as following: maternal
milk sample names were randomly permuted, while data and the infant sample names were
preserved, generating pseudo-mother-infant pairs, and the mean number of antimicrobial
resistance genes shared between the pseudo-mother-infant pairs was calculated. This was
repeated 1000 times, and the distribution of the mean values obtained from pseudo-mother-
infant pairs was compared to the mean value obtained from the real mother-infant pairs.

Statistics & Reproducibility

705 Statistical analysis was done in R[102] version 4.2.2 (2022-10-31). All figures if not indicated
706 otherwise were drawn with ggplot[103] version 3.4.1. All analyses have been performed with open

source software referenced in the Methods section. Statistical significance was set at $p < 0.05$, and Benjamini-Hochberg false discovery rate (FDR) correction was applied where appropriate to account for multiple testing. Unless otherwise stated, boxplots show the median as the center line, the interquartile range (25th to 75th percentile) as the box, whiskers extending to 1.5× the interquartile range, and outliers are not shown. No statistical method was used to predetermine sample size and no data randomization was performed. No data were excluded from the analyses. This was an observational study and participants received no interventions. Participants were not assigned to experimental groups; instead, they were grouped retrospectively based on metadata (e.g., breastfeeding status). The investigators were not blinded to allocation during experiments and outcome assessment.

Data availability

The raw metagenomic data generated in this study have been deposited in the NCBI Sequence Read Archive (SRA) database under BioProject accession numbers PRJNA1019702 and PRJNA1198101. The metadata associated with this study are provided in the Supplementary Information file. All supplementary tables are also available on Zenodo (https://zenodo.org/records/17317218).

Code availability

Code is available on github https://github.com/blekhmanlab/milk_infant_microbiome.

References

1. Legoux, F. *et al.* Microbial metabolites control the thymic development of mucosal-associated invariant T cells. *Science* **366**, 494–499 (2019).

2. Zegarra-Ruiz, D. F. *et al.* Thymic development of gut-microbiota-specific T cells. *Nature* **594**, 413–417 (2021).

3. LeBlanc, J. G. *et al.* Bacteria as vitamin suppliers to their host: a gut microbiota perspective. *Curr. Opin. Biotechnol.* **24**, 160–168 (2013).

4. Yao, Y. *et al.* The Role of Microbiota in Infant Health: From Early Life to Adulthood. *Front. Immunol.* **12**, 708472 (2021).

5. Wilkins, A. T. & Reimer, R. A. Obesity, Early Life Gut Microbiota, and Antibiotics. *Microorganisms* **9**, (2021).

6. Asnicar, F. *et al.* Studying Vertical Microbiome Transmission from Mothers to Infants by Strain-Level Metagenomic Profiling. *mSystems* **2**, (2017).

7. Korpela, K. *et al.* Selective maternal seeding and environment shape the human gut microbiome. *Genome Res.* **28**, 561–568 (2018).

8. Podlesny, D. & Fricke, W. F. Strain inheritance and neonatal gut microbiota development: A meta-analysis. *Int. J. Med. Microbiol.* **311**, 151483 (2021).

9. Duranti, S. *et al.* Maternal inheritance of bifidobacterial communities and bifidophages in infants through vertical transmission. *Microbiome* **5**, 66 (2017).

10. Enav, H., Bäckhed, F. & Ley, R. E. The developing infant gut microbiome: A strain-level view. *Cell Host Microbe* **30**, 627–638 (2022).

11. Ferretti, P. *et al.* Mother-to-Infant Microbial Transmission from Different Body Sites Shapes the Developing Infant Gut Microbiome. *Cell Host Microbe* **24**, 133–145.e5 (2018).

12. Song, S. J. *et al.* Naturalization of the microbiota developmental trajectory of Cesarean-born neonates after vaginal seeding. *Med* **2**, 951–964.e5 (2021).

13. Shao, Y. *et al.* Stunted microbiota and opportunistic pathogen colonization in caesarean-

section birth. *Nature* **574**, 117–121 (2019).

14. Mortensen, M. S. *et al.* Modeling transfer of vaginal microbiota from mother to infant in early

life. *Elife* **10**, e57051 (2021).

15. Mitchell, C. M. *et al.* Delivery Mode Affects Stability of Early Infant Gut Microbiota. *Cell Rep*

Med **1**, 100156 (2020).

16. CDC. Recommendations and benefits. *Centers for Disease Control and Prevention*

https://www.cdc.gov/nutrition/infantandtoddlernutrition/breastfeeding/recommendations-

benefits.html (2023).

17. Ferretti, P. *et al.* C. difficile may be overdiagnosed in adults and is a prevalent commensal

in infants. *Elife* **12**, (2023).

18. Ballard, O. & Morrow, A. L. Human milk composition: nutrients and bioactive factors.

Pediatr. Clin. North Am. **60**, 49–74 (2013).

19. Notarbartolo, V., Giuffrè, M., Montante, C., Corsello, G. & Carta, M. Composition of Human

Breast Milk Microbiota and Its Role in Children's Health. *Pediatr Gastroenterol Hepatol Nutr*

25, 194–210 (2022).

20. Das, L., Virmani, R., Sharma, V., Rawat, D. & Singh, Y. Human Milk Microbiota:

Transferring the Antibiotic Resistome to Infants. *Indian J. Microbiol.* **59**, 410–416 (2019).

21. Pärnänen, K. *et al.* Maternal gut and breast milk microbiota affect infant gut antibiotic

resistome and mobile genetic elements. *Nat. Commun.* **9**, 3891 (2018).

22. Xue, M. *et al.* Breastfeeding and risk of childhood asthma: a systematic review and meta-

analysis. *ERJ Open Res* **7**, (2021).

23. Dogaru, C. M., Nyffenegger, D., Pescatore, A. M., Spycher, B. D. & Kuehni, C. E.

Breastfeeding and childhood asthma: systematic review and meta-analysis. *Am. J.*

Epidemiol. **179**, 1153–1167 (2014).

24. Shenhav, L. *et al.* Microbial colonization programs are structured by breastfeeding and

guide healthy respiratory development. *Cell* **187**, 5431–5452.e20 (2024).

25. Uwaezuoke, S. N., Eneh, C. I. & Ndu, I. K. Relationship Between Exclusive Breastfeeding

and Lower Risk of Childhood Obesity: A Narrative Review of Published Evidence. *Clin.*

Med. Insights Pediatr. **11**, 1179556517690196 (2017).

26. Patelarou, E. *et al.* Current evidence on the associations of breastfeeding, infant formula,

and cow's milk introduction with type 1 diabetes mellitus: a systematic review. *Nutr. Rev.*

70, 509–519 (2012).

27. Kull, I., Wickman, M., Lilja, G., Nordvall, S. L. & Pershagen, G. Breast feeding and allergic

diseases in infants-a prospective birth cohort study. *Arch. Dis. Child.* **87**, 478–481 (2002).

28. Ruiz, L., García-Carral, C. & Rodriguez, J. M. Unfolding the Human Milk Microbiome

Landscape in the Omics Era. *Front. Microbiol.* **10**, 1378 (2019).

29. Williams, J. E. *et al.* Human milk microbial community structure is relatively stable and

related to variations in macronutrient and micronutrient intakes in healthy lactating women.

J. Nutr. **147**, 1739–1748 (2017).

30. Bogaert, D. *et al.* Mother-to-infant microbiota transmission and infant microbiota

development across multiple body sites. *Cell Host Microbe* **31**, 447–460.e6 (2023).

31. Qi, C. *et al.* Lactation-dependent vertical transmission of natural probiotics from the mother

to the infant gut through breast milk. *Food Funct.* **13**, 304–315 (2022).

32. Van Rossum, T., Ferretti, P., Maistrenko, O. M. & Bork, P. Diversity within species:

interpreting strains in microbiomes. *Nat. Rev. Microbiol.* **18**, 491–506 (2020).

33. Mantziari, A. & Rautava, S. Factors influencing the microbial composition of human milk.

Semin. Perinatol. **45**, 151507 (2021).

34. Saturio, S. *et al.* Role of bifidobacteria on infant health. *Microorganisms* **9**, 2415 (2021).

35. Fujimura, K. E. *et al.* Neonatal gut microbiota associates with childhood multisensitized

atopy and T cell differentiation. *Nat. Med.* **22**, 1187–1191 (2016).

36. Zhang, S. & Dang, Y. Roles of gut microbiota and metabolites in overweight and obesity of

children. *Front. Endocrinol. (Lausanne)* **13**, 994930 (2022).

37. Ennis, D., Shmorak, S., Jantscher-Krenn, E. & Yassour, M. Longitudinal quantification of

Bifidobacterium longum subsp. infantis reveals late colonization in the infant gut

independent of maternal milk HMO composition. *Nat. Commun.* **15**, 894 (2024).

38. Asakuma, S. *et al.* Physiology of consumption of human milk oligosaccharides by infant

gut-associated bifidobacteria. *J. Biol. Chem.* **286**, 34583–34592 (2011).

39. Henrick, B. M. *et al.* Bifidobacteria-mediated immune system imprinting early in life. *Cell*

184, 3884–3898.e11 (2021).

40. Blanco-Míguez, A. *et al.* Extending and improving metagenomic taxonomic profiling with

uncharacterized species using MetaPhlAn 4. *Nat. Biotechnol.* (2023) doi:10.1038/s41587-

023-01688-w.

41. Singh, P. *et al.* Unveiling the dynamics of the breast milk microbiome: impact of lactation

stage and gestational age. *J. Transl. Med.* **21**, 784 (2023).

42. Selma-Royo, M., Calvo Lerma, J., Cortés-Macías, E. & Collado, M. C. Human milk

microbiome: From actual knowledge to future perspective. *Semin. Perinatol.* **45**, 151450

(2021).

43. Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse

microbial communities with bioBakery 3. *Elife* **10**, (2021).

44. Bäckhed, F. *et al.* Dynamics and Stabilization of the Human Gut Microbiome during the

First Year of Life. *Cell Host Microbe* **17**, 852 (2015).

45. Lebeaux, R. M. *et al.* The infant gut resistome is associated with E. coli and early-life

exposures. *BMC Microbiol.* **21**, 201 (2021).

46. Kim, D.-W. & Cha, C.-J. Antibiotic resistome from the One-Health perspective:

understanding and controlling antimicrobial resistance transmission. *Exp. Mol. Med.* **53**,

301–309 (2021).

47. Chen, P.-W., Tseng, S.-Y. & Huang, M.-S. Antibiotic Susceptibility of Commensal Bacteria

from Human Milk. *Curr. Microbiol.* **72**, 113–119 (2016).

48. Behari, P., Englund, J., Alcasid, G., Garcia-Houchins, S. & Weber, S. G. Transmission of

methicillin-resistant Staphylococcus aureus to preterm infants through breast milk. *Infect.*

Control Hosp. Epidemiol. **25**, 778–780 (2004).

49. Samarra, A. *et al.* Maternal-infant antibiotic resistance genes transference: what do we

know? *Gut Microbes* **15**, 2194797 (2023).

50. Arango-Argoty, G. *et al.* DeepARG: a deep learning approach for predicting antibiotic

resistance genes from metagenomic data. *Microbiome* **6**, 23 (2018).

51. Kim, S. Y. & Yi, D. Y. Analysis of the human breast milk microbiome and bacterial

extracellular vesicles in healthy mothers. *Exp. Mol. Med.* **52**, 1288–1297 (2020).

52. Ruan, J.-W. *et al.* The composition of the maternal breastmilk microbiota influences the

microbiota network structure during early infancy. *J. Microbiol. Immunol. Infect.* **56**, 1084–

1097 (2023).

53. Li, Y. *et al.* The effect of breast milk Microbiota on the composition of infant gut Microbiota:

A cohort study. *Nutrients* **14**, 5397 (2022).

54. Kordy, K. *et al.* Contributions to human breast milk microbiome and enteromammary

transfer of Bifidobacterium breve. *PLoS One* **15**, e0219633 (2020).

55. Turroni, F. *et al.* Bifidobacteria and the infant gut: an example of co-evolution and natural

selection. *Cell. Mol. Life Sci.* **75**, 103–118 (2018).

56. Zhang, Q. *et al.* Optimization of metagenomic detection method for human breast milk

microbiome. *Microbiology* (2024).

57. Feehily, C. *et al.* Detailed mapping of Bifidobacterium strain transmission from mother to

infant via a dual culture-based and metagenomic approach. *Nat. Commun.* **14**, 3015

(2023).

58. Martín, R. *et al.* Isolation of bifidobacteria from breast milk and assessment of the

bifidobacterial population by PCR-denaturing gradient gel electrophoresis and quantitative

855   real-time PCR. *Appl. Environ. Microbiol.* **75**, 965–969 (2009).

856 59. Collado, M. C., Delgado, S., Maldonado, A. & Rodríguez, J. M. Assessment of the bacterial

857   diversity of breast milk of healthy women by quantitative real-time PCR. *Lett. Appl.*

858   *Microbiol.* **48**, 523–528 (2009).

859 60. Durazzi, F. *et al.* Comparison between 16S rRNA and shotgun sequencing data for the

860   taxonomic characterization of the gut microbiota. *Sci. Rep.* **11**, 3030 (2021).

861 61. Peterson, D. *et al.* Comparative Analysis of 16S rRNA Gene and Metagenome Sequencing

862   in Pediatric Gut Microbiomes. *Front. Microbiol.* **12**, 670336 (2021).

863 62. Sim, K. *et al.* Improved detection of bifidobacteria with optimised 16S rRNA-gene based

864   pyrosequencing. *PLoS One* **7**, e32543 (2012).

865 63. Meisel, J. S. *et al.* Skin Microbiome Surveys Are Strongly Influenced by Experimental

866   Design. *J. Invest. Dermatol.* **136**, 947–956 (2016).

867 64. Ward, T. L., Hosid, S., Ioshikhes, I. & Altosaar, I. Human milk metagenome: a functional

868   capacity analysis. *BMC Microbiol.* **13**, 116 (2013).

869 65. Zhang, B. *et al.* Impact of bead-beating intensity on the genus- and species-level

870   characterization of the gut microbiome using amplicon and complete 16S rRNA gene

871   sequencing. *Front. Cell. Infect. Microbiol.* **11**, 678522 (2021).

872 66. Shimamura, S. *et al.* Relationship between oxygen sensitivity and oxygen metabolism of

873   Bifidobacterium species. *J. Dairy Sci.* **75**, 3296–3306 (1992).

874 67. Ruiz, L. *et al.* Molecular clues to understand the aerotolerance phenotype of

875   Bifidobacterium animalis subsp. lactis. *Appl. Environ. Microbiol.* **78**, 644–650 (2012).

876 68. Schell, M. A. *et al.* The genome sequence of Bifidobacterium longum reflects its adaptation

877   to the human gastrointestinal tract. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 14422–14427 (2002).

878 69. Zhang, D. *et al.* Oxygen tolerance mechanism of Bifidobacterium animalis AR668-R1 based

879   on genomic and phenotypic analyses. *Lebenson. Wiss. Technol.* **215**, 117207 (2025).

880 70. Lopez Leyva, L., Brereton, N. J. B. & Koski, K. G. Emerging frontiers in human milk

microbiome research and suggested primers for 16S rRNA gene analysis. *Comput. Struct.*

Biotechnol. J. **19**, 121–133 (2021).

71. Heisel, T. *et al.* Bacterial, fungal, and interkingdom microbiome features of exclusively

breastfeeding dyads are associated with infant age, antibiotic exposure, and birth mode.

Front. Microbiol. **13**, 1050574 (2022).

72. Yassour, M. *et al.* Strain-Level Analysis of Mother-to-Child Bacterial Transmission during

the First Few Months of Life. *Cell Host Microbe* **24**, 146–154.e4 (2018).

73. Dorota, P. *et al.* Klebsiella pneumoniae in breast milk: a cause of sepsis in neonate. *Arch*

Med **9**, (2017).

74. Vasilyev, I. Y., Nikolaeva, I. V., Siniagina, M. N., Kharchenko, A. M. & Shaikhieva, G. S.

Multidrug-Resistant Hypervirulent Klebsiella pneumoniae Found Persisting Silently in Infant

Gut Microbiota. *Int. J. Microbiol.* **2020**, 4054393 (2020).

75. Liu, K., Zhang, L., Gu, X. & Qu, W. The Prevalence of Klebsiella spp. Associated With

Bovine Mastitis in China and Its Antimicrobial Resistance Rate: A Meta-Analysis. *Front Vet*

Sci **9**, 757504 (2022).

76. Wiciński, M., Sawicka, E., Gębalski, J., Kubiak, K. & Malinowski, B. Human milk

oligosaccharides: Health benefits, potential applications in infant formulas, and

pharmacology. *Nutrients* **12**, 266 (2020).

77. Masi, A. C. & Stewart, C. J. Untangling human milk oligosaccharides and infant gut

microbiome. *iScience* **25**, 103542 (2022).

78. Kijner, S., Kolodny, O. & Yassour, M. Human milk oligosaccharides and the infant gut

microbiome from an eco-evolutionary perspective. *Curr. Opin. Microbiol.* **68**, 102156

(2022).

79. Lou, Y. C. *et al.* Infant gut strain persistence is associated with maternal origin, phylogeny,

and traits including surface adhesion and iron acquisition. *Cell Rep. Med.* **2**, 100393 (2021).

80. Davis, E. C. *et al.* Gut microbiome and breast-feeding: Implications for early immune

development. *J. Allergy Clin. Immunol.* **150**, 523–534 (2022).

81. Kim, G. *et al.* Delayed establishment of gut Microbiota in infants delivered by Cesarean

section. *Front. Microbiol.* **11**, 2099 (2020).

82. Reyman, M. *et al.* Impact of delivery mode-associated gut microbiota dynamics on health in

the first year of life. *Nat. Commun.* **10**, 4997 (2019).

83. Tetracycline (class) (oral route, parenteral route). https://www.mayoclinic.org/drugs-

supplements/tetracycline-class-oral-route-parenteral-route/proper-use/drg-20069585

(2023).

84. Leo, S. *et al.* Metagenomics analysis of the neonatal intestinal resistome. *Front Pediatr* **11**,

1169651 (2023).

85. Vatanen, T. *et al.* Mobile genetic elements from the maternal microbiome shape infant gut

microbial assembly and metabolism. *Cell* **185**, 4921–4936.e15 (2022).

86. Reyes, S. M. *et al.* Pumping supplies alter the microbiome of pumped human milk: An in-

home, randomized, crossover trial. *Am. J. Clin. Nutr.* **114**, 1960–1970 (2021).

87. Fields, D. A. *et al.* Associations between human breast milk hormones and adipocytokines

and infant growth and body composition in the first 6 months of life. *Pediatr. Obes.* **12**

Suppl 1, 78–85 (2017).

88. Sadr Dadres, G. *et al.* Relationship of Maternal Weight Status Before, During, and After

Pregnancy with Breast Milk Hormone Concentrations. *Obesity* **27**, 621–628 (2019).

89. Whitaker, K. M. *et al.* Associations of Maternal Weight Status Before, During, and After

Pregnancy with Inflammatory Markers in Breast Milk. *Obesity* **26**, 1659–1660 (2018).

90. Johnson, K. E. *et al.* Human cytomegalovirus in breast milk is associated with milk

composition and the infant gut microbiome and growth. *Nat. Commun.* **15**, 6216 (2024).

91. Johnson, K. E. *et al.* Human milk variation is shaped by maternal genetics and impacts the

infant gut microbiome. *Cell Genom.* **4**, 100638 (2024).

92. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*

9, 357–359 (2012).

93. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

94. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

95. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

96. Andrews, S. & Others. FastQC: a quality control tool for high throughput sequence data.

Preprint at (2010).

97. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

98. Chen, S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication

using fastp. *Imeta* **2**, e107 (2023).

99. Gu, Z. Complex heatmap visualization. *Imeta* **1**, (2022).

100. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

101. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. *Nucleic Acids Res.* **49**, W293–W296 (2021).

102. Foundation for Statistical Computing, R. R. R: A language and environment for statistical

computing. *RA Lang Environ Stat Comput.*

103. Wickham, H. & Wickham, M. H. The ggplot package. *Google Scholar.*

Acknowledgements

We thank the MILk Study staff and participants (https://www.milkresearch.umn.edu), and members of the Albert and Blekhman labs for support and helpful discussions related to this project. This work could not have been completed without the scientific collaboration of the Health Partners Institute (Bloomington, MN) and the University of Minnesota Medical School, Department of Obstetrics, Gynecology, and Women's Health. This work was supported by the resources and staff at the Masonic Institute for the Developing Brain at the University of Minnesota (https://www.midb.umn.edu), the University of Oklahoma Health Sciences Center Harold Hamm Diabetes Center (https://www.ouhealth.com/find-a-location/ou-health-harold-hamm-diabetes-center) and the University of Minnesota Genomics Center (https://genomics.umn.edu). We are thankful for the computational resources provided by the Minnesota Supercomputing Institute (https://www.msi.umn.edu/) and by the University of Chicago Research Computing Center (https://rcc.uchicago.edu/). The MILk Study was supported by NIH/NICHD grant R01HD080444 (EWD and DAF). Additional support for this study was provided by the Masonic Cross Departmental Research Grant from the University of Minnesota Department of Pediatrics (FWA, RB, EWD, and CAG); the Masonic Children's Hospital Research Fund Award (CAG, EWD, and DK); NIH/NICHD grants R01HD109830 (RB, FWA, EWD, CAG, and KEJ), R21HD099473 (CAG and EWD), and F32HD105364 (KEJ); and a Faculty Research Development Grant from the University of Minnesota Office of Academic and Clinical Affairs (CAG, EWD, KEJ, and DK).

Author contributions

Conceptualization, R.B., C.A.G., E.W.D., and D.K.; Sample and metadata collection, E.W.D, S.G., and C.A.G; Samples processing and data analysis, P.F., M.A. and M.R.; Writing – original draft, P.F.; Writing – review & editing, P.F., M.A., K.J., M.R., T.H., D.A.F., F.W.A., C.A.G., E.W.D., and R.B.; Supervision, P.F. and R.B.; Funding acquisition, E.W.D., D.A.F., and R.B. All authors read and approved the final manuscript. The Authors declare no conflict of interest.

Competing Interests

The Authors declare no conflict of interest.

Figure legends

Figure 1. (A) Study design overview for the 507 samples collected from 195 mother-infant pairs.
Infant stool samples (n=334) were collected at one and six months of life. Maternal breast milk
samples (n=173) were collected one and three months after delivery. (B) Shannon diversity
distribution for infant stool and maternal milk samples over time. P-values calculated using
paired t-test (* for $p \leq 0.05$, ** for $p \leq 0.01$ and **** for $p \leq 0.0001$). Samples included: 122 for
milk at one month, 25 for milk three months, 175 for stool one month and 159 for stool at six
months of age. Taxonomic composition of (C) the most prevalent and abundant species found
in the human breast milk and (D) in the infant gut microbiome samples in relation to sample
collection time point, predominance group and other relevant infant metadata. The
predominance group identifies the most abundant species in each sample. Ordination plot
based on Bray-Curtis distance between samples for PCoA1 and PCoA2 (E) and PCoA1 and
PCoA3 (F), colored by body site of origin and sampling time. Percentage of variance explained
(PVE) is reported in parentheses.

1000

Figure 2. (A) Prevalence of each predominance group in milk and infant stool samples and (B)
the transition of samples between predominance groups over time. Each sample is assigned
one of four predominance groups indicated by the different colors. Numbers of samples
included: n=145 for milk at one month, n=25 for milk at three months, n=175 for stool samples
at one month and n=159 for stool samples at six months of age. (C) Prevalence of
Bifidobacterium species at one and six months of age. Prevalence at six months is further
stratified by exclusive breastfeeding status (n=121 for exclusive breastfeeding; n=34 for
mixed/formula diet). Samples lacking breastfeeding information at six months were excluded
from the stratified subgroups (n=4). Bonferroni adjusted p-values were calculated using Fisher's
exact test. (D) Relative abundance of *Bifidobacterium* species in infants at one and six months
of age. Relative abundances at six months of age are further stratified by exclusive
breastfeeding status. Relative abundances were calculated only on samples in which the
species was present. P-values calculated using Wilcoxon rank sum, * for $p \leq 0.05$, ** $p \leq 0.01$,
*** for $p \leq 0.001$ and **** for $p \leq 0.0001$. Reported p-values are adjusted using Bonferroni
method. (E) Species persistence in the infant gut across all samples and (F) stratified by
breastfeeding at six months. See Extended Data 7 for plots with separation between *BL. infantis*
and *BL. longum*. (G) Relative abundance of some of the differentially abundant species between
one and six months when divided by breastfeeding (BF) at six months. P-values calculated with

paired t-test and adjusted with BH correction. **** for P ≤ 0.0001, *** for P ≤ 0.001, ** for P ≤

0.01 and * for P ≤ 0.05.

1021

Figure 3. (A) Percentage of the infant stool microbiome that is shared with the respective

mother, across all mother-infant pairs. Species sharing between milk samples at one month

postpartum and infant stool samples at one month and six months of age are shown on the left

and right boxplot, respectively. T-test, * for p ≤ 0.05. Only mother-infant pairs for which milk at

one month and stool samples at one and six months were available were included (n=108

pairs). (B) Overview of the ten most frequently shared species between the infant stool samples

at one month of age and the respective mother's milk samples at one month postpartum. *BL.*

infantis = B. longum subsp*. infantis; BL. longum = B. longum* subsp*. longum*; (C) Mean

percentage of mother-infant pairs in species belonging to the *Bifidobacterium* genus were

shared between the milk and infant stool samples (values reported in gray), were found only in

the milk samples or were found only in the infant stool samples. Species sharing is calculated

comparing milk at one month versus infant stool samples at one and six months, and milk at 3

months versus infant stool samples at 6 months. Only mother-infant couples for which at least

one milk and one infant stool sample was collected are included. (D) Examples of strain sharing

between milk and infant stool samples for *B. longum* (SGB17248), *Phocaeicola vulgatus*

(SGB1814), *Streptococcus salivarius* (SGB8005) and *Klebsiella pneumoniae* (SGB10115),

highlighted in black. Instances of strains found within the same infant at both one and six

months of age (persistent strains) are highlighted with gray. (E) Percentage of strains shared

between mother-infant pairs divided by infant stool collection time point. (F) Taxonomic

composition of strains found to be persistent in the infant gut over time (until six months of age).

(G) Percentage of strains in infant stool at one month that persisted at six months divided by

delivery mode. For each infant, persistence was calculated as the number of strains shared

between the one- and six-month timepoints, divided by the total number of strains detected at

one month. Each dot is a mother-infant pair. P-value calculated with t-test, * for p ≤ 0.05.

1046

1047   **Figure 4.** Most abundant pathways identified in (A) breastmilk at one and three months

1048   postpartum and (B) infant gut at one and six months of life. (C) Relative abundance of pathways

1049   involved in the biosynthesis of essential amino acids across sample type, collection time point

1050   and breastfeeding (BF). Error bars represent a 95% confidence interval calculated by

1051   bootstrapping (1000 times). P-values calculated using t-test, * for p ≤ 0.05, ** p ≤ 0.01, *** for p

≤ 0.001 and **** for p ≤ 0.0001. Reported p-values are adjusted using Bonferroni method. (D) P-values of Spearman correlation between the abundance of all metabolic pathways shared between the maternal breast milk and infant stool samples for each mother-infant pair. P-values are corrected for multiple testing using Benjamini Hochberg correction and are shown in $\log_{10}$ scale. Only the top 25 mother-infant pairs are shown. Circle size denotes the correlation coefficient (rho), and the orange line denotes the significance threshold ($\log_{10}$ of p-value=0.01).

Figure 5. (A) Prevalence of the antimicrobial resistance genes (ARGs) classes predicted in maternal milk and the infant gut microbiome over time, calculated as percentage over the total number of detected ARGs for each sample type. (B) ARGs carriage in infant stool samples, divided by collection time point, infant metadata, predominance group and ARGs class. (C) Number of detected ARGs in infant stool samples dominated by bifidobacteria and non-bifidobacteria species at one and six months of life. P-values calculated using t-test, ** for p ≤ 0.0001, * for p<0.05. (D) Correlation between ARG carriage in maternal milk and infant stool samples, and (E) between infant stool samples at one month and six months. Each dot is a combination of a mother-infant pair and a predicted ARG class. Spearman's R-values (correlation coefficient) and p-values are reported. (F) Percentage of ARG genes shared between at least one maternal and one infant sample, for each mother-infant pair. Mean value is indicated by the orange line. The mother-infant pairs for which strain sharing events were identified are highlighted with the arrow. (G) Distribution of the percentage of shared ARGs between at least one maternal milk and one infant stool sample on permuted mother-infant pairs (pseudo-pairs). The mean value obtained from real mother-infant pairs is indicated by the orange line. (H) Percentage of mother-infant pairs sharing antimicrobial resistance genes, divided by ARG class. For each ARG class, percentage value was calculated on the total number of mother-infant pairs in which that ARG class was identified in at least one sample. The most frequently shared antimicrobial resistance genes per class are highlighted in orange.

A Infant stool - milk sharing

Species-level

% of taxa in the infant microbiome shared with the respective mother's milk

shared between: Milk 1M - Stool 1M, Milk 1M - Stool 6M

*

B Most shared species at 1 month postpartum

BL. infantis
BL. longum
S. epidermidis
B. breve
S. salivarius
S. aureus
S. mitis
P. vulgatus
S. hominis
R. mucilaginosa

% of mother-infant pairs

1M ↔ 1M

C Shared bifidobacteria (1M and 6M combined)

BL. infantis — 54.4%
BL. longum — 47.2%
B. breve — 30%
B.bifidum — 11%

mean % mother-infant pairs

Found: in both milk and infant stool | only milk | only infant

D

Strain-level

B. longum
Tree scale: 0.1

Infant 107
Infant 91
Infant 171
Infant 81
Infant 173
Infant 189
Mother-Infant Pair 1

S. salivarius
Tree scale: 0.1

Mother-Infant Pair 97
Infant 110
Infant 107
Infant 137
Mother-Infant Pair 144
Mother-Infant Pair 53

P. vulgatus
Tree scale: 0.1

Infant 25
Infant 126
Infant 195
Infant 109
Infant 65
Mother-Infant Pair 1

K. pneumoniae
Tree scale: 0.1

Infant 169
Infant 147
Mother-Infant Pair 173

Sample Type
Milk 1M
Stool 1M
Stool 6M

E Milk - gut strain sharing

Percentage of M-I strain sharing events %

shared Milk 1M - Stool 1M
shared Milk 1M - Stool 6M
shared Milk 1M - Stool 1M and 6M

F Taxonomy of the persistent strains in the infant gut

% of persistent strains in the infant gut between 1M and 6M

B. fragilis (8.5%)
P. vulgatus (8.5%)
B. bifidum (8.5%)
B. longum (13.4%)
E. coli (23.2%)
Other (37.8%)

G Persistent strains by delivery mode

Percentage of persistent strains per infant

*

Delivery mode: C-section (37), Vaginal (134)

A Infant Stool

Rel. abundance
10^1 — 10^{-2} — 10^{-3} — 0

Sample Type
Predominance group

Pathway Class

- peptidoglycan maturation
- glycolysis IV
- pentose phosphate pathway I
- L–valine biosynthesis
- pyruvate fermentation to isobutanol
- L–isoleucine biosynthesis I
- superpath. branched chain amino acid biosynthesis
- L–isoleucine biosynthesis III
- sucrose biosynthesis II
- glycogen degradation II
- guanosine ribonucleotides de novo biosynthesis
- L–lysine biosynthesis III
- L–lysine biosynthesis VI
- UMP biosynthesis I
- UDP–N–acetylmuramoyl–pentapeptide biosynth. II
- UDP–N–acetylmuramoyl–pentapeptide biosynth. I
- peptidoglycan biosynthesis I
- peptidoglycan biosynthesis III
- chorismate biosynthesis I
- chorismate biosynthesis from 3–dehydroquinate
- folate transformations II (plants)
- superpath. aromatic amino acid biosynthesis
- S–adenosyl–L–methionine salvage I
- 5–aminoimidazole ribonucleotide biosynthesis II
- superpath. 5–aminoimidazole ribonucleotide biosynth.
- L–arginine biosynthesis I
- superpath. L–isoleucine biosynthesis I
- superpath. L–threonine biosynthesis
- L–histidine biosynthesis
- dTDP–β–L–rhamnose biosynthesis

B Milk

Rel. abundance
10^1 — 10^{-3} — 10^{-4} — 0

Sample Type
Predominance group

- L–valine biosynthesis
- guanosine ribonucleotides de novo biosynth.
- sucrose biosynthesis II
- pentose phosphate pathway I
- pyruvate fermentation to isobutanol
- UDP–N–acetyl–D–glucosamine biosynth. I
- UMP biosynthesis I
- inosine–5'–phosphate biosynthesis I
- inosine–5'–phosphate biosynthesis II
- peptidoglycan maturation
- adenosine deoxyribonucleotides de novo biosynth. II
- guanosine deoxyribonucleotides de novo biosynth. II
- superpath. guanosine nucleotides de novo biosynth.
- 5–aminoimidazole ribonucleotide biosynthesis II
- superpath. 5–aminoimidazole ribonucleotide biosynth.
- 5–aminoimidazole ribonucleotide biosynthesis I
- L–isoleucine biosynthesis I
- superpath. branched chain amino acid biosynth.
- S–adenosyl–L–methionine salvage II
- adenine and adenosine salvage III

Sample Type
Stool 1M   Milk 1M
Stool 6M   Milk 3M

Predominance group
B. longum   B. bifidum
B. breve   Other

Class pathway
Cell structure   Amino acid biosynthesis
Cell metabolism   Nucleotide biosynthesis
  Other

C Essential aa biosynthesis

Rel. abundance

2×10^{-4}

1×10^{-4}

Milk		Stool		
1M	3M	1M	6M	6M
(146)	(27)	(175)	(121)	(34)
		Excl. BF	Excl. BF	Non excl. BF

D Correlation of abundances of shared pathways within mother-infant pairs

Spearman's correlation rho
−0.25, 0.00, 0.25, 0.50, 0.75

Mother-infant pair:
Pair 128, Pair 144, Pair 101, Pair 40, Pair 50, Pair 22, Pair 149, Pair 119, Pair 95, Pair 71, Pair 28, Pair 93, Pair 41, Pair 58, Pair 165, Pair 135, Pair 181, Pair 3, Pair 161, Pair 157, Pair 172, Pair 173, Pair 1

$\log_{10}$(adjusted p-value)
−20, −15, −10, −5, 0