Adaptive and pleiotropic effects of evolution in synonymous sugar environments.

Neetika Ahlawat[#,1], Pavithra Venkataraman[#,1,*], Anjali Mahilkar[1, 2], Raman Gulab Brajesh[1, 3], and Supreet Saini[1,*]

[1] Department of Chemical Engineering, Indian Institute of Technology Bombay, Mumbai 400 076

[2] Current address: Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI 41809

[3] Current address: Department of Biomedical Engineering and Bioinformatics, Swami Vivekanand Technical University, Durg 491107

[#] Authors contributed equally to this work.

* Corresponding Authors. Email: pavithrav@iitb.ac.in (PV); saini@che.iitb.ac.in (SS); Phone: +91 22 2576 7216

Running Title: Is pleiotropy predictable based on adaptation?

Abstract.

Adaptation to an environment is enabled by the accumulation of beneficial mutations.

When adapted populations are shifted to other environments, the byproduct or

pleiotropic fitness effects of these mutations can be wide-ranged. Since there exists

no molecular framework to quantify relatedness of environments, predicting

pleiotropic effects based on adaptation has been challenging. In this work, we ask if

evolution in highly similar environments elicits correlated adaptive and pleiotropic

responses. We evolve replicate populations of *Escherichia coli* in non-stressful

environments that contain either a mixture of glucose and galactose, lactose, or

melibiose as the source of carbon. We term these similar sugars as "synonymous",

since lactose and melibiose are disaccharides made up of glucose and galactose.

Therefore, the evolution environments differed only in the way carbon was presented

to the bacterial population. After 300 generations of evolution, we see that the

adaptive responses of these populations are not predictable. We investigate the

pleiotropic effects of adaptation in a range of non-synonymous environments, and

show that despite uncorrelated adaptive changes, the nature of pleiotropic effects is

largely predictable based on the fitness of the ancestor in the non-home

environments. Overall, our results highlight how subtle changes in the environment

can alter adaptation, but despite sequence-level variations, pleiotropy is qualitatively

predictable.

Keywords: Adaptation, pleiotropy, *E. coli*, synonymous sugars.

Introduction.

An evolving population accumulates mutations, and selection increases the frequency of beneficial mutations in the population. Mutational effects are contingent on the genetic background on which they occur, and the environment that the population is in. As a result, adaptation in one environment could lead to pleiotropic effects in the others. Pleiotropy was thought to operate only antagonistically[1,2]. However, many ecological and laboratory studies have shown the evolution of generalists, which are fit in a wide-range of environments. Thus, the potential beneficial effects of pleiotropy are important, especially to understand processes such as ecological specialization, genetic diversification, and movement on a rugged fitness landscape[3–9].

Several laboratory studies have focused on understanding the dynamics and variability of the pleiotropic effects of adaptation [10–20]. Two types of populations have been seen to evolve in these experiments – generalists and specialists. Generalists are populations that are viable in a range of environments, whereas specialists can grow well only in the home environment. While the evolution of specialists could be due to antagonistic pleiotropy, generalists could evolve due to mutations in genes that affect a small set of phenotypes [14,21–27]. The relatedness of the evolution environment with the environments in which the pleiotropic effects of adaptation are tested, is important. For example, Bakerlee et. al. show that yeast populations that adapted in temperature stress showed increased fitness in a highly saline environment [20]. However, the same population's fitness is much lower when tested in an environment colder than the evolution environment. Clearly, due to the lack of generality, adaptation to a stress does not guarantee an identical change in fitness when exposed to other stresses.

It is therefore not well understood how pleiotropic effects change with environment. Or, in other words, we do not know what kind of environments drive the evolution of generalists or specialists. How do subtle changes in the evolution environment alter adaptation and pleiotropic effects? Is the nature and number of genetic changes in generalists predictable? And most importantly, how general is the behaviour of generalists?

In this study, we propose to answer these questions by carrying out laboratory

evolution using *Escherichia coli* in three evolution environments. The environments

differ in the source of carbon available for the microbes to feed on. The three

evolution environments contained one of the following sugars - (a) a mixture of

glucose and galactose, (b) lactose, (c) melibiose. Glucose and galactose, when

linked by a β-1,4-glycosidic and an α-1,6-glucosidic bond form lactose and melibiose

respectively. Since these three sugars are different combinations of glucose and

galactose, we term them as "synonymous" environments. Essentially, the evolution

environments differed only in the way glucose and galactose were presented to the

evolving population.

We evolved six replicate populations in each of the synonymous environments for

300 generations, and assayed the fitness of the evolved populations in synonymous

and non-synonymous sugar environments. Our results allow us to identify the

pleiotropic effects of adaptation in a non-stressful environment, and subsequently

comment on the predictability of the evolution of specialists/generalists. Most

importantly, in a novel attempt, we show that minute changes in resource packaging

can alter adaptation and the consequent pleiotropic effects.

Results.

Adaptation in synonymous sugars is non-identical.

Replicate populations of *E. coli* were evolved in M9 minimal media containing 0.2% of glucose-galactose (0.1% of each), lactose, or melibiose. After 300 generations, two fitness parameters – growth rate in the exponential phase and carrying capacity – were calculated.

As shown in **Figure 1**, the biomass accumulated at the end of 16 hours increased for all the populations relative to the ancestor (*p<0.05, one-tailed t-test*), and other than two melibiose-evolved cells, growth rate increased in all cases (*p<0.05, one-tailed t-test*). Interestingly, two out of the six melibiose-evolved populations' growth rates are lower than that of the ancestor (*p<0.05, one-tailed t-test*).

Our results show that the selection pressure imposed by synonymous environments is non-identical. In fact, the melibiose evolved populations show a qualitative change in adaptive response compared to the lactose and glucose-galactose evolved populations, whose fitness differ from each other only quantitatively.

Figure 1. Selection acts on different traits in different evolution environments. Six replicate populations of *E. coli,* founded from a single clone, were evolved in three different, yet synonymous sugar environments – glucose-galactose mixture, lactose, and melibiose.

Populations with an increased growth rate (r) are thought to be fast growers, while populations with an increased carrying capacity (K) are considered better competitors[28]. While intuition suggests that evolving populations should exhibit r-K trade-offs, there exists very little empirical evidence in support. Recently, Marshall et al. showed theoretically how and why r and K may covary positively[29]. However, Wei et al. show that reduction in environment quality leads to such "trade-ups" in r and K[30]. Our results show that trade-ups in r and K are commonly observed in populations that evolved in different sources of carbon, and that the rate of occurrences of trade-ups or trade-offs is dependent on the evolution environment.

Pleiotropic effects of adaptation in synonymous environments.

Next, we shift these three sets of evolved populations to non-home environments, that contain synonymous sugars as the source of carbon. We calculate the growth r and K at the end of 16 hours, assuming exponential growth. Fitness changes of twelve populations in non-home synonymous environments, relative to the ancestor, are as shown in **Figure 2a** and **2b.**

Next, the fitness gains for each population relative to the ancestor, in home and non-home environments were compared. Each such comparison can be classified into one of the following three categories: (a) identical to the ancestor, in magnitude and type of effect, (b) identical to the ancestor, in the type of effect but are different in magnitude, or (c) different from the ancestor, in the type and magnitude of effect. **Figure 2c & 2d** shows the proportion of these cases (out of 12), for r and K, for the three sets of evolved populations, when grown in non-home synonymous environments.

Figure 2. Pleiotropic responses of evolved populations in synonymous environments.
We tested the fitness (r and K) of the three sets of evolved populations in away
environments, in which the source of carbon was a synonymous sugar, and checked if
fitness gains in home and away environments were correlated. For each set of evolved
population, there were two synonymous away environments in our experiment. **(a)** and **(b)**
show the relative gains in home and away environments, in r and K, respectively. We count
the number of distinct types of pleiotropic effects. As shown in **(c)** and **(d)**, the relative
frequencies of the three types of pleiotropic responses are different for different sets of
evolved populations.

Correlated fitness gains are the most common in the lactose-evolved populations,

followed by the glucose-galactose-evolved populations. The melibiose-evolved

populations, show the most diverse range of pleiotropic responses – growth rates of

all the melibiose-evolved populations were non-identical in the synonymous
environments, and the carrying capacity of more than half of them was different from
that in melibiose environment.

By and large, our results agree with the observations made by Ostrowski et. al., that
adaptation in a sugar environment confers positive pleiotropic effects in other
sugars[16]. However, despite being shifted to synonymous environments, our evolved
populations show considerable variability in their pleiotropic responses. What
happens to variability in pleiotropic responses in non-synonymous sugars?

Pleiotropic effects of adaptation in non-synonymous environments.

The three sets of populations which evolved independently in melibiose, lactose, and
glucose-galactose were shifted to non-synonymous sugar environments and tested
for fitness. These non-synonymous environments consisted of one of pentoses
(arabinose, xylose), a methyl pentose (rhamnose), sugar alcohols (glycerol, sorbitol),
hexoses (fructose, mannose), or a trisaccharide (raffinose) as the source of carbon.
Again, we measured r in exponential phase and K as cells enter stationary phase.
Fitness changes of the eighteen evolved populations, relative to that of the ancestor,
are shown in **Figure 3a** and **3b**. We found no environment in which the fitness of all
the evolved populations dropped compared to the ancestor. Arabinose and sorbitol
were two environments in which both r and K of all the evolved populations
increased relative to the ancestor. Therefore, all the populations that evolved in one
of the synonymous environments were fit in arabinose and sorbitol. However, we did
not find any set of evolved populations in which all the replicates performed
identically in home and any of the non-home environments. As a result, given a non-
home environment, we could only say if an evolved population is fitter than the
ancestor or not; we were unable to comment on the exact difference in fitness based
on the fitness of the evolved population in the home environment.

Figure 3. Pleiotropic responses of evolved populations in non-synonymous environments. We tested the fitness (r and K) of the three sets of evolved populations in away environments, in which the source of carbon was a non-synonymous sugar, and checked if fitness gains in home and away environments were correlated. For each set of evolved population, there were eight non-synonymous away environments in our experiment. **(a)** and **(b)** show the relative gains in home and away environments, in r and K, respectively. We count the number of distinct types of pleiotropic effects, and the results are as shown in **(c)** and **(d)**.

In the eight non-synonymous sources of carbon, glucose-galactose evolved populations showed the maximum number of similar fitness changes, followed by lactose-evolved populations. The melibiose-evolved populations continued to show

diverse responses in non-synonymous environments as well, as shown in **Figure 3c**

and **3d**.

Figure 4. Variability in pleiotropic responses. To test whether the pleiotropic fitness
changes (both *r* and *K*) of glucose-galactose, lactose, and melibiose evolved populations in
away environments (both synonymous and non-synonymous) follow any patterns, we
compared the relative frequencies of the three types of pleiotropic responses exhibited by
the three sets of evolved populations. Melibiose evolved populations exhibited the greatest
number of pleiotropic responses, followed by the lactose evolved populations. Overall, we
observe that even minute changes in the evolution environment play a significant role in
dictating the variability in pleiotropic responses.

Figure 4 shows the overall proportion of cases (considering both *r* and *K*, and

synonymous and non-synonymous environments) of pleiotropy for the three sets of

evolved populations. It is evident that the variability in the type of pleiotropic

responses depends on their evolution environment.

Fitness of the ancestor as a predictor of pleiotropic effects.

Global epistasis has enabled the prediction of the fitness effect of a beneficial

mutation as a function of the fitness of the background on which it occurs[31]. We tried

to identify if ancestral fitness can help predict pleiotropic effects. Specifically, we

checked if the pleiotropic fitness of an evolved population correlated with the fitness

of the ancestral population in any given environment. Therefore, we compared the

fitness changes in the evolved populations with that of the ancestor, in both synonymous and non-synonymous environments.

a.

b.

c.

d.

Figure 5. Fitness of the ancestor could act as a predictor of pleiotropic fitness gains. We test if the fitness of the ancestral population in an environment can act as a predictor of the fitness of an evolved population. So, we compare the fitness gain of an evolved population with ancestral fitness in away environments. **(a)** and **(b)** show these comparisons for r and K in synonymous environments, respectively. It must be noted that for each set of evolved populations, there were two synonymous away environments. We also plot relative fitness gains against ancestral fitness in eight non-synonymous environments. As shown in **(c)** and **(d)**, in most cases, there exists a trend – in non-synonymous environments, average pleiotropic fitness gains decrease with an increase in the fitness of the ancestor. However, there exists no such quantitative trend in the case of synonymous environments.

In synonymous environments, there existed no correlation between ancestral fitness and the nature, or number of pleiotropic responses, as shown in **Figure 5a and 5b**.

274 The three sets of evolved populations showed qualitatively similar pleiotropic

275 responses in the non-synonymous environments (refer to Supplementary Figures 1

276 & 2). As shown in **Figures 5c and 5d**, in non-synonymous environments, average

277 gains in r and K correlated negatively with ancestral fitness.

279 a.

  b.

281
Figure 6. Fitness of the ancestor could act as a predictor of the variability in
pleiotropic fitness gains. We showed that the fitness of the ancestor could as a rough
predictor of the average pleiotropic fitness gain. Next, to test if the variability in pleiotropic
responses can also be predicted in a similar fashion, we plot the range of fitness gains in r
(as shown in **(a)**) and K (as shown in **(b)**) against ancestral fitness in the eight non-
synonymous environments. Range was calculated as the difference between the maximum
and minimum fitness gains in the eighteen evolved populations. Variability in pleiotropic
responses, like mean pleiotropic fitness gain, also decreases with increase in ancestor's
fitness, in most cases.
291

We next check if the variability in pleiotropic responses of these evolved populations

is predictable based on ancestor's fitness in the non-home environment. To do so,

we consider the range of fitness gains (in r and K), calculated as the difference

between the maximum and minimum of the mean fitness gain, as a proxy for

variability in pleiotropic responses. As shown in **Figure 6a** and **6b**, as the fitness of

the ancestor increases in the non-home environment, the variability in pleiotropic

responses decreases in most cases.

Therefore, given the fitness of the ancestor in a non-home environment, we show

that a qualitative prediction of pleiotropic effects of an evolved population is possible.

The underlying genetic diversity explains the nature of pleiotropic responses.

We genome sequenced the eighteen evolved populations to identify the genetic changes accumulated in the three synonymous evolution environments. Mutation analyses reveal that all the mutations in the *rpo* genes were the most common among all sets of evolved populations. *Rpo* genes, that code for proteins that make up RNA polymerase, are known to facilitate adaptation in thermal, osmotic, and antibiotic stresses[32]. Other mutational targets included *yafW* which is a part of a toxin-antitoxin pair, *nanR* and *acrR*, which are transcription regulators[33,34], and the methionine transporter *metN*[35]. Overall, the melibiose-evolved populations had the highest number of distinct mutational targets, as shown in **Figure 7a** (Refer to Supplementary Table 1 for the detailed list of mutations). **Figure 7b** shows the correlated between the proportion of pleiotropic responses and the number of mutational targets.

a.

	Evolution environment		
	Glu-gal	Lactose	Melibiose
rpo(B/C/S	✓	✓	✓
nanR	✓	✓	✓
metN	✓	✓	✓
acrR	✓	✓	✓
yafW	✓	✓	✓
nagB	✓		
yhcH	✓		
yddA	✓		
ydgA	✓		
IS30		✓	
tar		✓	
rplI		✓	
tldD		✓	
rhsC		✓	✓
kgtP		✓	✓
ecpD			✓
relA			✓
rpsB			✓
kup			✓
pyrE			✓
rph			✓
aceE			✓
gor			✓
pstB			✓

Mutational target genes

b.

Figure 7. Genetic basis of pleiotropic effects. (a) The mutational targets in the three sets of evolved populations are shown. (b) The number of distinct mutational targets correlates with the proportion of pleiotropic responses. In other words, our results show that with an

We counted the number of instances where the three sets of evolved populations

showed any type of pleiotropy (change in magnitude and/or sign of fitness effect),

across the ten non-home environments (two synonymous, eight non-synonymous)

their fitness was measured in. Considering changes in both r and K, we see that

pleiotropic effects are the maximum for the melibiose-evolved populations, as shown

in **Figure 7b**. This proportion of pleiotropic responses, along with that of lactose-

evolved and glucose-galactose evolved populations, correlated predictably with the

number of distinct genes in which adaptive mutations occurred in replicate

populations.

Conclusions and Discussion.

Pleiotropic consequences of adaptation are expected to vary with changes in the duration of evolution[10,12,17,18,20]. It has been shown that evolving in a particular environment for a long period of time would lead to the rise of a specialist population, and not a generalist. However, the role of environment in dictating the evolution of a particular type of population - specialists or generalists - is not well understood. Also, there exists no method to quantify the relatedness of environments, making the prediction of pleiotropy challenging.

To probe the influence of environmental factors in dictating evolutionary trajectories and pleiotropic effects, we evolve six replicate populations of *E. coli* for a short period of time (300 generations) to capture the early onset of pleiotropy. The evolution environments were not harsh – the ancestral cell was viable in the three environments that we picked. The environments differed in the source of carbon present for the evolving population to utilize. In this study, we presented to the population two monosaccharides (glucose and galactose) packaged in three different ways – a mixture of glucose and galactose, lactose, and melibiose. These "synonymous" sugar environments allow us to maintain similarity in the evolution environments.

In a novel attempt, we show that evolution in synonymous sugars does not elicit identical adaptive responses. It must be noted that lactose and melibiose are utilized almost identically by *E. coli* – they are first imported into the cell with the help of a permease, a galactosidase then breaks them down into glucose and galactose, which are further metabolized [36,37]. Despite this similarity, we see that the pressure to accumulate mutations that improve biomass is the strongest in the melibiose environment. In fact, in some cases, evolution in melibiose led to a decrease in growth rate. This is one of the few cases where a trade-off between K and r has been empirically reported. It is noteworthy that this trade-off has been observed because of evolution in a non-stress environment. Commonly, growth rate is used as a proxy for fitness. Our observations underscore the need to look at other traits on which selection could be acting on evolving populations.

Fluctuating environments are known to generate genetic diversity[38], and alter gene

expression via mutations in global regulators[39]. By pointing at how minute changes in

the evolution environment can also generate diverse genotypic changes, we have

identified a simple yet significant driver of the process of generation of biodiversity in

prokaryotes.

We expected evolution in synonymous sugars to elicit positive pleiotropic responses

in synonymous environments[19]. But our fitness assays indicate clearly that positive

pleiotropic fitness changes are common, in both synonymous and non-synonymous

environments. In fact, we also find cases where the fitness effect in a non-home

environment is greater than that in the home environment. For example, the growth

rates of two melibiose-evolved populations decreased compared to the ancestor, in

the home environment. However, both these populations showed an increase in

growth rate in almost all the non-home environments. Interestingly, there were

scenarios were populations that adapted in a synonymous environment *B* performed

better in *A* than any of the populations that adapted in *A.* Therefore, adaptation in a

foreign environment could enable access to a fitness peak inaccessible to an

evolving population.

Almost all the melibiose-evolved populations show considerable fitness gains in

other synonymous and non-synonymous environments. However, why did these

mutations not fix in the other synonymous environments? Recently, Kosterlitz *et. al.*

reported that shifting hosts may aid adaptation of a gene, which are blocked due to

epistatic interactions in the native host[40]. Our results point at similar roadblocks to

adaptation posed by the environment, and indicate how minute changes in the

environment can help overcome them. It remains to be seen how the adaptive

trajectories of these evolved populations differ, when allowed to evolve in a non-

home environment.

The diversity in pleiotropic responses depended both on the home and the non-

home environment. For example, in glycerol, all the eighteen evolved populations,

despite having non-identical mutational targets, accumulated biomass different from

that of the ancestor. In seven out of the eight non-synonymous sugars that we tested

the fitness of the evolved populations in, over 50% of the populations showed

pleiotropic effects.

Across all sets, we found more generalists than specialists at the end of our

evolution experiment that lasted only 300 generations. We attempted to identify if

ancestral fitness can serve as an indicator of the exact pleiotropic effect.

Surprisingly, we found no correlation of pleiotropic responses with ancestor fitness in

synonymous environments, but there exists a pattern in non-synonymous

environments. Much like how the theory of diminishing returns explains the fitness

effect of new beneficial mutations given the background fitness, we show that given

the fitness of the ancestor in an environment, it is possible to comment on the

qualitative nature of the pleiotropic fitness of an evolved population despite variations

and stochasticity at the genetic level brought about by adaptation. The number of

instances where pleiotropic effects were observed (in non-home environments),

whether antagonistic or not, was understandably correlated with the number of

distinct mutational targets, as reported in the past [14]. Adaptive mutations occurred in

global regulators like the *rpo* genes, but different SNPs in a gene elicited variable

adaptive and pleiotropic effects. Therefore, to quantify the exact pleiotropic fitness

effects, our results show that a sequence level mapping of fitness effects and

environments is necessary.

Studies of pleiotropy have largely focused on studying the effects of evolving in one

stress environment in another. Our experimental approach provides a framework to

study evolution and its consequences in related environments, while evolving the

microbial population in a non-stress environment. Our attempt, in this work, has been

focused on being able to quantify and predict pleiotropy in related environments

(sugar environments), and not in highly dissimilar environments. We aim to identify

the byproduct effects of these adapted populations in unrelated environments

(temperature stress, osmotic stress, etc.) in the future, and test how changing the

qualitative nature of the environment alters pleiotropic responses. It is also unclear

how the trends and correlations we observe will change with time, and if and when

the generalist populations will become specialists. Therefore, we continue to

propagate these bacterial populations in their respective environments for future

studies.

Overall, our work motivates further probe into the temporal variability of adaptation in

synonymous environments. The results we report show that global epistasis renders

ancestral fitness as a predictor for early effects of pleiotropy, and point at the need to

perform high-throughput evolution experiments to build a framework that can help

decipher pleiotropic effects in a range of non-home environments.

Material and Methods.

Strains used and media composition.

E. coli K12 MG1655 (ATCC 47076) was used in this study.

Evolution experiments.

The evolution experiment was conducted in M9 minimal media (composition: $Na_2HPO_4.7H_2O$ (12.8 g/L), KH_2PO_4 (3g/L), NaCl (0.5g/L), NH_4Cl (1g/L), 1M $MgSO_4$ and 0.1M $CaCl_2$), containing either a mixture of glucose and galactose (0.1% each) or lactose (0.1%) or melibiose (0.1%).

Six independent replicate populations of *E. coli* were evolved at 37ºC and at 250 rpm in three environments - glucose-galactose mixture, lactose, or melibiose. Evolution experiment was performed by serially transferring 50µL of culture into 4.95mL of fresh M9 media containing appropriate sugar, every 12 hours.

Fitness measurements.

To measure fitness, cells were grown in LB (1% tryptone, 0.5% yeast extract, 0.05% NaCl) and incubated at 37ºC for 12 hours with shaking at 250 rpm. Thereafter, cells were sub-cultured 1:100 in fresh M9 media containing 0.2% glycerol for 18 hours. To measure fitness, cells were cultured in M9 media containing the appropriate carbon environment (glucose-galactose, lactose, melibiose, L-arabinose, D-xylose, D-mannose, D-fructose, D-sorbitol, D-raffinose) to an initial OD600 of 0.01.

Cells were then transferred to a 96-well plate (Costar) and the plate was covered with a breathable membrane (Breathe-Easy) to prevent evaporation. The 96-well plate was incubated at 37ºC with shaking in a microplate reader (Tecan Infinite Pro 200). OD600 was measured every 30 minutes until each culture reached stationary phase (~20 hours).

Growth rate was measured as described by Choudhury and Saini[41]. Maximum OD attained in each line was used as a measure for biomass accumulation.

Statistical tests

The error bars in the plots shown correspond to standard deviation, unless specified otherwise. A comparison of two means was done using a two-tailed *t* test. A one-

484 tailed *t* test was used to ascertain if one mean is greater than or lesser than another.
485 In all cases, the significance level was set to 0.05.

487 **Whole-genome sequencing.**

489 Genomic DNA of ancestral and evolved populations of *E. coli* was isolated using the
490 standard phenol/chloroform DNA isolation protocol[42]. DNA quality and concentration
491 were measured after DNA isolation using Nanodrop Spectrophotometer (Eppendorf),
492 and by gel electrophoresis. Genomic DNA samples were sent for paired-end
493 sequencing using Illumina NovaSeq 6000, with an average read-depth of 151bp.

495 Based on the quality report of fastq files, sequences were trimmed to retain only high-
496 quality sequences for analysis. The adapter trimmed reads were aligned to the
497 reference genome, *Escherichia coli* (ATCC 47076) using Burrows Wheeler Aligner
498 (BWA). Each sample had a minimum coverage of more than 30x. Variant calling was
499 done for the samples using GATK and further annotated using SnpEff. Variants that
500 were present in the ancestral strain were filtered out manually. The remaining SNPs
501 were used for further analysis. Raw sequencing data is available at
502 ncbi.nlm.nih.gov/sra/PRJNA1022868.

505 **Acknowledgements.**

507 This work was funded by a DBT/Wellcome Trust (India Alliance) grant (Award No.

508 IA/S/19/2/504632) to SS. PV is supported by the Prime Minister's Research Fellowship

509 (PMRF ID 1302050). NA is supported by the Institute Post-Doctoral Fellowship by IIT

510 Bombay.

References.

1. Stearns, S. C. Trade-Offs in Life-History Evolution. *Funct. Ecol.* **3**, 259 (1989).
2. Levins, R. A. *Evolution in Changing Environments: Some Theoretical Explorations*. (Princeton Univ. Press, Princeton, NJ, 1974).
3. Lee, C. E. Evolutionary genetics of invasive species. *Trends Ecol. Evol.* **17**, 386–391 (2002).
4. Bedhomme, S., Hillung, J. & Elena, S. F. Emerging viruses: why they are not jacks of all trades? *Curr. Opin. Virol.* **10**, 1–6 (2015).
5. Kassen, R. The experimental evolution of specialists, generalists, and the maintenance of diversity. *J. Evol. Biol.* **15**, 173–190 (2002).
6. Schluter, D. Evidence for Ecological Speciation and Its Alternative. *Science* **323**, 737–741 (2009).
7. Forister, M. L., Dyer, L. A., Singer, M. S., Stireman, J. O. & Lill, J. T. Revisiting the evolution of ecological specialization, with emphasis on insect–plant interactions. *Ecology* **93**, 981–991 (2012).
8. Mitchell-Olds, T., Willis, J. H. & Goldstein, D. B. Which evolutionary processes influence natural genetic variation for phenotypic traits? *Nat. Rev. Genet.* **8**, 845–856 (2007).
9. Savolainen, O., Lascoux, M. & Merilä, J. Ecological genomics of local adaptation. *Nat. Rev. Genet.* **14**, 807–820 (2013).
10. Cooper, V. S. & Lenski, R. E. The population genetics of ecological specialization in evolving Escherichia coli populations. *Nature* **407**, 736–739 (2000).
11. Dillon, M. M., Rouillard, N. P., Van Dam, B., Gallet, R. & Cooper, V. S. Diverse phenotypic and genetic responses to short-term selection in evolving *Escherichia coli* populations: DIVERSE RESPONSES TO SELECTION OF *E. coli* POPULATIONS. *Evolution* **70**, 586–599 (2016).
12. Bailey, S. F. & Kassen, R. Spatial Structure of Ecological Opportunity Drives Adaptation in a Bacterium. *Am. Nat.* **180**, 270–283 (2012).
13. Leiby, N. & Marx, C. J. Metabolic Erosion Primarily Through Mutation Accumulation, and Not Tradeoffs, Drives Limited Evolution of Substrate Specificity in Escherichia coli. *PLoS Biol.* **12**, e1001789 (2014).
14. Kinsler, G., Geiler-Samerotte, K. & Petrov, D. A. Fitness variation across subtle environmental perturbations reveals local modularity and global pleiotropy of adaptation. *eLife* **9**, e61271 (2020).
15. Jasmin, J.-N., Dillon, M. M. & Zeyl, C. The yield of experimental yeast populations declines during selection. *Proc. R. Soc. B Biol. Sci.* **279**, 4382–4388 (2012).
16. Ostrowski, E. A., Rozen, D. E. & Lenski, R. E. PLEIOTROPIC EFFECTS OF BENEFICIAL MUTATIONS IN ESCHERICHIA COLI. *Evolution* **59**, 2343–2352 (2005).
17. Meyer, J. R. *et al.* PARALLEL CHANGES IN HOST RESISTANCE TO VIRAL INFECTION DURING 45,000 GENERATIONS OF RELAXED SELECTION: EVOLUTION UNDER RELAXED SELECTION FOR RESISTANCE. *Evolution* no-no (2010) doi:10.1111/j.1558-5646.2010.01049.x.
18. Novak, M., Pfeiffer, T., Lenski, R. E., Sauer, U. & Bonhoeffer, S. Experimental Tests for an Evolutionary Trade-Off between Growth Rate and Yield in *E. coli*. *Am. Nat.* **168**, 242–251 (2006).
19. Jerison, E. R., Nguyen Ba, A. N., Desai, M. M. & Kryazhimskiy, S. Chance and necessity in the pleiotropic consequences of adaptation for budding yeast. *Nat. Ecol. Evol.* **4**, 601–611 (2020).
20. Bakerlee, C. W., Phillips, A. M., Nguyen Ba, A. N. & Desai, M. M. Dynamics and variability in the pleiotropic effects of adaptation in laboratory budding yeast populations. *eLife* **10**, e70918 (2021).
21. Crozat, E. *et al.* Parallel Genetic and Phenotypic Evolution of DNA Superhelicity in Experimental Populations of Escherichia coli. *Mol. Biol. Evol.* **27**, 2113–2128 (2010).
22. Fumasoni, M. & Murray, A. W. The evolutionary plasticity of chromosome metabolism allows adaptation to constitutive DNA replication stress. *eLife* **9**, e51963 (2020).
23. Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E. & Desai, M. M. The dynamics of molecular evolution over 60,000 generations. *Nature* **551**, 45–50 (2017).
24. Huang, C.-J., Lu, M.-Y., Chang, Y.-W. & Li, W.-H. Experimental Evolution of Yeast for High-Temperature Tolerance. *Mol. Biol. Evol.* (2018) doi:10.1093/molbev/msy077.
25. Venkataram, S. *et al.* Development of a Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast. *Cell* **166**, 1585-1596.e22 (2016).
26. Tenaillon, O. *et al.* The Molecular Diversity of Adaptive Convergence. *Science* **335**, 457–461 (2012).
27. Rodríguez-Verdugo, A., Gaut, B. S. & Tenaillon, O. Evolution of Escherichia coli rifampicin resistance in an antibiotic-free environment during thermal stress. *BMC Evol. Biol.* **13**, 50 (2013).
28. Pianka, E. R. On r- and K-Selection. *Am. Nat.* **104**, 592–597 (1970).
29. Marshall, D. J., Cameron, H. E. & Loreau, M. Relationships between intrinsic population growth rate, carrying capacity and metabolism in microbial populations. *ISME J.* **17**, 2140–2143 (2023).

30. Wei, X. & Zhang, J. Environment-dependent pleiotropic effects of mutations on the maximum
growth rate r and carrying capacity K of population growth. *PLOS Biol.* **17**, e3000121 (2019).
31. Kryazhimskiy, S., Rice, D. P., Jerison, E. R. & Desai, M. M. Microbial evolution. Global epistasis
makes adaptation predictable despite sequence-level stochasticity. *Science* **344**, 1519–1522
(2014).
32. Hews, C. L., Cho, T., Rowley, G. & Raivio, T. L. Maintaining Integrity Under Stress: Envelope
Stress Response Regulation of Pathogenesis in Gram-Negative Bacteria. *Front. Cell. Infect.*
Microbiol. **9**, 313 (2019).
33. Chu, D., Roobol, J. & Blomfield, I. C. A theoretical interpretation of the transient sialic acid toxicity
of a nanR mutant of Escherichia coli. *J. Mol. Biol.* **375**, 875–889 (2008).
34. Amores, G. R., de Las Heras, A., Sanches-Medeiros, A., Elfick, A. & Silva-Rocha, R. Systematic
identification of novel regulatory interactions controlling biofilm formation in the bacterium
Escherichia coli. *Sci. Rep.* **7**, 16768 (2017).
35. Kadaba, N. S., Kaiser, J. T., Johnson, E., Lee, A. & Rees, D. C. The high-affinity E. coli
methionine ABC transporter: structure and allosteric regulation. *Science* **321**, 250–253 (2008).
36. Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol. Biol.* **3**,
318–356 (1961).
37. Pardee, A. B. An inducible mechanism for accumulation of melibiose in Escherichia coli. *J.*
Bacteriol. **73**, 376–385 (1957).
38. Yamamichi, M., Letten, A. D. & Schreiber, S. J. Eco-evolutionary maintenance of diversity in
fluctuating environments. *Ecol. Lett.* **26**, (2023).
39. Mahilkar, A., Venkataraman, P., Mall, A. & Saini, S. Experimental Evolution of Anticipatory
Regulation in Escherichia coli. *Front. Microbiol.* **12**, 796228 (2021).
40. Kosterlitz, O. *et al.* Evolutionary "Crowdsourcing": Alignment of Fitness Landscapes Allows for
Cross-species Adaptation of a Horizontally Transferred Gene. *Mol. Biol. Evol.* **40**, msad237
(2023).
41. Choudhury, D. & Saini, S. Evolution of Escherichia coli in different carbon environments for 2,000
generations. *J. Evol. Biol.* **32**, 1331–1341 (2019).
42. He, F. E. coli Genomic DNA Extraction. *BIO-Protoc.* **1**, (2011).