Adaptive and pleiotropic effects of evolution in synonymous sugar environments.

Neetika Ahlawat[#,1], Pavithra Venkataraman[#,1,*], Raman Gulab Brajesh[1, 2], and Supreet

Saini[1,*]

[1] Department of Chemical Engineering, Indian Institute of Technology Bombay, Mumbai 400

076

[2] Current address: Department of Biomedical Engineering and Bioinformatics, Swami

Vivekanand Technical University, Durg 491107

[#] Authors contributed equally to this work.

* Corresponding Authors. Email: pavithrav@iitb.ac.in (PV); saini@che.iitb.ac.in (SS); Phone:

+91 22 2576 7216

Running Title: Is pleiotropy predictable based on adaptation?

Abstract.

Adaptation to an environment is enabled by the accumulation of beneficial mutations. When adapted populations are shifted to other environments, the byproduct or pleiotropic fitness effects of these mutations can be wide-ranged. Since there exists no molecular framework to quantify relatedness of environments, predicting pleiotropic effects based on adaptation has been challenging. In this work, we ask if evolution in highly similar environments elicits correlated adaptive and pleiotropic responses. We evolve replicate populations of *Escherichia coli* in non-stressful environments that contain either a mixture of glucose and galactose, lactose, or melibiose as the source of carbon. We term these similar sugars as "synonymous", since lactose and melibiose are disaccharides made up of glucose and galactose. Therefore, the evolution environments differed only in the way carbon was presented to the bacterial population. After 300 generations of evolution, we see that the adaptive responses of these populations are not predictable. We investigate the pleiotropic effects of adaptation in a range of non-synonymous environments, and show that despite uncorrelated adaptive changes, the nature of pleiotropic effects is largely predictable based on the fitness of the ancestor in the non-home environments. Overall, our results highlight how subtle changes in the environment can alter adaptation, but despite sequence-level variations, pleiotropy is qualitatively predictable.

*Keywords***:** Adaptation, pleiotropy, *E. coli*, synonymous sugars.

Lay Summary

In nature, evolution in "similar" environments is believed to elicit identical responses. For

example, the arctic fox and ptarmigan, which are two unrelated species living in the arctic,

have evolved to turn white in the winters. They did not evolve this ability because they from

the common ancestor, but because the environment favoured this trait. In this work, we ask

what happens to evolving populations if there are minute changes in the environment, and

what are the consequences of adapting in these environments that are "almost identical", or

as we call them, "synonymous".

We evolve replicate populations of the bacteria *E. coli* in three synonymous environments,

and quantify their ability to grow in both synonymous and non-synonymous environments.

We see that evolution does not proceed in an identical fashion in these populations, and that

each environment favours a different trait. However, interestingly, in non-synonymous

environments, these three sets of populations perform almost identically, and their growth is

qualitatively predictable.

Our results show that even simple and subtle changes in the environment can act as drivers

of biodiversity.

Introduction

An evolving population accumulates mutations, and selection increases the frequency of beneficial mutations in the population. Mutational effects are contingent on the genetic background on which they occur (Bateson et al., 1909), and the environment that the population evolves in (Plate, 1910). As a result, adaptation in one environment could lead to pleiotropic effects in the others. Pleiotropy was thought to operate only antagonistically (Levins, 1974; Stearns, 1989). However, many ecological and laboratory studies have shown the evolution of generalists, which are fit in a wide-range of environments (Bedhomme et al., 2015; Forister et al., 2012; Geiler-Samerotte et al., 2020; Hall et al., 2006; Kassen, 2002; C. E. Lee, 2002; Mitchell-Olds et al., 2007; Savolainen et al., 2013; Schluter, 2009; Wang et al., 2015). Thus, the potential effects of pleiotropy are important, especially to understand processes such as ecological specialization (Friberg et al., 2008; Hardy & Forister, 2023; Ostrowski et al., 2007), genetic diversification (D.-C. Lee et al., 2023; Noda-Garcia et al., 2019; Solovieff et al., 2013), and movement on a rugged fitness landscape (W. Huang et al., 2012; Martin & Lenormand, 2006). Pleiotropic alleles have also been implicated in a range of human diseases (Alzheimer's Disease Genetics Consortium et al., 2018; Bellou et al., 2020; Byars & Voskarides, 2020; Sivakumaran et al., 2011), and are known to confer antibiotic resistance (Hershberg, 2017; Pietsch et al., 2017; Schenk et al., 2015; Xia et al., 2017).

Several laboratory studies have focused on understanding the dynamics and variability of the pleiotropic effects of adaptation (Bailey & Kassen, 2012; Bakerlee et al., 2021; Cooper & Lenski, 2000; Dillon et al., 2016; Jasmin et al., 2012; Jerison et al., 2020; Kinsler et al., 2020; Leiby & Marx, 2014; Meyer et al., 2010; Novak et al., 2006; Ostrowski et al., 2005). Two types of populations have been seen to evolve in these experiments – generalists and specialists. Generalists are populations that are viable in a range of environments, whereas specialists incur significant fitness costs in non-home environments. While the evolution of specialists could be due to antagonistic pleiotropy, generalists could evolve due to mutations in genes that affect a small set of phenotypes (Crozat et al., 2010; Fumasoni & Murray, 2020; Good et al., 2017; C.-J. Huang et al., 2018; Kinsler et al., 2020; Rodríguez-Verdugo et al., 2013; Tenaillon et al., 2012; Venkataram et al., 2016). The fitness effect of a mutation in an environment which resembles the evolution environment is expected to be similar, because of relatedness in the growth conditions. For example, Bakerlee et. al. show that yeast populations that adapted in temperature stress showed increased fitness in a highly saline environment (Bakerlee et al., 2021). However, the same population's fitness is much lower when tested in an environment colder than the evolution environment. Clearly, due to

the lack of generality, adaptation to a stress does not guarantee an identical change in

fitness when exposed to other stresses.

It is therefore not well understood how pleiotropic effects change with environment. In other

words, we do not know what kind of environments drive the evolution of generalists or

specialists. How do subtle changes in the evolution environment alter adaptation and

pleiotropic effects? Is the nature and number of genetic changes in generalists predictable?

And most importantly, how general is the behaviour of generalists?

In this study, we propose to answer these questions by carrying out laboratory evolution

using *Escherichia coli* in three evolution environments (Supplement Figure 1). The

environments differ in the source of carbon available for the microbes to feed on. The three

evolution environments contained one of the following sugars - (a) a mixture of glucose and

galactose, (b) lactose, (c) melibiose. Glucose and galactose, when linked by a β-1,4-

glycosidic and an α-1,6-glucosidic bond form lactose and melibiose respectively. Since these

three sugars are different combinations of glucose and galactose, we term them as

"synonymous" environments. Essentially, the evolution environments differed only in the way

glucose and galactose were presented to the evolving population.

We evolved six replicate populations in each of the synonymous environments for 300

generations, and assayed the fitness of the evolved populations in synonymous and non-

synonymous sugar environments. Our results allow us to identify the pleiotropic effects of

adaptation in a not-so-harsh environment, and subsequently comment on the predictability of

the evolution of specialists/generalists. Most importantly, in a novel attempt, we show that

minute changes in resource packaging can alter adaptation and the consequent pleiotropic

effects.

Results.

Adaptation in synonymous sugars is non-identical.

Replicate populations of *E. coli* were evolved in M9 minimal media containing 0.2% of

glucose-galactose (0.1% of each), lactose, or melibiose. After 300 generations, two fitness

parameters – growth rate in the exponential phase and carrying capacity – were calculated.

As shown in **Figure 1**, the biomass accumulated at the end of 16 hours increased for all the

populations relative to the ancestor (*p<0.05, one-tailed t-test*), and other than two melibiose-

evolved cells, growth rate increased in all cases (*p<0.05, one-tailed t-test*). Interestingly, two

out of the six melibiose-evolved populations' growth rates are lower than that of the ancestor

(*p<0.05, one-tailed t-test*).

Our results show that the selection pressure imposed by synonymous environments is non-

identical. In fact, the melibiose evolved populations show a qualitative change in adaptive

response compared to the lactose and glucose-galactose evolved populations, whose

fitness differ from each other only quantitatively.

139
Figure 1. Selection acts on different traits in different evolution environments. Six replicate
populations of *E. coli,* founded from a single clone, were evolved in three different, yet synonymous
sugar environments – glucose-galactose mixture, lactose, and melibiose. After 300 generations of
evolution, fitness assays show that the growth rates (*r*) and carrying capacities (*K*) of almost all the
evolved populations increased relative to the ancestor. However, the melibiose-evolved populations
showed a much larger increase in carrying capacity than the other two sets of populations. In fact, two
of the melibiose-evolved populations' growth rates decreased relative to the ancestor. Therefore, our
results show that selection acting on a population changes even with a minor change in the
environment.

Populations with an increased growth rate (r) are fast growers, while populations with an

increased carrying capacity (K) are considered better competitors (Pianka, 1970). While

intuition suggests that evolving populations should exhibit r-K trade-offs, there exists very

little empirical evidence in support. Recently, Marshall et al. showed theoretically how and

why r and K may covary positively (Marshall et al., 2023). However, Wei et al. show that

reduction in environment quality leads to such "trade-ups" in r and K (Wei & Zhang, 2019).

Our results show that trade-ups in r and K are commonly observed in populations that

evolved in different sources of carbon, and that the rate of occurrences of trade-ups or trade-

offs is dependent on the exact nature of the evolution environment.

Pleiotropic effects of adaptation in synonymous environments.

Next, we shift these three sets of evolved populations to non-home environments, that

contain synonymous sugars as the source of carbon. We calculate the growth r and K at the

end of 16 hours, assuming exponential growth. Fitness changes of twelve populations in

non-home synonymous environments, relative to the ancestor, are as shown in **Figure 2a**

and **2b.**

Next, the fitness gains for each population relative to the ancestor, in home and non-home

environments were compared. Each such comparison can be classified into one of the

following three categories: (a) identical to the ancestor, in magnitude and type of effect, (b)

identical to the ancestor, in the type of effect but is different in magnitude, or (c) different

from the ancestor, in the type and magnitude of effect. **Figure 2c & 2d** shows the proportion

of these cases (out of 12), for r and K, for the three sets of evolved populations, when grown

in non-home synonymous environments.

a.

Figure 2. Pleiotropic responses of evolved populations in synonymous environments. We tested the fitness (r and K) of the three sets of evolved populations in away environments, in which the source of carbon was a synonymous sugar, and checked if fitness gains in home and away environments were correlated. For each set of evolved population, there were two synonymous away environments in our experiment. **(a)** and **(b)** show the relative gains in home and away environments, in r and K, respectively. We count the number of distinct types of pleiotropic effects. As shown in **(c)** and **(d)**, the relative frequencies of the three types of pleiotropic responses are different for different sets of evolved populations.

Correlated fitness gains are the most common in the lactose-evolved populations, followed by the glucose-galactose-evolved populations. The melibiose-evolved populations, show the most diverse range of pleiotropic responses – growth rates of all the melibiose-evolved

196   populations were non-identical in the synonymous environments, and the carrying capacity
197   of more than half of them was different from that in the home environment.
198
199   By and large, our results agree with the observations made by Ostrowski et. al., that
200   adaptation in a sugar environment confers positive pleiotropic effects in other sugars
201   (Ostrowski et al., 2005). However, despite being shifted to synonymous environments, our
202   evolved populations show considerable variability in their pleiotropic responses. What
203   happens to variability in pleiotropic responses in non-synonymous sugars?
204
205   **Pleiotropic effects of adaptation in non-synonymous environments.**
206
The three sets of populations which evolved independently in melibiose, lactose, and
glucose-galactose were shifted to non-synonymous sugar environments and tested for
fitness. These non-synonymous environments consisted of one of pentoses (arabinose,
xylose), a methyl pentose (rhamnose), sugar alcohols (glycerol, sorbitol), hexoses (fructose,
mannose), or a trisaccharide (raffinose) as the source of carbon. Again, we measured r in
exponential phase and K as cells enter stationary phase. Fitness changes of the eighteen
evolved populations, relative to that of the ancestor, are shown in **Figure 3a** and **3b**. We
found no environment in which the fitness of all the evolved populations dropped compared
to the ancestor. Arabinose and sorbitol were two environments in which both r and K of all
the evolved populations increased relative to the ancestor. Therefore, all the populations that
evolved in one of the synonymous environments were fit in arabinose and sorbitol. However,
we did not find any set of evolved populations in which all the replicates performed
identically in home and any of the non-home environments. As a result, given a non-home
environment, we could only say if an evolved population is fitter than the ancestor or not; we
were unable to comment on the exact difference in fitness based on the fitness of the
evolved population in the home environment.

Figure 3. Pleiotropic responses of evolved populations in non-synonymous environments. We tested the fitness (r and K) of the three sets of evolved populations in away environments, in which the source of carbon was a non-synonymous sugar, and checked if fitness gains in home and away environments were correlated. For each set of evolved population, there were eight non-synonymous away environments in our experiment. **(a)** and **(b)** show the relative gains in home and away environments, in r and K, respectively. We count the number of distinct types of pleiotropic effects, and the results are as shown in **(c)** and **(d)**.

In the eight non-synonymous sources of carbon, glucose-galactose evolved populations showed the maximum number of similar fitness changes, followed by the lactose-evolved populations. The melibiose-evolved populations continued to show diverse responses in non-synonymous environments as well, as shown in **Figure 3c** and **3d**.

Figure 4 shows the overall proportion of cases (considering both *r* and *K*, and synonymous and non-synonymous environments) of pleiotropy for the three sets of evolved populations. It is evident that the variability in the type of pleiotropic responses depends on their evolution environment.

Fitness of the ancestor as a predictor of pleiotropic effects.

Global epistasis has enabled the prediction of the fitness effect of a beneficial mutation as a function of the fitness of the background on which it occurs (Kryazhimskiy et al., 2014). We tried to identify if ancestral fitness can help predict pleiotropic effects. Specifically, we checked if the pleiotropic fitness of an evolved population correlated with the fitness of the ancestral population in any given environment. Therefore, we compared the fitness changes in the evolved populations with that of the ancestor, in both synonymous and non-synonymous environments.

Figure 5. Fitness of the ancestor could act as a predictor of pleiotropic fitness gains. We test if
the fitness of the ancestral population in an environment can act as a predictor of the fitness of an
evolved population. We compare the fitness gain of an evolved population with ancestral fitness in
away environments. **(a)** and **(b)** show these comparisons for r and K in synonymous environments,
respectively. It must be noted that for each set of evolved populations, there were two synonymous
away environments. We also plot relative fitness gains against ancestral fitness in eight non-
synonymous environments. As shown in **(c)** and **(d)**, in most cases, there exists a trend – in non-
synonymous environments, average pleiotropic fitness gains decrease with an increase in the fitness
of the ancestor. However, there exists no such quantitative trend in the case of synonymous
environments.

In synonymous environments, there existed no correlation between ancestral fitness and the

nature of pleiotropic responses, as shown in **Figure 5a and 5b**. The three sets of evolved

populations showed qualitatively similar pleiotropic responses in the non-synonymous

environments (refer to Supplement Figures 2 & 3). As shown in **Figures 5c and 5d**, in non-

synonymous environments, average gains in r and K correlated negatively with ancestral

fitness.

296
297
Figure 6. Fitness of the ancestor could act as a predictor of the variability in pleiotropic fitness
gains. We showed that the fitness of the ancestor could as a rough predictor of the average
pleiotropic fitness gain. Next, to test if the variability in pleiotropic responses can also be predicted in
a similar fashion, we plot the range of fitness gains in r (as shown in **(a)**) and K (as shown in **(b)**)
against ancestral fitness in the eight non-synonymous environments. Range was calculated as the
difference between the maximum and minimum fitness gains in the eighteen evolved populations.
Variability in pleiotropic responses, like mean pleiotropic fitness gain, also decreases with increase in
ancestor's fitness, in most cases.
306

We next check if the variability in pleiotropic responses of these evolved populations is

predictable based on ancestor's fitness in the non-home environment. To do so, we consider

the range of fitness gains (in r and K), calculated as the difference between the maximum

and minimum of the mean fitness gain, as a proxy for variability in pleiotropic responses. As

shown in **Figure 6a** and **6b**, as the fitness of the ancestor increases in the non-home

environment, the variability in pleiotropic responses decreases in most cases. But variability

did not correlate with average fitness changes. Therefore, given the fitness of the ancestor in

a non-home environment, we show that a qualitative prediction of pleiotropic effects of an

evolved population is possible.

The underlying genetic diversity explains the nature of pleiotropic responses.

We genome sequenced the eighteen evolved populations to identify the genetic changes

accumulated in the three synonymous evolution environments. Mutation analyses reveal that

mutations in the *rpo* genes were the most common among all sets of evolved populations.

Rpo genes, that code for proteins that make up RNA polymerase, are known to facilitate

adaptation in thermal, osmotic, and antibiotic stresses (Hews et al., 2019). Other mutational

targets included *yafW* which is a part of a toxin-antitoxin pair, *nanR* and *acrR,* which are

transcription regulators (Amores et al., 2017; Chu et al., 2008), and the methionine

transporter *metN* (Kadaba et al., 2008). Overall, the melibiose-evolved populations had the

highest number of distinct mutational targets, as shown in **Figure 7a** (Supplement Table 1

lists the mutations in each population). **Figure 7b** shows the correlated between the

proportion of pleiotropic responses and the number of mutational targets.

We counted the number of instances where the three sets of evolved populations showed

any type of pleiotropy (change in magnitude and/or sign of fitness effect), across the ten

non-home environments (two synonymous, eight non-synonymous) their fitness was

measured in. Considering changes in both r and K, we see that pleiotropic effects are the

maximum for the melibiose-evolved populations, as shown in **Figure 7b**. This proportion of

pleiotropic responses, along with that of lactose-evolved and glucose-galactose evolved

populations, correlated predictably with the number of distinct genes in which adaptive

mutations occurred in replicate populations.

346　　　　　a.　　　　　　　　　　　　　b.

348
349 **Figure 7. Genetic basis of pleiotropic effects.** (a) The mutational targets in the three sets of
350 evolved populations are shown. (b) The number of distinct mutational targets correlates with the
351 proportion of pleiotropic responses. In other words, our results show that with an increase in the
352 number of distinct types of mutational targets, proportion of pleiotropic effects of the evolved
353 populations increases.
354

Conclusions and Discussion.

Pleiotropic consequences of adaptation are expected to vary with changes in the duration of evolution (Bailey & Kassen, 2012; Bakerlee et al., 2021; Cooper & Lenski, 2000; Meyer et al., 2010; Novak et al., 2006). It has been shown that evolving in a particular environment for a long duration leads to the rise of a specialist population (Bakerlee et al., 2021). However, the role of the environment in dictating the evolution of a particular type of population - specialists or generalists - is not well understood. Also, there exists no method to quantify the relatedness of environments, making the prediction of pleiotropy challenging.

To probe the influence of environmental factors in dictating evolutionary trajectories and pleiotropic effects, we evolve six replicate populations of *E. coli* for a short period of time (300 generations) to capture the early onset of pleiotropy. The evolution environments were not harsh – the ancestral cell was viable in the three sugar environments that we picked. The evolution of *E. coli* in glycerol and lactate (propane-based organic compounds) has been shown to yield identical phenotypic response (Fong et al., 2005). How widely applicable is this result? In this study, we presented to the population two monosaccharides (glucose and galactose) packaged in three different ways – a mixture of glucose and galactose, lactose, and melibiose. These "synonymous" sugar environments allow us to maintain similarity in the evolution environments.

In a novel attempt, we show that evolution in synonymous sugars does not elicit identical adaptive responses. It must be noted that lactose and melibiose are utilized almost identically by *E. coli* – they are first imported into the cell with the help of a permease, a galactosidase then breaks them down into glucose and galactose, which are further metabolized (Jacob & Monod, 1961; Pardee, 1957). Despite this similarity, we see that the pressure to accumulate mutations that improve biomass is the strongest in the presence of melibiose. In fact, in some cases, evolution in melibiose led to a decrease in growth rate. This is one of the few cases where a trade-off between K and r has been empirically reported (Beardmore et al., 2011; Fitzsimmons et al., 2010; Meyer et al., 2015; Novak et al., 2006; Reding-Roman et al., 2017; Warringer et al., 2011; Wei & Zhang, 2019). Commonly, growth rate is used as a proxy for fitness. Our observations underscore the need to look at other traits of evolving populations on which selection could be acting.

Fluctuating environments are known to generate genetic diversity (Abdul-Rahman et al., 2021; Ellegren & Galtier, 2016; Yamamichi et al., 2023), and alter gene expression via mutations in global regulators (López-Maury et al., 2008; Mahilkar et al., 2021; Podrabsky &

Somero, 2004). By pointing at how minute changes in the evolution environment can also

generate diverse genotypic changes, we have identified a simple yet significant driver of the

process of generation of biodiversity in prokaryotes.

We expected evolution in synonymous sugars to elicit positive pleiotropic responses in

synonymous environments (Jerison et al., 2020). Our fitness assays indicate clearly that

positive pleiotropic fitness changes are common, in both synonymous and non-synonymous

environments. In fact, we also find cases where the fitness effect in a non-home environment

is greater than that in the home environment. For example, the growth rates of two

melibiose-evolved populations decreased compared to the ancestor, in the home

environment. However, both these populations showed an increase in growth rate in almost

all the non-home environments. Interestingly, there were scenarios were populations that

adapted in a synonymous environment *B* performed better in *A* than any of the populations

that adapted in *A.* Therefore, moving to a not-so-different foreign environment could enable

access to a fitness peak inaccessible to an evolving population.

Almost all the melibiose-evolved populations show considerable fitness gains in other

synonymous and non-synonymous environments. However, why did these mutations not fix

in the other synonymous environments? Recently, Kosterlitz *et. al.* reported that shifting

hosts may aid adaptation of a gene, which are blocked due to epistatic interactions in the

native host (Kosterlitz et al., 2023). Our results point at similar roadblocks to adaptation

posed by the environment, and indicate how minute changes in the environment can help

overcome them. It remains to be seen how the adaptive trajectories of these evolved

populations differ, when allowed to evolve in a non-home environment for a long period of

time.

The diversity in pleiotropic responses depended both on the home and the non-home

environment. For example, in glycerol, all the eighteen evolved populations accumulated

biomass different from that of the ancestor. In seven out of the eight non-synonymous

sugars that we tested the fitness of the evolved populations in, over 50% of the populations

showed pleiotropic effects.

Across all sets, we found more generalists than specialists at the end of our evolution

experiment that lasted 300 generations. We attempted to identify if ancestral fitness can

serve as an indicator of the exact pleiotropic effect. Surprisingly, we found no correlation of

pleiotropic responses with ancestor fitness in synonymous environments, but there exists a

pattern in non-synonymous environments. Much like how the theory of diminishing returns

explains the fitness effect of new beneficial mutations given the background fitness (Diaz-
Colunga et al., 2023), we show that given the fitness of the ancestor in an environment, it is
possible to comment on the qualitative nature of the pleiotropic fitness of an evolved
population despite variations and stochasticity at the genetic level brought about by
adaptation. The number of instances where pleiotropic effects were observed (in non-home
environments), whether antagonistic or not, was understandably correlated with the number
of distinct mutational targets, as reported in the past (Kinsler et al., 2020). Adaptive
mutations occurred in global regulators like the *rpo* genes, but different SNPs in a gene
elicited variable adaptive and pleiotropic effects. Therefore, to quantify the exact pleiotropic
fitness effects, our results show that a sequence level mapping of fitness effects and
environments is necessary.
441
Studies of pleiotropy have largely focused on studying the effects of evolving in one stress
environment in another. Our experimental approach provides a framework to study evolution
and its consequences in related environments, while evolving the microbial population in a
non-stress environment. Our attempt, in this work, has been focused on being able to
quantify and predict pleiotropy in related environments (sugar environments), and not in
highly dissimilar environments. We aim to identify the byproduct effects of these adapted
populations in unrelated environments (temperature stress, osmotic stress, etc.) in the
future, and test how changing the qualitative nature of the environment alters pleiotropic
responses. It is also unclear how the trends and correlations we observe will change with
time, and if and when the generalist populations will become specialists. Therefore, we
continue to propagate these bacterial populations in their respective environments for future
studies.
454
455 Overall, our work motivates further probe into the temporal variability of adaptation in
456 synonymous environments. The results we report show that global epistasis renders
457 ancestral fitness as a predictor for early effects of pleiotropy, and point at the need to
458 perform high-throughput evolution experiments to build a framework that can help decipher
459 pleiotropic effects in a range of non-home environments.
460

Material and Methods.

Strains used and media composition.

E. coli K12 MG1655 (ATCC 47076) was used in this study.

Evolution experiments.

The evolution experiment was conducted in M9 minimal media (composition: $Na_2HPO_4.7H_2O$ (12.8 g/L), KH_2PO_4 (3g/L), NaCl (0.5g/L), NH_4Cl (1g/L), 1M $MgSO_4$ and 0.1M $CaCl_2$), containing either a mixture of glucose and galactose (0.1% each) or lactose (0.1%) or melibiose (0.1%).

Six independent replicate populations of *E. coli* were evolved at 37°C and at 250 rpm in three environments - glucose-galactose mixture, lactose, or melibiose. Evolution experiment was performed by serially transferring 50µL of culture into 4.95mL of fresh M9 media containing appropriate sugar, every 12 hours.

Fitness measurements.

To measure fitness, cells were grown in LB (1% tryptone, 0.5% yeast extract, 0.05% NaCl) and incubated at 37°C for 12 hours with shaking at 250 rpm. Thereafter, cells were sub-cultured 1:100 in fresh M9 media containing 0.2% glycerol for 18 hours. To measure fitness, cells were cultured in M9 media containing the appropriate carbon environment (glucose-galactose, lactose, melibiose, L-arabinose, D-xylose, D-mannose, D-fructose, D-sorbitol, D-raffinose) to an initial OD600 of 0.01.

Cells were then transferred to a 96-well plate (Costar) and the plate was covered with a breathable membrane (Breathe-Easy) to prevent evaporation. The 96-well plate was incubated at 37°C with shaking in a microplate reader (Tecan Infinite Pro 200). OD600 was measured every 30 minutes until each culture reached stationary phase (~20 hours).

Growth rate was measured as described in the past (Choudhury & Saini, 2019). Maximum OD attained in each line was used as a measure for biomass accumulation. There was no observable lag phase in the glucose-galactose and lactose-evolved populations, as observed in another study (Choudhury & Saini, 2019). Therefore, we did not consider it as a measure of fitness change.

Statistical tests.

The error bars in the plots shown correspond to standard deviation, unless specified otherwise. A comparison of two means was done using a two-tailed *t* test. A one-tailed *t* test was used to ascertain if one mean is greater than or lesser than another. In all cases, the significance level was set to 0.05.

Whole-genome sequencing.

Genomic DNA of ancestral and evolved populations of *E. coli* was isolated using the standard phenol/chloroform DNA isolation protocol (He, 2011). DNA quality and concentration were measured after DNA isolation using Nanodrop Spectrophotometer (Eppendorf), and by gel electrophoresis. Genomic DNA samples were sent for paired-end sequencing using Illumina NovaSeq 6000, with an average read-depth of 151bp.

Based on the quality report of fastq files, sequences were trimmed to retain only high-quality sequences for analysis. The adapter trimmed reads were aligned to the reference genome, *Escherichia coli* (ATCC 47076) using Burrows Wheeler Aligner (BWA). Each sample had a minimum coverage of more than 30x. Variant calling was done for the samples using GATK and further annotated using SnpEff. Variants that were present in the ancestral strain were filtered out manually. The remaining SNPs were used for further analysis. Raw sequencing data is available at ncbi.nlm.nih.gov/sra/PRJNA1022868.

523 **Acknowledgements.**

525 This work was funded by a DBT/Wellcome Trust (India Alliance) grant (Award No. IA/S/19/2/504632)

526 to SS. PV is supported by the Prime Minister's Research Fellowship (PMRF ID 1302050). NA is

527 supported by the Institute Post-Doctoral Fellowship by IIT Bombay.

References.

Abdul-Rahman, F., Tranchina, D., & Gresham, D. (2021). Fluctuating environments maintain genetic diversity

through neutral fitness effects and balancing selection. *Molecular Biology and Evolution*, *38*(10),

4362–4375.

Alzheimer's Disease Genetics Consortium, Chung, J., Zhang, X., Allen, M., Wang, X., Ma, Y., Beecham, G.,

Montine, T. J., Younkin, S. G., Dickson, D. W., Golde, T. E., Price, N. D., Ertekin-Taner, N., Lunetta, K. L.,

Mez, J., Mayeux, R., Haines, J. L., Pericak-Vance, M. A., Schellenberg, G., … Farrer, L. A. (2018).

Genome-wide pleiotropy analysis of neuropathological traits related to Alzheimer's disease.

Alzheimer's Research & Therapy, *10*(1), 22. https://doi.org/10.1186/s13195-018-0349-z

Amores, G. R., de Las Heras, A., Sanches-Medeiros, A., Elfick, A., & Silva-Rocha, R. (2017). Systematic

identification of novel regulatory interactions controlling biofilm formation in the bacterium

Escherichia coli. *Scientific Reports*, *7*(1), 16768. https://doi.org/10.1038/s41598-017-17114-6

Bailey, S. F., & Kassen, R. (2012). Spatial Structure of Ecological Opportunity Drives Adaptation in a Bacterium.

The American Naturalist, *180*(2), 270–283. https://doi.org/10.1086/666609

Bakerlee, C. W., Phillips, A. M., Nguyen Ba, A. N., & Desai, M. M. (2021). Dynamics and variability in the

pleiotropic effects of adaptation in laboratory budding yeast populations. *eLife*, *10*, e70918.

https://doi.org/10.7554/eLife.70918

Bateson, W., Mendel, G., & Leighton, A. G. (1909). *Mendel's principles of heredity, by W. Bateson*. University

Press. https://doi.org/10.5962/bhl.title.1057

Beardmore, R. E., Gudelj, I., Lipson, D. A., & Hurst, L. D. (2011). Metabolic trade-offs and the maintenance of

the fittest and the flattest. *Nature*, *472*(7343), 342–346. https://doi.org/10.1038/nature09905

Bedhomme, S., Hillung, J., & Elena, S. F. (2015). Emerging viruses: Why they are not jacks of all trades? *Current*

Opinion in Virology, *10*, 1–6. https://doi.org/10.1016/j.coviro.2014.10.006

Bellou, E., Stevenson-Hoare, J., & Escott-Price, V. (2020). Polygenic risk and pleiotropy in neurodegenerative

diseases. *Neurobiology of Disease*, *142*, 104953.

Byars, S. G., & Voskarides, K. (2020). Antagonistic pleiotropy in human disease. *Journal of Molecular Evolution*,

88(1), 12–25.

Choudhury, D., & Saini, S. (2019). Evolution of Escherichia coli in different carbon environments for 2,000

generations. *Journal of Evolutionary Biology*, *32*(12), 1331–1341. https://doi.org/10.1111/jeb.13517

Chu, D., Roobol, J., & Blomfield, I. C. (2008). A theoretical interpretation of the transient sialic acid toxicity of a

nanR mutant of Escherichia coli. *Journal of Molecular Biology*, *375*(3), 875–889.

https://doi.org/10.1016/j.jmb.2007.10.073

Cooper, V. S., & Lenski, R. E. (2000). The population genetics of ecological specialization in evolving Escherichia

coli populations. *Nature*, *407*(6805), 736–739. https://doi.org/10.1038/35037572

Crozat, E., Winkworth, C., Gaffe, J., Hallin, P. F., Riley, M. A., Lenski, R. E., & Schneider, D. (2010). Parallel

Genetic and Phenotypic Evolution of DNA Superhelicity in Experimental Populations of Escherichia

coli. *Molecular Biology and Evolution*, *27*(9), 2113–2128. https://doi.org/10.1093/molbev/msq099

Diaz-Colunga, J., Skwara, A., Gowda, K., Diaz-Uriarte, R., Tikhonov, M., Bajic, D., & Sanchez, A. (2023). Global

epistasis on fitness landscapes. *Philosophical Transactions of the Royal Society B*, *378*(1877),

20220053.

Dillon, M. M., Rouillard, N. P., Van Dam, B., Gallet, R., & Cooper, V. S. (2016). Diverse phenotypic and genetic

responses to short-term selection in evolving *Escherichia coli* populations: DIVERSE RESPONSES TO

SELECTION OF *E. coli* POPULATIONS. *Evolution*, *70*(3), 586–599. https://doi.org/10.1111/evo.12868

Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*, *17*(7), 422–433.

Fitzsimmons, J. M., Schoustra, S. E., Kerr, J. T., & Kassen, R. (2010). Population consequences of mutational

events: Effects of antibiotic resistance on the r/K trade-off. *Evolutionary Ecology*, *24*(1), 227–236.

https://doi.org/10.1007/s10682-009-9302-8

Fong, S. S., Joyce, A. R., & Palsson, B. Ø. (2005). Parallel adaptive evolution cultures of Escherichia coli lead to

convergent growth phenotypes with different gene expression states. *Genome Research*, *15*(10),

1365–1372. https://doi.org/10.1101/gr.3832305

Forister, M. L., Dyer, L. A., Singer, M. S., Stireman, J. O., & Lill, J. T. (2012). Revisiting the evolution of ecological

specialization, with emphasis on insect–plant interactions. *Ecology*, *93*(5), 981–991.

https://doi.org/10.1890/11-0650.1

Friberg, M., Bergman, M., Kullberg, J., Wahlberg, N., & Wiklund, C. (2008). Niche separation in space and time

between two sympatric sister species—A case of ecological pleiotropy. *Evolutionary Ecology*, *22*, 1–

18.

Fumasoni, M., & Murray, A. W. (2020). The evolutionary plasticity of chromosome metabolism allows

adaptation to constitutive DNA replication stress. *eLife*, *9*, e51963.

https://doi.org/10.7554/eLife.51963

Geiler-Samerotte, K. A., Li, S., Lazaris, C., Taylor, A., Ziv, N., Ramjeawan, C., Paaby, A. B., & Siegal, M. L. (2020).

Extent and context dependence of pleiotropy revealed by high-throughput single-cell phenotyping.

PLoS Biology, *18*(8), e3000836.

Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E., & Desai, M. M. (2017). The dynamics of molecular

evolution over 60,000 generations. *Nature*, *551*(7678), 45–50. https://doi.org/10.1038/nature24287

Hall, M. C., Basten, C. J., & Willis, J. H. (2006). Pleiotropic quantitative trait loci contribute to population

divergence in traits associated with life-history variation in Mimulus guttatus. *Genetics*, *172*(3), 1829–

1844.

Hardy, N. B., & Forister, M. L. (2023). Niche specificity, polygeny, and pleiotropy in herbivorous insects. *The*

American Naturalist, *201*(3), 376–388.

He, F. (2011). E. coli Genomic DNA Extraction. *BIO-PROTOCOL*, *1*(14). https://doi.org/10.21769/BioProtoc.97

Hershberg, R. (2017). Antibiotic-Independent Adaptive Effects of Antibiotic Resistance Mutations. *Trends in*

Genetics, *33*(8), 521–528. https://doi.org/10.1016/j.tig.2017.05.003

Hews, C. L., Cho, T., Rowley, G., & Raivio, T. L. (2019). Maintaining Integrity Under Stress: Envelope Stress

Response Regulation of Pathogenesis in Gram-Negative Bacteria. *Frontiers in Cellular and Infection*

Microbiology, *9*, 313. https://doi.org/10.3389/fcimb.2019.00313

Huang, C.-J., Lu, M.-Y., Chang, Y.-W., & Li, W.-H. (2018). Experimental Evolution of Yeast for High-Temperature

Tolerance. *Molecular Biology and Evolution*. https://doi.org/10.1093/molbev/msy077

Huang, W., Werner, B., & Traulsen, A. (2012). The impact of random frequency-dependent mutations on the

average population fitness. *BMC Evolutionary Biology*, *12*, 160. https://doi.org/10.1186/1471-2148-

12-160

Jacob, F., & Monod, J. (1961). Genetic regulatory mechanisms in the synthesis of proteins. *Journal of Molecular*

Biology, *3*, 318–356. https://doi.org/10.1016/s0022-2836(61)80072-7

Jasmin, J.-N., Dillon, M. M., & Zeyl, C. (2012). The yield of experimental yeast populations declines during

selection. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1746), 4382–4388.

https://doi.org/10.1098/rspb.2012.1659

Jerison, E. R., Nguyen Ba, A. N., Desai, M. M., & Kryazhimskiy, S. (2020). Chance and necessity in the pleiotropic

consequences of adaptation for budding yeast. *Nature Ecology & Evolution*, *4*(4), 601–611.

https://doi.org/10.1038/s41559-020-1128-3

Kadaba, N. S., Kaiser, J. T., Johnson, E., Lee, A., & Rees, D. C. (2008). The high-affinity E. coli methionine ABC

transporter: Structure and allosteric regulation. *Science (New York, N.Y.)*, *321*(5886), 250–253.

https://doi.org/10.1126/science.1157987

Kassen, R. (2002). The experimental evolution of specialists, generalists, and the maintenance of diversity.

Journal of Evolutionary Biology, *15*(2), 173–190. https://doi.org/10.1046/j.1420-9101.2002.00377.x

Kinsler, G., Geiler-Samerotte, K., & Petrov, D. A. (2020). Fitness variation across subtle environmental

perturbations reveals local modularity and global pleiotropy of adaptation. *eLife*, *9*, e61271.

https://doi.org/10.7554/eLife.61271

Kosterlitz, O., Grassi, N., Werner, B., McGee, R. S., Top, E. M., & Kerr, B. (2023). Evolutionary "Crowdsourcing":

Alignment of Fitness Landscapes Allows for Cross-species Adaptation of a Horizontally Transferred

Gene. *Molecular Biology and Evolution*, *40*(11), msad237. https://doi.org/10.1093/molbev/msad237

Kryazhimskiy, S., Rice, D. P., Jerison, E. R., & Desai, M. M. (2014). Microbial evolution. Global epistasis makes

adaptation predictable despite sequence-level stochasticity. *Science (New York, N.Y.)*, *344*(6191),

1519–1522. https://doi.org/10.1126/science.1250939

Lee, C. E. (2002). Evolutionary genetics of invasive species. *Trends in Ecology & Evolution*, *17*(8), 386–391.

https://doi.org/10.1016/S0169-5347(02)02554-5

Lee, D.-C., Tai, J.-H., Lin, H.-F., Chao, T.-L., Ruan, Y., Cheng, Y.-W., Chou, Y.-C., Lin, Y.-Y., Chang, S.-Y., & Chen, P.-

J. (2023). Antagonistic pleiotropy plays an important role in governing the evolution and genetic

diversity of SARS-CoV-2. *bioRxiv*, 2023–02.

Leiby, N., & Marx, C. J. (2014). Metabolic Erosion Primarily Through Mutation Accumulation, and Not

Tradeoffs, Drives Limited Evolution of Substrate Specificity in Escherichia coli. *PLoS Biology*, *12*(2),

e1001789. https://doi.org/10.1371/journal.pbio.1001789

Levins, R. A. (1974). *Evolution in changing environments: Some theoretical explorations* (3. printing). Princeton

Univ. Press.

López-Maury, L., Marguerat, S., & Bähler, J. (2008). Tuning gene expression to changing environments: From

rapid responses to evolutionary adaptation. *Nature Reviews Genetics*, *9*(8), 583–593.

Mahilkar, A., Venkataraman, P., Mall, A., & Saini, S. (2021). Experimental Evolution of Anticipatory Regulation

in Escherichia coli. *Frontiers in Microbiology*, *12*, 796228. https://doi.org/10.3389/fmicb.2021.796228

Marshall, D. J., Cameron, H. E., & Loreau, M. (2023). Relationships between intrinsic population growth rate,

carrying capacity and metabolism in microbial populations. *The ISME Journal*, *17*(12), 2140–2143.

https://doi.org/10.1038/s41396-023-01543-5

Martin, G., & Lenormand, T. (2006). The fitness effect of mutations across environments: A survey in light of

fitness landscape models. *Evolution*, *60*(12), 2413–2427.

Meyer, J. R., Agrawal, A. A., Quick, R. T., Dobias, D. T., Schneider, D., & Lenski, R. E. (2010). PARALLEL CHANGES

IN HOST RESISTANCE TO VIRAL INFECTION DURING 45,000 GENERATIONS OF RELAXED SELECTION:

EVOLUTION UNDER RELAXED SELECTION FOR RESISTANCE. *Evolution*, no-no.

https://doi.org/10.1111/j.1558-5646.2010.01049.x

Meyer, J. R., Gudelj, I., & Beardmore, R. (2015). Biophysical mechanisms that maintain biodiversity through

trade-offs. *Nature Communications*, *6*, 6278. https://doi.org/10.1038/ncomms7278

Mitchell-Olds, T., Willis, J. H., & Goldstein, D. B. (2007). Which evolutionary processes influence natural genetic

variation for phenotypic traits? *Nature Reviews Genetics*, *8*(11), 845–856.

https://doi.org/10.1038/nrg2207

Noda-Garcia, L., Davidi, D., Korenblum, E., Elazar, A., Putintseva, E., Aharoni, A., & Tawfik, D. S. (2019). Chance

and pleiotropy dominate genetic diversity in complex bacterial environments. *Nature Microbiology*,

4(7), 1221–1230.

Novak, M., Pfeiffer, T., Lenski, R. E., Sauer, U., & Bonhoeffer, S. (2006). Experimental Tests for an Evolutionary

Trade-Off between Growth Rate and Yield in *E. coli*. *The American Naturalist*, *168*(2), 242–251.

https://doi.org/10.1086/506527

Ostrowski, E. A., Ofria, C., & Lenski, R. E. (2007). Ecological specialization and adaptive decay in digital

organisms. *The American Naturalist*, *169*(1), E1–E20.

Ostrowski, E. A., Rozen, D. E., & Lenski, R. E. (2005). PLEIOTROPIC EFFECTS OF BENEFICIAL MUTATIONS IN

ESCHERICHIA COLI. *Evolution*, *59*(11), 2343–2352. https://doi.org/10.1111/j.0014-

3820.2005.tb00944.x

Pardee, A. B. (1957). An inducible mechanism for accumulation of melibiose in Escherichia coli. *Journal of

Bacteriology*, *73*(3), 376–385. https://doi.org/10.1128/jb.73.3.376-385.1957

Pianka, E. R. (1970). On r- and K-Selection. *The American Naturalist*, *104*(940), 592–597.

https://doi.org/10.1086/282697

Pietsch, F., Bergman, J. M., Brandis, G., Marcusson, L. L., Zorzet, A., Huseby, D. L., & Hughes, D. (2017).

Ciprofloxacin selects for RNA polymerase mutations with pleiotropic antibiotic resistance effects.

Journal of Antimicrobial Chemotherapy, *72*(1), 75–84. https://doi.org/10.1093/jac/dkw364

Plate, L. (1910). Festschrift zum sechzigsten Geburtstag Richard Hertwigs. *Jena: G. Fischer*, *117*.

Podrabsky, J. E., & Somero, G. N. (2004). Changes in gene expression associated with acclimation to constant

temperatures and fluctuating daily temperatures in an annual killifish Austrofundulus limnaeus.

Journal of Experimental Biology, *207*(13), 2237–2254.

Reding-Roman, C., Hewlett, M., Duxbury, S., Gori, F., Gudelj, I., & Beardmore, R. (2017). The unconstrained

evolution of fast and efficient antibiotic-resistant bacterial genomes. *Nature Ecology & Evolution*, *1*(3),

0050. https://doi.org/10.1038/s41559-016-0050

Rodríguez-Verdugo, A., Gaut, B. S., & Tenaillon, O. (2013). Evolution of Escherichia coli rifampicin resistance in

an antibiotic-free environment during thermal stress. *BMC Evolutionary Biology*, *13*(1), 50.

https://doi.org/10.1186/1471-2148-13-50

Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation. *Nature Reviews*

Genetics, *14*(11), 807–820. https://doi.org/10.1038/nrg3522

Schenk, M. F., Witte, S., Salverda, M. L. M., Koopmanschap, B., Krug, J., & De Visser, J. A. G. M. (2015). Role of

pleiotropy during adaptation of TEM -1 $β$ -lactamase to two novel antibiotics. *Evolutionary*

Applications, *8*(3), 248–260. https://doi.org/10.1111/eva.12200

Schluter, D. (2009). Evidence for Ecological Speciation and Its Alternative. *Science*, *323*(5915), 737–741.

https://doi.org/10.1126/science.1160006

Sivakumaran, S., Agakov, F., Theodoratou, E., Prendergast, J. G., Zgaga, L., Manolio, T., Rudan, I., McKeigue, P.,

Wilson, J. F., & Campbell, H. (2011). Abundant pleiotropy in human complex diseases and traits. *The*

American Journal of Human Genetics, *89*(5), 607–618.

Solovieff, N., Cotsapas, C., Lee, P. H., Purcell, S. M., & Smoller, J. W. (2013). Pleiotropy in complex traits:

Challenges and strategies. *Nature Reviews Genetics*, *14*(7), 483–495.

Stearns, S. C. (1989). Trade-Offs in Life-History Evolution. *Functional Ecology*, *3*(3), 259.

https://doi.org/10.2307/2389364

Tenaillon, O., Rodríguez-Verdugo, A., Gaut, R. L., McDonald, P., Bennett, A. F., Long, A. D., & Gaut, B. S. (2012).

The Molecular Diversity of Adaptive Convergence. *Science*, *335*(6067), 457–461.

https://doi.org/10.1126/science.1212986

Venkataram, S., Dunn, B., Li, Y., Agarwala, A., Chang, J., Ebel, E. R., Geiler-Samerotte, K., Hérissant, L., Blundell,

J. R., Levy, S. F., Fisher, D. S., Sherlock, G., & Petrov, D. A. (2016). Development of a Comprehensive

Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast. *Cell*, *166*(6), 1585-1596.e22.

https://doi.org/10.1016/j.cell.2016.08.002

Wang, J., Atolia, E., Hua, B., Savir, Y., Escalante-Chong, R., & Springer, M. (2015). Natural variation in

preparation for nutrient depletion reveals a cost–benefit tradeoff. *PLoS Biology*, *13*(1), e1002041.

Warringer, J., Zörgö, E., Cubillos, F. A., Zia, A., Gjuvsland, A., Simpson, J. T., Forsmark, A., Durbin, R., Omholt, S.

W., Louis, E. J., Liti, G., Moses, A., & Blomberg, A. (2011). Trait variation in yeast is defined by

population history. *PLoS Genetics*, *7*(6), e1002111. https://doi.org/10.1371/journal.pgen.1002111

Wei, X., & Zhang, J. (2019). Environment-dependent pleiotropic effects of mutations on the maximum growth

rate r and carrying capacity K of population growth. *PLOS Biology*, *17*(1), e3000121.

https://doi.org/10.1371/journal.pbio.3000121

Xia, H., Tang, Q., Song, J., Ye, J., Wu, H., & Zhang, H. (2017). A *yigP* mutant strain is a small colony variant of *E.

coli* and shows pleiotropic antibiotic resistance. *Canadian Journal of Microbiology*, *63*(12), 961–969.

https://doi.org/10.1139/cjm-2017-0347

Yamamichi, M., Letten, A. D., & Schreiber, S. J. (2023). Eco-evolutionary maintenance of diversity in fluctuating

environments. *Ecology Letters*, *26*(S1). https://doi.org/10.1111/ele.14286